Characterizing the demographic history and prion protein variation to
infer susceptibility to chronic wasting disease in a naïve population of
white-tailed deer (*Odocoileus virginianus*)

Population and disease genomics of white-tailed deer.

Sarah E Haworth[1], Larissa Nituch[2], Joseph M Northrup[1,2], Aaron BA Shafer[1,3]*[⊠]

1 – Environmental and Life Sciences Graduate Program, Trent University, 7K9J 7B8 Peterborough, Canada

2 – Wildlife Research and Monitoring Section, Ontario Ministry of Natural Resources and Forestry, Trent University,
7K9J 7B8 Peterborough, Ontario, Canada

3 – Forensics Program, Trent University, K9J 7B8 Peterborough, Ontario, Canada

* – *corresponding author* aaronshafer@trentu.ca

⊠ – ORCID: https://orcid.org/0000-0001-7652-225X

Abstract

Assessments of the adaptive potential in natural populations are essential for understanding and
predicting responses to environmental stressors like climate change and infectious disease.
Species face a range of stressors in human-dominated landscapes, often with contrasting effects.
White-tailed deer (deer) are expanding in the northern part of their range following decreasing
winter severity and increasing forage availability. Chronic wasting disease (CWD), a prion disease
affecting cervids, is likewise expanding and represents a major threat to deer and other cervids.
We obtained tissue samples from free-ranging deer across their native range in Ontario, Canada
which has yet to detect CWD in wild populations of cervids. We used high-throughput
sequencing to assess neutral genomic variation, and variation in the PRNP gene that is partly
responsible for the protein misfolding when deer contract CWD. Neutral variation revealed a

high number of rare alleles and no population structure, and demographic models suggested a
rapid historical population expansion. Allele frequencies of PRNP variants associated with CWD
susceptibility and disease progression were evenly distributed across the landscape and
consistent with deer populations not infected with CWD. We then estimated the selection
coefficient of CWD, with simulations showing an observable and rapid shift in PRNP allele
frequencies that coincides with the start of a novel CWD epidemic. Sustained surveillance of
genomic and PRNP variation can be a useful tool for CWD-free regions where deer are managed
for ecological and economic benefits.

Keywords

RADseq, ungulate, PRNP, prion, Canadian wildlife, population genetics

Introduction

Human-induced environmental change has caused widespread alterations to ecological
and evolutionary processes (Harmon, Moran, & Ives, 2009; Pecl et al., 2017). Climate change is
expected to be the dominant driver of wildlife population declines and has been linked to broad-
scale biodiversity losses, but regional responses are often nuanced and context dependent (e.g.
Taylor et al., 2017; White, Gregovich, & Levi, 2017; Hashida et al., 2020). For example, climate
change in the Midwestern United States differentially favors the survival of two sympatric
populations of ungulates with similar selection pressures and life history traits, but differing
densities (Escobar, Moen, Craft, & VanderWaal, 2019; Weiskopf, Ledee, & Thompson, 2019). In
such instances, intraspecific genetic diversity and adaptive potential is crucial for long-term
population viability (Kardos & Shafer, 2018).

The emergence, spread, and persistence of infectious diseases in previously allopatric
populations is facilitated by climate change and other anthropogenic activity (Price et al., 2016;
Aguirre, 2017; Morand & Walther, 2020). The impacts of infectious disease on wildlife
populations are of interest to managers, especially if the affected species holds economic or
cultural value (Lambert et al., 2018; Weiskopf, Ledee, & Thompson, 2019). Preventing and
controlling diseases in free-ranging populations can, however, be complex and costly when they
are both naïve to the infectious disease and faced with climate change and other anthropogenic
activities (Herrera & Nunn, 2019; Miguel et al., 2020; Samuel et al., 2020). Selective pressures
are increased under these circumstances and populations are forced to respond to multiple
stressors simultaneously, or potentially face local extirpation (Fischer et al., 2020).

White-tailed deer (*Odocoileus virginianus*; deer) are the most widely distributed and
abundant ungulates in North America and hold significant economic and cultural value (Hewitt,
2011). The northern range of deer is primarily limited by snow but decreasing winter severity
has allowed deer to expand northward beyond their historical range limits (Dawe & Boutin,
2016; Kennedy-Slaney, Bowman, Walpole, & Pond, 2018). This expansion has implications for

ecosystems as, for example, deer herbivory alters long-term regional habitat characteristics and
plant communities (Frerker, Sabo, & Waller, 2017; Otsu, Iijima, & Nagaike, 2019; Kroeger et al.,
2020). Further, deer are an important prey species and predator populations increasing in
response to deer expansion has led to greater predation and apparent interspecific competition
with other ungulates (Latham et al., 2013; Barber-Meyer & Mech, 2016). Consequently,
northward expansions of deer are having profound impacts to ecosystems including facilitating
infectious pathogen and disease spread (Averill et al., 2018; Ferretti & Mori, 2020).

A widespread threat to deer in North America is the highly infectious and fatal
neurodegenerative prion disease called chronic wasting disease (CWD). CWD is the only prion
disease known to infect captive and free ranging species of cervids (family *Cervidae*) and has
been reported in North and South America, Europe, and South Korea (Haley et al., 2019). With
virtually no barriers to transmission and a lengthy infectious preclinical period, the local
prevalence of CWD in North America has been measured to be as high as 50% and 82% in wild
and captive populations, respectively (Miller et al., 2004; O'Rourke et al., 2004). Due to
constraints on CWD surveillance it is likely that the distribution and prevalence of CWD in wild
populations are underestimated (Escobar et al., 2020). It is clear the frequency and occurrence
of CWD has increased over time, in part driven by anthropogenic activities related to hunting
and wildlife farming (Osterholm et al., 2019).

Despite CWD being fatal there is inter-individual variation in susceptibility and clinical
progression. Susceptibility and clinical progression are associated with non-synonymous and
synonymous genetic variation in the functional prion protein gene (PRNP; Güere et al., 2020;
Chafin et al., 2020). Single nucleotide polymorphisms (SNPs) at nucleotide (nt) 60, nt153, nt285,
nt286, nt555, and nt676 in deer PRNP have been associated with altered CWD susceptibility or
pathogenic processes (Johnson et al., 2006a; Wilson et al., 2009; Brandt et al., 2015; Brandt
et al., 2018). The presence of CWD appears to affect population PRNP allele frequencies over space
and time due to selection (Robinson et al., 2012); however, altered CWD susceptibility and
pathogenic processes are clearly polygenic traits (Seabury et al. 2020) and disease spread is
different in structured populations, which might require different wildlife management practices
(Chafin et al., 2020).The efficacy of selection in the face of a novel pressure like CWD is
dependent on the effective population size (N_e). Common metrics to infer selection such as
Tajima's *D*, specifically measure shifts in allele frequencies across the site-frequency spectrum;
however, the frequency and proportion of rare alleles is sensitive to demographic processes
(Messer, Ellner, & Hairston Jr., 2016; Platt et al., 2019), and population genetics theory predicts
an excess of rare alleles in expanding populations (Gillespie, 2004).

Ontario, Canada reflects the northern leading edge of deer range in eastern North
America (Kennedy-Slaney, Bowman, Walpole, & Pond, 2018). The landscape of Ontario is
heterogenous and environmental clines exist around the Great Lakes region. Ontario has not
detected CWD in wild cervids, but CWD has been detected in farmed and captive cervids from

virtually every jurisdiction bordering Ontario, with the province using a weighted surveillance

approach to model CWD risk and strategically use resources for surveillance. Accordingly, at the

genome-level, we predicted an excess frequency of rare neutral and PRNP variants across our

study region given Ontario's deer population is expanding (Kennedy-Slaney, Bowman, Walpole,

& Pond, 2018). Based on large-scale distribution changes and recent population trends in deer

(Baldwin, Desloges, & Band, 2000; Latch et al., 2009), we predicted we would observe high

neutral genomic diversity (N_e) and demographic population expansion, indicating increased gene

flow and decreased population structure despite a heterogenous landscape. Since PRNP is not

under selection by CWD given the region is disease free, we predicted functional variation to

resemble regions most recently exposed to CWD (or still disease free). This is the first study to

characterize PRNP genetic variation and population genomic structure of wild deer, while also

determining the ancient and contemporary demographic and selection processes driving

patterns of diversity.

Materials and Methods

Study area and sample collection

We sampled white-tailed deer across Ontario, Canada (Figure 1). Between 2002-2018,

retropharyngeal lymph nodes were opportunistically extracted from hunter harvested deer

across Ontario through the CWD surveillance program managed by the Ontario Ministry of

Natural Resources and Forestry (OMNRF). Auxiliary data including year, sex, age-class, Mercator

grid cell unit (GCU; 10x10 km), and wildlife management unit (WMU) were also collected with

deer samples. North western and southern Ontario regions are geographically discontinuous for

deer (Figure 1), samples were therefore assigned to northern Ontario or southern Ontario

(Figure S1) for the purpose of analysis where sampling regions are compared. Preliminary

analysis of data did not warrant separating southeastern and southwestern Ontario as per

provincial management zones. Genomic DNA was extracted from deer samples using a silica-

based DNA extraction kit for tissue following manufacturers protocol (Qiagen, Cat. No. 69506)

and stored at -20°C. DNA quality was assessed by a spectrometer (NanoDrop 2000, Thermo

Scientific) and by 2% agarose GelRed gel electrophoresis.

Library preparation for PRNP genetic analysis

A 771 base-pair (bp) region of the deer prion protein precursor (PRNP) gene was

targeted and amplified using four degenerate primers (Table S1). Four replicate PCRs generated

a 460 base pair Fragment 1 and a 580 base pair Fragment 2 (Table S2). In a 2:1 ratio of Fragment

1 to Fragment 2, respectively, amplified DNA was added for each individual and then indexed

with standard Illumina multiplexing indices. Negative controls of UltraPure distilled water

(Invitrogen, 1897011) were used for each 96-well plate. The library was purified of artifacts

following manufacturers protocol for AMPure XP beads (Beckman Coulter, A63880) and
validated with a TapeStation D1000 kit (Agilent, 5067-5582). The library was sequenced on an
Illumina MiSeq platform at the University of Guelph Advanced Analysis Centre to generate 300
base pair (bp) pair-end reads for each sample.

Library preparation for RADseq genomic analysis

Restriction-site associated DNA sequencing (RADseq) libraries were generated using an
adapted protocol from Parchman et al. (2012) and Peterson et al. (2012) with Sbfl-HF and Msel
restriction enzymes. Samples were incubated, digested overnight, and heat-inactivated in 96-
well plates (Table S3). Negative controls of UltraPure distilled water (Invitrogen, 1897011) were
used for each 96-well plate. Restriction digested DNA was combined with 7 ul of ligation
mixture and 3 ul of one of the 24 available Sbfl adapters (1.0 uM). Adapters were ligated at 16°C
for 3 hours. DNA fragments were purified of artifacts following manufacturers protocol for
AMPure XP beads (Beckman Coulter, A63880). Adapter-ligated fragments were amplified in four
separate 10 ul reactions that incorporated barcodes. Reaction conditions and primers are
shown for ligation mixture and PCR in Table S4 and Table S5, respectively. Samples were pooled
and purification was performed following manufacturers protocol for QIAquick PCR Purification
kit (Qiagen, 28106) for a final elution to 42 ul. Size selection between 450 bp to 700 bp was
performed on 80 ul replicates of purified library and gel purification was performed following
manufacturers protocol for QIAquick Gel Extraction kit (Qiagen, 28706) for a final elution to 60
ul. The purified final library was validated with a TapeStation D1000 kit (Agilent, 5067-5582). The
libraries were sequenced at The Centre for Applied Genomics (TCAG) in The Hospital for Sick
Children (SickKids) on an Illumina HiSeq 2500 to produce 2x126 base pair paired end reads.

Bioinformatic pipeline and data analysis I – PRNP gene

The quality of reads was assessed using FastQC (Babraham Institute; v0.11.8). Samples
were excluded if at least one file in the pair-end files for a sample was less than 1kB in size or
failed to pass quality standards. A novel command line-based pipeline was developed to
assemble and genotype PRNP (accessible at https://gitlab.com/WiDGeT_TrentU). Our workflow
integrated Pullseq v1.0.2, BWA v0.7.17, SAMtools; v1.9, BCFtools v1.9, and VCFtools v0.1.16-15.
Briefly, for each sample, the pipeline extracted relevant reads based on the presence of primer
sequence, mapped the extracted reads to a 771 bp PRNP gene reference sequence. We
generated a consensus sequence and called single nucleotide polymorphisms (SNPs). SNP calls
were limited to positions where there was a minimum read depth of 30 and mapping quality
score of at least 30. Sanger sequencing of a subset of samples and their a priori called variants
were used to validate the bioinformatic pipeline.

The presence of asparagine (N) at aa138 (nt413A) indicates amplification of the
pseudogene (Brandt et al., 2015); we therefore filtered out all sequences with this site. SNPs

with a total frequency of occurrence of 1% or less were excluded from the analysis. A two-sided
Fisher's Exact Test was conducted on minor allele counts from either northern or southern
sampling regions at four well studied positions in the deer PRNP gene associated with CWD:
nucleotide (nt) 60, nt285, nt286, and nt676. Synonymous and non-synonymous sites were
identified using MEGA X v10.0.5. Haplotypes were estimated from unphased sequences with
PHASE v2.1.1 using a Markov chain Monte Carlo (MCMC) sampling approach with a minimum of
100,000 steps, with a discarded burn-in of 10,000, and samples were drawn every 100 MCMC
steps. Five repetitions were performed to verify consistent frequencies of haplotype assignment
(Brandt et al., 2018). Haplotypes with a frequency of less than 1% were removed. The genotype,
frequencies, and estimated standard deviations of the remaining haplotypes were analyzed as a
2x2 contingency table by sampling region.

Bioinformatic pipeline and data analysis II – RADseq

Fastq files were demultiplexed using process_radtags within the Stacks v2.3 module.
Parameters within process_radtags included the removal of any read with an uncalled base and
the discarding of reads with low quality scores. The demultiplexed sample files were aligned
against the deer genome (Genome Accession JAAVWD000000000) using BWA with samtools
used to sort, merge and compress BAM files. The referenced-based approaches on gstacks and
populations program within STACKs produced a variant call format (VCF) file with the restrictions
that the minimum percentage of individuals in a population required to process a locus for that
population was 90%. The VCF was filtered using VCFtools to only include reads with a minimum
read depth of 20. Population statistics, including F_{IS}, observed and estimated homozygosity, and
nucleotide diversity were calculated using the *populations* module. To analyze sources of
variation, we generated a principal component analysis (PCA) using the R v3.6.1 package
adegenet v2.1.3. A linear regression was run on principal component (PC) 1 and PC2 scores
against latitude and longitude. We estimated F_{ST} between north and south using StAMPP v1.6.1.
Population structure was detected using successive K-means clustering and a discriminant
analysis of principal components (DAPC) available in adegenet (Jombart, Devillard, & Balloux,
2010).

Demographic Analysis and Estimate of Effective Population Size

The final VCF was converted into 1D site frequency spectrum (SFS) for all of Ontario and
northern vs and southern designations, respectively, using vcf2dadi.py with projections for the
SFS estimated in easySFS. We applied a diffusion-based approach to demographic inference
through the Diffusion Approximation for Demographic Inference (δaδi) tool by Gutenkunst et al.,
(2009). Nine 1D models were assessed for Ontario as a single population. The optimum model
was selected as the lowest optimized log-likelihood of all successfully run models. δaδi was also
used to estimate the following summary statistics for the province: Watterson Theta (θ),
Tajima's *D* and the number of segregating sites. Using the mutation rate (μ) per site per

generation of a closely related species (*Rangifer tarandus* from Chen et al., 2019), total number
of sites (L), and the parameters estimated in the optimum model selected from δaδi, we
estimated the ancestral effective population size as $N_a = θ / 4μL$.

Estimation of selection on PRNP and allele frequency projections for a naïve population

We estimated the selection coefficient (*s*) at two sites (nt285 and nt286) using the
approach of Thompson et al. (2019) which accounts for number of generations and different
expression modes (i.e. recessive, dominant, codominant). Strength of selection against the less
resistant phenotype [i.e., the homozygous common allele (s_{AA})] can be estimated by calculating
values of s_{AA} that explained the estimated change in allele frequencies between positive and
negative animals from the same region. Here we used starting (negative CWD) allele frequencies
from Wilson et. al. (2009) and Kelly et al. (2009) and estimated *s* over *n* generations the
equation:

$p' = (1-s_{AA})p^2 + (1-s_{AB})p(1 − p)/(1-s_{AA})p^2 + (1-s_{AB})2p(1 − p) + (1-s_{BB})(1 − p)^2$

taken from Charlesworth and Charlesworth (2010). To account for uncertainty in time and allele
frequency estimates we ran 100 iterations with *n* ranging from 2-25 generations (~4-50 deer
years) and positive and negative allele frequencies (±1%) for those reported from both Wilson
et. al. (2009) and Kelly et al. (2009). Then using our estimated allele frequencies for Ontario, we
projected each allele frequencies into the next 25 generations using our estimated selection
coefficients, and the estimated *s* coefficients of 0.0103 and 0.074 from Robinson et al. (2012).
Calculations were conducted under three relative fitness scenarios as per Thompson et al.
(2019).

Results

PRNP Genetic analysis

A total of 631 Ontario deer samples were included in the PRNP genetic analysis (Figure
1). Nineteen SNPs were detected after filtering (Table 1), with 8 being non-synonymous
substitutions. Six of the detected variants in the PRNP gene have been linked to CWD
susceptibility or clinical progression, these include nt60, nt153, nt285, nt286, nt555, and nt676.
A two-sided Fisher's Exact test on the major and minor allele counts at the four important,
arguably the most studied, CWD-linked loci (nt60, nt285, nt286, and nt676) conducted between
Northern and Southern Ontario indicated that there was only a difference in frequency (p < 0.05)
at nt676 (Table S6).

There were 102 unique haplotypes with a count of at least one, with 12 haplotypes
having a frequency greater than 1% (Figure 2; Table S7). The two most common haplotypes,

Haplotype 3 (f=0.23) and Haplotype 1 (f=0.12) did not include any non-synonymous
substitutions. Haplotype A (f=0.30) and Haplotype B (f=0.25) reported by Brandt et al (2015,
2018) from northern Illinois were also detected as Haplotype 16 (f=0.09) and Haplotype 7
(f=0.05), respectively. The same Haplotype A (f=0.15) and Haplotype B (f=0.23) were reported by
Chafin et al., 2020 in deer from Arkansas, USA.

RADseq Genomic analysis

A total of 235 Ontario deer samples were sequenced in ddRADseq libraries. Following
quality control and quality assessment, including FastQC and line counts of demultiplexed files,
190 samples remained for downstream analysis (Figure 1). Estimated population diversity
statistics are summarized in Table 2. The PCA clearly separated northern and southern Ontario
along PC1 (Figure 3). A linear regression revealed that PC1 was strongly associated with
longitude (β = –0.79; adjusted R^2 = 0.77; p-value < 0.01). However, population structure between
northern and southern Ontario was weak (F_{ST}=0.02). A BIC based on the K-cluster analysis also
indicated that the most optimum number of clusters was 1.

The optimum 1D demographic model for Ontario was the BOTTLEGROWTH model,
which models an instantaneous size change followed by exponential change (Table S8; Figure 4).
From the 1D site frequency spectrum, the mean Tajima's D was estimated to be -2.126
consistent with a population expansion after a bottleneck. Using the calculated θ, we estimated
N_a to be ~20,000: this would place the timing of the population change (Tc) measured in $2N_a$
generations around the onset of the last glacial maximum (Figure 4).

PRNP selection and projection

We estimated the selection coefficients to be 0.08 (±0.06) and 0.11 (±0.07) for nt285
and nt286 under a dominance model. All simulated projections with our *s* values and the upper
reported value of Robinson et al (2012) showed a rapid shift in allele frequencies (Figure 5); the
majority of simulated trajectories did not overlap with the low *s* coefficient of 0.01 previously
reported by Robinson et al (2012).

Discussion

White-tailed deer in North America are intensively managed for hunter harvest and are
expanding their range northward due to climate change. The frequency and occurrence of CWD
infection in captive and free-ranging deer is likewise increasing (Osterholm et al., 2019; Rivera,
Brandt, Novakofski, & Mateus-Pinilla, 2019), but transmission and spread in free-ranging
populations are still poorly understood (Potapov, Merrill, Pybus, & Lewis, 2016). We observed
that the frequency of PRNP alleles in our naïve population differed from areas currently infected
with CWD or where CWD is endemic, including western Canada (Table S9; Kelly et al., 2008;
Wilson et al., 2009; Brandt et al., 2015; Brandt et al., 2018; Chafin et al., 2020).

The emergence, transmission, and persistence of highly infectious diseases in healthy
populations are often facilitated by climate change and exacerbated in areas with intense
anthropogenic activity (McKnight et al., 2017; Rizzoli et al., 2019). The introduction and re-
introduction of infectious diseases often results in rapid local population declines, reducing
species' adaptive potential and generating substantial economic losses (Belant & Deese, 2010;
Escobar, Moen, Craft, & VanderWaal, 2019). In Ontario deer, however, we observed an excess of
rare alleles with no evidence of strong population structure, including in southern Ontario where
an environmental cline exists around lake systems. An excess of rare alleles and high N_e and
absent population structure might provide the means for effective adaptation to selective
pressures, including climate change, infectious disease, and human activity.

Linking Population Demographics to Functional genes

We assessed contemporary and historical gene flow by examining neutral genomic
variation across tens of thousands of loci from deer across the province on Ontario. The data
were consistent with a large expanding population with a high level of genomic diversity and an
excess of rare alleles. These genomic patterns are also consistent with population simulations of
responses to climate change (Kennedy-Slaney, Bowman, Walpole, & Pond, 2018), and offers
some clues as to the potential adaptive response were CWD to arrive. Specifically, two
important features of the genomic data are noteworthy: the high number of a rare alleles and
the limited population structure across Ontario.

Demographic processes and life history strategies influence the proportion of rare
alleles, which are important to the adaptive process but are sensitive to recent evolutionary
processes (Peart et al. 2020). Evidence suggests the accumulation of rare alleles is independent
of taxa, but adaptation appears limited in low-diversity taxa (e.g., primates; Rousselle et al.,
2020). The deer genomic data evaluated here are consistent with high adaptive potential;
specifically, we calculated an N_a to be ~20,000, with the expansion estimated to be 100X
suggestive of a large species-wide N_e (derived either from π in Table 2 or the demographic model
in Figure 4).

Studies of populations infected with CWD often demonstrate a lack of allelic diversity at
the PRNP gene, which is thought to be due to CWD-driven selection positively selecting for
functionally relevant alleles (Haley et al., 2019). We found a high frequency of rare PRNP alleles
in our naïve population which differs from areas currently infected or where CWD is endemic,
including western Canada (Johnson et al., 2006a; Kelly et al., 2008; Wilson et al., 2009; Brandt et
al., 2015; Brandt et al., 2018). Furthermore, infectious diseases can cause population
fragmentation (i.e., structure), demographic changes, and genetic isolation (McKnight et al.,
2017). A pre-infection presence of a large proportion of rare alleles in both functional and
neutral genes, as observed in Ontario, supports surveillance data showing no CWD, which might
bode well for a long-term adaptive response to CWD infection and other stressors. Importantly,

even low selection coefficients should alter allele frequencies in a detectable manner (Figure 5).
However, it is not clear how this standing genetic variation compares to infected populations
prior to the arrival of CWD, creating some uncertainty in the potential adaptive response,
recognizing that loci outside PRNP are also involved (Seabury et al. 2020).

We observed no evidence of strong population structure despite the heterogeneous
landscape observed in Ontario, but there is a clear latitudinal cline in allele frequencies. This
suggests random mating is largely occurring a regional levels in spite of substantial
environmental changes including intense agricultural practices, substantial urbanization of the
landscape, and climate change (Schulte et al., 2007; Walter et al., 2009; Patton, Russell,
Windmuller-Campione, & Frelich, 2018). We might therefore expect that the spread of
advantageous rare alleles under selection may not be limited in the province. Conversely, we
might also expect the homogenous population to facilitate the spread of CWD if anthropogenic
activity were to introduce the disease into the focal population (Escobar et al., 2020). The
movement of infected wildlife might also pose a potential risk to human health at the wildlife-
human-livestock interface as CWD can cross species barriers (Igel-Egalon et al., 2020). Sustained
monitoring across CWD-free regions where deer are managed for sustainability or where food
security is threatened should continue, but consider population-level responses to climate
change (i.e. Kennedy-Slaney, Bowman, Walpole, & Pond, 2018) while integrating genetic
information beyond PRNP allele frequencies.

Conclusion

We gauged the adaptive potential of CWD naïve deer in Ontario, Canada by assessing
functional and neutral genetic diversity. The genome-wide data were consistent with a large
expanding population with a high neutral genomic diversity, no population structure, and an
excess of rare alleles, which is also consistent with non-genetic population and climate models
(Kennedy-Slaney, Bowman, Walpole, & Pond, 2018). We suggest these patterns might favour a
novel adaptive response to the arrival of CWD. Voluntary hunter-harvest based surveillance for
CWD will likely be able to detect the introduction of CWD in a naïve population; however, we
expect that there will be a lag in detection when prevalence is low since low densities likely limit
transmission (Gagnier, Laurion, & DeNicola, 2020). Unfortunately, a lag in detection will likely
permit the establishment of CWD and, over time, eradication becomes nearly impossible and
costs are high (Mysterud & Rolandsen, 2018). Sustained temporal monitoring of variation in
PRNP across CWD-free regions could be a detection tool as our simulations suggest detectable
shifts should occur with the arrival of the disease. By combining demographic patterns and
genotypes with current risk models, managers could improve risk-based detection efforts and
facilitate a more effective resource deployment plan as the disease alters the population.

Acknowledgements

Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grants (JMN -
RGPIN-2019-04330, ABAS RGPIN-2017-03934), NSERC CGS (SH), Canadian Foundation for
Innovation (JELF #66905), Compute Canada RRG (gme-665-AB), and the Ontario Animal Health
Network provided support for the project. We thank Cathy Cullingham for comments, and Anh
Dao for providing samples.

Data Accessibility

DNA sequences: SRA Accessions PRJNA565222 and SUB8436966

PRNP and RADseq analysis pipelines: https://gitlab.com/WiDGeT_TrentU/

376   **Tables**

377   **Table 1**. Allele and amino acids frequencies for reference/alternate alleles at Ontario
378   white-tailed deer (n=631) prion protein gene.

Locus	Allele	Codon	Amino acid	Major frequency	Minor frequency
60	C/T	20	D/-	0.9794	0.0206

136†	A/G	46	S/G	0.9873	0.0127
153	C/T	51	R/-	0.8605	0.1395
168†	A/G	56	G/-	0.9635	0.0365
174†	T/G	58	G/-	0.9842	0.0158
177†	C/G	59	G/-	0.9873	0.0127
178†	T/G	60	W/G	0.9160	0.0840
192†	T/G	64	H/Q	0.8653	0.1347
195†	A/G,T	65	G/-	0.7147	0.2853
198†	T/G	66	G/-	0.7021	0.2979
285	A/C	95	Q/H	0.9699	0.0301
286	G/A	96	G/S	0.6609	0.3391
324	A/G	108	P/-	0.9794	0.0206
365†	G/T	122	G/V	0.9651	0.0349
378	G/A	126	G/-	0.9651	0.0349
417	G/A	139	R/-	0.1696	0.8304
548†	T/A	183	V/D	0.9778	0.0222
555	C/T	185	I/-	0.4120	0.5880
676	C/A	226	Q/K	0.9620	0.0380

379   † indicates novel positions.

381   **Table 2**. Genome-wide population summary statistics including breakdown of sites,
382   number of individuals, nucleotide diversity estimate (π), individual genetic variance (I)
383   relative to the subpopulation genetic variance (F_{IS}), and the observed heterozygosity of
384   white-tailed deer in Ontario.

Group	Ontario	Northern	Southern

Number individuals	182±21	57±3	125±15
Total sites	13,439,671	14,042,841	12,885,176
Polymorphic sites	165,254	139,582	146,036
π	$5.0x10^{-4}$	$6.4x10^{-4}$	$5.7x10^{-4}$
Observed heterozygosity	$4.7x10^{-4}$	$6.1x10^{-4}$	$5.5x10^{-4}$
F_{IS}	$5.7x10^{-4}$	$3.9x10^{-4}$	$4.2x10^{-4}$

Figure 1. Distribution of free-ranging white-tailed deer samples obtained between 2003-2018 by the Ontario Ministry of Natural Resources and Forestry (OMNRF) that were used for the reduced representation genome analysis (n=190; cream) and the prion protein genetic analysis (n=631; orange *and* cream). The natural distribution of free-ranging white-tailed deer is shown for Ontario with a darker shade of gray.

Figure 2. A 771 bp region of the white-tailed deer prion protein gene was analyzed from free-ranging white-tailed deer in Ontario, Canada (n=631). The overlayed genotypes across 19 variable loci were organized by broad management in Ontario and are shown. Circles indicate loci previously described as variable in the white-tailed deer prion protein gene. Triangles indicate novel variable loci in the variable loci in the white-tailed deer prion protein gene.

Figure 3. A plot of PC1 and PC2 from the principal component analysis (PCA) on the
reduced representation white-tailed deer genome. PC1 and PC2 were able to explain a
total of 4.3% of the genomic variation observed. Gray circles represent scores from
samples in northern Ontario. Orange circles represent scores from samples in southern
Ontario.

Model	LL	Parameter	Estimate	BU
		nuB	548.48	35.37
BOTTLEGROWTH	-2034	nuF	139.97	30.83
		T	26.10	85.78

Figure 4. Demographic parameter estimates from δaδi for the optimal 1D model for white-tailed deer in Ontario. Parameter estimates are the ancient population size (N_a), the ratio of population size after instantaneous change to ancient population size (nuB); the ratio of contemporary to ancient population size (nuF); and the time in the past at which instantaneous change happened and growth began (T_C; in units of $2 \ast N_a$ generations). Included are the optimized log-likelihood (LL) and bootstrap uncertainties (BU).

Figure 5. Simulated allele frequency projections for nucleotide positions 285 (a) and 286 (b) in
the PRNP gene under selection. Green and red lines are the selection coefficients provided by
Robinson et al (2012); while the black dots are those from our simulations, with blue line
representing the mean selection coefficient.

References

Aguirre, A.A. (2017). Changing patterns of emerging zoonotic diseases in wildlife, domestic
animals, and humans linked to biodiversity loss and globalization. ILAR Journal, 58(3), 315-318.
doi: 10.1093/ilar/ilx035

Allen, C.D., Breshears, D.D., & McDowell, N.G. (2015). On underestimation of global vulnerability
to tree mortality and forest die-off from hotter drought in the Anthropocene. Ecosphere, 6(8),
129. doi:10.1890/ES15-00203.1

Averill, K.M., Mortensen, D.A., Smithwick, E.A.H., Kalisz, S., McShea, W.J., Bourg, N.A., Parker,
J.D., ... & Nuzzo, V.A. (2018). A regional assessment of white-tailed deer effects on plant
invasion. AoB PLANTS, 10(1), plx047. doi: 10.1093/aobpla/plx047

Baldwin, D.J.B., Desloges, J.R., & Band, L.E. (2000). Physical Geography of Ontario. In Perera,
A.H.; Euler, D.L.; Thompson, I.D. (1st Ed.). Ecology of a managed terrestrial landscape: Patterns
and processes of forest landscapes in Ontario. Ontario Ministry of Natural Resources, Toronto,
Ontario, and UBC Press, Vancouver, British Columbia.

Barber-Meyer, S.M., & Mech, L.D. (2016). White-tailed deer (Odocoileus virginianus) subsidize
gray wolves (Canis lupus) during a moose (Alces americanus) decline: A case of apparent
competition? Canadian Field Naturalist, 130(4), 308-314. doi: 10.22621/cfn.v130i4.1924

Belant, J.L., & Deese, A.R. (2010). Importance of wildlife disease surveillance. Human-Wildlife
Interaction, 4(2), 165-169. Retrieved August 11, 2010, from www.jstor.org/stable/24868837

Bourret, V., Albert, V., April, J., Côté, G., & Morissette, O. (2020). Past, present and future
contributions of evolutionary biology to wildlife forensics, management and conservation.
Evolutionary Applications, 13(6), 1420– 1434. doi: 10.1111/eva.12977

Brandt, A.L., Green, M.L., Ishida, Y., Roca, A.L., Novakofski, J., & Mateus-Pinilla, N.E. (2018).
Influence of the geographic distribution of prion protein gene sequence variation on patterns of
chronic wasting disease spread in white-tailed deer (Odocoileus virginianus). Prion, 12(3-4), 204–
215. doi: 10.1080/19336896.2018.1474671

Brandt, A.L., Kelly, A.C., Green, M.L., Shelton, P., Novakofski, J., & Mateus-Pinilla, N.E. (2015).
Prion protein gene sequence and chronic wasting disease susceptibility in white-tailed deer
(Odocoileus virginianus). Prion, 9(6), 449–462. doi: 10.1080/19336896.2015.1115179

Chafin, T.K., Douglas, M.R., Martin, B.T., Zbinden, Z.D., Middaugh, C.R., Ballard, J.R., Gray,
M.C.,...& Douglas, M.E. (2020). Age structuring and spatial heterogeneity in prion protein gene
(PRNP) polymorphism in white-tailed deer. bioRxiv. doi: 10.1101/2020.07.15.205039v2

Charlesworth, B., & Charlesworth, D. (2010). Elements of evolutionary genetics. Roberts &
Company. doi: 10.1017/S001667231000042X

Chen, L., Qui, Q., Jiang, Y., Wang, K., Lin, Z., Li, Z., ... et al. (2019). Large-scale ruminant genome
sequencing provides insights into their evolution and distinct traits. Science, 364(6446),
eaav6202. doi: 10.1126/science.aav6202

Dawe, K.L., & Boutin, S. (2016). Climate change is the primary driver of white-tailed deer
(Odocoileus virginianus) range expansion at the northern extent of its range; land use is
secondary. Ecology and Evolution, 6(18), 6435–6451. doi: 10.1002/ece3.2316

Escobar, L.E., Moen, R., Craft, M.E., & VanderWaal, K.L. (2019). Mapping parasite transmission
risk from white-tailed deer to a declining moose population. European Journal of Wildlife
Research, 65. doi: 10.1007/s10344-019-1297-z

Escobar, L.E., Pritzkow, S., Winter, S.N., Grear, D.A., Kirchgessner, M.S., Dominquez-Villegas, E.,
Machado, G., Peterson, A.T., & Soto, C. (2020). The ecology of chronic wasting disease in wildlife.
Biological Reviews, 95(2), 393-408. doi: 10.1111/brv.12568

Evira (2018). Moose Found Dead in Forest with Chronic Wasting Disease. Finnish Food Safety
Authority. [Website]. Retrieved from https://mmm.fi/en/-/evira-metsaan-kuolleella-hirvella-
naivetystauti

Ferretti, F., & Mori, E. (2020). Displacement interference between wild ungulate species: does it
occur? Ethology Ecology & Evolution, 32(1), 2-15. doi: 10.1080/03949370.2019.1680447

Fischer, K., Kreyling, J., Beaulieu, M., Beil, I., Bog, M., Bronte, D., Holm, S., ... & Gienapp, P.
(2020). Species-specific effects of thermal stress on the expression of genetic variation across a
diverse group of plant and animal taxa under experimental conditions. Heredity. doi:
10.1038/s41437-020-0338-4

Frerker, K., Sabo, A., & Waller, D. (2017). Correction: Long-term regional shifts in plant
community composition are largely explained by local deer impact experiments. PLoS One,
12(9), e0185037. doi: 10.1371/journal.pone.0185037

Gagnier, M., Laurion, I., & DeNicola, A.J. (2020). Control and surveillance operations to prevent
chronic wasting disease establishment in free-ranging white-tailed deer in Québec, Canada.
Animals, 10(283). doi: 10.3390/ani10020283

Gillespie, J.H. (2004). Population genetics: A concise guide (2nd ed.). Baltimore and London: The
Johns Hopkins University Press.

Güere, M. E., Våge, J., Tharaldsen, H., Benestad, S. L., Vikøren, T., Madslien, K., Hopp, P.,
Rolandsen, C. M., Røed, K. H., & Tranulis, M. A. (2020). Chronic wasting disease associated with
prion protein gene (PRNP) variation in Norwegian wild reindeer (Rangifer tarandus). Prion, 14(1),
1–10. doi: 10.1080/19336896.2019.1702446

Haley, N. J., Merrett, K., Buros Stein, A., Simpson, D., Carlson, A., Mitchell, G., Staskevicius, A.,
Nichols, T., Lehmkuhl, A. D., & Thomsen, B. V. (2019). Estimating relative CWD susceptibility and
disease progression in farmed white-tailed deer with rare PRNP alleles. PLoS One, 14(12),
e0224342. doi: 10.1371/journal.pone.0224342

Harmon, J.P., Moran, N.A., & Ives, A.R. (2009). Species response to environmental change:
Impacts of food web interactions and evolution, Science, 323(5919), 1347-1350. doi:
10.1126/science.1167396

Hashida, Y., Withey, J., Lewis, D.J., Newman, T., & Kline, J.D. (2020). Anticipating changes in
wildlife habitat induced by private forest owners' adaptation to climate change and carbon
policy. PLoS One, 15(4), e0230525. doi: 10.1371/journal.pone.0230525

Heaton, M.P., Leymaster, K.A., Freking, B.A., Hawk, D.A., Smith, T.P., Keele, J.W., Snelling, W.M.,
… & Laegreid, W.W. (2003). Prion gene sequence variation within diverse groups of U.S. sheep,
beef cattle, and deer. Mammalian Genome Society, 14(11), 765–777. doi: 10.1007/s00335-003-
2283-y

Herrera, J., & Nunn, C.L. (2019). Behavioural ecology and infectious disease: implications for
conservation of biodiversity. Philosophical transactions of the Royal Society of London Series B
Biological sciences, 374(1781), 20180054. doi: 10.1098/rstb.2018.0054

Hewitt, D.G. (2011). Biology and management of white-tailed deer. (ed.). Boca Raton, Fla: CRC
Press.

Igel-Egalon, A., Laferrière, F., Tixador, P., Moudjou, M., Herzog, L., Reine, F., Torres, J. M., … &
Béringue, V. (2020). Crossing species barriers relies on structurally distinct prion assemblies and
their complementation. Molecular Neurobiology, 57(6), 2572–2587. doi: 10.1007/s12035-020-
01897-3

Johnson, C., Johnson, J., Clayton, M., McKenzie, D., & Aiken, J. (2003). Prion protein gene
heterogeneity in free-ranging white-tailed deer within the chronic wasting disease affected
region of Wisconsin. Journal of Wildlife diseases, 39(3), 576–581. doi: 10.7589/0090-3558-
39.3.576

Johnson, C., Johnson, J., Vanderloo, J. P., Keane, D., Aiken, J.M., & McKenzie, D. (2006a). Prion
protein polymorphisms in white-tailed deer influence susceptibility to chronic wasting disease.
The Journal of General Virology, 87(7), 2109–2114. doi: 10.1099/vir.0.81615-0

Johnson, C.J., Phillips, K.E., Schramm, P.T., McKenzie, D., Aiken, J.M., & Pedersen, J.A. (2006b).
Prions adhere to soil minerals and remain infectious. PLoS Pathogens, 2(4), e32. doi:
10.1371/journal.ppat.0020032

Jombart, T., Devillard, S., & Balloux, F. (2010). Discriminant analysis of principal components: a
new method for the analysis of genetically structured populations. BMC Genetics, 11, 94. doi:
10.1186/1471-2156-11-94

Kardos, M., & Shafer, A.B.A. (2018). The peril of gene-targeted conservation. Trends in Ecology
and Evolution, 33(11), 827-839. doi: 10.1016/j.tree.2018.08.011.

Kelly, A.C., Mateus-Pinilla, N.E., Diffendorfer, J.,Jewell, E., Ruiz, M.O., Killefer, J., Shelton, P.,
Beissel, T., & Novakofski, J. (2008). Prion sequence polymorphisms and chronic wasting disease
resistance in Illinois white-tailed deer (Odocoileus virginianus). Prion, 2(1), 28–36. doi:
10.4161/pri.2.1.6321

Kennedy-Slaney, L., Bowman, J., Walpole, A.A., & Pond, B.A. (2018). Northward bound: The
distribution of white-tailed deer in Ontario under a changing climate. Wildlife Research, 45(3),
220-228. doi: 10.1071/WR17106

Lambert, S., Ezanno, P., Garel, M., & Gilot-Fromont, E. (2018). Demographic stochasticity drives
epidemiological patterns in wildlife with implications for diseases and population management.
Scientific Reports, 8, 16846. doi: 10.1038/s41598-018-34623-0

Latch, E.K., Heffelfinger, J.R., Fike, J.A., & Rhodes, O.E. (2009). Species-wide phylogeography of
North American mule deer (Odocoileus hemionus): cryptic glacial refugia and postglacial
recolonization. Molecular Ecology, 18(8), 1730-1745. doi: 10.1111/j.1365-294X.2009.04153.x

Latham, A.D.M., Latham, M.C., Knopff, K.H., Hebblewhite, M., & Bouting, S. (2013). Wolves,
white-tailed deer, and beaver: implications of seasonal prey switching for woodland caribou
declines. Ecography, 36(12), 1276-1290. doi: 10.1111/j.1600-0587.2013.00035.x

Mateus-Pinilla, N., Weng, H.Y., Ruiz, M.O., Shelton, P., & Novakofski, J. (2013). Evaluation of a
wild white-tailed deer population management program for controlling chronic wasting disease
in Illinois, 2003–2008. Preventive Veterinary Medicine, 110(3-4), 541-548. doi:
10.1016/j.prevetmed.2013.03.002

McKnight, D.T., Schwarzkopf, L., Alford, R.A., Bower, D.S., & Zenger, K.R. (2017). Effects of
emerging infectious disease on host population genetics: a review. Conservation Genetics, 18,
1235-1245. doi: 10.1007/s10592-017-0974-2

Messer, P.W., Ellner, S.P., & Hairson Jr., N.G. (2016). Can Population Genetics Adapt to Rapid
Evolution? Trends in Genetics, 32(7), 408-418. doi: 10.1016/j.tig.2016.04.005

Miguel, E., Grosbois, V., Caron, A., Pople, D., Roche, B., & Donnelly, C.A. (2020). A systemic
approach to assess the potential and risks of wildlife culling for infectious disease control.
Communications Biology, 3(353). doi: 10.1038/s42003-020-1032-z

Miller, M.W., & Williams, E.S. (2003). Prion disease: horizontal prion transmission in mule deer.
Nature, 425(6953), 35-36. doi: 10.1038/425035a

Miller, M.W., Williams, E.S., Hobbs, N.T., & Wolfe, L.L. (2004). Environmental sources of prion
transmission in mule deer. Emerging Infectious Diseases, 10(6), 1003-1006. doi:
10.3201/eid1006.040010

Miller, M.W., Williams, E.S., McCarty, C.W., Spraker, T.R., Kreeger, T.J., Larsen, C.T., & Thorne,
E.T. (2000). Epizootiology of chronic wasting disease in free-ranging cervids in Colorado and
Wyoming. Journal of Wildlife Diseases, 36(4), 676-690. doi: 10.7589/0090-3558-36.4.676

Miller, W.L., & Walter, D.W. (2019a). Can genetic assignment tests provide insight on the
influence of captive egression on the epizootiology of chronic wasting disease? Evolutionary
Applications, 13(4), 715-726. doi: 10.1111/eva.12895

Miller, W.M., & Walter, D.W. (2019b). Spatial heterogeneity of prion gene polymorphisms in an
area recently infected by chronic wasting disease. Prion, 13(1), 65-76. doi:
10.1080/19336896.2019.1583042

Miller, W.M., Miller-Butterworth, C.M., Diefenbach, D.R., & Walter, W.D. (2020). Assessment of
spatial genetic structure to identify populations at risk for infection of an emerging epizootic
disease. Ecology and Evolution, 10(9), 3977-3990. doi: 10.1002/ece3.6161

Morand, S., & Walther, B.A. (2020). The accelerated infectious disease risk in the Anthropocene:
more outbreaks and wider global spread. bioRxiv, 049866. doi: 10.1101/2020.04.20.049866

Mysterud, A., & Rolandsen, C.M. (2018). A reindeer cull to prevent chronic wasting disease in
Europe. Nature Ecology and Evolution, 2, 1343-1345. doi: 10.1038/s41559-018-0616-1

O'Rourke, K.I., Besser, T.E., Miller, M.W., Cline, T.F., Spraker, T.R., Jenny, A.L., Wild, M.A., ... &
Williams, E.S. (1998). PrP genotypes of captive and free-ranging Rocky Mountain elk (Cervus
elaphus nelsoni) with chronic wasting disease. The Journal of General Virology, 80(10), 2765–
2679. doi: 10.1099/0022-1317-80-10-2765

O'Rourke, K.I., Spraker, T.R., Hamburg, L.K., Besser, T.E., Brayton, K.A., & Knowles, D.P. (2004).
Polymorphisms in the prion precursor functional gene but not the pseudogene are associated
with susceptibility to chronic wasting disease in white-tailed deer. The Journal of General
Virology, 85(5), 1339–1346. doi: 10.1099/vir.0.79785-0

Osterholm, M.T., Anderson, C.J., Zabel, M.D., Scheftel, J.M., Moore, K.A., & Appleby, B.S. (2019).
Chronic wasting disease in cervids: Implications for prion transmission to humans and other
animal species. mBio, 10(4), e01091-19. doi: 10.1128/mBio.01091-19

Parchman, T.L., Gompert, Z., Mudge, J., Schilkey, F.D., Benkman, C.W., & Buerkle, C.A. (2012).
Genome-wide association genetics of an adaptive trait in lodgepole pine. Molecular Ecology,
21(12), 2991-3005. doi: 10.1111/j.1365-294X.2012.05513.x

Patton, S.R., Russell, M.B., Windmuller-Campione, M.A., & Frelich, L.E. (2018). Quantifying
impacts of white-tailed deer (Odocoileus virginianus Zimmerman) browse using forest inventory
and socio-environmental datasets. PLoS One, 13(8), e0201334. doi:
10.1371/journal.pone.0201334

Peart, C.R., Tusso, S., Pophaly, S.D., Botero-Castro, F., Wu, C.C., Aurioles-Gamboa, D., Baird, A.B.,
... & Wolf, J.B.W. (2020). Determinants of genetic variation across eco-evolutionary scales in
pinnipeds. Nature Ecology & Evolution, 4, 1095-1104. doi: 10.1038/s41559-020-1215-5

Pecl, G.T., Araújo, M.B., Bell, J.D., Blanchard, J., Bonebrake, T.C., Chen, I.C., Clark, T.D., Colwell,
R.K., ... & Williams, S.E. (2017). Biodiversity redistribution under climate change: Impacts on
ecosystems and human well-being. Science, 355(6332), eaai9214. doi: 10.1126/science.aai9214

Peterson, B.K., Weber, J.N., Kay, E.H., Fisher, H.S., & Hoekstra, H.E. (2012). Double digest
RADseq: an inexpensive method for de novo SNP discovery and genotyping in model and non-
model species. PLoS One, 7(5), e37135. doi: 10.1371/journal.pone.0037135

Potapov, A., Merrill, E., Pybus, M., & Lewis, M.A. (2016). Chronic wasting disease: transmission
mechanisms and the possibility of harvest management. PLoS ONE, 11(3): e0151039. doi:
10.1371/journal.pone.0151039

Price, S.J., Garner, T.W.J., Cunningham, A.A., Langton, T.E.S., & Nichols, R.A. (2016).
Reconstructing the emergence of a lethal infectious disease of wildlife supports a key role for
spread through translocations by humans. Proceedings of the Royal Society B Biological Sciences,
283(1839), 20160952. doi: 10.1098/rspb.2016.0952

Prusiner, S.B. (1982). Novel proteinaceous infectious particles cause scrapie. Science, 216(4542),
136-144. doi: 10.1126/science.6801762

Rivera, N. A., Brandt, A. L., Novakofski, J. E., & Mateus-Pinilla, N. E. (2019). Chronic Wasting
Disease in Cervids: Prevalence, Impact and Management Strategies. Veterinary medicine
(Auckland, N.Z.), 10, 123–139. doi: 10.2147/VMRR.S197404

Rizzoli, A., Taliapietra, V., Cagnacci, F., Marini, G., Arnoldi, D., Rosso, F., & Rosà, R. (2019).
Parasites and wildlife in a changing world: The vector-host-pathogen interaction as a learning
case. International Journal for Parasitology: Parasites and Wildlife, 9, 394-401. doi:
10.1016/j.ijppaw.2019.05.011

Robinson, S.J., Samuel, M.D., Johnson, C.J., Adams, M., & McKenzie, D.I. (2012). Emerging prion
disease drives host selection in a wildlife population. Ecological Applications, 22(3), 1050-1059.
doi: 10.1890/11-0907.1

Rousselle M., Simion P., Tilak M.K., Figuet E., Nabholz B., & Galtier, N. (2020) Is adaptation
limited by mutation? A timescale-dependent effect of genetic diversity on the adaptive
substitution rate in animals. PLoS Genetics, 16(4), e1008668. doi: 10.1371/journal.pgen.1008668

Russell, M.B., Woodall, C.W., Potter, K.M., Walters, B.F., Domke, G.M., & Oswalt, C.M. (2017).
Interactions between white-tailed deer density and the composition of forest understories in the
northern United States. Forest Ecology and Management, 384, 26-33. doi:
doi.org/10.1016/j.foreco.2016.10.038

Samuel, M.D., Liao, W., Atkinson, C.T., & LaPointe D.A. (2020). Facilitated adaptation for
conservation – Can gene editing save Hawaii's endangered birds from climate driven avian
malaria? Biological Conservation, 241, 108390. doi: j.biocon.2019.108390

Schulte, L.A., Mladenoff, D.J., Crow, T.R., Merrick, L.C., & Cleland, D.T. (2007). Homogenization of
northern U.S. Great Lakes forests due to land use. Landscape Ecology, 22, 1089–1103. doi:
10.1007/s10980-007-9095-5

Seabury, C.M., Oldeschulte, D.L., Bhattarai, E.K., Legare, D., Ferro, P.J., Metz, R.P., & Johnson,
C.D. (2020). Accurate genomic predictions for chronic wasting disease in U.S. white-tailed deer.
G3: Genes, Genomes, Genetics, 10(4), 1433-1441. doi: 10.1534/g3.119.401002

Taylor, P.D., Crewe, T.L., Mackenzie, S.A., Lepage, D., Aubry, Y., Crysler, Z., Finney, G., ... &
Woodworth, B.K. (2017). The motus wildlife tracking system: a collaborative research network to
enhance the understanding of wildlife movement. Avian Conservation and Ecology, 12(1), 8. doi:
10.5751/ACE-00953-120108

Thompson, T.Q., Bellinger, M.R., O'Rourke, S.M., Prince, D.J., Stevenson, A.E., Rodrigues, A.T., ...
& Miller, M.R. (2019). Anthropogenic habitat alteration leads to rapid loss of adaptive variation
and restoration potential in wild salmon populations. Proceedings of the National Academy of
Sciences of the United States of America, 116(1), 177-186. doi: 10.1073/pnas.1811559115

Vázquez-Miranda, H., & Zink, R.M. (2020). Geographic distribution of chronic wasting disease
resistant alleles in Nebraska, with comments on the evolution of resistance. Journal of Fish and
Wildlife Management, 11(1), 46-55. doi: 10.3996/012019-JFWM-002

Walter, W.D., VerCauteren, K.C., Campa, H., Clark, W.R., Fischer, J.W., Hygnstrom, S.E., ... &
Winterstein, S.R. (2009). Regional assessment on influence of landscape configuration and
connectivity on range size of white-tailed deer. Landscape Ecology, 24, 1405–1420. doi:
10.1007/s10980-009-9374-4

Weiskopf, S.R., Ledee, O.., & Thompson, L. M. (2019). Climate change effects on deer and moose
in the Midwest. The Journal of Wildlife Management, 83(4), 769-781. doi:10.1002/jwmg.21649

White, K.S., Gregovich, D.P., Levi, T. (2017). Projecting the future of an alpine ungulate under
climate change scenarios. Global Change Biology, 24(3), 1136-1149. doi: 10.1111/gcb.13919

Wilson, G.A., Nakada, S.M., Bollinger, T.K., Pybus, M.J., Merrill, E. H., & Coltman, D.W. (2009).
Polymorphisms at the PRNP gene influence susceptibility to chronic wasting disease in two
species of deer (Odocoileus Spp.) in western Canada. Journal of Toxicology and Environmental
Health Part A, 72(17-18), 1025–1029. doi: 10.1080/15287390903084264

Supplementary Materials

663 **Table S1**. Four degenerate primers used to amplify the functional white-tailed deer prion protein gene
664 region as two overlapping fragments are shown. Fragment 1 is 460 base-pairs in length and Fragment 2
665 is 590 base-pairs in length.

Primer	Fragment 1	Fragment 2
Forward (5'-3')	TCGTCGGCAGCGTCAGATGTGTATAA GAGACAGACRTGGGCATATGATGCTG AYACC	TCGTCGGCAGCGTCAGATGTGTATAAGAGAC AGTGGAGGCTGGGGTCAAGG
Reverse (5'-3')	GTCTCGTGGGCTCGGAGATGTGTATA AGAGACAGYTGCCAAAATGTATAAGA GG	GTCTCGTGGGCTCGGAGATGTGTATAAGAG ACAGACTACAGGGCTGCAGGTAGAYACT

680 **Table S2**. Thermal cycling conditions for PRNP PCR amplification for Fragment 1 and Fragment

Fragment 1	Fragment 2
98▯ for 2 min	
10 cycles of:	
98▯ for 15 sec	98▯ for 15 sec
60-55▯ for 30 sec[1]	65-60▯ for 30 sec[1]
72▯ for 45 sec	72▯ for 45 sec
98▯ for 15 sec	
20 cycles of:	
98▯ for 15 sec	98▯ for 15 sec
55▯ for 30 sec	60▯ for 30 sec
72▯ for 45 sec[2]	72▯ for 45 sec[2]
72▯ for 7 min	

1 – decrease by 0.5° per cycle; 2 – increasing by 5 sec per cycle

688 **Table S3**. Mixture for double restriction enzyme digestion. Samples and reaction mixture were incubated
689 at 37°C for 3 hours and then 25°C overnight on a thermal cycler with a heated lid. Samples and reaction
690 mixture were head-inactivated at 85°C for 30 minutes.

Reagent	1 X	Final concentration

10x Cutsmart	2.00 ul	1 X
H_2O	11.85 ul	
MseI (10 U/ µl)	0.10 ul	1 U
Sbfl-HF (20 U/ µl)	0.05 ul	1 U
Subtotal	14.00 ul	
DNA	x ul	~ 500 ng
H_2O	6.00 – x ul	
Subtotal	6.00 ul	
Final Volume	20.00 ul	

692 **Table S4**. Mixture for adapter ligation. The digested fragments were combined with 7 ul of the adapter
693 ligation mixture and 3 ul of a unique barcoded Sbfl adapter (1.0 uM). The ligation on ~ 30 ul reaction
694 mixture was performed in 16°C for 3 hours

Reagent	1 X	Final concentration
10x CutSmart	1.00	1 X
100 mM ATP	0.30	1 mM
H_2O	2.45	
MseI Y adapter (10 uM)	3.00	1 µM
T4 DNA ligase (400 U/µl)	0.25	100 U
Subtotal	7.00 ul	

697 **Table S5**. Illumina PCR mixture. The purified restriction-ligation DNA (3 ul) was combined with 7 ul of
698 PCR mixture and a PCR was performed on 10 ul. Four replicated per sample was performed. The thermal
699 cycler profile for this PCR was 98°C for 30 seconds; 20 cycles of 98°C for 20 seconds, 60°C for 30 seconds,
700 72°C for 45 seconds; and a final extension at 72°C for 5 minutes.

Reagent	1 X	Final concentration

H$_2$O	1.33	
KAPA HiFi ReadyMix	5.00	2 X
PCR primer mix (5 µM each)	0.67	0.5 µM
Subtotal	7.00 ul	

Table S6. The major/minor allele counts for four nucleotide (nt) positions of variation in the white-tailed deer prion protein gene that are linked to reduced susceptibility or reduced clinical progression of chronic wasting disease for Northern and Southern Ontario are shown. A two-sided Fisher's Exact test was conducted on the major and minor allele counts at the four chronic wasting disease-linked nucleotide positions. A p-value less than 0.05 was considered significant and indicated that there were significant differences between the groups.

Location	nt60	nt285	nt286	nt676
Northern	186/3	182/7	132/57	187/2
Southern	432/10	430/12	285/157	420/22
Southeastern	306/8	308/6	196/118	300/14
Southwestern	126/2	122/6	89/39	120/8
p-value	0.764	0.611	0.200	0.021

Table S7. Haplotypes were estimated with PHASE v2.1.1 set to the same parameters in Brandt et al., 2018: Markov chain Monte Carlo (MCMC) samples were taken from a minimum of 100,000 steps, with a discarded burn-in of 10,000; samples were drawn every 100 MCMC steps. Five repetitions were performed, and haplotype frequencies compared to verify consistent assignment. Included are estimates of population haplotypes with frequencies of greater than 1% (count=1262, number haplotypes = 151) and associated estimated standard deviations (S.E.; square root of the variance of the posterior distribution) at 19 variable positions, with 0 representing non-variants and 1 representing variants.

ID	f	S.E.	Codon	153	195	198	286	365	378	417	555

3	0.228	0.006	-	0	0	0	0	0	0	1	0
1	0.122	0.005	-/-	0	0	0	0	0	0	1	1
9	0.104	0.001	Ref	0	0	0	0	0	0	0	0
16	0.087	0.005	96S/-/-	0	0	0	1	0	0	1	1
7	0.050	0.001	-	0	0	0	0	0	0	0	1
252	0.041	0.003	-/-	1	0	0	0	0	0	1	0
18	0.033	0.004	96S/-	0	0	0	1	0	0	1	0
54	0.022	0.003	-/-/-	0	1	1	0	0	0	1	0
259	0.012	0.002	-/122V/-/-	1	0	0	0	1	1	1	0
27	0.012	0.002	-/-/-	0	0	1	0	0	0	1	1
28	0.011	0.003	-/96S/-/-	0	0	1	0	0	0	1	0
48	0.010	0.002	-/96S/-/-	0	1	0	1	0	0	1	1

Table S8. Optimized log-likelihood (LL) and bootstrap uncertainties (BU) obtained from 1D demographic models on Ontario white-tailed deer as a single population. Model specifics and parameters are outlined. The most optimal model for 1D is shown in bold. Modified 1D demographic models are indicated with an asterisk. All time estimates are reported in units of $2*Na$ generations. Migration rate (m) is reported in units of $2*Na*m$.

Model Name	LL	Parameter	Estimate	BU
SNM	-25746	n/a	n/a	41.3 $1.11x10^{-15}$

Model		Parameter		
				$7.77x10^{-16}$
TWO_EPOCH	-3015	nu	573.38	3.97
				40.69
		T	107.90	7.62
GROWTH	-8795	nu	21.57	217.02
				8.93
		T	6.58	3.79
BOTTLEGROWTH	-2034	nuB	548.48	35.37
		nuF	139.97	30.83
		T	26.10	85.78
BOTTLEPOP*	-8495	nu	54.48	287.41
		T	20.17	78.28
				46.48
TWOPOPCHANGES	-3129	nuB	$4.19x10^{-5}$	$5.16x10^{4}$
		nuF	0.32	
		TB	$5.14x10^{-3}$	$2.06x10^{-5}$
		TF	$6.04x10^{-2}$	0.11
GROWTHPLUSBOTTLE	-25631	nuB	8.47	$1.43x10^{5}$
		nuF	0.14	
		TB	0.92	2.74
		TF	4.28	$2.39x10^{-2}$
BOTTPLEPLUSGROWTH	-3544	nuB	397.36	8.75
		nuF	473.75	93.69

		T	97.40	169.23
THREE_EPOCH	-3274	nuB	$1.35x10^{-4}$	$1.99x10^{5}$
		nuF	$1.31x10^{-2}$	$1.46x10^{-5}$
		TB	$3.48x10^{-3}$	$2.64x10^{-3}$
		TF	$2.47x10^{-3}$	

Table S9. Nucleotide variations in free-ranging white-tailed deer prion protein gene that are associated
with chronic wasting disease are either protective (1), increase susceptibility (2), or are neutral. The
major and minor allele frequencies for each site across a 771 bp region of the prion protein gene in
white-tailed deer are reported for different regions, in descending order. The year CWD was found in
free-ranging cervids is reported for each location. The data are from free-ranging white-tailed deer
samples collected in: Alberta, Canada (AB); Colorado, USA (COL); Illinois, USA (IL); Ontario, Canada (ON);
Nebraska, USA (NE); Saskatchewan, Canada (SK); Wisconsin, USA (WI); and Wyoming, USA (WY).

Site	Role	Major	Minor	Region	CWD[+]	Citation
C60T	1	0.98	0.02	ON	n/a	Haworth et al., 2020
		0.98	0.02	WY	1985	Heaton et al., 2003

		0.94	0.06	AB, SK	2002,1996	Wilson et al., 2009
		0.92	0.08	IL	2002	Kelly et al., 2008
C153T	2	0.96	0.04	WY	1985	Heaton et al., 2003
		0.94	0.06	AB, SK	2002,1996	Wilson et al., 2009
		0.89	0.11	IL	2002	Kelly et al., 2008
		0.86	0.14	ON	n/a	Haworth et al., 2020
A285C	1	1.00	0.00	WI	2002	Johnson et al., 2006
		0.99	0.01	AB, SK	2002,1996	Wilson et al., 2009
		0.98	0.02	NE	1999	Vázquez-Miranda & Zink, 2020
		0.97	0.03	ON	n/a	Haworth et al., 2020
		0.94	0.06	IL	2002	Kelly et al., 2008
G286A	1	0.88	0.12	COL	1967	O'Rourke et al., 1998 (unpublished)
		0.86	0.14	IL	2002	Kelly et al., 2008
		0.83	0.17	NE	1999	Vázquez-Miranda & Zink, 2020
		0.81	0.19	WI	2002	Johnson et al., 2006
		0.66	0.34	ON	n/a	Haworth et al., 2020
A324G	1	0.98	0.02	ON	n/a	Haworth et al., 2020
		0.98	0.02	NE	1999	Vázquez-Miranda & Zink, 2020
		0.96	0.04	WY	1985	Heaton et al., 2003
		0.96	0.04	AB, SK	2002,1996	Wilson et al., 2009
		0.94	0.06	WY	1985	Heaton et al., 2003
G417A	3	0.99	0.01	AB, SK	2002,1996	Wilson et al., 2009
		0.79	0.21	COL	1967	O'Rourke et al., 1998
		0.17	0.83	ON	n/a	Haworth et al., 2020

C555T	1	0.89	0.11	n/a	n/a	Raymond et al., 2000
		0.65	0.35	AB, SK	2002,1996	Wilson et al., 2009
		0.58	0.42	IL	2002	Kelly et al., 2008
		0.41	0.59	ON	n/a	Haworth et al., 2020
C676A	1	0.99	0.01	IL	2002	Kelly et al., 2008
		0.98	0.02	AB, SK	2002,1996	Wilson et al., 2009
		0.97	0.03	WI	2002	Johnson et al., 2006
		0.96	0.04	ON	n/a	Haworth et al., 2020

Figure S1. The Canadian province of Ontario as managed by the Ontario Ministry of Natural Resources and Forestry. There are three broad regions Ontario is managed by: (A) Northern Ontario, (B) Southeastern Ontario, and (C) Southwestern Ontario. Collectively (B) and (C) form Southern Ontario. Outlined in red are the wildlife management units designated within each broad region.

Figure S2. The Bayesian information criterion (BIC) from a population cluster identification using successive K-means cluster assignment on the reduced representation white-tailed deer genome from identified one cluster as optimal.