Characterizing the demographic history and prion protein variation

to infer susceptibility to chronic wasting disease in a naïve

population of white-tailed deer (*Odocoileus virginianus*)

Population and disease genomics of white-tailed deer.

Sarah E Haworth[1], Larissa Nituch[2], Joseph M Northrup[1,2], Aaron BA Shafer[1,3*‡]

1 – Environmental and Life Sciences Graduate Program, Trent University, 7K9J 7B8 Peterborough, Canada

2 – Wildlife Research and Monitoring Section, Ontario Ministry of Natural Resources and Forestry, Trent

University, 7K9J 7B8 Peterborough, Ontario, Canada

3 – Forensics Program, Trent University, K9J 7B8 Peterborough, Ontario, Canada

* – *corresponding author* aaronshafer@trentu.ca

‡ – ORCID: https://orcid.org/0000-0001-7652-225X

Abstract

Assessments of the adaptive potential in natural populations are essential for

understanding and predicting responses to environmental stressors like climate

change and infectious disease. Species face a range of stressors in human-dominated

landscapes, often with contrasting effects. White-tailed deer (deer) are expanding in

the northern part of their range following decreasing winter severity and increasing

forage availability. Chronic wasting disease (CWD), a prion disease affecting cervids,

is likewise expanding and represents a major threat to deer and other cervids. We

obtained tissue samples from free-ranging deer across their native range in Ontario,

Canada which has yet to detect CWD in wild populations of cervids. We used high-

throughput sequencing to assess neutral genomic variation, and variation in the PRNP

gene that is partly responsible for the protein misfolding when deer contract CWD.

Neutral variation revealed a high number of rare alleles and no population structure,

and demographic models suggested a rapid historical population expansion. Allele

frequencies of PRNP variants associated with CWD susceptibility and disease

progression were evenly distributed across the landscape and consistent with deer

populations not infected with CWD. We then estimated the selection coefficient of

CWD, with simulations showing an observable and rapid shift in PRNP allele

frequencies that coincides with the start of a novel CWD epidemic. Sustained

surveillance of genomic and PRNP variation can be a useful tool for CWD-free regions

where deer are managed for ecological and economic benefits.

Keywords

RADseq, ungulate, PRNP, prion, Canadian wildlife, population genetics

Introduction

Human-induced environmental change has caused widespread alterations to ecological and evolutionary processes (Harmon, Moran, & Ives, 2009; Pecl et al., 2017). Climate change is expected to be the dominant driver of wildlife population declines and has been linked to broad-scale biodiversity losses, but regional responses are often nuanced and context dependent (e.g. Taylor et al., 2017; White, Gregovich, & Levi, 2017; Hashida et al., 2020). For example, climate change in the Midwestern United States differentially favors the survival of two sympatric populations of ungulates with similar selection pressures and life history traits, but differing densities (Escobar, Moen, Craft, & VanderWaal, 2019; Weiskopf, Ledee, & Thompson, 2019). In such instances, intraspecific genetic diversity and adaptive potential is crucial for long-term population viability (Kardos & Shafer, 2018).

The emergence, spread, and persistence of infectious diseases in previously allopatric populations is facilitated by climate change and other anthropogenic activity (Price et al., 2016; Aguirre, 2017; Morand & Walther, 2020). The impacts of infectious disease on wildlife populations are of interest to managers, especially if the affected species holds economic or cultural value (Lambert et al., 2018; Weiskopf, Ledee, & Thompson, 2019). Preventing and controlling diseases in free-ranging populations can, however, be complex and costly when they are both naïve to the infectious disease and faced with climate change and other anthropogenic activities (Herrera & Nunn, 2019; Miguel et al., 2020; Samuel et al., 2020). Selective pressures are increased under these circumstances and populations are forced to respond to

multiple stressors simultaneously, or potentially face local extirpation (Fischer et al.,

2020).

White-tailed deer (*Odocoileus virginianus*; deer) are the most widely

distributed and abundant ungulates in North America and hold significant economic

and cultural value (Hewitt, 2011). The northern range of deer is primarily limited by

snow but decreasing winter severity has allowed deer to expand northward beyond

their historical range limits (Dawe & Boutin, 2016; Kennedy-Slaney, Bowman,

Walpole, & Pond, 2018). This expansion has implications for ecosystems as, for

example, deer herbivory alters long-term regional habitat characteristics and plant

communities (Frerker, Sabo, & Waller, 2017; Otsu, Iijima, & Nagaike, 2019; Kroeger

et al., 2020). Further, deer are an important prey species and predator populations

increasing in response to deer expansion has led to greater predation and apparent

interspecific competition with other ungulates (Latham et al., 2013; Barber-Meyer &

Mech, 2016). Consequently, northward expansions of deer are having profound

impacts to ecosystems including facilitating infectious pathogen and disease spread

(Averill et al., 2018; Ferretti & Mori, 2020).

A widespread threat to deer in North America is the highly infectious and fatal

neurodegenerative prion disease called chronic wasting disease (CWD). CWD is the

only prion disease known to infect captive and free ranging species of cervids (family

Cervidae) and has been reported in North and South America, Europe, and South

Korea (Haley et al., 2019). With virtually no barriers to transmission and a lengthy

infectious preclinical period, the local prevalence of CWD in North America has been

measured to be as high as 50% and 82% in wild and captive populations, respectively

(Miller et al., 2004; O'Rourke et al., 2004). Due to constraints on CWD surveillance it

is likely that the distribution and prevalence of CWD in wild populations are

underestimated (Escobar et al., 2020). It is clear the frequency and occurrence of CWD

has increased over time, in part driven by anthropogenic activities related to hunting

and wildlife farming (Osterholm et al., 2019).

Despite CWD being fatal there is inter-individual variation in susceptibility and

clinical progression. Susceptibility and clinical progression are associated with non-

synonymous and synonymous genetic variation in the functional prion protein gene

(PRNP; Güere et al., 2020; Chafin et al., 2020). Single nucleotide polymorphisms

(SNPs) at nucleotide (nt) 60, nt153, nt285, nt286, nt555, and nt676 in deer PRNP have

been associated with altered CWD susceptibility or pathogenic processes (Johnson et

al., 2006a; Wilson et al., 2009; Brandt et al., 2015; Brandt et al., 2018). The presence

of CWD appears to affect population PRNP allele frequencies over space and time due

to selection (Robinson et al., 2012); however, altered CWD susceptibility and

pathogenic processes are clearly polygenic traits (Seabury et al. 2020) and disease

spread is different in structured populations, which might require different wildlife

management practices (Chafin et al., 2020).The efficacy of selection in the face of a

novel pressure like CWD is dependent on the effective population size (N_e). Common

metrics to infer selection such as Tajima's D, specifically measure shifts in allele

frequencies across the site-frequency spectrum; however, the frequency and

proportion of rare alleles is sensitive to demographic processes (Messer, Ellner, &

Hairston Jr., 2016; Platt et al., 2019), and population genetics theory predicts an

excess of rare alleles in expanding populations (Gillespie, 2004).

Ontario, Canada reflects the northern leading edge of deer range in eastern

North America (Kennedy-Slaney, Bowman, Walpole, & Pond, 2018). The landscape of

Ontario is heterogenous and environmental clines exist around the Great Lakes

region. Ontario has not detected CWD in wild cervids, but CWD has been detected in

farmed and captive cervids from virtually every jurisdiction bordering Ontario, with

the province using a weighted surveillance approach to model CWD risk and

strategically use resources for surveillance. Accordingly, at the genome-level, we

predicted an excess frequency of rare neutral and PRNP variants across our study

region given Ontario's deer population is expanding (Kennedy-Slaney, Bowman,

Walpole, & Pond, 2018). Based on large-scale distribution changes and recent

population trends in deer (Baldwin, Desloges, & Band, 2000; Latch et al., 2009), we

predicted we would observe high neutral genomic diversity (N_e) and demographic

population expansion, indicating increased gene flow and decreased population

structure despite a heterogenous landscape. Since PRNP is not under selection by

CWD given the region is disease free, we predicted functional variation to resemble

regions most recently exposed to CWD (or still disease free). This is the first study to

characterize PRNP genetic variation and population genomic structure of wild deer,

while also determining the ancient and contemporary demographic and selection

processes driving patterns of diversity.

Materials and Methods

Study area and sample collection

We sampled white-tailed deer across Ontario, Canada (Figure 1). Between

2002-2018, retropharyngeal lymph nodes were opportunistically extracted from

hunter harvested deer across Ontario through the CWD surveillance program

managed by the Ontario Ministry of Natural Resources and Forestry (OMNRF).

Auxiliary data including year, sex, age-class, Mercator grid cell unit (GCU; 10x10 km),

and wildlife management unit (WMU) were also collected with deer samples. North

western and southern Ontario regions are geographically discontinuous for deer

(Figure 1), samples were therefore assigned to northern Ontario or southern Ontario

(Figure S1) for the purpose of analysis where sampling regions are compared.

Preliminary analysis of data did not warrant separating southeastern and

southwestern Ontario as per provincial management zones. Genomic DNA was

extracted from deer samples using a silica-based DNA extraction kit for tissue

following manufacturers protocol (Qiagen, Cat. No. 69506) and stored at -20°C. DNA

quality was assessed by a spectrometer (NanoDrop 2000, Thermo Scientific) and by

2% agarose GelRed gel electrophoresis.

Library preparation for PRNP genetic analysis

A 771 base-pair (bp) region of the deer prion protein precursor (PRNP) gene

was targeted and amplified using four degenerate primers (Table S1). Four replicate

PCRs generated a 460 base pair Fragment 1 and a 580 base pair Fragment 2 (Table

S2). In a 2:1 ratio of Fragment 1 to Fragment 2, respectively, amplified DNA was added

for each individual and then indexed with standard Illumina multiplexing indices.

Negative controls of UltraPure distilled water (Invitrogen, 1897011) were used for

each 96-well plate. The library was purified of artifacts following manufacturers

protocol for AMPure XP beads (Beckman Coulter, A63880) and validated with a

TapeStation D1000 kit (Agilent, 5067-5582). The library was sequenced on an

Illumina MiSeq platform at the University of Guelph Advanced Analysis Centre to

generate 300 base pair (bp) pair-end reads for each sample.

Library preparation for RADseq genomic analysis

Restriction-site associated DNA sequencing (RADseq) libraries were

generated using an adapted protocol from Parchman et al. (2012) and Peterson et al.

(2012) with Sbfl-HF and Msel restriction enzymes. Samples were incubated, digested

overnight, and heat-inactivated in 96-well plates (Table S3). Negative controls of

UltraPure distilled water (Invitrogen, 1897011) were used for each 96-well plate.

Restriction digested DNA was combined with 7 ul of ligation mixture and 3 ul of one

of the 24 available Sbfl adapters (1.0 uM). Adapters were ligated at 16°C for 3 hours.

DNA fragments were purified of artifacts following manufacturers protocol for

AMPure XP beads (Beckman Coulter, A63880). Adapter-ligated fragments were

amplified in four separate 10 ul reactions that incorporated barcodes. Reaction

conditions and primers are shown for ligation mixture and PCR in Table S4 and Table

S5, respectively. Samples were pooled and purification was performed following

manufacturers protocol for QIAquick PCR Purification kit (Qiagen, 28106) for a final

elution to 42 ul. Size selection between 450 bp to 700 bp was performed on 80 ul

replicates of purified library and gel purification was performed following

manufacturers protocol for QIAquick Gel Extraction kit (Qiagen, 28706) for a final

elution to 60 ul. The purified final library was validated with a TapeStation D1000 kit

(Agilent, 5067-5582). The libraries were sequenced at The Centre for Applied

Genomics (TCAG) in The Hospital for Sick Children (SickKids) on an Illumina HiSeq

2500 to produce 2x126 base pair paired end reads.

Bioinformatic pipeline and data analysis I – PRNP gene

The quality of reads was assessed using FastQC (Babraham Institute; v0.11.8).

Samples were excluded if at least one file in the pair-end files for a sample was less

than 1kB in size or failed to pass quality standards. A novel command line-based

pipeline was developed to assemble and genotype PRNP (accessible at

https://gitlab.com/WiDGeT_TrentU). Our workflow integrated Pullseq v1.0.2, BWA

v0.7.17, SAMtools; v1.9, BCFtools v1.9, and VCFtools v0.1.16-15. Briefly, for each

sample, the pipeline extracted relevant reads based on the presence of primer

sequence, mapped the extracted reads to a 771 bp PRNP gene reference sequence. We

generated a consensus sequence and called single nucleotide polymorphisms (SNPs).

SNP calls were limited to positions where there was a minimum read depth of 30 and

mapping quality score of at least 30. Sanger sequencing of a subset of samples and

their a priori called variants were used to validate the bioinformatic pipeline.

The presence of asparagine (N) at aa138 (nt413A) indicates amplification of

the pseudogene (Brandt et al., 2015); we therefore filtered out all sequences with this

site. SNPs with a total frequency of occurrence of 1% or less were excluded from the

analysis. A two-sided Fisher's Exact Test was conducted on minor allele counts from

either northern or southern sampling regions at four well studied positions in the

deer PRNP gene associated with CWD: nucleotide (nt) 60, nt285, nt286, and nt676.

Synonymous and non-synonymous sites were identified using MEGA X v10.0.5.

Haplotypes were estimated from unphased sequences with PHASE v2.1.1 using a

Markov chain Monte Carlo (MCMC) sampling approach with a minimum of 100,000

steps, with a discarded burn-in of 10,000, and samples were drawn every 100 MCMC

steps. Five repetitions were performed to verify consistent frequencies of haplotype

assignment (Brandt et al., 2018). Haplotypes with a frequency of less than 1% were

removed. The genotype, frequencies, and estimated standard deviations of the

remaining haplotypes were analyzed as a 2x2 contingency table by sampling region.

Bioinformatic pipeline and data analysis II – RADseq

Fastq files were demultiplexed using process_radtags within the Stacks v2.3

module. Parameters within process_radtags included the removal of any read with an

uncalled base and the discarding of reads with low quality scores. The demultiplexed

sample files were aligned against the deer genome (Genome Accession

JAAVWD000000000) using BWA with samtools used to sort, merge and compress

BAM files. The referenced-based approaches on gstacks and populations program

within STACKs produced a variant call format (VCF) file with the restrictions that the

minimum percentage of individuals in a population required to process a locus for

that population was 90%. The VCF was filtered using VCFtools to only include reads

with a minimum read depth of 20. Population statistics, including F_{IS}, observed and

estimated homozygosity, and nucleotide diversity were calculated using the

populations module. To analyze sources of variation, we generated a principal

component analysis (PCA) using the R v3.6.1 package adegenet v2.1.3. A linear

regression was run on principal component (PC) 1 and PC2 scores against latitude

and longitude. We estimated F_{ST} between north and south using StAMPP v1.6.1.

Population structure was detected using successive K-means clustering and a

discriminant analysis of principal components (DAPC) available in adegenet (Jombart,

Devillard, & Balloux, 2010).

Demographic Analysis and Estimate of Effective Population Size

The final VCF was converted into 1D site frequency spectrum (SFS) for all of

Ontario and northern vs and southern designations, respectively, using vcf2dadi.py

with projections for the SFS estimated in easySFS. We applied a diffusion-based

approach to demographic inference through the Diffusion Approximation for

Demographic Inference (δaδi) tool by Gutenkunst et al., (2009). Nine 1D models were

assessed for Ontario as a single population. The optimum model was selected as the

lowest optimized log-likelihood of all successfully run models. δaδi was also used to

estimate the following summary statistics for the province: Watterson Theta (θ),

Tajima's *D* and the number of segregating sites. Using the mutation rate (μ) per site

per generation of a closely related species (*Rangifer tarandus* from Chen et al., 2019),

total number of sites (L), and the parameters estimated in the optimum model

selected from δaδi, we estimated the ancestral effective population size as N_a = θ /

4μL.

Estimation of selection on PRNP and allele frequency projections for a naïve population

We estimated the selection coefficient (*s*) at two sites (nt285 and nt286) using

the approach of Thompson et al. (2019) which accounts for number of generations

and different expression modes (i.e. recessive, dominant, codominant). Strength of

selection against the less resistant phenotype [i.e., the homozygous common allele

(s_{AA})] can be estimated by calculating values of s_{AA} that explained the estimated

change in allele frequencies between positive and negative animals from the same

region. Here we used starting (negative CWD) allele frequencies from Wilson et. al.

(2009) and Kelly et al. (2009) and estimated *s* over *n* generations the equation:

$p'=(1-s_{AA})p^2 + (1-s_{AB})p(1 - p)/(1-s_{AA})p^2 + (1-s_{AB})2p(1 - p) + (1-s_{BB})(1 - p)^2$

taken from Charlesworth and Charlesworth (2010). To account for uncertainty in

time and allele frequency estimates we ran 100 iterations with *n* ranging from 2-25

generations (~4-50 deer years) and positive and negative allele frequencies (±1%)

for those reported from both Wilson et. al. (2009) and Kelly et al. (2009). Then using

our estimated allele frequencies for Ontario, we projected each allele frequencies into

the next 25 generations using our estimated selection coefficients, and the estimated

s coefficients of 0.0103 and 0.074 from Robinson et al. (2012). Calculations were

conducted under three relative fitness scenarios as per Thompson et al. (2019).

Results

PRNP Genetic analysis

A total of 631 Ontario deer samples were included in the PRNP genetic analysis

(Figure 1). Nineteen SNPs were detected after filtering (Table 1), with 8 being non-

synonymous substitutions. Six of the detected variants in the PRNP gene have been

linked to CWD susceptibility or clinical progression, these include nt60, nt153, nt285,

nt286, nt555, and nt676. A two-sided Fisher's Exact test on the major and minor allele

counts at the four important, arguably the most studied, CWD-linked loci (nt60, nt285,

nt286, and nt676) conducted between Northern and Southern Ontario indicated that

there was only a difference in frequency ($p < 0.05$) at nt676 (Table S6).

There were 102 unique haplotypes with a count of at least one, with 12

haplotypes having a frequency greater than 1% (Figure 2; Table S7). The two most

common haplotypes, Haplotype 3 (f=0.23) and Haplotype 1 (f=0.12) did not include

any non-synonymous substitutions. Haplotype A (f=0.30) and Haplotype B (f=0.25)

reported by Brandt et al (2015, 2018) from northern Illinois were also detected as

Haplotype 16 (f=0.09) and Haplotype 7 (f=0.05), respectively. The same Haplotype A

(f=0.15) and Haplotype B (f=0.23) were reported by Chafin et al., 2020 in deer from

Arkansas, USA.

RADseq Genomic analysis

A total of 235 Ontario deer samples were sequenced in ddRADseq libraries.

Following quality control and quality assessment, including FastQC and line counts of

demultiplexed files, 190 samples remained for downstream analysis (Figure 1).

Estimated population diversity statistics are summarized in Table 2. The PCA clearly

separated northern and southern Ontario along PC1 (Figure 3). A linear regression

revealed that PC1 was strongly associated with longitude (β = −0.79; adjusted R^2 =

0.77; p-value < 0.01). However, population structure between northern and southern

Ontario was weak (F_{ST}=0.02). A BIC based on the K-cluster analysis also indicated that

the most optimum number of clusters was 1.

The optimum 1D demographic model for Ontario was the BOTTLEGROWTH

model, which models an instantaneous size change followed by exponential change

(Table S8; Figure 4). From the 1D site frequency spectrum, the mean Tajima's D was

estimated to be -2.126 consistent with a population expansion after a bottleneck.

Using the calculated θ, we estimated N_a to be ∼20,000: this would place the timing of

the population change (Tc) measured in $2N_a$ generations around the onset of the last

glacial maximum (Figure 4).

PRNP selection and projection

We estimated the selection coefficients to be 0.08 (±0.06) and 0.11 (±0.07) for

nt285 and nt286 under a dominance model. All simulated projections with our *s*

values and the upper reported value of Robinson et al (2012) showed a rapid shift in

allele frequencies (Figure 5); the majority of simulated trajectories did not overlap

with the low *s* coefficient of 0.01 previously reported by Robinson et al (2012).

Discussion

White-tailed deer in North America are intensively managed for hunter

harvest and are expanding their range northward due to climate change. The

frequency and occurrence of CWD infection in captive and free-ranging deer is

likewise increasing (Osterholm et al., 2019; Rivera, Brandt, Novakofski, & Mateus-

Pinilla, 2019), but transmission and spread in free-ranging populations are still poorly

understood (Potapov, Merrill, Pybus, & Lewis, 2016). We observed that the frequency

of PRNP alleles in our naïve population differed from areas currently infected with

CWD or where CWD is endemic, including western Canada (Table S8; Kelly et al.,

2008; Wilson et al., 2009; Brandt et al., 2015; Brandt et al., 2018; Chafin et al., 2020).

The emergence, transmission, and persistence of highly infectious diseases in

healthy populations are often facilitated by climate change and exacerbated in areas

with intense anthropogenic activity (McKnight et al., 2017; Rizzoli et al., 2019). The

introduction and re-introduction of infectious diseases often results in rapid local

population declines, reducing species' adaptive potential and generating substantial

economic losses (Belant & Deese, 2010; Escobar, Moen, Craft, & VanderWaal, 2019).

In Ontario deer, however, we observed an excess of rare alleles with no evidence of

strong population structure, including in southern Ontario where an environmental

cline exists around lake systems. An excess of rare alleles and high N_e and absent

population structure might provide the means for effective adaptation to selective

pressures, including climate change, infectious disease, and human activity.

Linking Population Demographics to Functional genes

We assessed contemporary and historical gene flow by examining neutral

genomic variation across tens of thousands of loci from deer across the province on

Ontario. The data were consistent with a large expanding population with a high level

of genomic diversity and an excess of rare alleles. These genomic patterns are also

consistent with population simulations of responses to climate change (Kennedy-

Slaney, Bowman, Walpole, & Pond, 2018), and offers some clues as to the potential

adaptive response were CWD to arrive. Specifically, two important features of the

genomic data are noteworthy: the high number of a rare alleles and the limited

population structure across Ontario.

Demographic processes and life history strategies influence the proportion of

rare alleles, which are important to the adaptive process but are sensitive to recent

evolutionary processes (Peart et al. 2020). Evidence suggests the accumulation of rare

alleles is independent of taxa, but adaptation appears limited in low-diversity taxa

(e.g., primates; Rousselle et al., 2020). The deer genomic data evaluated here are

consistent with high adaptive potential; specifically, we calculated an N_a to be

~20,000, with the expansion estimated to be 100X suggestive of a large species-wide

N_e (derived either from π in Table 2 or the demographic model in Figure 4).

Studies of populations infected with CWD often demonstrate a lack of allelic

diversity at the PRNP gene, which is thought to be due to CWD-driven selection

positively selecting for functionally relevant alleles (Haley et al., 2019). We found a

high frequency of rare PRNP alleles in our naïve population which differs from areas

currently infected or where CWD is endemic, including western Canada (Johnson et

al., 2006a; Kelly et al., 2008; Wilson et al., 2009; Brandt et al., 2015; Brandt et al.,

2018). Furthermore, infectious diseases can cause population fragmentation (i.e.,

structure), demographic changes, and genetic isolation (McKnight et al., 2017). A pre-

infection presence of a large proportion of rare alleles in both functional and neutral

genes, as observed in Ontario, supports surveillance data showing no CWD, which

might bode well for a long-term adaptive response to CWD infection and other

stressors. Importantly, even low selection coefficients should alter allele frequencies

in a detectable manner (Figure 5). However, it is not clear how this standing genetic

variation compares to infected populations prior to the arrival of CWD, creating some

uncertainty in the potential adaptive response, recognizing that loci outside PRNP are

also involved (Seabury et al. 2020).

We observed no evidence of strong population structure despite the

heterogeneous landscape observed in Ontario, but there is a clear latitudinal cline in

allele frequencies. This suggests random mating is largely occurring a regional levels

in spite of substantial environmental changes including intense agricultural practices,

substantial urbanization of the landscape, and climate change (Schulte et al., 2007;

Walter et al., 2009; Patton, Russell, Windmuller-Campione, & Frelich, 2018). We might

therefore expect that the spread of advantageous rare alleles under selection may not

be limited in the province. Conversely, we might also expect the homogenous

population to facilitate the spread of CWD if anthropogenic activity were to introduce

the disease into the focal population (Escobar et al., 2020). The movement of infected

wildlife might also pose a potential risk to human health at the wildlife-human-

livestock interface as CWD can cross species barriers (Igel-Egalon et al., 2020).

Sustained monitoring across CWD-free regions where deer are managed for

sustainability or where food security is threatened should continue, but consider

population-level responses to climate change (i.e. Kennedy-Slaney, Bowman, Walpole,

& Pond, 2018) while integrating genetic information beyond PRNP allele frequencies.

Conclusion

We gauged the adaptive potential of CWD naïve deer in Ontario, Canada by

assessing functional and neutral genetic diversity. The genome-wide data were

consistent with a large expanding population with a high neutral genomic diversity,

no population structure, and an excess of rare alleles, which is also consistent with

non-genetic population and climate models (Kennedy-Slaney, Bowman, Walpole, &

Pond, 2018). We suggest these patterns might favour a novel adaptive response to the

arrival of CWD. Voluntary hunter-harvest based surveillance for CWD will likely be

able to detect the introduction of CWD in a naïve population; however, we expect that

there will be a lag in detection when prevalence is low since low densities likely limit

transmission (Gagnier, Laurion, & DeNicola, 2020). Unfortunately, a lag in detection

will likely permit the establishment of CWD and, over time, eradication becomes

nearly impossible and costs are high (Mysterud & Rolandsen, 2018). Sustained

temporal monitoring of variation in PRNP across CWD-free regions could be a

detection tool as our simulations suggest detectable shifts should occur with the

arrival of the disease. By combining demographic patterns and genotypes with

current risk models, managers could improve risk-based detection efforts and

facilitate a more effective resource deployment plan as the disease alters the

population.

391   **Acknowledgements**

392   Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery

393   Grants (JMN - RGPIN-2019-04330, ABAS RGPIN-2017-03934), NSERC CGS (SH),

394   Canadian Foundation for Innovation (JELF #66905), Compute Canada RRG (gme-

395   665-AB), and the Ontario Animal Health Network provided support for the project.

396   We thank Cathy Cullingham for comments, and Anh Dao for providing samples.

397

398   **Data Accessibility**

399   DNA sequences: SRA Accessions PRJNA565222 and SUB8436966

400   PRNP and RADseq analysis pipelines: https://gitlab.com/WiDGeT_TrentU/

401

402

403

404

405

406

407

408

409

410

411

412

413

 Tables

 Table 1. Allele and amino acids frequencies for reference/alternate alleles at Ontario
 white-tailed deer (n=631) prion protein gene.

Locus	Allele	Codon	Amino acid	Major frequency	Minor frequency
60	C/T	20	D/-	0.9794	0.0206
136†	A/G	46	S/G	0.9873	0.0127
153	C/T	51	R/-	0.8605	0.1395
168†	A/G	56	G/-	0.9635	0.0365
174†	T/G	58	G/-	0.9842	0.0158
177†	C/G	59	G/-	0.9873	0.0127
178†	T/G	60	W/G	0.9160	0.0840
192†	T/G	64	H/Q	0.8653	0.1347
195†	A/G,T	65	G/-	0.7147	0.2853
198†	T/G	66	G/-	0.7021	0.2979
285	A/C	95	Q/H	0.9699	0.0301
286	G/A	96	G/S	0.6609	0.3391
324	A/G	108	P/-	0.9794	0.0206
365†	G/T	122	G/V	0.9651	0.0349
378	G/A	126	G/-	0.9651	0.0349
417	G/A	139	R/-	0.1696	0.8304
548†	T/A	183	V/D	0.9778	0.0222
555	C/T	185	I/-	0.4120	0.5880

| 676 | C/A | 226 | Q/K | 0.9620 | 0.0380 |

417   † indicates novel positions.

419   **Table 2**. Genome-wide population summary statistics including breakdown of sites,
420   number of individuals, nucleotide diversity estimate (π), individual genetic variance
421   (I) relative to the subpopulation genetic variance (F_{IS}), and the observed
422   heterozygosity of white-tailed deer in Ontario.

Group	Ontario	Northern	Southern
Number individuals	182±21	57±3	125±15
Total sites	13,439,671	14,042,841	12,885,176
Polymorphic sites	165,254	139,582	146,036
π	$5.0x10^{-4}$	$6.4x10^{-4}$	$5.7x10^{-4}$
Observed heterozygosity	$4.7x10^{-4}$	$6.1x10^{-4}$	$5.5x10^{-4}$
F_{IS}	$5.7x10^{-4}$	$3.9x10^{-4}$	$4.2x10^{-4}$

Figures

Figure 1. Distribution of free-ranging white-tailed deer samples obtained between
2003-2018 by the Ontario Ministry of Natural Resources and Forestry (OMNRF) that
were used for the reduced representation genome analysis (n=190; cream) and the
prion protein genetic analysis (n=631; orange *and* cream). The natural distribution of
free-ranging white-tailed deer is shown for Ontario with a darker shade of gray.

Figure 2. A 771 bp region of the white-tailed deer prion protein gene was analyzed
from free-ranging white-tailed deer in Ontario, Canada (n=631). The overlayed
genotypes across 19 variable loci were organized by broad management in Ontario
and are shown. Circles indicate loci previously described as variable in the white-
tailed deer prion protein gene. Triangles indicate novel variable loci in the variable
loci in the white-tailed deer prion protein gene.

Figure 3. A plot of PC1 and PC2 from the principal component analysis (PCA) on the
reduced representation white-tailed deer genome. PC1 and PC2 were able to explain
a total of 4.3% of the genomic variation observed. Gray circles represent scores from
samples in northern Ontario. Orange circles represent scores from samples in
southern Ontario.

Model	LL	Parameter	Estimate	BU
		nuB	548.48	35.37
BOTTLEGROWTH	-2034	nuF	139.97	30.83
		T	26.10	85.78

Figure 4. Demographic parameter estimates from δaδi for the optimal 1D model for white-tailed deer in Ontario. Parameter estimates are the ancient population size (N_a), the ratio of population size after instantaneous change to ancient population size (nuB); the ratio of contemporary to ancient population size (nuF); and the time in the past at which instantaneous change happened and growth began (T_C; in units of $2*N_a$ generations). Included are the optimized log-likelihood (LL) and bootstrap uncertainties (BU).

456
457 **Figure 5.** Simulated allele frequency projections for nucleotide positions 285 (a) and
458 286 (b) in the PRNP gene under selection. Green and red lines are the selection
459 coefficients provided by Robinson et al (2012); while the black dots are those from
460 our simulations, with blue line representing the mean selection coefficient.

References
Aguirre, A.A. (2017). Changing patterns of emerging zoonotic diseases in wildlife,
domestic animals, and humans linked to biodiversity loss and globalization. ILAR
Journal, 58(3), 315-318. doi: 10.1093/ilar/ilx035

Allen, C.D., Breshears, D.D., & McDowell, N.G. (2015). On underestimation of global
vulnerability to tree mortality and forest die-off from hotter drought in the
Anthropocene. Ecosphere, 6(8), 129. doi:10.1890/ES15-00203.1

Averill, K.M., Mortensen, D.A., Smithwick, E.A.H., Kalisz, S., McShea, W.J., Bourg, N.A.,
Parker, J.D., ... & Nuzzo, V.A. (2018). A regional assessment of white-tailed deer effects
on plant invasion. AoB PLANTS, 10(1), plx047. doi: 10.1093/aobpla/plx047

Baldwin, D.J.B., Desloges, J.R., & Band, L.E. (2000). Physical Geography of Ontario. In
Perera, A.H.; Euler, D.L.; Thompson, I.D. (1st Ed.). Ecology of a managed terrestrial
landscape: Patterns and processes of forest landscapes in Ontario. Ontario Ministry of
Natural Resources, Toronto, Ontario, and UBC Press, Vancouver, British Columbia.

Barber-Meyer, S.M., & Mech, L.D. (2016). White-tailed deer (Odocoileus virginianus)
subsidize gray wolves (Canis lupus) during a moose (Alces americanus) decline: A
case of apparent competition? Canadian Field Naturalist, 130(4), 308-314. doi:
10.22621/cfn.v130i4.1924

Belant, J.L., & Deese, A.R. (2010). Importance of wildlife disease surveillance. Human-
Wildlife Interaction, 4(2), 165-169. Retrieved August 11, 2010, from
www.jstor.org/stable/24868837

Bourret, V., Albert, V., April, J., Côté, G., & Morissette, O. (2020). Past, present and
future contributions of evolutionary biology to wildlife forensics, management and
conservation. Evolutionary Applications, 13(6), 1420– 1434. doi: 10.1111/eva.12977

Brandt, A.L., Green, M.L., Ishida, Y., Roca, A.L., Novakofski, J., & Mateus-Pinilla, N.E.
(2018). Influence of the geographic distribution of prion protein gene sequence
variation on patterns of chronic wasting disease spread in white-tailed deer
(Odocoileus   virginianus).   Prion,   12(3-4),   204–215.   doi:
10.1080/19336896.2018.1474671

Brandt, A.L., Kelly, A.C., Green, M.L., Shelton, P., Novakofski, J., & Mateus-Pinilla, N.E.
(2015). Prion protein gene sequence and chronic wasting disease susceptibility in
white-tailed   deer   (Odocoileus   virginianus).   Prion,   9(6),   449–462.   doi:
10.1080/19336896.2015.1115179

Chafin, T.K., Douglas, M.R., Martin, B.T., Zbinden, Z.D., Middaugh, C.R., Ballard, J.R.,
Gray, M.C.,...& Douglas, M.E. (2020). Age structuring and spatial heterogeneity in prion

protein gene (PRNP) polymorphism in white-tailed deer. bioRxiv. doi:
10.1101/2020.07.15.205039v2

Charlesworth, B., & Charlesworth, D. (2010). Elements of evolutionary genetics.
Roberts & Company. doi: 10.1017/S001667231000042X

Chen, L., Qui, Q., Jiang, Y., Wang, K., Lin, Z., Li, Z., … et al. (2019). Large-scale ruminant
genome sequencing provides insights into their evolution and distinct traits. Science,
364(6446), eaav6202. doi: 10.1126/science.aav6202

Dawe, K.L., & Boutin, S. (2016). Climate change is the primary driver of white-tailed
deer (Odocoileus virginianus) range expansion at the northern extent of its range;
land use is secondary. Ecology and Evolution, 6(18), 6435–6451. doi:
10.1002/ece3.2316

Escobar, L.E., Moen, R., Craft, M.E., & VanderWaal, K.L. (2019). Mapping parasite
transmission risk from white-tailed deer to a declining moose population. European
Journal of Wildlife Research, 65. doi: 10.1007/s10344-019-1297-z

Escobar, L.E., Pritzkow, S., Winter, S.N., Grear, D.A., Kirchgessner, M.S., Dominquez-
Villegas, E., Machado, G., Peterson, A.T., & Soto, C. (2020). The ecology of chronic
wasting disease in wildlife. Biological Reviews, 95(2), 393-408. doi:
10.1111/brv.12568

Evira (2018). Moose Found Dead in Forest with Chronic Wasting Disease. Finnish
Food Safety Authority. [Website]. Retrieved from https://mmm.fi/en/-/evira-
metsaan-kuolleella-hirvella-naivetystauti

Ferretti, F., & Mori, E. (2020). Displacement interference between wild ungulate
species: does it occur? Ethology Ecology & Evolution, 32(1), 2-15. doi:
10.1080/03949370.2019.1680447

Fischer, K., Kreyling, J., Beaulieu, M., Beil, I., Bog, M., Bronte, D., Holm, S., … & Gienapp,
P. (2020). Species-specific effects of thermal stress on the expression of genetic
variation across a diverse group of plant and animal taxa under experimental
conditions. Heredity. doi: 10.1038/s41437-020-0338-4

Frerker, K., Sabo, A., & Waller, D. (2017). Correction: Long-term regional shifts in plant
community composition are largely explained by local deer impact experiments. PLoS
One, 12(9), e0185037. doi: 10.1371/journal.pone.0185037

Gagnier, M., Laurion, I., & DeNicola, A.J. (2020). Control and surveillance operations to
prevent chronic wasting disease establishment in free-ranging white-tailed deer in
Québec, Canada. Animals, 10(283). doi: 10.3390/ani10020283

Gillespie, J.H. (2004). Population genetics: A concise guide (2nd ed.). Baltimore and
London: The Johns Hopkins University Press.

Güere, M. E., Våge, J., Tharaldsen, H., Benestad, S. L., Vikøren, T., Madslien, K., Hopp, P.,
Rolandsen, C. M., Røed, K. H., & Tranulis, M. A. (2020). Chronic wasting disease
associated with prion protein gene (PRNP) variation in Norwegian wild reindeer
(Rangifer tarandus). Prion, 14(1), 1–10. doi: 10.1080/19336896.2019.1702446

Haley, N. J., Merrett, K., Buros Stein, A., Simpson, D., Carlson, A., Mitchell, G.,
Staskevicius, A., Nichols, T., Lehmkuhl, A. D., & Thomsen, B. V. (2019). Estimating
relative CWD susceptibility and disease progression in farmed white-tailed deer with
rare PRNP alleles. PLoS One, 14(12), e0224342. doi: 10.1371/journal.pone.0224342

Harmon, J.P., Moran, N.A., & Ives, A.R. (2009). Species response to environmental
change: Impacts of food web interactions and evolution, Science, 323(5919), 1347-
1350. doi: 10.1126/science.1167396

Hashida, Y., Withey, J., Lewis, D.J., Newman, T., & Kline, J.D. (2020). Anticipating
changes in wildlife habitat induced by private forest owners' adaptation to climate
change and carbon policy. PLoS One, 15(4), e0230525. doi:
10.1371/journal.pone.0230525

Heaton, M.P., Leymaster, K.A., Freking, B.A., Hawk, D.A., Smith, T.P., Keele, J.W.,
Snelling, W.M., … & Laegreid, W.W. (2003). Prion gene sequence variation within
diverse groups of U.S. sheep, beef cattle, and deer. Mammalian Genome Society,
14(11), 765–777. doi: 10.1007/s00335-003-2283-y

Herrera, J., & Nunn, C.L. (2019). Behavioural ecology and infectious disease:
implications for conservation of biodiversity. Philosophical transactions of the Royal
Society of London Series B Biological sciences, 374(1781), 20180054. doi:
10.1098/rstb.2018.0054

Hewitt, D.G. (2011). Biology and management of white-tailed deer. (ed.). Boca Raton,
Fla: CRC Press.

Igel-Egalon, A., Laferrière, F., Tixador, P., Moudjou, M., Herzog, L., Reine, F., Torres, J.
M., … & Béringue, V. (2020). Crossing species barriers relies on structurally distinct
prion assemblies and their complementation. Molecular Neurobiology, 57(6), 2572–
2587. doi: 10.1007/s12035-020-01897-3

Johnson, C., Johnson, J., Clayton, M., McKenzie, D., & Aiken, J. (2003). Prion protein gene
heterogeneity in free-ranging white-tailed deer within the chronic wasting disease
affected region of Wisconsin. Journal of Wildlife diseases, 39(3), 576–581. doi:
10.7589/0090-3558-39.3.576

Johnson, C., Johnson, J., Vanderloo, J. P., Keane, D., Aiken, J.M., & McKenzie, D. (2006a).
Prion protein polymorphisms in white-tailed deer influence susceptibility to chronic
wasting disease. The Journal of General Virology, 87(7), 2109–2114. doi:
10.1099/vir.0.81615-0

Johnson, C.J., Phillips, K.E., Schramm, P.T., McKenzie, D., Aiken, J.M., & Pedersen, J.A.
(2006b). Prions adhere to soil minerals and remain infectious. PLoS Pathogens, 2(4),
e32. doi: 10.1371/journal.ppat.0020032

Jombart, T., Devillard, S., & Balloux, F. (2010). Discriminant analysis of principal
components: a new method for the analysis of genetically structured populations.
BMC Genetics, 11, 94. doi: 10.1186/1471-2156-11-94

Kardos, M., & Shafer, A.B.A. (2018). The peril of gene-targeted conservation. Trends in
Ecology and Evolution, 33(11), 827-839. doi: 10.1016/j.tree.2018.08.011.

Kelly, A.C., Mateus-Pinilla, N.E., Diffendorfer, J.,Jewell, E., Ruiz, M.O., Killefer, J., Shelton,
P., Beissel, T., & Novakofski, J. (2008). Prion sequence polymorphisms and chronic
wasting disease resistance in Illinois white-tailed deer (Odocoileus virginianus).
Prion, 2(1), 28–36. doi: 10.4161/pri.2.1.6321

Kennedy-Slaney, L., Bowman, J., Walpole, A.A., & Pond, B.A. (2018). Northward bound:
The distribution of white-tailed deer in Ontario under a changing climate. Wildlife
Research, 45(3), 220-228. doi: 10.1071/WR17106

Lambert, S., Ezanno, P., Garel, M., & Gilot-Fromont, E. (2018). Demographic
stochasticity drives epidemiological patterns in wildlife with implications for diseases
and population management. Scientific Reports, 8, 16846. doi: 10.1038/s41598-018-
34623-0

Latch, E.K., Heffelfinger, J.R., Fike, J.A., & Rhodes, O.E. (2009). Species-wide
phylogeography of North American mule deer (Odocoileus hemionus): cryptic glacial
refugia and postglacial recolonization. Molecular Ecology, 18(8), 1730-1745. doi:
10.1111/j.1365-294X.2009.04153.x

Latham, A.D.M., Latham, M.C., Knopff, K.H., Hebblewhite, M., & Bouting, S. (2013).
Wolves, white-tailed deer, and beaver: implications of seasonal prey switching for
woodland caribou declines. Ecography, 36(12), 1276-1290. doi: 10.1111/j.1600-
0587.2013.00035.x

Mateus-Pinilla, N., Weng, H.Y., Ruiz, M.O., Shelton, P., & Novakofski, J. (2013).
Evaluation of a wild white-tailed deer population management program for
controlling chronic wasting disease in Illinois, 2003–2008. Preventive Veterinary
Medicine, 110(3-4), 541-548. doi: 10.1016/j.prevetmed.2013.03.002

McKnight, D.T., Schwarzkopf, L., Alford, R.A., Bower, D.S., & Zenger, K.R. (2017). Effects
of emerging infectious disease on host population genetics: a review. Conservation
Genetics, 18, 1235-1245. doi: 10.1007/s10592-017-0974-2

Messer, P.W., Ellner, S.P., & Hairson Jr., N.G. (2016). Can Population Genetics Adapt to
Rapid Evolution? Trends in Genetics, 32(7), 408-418. doi: 10.1016/j.tig.2016.04.005

Miguel, E., Grosbois, V., Caron, A., Pople, D., Roche, B., & Donnelly, C.A. (2020). A
systemic approach to assess the potential and risks of wildlife culling for infectious
disease control. Communications Biology, 3(353). doi: 10.1038/s42003-020-1032-z

Miller, M.W., & Williams, E.S. (2003). Prion disease: horizontal prion transmission in
mule deer. Nature, 425(6953), 35-36. doi: 10.1038/425035a

Miller, M.W., Williams, E.S., Hobbs, N.T., & Wolfe, L.L. (2004). Environmental sources
of prion transmission in mule deer. Emerging Infectious Diseases, 10(6), 1003-1006.
doi: 10.3201/eid1006.040010

Miller, M.W., Williams, E.S., McCarty, C.W., Spraker, T.R., Kreeger, T.J., Larsen, C.T., &
Thorne, E.T. (2000). Epizootiology of chronic wasting disease in free-ranging cervids
in Colorado and Wyoming. Journal of Wildlife Diseases, 36(4), 676-690. doi:
10.7589/0090-3558-36.4.676

Miller, W.L., & Walter, D.W. (2019a). Can genetic assignment tests provide insight on
the influence of captive egression on the epizootiology of chronic wasting disease?
Evolutionary Applications, 13(4), 715-726. doi: 10.1111/eva.12895

Miller, W.M., & Walter, D.W. (2019b). Spatial heterogeneity of prion gene
polymorphisms in an area recently infected by chronic wasting disease. Prion, 13(1),
65-76. doi: 10.1080/19336896.2019.1583042

Miller, W.M., Miller-Butterworth, C.M., Diefenbach, D.R., & Walter, W.D. (2020).
Assessment of spatial genetic structure to identify populations at risk for infection of
an emerging epizootic disease. Ecology and Evolution, 10(9), 3977-3990. doi:
10.1002/ece3.6161

Morand, S., & Walther, B.A. (2020). The accelerated infectious disease risk in the
Anthropocene: more outbreaks and wider global spread. bioRxiv, 049866. doi:
10.1101/2020.04.20.049866

Mysterud, A., & Rolandsen, C.M. (2018). A reindeer cull to prevent chronic wasting
disease in Europe. Nature Ecology and Evolution, 2, 1343-1345. doi:
10.1038/s41559-018-0616-1

O'Rourke, K.I., Besser, T.E., Miller, M.W., Cline, T.F., Spraker, T.R., Jenny, A.L., Wild, M.A.,
... & Williams, E.S. (1998). PrP genotypes of captive and free-ranging Rocky Mountain
elk (Cervus elaphus nelsoni) with chronic wasting disease. The Journal of General
Virology, 80(10), 2765–2679. doi: 10.1099/0022-1317-80-10-2765

O'Rourke, K.I., Spraker, T.R., Hamburg, L.K., Besser, T.E., Brayton, K.A., & Knowles, D.P.
(2004). Polymorphisms in the prion precursor functional gene but not the
pseudogene are associated with susceptibility to chronic wasting disease in white-
tailed deer. The Journal of General Virology, 85(5), 1339–1346. doi:
10.1099/vir.0.79785-0

Osterholm, M.T., Anderson, C.J., Zabel, M.D., Scheftel, J.M., Moore, K.A., & Appleby, B.S.
(2019). Chronic wasting disease in cervids: Implications for prion transmission to
humans and other animal species. mBio, 10(4), e01091-19. doi:
10.1128/mBio.01091-19

Parchman, T.L., Gompert, Z., Mudge, J., Schilkey, F.D., Benkman, C.W., & Buerkle, C.A.
(2012). Genome-wide association genetics of an adaptive trait in lodgepole pine.
Molecular Ecology, 21(12), 2991-3005. doi: 10.1111/j.1365-294X.2012.05513.x

Patton, S.R., Russell, M.B., Windmuller-Campione, M.A., & Frelich, L.E. (2018).
Quantifying impacts of white-tailed deer (Odocoileus virginianus Zimmerman)
browse using forest inventory and socio-environmental datasets. PLoS One, 13(8),
e0201334. doi: 10.1371/journal.pone.0201334

Peart, C.R., Tusso, S., Pophaly, S.D., Botero-Castro, F., Wu, C.C., Aurioles-Gamboa, D.,
Baird, A.B., ... & Wolf, J.B.W. (2020). Determinants of genetic variation across eco-
evolutionary scales in pinnipeds. Nature Ecology & Evolution, 4, 1095-1104. doi:
10.1038/s41559-020-1215-5

Pecl, G.T., Araújo, M.B., Bell, J.D., Blanchard, J., Bonebrake, T.C., Chen, I.C., Clark, T.D.,
Colwell, R.K., ... & Williams, S.E. (2017). Biodiversity redistribution under climate
change: Impacts on ecosystems and human well-being. Science, 355(6332), eaai9214.
doi: 10.1126/science.aai9214

Peterson, B.K., Weber, J.N., Kay, E.H., Fisher, H.S., & Hoekstra, H.E. (2012). Double
digest RADseq: an inexpensive method for de novo SNP discovery and genotyping in
model and non-model species. PLoS One, 7(5), e37135. doi:
10.1371/journal.pone.0037135

Potapov, A., Merrill, E., Pybus, M., & Lewis, M.A. (2016). Chronic wasting disease:
transmission mechanisms and the possibility of harvest management. PLoS ONE,
11(3): e0151039. doi: 10.1371/journal.pone.0151039

Price, S.J., Garner, T.W.J., Cunningham, A.A., Langton, T.E.S., & Nichols, R.A. (2016).
Reconstructing the emergence of a lethal infectious disease of wildlife supports a key
role for spread through translocations by humans. Proceedings of the Royal Society B
Biological Sciences, 283(1839), 20160952. doi: 10.1098/rspb.2016.0952

Prusiner, S.B. (1982). Novel proteinaceous infectious particles cause scrapie. Science,
216(4542), 136-144. doi: 10.1126/science.6801762

Rivera, N. A., Brandt, A. L., Novakofski, J. E., & Mateus-Pinilla, N. E. (2019). Chronic
Wasting Disease in Cervids: Prevalence, Impact and Management Strategies.
Veterinary medicine (Auckland, N.Z.), 10, 123–139. doi: 10.2147/VMRR.S197404

Rizzoli, A., Taliapietra, V., Cagnacci, F., Marini, G., Arnoldi, D., Rosso, F., & Rosà, R.
(2019). Parasites and wildlife in a changing world: The vector-host-pathogen
interaction as a learning case. International Journal for Parasitology: Parasites and
Wildlife, 9, 394-401. doi: 10.1016/j.ijppaw.2019.05.011

Robinson, S.J., Samuel, M.D., Johnson, C.J., Adams, M., & McKenzie, D.I. (2012).
Emerging prion disease drives host selection in a wildlife population. Ecological
Applications, 22(3), 1050-1059. doi: 10.1890/11-0907.1

Rousselle M., Simion P., Tilak M.K., Figuet E., Nabholz B., & Galtier, N. (2020) Is
adaptation limited by mutation? A timescale-dependent effect of genetic diversity on
the adaptive substitution rate in animals. PLoS Genetics, 16(4), e1008668. doi:
10.1371/journal.pgen.1008668

Russell, M.B., Woodall, C.W., Potter, K.M., Walters, B.F., Domke, G.M., & Oswalt, C.M.
(2017). Interactions between white-tailed deer density and the composition of forest
understories in the northern United States. Forest Ecology and Management, 384, 26-
33. doi: doi.org/10.1016/j.foreco.2016.10.038

Samuel, M.D., Liao, W., Atkinson, C.T., & LaPointe D.A. (2020). Facilitated adaptation
for conservation – Can gene editing save Hawaii's endangered birds from climate
driven avian malaria? Biological Conservation, 241, 108390. doi:
j.biocon.2019.108390

Schulte, L.A., Mladenoff, D.J., Crow, T.R., Merrick, L.C., & Cleland, D.T. (2007).
Homogenization of northern U.S. Great Lakes forests due to land use. Landscape
Ecology, 22, 1089–1103. doi: 10.1007/s10980-007-9095-5

Seabury, C.M., Oldeschulte, D.L., Bhattarai, E.K., Legare, D., Ferro, P.J., Metz, R.P., &
Johnson, C.D. (2020). Accurate genomic predictions for chronic wasting disease in U.S.
white-tailed deer. G3: Genes, Genomes, Genetics, 10(4), 1433-1441. doi:
10.1534/g3.119.401002

Taylor, P.D., Crewe, T.L., Mackenzie, S.A., Lepage, D., Aubry, Y., Crysler, Z., Finney, G., …
& Woodworth, B.K. (2017). The motus wildlife tracking system: a collaborative
research network to enhance the understanding of wildlife movement. Avian
Conservation and Ecology, 12(1), 8. doi: 10.5751/ACE-00953-120108

Thompson, T.Q., Bellinger, M.R., O'Rourke, S.M., Prince, D.J., Stevenson, A.E., Rodrigues,
A.T., … & Miller, M.R. (2019). Anthropogenic habitat alteration leads to rapid loss of
adaptive variation and restoration potential in wild salmon populations. Proceedings
of the National Academy of Sciences of the United States of America, 116(1), 177-186.
doi: 10.1073/pnas.1811559115

Vázquez-Miranda, H., & Zink, R.M. (2020). Geographic distribution of chronic wasting
disease resistant alleles in Nebraska, with comments on the evolution of resistance.
Journal of Fish and Wildlife Management, 11(1), 46-55. doi: 10.3996/012019-JFWM-
002

Walter, W.D., VerCauteren, K.C., Campa, H., Clark, W.R., Fischer, J.W., Hygnstrom, S.E.,
… & Winterstein, S.R. (2009). Regional assessment on influence of landscape
configuration and connectivity on range size of white-tailed deer. Landscape Ecology,
24, 1405–1420. doi: 10.1007/s10980-009-9374-4

Weiskopf, S.R., Ledee, O.., & Thompson, L. M. (2019). Climate change effects on deer
and moose in the Midwest. The Journal of Wildlife Management, 83(4), 769-781.
doi:10.1002/jwmg.21649

White, K.S., Gregovich, D.P., Levi, T. (2017). Projecting the future of an alpine ungulate
under climate change scenarios. Global Change Biology, 24(3), 1136-1149. doi:
10.1111/gcb.13919

Wilson, G.A., Nakada, S.M., Bollinger, T.K., Pybus, M.J., Merrill, E. H., & Coltman, D.W.
(2009). Polymorphisms at the PRNP gene influence susceptibility to chronic wasting
disease in two species of deer (Odocoileus Spp.) in western Canada. Journal of
Toxicology and Environmental Health Part A, 72(17-18), 1025–1029. doi:
10.1080/15287390903084264

Supplementary Materials

741 **Table S1**. Four degenerate primers used to amplify the functional white-tailed deer prion
742 protein gene region as two overlapping fragments are shown. Fragment 1 is 460 base-pairs
743 in length and Fragment 2 is 590 base-pairs in length.

Primer	Fragment 1	Fragment 2
Forward (5'-3')	TCGTCGGCAGCGTCAGATGTGTA TAAGAGACAGACRTGGGCATAT GATGCTGAYACC	TCGTCGGCAGCGTCAGATGTGTATAA GAGACAGTGGAGGCTGGGGTCAAGG
Reverse (5'-3')	GTCTCGTGGGCTCGGAGATGTGT ATAAGAGACAGYTGCCAAAATG TATAAGAGG	GTCTCGTGGGCTCGGAGATGTGTATA AGAGACAGACTACAGGGCTGCAGGTA GAYACT

758 **Table S2**. Thermal cycling conditions for PRNP PCR amplification for Fragment 1 and
759 Fragment

Fragment 1	Fragment 2
98℃ for 2 min	
10 cycles of:	
98℃ for 15 sec	98℃ for 15 sec
60-55℃ for 30 sec[1]	65-60℃ for 30 sec[1]
72℃ for 45 sec	72℃ for 45 sec
98℃ for 15 sec	
20 cycles of:	
98℃ for 15 sec	98℃ for 15 sec
55℃ for 30 sec	60℃ for 30 sec
72℃ for 45 sec[2]	72℃ for 45 sec[2]
72℃ for 7 min	

1 – decrease by 0.5° per cycle; 2 – increasing by 5
sec per cycle

767 **Table S3**. Mixture for double restriction enzyme digestion. Samples and reaction mixture
768 were incubated at 37°C for 3 hours and then 25°C overnight on a thermal cycler with a heated
769 lid. Samples and reaction mixture were head-inactivated at 85°C for 30 minutes.

Reagent	1 X	Final concentration
10x Cutsmart	2.00 ul	1 X
H_2O	11.85 ul	
Msel (10 U/ μl)	0.10 ul	1 U
Sbfl-HF (20 U/ μl)	0.05 ul	1 U
Subtotal	14.00 ul	
DNA	x ul	$\sim$ 500 ng
H_2O	6.00 – x ul	
Subtotal	6.00 ul	
Final Volume	20.00 ul	

771 **Table S4**. Mixture for adapter ligation. The digested fragments were combined with 7 ul of
772 the adapter ligation mixture and 3 ul of a unique barcoded Sbfl adapter (1.0 uM). The ligation
773 on $\sim$ 30 ul reaction mixture was performed in 16°C for 3 hours

Reagent	1 X	Final concentration
10x CutSmart	1.00	1 X
100 mM ATP	0.30	1 mM
H_2O	2.45	
Msel Y adapter (10 uM)	3.00	1 μM
T4 DNA ligase (400 U/μl)	0.25	100 U
Subtotal	7.00 ul	

776   **Table S5**. Illumina PCR mixture. The purified restriction-ligation DNA (3 ul) was combined
777   with 7 ul of PCR mixture and a PCR was performed on 10 ul. Four replicated per sample was
778   performed. The thermal cycler profile for this PCR was 98°C for 30 seconds; 20 cycles of 98°C
779   for 20 seconds, 60°C for 30 seconds, 72°C for 45 seconds; and a final extension at 72°C for 5
780   minutes.

Reagent	1 X	Final concentration
H_2O	1.33	
KAPA HiFi ReadyMix	5.00	2 X
PCR primer mix (5 µM each)	0.67	0.5 µM
Subtotal	7.00 ul	

Table S6. The major/minor allele counts for four nucleotide (nt) positions of variation in the
white-tailed deer prion protein gene that are linked to reduced susceptibility or reduced
clinical progression of chronic wasting disease for Northern and Southern Ontario are shown.
A two-sided Fisher's Exact test was conducted on the major and minor allele counts at the
four chronic wasting disease-linked nucleotide positions. A p-value less than 0.05 was
considered significant and indicated that there were significant differences between the
groups.

Location	nt60	nt285	nt286	nt676
Northern	186/3	182/7	132/57	187/2
Southern	432/10	430/12	285/157	420/22
Southeastern	306/8	308/6	196/118	300/14
Southwestern	126/2	122/6	89/39	120/8
p-value	0.764	0.611	0.200	0.021

Table S7. Haplotypes were estimated with PHASE v2.1.1 set to the same parameters in Brandt et al., 2018: Markov chain Monte Carlo (MCMC) samples were taken from a minimum of 100,000 steps, with a discarded burn-in of 10,000; samples were drawn every 100 MCMC steps. Five repetitions were performed, and haplotype frequencies compared to verify consistent assignment. Included are estimates of population haplotypes with frequencies of greater than 1% (count=1262, number haplotypes = 151) and associated estimated standard deviations (S.E.; square root of the variance of the posterior distribution) at 19 variable positions, with 0 representing non-variants and 1 representing variants.

ID	f	S.E.	Codon	153	195	198	286	365	378	417	555
3	0.228	0.006	-	0	0	0	0	0	0	1	0
1	0.122	0.005	-/-	0	0	0	0	0	0	1	1
9	0.104	0.001	Ref	0	0	0	0	0	0	0	0
16	0.087	0.005	96S/-/-	0	0	0	1	0	0	1	1
7	0.050	0.001	-	0	0	0	0	0	0	0	1
252	0.041	0.003	-/-	1	0	0	0	0	0	1	0
18	0.033	0.004	96S/-	0	0	0	1	0	0	1	0
54	0.022	0.003	-/-/-	0	1	1	0	0	0	1	0
259	0.012	0.002	-/122V/-/-	1	0	0	0	1	1	1	0
27	0.012	0.002	-/-/-	0	0	1	0	0	0	1	1
28	0.011	0.003	-/96S/-/-	0	0	1	0	0	0	1	0
48	0.010	0.002	-/96S/-/-	0	1	0	1	0	0	1	1

806　**Table S8**. Optimized log-likelihood (LL) and bootstrap uncertainties (BU) obtained from 1D
807　demographic models on Ontario white-tailed deer as a single population. Model specifics and
808　parameters are outlined. The most optimal model for 1D is shown in bold. Modified 1D
809　demographic models are indicated with an asterisk. All time estimates are reported in units
810　of 2*Na generations. Migration rate (m) is reported in units of 2*Na*m.

Model Name	LL	Parameter	Estimate	BU
SNM	-25746	n/a	n/a	41.3
				$1.11x10^{-15}$
				$7.77x10^{-16}$
TWO_EPOCH	-3015	nu	573.38	3.97
				40.69
		T	107.90	7.62
GROWTH	-8795	nu	21.57	217.02
				8.93
		T	6.58	3.79
BOTTLEGROWTH	-2034	nuB	548.48	35.37
		nuF	139.97	30.83
		T	26.10	85.78
BOTTLEPOP*	-8495	nu	54.48	287.41
		T	20.17	78.28
				46.48
TWOPOPCHANGES	-3129	nuB	$4.19x10^{-5}$	$5.16x10^{4}$
		nuF	0.32	
		TB	$5.14x10^{-3}$	$2.06x10^{-5}$

		TF	$6.04x10^{-2}$	0.11
GROWTHPLUSBOTTLE	-25631	nuB	8.47	$1.43x10^5$
		nuF	0.14	
		TB	0.92	2.74
		TF	4.28	$2.39x10^{-2}$
BOTTPLEPLUSGROWTH	-3544	nuB	397.36	8.75
		nuF	473.75	93.69
		T	97.40	169.23
THREE_EPOCH	-3274	nuB	$1.35x10^{-4}$	$1.99x10^5$
		nuF	$1.31x10^{-2}$	$1.46x10^{-5}$
		TB	$3.48x10^{-3}$	$2.64x10^{-3}$
		TF	$2.47x10^{-3}$	

Table S9. Nucleotide variations in free-ranging white-tailed deer prion protein gene that are associated with chronic wasting disease are either protective (1), increase susceptibility (2), or are neutral. The major and minor allele frequencies for each site across a 771 bp region of the prion protein gene in white-tailed deer are reported for different regions, in descending order. The year CWD was found in free-ranging cervids is reported for each location. The data are from free-ranging white-tailed deer samples collected in: Alberta, Canada (AB); Colorado, USA (COL); Illinois, USA (IL); Ontario, Canada (ON); Nebraska, USA (NE); Saskatchewan, Canada (SK); Wisconsin, USA (WI); and Wyoming, USA (WY).

Site	Role	Major	Minor	Region	CWD+	Citation
C60T	1	0.98	0.02	ON	n/a	Haworth et al., 2020
		0.98	0.02	WY	1985	Heaton et al., 2003
		0.94	0.06	AB, SK	2002,1996	Wilson et al., 2009
		0.92	0.08	IL	2002	Kelly et al., 2008
C153T	2	0.96	0.04	WY	1985	Heaton et al., 2003
		0.94	0.06	AB, SK	2002,1996	Wilson et al., 2009
		0.89	0.11	IL	2002	Kelly et al., 2008
		0.86	0.14	ON	n/a	Haworth et al., 2020
A285C	1	1.00	0.00	WI	2002	Johnson et al., 2006
		0.99	0.01	AB, SK	2002,1996	Wilson et al., 2009
		0.98	0.02	NE	1999	Vázquez-Miranda & Zink, 2020
		0.97	0.03	ON	n/a	Haworth et al., 2020
		0.94	0.06	IL	2002	Kelly et al., 2008
G286A	1	0.88	0.12	COL	1967	O'Rourke et al., 1998 (unpublished)
		0.86	0.14	IL	2002	Kelly et al., 2008
		0.83	0.17	NE	1999	Vázquez-Miranda & Zink, 2020
		0.81	0.19	WI	2002	Johnson et al., 2006

		0.66	0.34	ON	n/a	Haworth et al., 2020
A324G	1	0.98	0.02	ON	n/a	Haworth et al., 2020
		0.98	0.02	NE	1999	Vázquez-Miranda & Zink, 2020
		0.96	0.04	WY	1985	Heaton et al., 2003
		0.96	0.04	AB, SK	2002,1996	Wilson et al., 2009
		0.94	0.06	WY	1985	Heaton et al., 2003
G417A	3	0.99	0.01	AB, SK	2002,1996	Wilson et al., 2009
		0.79	0.21	COL	1967	O'Rourke et al., 1998
		0.17	0.83	ON	n/a	Haworth et al., 2020
C555T	1	0.89	0.11	n/a	n/a	Raymond et al., 2000
		0.65	0.35	AB, SK	2002,1996	Wilson et al., 2009
		0.58	0.42	IL	2002	Kelly et al., 2008
		0.41	0.59	ON	n/a	Haworth et al., 2020
C676A	1	0.99	0.01	IL	2002	Kelly et al., 2008
		0.98	0.02	AB, SK	2002,1996	Wilson et al., 2009
		0.97	0.03	WI	2002	Johnson et al., 2006
		0.96	0.04	ON	n/a	Haworth et al., 2020

Figure S1. The Canadian province of Ontario as managed by the Ontario Ministry of Natural Resources and Forestry. There are three broad regions Ontario is managed by: (A) Northern Ontario, (B) Southeastern Ontario, and (C) Southwestern Ontario. Collectively (B) and (C) form Southern Ontario. Outlined in red are the wildlife management units designated within each broad region.

Value of BIC versus number of clusters

Figure S2. The Bayesian information criterion (BIC) from a population cluster identification using successive K-means cluster assignment on the reduced representation white-tailed deer genome from identified one cluster as optimal.