Singe cell oil (SCO) based bioactive compounds: II. Enzymatic synthesis of glucose fatty acid

esters using SCOs as acyl group-donors and their biological activities

Hatim A. El-Baz[a,b], Ahmed M. Elazzazy[c,d], Tamer S. Saleh[e], Marianna Dourou[f], Jazem A.

Mahyoub[g], Mohammed N. Baeshen[c], Hekmat R. Madian[h], and George Aggelis[c,f*]

[a] Department of Clinical Biochemistry, College of Medicine, University of Jeddah, Jeddah

21589, Saudi Arabia.

[b] Biochemistry Dept., Genetic Engineering and Biotechnology Division, National Research

Centre, Cairo 12622, Egypt

[c] Department of Biology, College of Science, University of Jeddah, Jeddah 21589, Saudi Arabia

[d] Department of Chemistry of Natural and Microbial Products, Division of Pharmaceutical and

Drug Industries, National Research Centre, Cairo 12622, Egypt.

[e] Department of Chemistry, College of Science, University of Jeddah, Jeddah 21589, Saudi

Arabia

[f] Unit of Microbiology, Division of Genetics, Cell and Developmental Biology, Department of

Biology, University of Patras, Patras 26504, Greece

[g] Department of Biological Sciences, Faculty of Science, King Abdulaziz University, Jeddah

21589, Saudi Arabia.

[h] Department of Processes Design & Development, Egyptian Petroleum Research Institute

(EPRI), Nasr City, Cairo 11727, Egypt

[*]Corresponding author: Professor George Aggelis, email address: George.Aggelis@upatras.gr

Unit of Microbiology, Division of Genetics, Cell and Developmental Biology, Department of

Biology, University of Patras, Patras 26504, Greece

Abstract

Sugar fatty acid esters, especially glucose fatty acid esters (GEs), have broad applications in food, cosmetic and pharmaceutical industries. In this research the fatty acid moieties derived from polyunsaturated fatty acid containing single cell oils (SCOs), i.e. those produced from *Cunninghamella echinulata* , *Umbelopsis isabellina* and *Nannochloropsis gaditana* as well as from olive oil and an eicosapentaenoic acid (EPA) concentrate were converted into GEs by enzymatic synthesis, using lipases as biocatalysts. The GE synthesis was monitored using thin-layer chromatography, FT-IR and in situ NMR. It was found that GE synthesis carried out using immobilized *Candida antarctica* B lipase was very effective reaching high yields, near to 100%. It was shown that EPA-GEs were very effective against several pathogenic bacteria and their activity can be attributed to their high EPA content. Furthermore, *C. echinulata*-GEs were more effective against pathogens comparing to *U. isabellina*-GEs, probably due to the presence of gamma linolenic acid (GLA) in the lipids of *C. echinulate*, which is known for its antimicrobial activity, in higher concentrations. *C. echinulata*-GEs also showed a strong insecticidal activity against *Aedes aegypti* larvae, followed by EPA-GEs, olive oil-GEs, and *N. gaditana*-GEs. All synthesized GEs induced apoptosis of the SKOV-3 ovarian cancer cell line, with the apoptotic rate increasing significantly after 48 h. A higher percentage of apoptosis was observed in the cells treated with EPA-GEs, followed by *C. echinulata*-GEs, *U. isabellina*-GEs and olive oil-GEs. We conclude that SCOs can be used in the synthesis of GEs with interesting biological properties.

Keywords: Immobilized lipases; Microbial oils; Fatty acid methyl esters; Glucose esters synthesis; Antimicrobial; Insecticidal; Anti-cancer activity

[2]

1. Introduction

Sugar fatty acid esters, so-called sugar esters (SEs), are biodegradable, odorless, non-irritating and non-toxic surfactants with broad applications in food, cosmetic and pharmaceutical industries [1–4]. Moreover, SEs have gained attention thanks to their anti-bacterial (e.g. against numerous pathogenic species of gram-positive and gram-negative bacteria) and anti-fungal activity [5], while they are also reported as insecticide and miticide [6].

SEs are synthesized from renewable resources, such as sugars and fatty acids (FAs). Different types of sugars (e.g. sucrose, fructose, glucose and lactose) can be used as acyl-acceptors to produce SEs by esterification with FAs or transesterification with FA esters as acyl-donors [7–9]. Glucose, a cheap and broadly available carbohydrate, having only one primary hydroxyl group, which predetermines a highly regioselective synthesis of SEs called glucose fatty acid esters (GEs) [10, 11]. Concerning acyl-donors, polyunsaturated fatty acids (PUFAs) can be used for this purpose, as it is known that PUFAs or PUFA derivatives possess interesting biological activities [12–14]. GEs can be synthesized both chemically and enzymatically [15, 16]. Lately, the main efforts in the processing of GEs have been focused on enzymatic synthesis because this is an advantageous method, requiring less energy and reducing solvent toxicity [11, 17], with lipases being the most important enzymes used for this synthesis [18, 19]. Commercially available immobilized lipases, such as *C. antarctica* B lipase, are used to catalyze the acylation of glucose with various FAs [20, 21].

Different sources of PUFAs can be used as sources of acyl groups, in the form of either FAs or FA esters, for GEs production. Whilst the terrestrial plant oils have been used as a source of PUFAs [22, 23], alternative and non-food supply sources, such as microalgae and fungi could be used thanks to their ability to produce microbial lipids rich in PUFAs of medicinal and

[4]

nutritional interest [24–26]. The microbial lipids, so-called single cell oils (SCOs), are

synthesized by oleaginous microorganisms that are capable of producing substantial amounts of

lipid stored within their cells [27, 28]. The genus of *Nannochloropsis* includes various marine

microalgae species able to efficiently grow under a variety of culture conditions on low quality

waters, even on wastewaters, and accumulate lipids rich in PUFAs, such as eicosapentaenoic

acid (EPA) [29–32]. Furthermore, oleaginous fungal species, such as *Cunninghamella echinulata*

and *Umbelopsis isabellina* are capable to synthesize PUFAs, especially γ-linolenic acid (GLA)

[13, 25], thus regarded as promising candidates for SCOs production [33–35].

In a previous paper [36] we produced FA amides (FAAs) using as acyl group-donors

lipids containing PUFAs in different percentages and we concluded that FAAs can be used as

bioactive compounds in various biological applications depending on their FA composition. The

aim of the current paper is to enzymatically synthesize GEs using similar to those used in FAAs

SCOs as acyl group-donors, either in the form of free fatty acids (FFAs) or in the form of fatty

acid methyl esters (FAMEs). The reaction was carried out under various conditions, using two

immobilized lipases as catalysts. SCOs, produced from various sources as describe in El-Baz et

al. [36], contained EPA, GLA or oleic acid in high percentages. The biological activity of the

aforementioned GEs against important human pathogens, the larvae of *Aedes aegypti* and the

SKOV-3 cancer cell line, was studied and compared with GEs synthesized using olive oil and an

EPA concentrate (i.e. a fish oil derivative containing EPA in very high percentages) as acyl

group-donors. We concluded that GEs derived from SCOs possess interesting biological

activities and can therefore be used in the production of pharmaceuticals in the future.

2. Materials and methods

2.1. Biological material and SCO production

The fungal strains *Cunninghamella echinulata* ATHUM 4411 and *Umbelopsis isabellina*

ATHUM 2935 (culture collection of National and Kapodistrian University of Athens, Greece)

and the microalga strain *Nannochloropsis gaditana* (culture collection CCAP 849/5) were used

as sources of SCOs. Culture conditions, cell mass harvesting, lipid extraction and purification,

FAMEs and FFAs preparation and gas chromatography analysis were described in El-Baz et al.

[36].

2.2. Enzymatic synthesis of GEs

GEs were synthesized by esterification (or transesterification) of glucose (Acros Organics,

Thermo Fisher Scientific, Waltham, MA) served as an acyl acceptor, with FFAs (or FAMEs)

served as acyl-group donors. The reaction was performed in 50-mL Erlenmeyer flasks using

different FA (or FAME):glucose molar ratios (i.e. 1:1, 1:2 or 1:3). The reactants were dissolved

in 25 mL of a solvent mixture consisted of 80% DMSO and 20% tert-amyl alcohol (Sigma

Aldrich Co.) in which 1 g of 3 Å molecular sieves was added and the mixture was sonicated for

20 min. The reaction was catalyzed by 0.25 g of immobilized *C. antarctica* B lipase (enzymatic

activity □5 Units/mg) or 0.25 g of immobilized *C. rugosa* lipase (enzymatic activity □0.1

Units/mg), both purchased from Sigma Aldrich Co., St. Louis, MO. The flasks were incubated at

$50 \pm 1\ ^{\circ}C$ in a shaking incubator at 100 rpm for 50 h.

After the incubation period the reaction mixture was filtered to remove the molecular

sieves and the immobilized lipase and the filtrates were evaporated under reduced pressure to

remove the solvent. The reaction residue was separated in ethyl acetate and distilled water (25

mL each). The organic layer, containing the synthesized GEs, was washed with 10 mL saturated

[6]

aqueous NaCl (Sigma Aldrich Co.), dried over $MgSO_4$ (Sigma Aldrich Co.), gravity-filtered and

the solvent removed under reduced pressure to get the crude product.

2.3. GE analysis

2.3.1. Thin-layer chromatography and FT-IR

Qualitative synthesis of GEs was monitored by thin-layer chromatography (TLC) as described in

El-Baz et al. [36] for FAAs. Equally, FT-IR spectra for FAMEs and the GE products were

recorded as described in the aforementioned paper. FT-IR spectra were used to detect the

formation of the ester carbonyl group confirming thus GE synthesis.

2.3.2. Quantitative determination of the reaction yields through in situ NMR monitoring

The % conversion of FAMEs to GEs was determined during the reaction via in situ NMR

monitoring. First, the protons NMR of both reactants individually were assigned and the progress

of the reaction was monitored by ^{1}H NMR at regular intervals of 10 h. The % conversion was

calculated according to the formula:

142   % Conversion = Ip / (Ir+Ip)

where Ip is the integration of the signal of the product and Ir is the integration of the signal of the

reactant.

Heteronuclear multiple bond correlation (HMBC) spectroscopy was used to identify the

ester product by determining the chemical shifts of carbon and hydrogen atoms and formation

the ester bond. NMR spectra were recorded at 298 K on a Bruker Avance III 400 (9.4 T, 400.13

MHz for 1H, 100.62 MHz for 13C) spectrometer (Bruker, Billerica, MA) with a 5-mm BBFO

[7]

probe. Chemical shifts (δ in ppm) were relative to internal standard DMSO-d6 (d 2.50) for 1H

NMR.

2.4. Biological activity of GEs

The antimicrobial, insecticidal and anticancer activities of the synthesized GEs were determined

using the protocols described in El-Baz et al. [36]. Briefly, the antimicrobial activity of GEs was

evaluated *in vitro* using the agar well diffusion assay [37], the minimum inhibitory concentration

(MIC) and minimum bactericidal concentration (MBC) [38] against human pathogens including

the Gram-negative *Escherichia coli* ATCC 25922, *Klebsiella pneumoniae* ATCC 700603,

Pseudomonas aeruginosa ATCC 15442, *Salmonella typhimurium* ATCC 14028 the Gram-

positive bacteria, *Bacillus subtilis* ATCC 6633, MRSA *Staphylococcus aureus* ATCC 4330, *S.*

aureus ATCC 25923 and the unicellular fungus *Candida albicans* ATCC 10221. The insecticidal

activity was evaluated by exposing early 4[th] instar larvae of a field strain of *Aedes aegypti* to

different concentrations of the GEs for 48 h, in glass beakers containing 100 mL of tap water.

Calculation of statistical parameters was performed using the Finney method [39]. The apoptotic

activity of the SKOV-3 ovarian cancer cell line in response to tested compounds (both FAMEs

and GEs) was determined by Annexin FITC, as per the manufacturer's instructions (BD

Biosciences, USA).

2.5. Statistical analysis

The acquired data were analyzed using SPSS 9.0 and the results were given as mean ± SD of

three replicates. The mean comparison between the various assessed groups was performed using

one-way analysis of variance (ANOVA). Statistical significance was defined when $p < 0.05$.

[8]

3. Results and discussion

3.1. Biomass and SCO production

The performance of the oleaginous microorganisms cultivated in flasks or bioreactors, as well as

the FA composition of the produced lipids, were previously presented [36]. Here, the essential

data are provided as supplementary material to facilitate reading (Tables S1, S2).

3.2. Optimization of the GE synthesis

For GE synthesis, the olive oil derived FFAs and FAMEs were used as model substrates to

optimize the lipase-catalyzed reaction of esterification and transesterification (Fig. 1). The

reaction was carried out in a solvent system consisted of tert-amyl alcohol and DMSO mixture as

describe elsewhere [11, 40, 41], for 50 h at 55 $^{\circ}$C with shaking at 100 rpm and the progress of the

reaction was monitored by TLC analysis.

The effect of various reaction conditions, such as the molar ratio of the reactants, the

reaction temperature and the type and the concentration of immobilized enzyme on the

conversion rate and yield was studied. Two immobilized lipases, lipase from *C. antarctica* B

(Lipase CA) and lipase from *C. rugosa* (Lipase CR) were used as catalysts for both esterification

and transesterification reactions for GEs production (Table 1). The employment of immobilized

lipases as catalysts for GEs synthesis has been proposed by several researchers [9, 21, 42–44].

Also, different molar ratios of olive oil FFAs (or FAMEs):glucose were tested (Table 1),

showing that the conversion yield increased with increasing the concentration of glucose,

attaining 100% conversion with lipase CA as catalyst and FAMEs:glucose ratio 1:3 (entry 6,

Table 1 and Fig. 2). Glucose can be considered as a good acyl acceptor for SE synthesis in non-

[9]

conventional media, ensuring high conversion yields, due to its relatively high solubility [45–

47]. Contrary, the use of sugars with higher degree of polymerization adversely affects the

conversion yield as a result of the low solubility of sugar in organic solvents.

Although the conversion yield obtained utilizing FFAs as acyl donor was high, this was

in all cases below to that obtained with FAMEs as substrate due to the water that was produced,

which may react with GE causing hydrolysis (Fig. 1). Contrary, the transesterification reaction

carried out using FAMEs is irreversible, with methanol as a byproduct acting as an internal water

scavenger, improving thus the conversion yield [48, 49]. Lipase CR showed lower conversion

yields (entries 7-12, Table 1) than lipase CA. Furthermore, the reusability of lipase CA was

checked for several reaction cycles for the synthesis of GE using olive oil FAMEs as substrate

under the optimized reaction conditions. Specifically, the enzyme was removed after the

completion of the reaction by filtration, washed with ethanol solvent in a Soxhlet extraction

apparatus and reused three times under the same reaction conditions. It was found that the

regenerated enzyme performed the reaction efficiently without loss of its catalytic activity, which

is important for the sustainability of the process. The reaction conditions were further optimized

using the lipase CA in different quantities as a catalyst and it was found that a 100% conversion

yield was obtained using 0.25 g of the lipase CA (Table 2, entry 4), while a higher enzyme

quantity was not necessary.

The results recorded in this study emphasized the feasibility of enzymatic synthesis of

GEs, the conversion yield of which reached 100% using lipase CA as biocatalyst. These results

are in agreement with those reported by Yan et al. [50] who demonstrated that glucose FA

monoesters were synthesized (with up to 93% yields) using lipase B from *C. antarctica*. High

yields for GE enzymatic synthesis were also reported by Sebatini et al. [51] using as catalyst

lipase-Fe_3O_4 nanoparticles. Furthermore, Findrik et al. [20] reported that the highest SE yield

was achieved using lipase CA as catalyst and glucose and palmitic acid as substrates.

3.3. Product identification and quantitative analysis

The percent conversion, which was taken as a scale to determine the optimum conditions, was

quantified via *in situ* NMR monitoring (Fig. 3). The [1]H NMR was noted new multiple signals at

4.17, 4.41 ppm, which matched the CH_2OCO grew concurrently with a decline in the intensity of

the glucose protons of C6 signals at 3.39 and 3.52 ppm. The latter signals disappeared after 50 h

of reaction when FAMEs:glucose 1:3 ratio was used, which means that 100% conversion was

achieved (Fig. 3). The main product obtained was identified as Glc (C-6)-OCOR, as indicated by

2D ([1]H-[13]C HMBC) NMR (Fig. 4a), showing correlation between the peak assigned to proton C-

6 glucose ester and the carbonyl function in olive oil FAMEs (Fig. 4b). The structure of the

obtained GEs was additionally confirmed by FT-IR analysis showing the appearance of a

broadband at 3380 cm[-1] due to the hydroxyl groups of glucose and the two stretching bands of O-

C bond at 1316 and 1015, in parallel with the disappearance of the band at 1743 cm[-1], due to the

consumption of the carbonyl group of FAMEs (Fig. 5).

3.4. GE synthesis using FAMEs from different origin

After optimization of the reaction conditions GEs were synthesized using FAMEs derived from

SCOs produced by *C. echinulata*, *U. isabellina* and *N. gaditana* and from an EPA concentrate

(Table 3). The GE conversion yield when EPA-FAMEs were used as substrate was excellent (i.e.

99%), followed by *N. gaditana*-FAMEs, *U. isabellina*-FAMEs and *C. echinulata*-FAMEs (i.e.

86, 85 and 80%, respectively – Fig. 6). The structures of the obtained GEs were confirmed on the

[11]

basis of their FT-IR spectra in which appearance of the broadband due to hydroxyl groups of

glucose was observed (Fig. 7 and Fig. S1-S5). According to the above presented data we can

conclude that the synthesis of GEs accomplished in this work was successful, while possible

large scale applications, using SCOs instead of traditional sources of PUFAs, will not interfere

with the food supply chain.

3.5. Antimicrobial activity of GEs

GEs derived from FAMEs of *C. echinulata*, *U. isabellina*, *N. gaditana* SCOs, olive oil and EPA

concentrate were tested against various human pathogens for their antimicrobial activity by the

agar well diffusion method, which resulted in the formation of a variable diameter zone of

inhibition (Table 4).

GEs produced in this work, except for *U. isabellina*-GEs, showed moderate to strong

inhibitory activity against all tested organisms. In details, *U. isabellina*-GEs showed weak

antimicrobial activity (inhibition zone ≈ 6 mm) against *S. typhimurium, P. aeruginosa* and *S.*

aureus. On the contrary, *C. echinulata*- GEs, *N. gaditana*-GEs and olive oil-GEs moderately

inhibited all tested organisms, while EPA-GEs showed the strongest antimicrobial activity

against all tested organisms, specifically against *C. albicans* (20.00 mm), *B. subtilis* and *S.*

aureus (17.00 mm). Finally, all GEs showed a higher antimicrobial activity against *C. albicans*

than against bacteria.

It was shown that EPA-GEs were very effective against all tested bacteria and their

activity can be attributed to their high EPA content. Furthermore, *C. echinulata*-GEs were more

effective against pathogens comparing to *U. isabellina*-GEs, probably due to the presence of

GLA in the lipids of *C. echinulata* in enhanced concentrations which is known for its

antimicrobial activity [13, 25].

The MIC and MBC values were determined for selected pathogens (Table 5). All tested

pathogenic strains are sensitive to all GEs being inhibited at low MIC, ranged between 6.3-50

µg/mL, and destroyed at MBC between 50-100 µg/mL. *C. echinulata*-GEs and EPA-GEs

significantly affected the growth of all the tested bacteria than *N. gaditana*-GEs and olive oil-

GEs.

The variability of the inhibitory effect found in this paper is in agreement with previous

papers reported that SEs exhibited a variable effect on different bacterial species [52, 53], while

depending on the conditions, SEs may inhibit Gram-negative [40, 54] or Gram-positive bacteria

[55, 56]. Also, depending on the dose, SEs can be either bactericidal [57] or bacteriostatic [58].

Wagh et al. [59] explained that, the inhibitory effect of SEs is dependent on the

esterification level, type (e.g. the length of aliphatic chain) and number of esterified FAs on the

sugar molecule and the nature of the carbohydrate. Furthermore, Karlová et al. [60] reported that

the antimicrobial effects of fructose esters decreased as the aliphatic chain increased. It seems

that the carbon chain length was the most important factor influencing the surface properties,

whereas degree of esterification and hydrophilic groups showed little effect [61].

The antimicrobial activity of glucose esters was tested against *E. coli*, *B. subtilis, B.*

megaterium and *B. cereus* [51, 56]. In addition, the unsaturated FAs lactose esters were shown to

exhibit antimicrobial activity against Gram-positive, Gram-negative microorganisms and fungi

[5]. The antimicrobial activity of SEs is due to autolysis caused by the interaction of the esters

with cell membranes of bacteria. The lytic action is thought to be due to the activation of

autolytic enzymes rather than the actual solubilization of the bacterial cell membrane [62].

[13]

The FAAs synthesized in El-Baz et al. [36] using lipids of similar FA composition as acyl

group-donors to those used in this paper, also exhibited a significant antimicrobial activity.

However, contrary to the results reported here, the FAAs containing oleic acid in high

percentages (i.e. derived from olive oil and *U. isabellina* oil) were more effective against human

pathogens than other FAAs.

3.6. Insecticidal activity of GEs

Aedes aegypti (the yellow fever mosquito) spreads dangerous human arboviruses including

dengue, Zika, and chikungunya. Consequently, control of yellow fever mosquitoes is a critical

public health priority [63].

The susceptibility of *A. aegypti* larvae to GEs under laboratory conditions was tested

using dipping methods. The larvicidal activity of a compound is usually improved by increasing

its concentration and exposure time, as Rodrigues et al. [64] reported for plant-derived bioactive

products, such as essential oils, ethanol extracts and FAMEs. In the current study, *C. echinulata*-

GEs showed a strong insecticidal activity against *A. aegypti* larvae with LC50 0.541 mg/L,

which could be probably attributed to the presence of GLA in significant concentrations,

followed by EPA-GEs, olive oil-GEs, and *N. gaditana*-GEs, demonstrating LC50 10.24, 12.88

and 16.92 mg/L, respectively. Contrary, *U. isabellina*-GEs were less active, presenting a LC50

equal to 39.62 mg/L (Table 6, Fig. 8). On the other hand, the RR values indicated that mosquito

of *A. aegypti* was much more susceptible to *C. echinulata*-GEs than to EPA-GEs, olive oil-GEs,

N. gaditana-GEs and *U. isabellina*-GEs by about 20 to 70 folds. Overall, most of the GEs

produced in this study, especially those of *C. echinulata*-GEs, had superior insecticidal activity.

Likewise, FAAs synthesized using GLA rich lipids produced by *C. echinulata*, displayed a

[14]

superior insecticidal activity against the same organism [36], suggesting that this FA is probably

a key molecule, responsible for the bioactivity of preparations.

Chortyk [65] reported that SEs are useful as effective, environmentally safe pesticides for

the control of soft-bodied arthropod pests. In addition, Puterka et al. [6] confirmed that the

majority of the SEs exhibited higher insecticidal activity than insecticide soap. The nature of

both the sugar and FA moieties determine the SE physicochemical properties, such as the

solubility in water and stability of emulsions, and their insecticidal activity. However, changing

the sugar or FA components from lower to higher carbon chains or whether the sugar is a

monosaccharide or disaccharide does not follow a consistent relationship to insecticide activity.

C18 FAs, such as oleic, elaidic, linoleic, and linoleic acids inhibited proliferation of malarial

parasites in mice infected with *Plasmodium vinckei petteri* or with *Plasmodium yoelii nigeriensis*

[66].

3.7. Quantitative analysis of ovarian cancer cell apoptosis induced by GEs

The ability of both FAMEs and GEs produced in this study to induce SKOV-3 cell apoptosis was

assessed by flow cytometry after Annexin FITC staining of cells (Fig. 9). The results show that

all GEs induced apoptosis of the SKOV-3 ovarian cancer cell line compared to untreated cells,

with the apoptotic rate increasing significantly after 48 h. A higher percentage of apoptosis was

observed in the cells treated with EPA-GEs (i.e. 43.1%), followed by *C. echinulata*-GEs, *U.*

isabellina-GEs and olive oil-GEs (i.e. 39.2, 34.0 and 33.5%, respectively). Similarly, FAAs

containing EPA in their structures in high percentages displayed a strong anticancer activity

against SKOV-3 ovarian cancer cell line [36].

Morin et al. [67] and Siena et al. [68] reported that, a variety of modified FAs are

promising molecules in the treatment of cancers. Also, there were several studies dealing with

anticancer, antimicrobial and anti-inflammatory activities of SE derivatives [69, 70]. Our results

correlated with those reported by An and Feng [71], whose evaluated the antitumor activity of

series of glucosyl esters derivatives against three cancer cells, human breast adenocarcinoma

(MCF-7), human colon carcinoma (K562) and human hepatoma (HepG2). They found that the

glucosyl esters exhibited significant anticancer activity in dose- and time-dependent fashion. The

structure - activity relationship analysis revealed that lipophilic properties might be an essential

parameter affecting their activity. Research to inhibit cancer cell proliferation has shown that SE

activity is linked to the nature of both sugar and fatty acyl chain [71].

4. Conclusions

Two immobilized lipases, especially lipase CA, efficiently catalyzed the synthesis of SEs using

glucose and FAMEs, derived from lipids of different origin including SCOs, as substrates. The

reaction of GE synthesis can be completed under environmentally friendly conditions using a

solvent mixture consisted of 80% DMSO and 20% tert-amyl alcohol in 24 h. The enzyme used in

the synthesis can be recycled at least three times without losing its catalytic activity. The

synthesized GEs displayed significant biological activities against important human pathogenic

microorganisms, the larvae of *A. aegypti* and the SKOV-3 ovarian cancer cell line, which are

related to their FA profile. Hence, we can conclude that SCOs, characterized by a wide diversity

in FA composition, can be considered as sources of acyl group-donors suitable for the production

of GEs of different bioactivity.

Author agreement

Hatim A. El-Baz, Ahmed M. Elazzazy, Tamer S. Saleh, Marianna Dourou, Jazem A. Mahyoub,

Mohammed N. Baeshen, Hekmat R. Madian and George Aggelis have all agreed to submission.

Declaration of Competing Interest

The authors declare that there are no conflicts of interest.

Acknowledgment

This work was funded by the *University of Jeddah*, Saudi Arabia, under grant No. (UJ-06-18-

ICP). The authors, therefore, acknowledge with thanks the University technical and financial

support.

References

1.   Furukawa, S., Akiyoshi, Y., O'Toole, G. A., Ogihara, H., & Morinaga, Y. (2010). Sugar

fatty acid esters inhibit biofilm formation by food-borne pathogenic bacteria. *International*

Journal of Food Microbiology, *138*(1–2), 176–180.

https://doi.org/10.1016/j.ijfoodmicro.2009.12.026

2.   Li, W., Das, S., Ng, K. Y., & Heng, P. W. S. (2011). Formulation, biological and

pharmacokinetic studies of sucrose ester-stabilized nanosuspensions of oleanolic acid.

Pharmaceutical Research, *28*(8), 2020–2033. https://doi.org/10.1007/s11095-011-0428-3

3.   Khan, N. R., & Rathod, V. K. (2015). Enzyme catalyzed synthesis of cosmetic esters and

its intensification: A review. *Process Biochemistry*, *50*(11), 1793–1806.

https://doi.org/10.1016/j.procbio.2015.07.014

4.   Neta, N. S., Teixeira, J. A., & Rodrigues, L. R. (2015). Sugar ester surfactants: Enzymatic

synthesis and applications in food industry. *Critical Reviews in Food Science and*

Nutrition. https://doi.org/10.1080/10408398.2012.667461

5.   Lucarini, S., Fagioli, L., Campana, R., Cole, H., Duranti, A., Baffone, W., … Casettari, L.

(2016). Unsaturated fatty acids lactose esters: cytotoxicity, permeability enhancement and

antimicrobial activity. *European journal of pharmaceutics and biopharmaceutics⬚:*

official journal of Arbeitsgemeinschaft fur Pharmazeutische Verfahrenstechnik e.V, *107*,

88–96. https://doi.org/10.1016/j.ejpb.2016.06.022

6.   Puterka, G. J., Farone, W., Palmer, T., & Barrington, A. (2003). Structure-function

relationships affecting the insecticidal and miticidal activity of sugar esters. *Journal of*

Economic Entomology. https://doi.org/10.1093/jee/96.3.636

7.   Zhao, K. H., Cai, Y. Z., Lin, X. S., Xiong, J., Halling, P. J., & Yang, Z. (2016). Enzymatic

synthesis of glucose-based fatty acid esters in bisolvent systems containing ionic liquids or

deep eutectic solvents. *Molecules*, *21*(10), 1–13.

https://doi.org/10.3390/molecules21101294

8.   de Lima, L. N., Mendes, A. A., Fernandez-Lafuente, R., Tardioli, P. W., & Camargo

Giordano, R. de L. (2018). Performance of different immobilized lipases in the syntheses

of short- and long-chain carboxylic acid esters by esterification reactions in organic

media. *Molecules*, *23*(4). https://doi.org/10.3390/molecules23040766

9.   Liang, M. Y., Chen, Y., Banwell, M. G., Wang, Y., & Lan, P. (2018). Enzymatic

preparation of a homologous series of long-chain 6- O -acylglucose esters and their

evaluation as emulsifiers. *Journal of Agricultural and Food Chemistry*.

https://doi.org/10.1021/acs.jafc.8b00913

10. Lin, X. S., Wen, Q., Huang, Z. L., Cai, Y. Z., Halling, P. J., & Yang, Z. (2015). Impacts of

ionic liquids on enzymatic synthesis of glucose laurate and optimization with superior

productivity by response surface methodology. *Process Biochemistry*.

https://doi.org/10.1016/j.procbio.2015.07.019

11. Ren, K., & Lamsal, B. P. (2017). Synthesis of some glucose-fatty acid esters by lipase

from *Candida antarctica* and their emulsion functions. *Food Chemistry*.

https://doi.org/10.1016/j.foodchem.2016.07.031

12. Mozaffarian, D., & Wu, J. H. Y. (2011). Omega-3 fatty acids and cardiovascular disease.

Journal of the American College of Cardiology, *58*(20), 2047–2067.

https://doi.org/10.1016/j.jacc.2011.06.063

13. Alakhras, R., Bellou, S., Fotaki, G., Stephanou, G., Demopoulos, N. A., Papanikolaou, S.,

& Aggelis, G. (2015). Fatty acid lithium salts from *Cunninghamella echinulata* have

cytotoxic and genotoxic effects on HL-60 human leukemia cells. *Engineering in Life*

Sciences, *15*(2), 243–253. https://doi.org/10.1002/elsc.201400208

14. Sayegh, F., Elazzazy, A., Bellou, S., Moustogianni, A., Elkady, A. I., Baeshen, M. N., &

Aggelis, G. (2016). Production of polyunsaturated single cell oils possessing antimicrobial

and anticancer properties. *Annals of Microbiology*, *66*(3), 937–948.

https://doi.org/10.1007/s13213-015-1176-0

15. Gumel, A. M., Annuar, M. S. M., Heidelberg, T., & Chisti, Y. (2011). Lipase mediated

synthesis of sugar fatty acid esters. *Process Biochemistry, 46*(11), 2079-2090.

https://doi.org/10.1016/j.procbio.2011.07.021

16. Plou, F. J., Cruces, M. A., Ferrer, M., Fuentes, G., Pastor, E., Bernabé, M., … Ballesteros,

A. (2002). Enzymatic acylation of di- and trisaccharides with fatty acids: choosing the

appropriate enzyme, support and solvent. *Journal of Biotechnology*, *96*(1), 55–66.

https://doi.org/10.1016/S0168-1656(02)00037-8

17. Jegannathan, K. R., & Nielsen, P. H. (2013). Environmental assessment of enzyme use in

industrial production-a literature review. *Journal of Cleaner Production*, *42*, 228–240.

https://doi.org/10.1016/j.jclepro.2012.11.005

18. Abdulmalek, E., Hamidon, N. F., & Abdul Rahman, M. B. (2016). Optimization and

characterization of lipase catalysed synthesis of xylose caproate ester in organic solvents.

Journal of Molecular Catalysis B: Enzymatic, *132*, 1-4.

https://doi.org/10.1016/j.molcatb.2016.06.010

19. Vescovi, V., dos Santos, J. B. C., & Tardioli, P. W. (2017). Porcine pancreatic lipase

hydrophobically adsorbed on octyl-silica: A robust biocatalyst for syntheses of xylose

fatty acid esters. *Biocatalysis and Biotransformation*, *35*(4), 298-305.

https://doi.org/10.1080/10242422.2017.1335717

20.   Findrik, Z., Megyeri, G., Gubicza, L., Bélafi-Bakó, K., Nemestóthy, N., & Sudar, M.

(2016). Lipase catalyzed synthesis of glucose palmitate in ionic liquid. *Journal of Cleaner*

Production, *112*, 1106-1111. https://doi.org/10.1016/j.clepro.2015.07.098

21.   Pöhnlein, M., Ulrich, J., Kirschhöfer, F., Nusser, M., Muhle-Goll, C., Kannengiesser, B.,

… Hausmann, R. (2015). Lipase-catalyzed synthesis of glucose-6-O-hexanoate in deep

eutectic solvents. *European Journal of Lipid Science and Technology*, *117*, 161-166.

https://doi.org/10.1002/ejlt.201400459

22.   Bell, J. G., Pratoomyot, J., Strachan, F., Henderson, R. J., Fontanillas, R., Hebard, A., …

Tocher, D. R. (2010). Growth, flesh adiposity and fatty acid composition of Atlantic

salmon (*Salmo salar*) families with contrasting flesh adiposity: Effects of replacement of

dietary fish oil with vegetable oils. *Aquaculture*, *306*, 225-232.

https://doi.org/10.1016/j.aquaculture.2010.05.021

23.   Eroldoğan, T., Turchini, G. M., Yılmaz, A. H., Taşbozan, O., Engin, K., Ölçülü, A., …

Mumoğullarında, P. (2012). Potential of cottonseed oil as fish oil replacer in European sea

bass feed formulation. *Turkish Journal of Fisheries and Aquatic Sciences*, *12*(4), 787-797.

https://doi.org/10.4194/1303-2712-v12_4_07

24.   Bellou, S., Baeshen, M. N., Elazzazy, A. M., Aggeli, D., Sayegh, F., & Aggelis, G.

(2014). Microalgal lipids biochemistry and biotechnological perspectives. *Biotechnology*

Advances, *32*(8), 1476–1493. https://doi.org/10.1016/j.biotechadv.2014.10.003

25.   Bellou, S., Triantaphyllidou, I. E., Aggeli, D., Elazzazy, A. M., Baeshen, M. N., &

Aggelis, G. (2016). Microbial oils as food additives: Recent approaches for improving

microbial oil production and its polyunsaturated fatty acid content. *Current Opinion in*

Biotechnology, *37*, 24–35. https://doi.org/10.1016/j.copbio.2015.09.005

26.   Kothri, M., Mavromati, M., Elazazy, A. M., Baeshen, M. N., Moussa, T. A. A., &

459   Aggelis, G. (2020). Microbial sources of polyunsaturated fatty acids (PUFAs) and the

460   prospect of organic residues and wastes as growth media for PUFA-producing

461   microorganisms. *FEMS Microbiology Letters*, *367*, fnaa028.

462   https://doi.org/10.1093/femsle/fnaa028

463   27.   Papanikolaou, S., & Aggelis, G. (2019). Sources of microbial oils with emphasis to

464   *Mortierella* (*Umbelopsis*) *isabellina* fungus. *World Journal of Microbiology and*

465   *Biotechnology*, *35*, 63. https://doi.org/10.1007/s11274-019-2631-z

466   28.   Athenaki, M., Gardeli, C., Diamantopoulou, P., Tchakouteu, S. S., Sarris, D.,

467   Philippoussis, A., & Papanikolaou, S. (2018). Lipids from yeasts and fungi: physiology,

468   production and analytical considerations. *Journal of Applied Microbiology*, *124*(2), 336–

469   367. https://doi.org/10.1111/jam.13633

470   29.   Bellou, S., & Aggelis, G. (2012). Biochemical activities in *Chlorella* sp. and

471   *Nannochloropsis salina* during lipid and sugar synthesis in a lab-scale open pond

472   simulating reactor. *Journal of Biotechnology*, *164*(2), 318–329.

473   https://doi.org/10.1016/j.jbiotec.2013.01.010

474   30.   Dourou, M., Tsolcha, O. N., Tekerlekopoulou, A. G., Bokas, D., & Aggelis, G. (2018).

475   Fish farm effluents are suitable growth media for *Nannochloropsis gaditana*, a

476   polyunsaturated fatty acid producing microalga. *Engineering in Life Sciences*, *18*(11),

477   851–860. https://doi.org/10.1002/elsc.201800064

478   31.   Campaña-Torres, A., Martínez-Córdova, L. R., Martínez-Porchas, M., López-Elías, J. A.,

479   & Porchas-Cornejo, M. A. (2012). Productive response of *Nannochloropsis oculata*,

cultured in different media and their efficiency as food for the rotifer brachionus

rotundiformis. *Phyton*, *81*, 45–50. ISSN 0031 9457

32. Rocha, J. M. S., Garcia, J. E. C., & Henriques, M. H. F. (2003). Growth aspects of the

marine microalga *Nannochloropsis gaditana. Biomolecular Engineering*, *20*(4–6), 237–

242. https://doi.org/10.1016/S1389-0344(03)00061-3

33. Ruan, Z., Hollinshead, W., Isaguirre, C., Tang, Y. J., Liao, W., & Liu, Y. (2015). Effects

of inhibitory compounds in lignocellulosic hydrolysates on *Mortierella isabellina* growth

and carbon utilization. *Bioresource Technology*, *183*, 18-24.

https://doi.org/10.1016/j.biortech.2015.02.026

34. Harde, S. M., Wang, Z., Horne, M., Zhu, J. Y., & Pan, X. (2016). Microbial lipid

production from SPORL-pretreated Douglas fir by *Mortierella isabellina. Fuel*, *175*, 64–

74. https://doi.org/10.1016/j.fuel.2016.02.023

35. Gardeli, C., Athenaki, M., Xenopoulos, E., Mallouchos, A., Koutinas, A. A., Aggelis, G.,

& Papanikolaou, S. (2017). Lipid production and characterization by *Mortierella*

(*Umbelopsis*) *isabellina* cultivated on lignocellulosic sugars. *Journal of Applied*

Microbiology, *123*(6), 1461–1477. https://doi.org/10.1111/jam.13587

36. El-Baz, H. A., Ahmed M. Elazzazy, Saleh, T. S., Dritsas, P., Mahyoub, J. A., Baeshen, M.

N., … Aggelis, G. (2020). Enzymatic synthesis of fatty acid amides using microbial lipids

as acyl group-donors and their biological activities. *bioRxiv*.

https://doi.org/10.1101/2020.09.13.295113

37. Perez, C., Pauli, M., & Bazerque, P. (1990). An antibiotic assay by the agar well diffusion

method. *Acta Biol Med Exp*. https://doi.org/10.5185/amlett.2015.5609

38. Wiegand, I., Hilpert, K., & Hancock, R. E. W. (2008). Agar and broth dilution methods to

determine the minimal inhibitory concentration (MIC) of antimicrobial substances. *Nature*

Protocols, *3*, 163–175. https://doi.org/10.1038/nprot.2007.521

39.   Finney, D. J. (1971). Probit analysis, 3rd edition, Cambridge University Press, Cambridge,

UK, pp 333.

40.   Ferrer, M., Soliveri, J., Plou, F. J., López-Cortés, N., Reyes-Duarte, D., Christensen, M.,

… Ballesteros, A. (2005). Synthesis of sugar esters in solvent mixtures by lipases from

Thermomyces lanuginosus and *Candida antarctica* B, and their antimicrobial properties.

Enzyme and Microbial Technology, *36*(4), 391–398.

https://doi.org/10.1016/j.enzmictec.2004.02.009

41.   Ferrer, M., Plou, F. J., Fuentes, G., Cruces, M. A., Andersen, L., Kirk, O., … Ballesteros,

A. (2002). Effect of the immobilization method of lipase from *Thermomyces lanuginosus*

on sucrose acylation. *Biocatalysis and Biotransformation*, *20*(1), 63–71.

https://doi.org/10.1080/10242420210153

42.   Habulin, M., Šabeder, S., & Knez, Ž. (2008). Enzymatic synthesis of sugar fatty acid

esters in organic solvent and in supercritical carbon dioxide and their antimicrobial

activity. *Journal of Supercritical Fluids*, *45*(3), 338-345.

https://doi.org/10.1016/j.supflu.2008.01.002

43.   Ganske, F., & Bornscheuer, U. T. (2005). Optimization of lipase-catalyzed glucose fatty

acid ester synthesis in a two-phase system containing ionic liquids and t-BuOH. *Journal of*

Molecular Catalysis B: Enzymatic, *36*(1–6), 40–42.

https://doi.org/10.1016/j.molcatb.2005.08.004

44.   Yu, J., Zhang, J., Zhao, A., & Ma, X. (2008). Study of glucose ester synthesis by

immobilized lipase from *Candida* sp. *Catalysis Communications*, *9*(6), 1369–1374.

https://doi.org/10.1016/j.catcom.2007.11.036

45.   Degn, P., & Zimmermann, W. (2001). Optimization of carbohydrate fatty acid ester

synthesis in organic media by a lipase from *Candida antarctica*. *Biotechnology and*

Bioengineering, *74*(6), 483-91. https://doi.org/10.1002/bit.1139

46.   H-Kittikun, A., Prasertsan, P., Zimmermann, W., Seesuriyachan, P., & Chaiyaso, T.

(2012). Sugar ester synthesis by thermostable lipase from *Streptomyces thermocarboxydus*

ME168. *Applied Biochemistry and Biotechnology*, *166*(8), 1969–1982.

https://doi.org/10.1007/s12010-012-9624-9

47.   Chaiyaso, T., H-kittikim, A., & Zimmermann, W. (2006). Biocatalytic acylation of

carbohydrates with fatty acids from palm fatty acid distillates. *Journal of Industrial*

Microbiology and Biotechnology, *33*(5), 338–342. https://doi.org/10.1007/s10295-005-

0073-0

48.   Davis, R. A., Lin, C. H., & Gervay-Hague, J. (2012). Chemoenzymatic synthesis of

cholesteryl-6-O-tetradecanoyl-α-d-glucopyranoside: A product of host cholesterol efflux

promoted by *Helicobacter pylori*. *Chemical Communications*, *48*, 9083-9085.

https://doi.org/10.1039/c2cc33948j

49.   Yang, Z., & Huang, Z. L. (2012). Enzymatic synthesis of sugar fatty acid esters in ionic

liquids. *Catalysis Science and Technology*, *2*(9), 1767–1775.

https://doi.org/10.1039/c2cy20109g

50.   Yan, Y., Bornscheuer, U. T., Stadler, G., Lutz-Wahl, S., Reuss, M., & Schmid, R. D.

(2001). Production of sugar fatty acid estrs by enzymatic esterification in a stirred-tank

membrane reactor: Optimization of parameters by response surface methodology. *Journal*

of the American Oil Chemists' Society, *78*(2), 147–153. https://doi.org/10.1007/s11746-

001-0235-x

51. Sebatini, A. M., Jain, M., Radha, P., Kiruthika, S., & Tamilarasan, K. (2016).

Immobilized lipase catalyzing glucose stearate synthesis and their surfactant properties

analysis. *3 Biotech*, *6*(2), 184. https://doi.org/10.1007/s13205-016-0501-z

52. Zhao, L., Zhang, H., Hao, T., & Li, S. (2015). In vitro antibacterial activities and

mechanism of sugar fatty acid esters against five food-related bacteria. *Food Chemistry*,

187, 370–377. https://doi.org/10.1016/j.foodchem.2015.04.108

53. Watanabe, T., Katayama, S., Matsubara, M., Honda, Y., & Kuwahara, M. (2000).

Antibacterial carbohydrate monoesters suppressing cell growth of *Streptococcus* mutans

in the presence of sucrose. *Current Microbiology*, *41*(3), 210–213.

https://doi.org/10.1007/s002840010121

54. Xiao, D., Ye, R., Davidson, P. M., Hayes, D. G., Golden, D. A., & Zhong, Q. (2011).

Sucrose monolaurate improves the efficacy of sodium hypochlorite against *Escherichia*

coli O157:H7 on spinach. *International Journal of Food Microbiology*, *145*(1), 64–68.

https://doi.org/10.1016/j.ijfoodmicro.2010.11.029

55. Devulapalle, K. S., Gómez de Segura, A., Ferrer, M., Alcalde, M., Mooser, G., & Plou, F.

J. (2004). Effect of carbohydrate fatty acid esters on *Streptococcus sobrinus* and

glucosyltransferase activity. *Carbohydrate Research*, *339*(6), 1029–1034.

https://doi.org/10.1016/j.carres.2004.01.007

56. Piao, J., Kawahara-Aoyama, Y., Inoue, T., & Adachi, S. (2006). Bacteriostatic activities

of monoacyl sugar alcohols against thermophilic sporeformers. *Bioscience, Biotechnology*

and Biochemistry, *70*(1), 263–265. https://doi.org/10.1271/bbb.70.263

57. de Lamo-Castellvi, S., Ratphitagsanti, W., Balasubramaniam, V. M., & Yousef, A. E.

(2010). Inactivation of *Bacillus amyloliquefaciens* spores by a combination of sucrose

laurate and pressure-assisted thermal processing. *Journal of Food Protection*, *73*(11),

2043–2052. https://doi.org/10.4315/0362-028X-73.11.2043

58. Shearer, A. E. H., Dunne, C. P., Sikes, A., & Hoover, D. G. (2000). Bacterial spore

inhibition and inactivation in foods by pressure, chemical preservatives, and mild heat.

Journal of Food Protection, *63*(11), 1503-1510. https://doi.org/10.4315/0362-028X-

63.11.1503

59. Wagh, A., Shen, S., Shen, F. A., Miller, C. D., & Walsha, M. K. (2012). Effect of lactose

monolaurate on pathogenic and nonpathogenic bacteria. *Applied and Environmental*

Microbiology, *78*(9), 3465–3468. https://doi.org/10.1128/AEM.07701-11

60. Karlová, T., Poláková, L., Šmidrkal, J., & Filip, V. (2010). Antimicrobial effects of fatty

acid fructose esters. *Czech Journal of Food Sciences*, *28*(2), 146–149.

https://doi.org/10.17221/37/2008-CJFS

61. Zhang, X., Wei, W., Cao, X., & Feng, F. (2015). Characterization of enzymatically

prepared sugar medium-chain fatty acid monoesters. *Journal of the Science of Food and*

Agriculture, *95*(8), 1631–1637. https://doi.org/10.1002/jsfa.6863

62. Wang, Y. J. (2004). Saccharides: Modifications and applications, in *Chemical and*

Functional Properties of Food Saccharides (Tomasik, P., ed.), Boca Raton, pp. 35-46.

63. Powell, J. R. (2018). Perspective piece mosquito-borne human viral diseases: Why *Aedes*

aegypti? *American Journal of Tropical Medicine and Hygiene*, *98*(6), 1563–1565.

https://doi.org/10.4269/ajtmh.17-0866

64. Rodrigues, A. M., Sampaio, C. de G., de Souza, J. S. N., Campos, A. R., da Silva, A. B.

R., de Morais, S. M., & Martins, V. E. P. (2019). Different susceptibilities of *Aedes*

[27]

aegypti and *Aedes albopictus* larvae to plant-derived products. *Revista da Sociedade*

Brasileira de Medicina Tropical, *52*, e20180197. https://doi.org/10.1590/0037-8682-

0197-2018

65. Chortyk, O. T. (1996). Chemically synthesized sugar esters for the control of soft bodied

arthropods. *World Intellectual Property Organization*. New York, USA: ACM Press.

https://doi.org/10.1145/634067.634234

66. Krugliak, M., Deharo, E., Shalmiev, G., Sauvain, M., Moretti, C., & Ginsburg, H. (1995).

Antimalarial effects of C18 fatty acids on *Plasmodium falciparum* in culture and on

Plasmodium vinckei petteri and *Plasmodium yoelii nigeriensis* in vivo. *Experimental*

Parasitology, *81*(1), 97–105. https://doi.org/10.1006/expr.1995.1097

67. Morin, C., Rousseau, É., & Fortin, S. (2013). Anti-proliferative effects of a new

docosapentaenoic acid monoacylglyceride in colorectal carcinoma cells. *Prostaglandins*

Leukotrienes and Essential Fatty Acids, *89*(4), 203–213.

https://doi.org/10.1016/j.plefa.2013.07.004

68. Siena, L., Cipollina, C., Di Vincenzo, S., Ferraro, M., Bruno, A., Gjomarkaj, M., & Pace,

E. (2018). Electrophilic derivatives of omega-3 fatty acids counteract lung cancer cell

growth. *Cancer Chemotherapy and Pharmacology*, *81*(4), 705-716.

https://doi.org/10.1007/s00280-018-3538-3

69. Campana, R., Merli, A., Verboni, M., Biondo, F., Favi, G., Duranti, A., & Lucarini, S.

(2019). Synthesis and evaluation of saccharide-based aliphatic and aromatic esters as

antimicrobial and antibiofilm agents. *Pharmaceuticals*, *12*(4), 186.

https://doi.org/10.3390/ph12040186

70. Petrova, K. T., Barros, M. T., Calhelha, R. C., Soković, M., & Ferreira, I. C. F. R. (2018).

Antimicrobial and cytotoxic activities of short carbon chain unsaturated sucrose esters. *Medicinal Chemistry Research*, *27*(3), 980–988. https://doi.org/10.1007/s00044-017-2121-5

71.   An, D., & Feng, D. (2019). Enzymic synthesis, physicochemical, and cell activity of glucosyl ester derivatives based on N-fatty acyl amino acid. *Chemical Papers*, *73*(3), 653–662. https://doi.org/10.1007/s11696-018-0614-4

625   Table 1

626   Comparison of GE synthesis using two immobilized lipases and olive oil derived FFAs or

627   FAMEs as substrates at different molar ratio of the reactants.

Entry	Substrate	Immobilized enzyme	Olive oil: Glucose Ratio	Product	Conversion (%)
1	FFAs	Lipase CA	1:1	Still contain FFAs	50.5 ± 2.5
2			1:2	Still contain FFAs	70.7 ± 3.5
3			1:3	Almost 100% ester	95.2 ± 4.8
4	FAMEs		1:1	Still contain FAMEs	70.1 ± 3.5
5			1:2	Still contain FAMEs	96.1 ± 3.5
6			1:3	100% ester	100.0 ± 5.0
7	FFAs	Lipase CR	1:1	Still contain FFAs	30 ± 1.5
8			1:2	Still contain FFAs	63 ± 3.2
9			1:3	Still contain FFAs	89 ± 4.5
10	FAMEs		1:1	Still contain FAMEs	64 ± 3.2
11			1:2	Still contain FAMEs	87 ± 4.4
12			1:3	Still contain FAMEs	92 ± 4.6

630   Table 2

631   GE synthesis using olive oil derived FAMEs (at a ratio FAMEs:Glc 1:3), utilizing lipase CA as

632   catalyst at different concentrations, in a solvent mixture consisted of 80% DMSO and 20% tert-

633   amyl alcohol and for a duration of incubation 50 h.

Entry	Lipase CA (g/25 mL)	Conversion (%)
1	0.10	80.3 ± 4.0
2	0.15	92.1 ± 4.6
3	0.20	96.3 ± 4.8
4	0.25	100.3 ± 5.0
5	0.30	100.2 ± 5.0

640   Table 3

641   GE synthesis by lipase CA in a solvent mixture consisted of 80% DMSO and 20% tert-amyl

642   alcohol) utilizing FAMEs of different origin as substrate.

Entry	Source of FAMEs	Time (h)	Conversion (%)
1	*Cunninghamella echinulata*	24	80.3 ± 4.0
2	*Umbelopsis isabellina*	18	85.1 ± 4.3
3	*Nannochloropsis gaditana*	18	86.3 ± 4.3
4	EPA concentrate	18	99.1 ± 5.0

644 Table 4

645 GEs antimicrobial activity using human pathogens as tested organisms. Data represent the mean

646 of three replicates ± SD of the diameter of the inhibition zones.

	GEs synthesized using FAMEs derived from:				
	C. echinulata	*U. isabellina*	*N. gaditana*	Olive oil	EPA concentrate
	Inhibition zone (mm)				
Escherichia coli (ATCC 25922)	9.4 ± 0.3	0.0	10.0 ± 0.4	11.5 ± 0.4	11.1 ± 0.0
Klebsiella pneumoniae (ATCC 700603)	10.5 ± 0.5	0.0	11.1 ± 0.4	10.3 ± 0.6	14.2 ± 0.1
Salmonella typhimurium (ATCC 14028)	5.6 ± 0.5	5.8 ± 0.7	6.1 ± 0.3	10.1 ± 0.7	15.0 ± 0.0
Pseudomonas aeruginosa (ATCC 15442)	6.8 ± 0.8	5.5 ± 0.8	10.5 ± 0.5	10.1 ± 0.4	13.2 ± 0.0
Bacillus subtilis (ATCC 6633)	14.1 ± 0.5	0.00	9.0 ± 0.0	9.8 ± 0.7	17.0 ± 0.5
MRSA *Staphylococcus aureus* (ATCC 4330)	11.2 ± 0.1	5.8 ± 1.0	8.5 ± 0.5	10.1 ± 0.2	16.0 ± 0.1
Staphylococcus aureus (ATCC 25923)	12.3 ± 0.2	0.0	8.6 ± 0.3	12.0 ± 0.6	17.0 ± 0.2
Candida albicans (ATCC 10231)	14.3 ± 0.0	13.3 ± 0.0	14.00 ± 0.0	15.00 ± 0.3	20.0 ± 0.1

648   Table 5

649   Minimum inhibitory concentration (MIC, µg/mL) and minimum bactericidal concentration

650   (MBC, µg/mL) of the synthesized GEs against pathogens.

Test organisms	Source of FAMEs							
	C. echinulata		*N. gaditana*		Olive oil		EPA concentrate oil	
	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC
Klebsilla pneumoniae (ATCC 700603)	25.0 ± 0.0	100.0 ± 0.0	50.0 ± 0.0	100.0 ± 0.0	66.7 ± 28.8	83.3 ± 28.8	33.3 ± 14.4	100.0 ± 0.0
Pseudomonas aeruginosa (ATCC 15442)	41.7 ± 14.4	100.0 ± 0.0	50.0 ± 0.0	100.0 ± 0.0	25.0 ± 0.0	83.3 ± 28.8	20.8 ± 7.2	41.7 ± 14.4
Bacillus subtilis (ATCC 6633)	16.7 ± 7.3	50.0 ± 0.0	20.8 ± 7.3	100.0 ± 0.0	33.3 ± 14.4	100.0 ± 0.0	10.4 ± 3.6	41.7 ± 14.4
Staphylococcus aureus (ATCC 25923)	12.5 ± 0.0	50.0 ± 0.0	50.0 ± 0.0	83.3 ± 28.8	83.3 ± 28.7	100.0 ± 0.0	8.3 ± 3.6	33.3 ± 14.4

652   Table 6

653   Susceptibility of *Aedes aegypti* larvae to GEs under laboratory conditions by using dipping

654   methods. Data represent the mean values of six replicates.

Source of GEs	LC50 (mg/L)	Lower limit	Upper limit	RR
Cunninghamella echinulata	0.54	0.47	0.64	1.00
Umbelopsis isabellina	39.62	33.75	46.10	73.23
Nannochloropsis gaditana	16.92	12.60	25.82	31.27
Olive oil	12.88	10.08	17.54	23.81
EPA concentrate	10.24	7.84	14.00	18.93

Figure captions

Fig. 1

Enzyme-catalyzed synthesis of glucose esters using free fatty acids (FFAs) or fatty acid methyl

esters (FAMEs) as substrates.

Fig. 2

Reaction course over time of olive oil FAMEs conversion to glucose esters in different olive oil

FAMEs:glucose ratios.

Fig. 3

NMR monitoring of glucose ester synthesis. The signal intensity at 4.17 and 4.41 ppm derived

from the CH_2OCO group increased simultaneously with a decrease of the signal intensity derived

from the glucose protons of C6 at 3.39 and 3.52 ppm.

Fig. 4

Identification of the ester produced as Glc (C-6)-OCOR by 2D (^{1}H-^{13}C HMBC) NMR (a) and

correlation between the protons of C-6 glucose ester and the carbonyl function in olive oil

FAMEs (b).

Fig. 5

FT-IR of olive oil FAMEs and their glucose esters.

Fig. 6

Reaction course over time of conversion of FAMEs derived from different origins to glucose

esters.

Fig.7

FT-IR analysis of EPA concentrate FAMEs and their glucose esters.

Fig. 8

The larval mortality effect of GEs derived from *C. echinulata*, EPA concentrate, olive oil, *U.*

isabellina and *N. gaditana* at different concentrations against *Aedes aegypti* after continuous

exposure for 48 h.

Fig. 9

Effect of FAMEs and GEs on SKOV-3 cancer cell line. Flow cytometry analysis of apoptosis in

SKOV-3 cells either untreated or treated with 10 µg/mL of every compound for 48 h. After the

treatment period, the cells were stained with Annexin FITC and subsequently analyzed by flow

cytometry.

Abbreviations: *C. e.*, *C. echinulata*; *U. i., U. isabellina*; *N. g.*, *N. gaditana*; O. o., Olive oil; EPA,

EPA concentrate

time = 50 h

time = 20 h

(a)

(b)

FT-IR of olive oil methyl ester

2853

2923

1743 cm⁻¹ (RCOO)

FT-IR of olive oil-GE

3380 cm⁻¹ (OH)

FT-IR of EPA concentrate methyl ester

3012

2853

2923

1741 cm^{-1} (RCOO)

FT-IR of EPA concentrate-GE

3416 cm^{-1} (OH)