Circadian key component CLOCK/BMAL1 interferes with segmentation clock in mouse
embryonic organoids

Authors: Yasuhiro Umemura[a], Nobuya Koike[a], Yoshiki Tsuchiya[a], Hitomi Watanabe[b], Gen
Kondoh[b], Ryoichiro Kageyama[c], and Kazuhiro Yagita[a,*].

Affiliations:

[a]Department of Physiology and Systems Bioscience, Kyoto Prefectural University of Medicine,
Kawaramachi-Hirokoji, Kyoto 602-8566, Japan.

[b]Laboratory of Integrative Biological Science, Institute for Frontier Life and Medical Sciences,
Kyoto University, Kyoto 606-8501, Japan.

[c]Laboratory of Growth Regulation System, Institute for Frontier Life and Medical Sciences, Kyoto
University, Kyoto 606-8507, Japan.

*Correspondence to: Kazuhiro Yagita

Email: kyagita@koto.kpu-m.ac.jp

Author Contributions: Y.U. and K.Y. designed the research; Y.U., Y.T., H.W., G.K., and K.Y.
performed the research; Y.U., N.K., Y.T., R.K., and K.Y. analyzed the data; Y.U., N.K., and K.Y.
wrote the paper

Competing Interest Statement: The authors declare no competing interests

Abstract

In mammals, circadian clocks are strictly suppressed during early embryonic stages as well as pluripotent stem cells, by the lack of CLOCK/BMAL1 mediated circadian feedback loops. During ontogenesis, the innate circadian clocks emerge gradually at a late developmental stage, then, with which the circadian temporal order is invested in each cell level throughout a body. Meanwhile, in the early developmental stage, a segmented body plan is essential for an intact developmental process and somitogenesis is controlled by another cell-autonomous oscillator, the segmentation clock, in the posterior presomitic mesoderm (PSM). In the present study, focusing upon the interaction between circadian key components and the segmentation clock, we investigated the effect of the CLOCK/BMAL1 on the segmentation clock *Hes7* oscillation, revealing that the expression of functional CLOCK/BMAL1 severely interferes with the ultradian rhythm of segmentation clock in induced PSM and gastruloids. RNA sequencing analysis showed that the premature expression of CLOCK/BMAL1 affects the *Hes7* transcription and its regulatory pathways. These results suggest that the suppression of CLOCK/BMAL1-mediated transcriptional regulation during the somitogenesis may be inevitable for intact mammalian development.

Introduction

The circadian clock is the cell-autonomous time-keeping system generating the orderly regulated various physiological functions, which enables cells, organs, and systems to adapt to the cyclic environment of the rotating Earth (1-5). The core architecture of the circadian molecular clock consists of negative transcriptional/translational feedback loops (TTFLs) composed of a set of circadian clock genes, including *Bmal1*, *Clock*, *Period* (*Per1*, *2*, *3*), and *Cryptochrome* (*Cry1*, *2*), functioning under the control of E-box elements (2, 6). The kernel of TTFLs is composed of heterodimerized CLOCK/BMAL1 key transcriptional factors that positively regulate the circadian output genes, as well as *Per* and *Cry* genes via E-box. PERs and CRYs inhibit CLOCK/BMAL1 transcriptional activity, and the negative feedback loops between these genes generate oscillations of approximately 24 hr.

In mammalian development, it has been demonstrated that early embryos and pluripotent stem cells have no apparent circadian molecular oscillations (7-11), whereas the innate circadian clock develops during ontogenesis and is established at a late developmental stage (12-15). Regarding the mechanisms regulating circadian clock development, using an *in vitro* model of embryonic stem cell (ESC) differentiation and mouse embryos, it was shown that prolonged posttranscriptional mechanisms, such as suppressed translation of CLOCK protein and

predominant cytoplasmic localization of PER proteins, inhibit the establishment of the circadian TTFL cycle (16-18). Although it was revealed that the multiple mechanisms strictly suppress the circadian molecular clock in the undifferentiated cells and early stage embryos, the biological and physiological significance of the delayed emergence of circadian clock oscillation in mammalian embryos has been unknown.

In the early developmental stages, a segmented body plan is essential for an intact developmental process. Somitogenesis is related to another cell-autonomous oscillator, the segmentation clock, in the posterior presomitic mesoderm (PSM) (19, 20). The mouse segmentation clock is underlain by a negative feedback loop involving *Hes7* oscillation (21, 22). HES7 is a key transcriptional factor that represses its own expression and oscillates through a negative feedback loop in a period of 2–3 hr in mouse and 4–5 hr in humans. The NOTCH, WNT, and fibroblast growth factor signaling pathways are involved in the regulation of the *Hes7* oscillator and its intercellular synchronization (20). In mammals, two different types of rhythm sequentially emerge during the developmental process, however, there is a lack of knowledge about the biological significance of the rhythm conversion during development.

In this study, focusing on the relationship between circadian clock and segmentation clock, we investigated the effect of the premature expression of CLOCK/BMAL1 on the segmentation clock oscillation, and revealed severe interference with the ultradian rhythm of segmentation clock in iPSM and gastruloids. RNA sequencing analysis showed that CLOCK/BMAL1 affects the *Hes7* transcription and its regulatory pathways. These findings highlight that the suppression of functional CLOCK/BMAL1, which leads to arrest the circadian clock oscillation, during the early to mid-developmental stage may be inevitable for the intact process of mammalian embryogenesis.

Results

A circadian clock gene, *Per1*, and a segmentation clock gene, *Hes7*, are adjacent genes in the mammalian genome

In mammals, the temporal relationship between the segmentation clock and circadian clock appears to be mutually exclusive (**Fig. S1**) (12-15, 19, 20). To explore the functional interaction between these two biological rhythms with different frequencies during the developmental process,

we focused on the genomic architecture of genes comprising circadian and segmentation clocks,

respectively. Intriguingly, one of the core circadian clock genes, *Per1*, is physically adjacent to an

essential component of the segmentation clock, *Hes7*, in a genomic region conserved in higher

vertebrates, including mice and humans. The *Per1* homolog *Per2* is adjacent to the *Hes7* homolog

Hes6 in the genome (**Fig. 1A**). Since *Hes7* exhibits the essential characteristics of a segmentation

clock (23) and neighboring genes can influence the expressions with each other during

somitogenesis in zebrafish (24), we focused on the effect of the regulation mechanism of the

circadian clock on segmentation clock oscillation. Therefore, we investigated the effect of the

CLOCK/BMAL1-mediated activation of *Per1* transcription on the segmentation clock oscillation

in induced presomitic mesoderm (iPSM), an *in vitro* recapitulating model of a segmentation clock,

using ESCs carrying the *Hes7*-promoter-driven luciferase reporter (*pHes7-luc*) (25) (**Fig. 1B**). In

the iPSM, the pluripotent markers have not yet been down-regulated sufficiently as previously

reported (25), and the iPSM differentiated from *Per2^Luc* ESCs showed no apparent circadian clock

oscillation (**Fig. S2A and B**). In addition, the immunostaining pattern of CLOCK, BMAL1, and

PER1 in the iPSMs was quite similar to that in the undifferentiated ESCs (**Fig. S2C**) (16, 17),

confirming that the circadian TTFL was not established and circadian clock oscillation was also

strictly suppressed in the iPSM by the common inhibitory molecular mechanisms to the

undifferentiated ESCs.

Because CLOCK/BMAL1 is key transcription regulator of circadian TTFL, and the

expression of CLOCK protein is suppressed post-transcriptionally in iPSM as well as ESCs and

early embryos (17), we established two ESC lines carrying both the doxycycline (dox)-inducible

Clock and *Bmal1* genes (**Fig. 1C**, **Fig. S3**). In iPSM differentiated from ESCs, the expression of

both *Clock/Bmal1* mRNA and CLOCK/BMAL1 proteins was confirmed after the addition of dox

(**Fig. 1 D and E**), and we found that overexpression of both *Clock* and *Bmal1* successfully

activated the expression of core clock genes (**Fig. 2A**). As the dominant negative mutant of *Bmal1*

(*Bmal1DN*) (26) co-expressed with *Clock* did not activate the *Per1/2* and *Cry1/2* genes, we

concluded that CLOCK/BMAL1 specifically activated the expression of these clock genes via an

E-box (**Fig. 1 C–E, 2A**). We then examined the expression of genes in *Hes7*, which is proximal to

Per1, and *Hes6*, which is proximal to *Per2* in the genome. The expression of *Clock/Bmal1* induced

by dox in the iPSM induced significant upregulation of the expression of the *Hes7*, but not *Hes6*,

gene (**Fig. 2B**). Similarly, we also observed the upregulation of *Hes7* expression by *Clock/Bmal1*

induction in the undifferentiated ESCs (**Fig. S4 A and B**). These results indicate that the circadian

components CLOCK/BMAL1 also affect the segmentation clock gene *Hes7*, as well as *Per1*.

Inhibition of *Hes7* ultradian rhythm by CLOCK/BMAL1 in iPSM

We next performed a functional analysis using the *in vitro* recapitulation model of a segmentation clock oscillation in iPSM (25). The oscillations in bioluminescence from *pHes7-luc* reporters were observed using a photomultiplier tube device (PMT) and an EM-CCD camera (**Fig. 3A**). We confirmed an oscillation of *Hes7*-promoter-driven bioluminescence with a period of approximately 2.5–3 hr in control iPSM with or without dox using PMT and the EM-CCD camera (**Fig. 3 B and C**). Traveling waves of *pHes7-luc* bioluminescence were observed, indicating that the segmentation clock oscillation in iPSM was successfully recapitulated, consistent with a previous report (25). Using this iPSM-based segmentation clock system, we investigated the effect of *Clock/Bmal1* expression on *Hes7*-promoter-driven oscillation. The expression of *Clock/Bmal1* genes (Dox+) resulted in defects of the oscillation in *Hes7* promoter activity, whereas *pHes7-luc* bioluminescence continued to oscillate under Dox– conditions. Oscillation of the segmentation clock was observed even during the induction of *Clock/Bmal1DN* (**Fig. 3D**), indicating that the CLOCK/BMAL1-mediated mechanism interfered with the transcriptional oscillation of *Hes7*. A traveling wave of *Hes7* promoter activity disappeared with the expression of *Clock/Bmal1* (**Fig. 3 E and F**), and dox-dependent arrest of *pHes7-luc* traveling wave (**Fig. 3 G and H**) clearly demonstrated the CLOCK/BMAL1-mediated interference with *Hes7*-driven segmentation clock oscillation in iPSM.

Interference with somitogenesis-like segmentation by induction of CLOCK/BMAL1 in gastruloids

In addition, to explore the effect of CLOCK/BMAL1 expression on somitogenesis, we established the ESC-derived embryonic organoids, gastruloids, recapitulating an embryo-like organization, including somitogenesis-like process *in vitro* (27) (**Fig. 4A**). The *pHes7-luc* bioluminescence represented a traveling wave accompanied by the formation of segment-like structures with anteroposterior polarity, in which the gastruloids were stained with stripes of a somite marker, *Uncx4.1*, by *in situ* hybridization (**Fig. 4 B–D**). Only dox treatment in control gastruloids induced no change in the *pHes7-luc* bioluminescence oscillation and somitogenesis-like process (**Fig. 4 E–G**). The dox-inducible *Clock/Bmal1* ESC line carrying *pHes7-luc* was differentiated *in vitro* into gastruloids and produced the somitogenesis-like process without dox (**Fig. 4 H–J**). In contrast, the dox-dependent induction of *Clock/Bmal1* expression in the gastruloids interrupted the *pHes7-luc* oscillation and disrupted the somitogenesis-like structures (**Fig. 4 K–M**). In gastruloids, the expression of both *Clock/Bmal1* mRNA was confirmed after the

150 addition of dox (**Fig. S5**). These results suggest that the premature expression of the circadian key
151 transcriptional regulator CLOCK/BMAL1 critically interferes with not only *Hes7* oscillation, but
152 also somitogenesis.

CLOCK/BMAL1-mediated interference in *Hes7* regulatory network

Next, to examine the perturbation mechanisms of the segmentation clock oscillation by the
circadian components CLOCK/BMAL1, we analyzed the RNA sequencing (RNA-seq) data
obtained from the total RNA of iPSM colonies. We extracted 509 upregulated and 88
downregulated differentially expressed genes (DEGs) after the induction of *Clock/Bmal1* gene
expression in iPSM colonies (**Fig. 5A**). A KEGG pathway enrichment analysis for the DEGs
revealed enrichment of the WNT, MAPK, and NOTCH signaling pathways related to *Hes7*
oscillation (28) (**Fig. 5B**). Almost all other ranked pathways also included the WNT, MAPK, and
NOTCH signaling pathway-related genes (**Fig. 5B**). Similarly, enrichment of the WNT, MAPK,
and NOTCH signaling pathways by *Clock/Bmal1* induction was also observed in the
undifferentiated ESCs (**Fig. S6 A and B**). These findings indicate that the expression of
CLOCK/BMAL1 affects the *Hes7*-related signaling pathways, which interferes with the feedback
loop regulating *Hes7* oscillation. Intriguingly, in addition to *Hes7* gene expression, the expressions
of *Aloxe3* in iPSM and *Aloxe3* and *Vamp2* in ESCs, the other contiguous genes with *Per1*, were
upregulated with the induction of *Clock/Bmal1* expression, and this result was confirmed by
quantitative RT-PCR (qPCR) (**Fig. 5 C–E**, **Fig. S6 C–E**) suggesting that forced expression of
CLOCK/BMAL1 also affects a wide region around the *Hes7* gene locus on the same chromosome.
These results suggest that the premature expression of the circadian components CLOCK/BMAL1
interfered with *Hes7* oscillation and somitogenesis by perturbing the *Hes7* expressions through
indirect regulatory pathways (**Fig. 5F**). Because the loss of the *Hes7* ultradian expression rhythm
in the mouse cause segmentation defects (22, 29), the oscillatory expression of *Hes7* is essential
for mammalian development. Therefore, the results in this study suggest that it may be imperative
that CLOCK/BMAL1 function is suppressed until the completion of segmentation and other
related developmental events.

Discussion

Our present study showed that premature expression of circadian key components
CLOCK/BMAL1 severely interferes with the ultradian rhythm of the segmentation clock in iPSM
and gastruloids.

We have previously reported that during the early to mid-developmental stage, there are
multiple molecular mechanisms that underlie the strict suppression of circadian TTFLs, such as
the post-transcriptional suppression of CLOCK protein (17, 18) and the exclusive cytoplasmic
localization of PER proteins (16). Furthermore, we have also reported that the maternal circadian
clock cannot entrain the fetus until the establishment of the fetal circadian clock itself (17). These
results suggest that the circadian rhythm in mammalian embryos is rigorously suppressed by the
multilayered inhibitory mechanisms during the early to mid-developmental stage. During the
multilayered suppression of circadian clock oscillation, the ultradian temporal oscillation of *Hes7*
expression, segmentation clock, proceeds and forms the spatial repetitive structure of somites.

In the present study, we investigated the effect of the CLOCK/BMAL1-mediated activation
of *Per1* transcription on the segmentation clock oscillation by using the iPSM differentiated from
ESCs. It was suggested that, similar to the undifferentiated ESCs, circadian clock oscillation is
suppressed in the iPSM by the common mechanisms to the ESCs and early embryos (**see Fig. S2**).
Recently, it was reported that hundreds of genes including *Per1* also oscillates in the same phase
as *Hes7* ultradian rhythm in *in vitro*-PSM of both mouse and humans (30), suggesting that *Per1* is
deviated and free from the circadian gene regulatory mechanism of TTFL. These findings are
consistent with the previously reported observations indicating that the multilayered inhibitory
mechanisms including post-transcriptional inhibition of CLOCK and the predominant cytoplasmic
accumulation of PER1 do not allow the oscillation of circadian TTFL (17, 18). Interestingly,
although the expression of BMAL1 protein was observed even in ESCs (17), the dox-induced
CLOCK sole expression in ESCs resulted in the only partial upregulation of E-box driven circadian
clock genes (**Fig. S3**), raising the possibility that the endogenously expressed BMAL1 might be
post-translationally modified to not function. Therefore, in this study, we used ESC lines carrying
both the dox-inducible *Clock* and *Bmal1* genes as a model system of premature expression of
CLOCK/BMAL1 (**see Fig. 1C**).

We demonstrated that the expression of CLOCK/BMAL1 affected the WNT, MAPK, and
NOTCH signaling pathways related to *Hes7* oscillation in iPSM (**see Fig. 5 A and B**). In addition,
the premature expression of CLOCK/BMAL1 resulted in not only the up-regulation of *Per1*
expression but also the expressions of *Hes7, Aloxe3*, and *Vamp2*, localized adjacently on the *Per1*
genomic locus (**see Fig. 2 A and B, Fig. 5 C–E, Fig. S6 C–E**). In the iPSM, the up-regulation of
these gene expressions has already been induced after the 2-hour dox treatment (**see Fig. 2 A and**

B, Fig. 5 C–E). Considering that the *Per1* promoter harbors E-box elements with which CLOCK/BMAL1 heterodimer has a much higher affinity than the other genomic region (31), the immediate up-regulation of genes near *Per1* gene locus after the induction of CLOCK/BMAL1 expressions could be caused by the ripple effect (32). On the other hand, the bioluminescence from *Hes7*-promoter driven luciferase reporters in the iPSMs not only lost cycling but also decreased signal intensity in approximately 2 hours after the dox addition (**see Fig. 3D**), indicating that the expression of CLOCK/BMAL1 in the iPSM has also inhibitory effects on the *Hes7* gene expressions. Among components involved in the *Hes7*-regulatory signaling pathways, expression of CLOCK/BMAL1 induced some negative regulators, such as the *Dusp* phosphatase family (33) in the MAPK signaling pathway, *Sfrp* in the WNT signaling pathway (34), and *Lfng* in the NOTCH signaling pathway (35) (**see Fig. 5B**). Therefore, the premature expression of CLOCK/BMAL1 first may upregulate *Hes7* transcription and induce subsequent downregulation of *Hes7* gene expression by the induction of the negative regulators in addition to the HES7 autoinhibition. Consequently, the premature expression of the circadian components CLOCK/BMAL1 interfered with *Hes7* oscillation by perturbing the *Hes7* expression through various pathways.

In this study, we used a mouse embryonic organoid, gastruloids, as an *in vitro* recapitulation model of somitogenesis-like process (27). The premature expression of CLOCK/BMAL1 in the gastruloids disrupted not only the *Hes7* oscillation but also the striped structure of the somite marker, *Uncx4.1* (**see Fig. 4M**). Because the RNA-seq analysis data showed that hundreds of genes were affected by the induction of CLOCK/BMAL1 (**see Fig. 5A**), the possibility cannot be denied that the premature expression of CLOCK/BMAL1 affects cell fates or characters. However, the posterior structure in the gastruloids was held even after the induction of CLOCK/BMAL1 and then continued to extend, concomitant with the decrease of *Hes7* bioluminescence signals and the arrest of the *Hes7* oscillation (**see Fig. 4K**), suggesting that the premature expression of CLOCK/BMAL1 interfered with the somitogenesis process by perturbing *Hes7* oscillation of the segmentation clock.

In vitro recapitulation of embryonic process using iPSM and gastruloids has differences such as no brain tissues comparing with in vivo process. However, key regulators of somitogenesis we focus on in this study are expressed similarly between embryos and gastruloids using single-cell RNA sequencing and spatial transcriptomics (27), and the in vitro recapitulation model enables to analyze the *Hes7* oscillation in more detail using real-time imaging without maternal effects.

Our findings shown in this study indicated that the CLOCK/BMAL1, key components regulating the circadian TTFL, affected and interfered with the segmentation clock. Considering that transcriptional activation of CLOCK/BMAL1 is essential for the circadian regulatory

248 networks, these results suggest that the strict suppression of circadian molecular oscillatory
249 mechanisms during the early stage embryos is inevitable for the intact developmental process in
250 mammals. Therefore, this may be the biological and physiological significance of the delayed
251 emergence of circadian clock oscillation and the rhythm conversion observed in mammalian
252 development.

Materials and Methods

Cell culture

KY1.1 ESCs (7), referred to as ESC in the text, and *Per2^Luc* ESCs (5, 36) were maintained as
described previously (17). E14TG2a ESCs carrying *Hes7*-promoter-driven luciferase reporters
(25), referred to as *pHes7-luc* ESCs in the text, were maintained without feeder cells in DMEM
(Nacalai) supplemented with 15% fetal bovine serum (Hyclone), 2 mM L-glutamine (Nacalai), 1
mM nonessential amino acids (Nacalai), 100 µM StemSure® 2-mercaptoethanol solution (Wako),
1 mM sodium pyruvate (Nacalai), 100 units/mL penicillin and streptomycin (Nacalai), 1000
units/mL leukemia inhibitory factor (Wako), 3 µM CHIRON99021 (Wako or Tocris Biosciences),
and 1 µM PD0325901 (Wako) with 5% CO_2 at 37°C.

Transfection and establishment of cell lines

ESCs stably expressing dox-inducible *Clock/Bmal1* or *Clock/Bmal1DN* (I584X) were
established as described previously (17). For *TetO-Clock/Bmal1* or *TetO-Clock/Bmal1DN* ESCs,
KY1.1 ESCs or *pHes7-luc* ESCs were transfected using 10.5 µl of FuGENE 6 mixed with 1 µg of
pCAG-PBase, 1 µg of PB-TET-Clock (17), 1 µg of PB-TET-Bmal1 or PB-TET-Bmal1DN
(I584X), 1 µg of PB-CAG-rtTA Adv, and 0.5 µg of puromycin selection vector. The transfected
cells were grown in a culture medium supplemented with 2 µg/mL puromycin for two days. The
ESC colonies were picked and checked by qPCR after treatment with 500 ng/mL dox. For PB-
TET-Bmal1 and PB-TET-Bmal1DN (I584X), Bmal1 cDNA and Bmal1DN (I584X) cDNA (26)
were cloned into a PB-TET vector (37). For the *TetO-Clock Per2^Luc* ESCs, *Per2^Luc* ESCs were
established as described previously (17).

Bioluminescence imaging

The iPSM colonies were differentiated from the *pHes7-luc* ESCs and *Per2^{Luc}* ESCs as described previously (25). The *Per2^{Luc}* ESCs were cultured without feeder cells in the ES medium containing 3 µM CHIRON99021 and 1 µM PD0325901 before in vitro differentiation. Bioluminescence imaging of single *pHes7-luc* iPSM colonies and *Per2^{Luc}* iPSM colonies was performed in gelatin-coated 24-well black plates or 35-mm dishes (26). DMEM was used that was supplemented with 15% Knock-out Serum Replacement (KSR), 2 mM L-glutamine, 1 mM nonessential amino acids, 1 mM sodium pyruvate, 100 units/mL penicillin and streptomycin, 0.5% DMSO, 1 µM CHIRON99021, and 0.1 µM LDN193189 (Sigma) containing 1 mM luciferin and 10 mM HEPES. For live imaging of single iPSM colonies using an EM-CCD camera, each iPSM colony was cultured on a fibronectin-coated glass base dish for 6 h, and images were acquired every 5 min with an exposure time of 10 sec (control) or 2.5 sec (*Clock/Bmal1* induction) under 5% CO_2 using an LV200 Bioluminescence Imaging System (Olympus).

Gastruloids were generated as described in a previous report (27). In total, 200–250 live cells were plated in 40 µl of N2B27 medium into each well of a U-bottomed nontissue culture-treated 96-well plate (Greinier 650185). After a 96-hr cultivation, the gastruloids were embedded in 10% Matrigel (Corning 356231) containing 1 mM luciferin. For live imaging of single gastruloids, the images were acquired every 5 min with an exposure time of 3.5 sec (*Clock/Bmal1* induction) or 10 sec (control) under 5% CO_2 using the LV200 system. The Videos were analyzed using the ImageJ software (38). Kymographs of the averaged bioluminescence intensity along the straight or segmented line of 5-pixel width were generated using the plug-in KymoResliceWide.

In situ hybridization

Hybridization chain reaction (HCR) v3 was performed as described previously (27, 39) using reagents procured from Molecular Instruments. *Uncx4.1* HCR probe (Accession NM_013702.3, hairpin B1) was labeled with Alexa Fluor 488.

Quantitative RT-PCR

The iPSM colonies, ESCs, and gastruloids were washed with ice-cold PBS, and total RNA was extracted using Isogen reagent (Nippon Gene) or miRNeasy Mini Kits (QIAGEN) according to the manufacturer's instructions. To remove the feeder cells from ESCs cultured on a feeder

layer, the cells were treated with trypsin, and then the mixed cell populations were seeded on
gelatin-coated dishes and incubated for 25 min at 37°C three times in ES cell medium. Non-
attached ESCs were seeded in a gelatin-coated dish overnight and then treated with or without 500
ng/mL doxycycline for 6 hr. The iPSM colonies and gastruloids were treated with or without 1000
ng/mL doxycycline for 2 hr. First-strand cDNAs were synthesized with 1000 or 280 ng of total
RNA using M-MLV reverse transcriptase (Invitrogen) according to the manufacturer's
instructions. Quantitative PCR analysis was performed using the StepOnePlus™ Real-Time PCR
system (Applied Biosystems) and iTaq™ Universal SYBR Green Supermix (Bio-Rad
Laboratories). Standard PCR amplification protocols were applied, followed by dissociation-curve
analysis to confirm specificity. Transcription levels were normalized to the level of β-actin. The
following primer sequences were used:

320	*Bmal1*	Forward (F)	CCACCTCAGAGCCATTGATACA
321	*Bmal1*	Reverse (R)	GAGCAGGTTTAGTTCCACTTTGTCT
322	*Clock*	F	ATTTCAGCGTTCCCATTTGA
323	*Clock*	R	TGCCAACAAATTTACCTCCAG
324	*Per1*	F	CCCAGCTTTACCTGCAGAAG
325	*Per1*	R	ATGGTCGAAAGGAAGCCTCT
326	*Per2*	F	CAGCACGCTGGCAACCTTGAAGTAT
327	*Per2*	R	CAGGGCTGGCTCTCACTGGACATTA
328	*Cry1*	F	TGAGGCAAGCAGACTGAATATTG
329	*Cry1*	R	CCTCTGTACCGGGAAAGCTG
330	*Cry2*	F	CTGGCGAGAAGGTAGAGTGG
331	*Cry2*	R	GACGCAGAATTAGCCTTTGC
332	*Dbp*	F	CGAAGAACGTCATGATGCAG
333	*Dbp*	R	GGTTCCCCAACATGCTAAGA
334	*Hes6*	F	CAACGAGAGTCTTCAGGAGCTGCG
335	*Hes6*	R	GCATGCACTGGATGTAGCCAGCAG

336	*Hes7*	F	GAGAGGACCAGGGACCAGA
337	*Hes7*	R	TTCGCTCCCTCAAGTAGCC
338	*Vamp2*	F	GAGCTGGATGACCGTGCAGATG
339	*Vamp2*	R	ATGGCGCAGATCACTCCCAAGA
340	*Aloxe3*	F	AAGCCCGCCAAGAATGTTATC
341	*Aloxe3*	R	CGGTTCCCAGAGTTGTCATCC
342	*Actb*	F	GGCTGTATTCCCCTCCATCG
343	*Actb*	R	CCAGTTGGTAACAATGCCATGT

RNA-seq

The iPSM colonies and ESCs were washed with ice-cold PBS, and total RNA was extracted
using miRNeasy Mini Kits (QIAGEN) according to the manufacturer's instructions. Total RNA
sequencing was conducted by Macrogen Japan on an Illumina NovaSeq 6000 with 101-bp paired-
end reads. After trimming the adaptor sequences using Trimmomatic (40), the reads that mapped
to ribosomal DNA (GenBank: BK000964.1) (41) were filtered out and the sequence reads were
mapped to the mouse genome (GRCm38/mm10) using STAR (42), as described previously (16).
To obtain reliable alignments, reads with a mapping quality of less than ten were removed using
SAM tools (43). The known canonical genes from GENCODE VM23 (44) were used for
annotation, and the reads mapped to the gene bodies were quantified using Homer (45). The
longest transcript for each gene was used for gene-level analysis. We assumed that a gene was
expressed when there were more than 20 reads mapped on average to the gene body. Differential
gene expression in the RNA-seq data was determined using DESeq2 with thresholds of FDR <
0.05, fold change > 1.5, and expression level cutoff > 0.1 FPKM (46). WebGestalt was used for
KEGG pathway enrichment analysis (47). In the RNA-seq data using iPSM colonies, the reads
mapped in the promoter (chr11:69115096-69120473) and 3′ UTR (chr11:69122995-69123324)
of *Hes7* were filtered out to eliminate transcripts from the *pHes7-luc* reporter transgene. The
heatmaps of gene expression and KEGG pathways were generated with R using the pheatmap and
pathview packages, respectively.

Immunostaining

Immunostaining

The iPSM colonies were fixed in cold methanol for 15 min at room temperature. The fixed

iPSM was blocked with 1% BSA or 5% skim milk overnight at 4°C and then incubated with anti-

CLOCK mouse antibody (CLSP4) (48), anti-BMAL1 mouse antibody (MBL, JAPAN), anti-

BMAL1 guinea pig antibody (16), or anti-PER1 rabbit antibody (AB2201, Millipore) overnight at

4°C. After washing in 1% BSA, the iPSM colonies were incubated with a CF™488A-conjugated

donkey anti-mouse IgG (Nacalai), Cy3-conjugated goat anti-guinea pig IgG (Jackson),

DyLight™488-conjugated donkey anti-rabbit IgG (Jackson) for 2 hr at 4°C, and the nuclei were

stained with TO-PRO®-3 1:1000 (Thermo Fisher Scientific, USA) for 10–20 min. The iPSM

colonies were washed in 1% BSA and observed using an LSM510 or 900 confocal laser scanning

microscope (Zeiss).

Data availability

Data availability

RNA sequence data are available at the Gene Expression Omnibus. All other datasets generated in

this study are available from the corresponding author upon reasonable request.

Acknowledgments: We thank the Yagita lab members for technical assistance. This work was

supported in part by grants-in-aid for scientific research from the Japan Society for the Promotion

of Science to Y.U. (19K06679) and K.Y. (18H02600), the Cooperative Research Program (Joint

Usage/Research Center program) of the Institute for Frontier Life and Medical Sciences, Kyoto

University (K.Y. and G.K.)

References

References

1.   A. Balsalobre, F. Damiola, U. Schibler, A serum shock induces circadian gene expression

in mammalian tissue culture cells. *Cell* **93**, 929-937 (1998).

2.   J. S. Takahashi, Transcriptional architecture of the mammalian circadian clock. *Nat Rev*

Genet **18**, 164-179 (2017).

3.   K. Yagita, F. Tamanini, G. T. van Der Horst, H. Okamura, Molecular mechanisms of the
biological clock in cultured fibroblasts. *Science* **292**, 278-281 (2001).

4.   S. Yamazaki *et al.*, Resetting central and peripheral circadian oscillators in transgenic rats.
Science **288**, 682-685 (2000).

5.   S. H. Yoo *et al.*, PERIOD2::LUCIFERASE real-time reporting of circadian dynamics
reveals persistent circadian oscillations in mouse peripheral tissues. *Proc Natl Acad Sci U*
S A **101**, 5339-5346 (2004).

6.   J. B. Hogenesch, H. R. Ueda, Understanding systems-level properties: timely stories from
the study of clocks. *Nat Rev Genet* **12**, 407-416 (2011).

7.   K. Yagita *et al.*, Development of the circadian oscillator during differentiation of mouse
embryonic stem cells in vitro. *Proc Natl Acad Sci U S A* **107**, 3846-3851 (2010).

8.   J. D. Alvarez, D. Chen, E. Storer, A. Sehgal, Non-cyclic and developmental stage-specific
expression of circadian clock proteins during murine spermatogenesis. *Biol Reprod* **69**, 81-
91 (2003).

9.   T. Amano *et al.*, Expression and functional analyses of circadian genes in mouse oocytes
and preimplantation embryos: Cry1 is involved in the meiotic process independently of
circadian clock regulation. *Biol Reprod* **80**, 473-483 (2009).

10. E. Kowalska, E. Moriggi, C. Bauer, C. Dibner, S. A. Brown, The circadian clock starts
ticking at a developmentally early stage. *J Biol Rhythms* **25**, 442-449 (2010).

11. D. Morse, N. Cermakian, S. Brancorsini, M. Parvinen, P. Sassone-Corsi, No circadian
rhythms in testis: Period1 expression is clock independent and developmentally regulated
in the mouse. *Mol Endocrinol* **17**, 141-151 (2003).

12. F. C. Davis, R. A. Gorski, Development of hamster circadian rhythms: role of the maternal
suprachiasmatic nucleus. *J Comp Physiol A* **162**, 601-610 (1988).

13. C. Jud, U. Albrecht, Circadian rhythms in murine pups develop in absence of a functional
maternal circadian clock. *J Biol Rhythms* **21**, 149-154 (2006).

14. S. M. Reppert, W. J. Schwartz, Maternal suprachiasmatic nuclei are necessary for maternal
coordination of the developing circadian system. *J Neurosci* **6**, 2724-2729 (1986).

15. V. Carmona-Alcocer *et al.*, Ontogeny of Circadian Rhythms and Synchrony in the
Suprachiasmatic Nucleus. *J Neurosci* **38**, 1326-1334 (2018).

16. Y. Umemura *et al.*, Transcriptional program of Kpna2/Importin-alpha2 regulates cellular
differentiation-coupled circadian clock development in mammalian cells. *Proc Natl Acad*
Sci U S A **111**, E5039-5048 (2014).

17. Y. Umemura *et al.*, Involvement of posttranscriptional regulation of Clock in the
emergence of circadian clock oscillation during mouse development. *Proc Natl Acad Sci*
U S A **114**, E7479-E7488 (2017).

18. Y. Umemura, I. Maki, Y. Tsuchiya, N. Koike, K. Yagita, Human Circadian Molecular
Oscillation Development Using Induced Pluripotent Stem Cells. *J Biol Rhythms*
10.1177/0748730419865436, 748730419865436 (2019).

19. Y. Harima, I. Imayoshi, H. Shimojo, T. Kobayashi, R. Kageyama, The roles and
mechanism of ultradian oscillatory expression of the mouse Hes genes. *Semin Cell Dev*
Biol **34**, 85-90 (2014).

20. A. Hubaud, O. Pourquie, Signalling dynamics in vertebrate segmentation. *Nat Rev Mol*
Cell Biol **15**, 709-721 (2014).

21. Y. Bessho, H. Hirata, Y. Masamizu, R. Kageyama, Periodic repression by the bHLH factor
Hes7 is an essential mechanism for the somite segmentation clock. *Genes Dev* **17**, 1451-
1456 (2003).

22. Y. Takashima, T. Ohtsuka, A. Gonzalez, H. Miyachi, R. Kageyama, Intronic delay is
essential for oscillatory expression in the segmentation clock. *Proc Natl Acad Sci U S A*
108, 3300-3305 (2011).

23. Y. Bessho *et al.*, Dynamic expression and essential functions of Hes7 in somite
segmentation. *Genes Dev* **15**, 2642-2647 (2001).

24. O. Q. H. Zinani, K. Keseroglu, A. Ay, E. M. Ozbudak, Pairing of segmentation clock genes
drives robust pattern formation. *Nature* **589**, 431-436 (2021).

25. M. Matsumiya, T. Tomita, K. Yoshioka-Kobayashi, A. Isomura, R. Kageyama, ES cell-
derived presomitic mesoderm-like tissues for analysis of synchronized oscillations in the
segmentation clock. *Development* **145** (2018).

26.   Y. B. Kiyohara *et al.*, The BMAL1 C terminus regulates the circadian transcription
451   feedback loop. *Proc Natl Acad Sci U S A* **103**, 10074-10079 (2006).

452   27.   S. C. van den Brink *et al.*, Single-cell and spatial transcriptomics reveal somitogenesis in
453   gastruloids. *Nature* **582**, 405-409 (2020).

454   28.   M. Kanehisa, S. Goto, KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids*
455   *Res* **28**, 27-30 (2000).

456   29.   Y. Niwa *et al.*, The initiation and propagation of Hes7 oscillation are cooperatively
457   regulated by Fgf and notch signaling in the somite segmentation clock. *Dev Cell* **13**, 298-
458   304 (2007).

459   30.   M. Matsuda *et al.*, Recapitulating the human segmentation clock with pluripotent stem
460   cells. *Nature* **580**, 124-129 (2020).

461   31.   N. Koike *et al.*, Transcriptional architecture and chromatin landscape of the core circadian
462   clock in mammals. *Science* **338**, 349-354 (2012).

463   32.   M. Ebisuya, T. Yamamoto, M. Nakajima, E. Nishida, Ripples from neighbouring
464   transcription. *Nat Cell Biol* **10**, 1106-1113 (2008).

465   33.   D. M. Owens, S. M. Keyse, Differential regulation of MAP kinase signalling by dual-
466   specificity protein phosphatases. *Oncogene* **26**, 3203-3213 (2007).

467   34.   R. T. Moon, J. D. Brown, J. A. Yang-Snyder, J. R. Miller, Structurally related receptors
468   and antagonists compete for secreted Wnt ligands. *Cell* **88**, 725-728 (1997).

469   35.   J. K. Dale *et al.*, Periodic notch inhibition by lunatic fringe underlies the chick
470   segmentation clock. *Nature* **421**, 275-278 (2003).

471   36.   Z. Chen *et al.*, Identification of diverse modulators of central and peripheral circadian
472   clocks by high-throughput chemical screening. *Proc Natl Acad Sci U S A* **109**, 101-106
473   (2012).

474   37.   Y. Inada *et al.*, Cell and tissue-autonomous development of the circadian clock in mouse
475   embryos. *FEBS Lett* **588**, 459-465 (2014).

476   38.   C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image
477   analysis. *Nat Methods* **9**, 671-675 (2012).

478   39.   H. M. T. Choi *et al.*, Third-generation in situ hybridization chain reaction: multiplexed,
479   quantitative, sensitive, versatile, robust. *Development* **145** (2018).

480   40.   A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
481   sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

482   41.   N. R. Coordinators, Database resources of the National Center for Biotechnology
483   Information. *Nucleic Acids Res* **46**, D8-D13 (2018).

484   42.   A. Dobin *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
485   (2013).

486   43.   H. Li *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-
487   2079 (2009).

488   44.   A. Frankish *et al.*, GENCODE reference annotation for the human and mouse genomes.
489   *Nucleic Acids Res* **47**, D766-D773 (2019).

490   45.   S. Heinz *et al.*, Simple combinations of lineage-determining transcription factors prime
491   cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**, 576-
492   589 (2010).

493   46.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
494   RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

495   47.   Y. Liao, J. Wang, E. J. Jaehnig, Z. Shi, B. Zhang, WebGestalt 2019: gene set analysis
496   toolkit with revamped UIs and APIs. *Nucleic Acids Res* **47**, W199-W205 (2019).

497   48.   H. Yoshitane *et al.*, Roles of CLOCK phosphorylation in suppression of E-box-dependent
498   transcription. *Mol Cell Biol* **29**, 3675-3686 (2009).

Fig. 1. Establishment of ESC lines carrying both the dox-inducible *Clock* and *Bmal1* genes.
(**A**) Human genomic locus of circadian clock genes, *PER1*, and the essential segmentation clock gene, *HES7*, are highly conserved in higher vertebrates. The *PER1* homolog *PER2* is also located adjacent to the *HES7* homolog *HES6* in the genome. (**B**) ESCs were differentiated into iPSM for 96 hr *in vitro*, and then the iPSM colonies were treated with or without dox. (**C**) qPCR of *Clock* and *Bmal1* mRNA in the indicated ESCs. 500 ng/mL dox treatment for 6 hr (red) or not (blue). Each number indicates clone number. Mean ± SD (n = 3 biological replicates). (**D**) qPCR of *Clock* and *Bmal1* mRNA in the indicated iPSM colonies. 1000 ng/mL dox treatment for 2 hr (red) or not (blue). Mean ± SD (n = 2 technical replicates). The average expression level of *pHes7-luc* ESCs or iPSM colonies without dox was set to 1. (**E**) Representative maximum intensity projection of

510   the immunostaining of iPSM colonies treated with 1000 ng/mL dox for 2 hr or not. n = 2–3

511   biological replicates. Scale = 250 μm.

Fig. 2. CLOCK/BMAL1 expressions upregulated not only circadian clock genes but also *Hes7* gene expressions in the iPSM. (A, B) qPCR of core circadian clock genes (**A**) and *Hes*7 or *Hes6* gene (**B**) in the indicated iPSM colonies. 1000 ng/mL dox treatment for 2 hr (red) or not (blue). Mean ± SD (n = 2 technical replicates). The average expression level of iPSM colonies without dox was set to 1. Two-tailed t-test, *P < 0.05, **P < 0.01.

Fig. 3. CLOCK/BMAL1 expressions arrested the autonomous oscillations of *Hes7* in the iPSM colonies. (**A**) Bioluminescence of each dox-inducible *Clock/Bmal1* or *Clock/Bmal1DN pHes7-luc* iPSM colony was observed using PMT or an EM-CCD camera without or with dox. (**B, C**) Representative bioluminescence traces (**B**, n = 25 biological replicates) and live imaging (**C**, n = 3 biological replicates) of single *pHes7-luc* iPSM colony with or without dox. The kymograph of the imaging along the arrow is shown. (**D**) Representative bioluminescence traces of the single indicated iPSM colony with and without 1000 ng/mL dox. n = 10-54 biological replicates. (**E–H**) Live imaging of the single *tetO-Clock/Bmal1 pHes7-luc* iPSM colony with and without dox. Dox-containing medium or only medium was added at the indicated time points at the final dox concentration of 1000 ng/mL (**G, H**). Each kymograph along the arrow is shown. Scales = 250 μm. n = 2-4 biological replicates.

Fig. 4. CLOCK/BMAL1 expressions interfered with the autonomous oscillations of *Hes7* and somitogenesis-like process in the gastruloids. (**A**) Dox-inducible *Clock/Bmal1 pHes7-luc* ESCs were differentiated into gastruloids for 96 hr *in vitro*, and then the gastruloids embedded in 10% Matrigel were treated with or without dox. *pHes7-luc* bioluminescence was observed using an EM-CCD camera without or with dox. (**B–M**) Time-lapse bioluminescence (red) and bright field imaging of the single *pHes7-luc* gastruloid or *tetO-Clock/Bmal1 pHes7-luc* gastruloid without and with dox. Dox-containing medium was added at the indicated time points at the final dox concentration of 1000 ng/mL (**C, F, I, L**). Each kymograph is shown along the yellow lines in **B, E, H, K**. *In situ* hybridization of *Uncx4.1* in the gastruloids after the live cell imaging (**D, G, J, M**). Scales = 250 μm. n = 2-4 biological replicates.

Fig. 5. CLOCK/BMAL1 expressions in the iPSM colonies affected *Hes7*-related signaling pathways and upregulated the expression of contiguous genes, *Per1*, *Hes7*, and *Aloxe3*. (A) Upregulated and downregulated DEGs in the indicated iPSM colonies treated with dox. (B) KEGG

pathway analysis of the DEGs. Each pathway was indicated with each transformed p-value. The

ranked pathways contained several common genes in the WNT, MAPK, and NOTCH signaling

pathways. (**C**) UCSC genome browser views of RNA-seq data of the contiguous genes *Vamp2*,

Per1, *Hes7*, and *Aloxe3*. The reads shown are normalized average reads per 10 million total reads

in 10-bp bins. (**D**) mRNA expression of *Vamp2*, *Per1*, *Hes7*, and *Aloxe3* in the indicated iPSM

colonies according to RNA-seq. (**E**) Validation of *Vamp2* and *Aloxe3* gene expression levels in

the indicated iPSM colonies using qPCR. Colored boxes indicate 1000 ng/mL dox treatment for 2

hr (red) or no treatment (blue). Mean ± SD (n = 2 technical replicates). The averaged expression

level of *pHes7-luc* iPSM colonies without dox was set to 1. Two-tailed t-test, *P < 0.05, **P <

0.01. (**F**) The premature expression of CLOCK/BMAL1 in the iPSM and gastruloids interfered

with the segmentation clock oscillation and somitogenesis-like process.

Supporting information for

Circadian key component CLOCK/BMAL1 interferes with segmentation clock in mouse embryonic organoids

Yasuhiro Umemura, Nobuya Koike, Yoshiki Tsuchiya, Hitomi Watanabe, Gen Kondoh, Ryoichiro Kageyama, and Kazuhiro Yagita

Correspondence to: kyagita@koto.kpu-m.ac.jp

Fig. S1. Mutually exclusive appearance of segmentation clock and circadian clock. In

mammals, two different types of rhythm appear sequentially during the developmental process.

One is the ultradian rhythm by segmentation clock, which controls somitogenesis. The other one

is the circadian oscillation, which regulates the predictive adaptation of physiological functions

to the day–night environmental cycle.

Fig. S2. The iPSM has no apparent circadian clock oscillation. (A) The expression levels of
pluripotent marker genes, *Nanog*, *Sox2*, and *Oct3/4,* were measured using RNA-seq in ESCs and
pHes7-luc iPSM. The data were shown with *in vitro* 0, 7-, and 28-day differentiated WT ESCs and
Per2^{Luc} ESCs (GSE61184). No circadian clock oscillates in the *in vitro* 7-day differentiated ESCs
and the circadian clock oscillation starts to emerge after 14 days of differentiation culture (16, 17).
(B) Bioluminescence traces of the iPSM differentiated from *Per2^{Luc}* ESCs. The iPSM was
stimulated with 10 µM forskolin (Fsk, middle) or 100 nM dexamethasone (Dex, bottom). n = 3
biological replicates. **(C)** Representative immunostaining of CLOCK, BMAL1, and PER1 in the
iPSM. The immunostaining represented the suppression of CLOCK proteins in contrast to the
expression of BMAL1. Furthermore, although the nuclei indicated the quite faint signals, PER1
protein in the iPSM is still predominantly accumulated in the cytoplasm.

Fig. S3. The core circadian clock gene expressions in the ESCs harboring dox-inducible *Clock*.

Although the expression of BMAL1 protein was observed even in ESCs (17), dox-dependent sole

expression of *Clock* was insufficient for the E-box-driven expression of clock genes such as *Per1/2*

and *Cry1/2* in undifferentiated ESCs, raising the possibility that the endogenously expressed

BMAL1 might be post-translationally modified to not function. Colored boxes indicate 500 ng/mL

dox treatment for 6 hr (red) or no treatment (blue). Each number indicates clone number. Mean ±

SD (n = 3 biological replicates). The averaged expression level of *Per2^Luc* ESCs without dox was

set to 1. Two-tailed t-test, *P < 0.05, **P < 0.01, ***P < 0.001.

Fig. S4. *Clock/Bmal1* gene expressions in ESCs upregulated not only circadian clock genes but also *Hes7* gene expression. (A, B) qPCR of core circadian clock gene (A) and *Hes*7 or *Hes6* gene (B) expression in ESCs harboring dox-inducible *Clock* and *Bmal1* or *Clock* and *Bmal1DN*. Colored boxes indicate the presence (red) or absence (blue) of 500 ng/mL dox treatment for 6 hr. Each number indicates clone number. Mean ± SD (n = 2–3 biological replicates). The average expression level of ESCs without dox was set to 1.

Fig. S5. qPCR of *Clock* and *Bmal1* mRNA in the indicated gastruloids. Colored boxes indicate 1000 ng/mL dox treatment for 2 hr (red) or no treatment (blue). Each number indicates clone number. Mean ± SD (n = 3 biological replicates). The average expression level of *pHes7-luc* gastruloids without dox was set to 1.

Fig. S6. *Clock/Bmal1* **gene expressions in ESCs affected the** *Hes7***-related signaling pathways**
and upregulated the expression of the contiguous genes *Vamp2***,** *Per1***,** *Hes7***, and** *Aloxe3***.** (**A**)
Upregulated and downregulated differentially expressed genes in the indicated ESCs treated with
dox (FDR < 0.05, FC > 1.5). (**B**) KEGG pathway analysis of the DEGs. Each pathway was
indicated with each transformed p-value. The ranked pathways contained several common genes
in the WNT, MAPK, and NOTCH signaling pathways. (**C**) UCSC genome browser views of RNA-
seq data of contiguous genes, including *Vamp2*, *Per1*, *Hes7*, and *Aloxe3*. The reads shown are
normalized average reads per 10 million total reads in 10-bp bins. (**D**) mRNA expression of *Vamp2*,
Per1, *Hes7*, and *Aloxe3* in the indicated ESCs measured using RNA-seq. (**E**) Validation of *Vamp2*

52   and *Aloxe3* in the indicated ESCs by qPCR. Colored boxes indicate the use of 500 ng/mL dox

53   treatment for 6 hr (red) or untreated cells (blue). Each number indicates clone number. Mean ± SD

54   (n = 3 biological replicates). Two-tailed t-test, *P < 0.05, **P < 0.01, ***P < 0.001, ****P <

55   0.0001, *****P < 0.00001.