A non-olfactory shark adenosine receptor activates CFTR with unique pharmacology
and structural features

Sumeet Bhanot, Gabriele Hemminger, Cole L. Martin, Stephen G. Aller and John N. Forrest Jr.

Renal Section, Department of Medicine, Yale University School of Medicine, New Haven CT 06510
The Mount Desert Island Biological Laboratory, Salisbury Cove, ME 04672
The Department of Pharmacology & Toxicology, University of Alabama at Birmingham, Birmingham, AL 35205

Running Head: Characterization of a novel adenosine receptor (A_0) from the shark
rectal gland (*Squalus acanthias*)

Corresponding Authors: Stephen G. Aller, Department of Pharmacology & Toxicology, University of
Birmingham at Alabama, Birmingham, AL 35205, sgaller@uab.edu
and John N. Forrest Jr., Department of Medicine, Yale University School of Medicine, 367 Cedar St.,
New Haven CT 06510, john.forrest@yale.edu

Abstract

     Adenosine receptors (ADORs) are G-protein coupled purinoceptors that have several functions including regulation of chloride secretion via CFTR in human airway and kidney. We cloned an ADOR from *Squalus acanthias* (shark) that likely regulates CFTR in the rectal gland. Phylogenic- and expression- analyses indicate that elasmobranch ADORs are non-olfactory, and appear to represent extant predecessors of mammalian ADORs. We therefore designate the shark ADOR as the A_0 receptor. We co-expressed A_0 with CFTR in *Xenopus laevis* oocytes and characterized the coupling of A_0 to the chloride channel. Two electrode voltage clamping was performed and current-voltage (I-V) responses were recorded to monitor CFTR status. Only in A_0- and CFTR- co-injected oocytes did adenosine analogs produce a significant concentration-dependent activation of CFTR consistent with its electrophysiological signature. A pharmacological profile for A_0 was obtained for ADOR agonists and antagonists that differed markedly from all mammalian ADOR subtypes (agonists: R-PIA > S-PIA > CGS21680 > CPA > 2ClADO > CV1808 = DPMA > NECA) and (antagonists: DPCPX > PD115199 > 8PT > CGC > CGS15943). Structures of human ADORs permitted a high-confidence homology model of the shark A_0 core which revealed unique structural features of ancestral receptors. We conclude: (1) A_0 is a novel and unique adenosine receptor ancestor by functional and structural criteria; (2) A_0 likely activates CFTR *in vivo* and this receptor activates CFTR in oocytes indicating an evolutionary coupling between ADORs and chloride secretion; and (3) A_0 appears to be a non-olfactory evolutionary ancestor of all four mammalian ADOR subtypes.

Keywords

Adenosine receptor, molecular evolution, shark, structural biology, CFTR, chloride secretion

Significance Statement

We have cloned and characterized an ancient adenosine receptor from sharks that is unlikely to be olfactory in function. The shark receptor, which we designate as A_0, has a unique pharmacological profile, characteristic structural features, and is also highly likely to be the dominant ADOR regulator of the shark ancient ortholog of the Cystic Fibrosis chloride channel, called CFTR.

Introduction

58
59

Two broad classes of purinoceptors, P_1 & P_2, recognize and bind adenosine-based nucleosides and
nucleotides, and can be distinguished by two anti-parallel pharmacological profiles (P_1: Adenosine > AMP
> ADP > ATP & P_2: ATP > ADP > AMP > Adenosine) (1). Receptors specific for adenosine (ADORs) are
divided into four non-olfactory subtypes including A_1, A_{2a}, A_{2b} & A_3, which couple to $G_{o,i(1-3)}$, G_s, G_s
and $G_{i(2,3),q}$-like G proteins, respectively (2-7). Typically, all ADORs modulate the activity of at least one
of a number of different isoforms of adenylyl cyclase through the coupling to a G protein. The A_1 ADOR
can also modulate other effector systems, namely potassium channels, calcium channels, phospholipase A
and C (7).

One major unifying theme in the function of ADORs in mammals appears to be a feedback mechanism to
either curb metabolic activity of the cell or to oppositely vasodilate for increasing oxygen supply and meet
higher metabolic demand (8). Relatedly, in sharks, which evolved prior to any land-based vertebrate, low
concentrations of adenosine released from metabolic activity by the highly active shark rectal gland of
Squalus acanthias, serve as an autocoid feedback inhibitor to slow ion transport (9). Curiously, high
concentrations of adenosine (~10 µM) increased ion transport activity at both the apical- and basolateral-
membranes, and stimulated chloride secretion in rectal gland via the shark ortholog of the Cystic Fibrosis
transmembrane Conductance Regulator (CFTR) (10,11). Shark CFTR is an impressive 70% identical to
human CFTR at the amino acid residue sequence level (12), and an ancient evolutionary link in ADOR
signaling of CFTR activity in both fish and mammals is now well established.

The four main ADOR subtypes (A_1, A_{2a}, A_{2b} & A_3), first identified in many different mammals and now
observed in the genomes of birds, reptiles, amphibians and fish, can be distinguished at phylogenetic- and
pharmacological levels. For the A_1 receptor, the adenosine analogs CPA and R-PIA are the most potent,
while CV1808 and CGS21680 are weaker agonists. For A_{2a}, NECA, CV1808, and CGS21680 are the most
potent agonists, and CPA is less potent. The A_{2b} receptor subtype responds with high potency to NECA but
exhibits little or no response to CV1808 and CGS21680. A fifth ADOR subtype, designated A2c, was more
recently identified and was shown to be an olfactory ADOR that allows fish to sense adenosine released into
the water by potential prey (13).

We sought to examine ADORs in *Squalus acanthias* and we identified a single receptor with highest
expression in the rectal gland and stomach, but very low levels in brain and other tissues. Phylogenetic
analysis shows that the *Squalus acanthias* ADOR, as well as putative ADORs identified by genome
sequencing from other sharks, amphibians and reptiles, form a separate clade that is distinct from the other
five ADOR subtypes. We used the *Xenopus laevis* oocyte system to co-express the *Squalus acanthias*
ADOR with CFTR to demonstrate the coupling of receptor to chloride channel. Co-injected oocytes
revealed a pharmacological profile of *Squalus acanthias* ADOR that is unique from all four major subtypes
of ADORs. We argue that the *Squalus acanthias* ADOR is not olfactory and that the clade containing the
elasmobranch ADORs is ancestral to the four major subtypes. We therefore designated the shark ADOR as
A_0-ADOR, and characterize this ancestral receptor here including novel structural features revealed by
three-dimensional atomic homology modeling.

Materials and Methods
Molecular Biology

Tissues were obtained from male dogfish sharks, *Squalus acanthias*, weighing 2-4 kg, which were caught by gill nets in Frenchman's Bay, ME, and kept in marine live cars until use, usually within 3 days of capture. One male dogfish shark was killed by pithing the spinal cord. Details describing the molecular biology (preparation of total RNA from shark tissue, degenerate PCR, library screening, RACE PCR, Northern blot, qRT-PCR and electrophysiology in oocytes) are presented in the supplementary material (see supplementary figs. S1-S5, https://www.biorxiv.org/content/biorxiv/early/2020/11/02/2020.11.01.363762/DC1/embed/media-1.pdf?download=true). Human CFTR was obtained from Dr. Carol Semrad in the pKS-βglobin-CFTR-polyAC plasmid and human adenosine receptors were provided by Dr. Joel Linden. Capped messenger RNA for all receptors was synthesized from linearized plasmid templates using the Ambion mMessage mMachine T7 *in vitro* transcription kit according to the manufacturer's protocol.

To retrieve putative adenosine receptor coding sequences with good representation from mammals, birds, reptiles, amphibians and fish, multiple database searches were carried out using the NCBI BlastP Server against the full nonredundant dataset. To find potential ancestral receptor sequences, additional searches using shark A_0 sequence were performed excluding the Mammalia taxid (40674). Global alignments were performed using the Clustal method (14) within the Lasergene software package (DNASTAR, Inc) and similar results were obtained using the implementation of Clustal Omega (15) in the EMBOSS package (16) and standard parameters (17). Alignments for phylogenetic analyses including the production of the phylogenetic tree in figure 1 were conducted with MEGA7 (18).

Xenopus laevis expression system

Adult female *Xenopus laevis* were obtained from Nasco and Xenopus 1, and housed in the animal care facilities at Yale University. Surgeries to harvest oocytes were performed 4-7 days prior to experiments according to a protocol approved by the Yale Animal Care Committee. Injection pipets pulled on a Sachs-Flaming Micropipette puller Model PC-84, using Drummond 210/310/510 glass pipettes were beveled to a tip diameter of 10-15 μm. The pipettes were back-filled with mineral oil and attached to a Drummond microinjector, in preparation for cRNA aspiration and oocyte injection. Oocytes were injected with 50 nl (~15 ng) of cRNA encoding shark A_0- or human A_1, A_{2a}, A_{2b}, or A_3 receptors and 5-10 ng of human CFTR, and subsequently incubated for 3-5 days at 18-20 °C before experiments.

Oocytes were prepared for two electrode voltage clamping by inserting two 0.6-0.9 MΩ, 3 M KCl filled Ag electrodes, pulled on a Sachs-Flaming micro-pipet puller model PC-84, using borosilicated glass obtained from Sutter Instrument (BF120-69-10). Oocytes were perifused with frog Ringer's, FR, (98 mM NaCl, 2 mM KCl, 1.8 mM $CaCl_2$, 1 mM $MgCl_2$ and 5 mM HEPES. Adjusted to pH 7.4), which was also used in the preparation of solutions for the pharmacological agents tested. Oocytes were voltage clamped using the Dagan TEV-200 amplifier along with the Dagan LM-12 Lab Interface, on a PC computer running pClamp Version 5.5 software. The Soltec 1242 plotter was used to acquire the data. Oocytes were clamped at -60 mV and the current was measured as a function of time. Moreover, frequent voltage pulses (-120mV to +60mV) were performed to determine the current-voltage (I-V) relationship at any given timepoint. Data were corrected for any capacitive current incurred due to the voltage pulses. Inconclusive data were obtained with the human A_{2a} and A_3 receptors, which presumably do not couple efficiently to frog G proteins or the proper adenylyl

cyclase(s) for CFTR activation, and were omitted from the analysis.

Data analysis was performed using Microsoft Excel. Some of the data were analyzed using an

Excel macro provided by the lab of Dr. David Dawson, which allowed the subtraction of the capacitive

current from the current voltage relationship, incurred during the voltage pulses. Statistical analysis was

limited to standard deviation performed on the concentration response of the shark A_0 receptor.

Homology model of the shark A_0

At the time of this writing there were 48 high-resolution structures of adenosine receptors in

the Protein Data Bank (PDB) as determined by either x-ray crystallography or cryo-electron

microscopy. All structures are human in origin and only two of the four known subtypes were

represented, A_{2a} (n=45) and A_1 (n=3). We chose a small subset for consideration as templates for

homology modeling of the shark A_0 receptor: A_{2a} with adenosine bound (PDB code 2ydv) (19), A_1

with adenosine (6d9h) (20), A_{2a} with N-ethyl-5'-carboxamido adenosine, NECA (6gdg) (21), and

A_{2a} with CGS21680 (4ug2) (22). Primary amino acid sequence alignments between shark A_0

sequence and the selected subset were performed using NCBI Cobalt (23). Amino acid sequences

present in the human receptors but not present in shark A_0 were removed (see full alignment

between shark A_0 and human A_{2a} in supplementary fig. S6,

https://www.biorxiv.org/content/biorxiv/early/2020/11/02/2020.11.01.363762/DC1/embed/media-1.pdf?download=true).

The human A_{2a} receptor appeared to be a sufficiently good template for A_0 modeling as it shares

45% identical amino acid sequence with A_0 and exhibited an NCBI Cobalt 3-bit amino acid

chemical similarity of 81% after removing disordered sequences from the A_{2a} structure, and after

removing sequences present in either receptor but not the other (supplementary fig S7,

https://www.biorxiv.org/content/biorxiv/early/2020/11/02/2020.11.01.363762/DC1/embed/media-1.pdf?download=true).

Using the A_{2a} x-ray crystal structure template (PDB code 2ydv), A_{2a} insertion sequences were

removed and mutagenesis to A_0 was performed with scripts written in perl and PyMOL version

2.0.6 (Schrödinger, LLC). Two short amino acid sequences of shark A_0 that are not present in the

human receptors (i.e. A_0-Arg95 and A_0-Asn228-Ser230), were inserted into the 3-dimensional

model using Coot version 0.8.9 (24) and model minimization/regularization. Refinements using

phenix (25) of A_0 against A_{2a} dataset (PDB code 2ydv) were performed by replacing the human

receptor with the shark starting model. Three rounds of rigid body, XYZ-reciprocal space and B-

factor refinement were performed including secondary-structure restrains and

Ramachandran/rotamer optimization, all with resolution cutoff set to 4.0 Å resolution.

Parameterization of the R-PIA ligand was accomplished using phenix (25) starting with the

Simplified Molecular-Input Line-Entry System (SMILES) string for the ligand (26):

C[C@H](Cc1ccccc1)Nc2ncnc3n(cnc23)[C@@H]4O[C@H](CO)[C@@H](O)[C@H]4O

The position of the lipid bilayer was predicted based on the location of conserved aromatic amino

acids known to interact with lipid head groups spaced ~35 Å apart (27).

Results

Identification of shark A_0 ADOR

Since the *Squalus acanthias* genome is not yet available and there is some uncertainty associated

with the search for *bona fide* protein-coding sequences from genomic data, we sought to clone shark

adenosine receptors using reverse-transcription procedures from mRNA/cDNA prepared from shark

tissues. PCR of shark rectal gland cDNA using degenerate primers to conserved regions of

transmembrane domains 5 and 7 of previously cloned ADORs yielded a single 352 bp product (see

supplementary fig. S1). The translated amino acid sequence of this fragment (P-SH12) had highest
homology to ADORs that was only 39-45% identical to the human receptor sub-types (A_1, A_{2a}, A_{2b}
and A_3). Northern blot and quantitative RT-PCR (qRT-PCR) revealed strongest expression of the
putative shark ADOR in the rectal gland and stomach with weak levels in other tissues including the
brain (supplementary figs. S2 and S3). To control for the quality of cDNA template used in the qRT-
PCR experiments, shark CFTR was amplified from the same cDNA templates prepared from the various
shark tissues. Results showed abundant CFTR expression in rectal gland, intestine, brain and testis
(supplementary fig. S4).
Screening of a shark rectal gland cDNA library using ^{32}P-labeled P-SH12 yielded more
nucleotide sequence from a single putative ADOR and allowed the design of primers for Rapid
Amplification of cDNA Ends-PCR (RACE-PCR). Extensive rounds of both cDNA library screening and
RACE-PCR using total RNA from shark rectal gland revealed only a single putative ADOR entity
which we designate here as the shark A_0 receptor (for details see supplementary material).
Hydropathy analysis of the putative shark A_0 ADOR amino acid sequence revealed seven
predicted transmembrane regions and a predicted extracellular orientation of the N-terminus that was
consistent with all known G protein-coupled receptors (see supplementary fig. S5). Phylogenetic
analysis of known- and putative- ADORs revealed an ancestral branch that included dogfish shark A_0
and other putative ADORs from non-mammalian species that are distinct from the recently discovered
olfactory ADORs (Wakisaka et al. 2017; figure 1). Ancestral olfactory- and non-olfactory receptors
could be distinguished from the four specialized mammalian subtypes (i.e. A_1, A_{2a}, A_{2b} and A_3) as
having an abbreviated second extracellular loop (ECL2) at 9 amino acid (aa) residues (supplementary
fig. S8, https://www.biorxiv.org/content/biorxiv/early/2020/11/02/2020.11.01.363762/DC1/embed/media-
1.pdf?download=true). ECL2 of A_1 is elongated by 4-17 aa (n=42 receptor species examined), ECL2 of
A_{2a} is elongated by 8-14 aa (n=22), ECL2 of A_{2b} is elongated by 10-19 aa (n=28) and ECL2 of A_3 is
elongated by 1-7 aa (n=18, see supplementary fig. S9,
https://www.biorxiv.org/content/biorxiv/early/2020/11/02/2020.11.01.363762/DC1/embed/media-1.pdf?download=true).
Assuming mutagenesis rates of all adenosine receptors are similar amongst species, we interpret the
phylogram to indicate that the elasmobranch A_0 ADOR belongs to an evolutionary ancestral branch of
the specialized adenosine receptor subtypes. We next surmised that shark A_0 is responsive to adenosine
analogs with potentially unique properties compared to known receptor subtypes, and we sought to test
this hypothesis using functional and pharmacological characterization in *Xenopus laevis* oocytes.
219
CFTR expression
Expression studies were conducted in oocytes co-injected with human CFTR and shark A_0
cRNA, and current-voltage (I-V) relationships were determined by imposing voltage pulses (-120 mV to
+60 mV) and measuring the clamping current. Expression of CFTR was determined by the use of
forskolin, a well-known activator of adenylyl cyclase that increases intracellular cAMP levels and
activates CFTR by phosphorylation of the R-domain by a cAMP dependent protein kinase A (Dulhanty
and Riordan 1994). Figure 2 illustrates the I-V relationships of a single representative experiment on an
oocyte 3 days after co-injection. Addition of 5 μM forskolin produced an increase in conductance
(increase in the slope of the I-V relationship) from the basal level, an increase in the magnitude of the
clamping current at -60 mV from -76 nA to -563 nA, and a rightward shift in the reversal potential to -
29.4 ± 0.75 mV (n=3 I-V ramps $\pm$ SD) from a basal value of -38.0 ± 3.8 mV (n=5 ramps $\pm$ SD). This
negative, non-rectifying current with a reversal potential close to that of Cl^- ($Cl^-_{rev} = \sim$ -30 mV) has the
electrophysiological characteristics of a CFTR Cl^- current. Furthermore, this response was sensitive to

300 µM glibenclamide, an inhibitor of the CFTR Cl⁻ channel (28) (n=1, data not shown). The response

to forskolin was also inhibited by the PKA antagonist H-89 (25 µM, n=3) in oocytes injected with

human CFTR cRNA (data not shown), suggesting that CFTR is activated by a cAMP dependent protein

kinase (PKA) in this system. To confirm, we measured total cAMP in oocytes expressing the shark A_0

receptor and observed an increase in total cAMP that paralleled the order of agonist potency we

observed in electrophysiology experiments (see supplementary fig. S10). The negative, non-rectifying

CFTR current was absent in un-injected- and water injected- oocytes (n=4) under identical conditions,

and were reproducible both in the same oocyte and in all oocytes injected with human CFTR cRNA.

These data established expression of CFTR in our system.

Addition of 10 µM 2-chloroadenosine in the representative oocyte (figure 2) also

produced an increase in total current by a similar non-rectifying negative current of the

magnitude ~ -500 nA and a reversal potential of approximately -30.5 ± 0.28 mV (n=3 ramps ±

SD). This response was reversibly inhibited by 10 µM DPCPX, an adenosine receptor

antagonist, where the clamping current at -60 mV was ~ -90 nA with a reversal potential of -

39.4 ± 4.8 mV (n=3 ramps ± SD), both of which are reminiscent of the basal state (figure 2).

This response suggested that the expression and stimulation of the shark A_0 receptor is

associated with an increase in Cl⁻ current which has characteristics consistent with CFTR Cl⁻

currents. Our results also established a coupling of the shark A_0 adenosine receptor and human

CFTR in the *Xenopus* oocyte expression system.

Pharmacological characterization of shark A_0

Determination of the pharmacological profiles of adenosine receptor agonists and

antagonists in the oocyte necessitated the development of protocols in the *Xenopus laevis* oocyte

expression system. We first compared three agonists in a series of identical experiments to provide

quantitative data for statistical analysis. We then developed a second protocol to determine a

relative response to a larger number of agonists and antagonists to provide a more comprehensive

profile for comparison with other receptor profiles in the literature.

<u>**Quantitative concentration response to agonists**</u>

Based on preliminary data, NECA, 2-chloroadenosine (2ClAdo) and R-PIA were selected to

determine potency order in a quantitative manner. The concentration response to each agonist was

measured as in the representative experiment shown in figure 3. Oocytes were first stimulated with

sub-maximal 5 µM forskolin, and subsequently brought to a basal state with frog Ringer's (FR).

The forskolin concentration was chosen to avoid desensitization of the adenylyl cyclase/CFTR

pathway since each oocyte was stimulated multiple times by forskolin and adenosine analogs. Next,

the oocytes were exposed to increasing concentrations of NECA (0.01 µM, 0.1 µM, 1 µM & 10

µM), each incremental concentration of the agonist was added once the response to the prior

concentration had peaked (n=3). This sequence was followed by an identical protocol for 2-

chloroadenosine (n=3) and R-PIA (n=6). Variation in the sequence of the agonists tested gave

similar results.

The initial 5µM forskolin response was used as an internal control (comparison of the

agonist response to forskolin) in these studies because of the variability of the expression of CFTR

secondary to imprecision in injection of RNA, inter-oocyte variability in CFTR expression and

inter-frog and intra- frog variability in the quality of the oocytes due to environmental factors. Data

were normalized to the response to 5µM forskolin preceding each concentration response.

Results from the concentration response experiments are summarized in figure 4. The peak
normalized stimulation of the control oocytes (injected with human CFTR cRNA only) at $10\,\mu M$ of
NECA, 2-chloroadenosine and R-PIA are -0.01%, 0.04% and 0.03% respectively (normalized to 5
μM forskolin responses, n=2 each, with mean$\pm$SD of 191 ± 26 nA, 231 ± 27 nA and 295 ± 96 nA,
respectively), showing very poor coupling of any endogenous oocyte adenosine receptor to human
CFTR. The responses of oocytes injected with human CFTR and shark A_0 cRNA, at concentrations
of 0.01 μM of NECA, 2-chloroadenosine and R-PIA, and 0.1 μM NECA and 2-chloroadenosine
were not significantly different from un-injected controls ($p>0.05$). The response to R-PIA at 0.1
μM (55.6%) was greater than NECA and 2-chloroadenosine (2.7% and 4.2% respectively). At an
agonist concentration of 1 μM the differences between the three agonists NECA, 2-chloroadenosine
and R-PIA (13.0%, 35.1% and 196%, respectively) were appreciable and significant ($p<0.05$). The
response at $10\,\mu M$ (66.5%, 151% and 210% respectively) were markedly different ($p<0.05$). The
probability values for determining a significant main effect for each of these observations were
obtained using an ANOVA. These data established the potency order of R-PIA>2- chloro-
adenosine>NECA for the shark A_0 adenosine receptor in *Xenopus laevis* oocytes.
293
Qualitative profile for agonists and antagonists
With the above profile established for the three agonists, a more extensive profile was
ascertained for 8 agonists (NECA, DPMA, CV1808, 2-chloroadenosine, CPA, S-PIA, CGS21680,
R-PIA) and 5 antagonists (CSC, 8PT, DPCPX, CGS15943, PD115199) to allow comparison with
profiles established for the presently cloned adenosine receptors. The analysis was carried out at an
agonist concentration of 1 μM and an antagonist concentration of 10 μM to optimize the analysis.
The protocol required oocyte perifusion with one agonist at 1 μM and substitution with a
second after stabilization of the signal. The responses were judged as either an increase, a decrease
or equivocal. The data were compiled into the agonist profile for the shark A_0 receptor of R-
PIA>S-PIA>CGS21680>CPA>2-chloroadenosine>CV1808=DPMA>NECA (n=7). Varying the
order of agonists produced highly consistent results, in that the profiles conducted in order of
increasing agonist potency were usually no different from profiles conducted in order of decreasing
agonist potency.
The antagonist order of potency was carried out using a similar protocol. Oocytes were
perifused with 1 μM R-PIA to generate a stable stimulated state, followed by perifusion with 1 μM R-
PIA and 10 μM of the various antagonists. The antagonist profile obtained for the shark A_0 receptor was
DPCPX>PD115199>8PT>CSC>CGS15943 (n=5). The relative potency order of agonists- and
antagonists- were determined qualitatively in 10 different oocytes including 186 pairwise comparisons
(please see supplementary tables S3,
https://www.biorxiv.org/content/biorxiv/early/2020/11/02/2020.11.01.363762/DC1/embed/media-1.pdf?download=true),
and 91% (n=169) of all pairwise observations agree with our final conclusion of potency order.
315
Protocol validation using human ADORs
The human A_1, A_{2a}, A_{2b} and A_3 receptors, obtained from the Linden lab at the University
of Virginia, were also examined in our oocyte system using the qualitative protocol to provide
validation for this form of analysis. Although four human ADORs were expressed in the oocyte,
reproducible profiles were obtained for two receptors, the human A_1 (agonist profile, n=3) and
human A_{2b} (agonist profile, n=4, antagonist profile, n=3) (table 1). The human A_1 and A_{2b}
receptor agonist profiles determined in oocytes are in excellent agreement with binding studies

and functional studies (see supplementary table S4,
).
However, the A_{2b} antagonist profile shows an apparent discrepancy in the potency order of
CGS15943 and 8PT. This difference may be attributable to a cross species comparison (rat vs
human). Taken together, this data provides strong support for the validity of our protocols used in
pharmacological characterization of human- and shark- ADORs.
329
Homology model of shark A_0
Given the reasonable sequence homology of shark A_0 between ADORs that have been well-
characterized and for which structures are known (supplementary fig. S11,
),
we conducted exploratory refinements of the A_0 starting homology model against the deposited
structure factors of the human A_{2a} x-ray crystal structure (PDB 2ydv), replacing the coordinates of
A_{2a} with A_0. Refinement was restricted to low-resolution (4.0 Å) to effectively down-weight the
contribution of mis-matched side chain sequences between the two species while providing a data-
driven constraint for backbone atoms and secondary structure. The resulting model exhibited good
protein Ramachandran- and side chain rotameric geometry (supplementary table S5,
)
and had convincing secondary structure for the transmembrane helices and extramembrane
segments (figure 5). Both sequence alignments and the shark A_0 homology model revealed an
abbreviated extracellular loop 2 (ECL2) in ancestral receptors compared to the more specialized
subtypes. While it is not yet known what, if any, functional consequence arises from different
lengths of the ECL2 specifically, more than 30 ancestral receptors in the database have a minimized
ECL2 (supplementary fig. S8). Thus, it would seem that ECL2 of ancestral receptors is at the
minimum length necessary to maintain the structure and/or function of the adenosine receptor
architecture.
349
Conserved Ligand-Receptor Interactions
To explore the extent to which our homology model recapitulates known fundamental
interactions between ligand and receptor, we separately modeled adenosine (ADO), 2-
chloroadenosine and the most potent ligand in this work, R-phenyl-isopropyl adenosine (R-PIA),
into the A_0 binding pocket, conducted refinements and inspected the receptor-ligand interactions.
While we recognize there will be error in the process of homology modeling A_0 in the absence of
an actual structure, potential fundamental interactions between A_0 and ligands are well conserved
in the refined A_0 model compared to human ADOR structures (figure 6 and supplementary table
S6,
).
Among the most important known interactions between receptor and ligand occur between
ADOR and the purine pharmacophore of adenosine- and adenosine analogs. The structures of
human A_{2a}- and A_1 reveal that an aromatic residue (F168 and F171, respectively), protrudes down
into the ligand binding pocket and contribute an intimate π-π stacking interaction with the purine
ring of adenosine (29). Our refined homology model of shark preserves this interaction
convincingly well, with the corresponding A_0 aromatic residue being F179 (figure 6). A second
critical interaction revealed by the structures of A_{2a}- and A_1 is a strong polar interaction, most
likely hydrogen bonding, between an asparagine residue (N253 and N254, respectively) and the N7

nitrogen and the amino group of C6 of the purine ring. The corresponding A_0 asparagine is N267
(figure 6). These two interactions form the fundamental basis for recognizing the adenosine
pharmacophore and are invariant in all ADORs we have examined (see multiple sequence
alignments in supplementary figs. S12 and S13,
https://www.biorxiv.org/content/biorxiv/early/2020/11/02/2020.11.01.363762/DC1/embed/media-1.pdf?download=true). In
totality, nine amino acid residues in close proximity to the adenosine analog in our shark A_0
homology model are highly conserved in all known- and putative ADORs, including the ancestral
branch (see also supplementary table S6).
376
Discussion
Co-expression of shark A_0 and human CFTR
We successfully co-expressed the shark A_0 adenosine receptor with human CFTR in
Xenopus laevis oocytes, which, to our knowledge, represents the first heterologous co-expression of
any adenosine receptor with CFTR in the literature. Oocytes appear to be an ideal system to
characterize shark A_0, since neither frog ADORs nor frog CFTR appears to be active or coupled.
We coupled an adenosine receptor from an elasmobranch shark, a subclass of Chondrichthyes fish
with a temporal range dating back to at least the Devonian period (~419-360 million years ago), to
human CFTR using the endogenous G-proteins and cellular enzymes of the oocyte. Other
investigators have in the past exploited CFTR for co-expression studies, specifically with alpha 2-
adrenergic-, beta 2-adrenergic-, delta-opioid-, and the 5HT1A receptors ((30), (31)), however those
receptors were mammalian in origin, which would not be viewed nearly as long-lived as receptors
identified in the shark. A_0 function in oocytes suggests that the interaction between receptors and
frog G-proteins is maintained despite the ancestral properties and structural differences A_0 has with
respect to more specialized ADORs. This highlights a likely conservation of the interface between
G-proteins and receptors with which they interact, across species and through evolution. Our results
also emphasize the versatility and value of the *Xenopus laevis* expression system for studies of
GPCR molecular evolution.
395
A_0 exhibits unique ADOR pharmacology
We characterized the shark A_0 receptor in *Xenopus* oocytes using a novel protocol, in
contrast to previous studies using binding assays, *in vivo* functional studies and cAMP
measurements. To this end it was necessary to couple the shark A_0 with a reporter target protein, in
this case human CFTR, to permit measurement of expression by electrophysiology. The data from
these studies revealed a unique profile for the shark A_0 receptor. Table 2 summarizes the
comparison of the A_0 receptor to other adenosine receptors with respect to agonist potency.
The shark A_0 differs: 1) from A_1 in its potency order of 2ClAdo and NECA, 2) from A_{2a}
and A_{2b} which have the reverse potency order and 3) from A_3 in which the least potent shark A_0
agonist, NECA, and the most potent shark A_0 agonist, R-PIA, have equal potencies. Therefore, the
shark A_0 appears to have a unique agonist profile judging from the potency order of R-PIA >
2ClAdo > NECA.
Subsequent qualitative studies further established the unique nature of the shark A_0. Table 3
compares the pharmacological profile for the shark A_0 to mammalian A_1, A_{2a}, A_{2b} and A_3
adenosine receptor subtypes. Analysis reveals the A_1 specific agonist R-PIA is the most potent in
the shark, however the A_1 selective agonist CPA is of intermediate potency. While S-PIA and

CGS21680 are more potent in the shark A_0 than in the A_1 receptors. All of the above are
inconsistent with an A_1-like receptor profile for A_0.
The shark A_0 receptor is dissimilar to A_{2a} or A_{2b} with respect to its low potency to the A_2-
preferring agonist NECA. Furthermore, response of A_0 to S-PIA was unexpectedly high
considering that S-PIA is not relatively potent at A_1, A_{2a} or A_{2b} receptors. Interestingly, the A_{2a}
preferring agonist CGS21680 has relatively high potency in the shark A_0 profile, while CV1808,
also A_{2a} preferring, has a much lower potency in the profile. The shark A_0 profile bears little
resemblance to that of A_{2b} receptors especially with respect to the position of CGS21680, NECA,
R-PIA and S-PIA. Therefore, the shark A_0 resembles neither the A_{2a} nor the A_{2b} adenosine
receptors (table 3). Similarly, it is quite different from the known A_3 adenosine receptor profiles
(table 3), with low potency of NECA (intermediate potency in A_3), relatively high potency of
CGS21680 and 5-PIA (lower in the profile for A_3) and the large difference in potency of R-PIA
and NECA, which are of equal potency in the A_3 adenosine receptor profiles.
A comparison of the antagonist profiles yields similar results. Although the shark A_0
antagonist profile is reminiscent of the A_1 antagonist functional profile (table 3), when compared to
binding studies for the rat A_1 (supplementary table S7,
https://www.biorxiv.org/content/biorxiv/early/2020/11/02/2020.11.01.363762/DC1/embed/media-1.pdf?download=true),
the low potency of CGS15943 in the shark A_0 profile is striking. The relative potency of
CGS15943 is also incongruent with the rat A_{2a} and rat A_{2b} profiles (supplementary table S8,
https://www.biorxiv.org/content/biorxiv/early/2020/11/02/2020.11.01.363762/DC1/embed/media-1.pdf?download=true).
Moreover, the potency order of DPCPX>PDl15199 in the shark A_0, is reversed for A_{2a} receptors
and A_{2b} receptors. It is clear that shark A_0 also possess a unique antagonist profile.
In summary the agonist- and antagonist- profiles do not allow the classification of this
receptor into any known receptor subtype, providing functional evidence that A_0 is a unique
ADOR.
437
A_0 is an extant ADOR ancestor
Since the initial characterization of specialized adenosine receptor (ADOR) subtypes ~40
years ago, hundreds of ADORs have been identified. Many examples arise from genomic
sequencing but have not been characterized structurally or functionally. Phylogenetic analysis
shows that examples of specialized ADORs (i.e. A_1, A_{2a}, A_{2b} and A_3) are apparent throughout the
animal kingdom, including fish, amphibians, reptiles, birds and mammals. However, a separate
phylogenetic branch emerges from the analysis that contains ADORs in reptiles, amphibians and
fish, including the shark A_0 ADOR we have characterized here. Shark A_0, and the related
members on the branch, appear to be ancestral in nature. We draw this conclusion based on the
following: 1) amino acid sequence similarity of A_0 is roughly equally different from any specific
specialized subtype (A_1, A_{2a}, A_{2b} and A_3; see supplementary table S9,
https://www.biorxiv.org/content/biorxiv/early/2020/11/02/2020.11.01.363762/DC1/embed/media-1.pdf?download=true); 2)
the pharmacological profile of A_0 is significantly different from the profiles of any subtype; 3) A_0
belongs to a phylogenetic branch that is distinct from the branches formed by the other specialized
subtypes; and 4) ADOR members of the ancestor branch contains structural features unique only to
themselves, such as a minimal ECL2.
A_1, A_{2a}, A_{2b} and A_3 appear to be specialized for several reasons: 1) they have distinct and
consistent pharmacological profiles amongst their respective sub-branches; 2) they prefer to couple
to one G-protein alpha subunit (e.g. inhibitory $G_i\alpha$ or stimulatory $G_s\alpha$ on adenylyl cyclase activity),

and 3) they are abundant in birds and mammals which are thought to arrive later in evolution. The latter point highlights an interesting observation: specialized ADORs are prevalent in mammals which arrived more recently in evolution time-scales, but the ancestral receptors, including olfactory- and non-olfactory ADORs, appear to have been lost in mammals.

Conclusion

We have cloned, expressed and characterized an adenosine receptor (A_0) from the dogfish shark (*Squalus acanthias*) that exhibits novel pharmacology compared to all known subtypes (A_1, A_{2a}, A_{2b} and A_3). A_0 and its closest relatives form a distinct ancestral phylogenetic branch from the known subtypes and do not appear have representation in birds or mammals. All ancestral ADORs appear to have distinguishing structural elements, such as a minimally short extracellular loop 2. Shark A_0 expression is strong in rectal gland where it likely regulates chloride secretion but its expression is weak in brain, allowing us to conclude that is not olfactory in primary function. A non-olfactory ancestral branch of ADORs, including A_0 from elasmobranchs, amphibians, and reptiles is now apparent from the wealth of genomic sequencing data.

In contrast to the recently identified olfactory ADOR (A2c) found exclusively in fish (13), A_0 was cloned from the shark rectal gland, a non-olfactory epithelial organ known for diverse GPCR-mediated regulation of chloride secretion via the CFTR chloride channel (32). Shark CFTR is more than 70% identical to human CFTR, and ADOR regulation of CFTR in sharks and mammals is well described (33,34). Taken together, a long-standing evolutionary functional coupling between ADORs and CFTR-mediated chloride secretion in epithelial tissues is evident, and the elasmobranch A_0 receptor emerges as a non-olfactory extant ancestor of specialized ADOR subtypes that most likely regulates CFTR in the shark.

482   **Acknowledgements**
483   We wish to thank Drs. Stephanie Halene and Kristina Lübbe for their important
484   contributions toward the manuscript. This work was supported by a Howard Hughes Medical Institute
485   Grant to SB, internal UAB funding to SGA, and by NIH grants R01 DK034208, NIEHS P30 ES003828
486   and NSF grant DBI-0453391 to JNF.

488   **Data Availability**
489   We agree to deposit the shark adenosine receptor nucleotide sequence to public repositories
490   (NCBI) or make it available upon request.

491
Figure 1. Phylogenetic relationships of ADOR taxa. Evolutionary history is inferred using the Neighbor-Joining method (35). The optimal tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the tree. Distances were calculated using the Poisson correction method (36) and are in the units of the number of amino acids substitutions per site. A_1, A_{2a}, A_{2b}, A_3 and olfactory (A_{2c}) ADORs shown in Wakisaka, et al. 2017 are shown using the same names as in the paper. The first characterized A2c (Zebrafish, KY938008) by Wakisaka et al. is also shown in the figure. Additional putative ADORs found in this work are shown by species and Accession. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test using 500 replicates (Felsenstein 1985) are shown at selected branchpoints. Analyses were conducted using MEGA7 (Kumar, et al, 2016) using only full-length sequences for all receptors.

Figure 2. Current-Voltage (I-V) relationship illustrating the effects of forskolin, 2-chloroadenosine (a
deaminase-resistant form of adenosine) and 2-chloroadenosine & DPCPX (a known adenosine receptor
antagonist) in a representative oocyte co-expressing the shark A_0 receptor and human CFTR Cl$^-$ channel.
Panel A: selected single I-V plots for basal, forskolin, 2ClAdo and 2ClAdo+DPCPX. Panel B:
Corresponding total oocyte current at -60 mV clamping voltage for the plots in panel A. Panel C: Reversal
potentials (mean ± SD) were obtained for basal (-38.0 ± 3.8, n=5), forskolin ("For", -29.5 ± 0.74, n=3),
2ClAdo (-30.5 ± 0.28, n=4) and 2ClAdo+DPCPX (-39.4 ± 4.8, n=3).
510
511

512
Figure 3. Representative tracing for the determination of the concentration response profile for the shark
A_0 adenosine receptor. A total of four oocytes were used for the determination of the concentration
response reported in the main text. Three oocytes, including results from one representative shown in the
figure, were exposed to multiple sequences of incremental concentrations of NECA, 2-chloroadenosine
and R-PIA (not shown in the figure), while the fourth was used for the study of R-PIA exclusively.
Following any stimulation of the oocytes (with For or adenosine analogs), oocytes were returned to basal
levels by perifusion with FR.
520
521

Figure 4. Concentration response of the shark A_0 adenosine receptor in both A_0- and CFTR- co-injected
oocytes as well as CFTR only-injected oocytes (labeled "Control" in the figure). The controls were conducted
in three oocytes, with each agonist tested in each of the three oocytes (see asterisks (*) on the x-axis). The
experimental group was tested in four oocytes (NECA, n=3; 2-chloroadenosine, n=3; R-PIA, n=6; and are
highlighted in open squares, open triangles and open triangles, respectively). The y-axis is expressed in terms
of the percent response to 5 µM forskolin in total oocyte current (nA).

532
533
Figure 5. Overall view of the shark A_0 homology model compared to human A_{2a} and A_1 receptors (PDB
codes 2ydv and 6d9h). The approximate position of the hydrophobic stretch of the lipid bilayer is depicted
with lines separated by ~ 35 Å using conserved aromatic side chains as fiducials (27). A striking feature
of shark A_0 compared to the more specialized ADORs is an abbreviated extracellular loop 2 (ECL2,
labeled) and the relatively low profile that the A_0 receptor presents to the extracellular milieu. Multiple
sequence alignment of all known- and putative receptor orthologs (supplementary figs. S9 and S10)
reveals that an abbreviated ECL2 appears to be unique feature of ancestral ADORs.
541
542

543
544
Figure 6. Binding pockets for adenosine and adenosine agonists. Panel A shows a potential binding mode
of R-phenyl-isopropyl adenosine (R-PIA) as it appears in our refined homology model of the shark A_0
receptor. The purpose is not to highlight precise interactions, bonds or geometry, but to indicate that the
major known adenosine-binding ligands are conserved in the shark A_0 ancestor receptor and are well
positioned to potentially interact with ligands. For comparison, the location of N-ethyl-carboxyl
adenosine (NECA) binding in the 2.6 Å x-ray structure of human A_{2a} receptor (PDB code 2ydv) is shown
in panel B. Panel C shows the location of adenosine (ADO) binding in the 3.6 Å cryo-EM structure of
the human A_1 receptor (PDB code 6d9h).
553
554

References

References

1.   Burnstock, G. (1978) *A basis for distinguishing two types of purinergic receptor.*, Raven Press,
New York

2.   Olah, M. E. (1997) Identification of A2a adenosine receptor domains involved in selective
coupling to Gs. Analysis of chimeric A1/A2a adenosine receptors. *J Biol Chem* **272**, 337-344

3.   Sweeney, M. I., and Dolphin, A. C. (1995) Adenosine A1 agonists and the Ca2+ channel agonist
bay K 8644 produce a synergistic stimulation of the GTPase activity of Go in rat frontal cortical
membranes. *J Neurochem* **64**, 2034-2042

4.   Palmer, T. M., and Stiles, G. L. (1995) Adenosine receptors. *Neuropharmacology* **34**, 683-694

5.   Jockers, R., Linder, M. E., Hohenegger, M., Nanoff, C., Bertin, B., Strosberg, A. D., Marullo, S.,
and Freissmuth, M. (1994) Species difference in the G protein selectivity of the human and
bovine A1-adenosine receptor. *J Biol Chem* **269**, 32077-32084

6.   Marala, R. B., and Mustafa, S. J. (1993) Direct evidence for the coupling of A2-adenosine
receptor to stimulatory guanine nucleotide-binding-protein in bovine brain striatum. *J*
Pharmacol Exp Ther **266**, 294-300

7.   Freholm, B. B. e. a. (1994) Nomenclature and Classification of Purinoceptors. *Pharmacological*
Reviews **46**, 143-156

8.   Burns, R. F., Lu, G. H., and Pugsley, T. A. (1987) Adenosine Receptor Subtypes: Binding Studies.
in *Topics and Prespectives in Adenosine Research*, Springer-Verlag, Berlin, Heidelberg. pp 59-73

9.   Kelley, G., Aassar, O., and Forrest, J. J. (1991) Endogenous adenosine is an autocoid feedback
inhibitor f chloride transport in the shark rectal gland. *Journal of clinical investigation* **88**, 1933-
1939

10. Forrest, J. N. J. (1996) Cellular and molecular biology of chloride secretion in the shark rectal
gland: regulation by adenosine receptors. [Review]. *Kidney International* **49(6)**, 1557-1562

11. Forrest, J. N., Jr., Aller, S. G., Wood, S. J., Ratner, M. A., Forrest, J. K., and Kelley, G. G. (1997)
Cadmium disrupts the signal transduction pathway of both inhibitory and stimulatory receptors
regulating chloride secretion in the shark rectal gland. *J Exp Zool* **279**, 530-536

12. Marshall, J., Martin, K., Picciotto, M., HOCkfield, S., Nairn, A., and Kaczmarek, L. (1991)
Identification and localization of a dogfish homolog of human cyctic fibrosis transmembrane
conductance regulator. *The journal of biological chemistry* **266**, 22749-22754

13. Wakisaka, N., Miyasaka, N., Koide, T., Masuda, M., Hiraki-Kajiyama, T., and Yoshihara, Y. (2017)
An Adenosine Receptor for Olfaction in Fish. *Curr Biol* **27**, 1437-1447 e1434

14. Thompson, J. D., Higgins, D. G., and Gibson, T. J. (1994) CLUSTAL W: improving the sensitivity of
progressive multiple sequence alignment through sequence weighting, position-specific gap
penalties and weight matrix choice. *Nucleic Acids Res* **22**, 4673-4680

15. Madeira, F., Park, Y. M., Lee, J., Buso, N., Gur, T., Madhusoodanan, N., Basutkar, P., Tivey, A. R.
N., Potter, S. C., Finn, R. D., and Lopez, R. (2019) The EMBL-EBI search and sequence analysis
tools APIs in 2019. *Nucleic Acids Res* **47**, W636-W641

16. Rice, P., Longden, I., and Bleasby, A. (2000) EMBOSS: the European Molecular Biology Open
Software Suite. *Trends Genet* **16**, 276-277

17. Nordstrom, K. J., Sallman Almen, M., Edstam, M. M., Fredriksson, R., and Schioth, H. B. (2011)
Independent HHsearch, Needleman--Wunsch-based, and motif analyses reveal the overall
hierarchy for most of the G protein-coupled receptor families. *Mol Biol Evol* **28**, 2471-2480

18. Kumar, S., Stecher, G., and Tamura, K. (2016) MEGA7: Molecular Evolutionary Genetics Analysis
Version 7.0 for Bigger Datasets. *Mol Biol Evol* **33**, 1870-1874

19.   Lebon, G., Warne, T., Edwards, P. C., Bennett, K., Langmead, C. J., Leslie, A. G., and Tate, C. G.
(2011) Agonist-bound adenosine A2A receptor structures reveal common features of GPCR
activation. *Nature* **474**, 521-525

20.   Draper-Joyce, C. J., Khoshouei, M., Thal, D. M., Liang, Y. L., Nguyen, A. T. N., Furness, S. G. B.,
Venugopal, H., Baltos, J. A., Plitzko, J. M., Danev, R., Baumeister, W., May, L. T., Wootten, D.,
Sexton, P. M., Glukhova, A., and Christopoulos, A. (2018) Structure of the adenosine-bound
human adenosine A1 receptor-Gi complex. *Nature* **558**, 559-563

21.   Garcia-Nafria, J., Lee, Y., Bai, X., Carpenter, B., and Tate, C. G. (2018) Cryo-EM structure of the
adenosine A2A receptor coupled to an engineered heterotrimeric G protein. *Elife* **7**

22.   Lebon, G., Edwards, P. C., Leslie, A. G., and Tate, C. G. (2015) Molecular Determinants of
CGS21680 Binding to the Human Adenosine A2A Receptor. *Mol Pharmacol* **87**, 907-915

23.   Sayers, E. (2017) Database Resources of the National Center for biotechnology Information.
Nucleic Acids Res **45**, D12-D17

24.   Emsley, P., Lohkamp, B., Scott, W. G., and Cowtan, K. (2010) Features and development of Coot.
Acta Crystallogr D Biol Crystallogr **66**, 486-501

25.   Adams, P. D., Afonine, P. V., Bunkoczi, G., Chen, V. B., Davis, I. W., Echols, N., Headd, J. J., Hung,
L. W., Kapral, G. J., Grosse-Kunstleve, R. W., McCoy, A. J., Moriarty, N. W., Oeffner, R., Read, R.
J., Richardson, D. C., Richardson, J. S., Terwilliger, T. C., and Zwart, P. H. (2010) PHENIX: a
comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr D
Biol Crystallogr* **66**, 213-221

26.   Weininger, D. (1988) SMILES, a chemical language and information system. *J Chem Inform
Computer Sci* **28**, 31-36

27.   de Planque, M. R., Kruijtzer, J. A., Liskamp, R. M., Marsh, D., Greathouse, D. V., Koeppe, R. E.,
2nd, de Kruijff, B., and Killian, J. A. (1999) Different membrane anchoring positions of
tryptophan and lysine in synthetic transmembrane alpha-helical peptides. *J Biol Chem* **274**,
20839-20846

28.   Devor, D. C., Forrest, J. N., Jr., Suggs, W. K., and Frizzell, R. A. (1995) cAMP-activated Cl-
channels in primary cultures of spiny dogfish (Squalus acanthias) rectal gland. *Am J Physiol* **268**,
C70-79

29.   Xu, F., Wu, H., Katritch, V., Han, G. W., Jacobson, K. A., Gao, Z. G., Cherezov, V., and Stevens, R.
C. (2011) Structure of an agonist-bound human A2A adenosine receptor. *Science* **332**, 322-327

30.   Birnbaum, A. K., Wotta, D. R., Law, P. Y., and Wilcox, G. L. (1995) Functional expression of
adrenergic and opioid receptors in Xenopus oocytes: interaction between alpha 2- and beta 2-
adrenergic receptors. *Brain Res Mol Brain Res* **28**, 72-80

31.   Uezono, Y., Bradley, J., Min, C., McCarty, N. A., Quick, M., Riordan, J. R., Chavkin, C., Zinn, K.,
Lester, H. A., and Davidson, N. (1993) Receptors that couple to 2 classes of G proteins increase
cAMP and activate CFTR expressed in Xenopus oocytes. *Receptors Channels* **1**, 233-241

32.   Forrest, J. N., Jr. (2016) The Shark Rectal Gland Model: A Champion of Receptor Mediated
Chloride Secretion through Cftr. *Trans Am Clin Climatol Assoc* **127**, 162-175

33.   Clancy, J. P., Ruiz, F. E., and Sorscher, E. J. (1999) Adenosine and its nucleotides activate wild-
type and R117H CFTR through an A2B receptor-coupled pathway. *Am J Physiol* **276**, C361-369

34.   Kelley, G. G., Poeschla, E. M., Barron, H. V., and Forrest, J. N., Jr. (1990) A1 adenosine receptors
inhibit chloride transport in the shark rectal gland. Dissociation of inhibition and cyclic AMP. *J
Clin Invest* **85**, 1629-1636

35.   Saitou, N., and Nei, M. (1987) The neighbor-joining method: a new method for reconstructing
phylogenetic trees. *Mol Biol Evol* **4**, 406-425

36.   Zuckerkandl, E. a. P., L. (1965) Evolutionary divergence and convergence in proteins. in *Evolving*
647   *Genes and Proteins*, Academic Press, New York. pp 97-166
648