Using Community Science to Reveal the Global Chemogeography of River Metabolomes

Vanessa A Garayburu-Caruso[1,†], Robert E Danczak[1,†], James C Stegen[1], Lupita Renteria[1], Marcy McCall[1], Amy E Goldman[1], Rosalie K Chu[2], Jason Toyoda[2], Charles T. Resch[1], Joshua M Torgeson[1], Jacqueline Wells[3], Sarah Fansler[1], Swatantar Kumar[1] and Emily B Graham[1,*].

[1] Pacific Northwest National Laboratory, Richland, WA

[2] Environmental Molecular Sciences Laboratory, Richland, WA

[3] Oregon State University, Corvallis, OR

[†] Denotes equal contribution

[*] Correspondence: emily.graham@pnnl.gov

Abstract: River corridor metabolomes reflect organic matter (OM) processing that drives aquatic biogeochemical cycles. Recent work highlights the power of ultrahigh-resolution mass spectrometry for understanding metabolome composition and river corridor metabolism. Yet, there have been no studies on the global chemogeography of surface water and sediment metabolomes using ultrahigh-resolution techniques. Here, we describe a community science effort from the Worldwide Hydrobiogeochemistry Observation Network for Dynamic River Systems (WHONDRS) consortium to characterize global metabolomes in surface water and sediment that span multiple stream orders and biomes. We describe the distribution of key aspects of metabolomes including elemental groups, chemical classes, indices, and inferred biochemical transformations. We show that metabolomes significantly differ across surface water and sediment, and that surface water metabolomes are more rich and variable. We also use inferred biochemical transformations to identify core metabolic processes shared among surface water and sediment. Finally, we observe significant spatial variation in sediment metabolites between rivers in the eastern and western portions of the continental United States. Our work not only provides a basis for understanding

27 global patterns in river corridor biogeochemical cycles but also demonstrates that community

28 science endeavors can enable global research projects that are unfeasible with traditional research

29 models.

30 **Keywords:** environmental metabolomics; river corridor; sediment organic matter; WHONDRS;

31 CONUS; carbon character; dissolved organic matter

---

1. Introduction

Organic matter (OM) transformations in aquatic ecosystems are a critical source of uncertainty

in global biogeochemical cycles [1–4]. More than half of OM inputs to freshwater ecosystems are

metabolized before reaching the oceans [1,2,4]; yet while several studies have focused on quantifying

OM uptake and export rates [1,5,6], the processes driving river corridor OM transformations across

spatial scales remain poorly understood.

River corridor OM pools contain an extensive variety of molecules that are both produced and

metabolized by microorganisms, processes that are reflected in the composition of sediment and

surface water metabolomes [2,7,8]. Metabolic transformations of OM in freshwater ecosystems have

been traditionally estimated by a combination of laboratory incubations and in-stream tracer

additions [9–12]. However, results from incubation experiments are challenging to scale beyond

laboratory conditions [9,10] and in-stream tracer processing often does not reflect ambient

biogeochemical processes, as the naturally-occurring metabolome is more chemically diverse than

the tracer added to the stream [11,12]. Several studies have shown that OM pool composition can

influence microbial activity and highlight complexities in the metabolic processes that determine OM

transformations [13–18]. Consequently, determining mechanisms underlying river corridor

metabolome composition at a large scale remains challenging.

Environmental metabolomics on field-derived samples, enabled by ultrahigh-resolution mass

spectrometry, has allowed us to reveal connections between OM character, reactivity, and

biochemical transformations within and across river ecosystems [15,17–23]. These advances have

vastly improved our understanding of the mechanisms governing OM bioavailability and

biochemical transformations at a global scale. For instance, previous studies have used ultrahigh-

resolution metabolomics from river water across different climatic regions, to find common

compositional features that would inform global carbon dynamics [24] and to investigate

environmental drivers affecting OM composition, bioavailability, and transport of OM [25]. In

addition, recent studies show that OM thermodynamics influence aerobic respiration under carbon-

limited scenarios [16], that biogeochemical hotspots are influenced by OM nitrogen content [17], and

that hyporheic zone mixing induces OM metabolism via a priming effect [15]. These detailed

metabolome characterizations have the potential to enable global-scale inferences about watershed

features (e.g., vegetation, lithology, hydrology, microbiology, climate) that govern the reactivity and

fate of OM across river corridors [23,26]. In turn, metabolomics can enhance our predictive

capabilities of global river corridor biogeochemical cycles by helping to improve the representation

of biochemical mechanisms in numerical models, such as reactive transport codes [27,28]. For

example, an emerging substrate-explicit model uses thermodynamic theory to explicitly account for

the chemical composition of all metabolites in OM pools to improve the predictive capacity of

biogeochemical models [28].

Characterizing metabolomes across global spatiotemporal scales requires a way to collect

multiple data types across diverse locations in such a way that they can be analyzed together. This

goal can be facilitated by a framework that requires studies to **I**ntegrate biological, physical, and

chemical processes across scales; **C**oordinate with consistent methods; are **O**pen across the research

lifecycle; and **N**etwork with global collaborators to reduce the burden on a single team (ICON)

[29,30]. When ICON principles are applied, they allow for distributed sampling in ways that have

historically been difficult to achieve.

The Worldwide Hydrobiogeochemistry Observation Network for Dynamic River Systems

(WHONDRS) is a global consortium of researchers based out of Pacific Northwest National

Laboratory that uses an ICON-based approach to understand coupled hydrologic, biogeochemical,

and microbial functions in river corridors [23]. ICON principles allow WHONDRS to collect open,

globally distributed data through collaboration with the scientific community. The WHONDRS

consortium designs sampling campaigns that target specific spatial and temporal scales, modifies its

approach based on community input, and then sends free sampling kits to collaborators. All

WHONDRS data are openly accessible through environmental Systems Science Data Infrastructure

for a Virtual Ecosystem (ESS-DIVE-https://data.ess-dive.lbl.gov/) and the National Center for

Biotechnology Information (NCBI), and the WHONDRS consortium ascribes to FAIR data principles

(findable, accessible, interoperable, reusable) [31]. This approach enables WHONDRS to collect,

analyze, and distribute ultrahigh resolution metabolomic data to the global scientific community.

Here, we describe a community science effort conducted by the WHONDRS consortium during July-

August 2019 that used Fourier-transform ion cyclotron resonance mass spectrometry (FTICR-MS) to

characterize metabolomes in global surface water and sediment spanning a range of biomes (e.g.,

desert-like in the Columbia Plateau, subtropical in southern Florida, temperate forests in the Mid-

Atlantic) and stream orders [32]. We describe key metabolome characteristics of surface water and

sediment and explore spatial variation of these characteristics within the United States. We focus on

central aspects of metabolomes including assigned elemental groups, chemical classes, descriptor

indices, and biochemical transformations. This paper provides a benchmark for studying integrated

surface water and sediment river corridor metabolomes and highlights the need to engage a wider

scientific community in order to expand the reach and impact of scientific advancements.

2. Results and Discussion

2.1. Surface water metabolome is more unsaturated, aromatic, oxidized, rich, and variable than sediment

metabolome

In order to assess patterns in global metabolome composition, we derived a number of

descriptive metrics that summarize FTICR-MS metabolomic profiles. Specifically, we compared

double-bond equivalents (DBE), modified aromaticity index (AI_{Mod}), nominal oxidation state of

carbon (NOSC), inferred chemical classes (e.g., lignin-like, protein-like) and elemental groups (e.g.,

CHO, CHON, CHOSP) of surface water and sediment metabolomes. The double-bond equivalents

metric (DBE) describes the degree of chemical unsaturation of bonds in a particular metabolite [33,34],

AI_{Mod} quantifies the degree of aromaticity (i.e., ring-like shape) of a metabolite [33–35], and NOSC

indicates the energy required to oxidize different metabolomes [36]. High values of AI_{Mod} can denote

the existence of either aromatic ($AI_{Mod} > 0.5$) or condensed aromatic structures ($AI_{Mod} \geq 0.67$), and high

DBE indicates more saturated compounds. NOSC is inversely correlated with the Gibbs free energy

of carbon oxidation. Higher NOSC corresponds to metabolites that are more oxidized and

thermodynamically favorable [15–18,36,37]. Chemical class assignments for each metabolite were

predicted using oxygen-to-carbon and hydrogen-to-carbon ratios (i.e., Van Krevelen classes [38]).

Finally, we used the molecular formula assigned to each metabolite to describe relative abundance

of different heteroatom combinations associated with CHO groups (i.e., differences in -N, -S, -P). We

then compared metrics across all metabolites found in any surface water sample vs. all metabolites

found in any sediment sample. All analyses were conducted only on FTICR-MS peaks that were able

to be assigned a molecular formula. Other metrics describing metabolome composition are reported

in the SI (Table S1).

Figure 1. Density plots comparing the properties of all molecular formulas found in surface water and sediment samples. AI_{Mod} (modified aromaticity index) is a measure of the potential ring-like structure in a given molecular formula. DBE (double-bond equivalents) is an approximation of potential unsaturation. NOSC (nominal oxidation state of carbon) represents the degree of oxidation/reduction of a given molecular formula. Significance values obtained via a two-sided Mann Whitney U test to compare sample type distributions are denoted in the upper right corner of each panel.

Surface water metabolomes were composed of comparatively more unsaturated and aromatic compounds with higher nominal oxidation state than sediment. This was denoted by significantly higher AI_{Mod}, DBE, and NOSC than sediment metabolomes (**Figure 1,** p-value < 0.001). In addition, we observed higher relative abundances of lignin-like, tannin-like, and condensed-hydrocarbon-like metabolites in surface water versus sediment (**Figure 2,** all p-values < 0.001). These classes of metabolites are characteristic of terrestrial OM [39], and their prevalence in surface water

metabolomes indicate a larger contribution of terrestrial OM in surface water relative to sediment.

This may also indicate greater contributions of microbially processed OM in sediment, as has been

observed previously in comparisons between surface water and hyporheic zone porewater [40].

The higher relative abundance of unsaturated and aromatic metabolites in surface water

contrasts previous studies that have observed that these compounds are more common in sediment

porewater than in lake surface water or aquifer recharge water [41,42]. These studies inferred low

physical, chemical, and/or biological transformation of porewater sediment-associated OM. This

deviation might be connected to the systems studied. For example, Pracht et al. [41] examined a

system where sediment OM was protected due to physical and/or chemical constraints such as

mineral sorption and hydrophobic encapsulation [41]. We studied rivers where the shallow benthic

layer and the hyporheic zone are known to enhance biogeochemical reactions [43–47]. In turn, we

hypothesize that very high rates of biological activity in riverbed sediment [48,49] could be

responsible for lower AI_{Mod}, DBE, and NOSC values of sediment metabolomes relative to surface

water, in contrast to previous work in potentially less active lake and aquifer systems.

Figure 2. Box plots comparing the relative abundances of metabolites belonging to a specific elemental

groups (a) and chemical classes (b) between sediment and surface water. As in Figure 1, these values

were obtained from all metabolites assigned molecular formulas in sediment and surface water.

Significance values were obtained via a two-sided Mann Whitney U test ($0.05 > * > 0.01 > ** > 0.001 >$

***).

In addition, the relative abundances of lipid-like and protein-like metabolites were significantly

higher in sediment than in surface water (p-value < 0.001). More lipid-like compounds in sediment

could reflect higher microbial biomass [50] and further supports our inference that sediment

metabolomes were influenced by microbial processes to a greater extent than surface water

metabolomes. This highlights the key role played by riverbed sediment and associated hyporheic

zones in river corridor biogeochemistry that likely influences global elemental cycles but is not

captured in current Earth system models.

Conversely, elemental groups of metabolites were similar across surface water and sediment.

The median abundance of each elemental group did not vary more than 5% between the two

environments, except for CHO (~9%) and CHONSP (~0.2%, not statistically significant) groups. Bulk

similarities in elemental groups, in contrast to chemical classes, could indicate that the presence or

absence of heteroatoms alone is insufficient to distinguish metabolomes and that elemental

stoichiometry of the entire metabolite (the basis for chemical class assignment) may be more

important for distinguishing metabolomes. This is important because the elemental stoichiometry of

metabolites can mechanistically connect OM thermodynamics to biogeochemical reactions and rates

[28].

Metabolites found in surface water were distinct from and showed more among-sample

variation than those in sediment. To evaluate compositional differences, we conducted a principal

component analysis (PCA) and a beta-dispersion analysis (**Figure 3**). The PCA, in conjunction with a

PERMANOVA comparison, indicated that surface water and sediment metabolomes significantly

diverged in composition (p-value < 0.001). The beta-dispersion analysis further indicated that surface

water metabolomes were more dispersed in multivariate space than sediment metabolomes (**Figure**

3; p-value < 0.001). Additionally, surface water metabolomes had higher richness (i.e., more peaks

with assigned formulas detected on average) than sediment metabolomes (**Figure 1**). These patterns

indicate that metabolomes in surface water and sediment may be shaped by distinct processes that

likely span differences in inputs, rates of microbial activity, and abiotic constraints.

We hypothesize that higher richness and greater among-sample variation in surface water

metabolomes could reflect more heterogeneous environmental pressures. For example, we sampled

across a broad range of latitudes and stream orders that likely led to among-site variation in light

exposure (Table S2 [32]). This may, in turn, lead to variation in surface water temperatures and

surface water metabolite photodegradation, thereby increasing metabolome variability and richness

[51]. Hydrology could also contribute to metabolome richness in surface water as precipitation events

and associated runoff transport large amounts of terrestrial OM into rivers [52]. For example, precipitation has been shown to increase aromatic OM and decrease more labile OM in surface water [53–55]. We hypothesize that more immediate connectivity between surface water and terrestrial systems, relative to connectivity between sediment and terrestrial systems, is at least partially responsible for greater variation and higher richness in surface water metabolomes.

In addition, lower sediment metabolome variation and richness could be due to comparatively higher rates of microbial activity in sediment that degrade polymeric OM into a limited set of less chemically complex metabolites. This would result in a reduction in the number of distinct metabolites present by collapsing a diverse pool of OM into microbial exudates. Sediment metabolomes may also be constrained by interactions with sediment mineral surfaces, especially considering that we studied only the water extractable metabolome. This subset of the full sediment metabolome may inherently be composed of a restricted suite of metabolites [56], leading to lower among-sample variation and lower richness. We nonetheless hypothesize that among-site variation in mineralogy could contribute to some of the observed sediment metabolome variation. The WHONDRS consortium is currently generating the mineralogy data to test this hypothesis.

Together, the observations presented in this section indicate that there are significant differences across global surface water and sediment metabolomes, where surface water metabolomes are more unsaturated, aromatic, oxidized, rich, and variable. These characteristics suggest that surface water metabolomes are more dynamic due to a variety of watershed and river corridor processes (discussed above), while sediment metabolomes may be more stable integrators of localized processes (e.g., mineral interactions and microbial processing of OM).

Figure 3. Principal component analysis (PCA) of the molecular formula data (a). Differences between

surface water and sediment metabolomes were significant per a Euclidean distance-based

PERMANOVA (p-value < 0.001). The degree of among-sample variation was evaluated by

quantifying beta-dispersion. Surface water had higher beta-dispersion per a two-sided Mann-

Whitney U test (p-value < 0.001) (b).

2.2. Phosphorous, nitrogen and sulfur-containing transformations vary across surface water and sediment

metabolomes

We evaluated how potential reactions in metabolomes varied across the globe by inferring

biochemical transformations as per Bailey et al. [50], Kaling et al. [57], Moritz et al. [58], Graham et

al. [17,18], Garayburu-Caruso et al. [16], Danczak et al. [26], and Stegen et al. [15]. This method

leverages the ultrahigh-resolution of FTICR-MS to compare mass differences between detected peaks

to a database of common biochemical transformations. Identified biochemical transformations

provide information regarding the frequency at which a specific molecule could have been gained or

lost during metabolism. Resulting transformation counts can then be separated based upon their

chemical properties to study the potential role of the molecule gained or loss in metabolome

composition. Unlike the analyses described in the previous section, where formula assignments of

metabolites is necessary, this method allows for the incorporation of all detected metabolites into

downstream analyses.

We observed that biochemical transformations involving molecules containing nitrogen (N),

sulfur (S), or phosphorous (P) exhibited divergent patterns between surface water and sediment

metabolomes (**Figure 4**). Specifically, surface water had a significantly higher relative abundance of

N-containing transformations while sediment metabolomes had more S- and P-containing

transformations (p-value < 0.001 in all cases). This contrast between surface water and sediment may

occur due to variation in nutrient requirements within the water column and sediment, as

biochemical transformations have been inferred to reflect nutrient limitations in other systems. For

example, Garayburu-Caruso et al. [16] found an increase in N-containing transformations under

nutrient limited conditions, but only when N-containing OM was introduced. More N-based

transformations in surface water as compared to sediment could therefore reflect microbial N mining

in surface water through preferential decomposition of N-containing OM [59,60]. In addition, higher

abundance of P- and S-containing biochemical transformations in sediment further suggest that

microbial metabolism is limited by different factors between surface water and sediment

environments, potentially associated with nutrient assimilation processes [61,62].

Figure 4. Boxplots displaying the patterns of CHO-only transformations (a), N-containing transformations (b), S-containing transformations (c), and P-containing transformations (d). FDR-corrected two-sided Mann-Whitney U test p-values are provided in either the top right or the bottom left corner of panels with significant comparisons.

Interestingly, the higher abundance of N-containing transformations in surface water observed in this study is contrary to past work, where porewater had a higher relative abundance of N-containing transformations than surface water [15,26]. Given that these studies were collected from the Pacific Northwest region of the United States (e.g., eastern Washington and Oregon), this highlights the necessity to expand research beyond individual test systems or geographic regions and emphasizes the utility of global studies through efforts like WHONDRS.

In contrast, biochemical transformations that did not involve N-, S-, or P-containing molecules

were not significantly different between surface water and sediment metabolomes. Similar patterns

have been observed in other studies [26]. These results indicate the presence of ubiquitous

biochemical transformations that occur in both surface water and sediment (**Figure 4**). Based on these

results we hypothesize that N-, S-, and P-containing transformations may have a stronger

dependency than CHO-only transformations on nutrient status. That is, changes in nutrient

availability across surface water and sediment environments may drive shifts in N-, S-, and/or P-

containing transformations but not influence transformations that do not involve these nutrients.

Additional data on variation in nutrient limitation and availability will be required to test this

hypothesis.

2.3. Sediment metabolomes are more spatially variable than surface water metabolomes

Because most of our sampling locations were in the continental United States (CONUS), we used

CONUS data to resolve potential spatial patterns in metabolomes (**Figures 5 and 6**). In order to

uncover site-by-site metabolomic variation, we calculated the mean value for each derived metric

(e.g., AI_{Mod}, NOSC, DBE) and calculated the relative abundance of elemental groups and chemical

classes (e.g., CHO and lignin-like) for each sample.

Overall, differences between the mean properties of CONUS surface water and sediment

metabolomes were generally consistent with differences between global surface water and sediment

metabolomes reported in **Figure 1**. For instance, surface water metabolomes displayed higher AI_{Mod},

DBE, and NOSC than sediment metabolomes (**Figure 5,** p-value < 0.001 for all, Table S3). We also

observed similar patterns in both the relative abundances of specific elemental groups (**Figure 6,** p-

value < 0.001 for all, Table S3) and chemical classes (p-value < 0.001 for all, Table S3 and File S1). In

order to expand our analyses, we investigated spatial patterns in individual metabolomic features

across the CONUS by comparing sites that were east (hereafter "East", surface water n = 34, sediment

n = 33) vs. west (hereafter "West", surface water n = 45, sediment n = 38) of the Mississippi River.

Figure 5. Maps of the United States revealing the spatial variability of average NOSC in sediment (a)

and surface water (b) metabolomes. East vs. West spatial patterns for various derived metrics are

displayed in panel (c). Statistically significant differences identified via a two-sided Mann Whitney U

test are indicated by p-values listed in each comparison. The red-dashed line represents the longitude

of the Mississippi River at St. Louis, MO, USA -- the dividing line between East and West samples.

In general, we observed greater spatial patterning in sediment metabolomes than in surface

water metabolomes. Average NOSC and AI_{Mod} values of sediment metabolomes were higher in the

East than the West (p-value < 0.001 for both). Metabolites containing CHONP, CHONS, or CHOSP

constituted a significantly lower relative abundance of metabolomes in the East sediment, relative to

the West (p-value < 0.001 for all). Lastly, lignin- and tannin-like metabolites constituted a higher

relative abundance of sediment metabolomes in the East, while protein-, condensed hydrocarbon-,

and unsaturated hydrocarbon-like metabolites were more abundant in the West (p-value < 0.001 for

all, Table S3 and File S1).

In contrast to spatial patterns in sediment metabolomes, surface water showed less spatial

structure with only CHO and CHOS metabolites showing significant shifts between East and West

(p-value < 0.001 for both). CHO and CHOS metabolites comprised higher and lower relative

abundances in the West than in the East, respectively (**Figure 6**). No spatial patterns were observed

in NOSC (p-value = 0.4, Table S3), DBE (p-value = 0.07, Table S3), or AI_{Mod} (p-value = 0.93, Table S3) in

surface water (**Figure 5**). Results from the remainder of molecular indices, elemental groups, and

chemical class comparisons are shown in Table S3 and File S1.

Figure 6. Maps of the United States revealing the spatial variability of the relative abundance of

metabolites containing only CHO with sediment (a) and surface water (b). East vs. West spatial

patterns for different elemental groups relative abundances are displayed in panel (c). Statistically

significant differences identified via a two-sided Mann Whitney U test are indicated by p-values listed

in each comparison. The red-dashed line represents the longitude of the Mississippi River at St. Louis,

MO, USA -- the dividing line between East and West samples.

Together, these patterns suggest that spatial differences in the metabolic processes driving OM

cycling in the East versus the West have a stronger influence on sediment metabolomes than surface

water metabolomes. While our non-spatial analyses showed greater among-sample and among-site

variability in surface water metabolomes (**Figure 3**), the lack of spatial structure across the CONUS

suggests that this variability is not driven by factors that are spatially structured at the continental

scale. Instead, we hypothesize that surface water metabolomes are more temporally variable due to

fluctuating inputs from precipitation events. To more directly evaluate spatial structure in surface

water metabolomes it is likely necessary to control for precipitation history and hydrologic

connectivity to terrestrial systems. These inferences are supported by previous studies addressing

spatial dissolved OM chemography dynamics showing that longitudinal patterns of dissolved OM

in surface water are sensitive to hydrologic events [63–65]. However, there is a complex interaction

between hydrology and space, as the sources and quality of OM from different regions may respond

differently to hydrological variation [66].

The contrasting patterns in surface water and sediment metabolome characteristics across the

East-West gradient could be the result of many factors, including vegetation cover [67], underlying

lithology [68], photoreactivity [51], climate and precipitation regime [68], and/or microbial

metabolism [69]. Additional data and analyses will be required to disentangle the relative

contributions of these potential drivers. Pursuing this knowledge is important for explaining and

ultimately predicting OM transformations. In turn, representing these processes and their impacts on

biogeochemical cycles in processed-based models has the potential to improve the accuracy of

biogeochemical predictions across the globe.

3. Materials and Methods

3.1 WHONDRS Summer 2019 Sampling Campaign

In July and August 2019, the WHONDRS consortium initiated a study of global river corridors

to evaluate interactions between ecosystem features, microbial communities, and metabolomes in

surface water and shallow sediment. To design the study, the WHONDRS consortium held multiple

webinars with collaborators who volunteered to collect samples. The webinars allowed for

community input on sampling protocol and data collected. More details are available at

https://whondrs.pnnl.gov.

Briefly, WHONDRS developed sampling protocols and videos in coordination with the scientific

community that were made openly available via YouTube, sent free sampling kits to collaborators,

and conducted a suite of biogeochemical analyses on surface water and sediment. All data will be

made open access following QA/QC at https://data.ess-dive.lbl.gov/. Preliminary data are available

on a Google Drive linked via https://whondrs.pnnl.gov as they become available.

The 2019 study collected samples and metadata associated with stream order, climate,

vegetation, and geomorphological features from 97 river corridors in 8 countries within a 6-week

period, from July 29th to September 19th (Table S2, [32]). Stream order information (Table S2) was

acquired for sites within the continental United States through the EPA National Hydrography

Dataset Plus (https://www.epa.gov/waterdata/nhdplus-national-hydrography-dataset-plus) and

stream orders for a couple of Canada sites were acquired through British Columbia Data Catalogue

(https://catalogue.data.gov.bc.ca/dataset/75299593-3222-40f9-879f-29e9824fc978).

This paper focuses on surface water (95 sites) and sediment (78 sites), collected across biomes

(i.e., desert, tropical, temperate forests), Toyoda et al. [32] provides additional metadata associated

to specific site characteristics (e.g., hydrogeomorphology, vegetation, temperature, discharge). We

use these data to describe the global distribution of key aspects of metabolomes including inferred

elemental groups, chemical classes, descriptor indices, and biochemical transformations.

3.2 Sample collection and laboratory pre-processing

At each location, collaborators selected sampling sites within 100 m of a station that measured

river discharge, height, or pressure. Within each site, 3 depositional zones were identified for

sediment collection following NEON's protocol (NEON.DOC.001193; [70]) and labeled as upstream,

midstream, or downstream. The depositional zones were situated within 10 m of each other. Surface

water was sampled in triplicate prior to sediment sampling to minimize the potential collection of

sediment debris from upstream sites mobilized during sampling. Surface water was collected only at

the downstream site. Sediment were collected from all three zones.

Surface water was collected using a 60 mL syringe and was filtered through a 0.22 um sterivex

filter (EMD Millipore) into 40 mL glass vial (I-Chem amber VOA glass vials; ThermoFisher, pre-

acidified with 10 uL of 85% phosphoric acid). Subsequently, 125 mL of surface sediment (1-3 cm

depth) were sampled from a ~1m^2 area at each depositional zone with a stainless-steel scoop, making

sure the sediments were saturated upon collection. All samples were shipped to Pacific Northwest

National Laboratory on blue ice within 24 h of collection.

Surface water samples were immediately frozen at -20°C upon receiving. Sediment from each

depositional zone were individually sieved to < 2 mm and subsampled into proteomic friendly tubes

(Genesee Scientific), and stored at -20°C for FTICR-MS analysis.

3.3 Fourier transform ion cyclotron resonance mass spectrometry (FTICR-MS)

Surface water samples were thawed in the dark at 4°C for 72 hours. Non-purgeable organic

carbon (NPOC) was determined using a 5 mL aliquot of the acidified water sample by a Shimadzu

combustion carbon analyzer TOC-L CSH/CSN E100V with ASI-L auto sampler. NPOC

concentrations (Table S4) were normalized to 1.5 mg C L⁻¹ across all samples to allow for data

comparison across sites within this study and other WHONDRS sampling campaigns. Diluted

samples were acidified to pH 2 with 85% phosphoric acid and extracted with PPL cartridges (Bond

Elut), following Dittmar et al. [71].

Sediment samples were thawed overnight in the dark at 4°C. Then, sediment organic matter was

extracted in proteomic friendly tubes (Genesee Scientific) with a 1:2 ratio of sediment to water (5 g of

sediment to 10 mL of milli-Q water). During the extraction, tubes were continuously shaken in the

dark at 375 rpm and 21°C for 2 h, after which the tubes were centrifuged at 6000 rcf and 21°C for 5

min. The supernatant was collected and filtered through 0.22 μm polyethersulfone membrane filter

(Millipore Sterivex, USA) into borosilicate glass vials. NPOC (Shimadzu combustion carbon analyzer

TOC–Vcsh with ASI– V auto sampler) was determined using a 5 mL aliquot from the filtered

supernatant. As with the water samples, this superpernatant was normalized to a standard NPOC

concentration (Table S4) of 1.5 mg C L⁻¹, acidified to pH 2 with 85% phosphoric acid and extracted

with PPL cartridges following the same methods described above.

A 12 Tesla (12T) Bruker SolariX Fourier transform ion cyclotron mass spectrometer (FTICR-MS;

Bruker, SolariX, Billerica, MA) located at the Environmental Molecular Sciences Laboratory in

Richland, WA, was used to collect ultrahigh-resolution mass spectra of surface water and sediment

OM pools. Resolution was 220K at 481.185 m/z. The FTICR-MS was outfitted with a standard

electrospray ionization (ESI) source, and data was acquired in negative mode with the voltage set to

+4.2kV. Data were collected with an ion accumulation of 0.05 sec for surface water and 0.1 or 0.2 sec

for sediment from 100 – 900 m/z at 4M. One hundred forty-four scans were co-added for each sample

and internally calibrated using an OM homologous series separated by 14 Da (–CH2 groups). The

mass measurement accuracy was typically within 1 ppm for singly charged ions across a broad m/z

range (100 m/z - 900 m/z). BrukerDaltonik Data Analysis (version 4.2) was used to convert raw spectra

to a list of m/z values by applying the FTMS peak picker module with a signal-to-noise ratio (S/N)

threshold set to 7 and absolute intensity threshold to the default value of 100. Chemical formulas

were assigned using Formularity (Formularity [72]) following the Compound Identification

Algorithm based on the following criteria: S/N >7, and mass measurement error < 0.5 ppm, taking

into consideration the presence of C, H, O, N, S, and P and excluding other elements. Formularity

[72] was also used to align spectra using a 0.5 ppm threshold.

3.4 FTICR-MS Data Analysis

All FTICR-MS analyses were performed using R v4.0.0 [73], and all plots were generated using

the ggplot2 package (v3.2.2) [74]. The R package 'ftmsRanalysis' [75] was used to (1) remove peaks

outside of a high confidence m/z range (200 m/z - 900 m/z) and/or with a 13C isotopic signature, (2)

calculate molecular formula properties (i.e., Kendrick defect, double-bond equivalent, modified

aromaticity index, nominal oxidation state of carbon, standard Gibbs Free Energy of carbon

oxidation), and (3) to determine to which chemical class a given metabolite belonged [33,34,36,38].

In order to evaluate bulk variation across sample types, a Mann Whitney U test (*wilcox.test*) with a

false discovery rate (FDR) p-value adjustment (*p.adjust*) was used to evaluate the divergence in

molecular properties of all metabolites observed in either surface water or sediment. Differences in

elemental group and chemical class relative abundances within samples between surface water and

sediment were evaluated using the same approach. A principal component analysis (PCA; *prcomp*)

was used to visualize differences between surface water and sediment metabolomes after a

presence/absence transformation. A Euclidean distance matrix was obtained (*vegdist*, vegan package

v2.5-6) and evaluated using a PERMANOVA (*adonis*, vegan package v2.5-6) in order to assess

multivariate differences between sample types [76]. Inter-sample type variability was evaluated

using the same Euclidean distance matrix in a beta-dispersion analysis (*betadisper*, vegan package

v2.5-6); divergence in distance to centroid values were then evaluated using a Mann Whitney U test

[76].

To determine CONUS-scale patterns, sites were divided into eastern and western US based on

their position relative to the location of the Mississippi River at St. Louis, Missouri. Replicates at each

site were merged such that if a metabolite was observed in one replicate, it was considered present

at the site. Average molecular properties were then calculated, and elemental group/chemical class

relative abundances were determined for each site/sample type combination based upon the

metabolites present. This resulted in a single value for any given variable in surface water or sediment

at a given site. Differences between these values across the East vs. West CONUS were then assessed

using a Mann Whitney U test with FDR correction. Maps were generated using ggplot to visualize

spatial variance. All maps can be found in File S1.

3.5 Biochemical Transformation Analysis

We inferred biochemical transformations in sediment and surface water metabolomes as per

Bailey et al. [50], Kaling et al. [57], Moritz et al. [58], Graham et al. [17,18], Garayburu-Caruso et al.

[16], Danczak et al. [26], and Stegen et al. [15] to estimate the gain or loss of specific molecules (e.g.,

glucose, valine, glutamine). Briefly, pairwise mass differences between all peaks in a sample were

compared with a reference list of 1255 commonly observed biochemical reactions of organic matter

(Table S5). For mass differences matching to compounds in the reference list (within 1 ppm), we

inferred the gain or loss of that compound via a biochemical transformation. For example, if a mass

difference between two peaks corresponded to 71.0371, that would correlate to the loss or gain of the

amino acid alanine, while a mass difference of 79.9662 would correspond to loss or gain of a

phosphate. Transformations were separated in 4 different groups based upon their labels: CHO-only,

N-containing, S-containing, and P-containing. Differences in the relative abundance of

transformations across samples were identified using a Mann Whitney U test with FDR correction.

3.6 Data Availability

Original and expanded metadata as well as surface water metabolome data used in this study

are publicly available on the Department of Energy data archive site ESS-DIVE [32]. Sediment

metabolomes data provided in the SI are available on the WHONDRS consortium preliminary data

Google Drive [https://drive.google.com/open?id=1I2Hz3PuXPGVmu4yjOWP36so1x26zUnjR] and

will soon be published on ESS-DIVE. All scripts used in this study are available on GitHub at

[https://github.com/danczakre/GlobalRiverMetabolomes].

4. Conclusions

We leveraged community science facilitated by the WHONDRS consortium to present the first

ultrahigh resolution analysis of global river corridor metabolomes of both surface water and

sediment. Our data showed strong divergence between surface water and sediment metabolomes,

consistent with previous work within local systems. Surface water metabolomes were more rich and

variable and contained more unsaturated and aromatic metabolites than sediment, possibly

suggesting higher influence from terrestrial inputs or lower microbial processing. Further, surface

water and sediment metabolomes had a consistent set of core biochemical transformations (CHO-

only) but differed in N-, S-, and P-containing transformations that may be more influenced by

nutrient limitations. Finally, we hypothesize the presence of systematic, spatially structured drivers

influencing sediment metabolomes more strongly than surface water, as sediment metabolomes

showed larger differences between the eastern and western parts of the continental United States.

While there are many potential explanations for these patterns, the publicly available datasets

being actively compiled by WHONDRS are well-suited for follow-on analyses to identify factors

underlying metabolome variability. Given that the WHONDRS sampling campaign spanned 1st to

9th stream orders across multiple biomes (e.g., desert-like in the Columbia Plateau, subtropical in

southern Florida, temperate forests in the Mid-Atlantic), outcomes of current and future data

analyses and modeling efforts will enable transferable knowledge that can be applied throughout the

world. To expand the breadth of questions that can be pursued with the data, WHONDRS is currently

collecting information pertaining to mineralogy, geochemistry (e.g., anion and total N

concentrations), microbiology (e.g., metagenomics, metatranscriptomics, flow cytometric cell

counts), and various remote sensing data types (e.g., vegetation cover). Future questions might, for

example, involve exploring spatial patterns of metabolomes across stream orders, correlating N-, S-,

and P-containing transformations with land use, mineralogy and vegetation, or investigating

relationships between microbial activity and metabolome composition. We encourage the scientific

community to explore WHONDRS datasets and combine them with additional data products to

pursue novel scientific questions at local to global scales, and to further engage with and pursue

science that embodies the ICON principles.

Supplementary Materials:

Table S1: This table contains by-sample mean, median, and standard deviation for molecular characteristics (i.e.,

aromaticity index, H:C ratios, etc.; Sheet 1), the number of metabolites belonging to a given elemental group (i.e.,

CHON, CHO; Sheet 2), and the number of metabolites belonging to some compound class (i.e., %lignin,

%protein, etc.; Sheet 3). Sheet 4 contains information about how each of the different measurements were

calculated.

Table S2: This spreadsheet contains meta-data for each site from which samples were collected, including (but

not limited to) latitude, longitude, stream order, and sampling date.

Table S3: This table contains the results of the FDR-corrected, two-sided Mann Whitney statistics performed to

evaluate spatial variability across the continental United States of America.

Table S4: Table of non-purgeable organic carbon (NPOC) concentrations for surface water and sediment water

extractions.

Table S5: The file is the database of transformations used in the transformation analysis. The first column

represents the transformation label, while the second column is the corresponding mass difference. There are

two types of transformations listed in this file: 1) the gain or loss of the listed molecular formula (e.g.,

C1H1O1N1) with numeric values indicating the number of atoms associated with the element that precedes the

numeric value, and 2) a substitution reaction denoted by an underscore (e.g., C1H1N1O_1). In the case of a

substitution reaction the underscore connects the element lost to the number of atoms lost. For example, C1H1N1O_1 indicates a molecule gained C1H1N1 and lost one O atom. Some substitution reactions include multiple elements that are lost such that there are multiple underscores. In all cases, an underscore connects the element lost to the number of atoms lost. In all cases, atoms are gained if they are not followed immediately by an underscore. For example, C_1H_4O2 indicates loss of one C, loss of four H, and gain of two O. If no numeric value follows an element, it indicates there is a gain of a single atom of that element (e.g., CH2 indicates one atom of C).

File S1: A compressed file containing all of the maps generated during the spatial analysis of the continental United States of America.

Author Contributions: V.A.G.-C., R.E.D., E.B.G, and J.C.S., conceptualized the study; V.A.G.-C., L.R., J.W., J.M.T., M.M., A.E.G, S.K and S.F. carried out sample processing; C.T.R., R.K.C., and J.T. conducted the instrumental analyses; V.A.G.-C., R.E.D, J.C.S and E.B.G drafted the manuscript; and all authors contributed to the writing.

Funding: This research was supported by the U.S. Department of Energy (DOE), Office of Biological and Environmental Research (BER), as part of Subsurface Biogeochemical Research Program's Scientific Focus Area (SFA) at the Pacific Northwest National Laboratory (PNNL). Data were generated under EMSL User Proposal 51180. A portion of the research was performed at Environmental Molecular Science Laboratory User Facility. PNNL is operated for DOE by Battelle under Contract DE-AC06-76RLO 1830.

Acknowledgments: The authors would like to thank the WHONDRS consortium.

References

1. Battin, T.J.; Luyssaert, S.; Kaplan, L.A.; Aufdenkampe, A.K.; Richter, A.; Tranvik, L.J. The boundless carbon cycle. *Nat. Geosci.* **2009**, *2*, 598–600, doi:10.1038/ngeo618.

2. Cole, J.J.; Prairie, Y.T.; Caraco, N.F.; McDowell, W.H.; Tranvik, L.J.; Striegl, R.G.; Duarte, C.M.; Kortelainen, P.; Downing, J.A.; Middelburg, J.J.; et al. Plumbing the Global Carbon Cycle: Integrating Inland Waters into the Terrestrial Carbon Budget. *Ecosystems* **2007**, *10*, 172–185, doi:10.1007/s10021-006-9013-8.

3. Marín-Spiotta, E.; Gruley, K.E.; Crawford, J.; Atkinson, E.E.; Miesel, J.R.; Greene, S.; Cardona-Correa, C.; Spencer, R.G.M. Paradigm shifts in soil organic matter research affect interpretations of aquatic carbon cycling: transcending disciplinary and ecosystem boundaries. *Biogeochemistry* **2014**, *117*, 279–297, doi:10.1007/s10533-013-9949-7.

4. Regnier, P.; Friedlingstein, P.; Ciais, P.; Mackenzie, F.T.; Gruber, N.; Janssens, I.A.; Laruelle, G.G.; Lauerwald, R.; Luyssaert, S.; Andersson, A.J.; et al. Anthropogenic perturbation of the carbon fluxes from land to ocean. *Nat. Geosci.* **2013**, *6*, 597–607, doi:10.1038/ngeo1830.

5. Hotchkiss, E.R.; Hall Jr, R.O.; Sponseller, R.A.; Butman, D.; Klaminder, J.; Laudon, H.; Rosvall, M.; Karlsson, J. Sources of and processes controlling CO 2 emissions change with the size of streams and rivers. *Nat. Geosci.* **2015**, *8*, 696–699, doi:10.1038/ngeo2507.

6. Catalán, N.; Casas-Ruiz, J.P.; Arce, M.I.; Abril, M.; Bravo, A.G.; Campo, R. del; Estévez, E.; Freixa, A.; Giménez-Grau, P.; González-Ferreras, A.M.; et al. Behind the Scenes: Mechanisms Regulating Climatic Patterns of Dissolved Organic Carbon Uptake in Headwater Streams. *Glob. Biogeochem. Cycles* **2018**, *32*, 1528–1541, doi:10.1029/2018GB005919.

7. Aufdenkampe, A.K.; Mayorga, E.; Raymond, P.A.; Melack, J.M.; Doney, S.C.; Alin, S.R.; Aalto, R.E.; Yoo, K. Riverine coupling of biogeochemical cycles between land, oceans, and atmosphere. *Front. Ecol. Environ.* **2011**, *9*, 53–60, doi:10.1890/100014.

8. Raymond, P.A.; Hartmann, J.; Lauerwald, R.; Sobek, S.; McDonald, C.; Hoover, M.; Butman, D.; Striegl, R.; Mayorga, E.; Humborg, C.; et al. Global carbon dioxide emissions from inland waters. *Nature* **2013**, *503*, 355–359, doi:10.1038/nature12760.

9. Moody, C.S.; Worrall, F.; Evans, C.D.; Jones, T.G. The rate of loss of dissolved organic carbon (DOC) through a catchment. *J. Hydrol.* **2013**, *492*, 139–150, doi:10.1016/j.jhydrol.2013.03.016.

10. Cory, R.M.; Ward, C.P.; Crump, B.C.; Kling, G.W. Sunlight controls water column processing of carbon in arctic fresh waters. *Science* **2014**, *345*, 925, doi:10.1126/science.1253119.

11. Newbold, J.D.; Bott, T.L.; Kaplan, L.A.; Dow, C.L.; Jackson, J.K.; Aufdenkampe, A.K.; Martin, L.A.; Horn, D.J.V.; Long, A.A. Uptake of nutrients and organic C in streams in New York City drinking-water-supply watersheds. *Freshw. Sci.* **2006**, *25*, 998–1017, doi:10.1899/0887-3593(2006)025[0998:UONAOC]2.0.CO;2.

12. Bernhardt, E.S.; McDowell, W.H. Twenty years apart: Comparisons of DOM uptake during leaf leachate releases to Hubbard Brook Valley streams in 1979 versus 2000. *J. Geophys. Res. Biogeosciences* **2008**, *113*, doi:10.1029/2007JG000618.

580   13.   Koehler, B.; Wachenfeldt, E. von; Kothawala, D.; Tranvik, L.J. Reactivity continuum of dissolved organic

581   carbon decomposition in lake water. *J. Geophys. Res. Biogeosciences* **2012**, *117*, doi:10.1029/2011JG001793.

582   14.   Berggren, M.; Giorgio, P.A. del Distinct patterns of microbial metabolism associated to riverine dissolved

583   organic carbon of different source and quality. *J. Geophys. Res. Biogeosciences* **2015**, *120*, 989–999,

584   doi:10.1002/2015JG002963.

585   15.   Stegen, J.C.; Johnson, T.; Fredrickson, J.K.; Wilkins, M.J.; Konopka, A.E.; Nelson, W.C.; Arntzen, E.V.;

586   Chrisler, W.B.; Chu, R.K.; Fansler, S.J.; et al. Influences of organic carbon speciation on hyporheic corridor

587   biogeochemistry and microbial ecology. *Nat. Commun.* **2018**, *9*, 1–11, doi:10.1038/s41467-018-02922-9.

588   16.   Garayburu-Caruso, V.A.; Stegen, J.C.; Song, H.-S.; Renteria, L.; Wells, J.; Garcia, W.; Resch, C.T.; Goldman,

589   A.E.; Chu, R.K.; Toyoda, J.; et al. Carbon Limitation Leads to Thermodynamic Regulation of Aerobic

590   Metabolism. *Environ. Sci. Technol. Lett.* **2020**, *7*, 517–524, doi:10.1021/acs.estlett.0c00258.

591   17.   Graham, E.B.; Crump, A.R.; Kennedy, D.W.; Arntzen, E.; Fansler, S.; Purvine, S.O.; Nicora, C.D.; Nelson,

592   W.; Tfaily, M.M.; Stegen, J.C. Multi 'omics comparison reveals metabolome biochemistry, not microbiome

593   composition or gene expression, corresponds to elevated biogeochemical function in the hyporheic zone.

594   *Sci. Total Environ.* **2018**, *642*, 742–753, doi:10.1016/j.scitotenv.2018.05.256.

595   18.   Graham, E.B.; Tfaily, M.M.; Crump, A.R.; Goldman, A.E.; Bramer, L.M.; Arntzen, E.; Romero, E.; Resch,

596   C.T.; Kennedy, D.W.; Stegen, J.C. Carbon Inputs From Riparian Vegetation Limit Oxidation of Physically

597   Bound Organic Carbon Via Biochemical and Thermodynamic Processes. *J. Geophys. Res. Biogeosciences* **2017**,

598   *122*, 3188–3205, doi:10.1002/2017JG003967.

599   19.   Lam, B.; Baer, A.; Alaee, M.; Lefebvre, B.; Moser, A.; Williams, A.; Simpson, A.J. Major Structural

600   Components in Freshwater Dissolved Organic Matter. *Environ. Sci. Technol.* **2007**, *41*, 8240–8247,

601   doi:10.1021/es0713072.

602   20.   Jaffé, R.; McKnight, D.; Maie, N.; Cory, R.; McDowell, W.H.; Campbell, J.L. Spatial and temporal variations

603   in DOM composition in ecosystems: The importance of long-term monitoring of optical properties. *J.*

604   *Geophys. Res. Biogeosciences* **2008**, *113*, doi:10.1029/2008JG000683.

605   21.   Gonsior, M.; Peake, B.M.; Cooper, W.T.; Podgorski, D.; D'Andrilli, J.; Cooper, W.J. Photochemically

606   induced changes in dissolved organic matter identified by ultrahigh resolution fourier transform ion

607   cyclotron resonance mass spectrometry. *Environ. Sci. Technol.* **2009**, *43*, 698–703, doi:10.1021/es8022804.

608 22. Lu, Y.; Li, X.; Mesfioui, R.; Bauer, J.E.; Chambers, R.M.; Canuel, E.A.; Hatcher, P.G. Use of ESI-FTICR-MS

609   to Characterize Dissolved Organic Matter in Headwater Streams Draining Forest-Dominated and Pasture-

610   Dominated Watersheds. *PLOS ONE* **2015**, *10*, e0145639, doi:10.1371/journal.pone.0145639.

611 23. Stegen, J.C.; Goldman, A.E. WHONDRS: a Community Resource for Studying Dynamic River Corridors.

612   *mSystems* **2018**, *3*, doi:10.1128/mSystems.00151-18.

613 24. Jaffé, R.; Yamashita, Y.; Maie, N.; Cooper, W.T.; Dittmar, T.; Dodds, W.K.; Jones, J.B.; Myoshi, T.; Ortiz-

614   Zayas, J.R.; Podgorski, D.C.; et al. Dissolved Organic Matter in Headwater Streams: Compositional

615   Variability across Climatic Regions of North America. *Geochim. Cosmochim. Acta* **2012**, *94*, 95–108,

616   doi:10.1016/j.gca.2012.06.031.

617 25. Wagner, S.; Riedel, T.; Niggemann, J.; Vähätalo, A.V.; Dittmar, T.; Jaffé, R. Linking the Molecular Signature

618   of Heteroatomic Dissolved Organic Matter to Watershed Characteristics in World Rivers. *Environ. Sci.*

619   *Technol.* **2015**, *49*, 13798–13806, doi:10.1021/acs.est.5b00525.

620 26. Danczak, R.E.; Goldman, A.E.; Chu, R.K.; Toyoda, J.G.; Garayburu-Caruso, V.A.; Tolić, N.; Graham, E.B.;

621   Morad, J.W.; Renteria, L.; Wells, J.R.; et al. Ecological theory applied to environmental metabolomes reveals

622   compositional divergence despite conserved molecular properties. *bioRxiv* **2020**, 2020.02.12.946459,

623   doi:10.1101/2020.02.12.946459.

624 27. Steefel, C.I.; Appelo, C.A.J.; Arora, B.; Jacques, D.; Kalbacher, T.; Kolditz, O.; Lagneau, V.; Lichtner, P.C.;

625   Mayer, K.U.; Meeussen, J.C.L.; et al. Reactive transport codes for subsurface environmental simulation.

626   *Comput. Geosci.* **2015**, *19*, 445–478, doi:10.1007/s10596-014-9443-x.

627 28. Song, H.-S.; Stegen, J.C.; Graham, E.B.; Lee, J.-Y.; Garayburu-Caruso, V.A.; Nelson, W.C.; Chen, X.;

628   Moulton, J.D.; Scheibe, T.D. Representing Organic Matter Thermodynamics in Biogeochemical Reactions

629   via Substrate-Explicit Modeling. *Frontiers in Microbiology* **2020**, 11 https://doi.org/10.3389/fmicb.2020.531756

630 29. Stegen, J.; Brodie, E.; Wrighton, K.; Bayer, P.; Lesmes, D.; Emani, S.; Moerman, J. *Open Watershed Science by*

631   *Design: Leveraging Distributed Research Networks to Understand Watershed Systems: Workshop Report*; USDOE

632   Office of Science (SC)(United States), 2019;

633 30. Uhlmann, E.L.; Ebersole, C.R.; Chartier, C.R.; Errington, T.M.; Kidwell, M.C.; Lai, C.K.; McCarthy, R.J.;

634   Riegelman, A.; Silberzahn, R.; Nosek, B.A. Scientific Utopia III: Crowdsourcing Science: *Perspect. Psychol.*

635   *Sci.* **2019**, doi:10.1177/1745691619850561.

636   31.   Wilkinson, M.D.; Dumontier, M.; Aalbersberg, Ij.J.; Appleton, G.; Axton, M.; Baak, A.; Blomberg, N.; Boiten,

637   J.-W.; da Silva Santos, L.B.; Bourne, P.E.; et al. The FAIR Guiding Principles for scientific data management

638   and stewardship. *Sci. Data* **2016**, *3*, 160018, doi:10.1038/sdata.2016.18.

639   32.   Toyoda, J.G.; Goldman, A.E.; Chu, R.K.; Danczak, R.E.; Daly, R.A.; Garayburu-Caruso, V.A.; Graham, E.B.;

640   Lin, X.; Moran, J.J.; Ren, H. *WHONDRS Summer 2019 Sampling Campaign: Global River Corridor Surface Water*

641   *FTICR-MS and Stable Isotopes*; Environmental System Science Data Infrastructure for a Virtual Ecosystem …,

642   2020;

643   33.   Koch, B.P.; Dittmar, T. From mass to structure: an aromaticity index for high-resolution mass data of

644   natural organic matter. *Rapid Commun. Mass Spectrom.* **2006**, *20*, 926–932, doi:10.1002/rcm.2386.

645   34.   Koch, B.P.; Dittmar, T. From mass to structure: an aromaticity index for high-resolution mass data of

646   natural organic matter. *Rapid Commun. Mass Spectrom.* **2016**, *30*, 250–250, doi:10.1002/rcm.7433.

647   35.   Willoughby, A.S.; Wozniak, A.S.; Hatcher, P.G. A molecular-level approach for characterizing water-

648   insoluble components of ambient organic aerosol particulates using ultrahigh-resolution mass

649   spectrometry. *Atmospheric Chem. Phys.* **2014**, *14*, 10299–10314, doi:https://doi.org/10.5194/acp-14-10299-

650   2014.

651   36.   LaRowe, D.E.; Van Cappellen, P. Degradation of natural organic matter: A thermodynamic analysis.

652   *Geochim. Cosmochim. Acta* **2011**, *75*, 2030–2042, doi:10.1016/j.gca.2011.01.020.

653   37.   Boye, K.; Noël, V.; Tfaily, M.M.; Bone, S.E.; Williams, K.H.; Bargar, J.R.; Fendorf, S. Thermodynamically

654   controlled preservation of organic carbon in floodplains. *Nat. Geosci.* **2017**, *10*, 415–419,

655   doi:10.1038/ngeo2940.

656   38.   Kim, S.; Kramer, R.W.; Hatcher, P.G. Graphical Method for Analysis of Ultrahigh-Resolution Broadband

657   Mass Spectra of Natural Organic Matter, the Van Krevelen Diagram. *Anal. Chem.* **2003**, *75*, 5336–5344,

658   doi:10.1021/ac034415p.

659   39.   Hedges, J.I.; Mann, D.C. The lignin geochemistry of marine sediments from the southern Washington coast.

660   *Geochim. Cosmochim. Acta* **1979**, *43*, 1809–1818, doi:10.1016/0016-7037(79)90029-2.

661   40.   Stegen, J.C.; Johnson, T.; Fredrickson, J.K.; Wilkins, M.J.; Konopka, A.E.; Nelson, W.C.; Arntzen, E.V.;

662   Chrisler, W.B.; Chu, R.K.; Fansler, S.J.; et al. Influences of organic carbon speciation on hyporheic corridor

663   biogeochemistry and microbial ecology. *Nat. Commun.* **2018**, *9*, 585, doi:10.1038/s41467-018-02922-9.

41. Pracht, L.E.; Tfaily, M.M.; Ardissono, R.J.; Neumann, R.B. Molecular characterization of organic matter

mobilized from Bangladeshi aquifer sediment: tracking carbon compositional change during microbial

utilization. *Biogeosciences* **2018**, *15*, 1733–1747, doi:https://doi.org/10.5194/bg-15-1733-2018.

42. Valle, J.; Harir, M.; Gonsior, M.; Enrich-Prast, A.; Schmitt-Kopplin, P.; Bastviken, D.; Hertkorn, N.

Molecular differences between water column and sediment pore water SPE-DOM in ten Swedish boreal

lakes. *Water Res.* **2020**, *170*, 115320, doi:10.1016/j.watres.2019.115320.

43. Boulton, A.J.; Findlay, S.; Marmonier, P.; Stanley, E.H.; Valett, H.M. The functional significance of the

hyporheic zone in streams and rivers. *Annu. Rev. Ecol. Syst.* **1998**, *29*, 59–81,

doi:10.1146/annurev.ecolsys.29.1.59.

44. Boano, F.; Harvey, J.W.; Marion, A.; Packman, A.I.; Revelli, R.; Ridolfi, L.; Wörman, A. Hyporheic flow and

transport processes: Mechanisms, models, and biogeochemical implications. *Rev. Geophys.* **2014**, *52*, 603–

679, doi:10.1002/2012RG000417.

45. Harvey, J.; Gooseff, M. River corridor science: Hydrologic exchange and ecological consequences from

bedforms to basins. *Water Resour. Res.* **2015**, *51*, 6893–6922, doi:10.1002/2015WR017617.

46. Gomez-Velez, J.D.; Harvey, J.W.; Cardenas, M.B.; Kiel, B. Denitrification in the Mississippi River network

controlled by flow through river bedforms. *Nat. Geosci.* **2015**, *8*, 941–945, doi:10.1038/ngeo2567.

47. Knapp, J.L.A.; González-Pinzón, R.; Drummond, J.D.; Larsen, L.G.; Cirpka, O.A.; Harvey, J.W. Tracer-based

characterization of hyporheic exchange and benthic biolayers in streams. *Water Resour. Res.* **2017**, *53*, 1575–

1594, doi:10.1002/2016WR019393.

48. Fischer, H.; Kloep, F.; Wilzcek, S.; Pusch, M.T. A river's liver–microbial processes within the hyporheic

zone of a large lowland river. *Biogeochemistry* **2005**, *76*, 349–371.

49. Battin, T.J.; Besemer, K.; Bengtsson, M.M.; Romani, A.M.; Packmann, A.I. The ecology and biogeochemistry

of stream biofilms. *Nat. Rev. Microbiol.* **2016**, *14*, 251.

50. Bailey, V.L.; Smith, A.P.; Tfaily, M.; Fansler, S.J.; Bond-Lamberty, B. Differences in soluble organic carbon

chemistry in pore waters sampled from different pore size domains. *Soil Biol. Biochem.* **2017**, *107*, 133–143,

doi:10.1016/j.soilbio.2016.11.025.

51. Cory, R.M.; Kling, G.W. Interactions between sunlight and microorganisms influence dissolved organic

matter degradation along the aquatic continuum. *Limnol. Oceanogr. Lett.* **2018**, *3*, 102–116,

doi:10.1002/lol2.10060.

52.   Bao, H.; Niggemann, J.; Huang, D.; Dittmar, T.; Kao, S.-J. Different Responses of Dissolved Black Carbon
694   and Dissolved Lignin to Seasonal Hydrological Changes and an Extreme Rain Event. *J. Geophys. Res.*
695   *Biogeosciences* **2019**, *124*, 479–493, doi:10.1029/2018JG004822.

696   53.   Fellman, J.B.; Hood, E.; Edwards, R.T.; D'Amore, D.V. Changes in the concentration, biodegradability, and
697   fluorescent properties of dissolved organic matter during stormflows in coastal temperate watersheds. *J.*
698   *Geophys. Res. Biogeosciences* **2009**, *114*, doi:10.1029/2008JG000790.

699   54.   Hood, E.; Gooseff, M.N.; Johnson, S.L. Changes in the character of stream water dissolved organic carbon
700   during flushing in three small watersheds, Oregon. *J. Geophys. Res. Biogeosciences* **2006**, *111*,
701   doi:10.1029/2005JG000082.

702   55.   Ward, N.D.; Richey, J.E.; Keil, R.G. Temporal variation in river nutrient and dissolved lignin phenol
703   concentrations and the impact of storm events on nutrient loading to Hood Canal, Washington, USA.
704   *Biogeochemistry* **2012**, *111*, 629–645, doi:10.1007/s10533-012-9700-9.

705   56.   Tfaily, M.M.; Chu, R.K.; Toyoda, J.; Tolić, N.; Robinson, E.W.; Paša-Tolić, L.; Hess, N.J. Sequential extraction
706   protocol for organic matter from soils and sediments using high resolution mass spectrometry. *Anal. Chim.*
707   *Acta* **2017**, *972*, 54–61, doi:https://doi.org/10.1016/j.aca.2017.03.031.

708   57.   Kaling, M.; Schmidt, A.; Moritz, F.; Rosenkranz, M.; Witting, M.; Kasper, K.; Janz, D.; Schmitt-Kopplin, P.;
709   Schnitzler, J.-P.; Polle, A. Mycorrhiza-Triggered Transcriptomic and Metabolomic Networks Impinge on
710   Herbivore Fitness. *Plant Physiol.* **2018**, *176*, 2639–2656, doi:10.1104/pp.17.01810.

711   58.   Moritz, F.; Kaling, M.; Schnitzler, J.-P.; Schmitt-Kopplin, P. Characterization of poplar metabotypes via
712   mass difference enrichment analysis. *Plant Cell Environ.* **2017**, *40*, 1057–1073, doi:10.1111/pce.12878.

713   59.   Craine, J.M.; Morrow, C.; Fierer, N. Microbial Nitrogen Limitation Increases Decomposition. *Ecology* **2007**,
714   *88*, 2105–2113, doi:10.1890/06-1847.1.

715   60.   Moorhead, D.L.; Sinsabaugh, R.L. A Theoretical Model of Litter Decay and Microbial Interaction. *Ecol.*
716   *Monogr.* **2006**, *76*, 151–174, doi:10.1890/0012-9615(2006)076[0151:ATMOLD]2.0.CO;2.

717   61.   Zhao, F.J.; Wu, J.; McGrath, S.P. Chapter 12 - Soil Organic Sulphur and its Turnover. In *Humic Substances in*
718   *Terrestrial Ecosystems*; Piccolo, A., Ed.; Elsevier Science B.V.: Amsterdam, 1996; pp. 467–506 ISBN 978-0-444-
719   81516-3.

720   62.   Reddy, K.R.; Kadlec, R.H.; Flaig, E.; Gale, P.M. Phosphorus Retention in Streams and Wetlands: A Review.
721   *Crit. Rev. Environ. Sci. Technol.* **1999**, *29*, 83–146, doi:10.1080/10643389991259182.

63.   Freixa, A.; Ejarque, E.; Crognale, S.; Amalfitano, S.; Fazi, S.; Butturini, A.; Romaní, A.M. Sediment microbial

communities rely on different dissolved organic matter sources along a Mediterranean river continuum.

Limnol. Oceanogr. **2016**, *61*, 1389–1405, doi:10.1002/lno.10308.

64.   Vazquez, E.; Amalfitano, S.; Fazi, S.; Butturini, A. Dissolved organic matter composition in a fragmented

Mediterranean fluvial system under severe drought conditions. *Biogeochemistry* **2011**, *102*, 59–72,

doi:10.1007/s10533-010-9421-x.

65.   Butturini, A.; Guarch, A.; Romaní, A.M.; Freixa, A.; Amalfitano, S.; Fazi, S.; Ejarque, E. Hydrological

conditions control in situ DOM retention and release along a Mediterranean river. *Water Res.* **2016**, *99*, 33–

45, doi:10.1016/j.watres.2016.04.036.

66.   Ejarque, E.; Freixa, A.; Vazquez, E.; Guarch, A.; Amalfitano, S.; Fazi, S.; Romaní, A.M.; Butturini, A. Quality

and reactivity of dissolved organic matter in a Mediterranean river across hydrological and spatial

gradients. *Sci. Total Environ.* **2017**, *599–600*, 1802–1812, doi:10.1016/j.scitotenv.2017.05.113.

67.   Robinson, N.P.; Allred, B.W.; Jones, M.O.; Moreno, A.; Kimball, J.S.; Naugle, D.E.; Erickson, T.A.;

Richardson, A.D. A dynamic Landsat derived normalized difference vegetation index (NDVI) product for

the conterminous United States. *Remote Sens.* **2017**, *9*, 863.

68.   *A new map of standardized terrestrial ecosystems of the conterminous United States*; Sayre, R., Ed.; Professional

paper / U.S. Department of the Interior, U.S. Geological Survey; U.S. Dept. of the Interior, U.S. Geological

Survey: Reston, Va, 2009; ISBN 978-1-4113-2432-9.

69.   Fierer, N.; Ladau, J.; Clemente, J.C.; Leff, J.W.; Owens, S.M.; Pollard, K.S.; Knight, R.; Gilbert, J.A.; McCulley,

R.L. Reconstructing the Microbial Diversity and Function of Pre-Agricultural Tallgrass Prairie Soils in the

United States. *Science* **2013**, *342*, 621–624, doi:10.1126/science.1243768.

70.   Jensen, B. *AOS Protocol and Procedure: Sediment Chemistry Sampling in Wadeable Streams (NEON.DOC.001193)*

http://data.neonscience.org/documents; 2019;

71.   Dittmar, T.; Koch, B.; Hertkorn, N.; Kattner, G. A simple and efficient method for the solid-phase extraction

of dissolved organic matter (SPE-DOM) from seawater: SPE-DOM from seawater. *Limnol. Oceanogr.*

Methods **2008**, *6*, 230–235, doi:10.4319/lom.2008.6.230.

72.   Tolić, N.; Liu, Y.; Liyu, A.; Shen, Y.; Tfaily, M.M.; Kujawinski, E.B.; Longnecker, K.; Kuo, L.-J.; Robinson,

E.W.; Paša-Tolić, L.; et al. Formularity: Software for Automated Formula Assignment of Natural and Other

Organic Matter from Ultrahigh-Resolution Mass Spectra. *Anal. Chem.* **2017**, *89*, 12659–12665,

doi:10.1021/acs.analchem.7b03318.

73.   R Core Team *R: A language and environment for statistical computing.*; R Foundation for Statistical Computing:

753   Vienna, Austria.;

754   74.   Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*; Springer-Verlag: New York, 2016.;

755   75.   Bramer, L.M.; White, A.M.; Stratton, K.G.; Thompson, A.M.; Claborne, D.; Hofmockel, K.; McCue, L.A.

756   ftmsRanalysis: An R package for exploratory data analysis and interactive visualization of FT-MS data.

757   *PLOS Comput. Biol.* **2020**, *16*, e1007654, doi:10.1371/journal.pcbi.1007654.

758   76.   Oksanen, J.; Blanchet, F.G.; Friendly, M.; Kindt, R.; Legendre, P.; McGlinn, D.; Minchin, P.R.; O'hara, R.;

759   Simpson, G.L.; Solymos, P. vegan: Community Ecology Package. R package version 2.5-6. *Vienna R Found.*

760   *Stat. Comput. Sch.* **2019**.