Follicular fluid metabolome and cytokinome profiles in poor ovarian

responders and the impact of dehydroepiandrosterone supplementation

Veronique Viardot-Foucault[a,*], Jieliang Zhou[b,*], Dexi Bi[c], Yoshihiko Takinami[d], Heng

Hao Tan[a,e], Jerry.K.Y. Chan[a,e,†], Yie Hou Lee[b,e,†]

[a]Department of Reproductive Medicine, KK Women's and Children's Hospital, 100

Bukit Timah Road, Singapore 229899

[b]Translational 'Omics and Biomarkers Group, KK Research Centre, KK Women's

and Children's Hospital, 100 Bukit Timah Road, Singapore 229899

[c]Department of Pathology, Shanghai Tenth People's Hospital, Tongji University

School of Medicine, Shanghai 200072, China

[d]Bruker Japan, 3-9 Yokohama City, Kanagawa, Japan 220-0022

[e]Obstetrics and Gynaecology Academic Clinical Program, Duke-NUS Medical

School, 8 College Road, Singapore 169857.

Emails:

VVF: veronique.viardot@singhealth.com.sg;

JLZ: Zhou.Jieliang@kkh.com.sg

DB: bidexi2@163.com

YT : Yoshihiko.Takinami@bruker.com

HHT: tan.heng.hao@singhealth.com.sg

JKYC: jerrychan@duke-nus.edu.sg

YHL: gmsleeyh@nus.edu.sg

[†]***Address correspondence to:***

Jerry Chan MD, Ph.D

Phone: +65 6394 1051

E-mail: jerrychan@duke-nus.edu.sg

Lee Yie Hou Ph.D.

Phone: 65-6394 8122

Fax: 65-6293 1618

E-mail: gmsleeyh@nus.edu.sg

Conflict of interest: The authors have declared that no conflict of interest exists.

*These authors contributed equally.

Keywords: metabolomics, follicular fluid, DHEA, poor ovarian responder, cytokines

Abstract

Poor ovarian responders (POR) are women undergoing in-vitro fertilization who respond poorly to ovarian stimulation, resulting in the retrieval of lower number of oocytes, and subsequently lower pregnancy rates. The follicular fluid (FF) provides a crucial microenvironment for the proper development of follicles and oocytes. Conversely, dysregulated FF metabolome and cytokinome could have detrimental effects on oocytes in POR. Androgens such as dehydroepiandrosterone (DHEA) have been proposed to alter the POR follicular microenvironment but its effects on the FF metabolome and cytokine profiles is unknown. In this study, untargeted LC-MS/MS metabolomics was performed on FF of POR patients with DHEA supplementation (DHEA+) and without (DHEA-) in a randomized clinical trial (*N*=52). Untargeted metabolomics identified 118 FF metabolites of diverse chemistries, which included lipids, steroids, amino acids, hormones, among others. FF metabolomes were different between DHEA+ and DHEA- groups. Specifically, glycerophosphocholine, linoleic acid, progesterone, and valine were significantly lower in DHEA+ relative to DHEA-. Among cytokines, MCP1, IFNγ, LIF and VEGF-D were significantly lower in DHEA+ relative to DHEA. Collectively, our data suggest a role of DHEA on these metabolic and cytokines pathways, and these FF metabolites could be used to guide future studies in DHEA supplementation regimen.

(197 words)

Introduction

Poor ovarian responders (POR) are a sub-group of infertile women that account for 9-26% of *in vitro* fertilization (IVF) indications.[1,2] In patients designated as "poor responders," so-called due to poor response to ovarian stimulation given during IVF workup, the limited number of obtained oocytes remains the major problem in optimizing the live birth rates.[3] In fact, as a result of a lower number of oocytes retrieved, there are fewer embryos to select and transfer, and subsequently these patients have lower pregnancy rates per transfer and lower cumulative pregnancy rates per started cycle compared with normal responders. In PORs the mechanism of ovarian insufficiency can be multifactorial with causes such as ovarian surgery especially in case of endometrioma,[4,5] uterine artery embolization for the treatment of uterine leiomyoma,[6,7] genetic defects, chemotherapy, radiotherapy, autoimmune disorders, single ovary, chronic smoking,[8,9] or linked to diseases such as diabetes mellitus Type I.[10] However, in most cases, follicular depletion plausibly reflecting premature ovarian aging,[11] which clinically translates into a reduction of implantation rates, an increase of early pregnancy loss, and disappointingly low IVF success.[12,13]

In each menstrual cycle, human ovaries produce a single dominant follicle. Growth of the dominant follicle encompasses enlargement of the oocyte, replication of follicular cells, and formation and expansion of a fluid-filled follicular antrum or cavity, providing a specialized microenvironment for the development of oocytes. Follicular fluid (FF) that fills the antrum cavity is derived from the surrounding theca capillaries, abundant and easily accessible during IVF procedures due to ample volume being produced during follicle maturation.[14] FF are rich in metabolites, notably hormones, and proteins that are critical for oocyte growth and development, which determines subsequent potential to achieve fertilization and embryo development. As such,

constituents of the FF surrounding the oocyte provides a unique biochemical window

about the growth and differentiation of the oocyte.[15] To divulge into the biochemical

space of human FF, gas chromatography-mass spectrometric (GC-MS) and proton

nuclear magnetic resonance (^{1}H NMR) metabolomic analyses have been previously

conducted,[16–19] as were proteomic analyses.[20–25] These studies mainly report the

IVF FF profile, whereas the FF metabolome of POR remains poorly characterized.

Dehydroepiandrosterone (DHEA) is a steroid produced in the adrenal cortex and the

ovarian theca cells in women that is converted into more active forms of androgens

such as testosterone.[5] It has been suggested that DHEA supplementation may

increase the number of available follicles in PORs through an increased serum level

of insulin-like growth factor, increased follicular response to follicle stimulating

hormone (FSH), and improved quality of oocytes.[26] However, the efficacy of DHEA

pre-treatment has been controversial, with partial to low clinical evidence being

observed.[9,13-14,16] Here, a randomized clinical trial was conducted to evaluate the

effects of DHEA on IVF outcomes in our patient population, and a metabolomics

approach was used to understand the effects of DHEA on the FF metabolome.

Given that exogenous stimulus such as DHEA supplementation might alter the FF

metabolic profile, we hypothesized that DHEA effects could be improved through

uncovering 'responder' FF metabolites, which could be tapped to improve DHEA

dose regime. With the advent of LC-MS/MS, LC-MS global (untargeted)

metabolomics analysis has provided the ability to reveal biologically relevant

changes within a system, even at sensitive ranges before the precedence of gross

morphological   or   phenotypical   changes.[28]   Furthermore,   DHEA   has

immunoregulatory functions, and the large-scale study of cytokines plausibly reveals

DHEA immunomodulatory targets.[29] A better understanding of the molecular

mechanisms implicated in the conversion of DHEA to testosterone will offer

investigators the opportunity to design individualized treatments for infertile women

with POR, which are tailored to patients' unique metabolic profiles. Therefore, in this

study we mapped the FF metabolome and cytokine profile of POR with and without

DHEA supplementation.

Material and methods

Ethical approval

The study was approved by the Institutional Review Board at the Singapore Health Systems, and written informed consent was obtained from each participant. The study was registered with the National Institutes of Health (NIH) clinical trial site (clinicaltrials.gov: NCT01535872).

Study population and design

The prospective open-label randomized controlled trial (RCT) was conducted to evaluate the effect of DHEA administration in women, below the age of 42 starting their IVF treatment who met one of the two following features of POR (an abnormal ovarian reserve test and/or a previous poor response to ovarian stimulation in an IVF cycle) were assessed for eligibility.[3] We defined POR in this study as women with diminished ovarian reserves (AMH <1.0 ng/mL or Day 2 or 3 FSH >10 IU/L), or women with fewer than four oocytes retrieved with either standard long or antagonist protocols.[3] Inclusion criteria included women with diminished ovarian reserves (anti-müllerian hormone <1.0 ng/mL or D2/3 follicle stimulating hormone >10 IU/L), or women with fewer than four oocytes retrieved with either standard long or antagonist protocols. Exclusion criteria included women with previous or current DHEA supplementation, use of corticosteroids within the past three months, major systemic illnesses, and allergy to DHEA. Eligible patients were randomized to either the DHEA treatment group (DHEA+, *N*=28) and received DHEA (Pharma Natural, USA) at the dose of 75 mg/day for three to eight months prior starting their controlled ovarian stimulation (COS), or the control group, receiving no treatment (DHEA-, *N*=24)

(**Table 1**). Average age (DHEA-, 36 years; DHEA+, 37 years) and body mass index

(DHEA-, 24 kg/m^2; DHEA+, 23 kg/m^2) were similar in both groups (p>0.05).

IVF/ICSI protocol

All individuals received the same stimulation protocol, same starting dose of

gonadotropin, and fertilization technique. Briefly, the IVF/ICSI treatment cycle

involved an antagonist based COS protocol consisting of daily sub-cutaneous

injections of recombinant-FSH (Puregon, Follitropin beta, 300iu; MSD, USA) and

highly-purified human menopausal gonadotropin (Menopur; Menotropin, 150 IU;

Ferring Pharmaceuticals, Germany) with initiation of gonadotropin releasing

hormone antagonist (Ganirelix, Orgalutan, 0.25 mg s/c; MSD, USA) on day 5 of

COS. The dose of Menopur and Puregon could be further increased depending on

individual ovarian response. All patients had this standardized antagonist (short)

protocol: no agonist (long) protocol was used. Human chorionic gonadotropin (i.m

10,000 IU hCG; Pregnyl; MSD, USA) was administered when at least one follicle

measured ≥ 17 mm in diameter (averaged orthogonal measurements). The

endometrial thickness, peak estradiol and progesterone levels were assessed on the

day of human chorionic gonadotropin (hCG) trigger. Ultrasound-guided trans-vaginal

oocyte retrieval was performed 36 hours after hCG administration. The effect of

DHEA supplementation on the markers of ovarian reserve (anti-müllerian hormone;

AMH), and follicular function (IGF-1)[30], as well as ovarian follicular levels of estradiol,

testosterone, and DHEA collected from the lead follicle at the time of OPU were

assessed through ELISA as previously described.[31]

The embryo transfer was performed on day 2 or day 3 of embryo-culture, and luteal

phase support was achieved with vaginal progesterone (micronized progesterone,

Utrogestan, 200 mg three times a day, Besins-International, France). Pregnancy was

established by serum beta-hCG seventeen days post embryo transfer. Clinical

pregnancy will be established by a transvaginal ultrasound four weeks after embryo

transfer. IVF/ICSI primary and secondary outcomes are shown in **Table 3**.

Sample preparation

For untargeted metabolomics analysis, sample preparation followed a previously

published report with some modifications.[32] FF were available for metabolomics and

cytokine analyses (DHEA+, *N*=18 and DHEA-, *N*=16). A volume of 50 µL from each

FF sample was thawed at 4°C, and FF proteins were precipitated with 200 µL ice-

cold methanol. After vortexing, the mixture was centrifuged at 16,000 rpm for 10 min

at 4°C and the supernatant was collected and evaporated to dryness in a

speedvacuum evaporator. The dry extracts were then re-dissolved in 200 µL of

water/methanol (98:2; v/v) for liquid chromatography-tandem mass spectrometry

(LC-MS/MS) analysis.

A pooled quality control (QC) sample was generated to allow comparison of analytic

behavior over long periods of time. The pooled reference samples were for the

purposes of quality control (i.e., to ensure relative consistency among identical

samples within days) and for quality assurance (i.e., to ensure consistent results

between days). They did not contribute data to downstream statistical analysis.

Liquid Chromatography-Tandem Mass Spectrometry-based Metabolomics

The supernatant fraction from sample preparation step was analyzed using Agilent

1290 ultra-high pressure (performance) liquid chromatography system (Waldbronn,

Germany) equipped with a Bruker impact II Q-TOF mass spectrometer with its

normal electrospray ionization (ESI) ion source (Bruker Daltonics). 2.5 µL of samples

was injected and were separated using Waters Acquity HSS T3 (2.1 mm i.d. x 100

mm, 1.8 μm) at a flow rate of 0.2 mL/min. The oven temperature was set at 50°C.

The gradient elution involved a mobile phase consisting of (A) 0.1% formic acid in

water and (B) 0.1% formic acid in methanol. The initial condition was set at 5% B. A

5.5 min linear gradient to 60% B was applied, followed by a 13 min gradient to 98%

B (total 24 min including wash and re-equilibration) at a flow rate of 0.4 ml/min. The

ion spray voltage was set at 4,500 V, and the Dry Temperature was maintained at

°C. The drying nitrogen gas flow rate and the nebulizer gas pressure were set at 8.0

L/min and 26 psi, respectively. Calibration of the system was performed using

sodium formate clusters before data acquisition. The stability of the LC-MS method

was examined and evaluated by sodium formate clusters (1 mM NaOH, 0.1% formic

acid, 50% 2-propanol) infused into the system.

The ESI mass spectra were acquired in positive ion mode. Mass data were collected

between *m/z* 100 and 1000 at a rate of three scans per second. Auto MS/MS was

triggered at 8 Hz with duty cycle of 1.5 s. Threshold was set at 1500 counts, with

active exclusion activated after 3 spectra, released after 0.3 min and overwritten if

the current or previous intensity changes. MS/MS spectra were acquired at collision

energy of 20–50 eV automatically varied by the charge states and the intensities of

the selected precursors. Fragment spectra acquisition was carried out at a scan rate

dependent on the MS precursor intensities - MS/MS spectra for high-intensity

precursors were acquired for a shorter time (90000 counts, 12 Hz) than low-intensity

precursor ions (10000 counts, 6 Hz) thus allowing for a balancing of maximal scan

time and MS/MS spectral quality.

Compound identification

Structure identification was achieved via the following in MetaboScape (version 2.0;

Figure 1): elemental composition was predicted via isotopic pattern following the

rules (i) mSigma of MS1: 20 with tolerance of 5 ppm and (ii) MS2: 50 with tolerance

of 2 mDa of the differential metabolites was searched against Bruker HMDB (Human

Metabolome Database) using a precursor match of ±10 mDa, minimum score of 400

and minimum match score of 250. Progesterone, glycerophosphocholinelinoleic acid

and valine were structurally confirmed using chemical standards.

Multiplex immunoassay analysis

45 cytokines were detected and measured using ProCartaplex (EBioscience, CA,

USA) as previously reported [BDNF, EGF, Eotaxin (CCL11), FGF-2 (FGF basic),

GM-CSF, CXCL1 (GROα), HGF, IFNγ, IFNα, IL-1RA, IL-1β , IL-1α, IL-2, IL-4, IL-5,

IL-6, IL-7, CXCL8 (IL-8), IL-9, IL-10, IL-12 p70, IL-13, IL-15, IL-17A, IL-18, IL-21, IL-

22, IL-23, IL-27, IL-31, CXCL10 (IP-10), LIF, CCL2 (MCP-1), CCL3 (MIP-1α), CCL4

(MIP-1β), βNGF, PDGF-BB, PLGF, CCL5 (RANTES), SCF, CXCL12 (SDF1α),

TNFα, LTA (TNFβ), VEGF-A, VEGF-D].[33] Briefly, 5 μL of FFs were diluted with 5 μL

Universal Dilution Buffer, and mixed with 50 μL of antibody-conjugated, magnetic

beads in a 96 well DropArray plate (Curiox Biosystems, Singapore) and rotated at

450 rpm for 120 min at 25°C while protected from light. Beads were internally dyed

with different concentrations of two spectrally distinct fluorophores and covalently

conjugated to antibodies against the 45 cytokines, chemokines and growth factors.

The plate was washed three times with wash buffer (PBS, 0.05% Tween-20) on the

LT210 Washing Station (Curiox) before adding 10 μL of secondary antibody and

rotating at 450 rpm for 30 min at 25°C protected from light. Subsequently, the plate

was washed three times with wash buffer, and 10 μL of streptavidin-phycoerythrin

added and rotated at 450 rpm for 30 min at 25°C protected from light. The plate was

again washed thrice with wash buffer; 60 μL of reading buffer was then added and

the samples read using the Bio-Plex Luminex 200 (BioRad). The beads were

classified by the red classification laser (635 nm) into its distinct sets, while a green

reporter laser (532 nm) excites the phycoerythrin, a fluorescent reporter tag bound to

the detection antibody. Quantitation of the 45 cytokines was then determined by

extrapolation to a six or seven-point standard curve using five-parameter logistic

regression modelling. Calibrations and validations were performed prior to runs and

on a monthly basis respectively.

Statistical analysis

GraphPad Prism 6 (GraphPad Software Inc.) was used for performing all statistical

analyses. Data were checked for normal distribution using Kolmogorov-Smirnov test.

Unpaired or paired t-test was performed, as appropriate, to determine statistical

significance between groups form normally distributed data. Mann-Whitney U test

was used for non-normally distributed data. For comparing more than three groups,

the data were analyzed using ANOVA test, followed by the *t*-test with Bonferroni

adjustment. *P*<0.05 was considered significant. Metabolomic data was further

analyzed by Principal Component Analysis (PCA) and Partial Least Squares

Regression (PLSR) modelling (Unscrambler X version 10.1) after the normalization

of data by first centering the data to the median and scaling it by division with the

standard deviation. Full cross-validation was applied in PLSR to increase model

performance and for the calculation of coefficient regression values.

Results

LC-MS/MS metabolomics system evaluation

In order to obtain reliable metabolic profiles of the samples, we evaluated the

stability and reproducibility of the LC-MS method by performing Principal Component

Analysis (PCA) on all the samples including the eight pooled quality control samples.

As shown in **Figure S1**, the QC samples clustered in PCA scores plots, and together

with retention time CV% < 0.1 min, peak *m/z* values 3 mDa, and relative standard

deviations of peak areas < 20% (**Table 1**), there was good system stability, mass

accuracy and reproducibility of the chromatographic separation during the whole LC-

MS/MS sequence. In addition, intensity CV% of the identified compounds in pooled

quality control samples are low (average 6%; **Table 1**). PCA hotelling (T^2) revealed

one DHEA+ subject as an outlier (D4) and was removed from further analysis

(**Figure S1**). From a total 2717 time-aligned features, an average of 903 features

was chosen for auto MS/MS mode. From these, a total of 100 metabolites were

identified via chemical standard confirmed HMDB.[34] An average of 65 MS/MS

confirmed metabolites per patient was identified, which was similar in terms of

metabolite identified in either DHEA- or DHEA+ subjects (*p*=0.8; range:59-76

metabolites; **Table S1**).

POR follicular fluid metabolome

In POR subjects, the FF metabolome spanned three orders of magnitude, and was

composed of a diversity of metabolites including, glycerophospholipids and

derivatives (glycerophosphocholine, phosphatidylcholines), fatty acids

(heptadecanoic acid, linoleic acid, vaccenic acid, myristic acid), cholesterols

(isocaproic acid, 7-ketocholesterol), glucocorticoids (11-deoxycortisol or cortexolone,

cortisol, corticosterone), hormones (17-hydroxyprogesterone, deoxycorticosterone,

11α-hydroxyprogesterone, 16-dehydroprogesterone, androstenedione,

epitestosterone, progesterone, pregnenolone). Other metabolites included bile acids

(3b-hydroxy-5-cholenoic acid, 3-oxocholic acid, glycocholic acid), peptides and

derivatives (3-indolepropionic acid), lactones (delta-hexanolactone/caprolactone),

lactic acid, vitamin D3 and sphingosine (**Table 2**).

Altered follicular fluid metabolome in response to DHEA

Between the DHEA+ and DHEA- groups, the FF metabolome profiles were distinct.

Based on the FF metabolomes, PLSR modelling distinguished the DHEA+ subjects

from the controls at 2 PLS components, describing 55% of the X-variance and 38%

of the Y-variance (**Figure 1**). In DHEA- subjects, progesterone was the most

abundant FF metabolite, followed by L-alanine, L-phenylalanine, pyridine, L-leucine,

which the top five metabolites collectively made up close to half (48%) of the FF

metabolome (**Figure 2A**). In DHEA+ subjects, the FF metabolome profile of highly

abundant metabolites was different, with cortisol as the most abundance metabolite,

followed by L-alanine, L-phenylalanine, pyridine, L-isoleucine and L-leucine. These

top six metabolites collectively made up ~49.5% of the DHEA+ FF metabolome

(**Figure 2B**). Interestingly, pyridine considered a non-endogenous metabolite

(HMDB0000926), was found in such abundance which suggested it came from the

synthesis of DHEA.[35] Notably, the observed MS/MS spectra of pyridine at various eV

matched very well with HMDB database (**Figure S2A**), which suggested its correct

identification. As a precursor to testosterone and estrogen, DHEA could be

converted to testosterone, and aromatized to estrogen; in the case of POR,

exogenous DHEA was proposed to increase androgens in promoting folliculogenesis

and potentiate the effects of gonadotropins.[8,36,37] FF testosterone was detected in

our metabolomics profiling, although the differences between DHEA+ and DHEA-

group were small [DHEA-: mean signal intensity= 2294.5; DHEA+: mean signal

intensity=2267.75 (testosterone), $p>0.05$; **Figure S2B**].

Four FF metabolites, namely, glycerophosphocholine, linoleic acid, progesterone,

and L-valine were significantly lower in DHEA+ relative to DHEA- ($p<0.05$-0.005;

Figure 3A-D). Interestingly, pregnenolone, a cholesterol metabolite and steroid that

is upstream of DHEA metabolism, was detected only in DHEA+ (6/18 subjects), and

not DHEA- (0/16 subjects). Receiver operating characteristic (ROC) analyses of the

four metabolites revealed area under the curve (AUC) ranging from 0.711

(progesterone), 0.730 (glycerophosphocholine), 0.785 (linoleic acid) and 0.818 (L-

valine) ($p<0.05$-0.01; **Figure 3E-H**), suggesting the utility of these FF metabolites in

monitoring DHEA dose modulation. Linoleic acid and L-valine remained significantly

lower in DHEA+ ($p<0.05$, $p<0.001$ for both) when women with endometriosis (N=5)

were removed from analysis, strongly suggesting the significant effect of DHEA on

these metabolites (**Figure S3**).

FF cytokine profile in response to DHEA

Among the 45 immunomodulatory proteins (cytokines, chemokines and growth

factors), 22 were detected in human FF, namely 10 cytokines (IFNγ, IL12p70, IL13,

IL1b, TNFα, IL1Ra, IL5, IL7, IL10, IL18), 6 chemokines (eotaxin, IP-10, MCP1,

MIP1β, SCF, SDF-1α) and 8 growth factors (bNGF, BDNF, EGF, HGF, LIF, PIGF,

VEGF-A, VEGF-D). Among them, FF MCP1, IFNγ, LIF and VEGF-D were significant

lower in DHEA+ compared to DHEA- (p=0.03, 0.014, 0.031, 0.0161 respectively;

Figure 4). No correlation was found between the significant cytokines and

metabolites.

Clinical observations

Because of the treatment, DHEA-sulphate and free testosterone were significantly

higher in the DHEA+ group compared to the DHEA- group (**Figure S4**). **Table 3**

summarizes the study primary and secondary clinical outcomes. There were no

significant differences for other of the matched criteria, consistent with other studies

[38–40], although there were trends of increased retrieved oocytes (5.27 ± 2.9 versus

4.5 ± 3.6) and metaphase II oocytes (4.0 ± 2.5 versus 3.5 ± 2.2) in DHEA+ group.

Discussion

The follicular milieu provides oocytes with a specialized microenvironment that

promotes the developmental competence of oocytes. It has been proposed that

exogenous DHEA supplementation in POR patients with reduced follicle

responsiveness to FSH, can optimize their response to ovarian stimulation for IVF

among POR patients; however, the effect remains controversial.[4,13,41] This LC-

MS/MS metabolomics study extends the human FF metabolome space in terms of

characterization of its constituents, providing new insights into the complexities of

oocyte development especially in POR women,[42] as well as with DHEA

supplementation.[43,44] [16–18] Aside from previously reported constituents of FF such as

linoleic acid,[16] amino acids,[17] and steroids including progesterone, testosterone[45],

this study also captured metabolic products of ovarian steroidogenesis, cholesterols

and glucocorticoids, in the FF. Further, our FF metabolomics distinguished the

DHEA+ group from the DHEA- group. This is despite the paucity of DHEA

supplementation leading to clinical outcomes in this study, which was also observed

by others,[38–40] and that our identification of four significantly changed metabolites

(glycerophosphocholine, linoleic acid, progesterone, and valine) which may alert us

to the metabolic effects of exogenous DHEA supplementation.

Choline and derivatives are an emerging class of metabolites critical in

developmental competence of fertilized oocytes,[18] and glycerophosphorylcholine

was found increased in the DHEA+ group. Glycerophosphorylcholine is formed from

the breakdown of phosphatidylcholine, and is an organic osmolyte, plausibly

affecting concentrations of other constitutes of FF,[15] and regulation of the diffusion of

compounds into FF necessary for folliculogenesis and oogenesis.[46] PORs are known

to exhibit a low diffusion of exogenous gonadotropin into FF, which is correlated with

poor IVF outcomes.[47] It is conceivable that DHEA induced the metabolism of

phosphatidylcholine to glycerophosphorylcholine, and that a different DHEA

supplementation regimen might lead to changes in FF composition, albeit at

unknown effects on clinical outcomes. Progesterone is one of the key hormones for

the progress of the first metiotic division in oocyte maturation, but changes to

progesterone levels with DHEA supplementation has been fraught with

controversies[48]. Our metabolomics study revealed for the first time that DHEA

supplementation led to a decrease in FF progesterone levels, but to what impact

lower progesterone induced by DHEA supplementation has on PORs remains to be

investigated. Valine degradation has been previously reported in a iTRAQ

proteomics study comparing competent versus incompetent buffalo oocyte

proteome.[49] In a metabolomics study, valine metabolism was also identified in bovine

cumulus and cumulus-oocyte-complex-conditioned media that undergo oocyte

maturation;[50] although in both studies, valine was not directly detected in the omics

profiling. In humans, degenerate oocytes or germinal vesicles that failed meiotically

to reach metaphase II deplete valine more than competent oocytes, in other words

lower valine levels in media which is consistent with our results, and suggest

plausible biological roles of valine in oocyte maturation. Interestingly, we noted a

segregation of DHEA+ patients with low and high level of valine, with the high valine

group approaching concentrations of the DHEA- group. Together with valine's high

AUC value, it is tempting to speculate that valine might be a strong biomarker for

monitoring individual DHEA supplementation. Linoleic acid is the most abundant

polyunsaturated fatty acid in the bovine[51] and human FF (**Figure 2**), and varying

concentrations of linoleic acid have reportedly different effects on oocyte maturation.

At a concentration of 100 µM, linoleic acid added to maturation media inhibits bovine

oocyte maturation and subsequent blastocyst development through increasing

prostaglandin E_2 concentration in the medium, decreasing intracellular cAMP,

decreasing phosphorylation of the MAPK1 and AKT and inhibited germinal vesicle

breakdown.[51,52] Conversely, at concentrations at 50 µM or below, linoleic acid

improved oocyte quality by increasing the content of neutral lipids stored in lipid

droplets.[52] Our metabolomics results suggest that FF linoleic acid is another key

biomarker for titrating and monitoring individual DHEA supplementation. Therefore,

from published literature and our data suggest that DHEA might exert protective

effects on oocytes in POR patients, albeit at an insufficient dose or treatment

duration.

The elevated DHEA-sulphate levels coupled with a lack of difference in FF

testosterone with DHEA supplementation suggest the following possibilities in POR

patients: (i) inadequate DHEA conversion to testosterone due to polymorphism in

SULT2A1, CYP19A1 and FMR1 genes,[53] or (ii) long CAG repeats in androgen

receptor gene which is linked to its lower transcriptional activity at the promoters of

genes involved in the metabolism of DHEA to testosterone.[54] The former is unlikely:

in a case-control study involving 94 subjects, androgen secretion was not impaired in

pre-ovulatory follicles of POR compared to normal responders, and similar levels of

follicular testosterone levels was reported.[55] However, ethnicity and genetic

predispositions might play a role as Chinese women are reported to have higher free

androgens and African American women lower,[56] which might explain their

differences in pregnancy rates in association with IVF than those observed among

other ethnic groups. Conversely, long CAG repeats is associated with risk of POR

and oocyte insensitivity to androgenic stimulation,[57] thus hinting a tenable rationale

on the observed similar FF androgen levels between the DHEA- and DHEA+

subjects in this study. The abundance of cortisol in the DHEA+ subjects is an

interesting finding, in particular that DHEA reduces circulating cortisol,[58] indicating

follicular versus systemic difference in how DHEA affects cortisol levels. *In vitro*, it

was noted that DHEA suppresses cortisol activity,[59] including the antagonist effects

of DHEA on the anti-inflammatory responses induced by cortisol via glucocorticoid

receptor-mediated pathways.[60] It is noteworthy that high FF cortisol levels found in

fertilized IVF individuals compared to unfertilized individuals led to the postulation

that oocyte exposure to cortisol is required with oocyte maturation.[61] The higher

levels of FF cortisol observed in DHEA+ subjects therefore argues for a

compensatory response to modulate the ratio of the two hormones to maintain for a

favourable FF response to mature oocytes.[60]

Due to the highly confident identification based on MS/MS, and mass accuracy of

LC-MS/MS-based metabolomics, we were able to distinguish progesterone from

DHEA, an advantage over interference-prone immunoassays that face a cross-

reactivity bioanalytical problem.[48] Similarly, LC-MS/MS-based determination of

androgens was preferred over immunoassays due to strong interference from

DHEA.[62] We did not detect E1 and E2; because for phenolic hydroxyl group of

estrogens to act as proton donors, the signal would be more sensitive in the negative

ion mode electrospray ionization[63] than in the positive ion mode which was used in

this study. Alternatively, derivatization with dansyl chloride, an amine-containing

sulfonyl halide[64,65] can be used for increased sensitivity but is more amenable to

targeted LC-MS/MS.

In mouse models of polycystic ovary syndrome, treatment with DHEA resulted in

increased production of cytokines such as serum TNFα, IL-6, IL12p70, and IFNγ.[66,67]

In this study, DHEA supplementation lead to the reduction of FF IFNγ, LIF, MCP-1,

and VEGF-D levels. It appears that DHEA modulates chemokines and growth factors

in POR FF without a clear Th1 or Th2 immune response as proposed.[40] LIF or

leukemia inhibitory factor is expressed in the ovary and controls follicular growth.[68] It

was reported that LIF suppressed the growth of primary, secondary, and early antral

follicles in cultured ovarian tissues.[69] The authors postulated that LIF produced in the

late antral or graafian follicles is secreted to suppress the growth of the neighbouring

primary, secondary, and early antral follicles as part of follicular growth.[15]

Interestingly, when hCG is administered in rhesus macaques, at 12 h follicular LIF

levels increase and induce follicle rupture and ovulation and decrease at 24 h.[70] In

our study, the number of MII oocytes and oocytes trended higher in the DHEA+

group, suggesting that the biological roles of LIF might have been achieved (follicular

maturation and rupture) but inadequate to generate a clinically significant outcome.

In vitro results suggested that follicles produce VEGF-A, with VEGF-A inducing the

expanding vasculature to support the increased needs growing follicles.[71] The

decrease in VEGF-A in DHEA+ individuals is intriguing. Fisher *et al.*, described that

in cultured follicles, the rise in VEGF-A levels in faster-growing follicles are

dependent on FSH dose and oxygen tension.[72] There have been reports that DHEA

inhibits oxygen consumption in neurons,[73] tempting the postulation that DHEA

inhibited oxygen consumption in follicle that subsequently led to lower production of VEGF-A in DHEA+ individuals. Further, the lack of correlation between the significant cytokines and metabolites suggest that DHEA converting to steroids which then modulate cytokine production within the follicular microenvironment is more complex than originally thought.

In conclusion, our study provided new insights to POR FF at the metabolome level, and as indicated from the FF metabolome analysis, exogenous DHEA to these patients altered the overall metabolome coverage and abundance to four metabolites. Hypotheses generated from this study included plausible mechanisms underlying DHEA metabolism, and the potential utility of glycerophosphocholine, linoleic acid, progesterone, and L-valine as markers to assess DHEA supplementation. Therefore, future directions include targeted quantitative LC-MS/MS approaches to be developed to detect and quantify four "responder" metabolites in approaches similar to those previously conducted on human peritoneal fluids and sera.[74–76] to design treatment based on metabolomics profiles. Steroid hormones including testosterone should also be quantified via LC-MS/MS to establish baseline levels before commencing DHEA supplementation. Further, comparing POR and normal responders will provide further insights to the alteration of the FF metabolome, and reach a deeper understanding of underpinning pathophysiology to POR.

Acknowledgements

This work was supported by SingHealth Foundation (SHF/CTG034/2010) to VVF, and the National Medical Research Council Centre Grant Programme (NMRC/CG/M003/2017) to YHL. JKYC received salary support from Singapore's Ministry of Health's National Medical Research Council (NMRC/CSA(SI)/008/2016).

Disclosure of conflict-of-interest statement

The authors have nothing to disclose.

Authors contribution

VVF, HHT and JCKY recruited patients. JZ performed metabolomics and cytokine

analyses. DB performed statistical analysis. YT provided expert help on

metabolomics. LYH and JCKY supervised the work and resources. All authors

agreed to submission of the manuscript and have agreed to the order of authors.

Reference

(1)   Surrey, E. S.; Schoolcraft, W. B. Evaluating Strategies for Improving Ovarian

Response of the Poor Responder Undergoing Assisted Reproductive

Techniques. *Fertility and Sterility*. 2000, pp 667–676.

https://doi.org/10.1016/S0015-0282(99)00630-5.

(2)   Devine, K.; Mumford, S. L.; Wu, M.; DeCherney, A. H.; Hill, M. J.; Propst, A.

Diminished Ovarian Reserve in the United States Assisted Reproductive

Technology Population: Diagnostic Trends among 181,536 Cycles from the

Society for Assisted Reproductive Technology Clinic Outcomes Reporting

System. In *Fertility and Sterility*; 2015; Vol. 104, pp 612-619.e3.

https://doi.org/10.1016/j.fertnstert.2015.05.017.

(3)   Ferraretti, A. P.; La Marca, A.; Fauser, B. C. J. M.; Tarlatzis, B.; Nargund, G.;

Gianaroli, L.; ESHRE working group on Poor Ovarian Response Definition.

ESHRE Consensus on the Definition of "poor Response" to Ovarian

Stimulation for in Vitro Fertilization: The Bologna Criteria. *Hum. Reprod.* **2011**,

26 (7), 1616–1624. https://doi.org/10.1093/humrep/der092.

(4)   Kolibianakis, E. M.; Venetis, C. A.; Diedrich, K.; Tarlatzis, B. C.; Griesinger, G.

Addition of Growth Hormone to Gonadotrophins in Ovarian Stimulation of Poor

Responders Treated by In-Vitro Fertilization: A Systematic Review and Meta-

Analysis. *Human Reproduction Update.* 2009, pp 613–622.

https://doi.org/10.1093/humupd/dmp026.

(5) Haning, R. V; Flood, C. A.; Hackett, R. J.; Loughlin, J. S.; McClure, N.;

Longcope, C. Metabolic Clearance Rate of Dehydroepiandrosterone Sulfate,

Its Metabolism to Testosterone, and Its Intrafollicular Metabolism to

Dehydroepiandrosterone, Androstenedione, Testosterone, and

Dihydrotestosterone in Vivo. *J. Clin. Endocrinol. Metab.* **1991**, *72* (5), 1088–

1095. https://doi.org/10.1210/jcem-72-5-1088.

(6) Casson, P. R.; Santoro, N.; Elkind-Hirsch, K.; Carson, S. A.; Hornsby, P. J.;

Abraham, G.; Buster, J. E. Postmenopausal Dehydroepiandrosterone

Administration Increases Free Insulin-like Growth Factor-I and Decreases

High-Density Lipoprotein: A Six- Month Trial. *Fertil. Steril.* **1998**, *70* (1), 107–

110. https://doi.org/10.1016/S0015-0282(98)00121-6.

(7) Barad, D.; Gleicher, N. Effect of Dehydroepiandrosterone on Oocyte and

Embryo Yields, Embryo Grade and Cell Number in IVF. *Hum. Reprod.* **2006**,

21 (11), 2845–2849. https://doi.org/10.1093/humrep/del254.

(8) Casson, P. R. Dehydroepiandrosterone Supplementation Augments Ovarian

Stimulation in Poor Responders: A Case Series. *Hum. Reprod.* **2000**, *15* (10),

2129–2132. https://doi.org/10.1093/humrep/15.10.2129.

(9) Burger, H. G. Androgen Production in Women. *Fertil. Steril.* **2002**, *77* (4), S3–

S5. https://doi.org/10.1016/S0015-0282(02)02985-0.

(10) Roy, S.; Mahesh, V. B.; Greenblatt, R. B. Effect of Dehydroepiandrosterone

and Δ4-Androstenedione on the Reproductive Organs of Female Rats:

Production of Cystic Changes in the Ovary. *Nature* **1962**, *196* (4849), 42–43.

https://doi.org/10.1038/196042a0.

(11) Broekmans, F. J.; Soules, M. R.; Fauser, B. C. Ovarian Aging: Mechanisms

and Clinical Consequences. *Endocr. Rev.* **2009**, *30* (5), 465–493.

https://doi.org/10.1210/er.2009-0006.

(12) Barad, D.; Brill, H.; Gleicher, N. Update on the Use of Dehydroepiandrosterone

Supplementation among Women with Diminished Ovarian Function. In *Journal*

of Assisted Reproduction and Genetics; 2007; Vol. 24, pp 629–634.

https://doi.org/10.1007/s10815-007-9178-x.

(13) Kyrou, D.; Kolibianakis, E. M.; Venetis, C. A.; Papanikolaou, E. G.; Bontis, J.;

Tarlatzis, B. C. How to Improve the Probability of Pregnancy in Poor

Responders Undergoing in Vitro Fertilization: A Systematic Review and Meta-

Analysis. *Fertil. Steril.* **2009**, *91* (3), 749–766.

https://doi.org/10.1016/j.fertnstert.2007.12.077.

(14) Bächler, M.; Menshykau, D.; De Geyter, C.; Iber, D. Species-Specific

Differences in Follicular Antral Sizes Result from Diffusion-Based Limitations

on the Thickness of the Granulosa Cell Layer. *MHR Basic Sci. Reprod. Med.*

2014, *20* (3), 208–221. https://doi.org/10.1093/molehr/gat078.

(15) Rodgers, R. J.; Irving-Rodgers, H. F. Formation of the Ovarian Follicular

Antrum and Follicular Fluid. *Biol. Reprod.* **2010**, *82* (6), 1021–1029.

https://doi.org/10.1095/biolreprod.109.082941.

(16) O'Gorman, A.; Wallace, M.; Cottell, E.; Gibney, M. J.; McAuliffe, F. M.;

Wingfield, M.; Brennan, L. Metabolic Profiling of Human Follicular Fluid

Identifies Potential Biomarkers of Oocyte Developmental Competence.

Reproduction **2013**, *146* (4), 389–395. https://doi.org/10.1530/REP-13-0184.

(17) Lan Xia; Zhao, X.; Sun, Y.; Hong, Y.; Yuping Gao, S. H. Metabolomic Profiling

of Human Follicular Fluid from Patients with Repeated Failure of in Vitro

Fertilization Using Gas Chromatography/Mass Spectrometry. *Int. J. Clin.*

Experimenal Pathol. **2014**, *7* (10), 7220–7229.

(18) Wallace, M.; Cottell, E.; Gibney, M. J.; McAuliffe, F. M.; Wingfield, M.;

Brennan, L. An Investigation into the Relationship between the Metabolic

Profile of Follicular Fluid, Oocyte Developmental Potential, and Implantation

Outcome. *Fertil. Steril.* **2012**, *97* (5), 1078-1084.e8.

https://doi.org/10.1016/j.fertnstert.2012.01.122.

(19) Piñero-Sagredo, E.; Nunes, S.; de los Santos, M. J.; Celda, B.; Esteve, V.

NMR Metabolic Profile of Human Follicular Fluid. *NMR Biomed.* **2010**, *23* (5),

485–495. https://doi.org/10.1002/nbm.1488.

(20) Zamah, A. M.; Hassis, M. E.; Albertolle, M. E.; Williams, K. E. Proteomic

Analysis of Human Follicular Fluid from Fertile Women. *Clin. Proteomics* **2015**,

12 (1), 5. https://doi.org/10.1186/s12014-015-9077-6.

(21) Angelucci, S.; Ciavardelli, D.; Di Giuseppe, F.; Eleuterio, E.; Sulpizio, M.;

Tiboni, G. M.; Giampietro, F.; Palumbo, P.; Di Ilio, C. Proteome Analysis of

Human Follicular Fluid. *Biochim. Biophys. Acta - Proteins Proteomics* **2006**,

1764 (11), 1775–1785. https://doi.org/10.1016/j.bbapap.2006.09.001.

(22) Jarkovska, K.; Martinkova, J.; Liskova, L.; Halada, P.; Moos, J.; Rezabek, K.;

Gadher, S. J.; Kovarova, H. Proteome Mining of Human Follicular Fluid

Reveals a Crucial Role of Complement Cascade and Key Biological Pathways

in Women Undergoing in Vitro Fertilization. *J. Proteome Res.* **2010**, *9* (3),

1289–1301. https://doi.org/10.1021/pr900802u.

(23) Hanrieder, J.; Nyakas, A.; Naessén, T.; Bergquist, J. Proteomic Analysis of

Human Follicular Fluid Using an Alternative Bottom-Up Approach. *J. Proteome*

Res. **2008**, *7* (1), 443–449. https://doi.org/10.1021/pr070277z.

(24) Estes, S. J.; Ye, B.; Qiu, W.; Cramer, D.; Hornstein, M. D.; Missmer, S. A. A

Proteomic Analysis of IVF Follicular Fluid in Women ≤32 Years Old. *Fertil.*

Steril. **2009**, *92* (5), 1569–1578.

https://doi.org/10.1016/j.fertnstert.2008.08.120.

(25) Chen, F.; Spiessens, C.; D'Hooghe, T.; Peeraer, K.; Carpentier, S. Follicular

Fluid Biomarkers for Human in Vitro Fertilization Outcome: Proof of Principle.

Proteome Sci. **2016**, *14* (1), 17. https://doi.org/10.1186/s12953-016-0106-9.

(26) Yakin, K.; Urman, B. DHEA as a Miracle Drug in the Treatment of Poor

Responders; Hype or Hope? *Hum. Reprod.* **2011**, *26* (8), 1941–1944.

https://doi.org/10.1093/humrep/der150.

(27) Sönmezer, M.; Özmen, B.; Çil, A. P.; Özkavukçu, S.; Taşçi, T.; Olmuş, H.;

Atabeko□lu, C. S. Dehydroepiandrosterone Supplementation Improves

Ovarian Response and Cycle Outcome in Poor Responders. *Reprod. Biomed.*

Online **2009**, *19* (4), 508–513. https://doi.org/10.1016/j.rbmo.2009.06.006.

(28) Cui, L.; Lu, H.; Lee, Y. H. Challenges and Emergent Solutions for LC-MS/MS

Based Untargeted Metabolomics in Diseases. *Mass Spectrom. Rev.* **2018**.

https://doi.org/10.1002/mas.21562.

(29) Hazeldine, J.; Arlt, W.; Lord, J. M. Dehydroepiandrosterone as a Regulator of

Immune Cell Function. *J. Steroid Biochem. Mol. Biol.* **2010**, *120* (2–3), 127–

136. https://doi.org/10.1016/j.jsbmb.2009.12.016.

(30) Mason, H. D.; Margara, R.; Winston, R. M.; Seppala, M.; Koistinen, R.; Franks,

S. Insulin-like Growth Factor-I (IGF-I) Inhibits Production of IGF-Binding

Protein-1 While Stimulating Estradiol Secretion in Granulosa Cells from

Normal and Polycystic Human Ovaries. *J. Clin. Endocrinol. Metab.* **1993**, *76*

(5), 1275–1279. https://doi.org/10.1210/jcem.76.5.7684393.

(31)   Lee, Y. H.; Goh, W. W. Bin; Ng, C. K.; Raida, M.; Wong, L.; Lin, Q.; Boelsterli,

604   U. a; Chung, M. C. M. Integrative Toxicoproteomics Implicates Impaired

605   Mitochondrial Glutathione Import as an Off-Target Effect of Troglitazone. *J.*

606   *Proteome Res.* **2013**, *12* (6), 2933–2945. https://doi.org/10.1021/pr400219s.

607   (32)   Cui, L.; Fang, J.; Ooi, E. E.; Lee, Y. H. Serial Metabolome Changes in a

608   Prospective Cohort of Subjects with Influenza Viral Infection and Comparison

609   with Dengue Fever. *J. Proteome Res.* **2017**, *16* (7), 2614–2622.

610   https://doi.org/10.1021/acs.jproteome.7b00173.

611   (33)   Peter Durairaj, R. R.; Aberkane, A.; Polanski, L.; Maruyama, Y.; Baumgarten,

612   M.; Lucas, E. S.; Quenby, S.; Chan, J. K. Y.; Raine-Fenning, N.; Brosens, J. J.;

613   Van de Velde, H.; Lee, Y. H. Deregulation of the Endometrial Stromal Cell

614   Secretome Precedes Embryo Implantation Failure. *MHR Basic Sci. Reprod.*

615   *Med.* **2017**, 1–10. https://doi.org/10.1093/molehr/gax023.

616   (34)   Wishart, D. S.; Tzur, D.; Knox, C.; Eisner, R.; Guo, A. C.; Young, N.; Cheng,

617   D.; Jewell, K.; Arndt, D.; Sawhney, S.; Fung, C.; Nikolai, L.; Lewis, M.;

618   Coutouly, M.-A.; Forsythe, I.; Tang, P.; Shrivastava, S.; Jeroncic, K.; Stothard,

619   P.; Amegbey, G.; Block, D.; Hau, D. D.; Wagner, J.; Miniaci, J.; Clements, M.;

620   Gebremedhin, M.; Guo, N.; Zhang, Y.; Duggan, G. E.; Macinnis, G. D.; Weljie,

621   A. M.; Dowlatabadi, R.; Bamforth, F.; Clive, D.; Greiner, R.; Li, L.; Marrie, T.;

622   Sykes, B. D.; Vogel, H. J.; Querengesser, L. HMDB: The Human Metabolome

623   Database. *Nucleic Acids Res.* **2007**, *35* (Database issue), D521-6.

624   https://doi.org/10.1093/nar/gkl923.

625   (35)   Williams, J. R.; Boehm, J. C. Studies on the Synthesis of

626   Dehydroepiandrosterone (DHEA) Phosphatide. *Steroids* **1995**, *60* (4), 333–

336.

(36)   Nielsen, M. E.; Rasmussen, I. A.; Kristensen, S. G.; Christensen, S. T.;

Mollgard, K.; Wreford Andersen, E.; Byskov, A. G.; Yding Andersen, C. In

Human Granulosa Cells from Small Antral Follicles, Androgen Receptor MRNA

and Androgen Levels in Follicular Fluid Correlate with FSH Receptor MRNA.

Mol. Hum. Reprod. **2011**, *17* (1), 63–70.

https://doi.org/10.1093/molehr/gaq073.

(37)   Gleicher, N.; Weghofer, A.; Barad, D. H. The Role of Androgens in Follicle

Maturation and Ovulation Induction: Friend or Foe of Infertility Treatment?

Reprod. Biol. Endocrinol. **2011**, *9* (1), 116. https://doi.org/10.1186/1477-7827-

9-116.

(38)   Wiser, A.; Gonen, O.; Ghetler, Y.; Shavit, T.; Berkovitz, A.; Shulman, A.

Addition of Dehydroepiandrosterone (DHEA) for Poor-Responder Patients

before and during IVF Treatment Improves the Pregnancy Rate: A

Randomized Prospective Study. *Hum. Reprod.* **2010**, *25* (10), 2496–2500.

https://doi.org/10.1093/humrep/deq220.

(39)   Hu, Q.; Hong, L.; Nie, M.; Wang, Q.; Fang, Y.; Dai, Y.; Zhai, Y.; Wang, S.; Yin,

C.; Yang, X. The Effect of Dehydroepiandrosterone Supplementation on

Ovarian Response Is Associated with Androgen Receptor in Diminished

Ovarian Reserve Women. *J. Ovarian Res.* **2017**, *10* (1), 32.

https://doi.org/10.1186/s13048-017-0326-3.

(40)   Zhang, J.; Qiu, X.; Gui, Y.; Xu, Y.; Li, D.; Wang, L. Dehydroepiandrosterone

Improves the Ovarian Reserve of Women with Diminished Ovarian Reserve

and Is a Potential Regulator of the Immune Response in the Ovaries. *Biosci.*

Trends **2015**, *9* (6), 350–359. https://doi.org/10.5582/bst.2015.01154.

(41)   Duffy, J. M.; Ahmad, G.; Mohiyiddeen, L.; Nardo, L. G.; Watson, A. Growth

Hormone for in Vitro Fertilization. *Cochrane database Syst. Rev.* **2010**, No. 1,

CD000099. https://doi.org/10.1002/14651858.CD000099.pub3.

(42)   Emori, M. M.; Drapkin, R. The Hormonal Composition of Follicular Fluid and Its

Implications for Ovarian Cancer Pathogenesis. *Reprod. Biol. Endocrinol.* **2014**,

12 (1), 60. https://doi.org/10.1186/1477-7827-12-60.

(43)   Bedaiwy, M.; Shahin, A. Y.; AbulHassan, A. M.; Goldberg, J. M.; Sharma, R.

K.; Agarwal, A.; Falcone, T. Differential Expression of Follicular Fluid

Cytokines: Relationship to Subsequent Pregnancy in IVF Cycles. *Reprod.*

Biomed. Online **2007**, *15* (3), 321–325. https://doi.org/10.1016/S1472-

6483(10)60346-X.

(44)   Baskind, N. E.; Orsi, N. M.; Sharma, V. Follicular-Phase Ovarian Follicular

Fluid and Plasma Cytokine Profiling of Natural Cycle in Vitro Fertilization

Patients. *Fertil. Steril.* **2014**, *102* (2), 410–418.

https://doi.org/10.1016/j.fertnstert.2014.04.032.

(45)   Wen, X.; Li, D.; Tozer, A. J.; Docherty, S. M.; Iles, R. K. Estradiol,

Progesterone, Testosterone Profiles in Human Follicular Fluid and Cultured

Granulosa Cells from Luteinized Pre-Ovulatory Follicles. *Reprod. Biol.*

Endocrinol. **2010**, *8* (1), 117. https://doi.org/10.1186/1477-7827-8-117.

(46)   Hsueh, A. J. W.; Kawamura, K.; Cheng, Y.; Fauser, B. C. J. M. Intraovarian

Control of Early Folliculogenesis. *Endocr. Rev.* **2015**, *36* (1), 1–24.

https://doi.org/10.1210/er.2014-1020.

(47)   Nagata, Y.; Honjou, K.; Sonoda, M.; Sumii, Y.; Inoue, Y.; Kawarabayashi, T.

Pharmacokinetics of Exogenous Gonadotropin and Ovarian Response in in

Vitro Fertilization. *Fertil. Steril.* **1999**, *72* (2), 235–239.

https://doi.org/10.1016/S0015-0282(99)00228-9.

(48)   Forman, E. J.; Franasiak, J. M.; Scott, R. T. Elevated Progesterone Levels in
        Women on DHEA Supplementation Likely Represent Assay Interference. *J.
        Assist. Reprod. Genet.* **2015**, *32* (4), 661–661. https://doi.org/10.1007/s10815-
        015-0442-1.

(49)   Chen, L.; Zhai, L.; Qu, C.; Zhang, C.; Li, S.; Wu, F.; Qi, Y.; Lu, F.; Xu, P.; Li,
        X.; Shi, D. Comparative Proteomic Analysis of Buffalo Oocytes Matured in
        Vitro Using ITRAQ Technique. *Sci. Rep.* **2016**, *6* (1), 31795.
        https://doi.org/10.1038/srep31795.

(50)   Uhde, K.; van Tol, H. T. A.; Stout, T. A. E.; Roelen, B. A. J. Metabolomic
        Profiles of Bovine Cumulus Cells and Cumulus-Oocyte-Complex-Conditioned
        Medium during Maturation in Vitro. *Sci. Rep.* **2018**, *8* (1), 9477.
        https://doi.org/10.1038/s41598-018-27829-9.

(51)   Marei, W. F.; Wathes, D. C.; Fouladi-Nashta, A. A. Impact of Linoleic Acid on
        Bovine Oocyte Maturation and Embryo Development. *Reproduction* **2010**, *139*
        (6), 979–988. https://doi.org/10.1530/REP-09-0503.

(52)   Carro, M.; Buschiazzo, J.; Ríos, G. L.; Oresti, G. M.; Alberio, R. H. Linoleic
        Acid Stimulates Neutral Lipid Accumulation in Lipid Droplets of Maturing
        Bovine Oocytes. *Theriogenology* **2013**, *79* (4), 687–694.
        https://doi.org/10.1016/j.theriogenology.2012.11.025.

(53)   Shohat-Tal, A.; Sen, A.; Barad, D. H.; Kushnir, V.; Gleicher, N. Genetics of
        Androgen Metabolism in Women with Infertility and Hypoandrogenism. *Nat.
        Rev. Endocrinol.* **2015**, *11* (7), 429–441.
        https://doi.org/10.1038/nrendo.2015.64.

(54)   Borgbo, T.; Macek, M.; Chrudimska, J.; Jeppesen, J. V.; Hansen, L. L.;

Andersen, C. Y. Size Matters: Associations between the Androgen Receptor CAG Repeat Length and the Intrafollicular Hormone Milieu. *Mol. Cell. Endocrinol.* **2016**, *419*, 12–17. https://doi.org/10.1016/j.mce.2015.09.015.

(55) De Los Santos, M. J.; García-Laez, V.; Beltrán, D.; Labarta, E.; Zuzuarregui, J. L.; Alamá, P.; Gámiz, P.; Crespo, J.; Bosch, E.; Pellicer, A. The Follicular Hormonal Profile in Low-Responder Patients Undergoing Unstimulated Cycles: Is It Hypoandrogenic? *Hum. Reprod.* **2013**, *28* (1), 224–229. https://doi.org/10.1093/humrep/des349.

(56) Sutton-Tyrrell, K. Sex Hormone-Binding Globulin and the Free Androgen Index Are Related to Cardiovascular Risk Factors in Multiethnic Premenopausal and Perimenopausal Women Enrolled in the Study of Women Across the Nation (SWAN). *Circulation* **2005**, *111* (10), 1242–1249. https://doi.org/10.1161/01.CIR.0000157697.54255.CE.

(57) Lledo, B.; Llácer, J.; Ortiz, J. A.; Martinez, B.; Morales, R.; Bernabeu, R. A Pharmacogenetic Approach to Improve Low Ovarian Response: The Role of CAG Repeats Length in the Androgen Receptor Gene. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **2018**, *227*, 41–45. https://doi.org/10.1016/j.ejogrb.2018.06.001.

(58) Kroboth, P. D.; Amico, J. A.; Stone, R. A.; Folan, M.; Frye, R. F.; Kroboth, F. J.; Bigos, K. L.; Fabian, T. J.; Linares, A. M.; Pollock, B. G.; Hakala, C. Influence of DHEA Administration on 24-Hour Cortisol Concentrations. *J. Clin. Psychopharmacol.* **2003**, *23* (1), 96–99. https://doi.org/10.1097/00004714-200302000-00014.

(59) Buoso, E.; Lanni, C.; Molteni, E.; Rousset, F.; Corsini, E.; Racchi, M. Opposing Effects of Cortisol and Dehydroepiandrosterone on the Expression of the

Receptor for Activated C Kinase 1: Implications in Immunosenescence. *Exp.*

Gerontol. **2011**, *46* (11), 877–883. https://doi.org/10.1016/j.exger.2011.07.007.

(60) Kamin, H. S.; Kertes, D. A. Cortisol and DHEA in Development and

Psychopathology. *Horm. Behav.* **2017**, *89*, 69–85.

https://doi.org/10.1016/j.yhbeh.2016.11.018.

(61) Keay, S. D. Higher Cortisol:Cortisone Ratios in the Preovulatory Follicle of

Completely Unstimulated IVF Cycles Indicate Oocytes with Increased

Pregnancy Potential. *Hum. Reprod.* **2002**, *17* (9), 2410–2414.

https://doi.org/10.1093/humrep/17.9.2410.

(62) Heald, A. H.; Butterworth, A.; Kane, J. W.; Borzomato, J.; Taylor, N. F.; Layton,

T.; Kilpatrick, E. S.; Rudenski, A. Investigation into Possible Causes of

Interference in Serum Testosterone Measurement in Women. *Ann. Clin.*

Biochem. **2006**, *43* (3), 189–195.

https://doi.org/10.1258/000456306776865106.

(63) Guo, T.; Gu, J.; Soldin, O. P.; Singh, R. J.; Soldin, S. J. Rapid Measurement of

Estrogens and Their Metabolites in Human Serum by Liquid Chromatography-

Tandem Mass Spectrometry without Derivatization. *Clin. Biochem.* **2008**, *41*

(9), 736–741. https://doi.org/10.1016/j.clinbiochem.2008.02.009.

(64) Nelson, R. E. Liquid Chromatography-Tandem Mass Spectrometry Assay for

Simultaneous Measurement of Estradiol and Estrone in Human Plasma. *Clin.*

Chem. **2004**, *50* (2), 373–384. https://doi.org/10.1373/clinchem.2003.025478.

(65) Anari, M. R.; Bakhtiar, R.; Zhu, B.; Huskey, S.; Franklin, R. B.; Evans, D. C.

Derivatization of Ethinylestradiol with Dansyl Chloride To Enhance

Electrospray Ionization: Application in Trace Analysis of Ethinylestradiol in

Rhesus Monkey Plasma. *Anal. Chem.* **2002**, *74* (16), 4136–4144.

https://doi.org/10.1021/ac025712h.

(66) Sander, V.; Luchetti, C. G.; Solano, M. E.; Elia, E.; Di Girolamo, G.; Gonzalez,

C.; Motta, A. B. Role of the N, N'-Dimethylbiguanide Metformin in the

Treatment of Female Prepuberal BALB/c Mice Hyperandrogenized with

Dehydroepiandrosterone. *Reproduction* **2006**, *131* (3), 591–602.

https://doi.org/10.1530/rep.1.00941.

(67) Solano, M. E.; Sander, V. A.; Ho, H.; Motta, A. B.; Arck, P. C. Systemic

Inflammation, Cellular Influx and up-Regulation of Ovarian VCAM-1

Expression in a Mouse Model of Polycystic Ovary Syndrome (PCOS). *J.*

Reprod. Immunol. **2011**, *92* (1–2), 33–44.

https://doi.org/10.1016/j.jri.2011.09.003.

(68) Abir, R.; Fisch, B.; Jin, S.; Barnnet, M.; Freimann, S.; Van den Hurk, R.;

Feldberg, D.; Nitke, S.; Krissi, H.; Ao, A. Immunocytochemical Detection and

RT-PCR Expression of Leukaemia Inhibitory Factor and Its Receptor in Human

Fetal and Adult Ovaries. *Mol. Hum. Reprod.* **2004**.

https://doi.org/10.1093/molehr/gah047.

(69) Komatsu, K.; Koya, T.; Wang, J.; Yamashita, M.; Kikkawa, F.; Iwase, A.

Analysis of the Effect of Leukemia Inhibitory Factor on Follicular Growth in

Cultured Murine Ovarian Tissue. *Biol. Reprod.* **2015**, *93* (1).

https://doi.org/10.1095/biolreprod.115.128421.

(70) Murphy, M. J.; Halow, N. G.; Royer, P. A.; Hennebold, J. D. Leukemia

Inhibitory Factor Is Necessary for Ovulation in Female Rhesus Macaques.

Endocrinology **2016**, *157* (11), 4378–4387. https://doi.org/10.1210/en.2016-

1283.

(71) Xu, J.; Bernuci, M. P.; Lawson, M. S.; Yeoman, R. R.; Fisher, T. E.; Zelinski,

M. B.; Stouffer, R. L. Survival, Growth, and Maturation of Secondary Follicles

from Prepubertal, Young, and Older Adult Rhesus Monkeys during

Encapsulated Three-Dimensional Culture: Effects of Gonadotropins and

Insulin. *Reproduction* **2010**. https://doi.org/10.1530/REP-10-0284.

(72) Fisher, T. E.; Molskness, T. A.; Villeda, A.; Zelinski, M. B.; Stouffer, R. L.; Xu,

J. Vascular Endothelial Growth Factor and Angiopoietin Production by Primate

Follicles during Culture Is a Function of Growth Rate, Gonadotrophin Exposure

and Oxygen Milieu. *Hum. Reprod.* **2013**, *28* (12), 3263–3270.

https://doi.org/10.1093/humrep/det337.

(73) Safiulina, D.; Peet, N.; Seppet, E.; Zharkovsky, A.; Kaasik, A.

Dehydroepiandrosterone Inhibits Complex I of the Mitochondrial Respiratory

Chain and Is Neurotoxic In Vitro and In Vivo at High Concentrations. *Toxicol.*

Sci. **2006**, *93* (2), 348–356. https://doi.org/10.1093/toxsci/kfl064.

(74) Lee, Y. H.; Yang, J. X.; Allen, J. C.; Tan, C. S.; Chern, B. S. M.; Tan, T. Y.;

Tan, H. H.; Mattar, C. N. Z.; Chan, J. K. Y. Elevated Peritoneal Fluid

Ceramides in Human Endometriosis-Associated Infertility and Their Effects on

Mouse Oocyte Maturation. *Fertil. Steril.* **2018**.

https://doi.org/10.1016/j.fertnstert.2018.05.003.

(75) Lee, Y. H.; Tan, C. W.; Venkatratnam, A.; Tan, C. S.; Cui, L.; Loh, S. F.;

Griffith, L.; Tannenbaum, S. R.; Chan, J. K. Y. Dysregulated Sphingolipid

Metabolism in Endometriosis. *J. Clin. Endocrinol. Metab.* **2014**, *99* (10),

E1913-21. https://doi.org/10.1210/jc.2014-1340.

(76) Lee, Y. H.; Cui, L.; Fang, J.; Chern, B. S. M.; Tan, H. H.; Chan, J. K. Y. Limited

Value of Pro-Inflammatory Oxylipins and Cytokines as Circulating Biomarkers

in Endometriosis – a Targeted 'omics Study. *Sci. Rep.* **2016**, *6*, 26117.

https://doi.org/10.1038/srep26117.

Tables

Table 1. Baseline characteristics of patients in this study

Table 2. List of identified follicular fluid metabolites

Table 3. Table 2. Primary and secondary outcomes between DHEA+ and DHEA-control groups.

Figures

Figure 1. Partial Least Squares Scores plot of DHEA- and DHEA+ follicular fluid metabolome. Metabolomic data was median centred, and scaled by division with the standard deviation. The follicular fluid metabolome distinguished poor ovarian response subjects on DHEA supplementation (DHEA+, red) and without DHEA supplementation (DHEA-, blue).

Figure 2. Histogram of follicular fluid metabolites in poor ovarian responders with and without DHEA supplementation. Follicular metabolome coverage and metabolite abundance as quantified by untargeted LC-MS/MS metabolomics in (A) DHEA- and (B) DHEA+ poor ovarian response subjects. Metabolites were ranked according to their intensity counts.

Figure 3. Significantly different follicular fluid metabolites in DHEA+ and DHEA- patients. (A-D) Dot plots of significantly different FF metabolites. Student's t-tests were performed and $p<0.05$ is considered statistically significant. (E-F) corresponding receiver operating curve (ROC) analyses of the metabolites. Area under curve (AUC) of the metabolites and their P-values are reported.

Figure 4. Significantly changed follicular fluid cytokines in DHEA+ and DHEA-patients. Among 45 cytokines, chemokines and growth factors measured by multiplex immunoassay, (A) MCP-1, (B) IFNγ, (C) LIF and (D) VEGF-D were

significantly lower in poor ovarian response subjects with DHEA supplementation.

Student's t-tests were performed and p<0.05 is considered statistically significant.

Supplementary Figures

Supplementary Figure 1. Principle component analysis reveals DHEA+4 (arrow) as

a potential outlier and was removed from subsequent analysis.

Supplementary Figure 2. (a) MS/MS spectra of pyridine at increasing eV. (b)

Follicular fluid testerosterone levels as measured by metabolomics. DHEA+, POR

subjects on DHEA supplementation and DHEA- without DHEA supplementation.

Supplementary Figure 3. (a) Dot Plots of Linoleic acid and L-Valine after removal

of women with endometriosis (N=5), (b) ROC curves of Linoleic acid and L-Valine

after removal of women with endometriosis (N=5).

Supplementary Figure 4. Histograms of estradiol, anti-müllerian hormone (AMH),

DHEA-sulphate and insulin Growth Factor-1 (IGFBP-1) concentrations as

determined by immunoassay. NS, not significant.

Figure 1. Partial Least Squares Scores plot of DHEA- and DHEA+ follicular fluid metabolome. Metabolomic data was median centred and scaled by division with the standard deviation. The follicular fluid metabolome distinguished POR subjects on DHEA supplementation (DHEA+, red) and without DHEA supplementation (DHEA-, blue).

Figure 2. Histogram of follicular fluid metabolites in poor ovarian responders with and without DHEA supplementation. Follicular metabolome coverage and metabolite abundance as quantified by untargeted LC-MS/MS metabolomics in (a) DHEA- and (b) DHEA+ subjects, and were ranked according to their intensity counts. The top 50% metabolites are shown, with progesterone being the major differentiating metabolite between the DHEA- and DHEA+ subjects.

Figure 3. Significantly changed follicular fluid metabolites in DHEA+ and DHEA- patients. (a-d) Dot plots of significantly changed metabolites in poor ovarian responder patients. (e-f) Corresponding receiver operating curve (ROC) analyses of the metabolites. Area under curve (AUC) of the metabolites and their *P*-values are reported.

Table 1. Baseline characteristics of patients in this study

	DHEA group (*n* = 28)	Control group (*n* = 24)
Age (year), mean (SD)	37 (3)	36 (4)
BMI (kg/m^2), mean (SD)	23 (5)	24 (5)
Race, *n* (%)		
Chinese	21 (75.0)	14 (58.3)
Malay	0 (0.0)	8 (33.3)
Indian	1 (3.6)	2 (8.4)
Others	6 (21.4)	0 (0.0)
Primary infertility, *n* (%)		
Yes	19 (67.9)	13 (54.2)
No	9 (32.1)	11 (45.8)
Infertility duration (years), median (range)	4 (1-16)	4 (1-11)
Primary infertility diagnosis, *n* (%)		
Male factor	22 (78.6)	17 (70.8)
Tubal factor	1 (3.6)	1 (4.2)
Endometriosis	1 (3.6)	2 (8.3)
Low ovarian reserve	3 (10.6)	4 (16.7)
Others	1 (3.6)	0 (0.0)
Cycle number *n* (%)		
Cycle 1	4 (14.3)	12 (50.0)
Cycle 2 & above	22 (78.6)	12 (50.0)
Basal FSH (IU/L) , mean (range)*	8.4 (4.8-27.7)	6.5 (3.6-16.6)
Estradiol (pmol/L) , mean (range)*	98.7 (37.0-320.0)	92.4 (37.0-273.0)
AMH (ng/ml) , mean (range)*	0.7 (0.2-2.8)	0.8 (0.2-2.7)
Free Testosterone (pmol/L), mean (range)*	2.0 (0.5-14.3)	1.7 (0.9-2.9)
DHEAs (µmol/L), mean (range)*	4.0 (0.5-17.3)	3.8 (0.9-10.0)
Antral follicle count, mean (range)*	4.6 (1-15)	5.1 (0-12)
Ovarian volume RO (cm^3), mean (range)*	7.3 (1.9-24.0)	7.7 (2.0-29.5)
Ovarian volume LO (cm^3), mean (range)*	5.1 (1.8-16.6)	8.2 (2.4-26.7)

*Blood baseline parameters, antral follicle count and ovarian volumes were obtained in 27 patients in the DHEA group and 20 patients in the control group and were summarized by geometric mean (range).

Table 2. List of identified follicular fluid metabolites.

No.	HMDB No.	Accurate mass	Theoretical mass	Compound Name	Chemical formula	Pathway
1	HMDB01859	151.0626	151.0633	Acetaminophen	C8H9NO2	Acetaminophen metabolism
2	HMDB04987	261.1206	261.1325	Alpha-Aspartyl-lysine	C10H19N3O5	Amino acid metabolism
3	HMDB03423 /HMDB0064 1	146.0687	146.0691	D-Glutamine/L-Glutamine	C5H10N2O3	Amino acid metabolism
4	HMDB00714	179.0576	179.0582	Hippuric acid	C9H9NO3	Amino acid metabolism
5	HMDB00161 /HMDB0131 0/ HMDB00271	89.0465	89.0477	L-Alanine/D-Alanine/Sarcosine	C3H7NO2	Amino acid metabolism
6	HMDB00641 /HMDB0342 3	146.0687	146.0691	L-Glutamine/ D-Glutamine	C5H10N2O3	Amino acid metabolism
7	HMDB00687 /HMDB0055 7/HMDB001 72/HMDB01 645	131.0941	131.0946	L-Leucine/L-Alloisoleucine/ L-Isoleucine/L-Norleucine	C6H13NO2	Amino acid metabolism
8	HMDB00696	149.0504	149.0510	L-Methionine	C5H11NO2S	Amino acid metabolism
9	HMDB00162	115.0629	115.0633	L-Proline	C5H9NO2	Amino acid metabolism
10	HMDB00883 /HMDB0004 3	117.0784	117.0790	L-Valine/Betaine	C5H11NO2	Amino acid metabolism
11	HMDB00064 /HMDB0076 6	131.0688	131.0695	Creatine/ N-Acetyl-L-alanine	C4H9N3O2	Arginine, proline, glycine and serine metabolism
12	HMDB00043 /HMDB0214 1	117.0786	117.0790	Betaine/ N-Methyl-a-aminoisobutyric acid	C5H11NO2	Betaine Metabolism

13	HMDB01847	194.0796	194.0804	Caffeine	C8H10N4O2	Caffeine metabolism
14	HMDB01860	180.0642	180.0647	Paraxanthine	C7H8N4O2	Caffeine metabolism
15	HMDB02825	180.0638	180.0647	Theobromine	C7H8N4O2	Caffeine metabolism
16	HMDB00062	161.1048	161.1052	L-Carnitine	C7H15NO3	Fatty acid metabolism
17	HMDB00267 /HMDB00805	129.0419	129.0426	Pyroglutamic acid/ Pyrrolidonecarboxylic acid	C5H7NO3	Glutathione metabolism
18	HMDB00017	183.0528	183.0532	4-Pyridoxic acid	C8H9NO4	Vitamin B6 metabolism
19	HMDB00995	312.2087	312.2089	16-Dehydroprogesterone	C21H28O2	Lipid metabolism
20	HMDB00502	388.2485	406.2719	3-Oxocholic acid	C24H38O5	Lipid metabolism
21	HMDB00308	356.2708	374.2821	3b-Hydroxy-5-cholenoic acid	C5H4N4O3	Lipid metabolism
22	HMDB00501	400.3339	400.3341	7-Ketocholesterol	C27H44O2	Lipid metabolism
23	HMDB00503	372.2659	390.2770	7a-Hydroxy-3-oxo-5b-cholanoic acid	C24H38O4	Lipid metabolism
24	HMDB00784	188.1041	188.1049	Azelaic acid	C9H16O4	Lipid metabolism
25	HMDB00015 /HMDB01547	346.2131	346.2144	Cortexolone/Corticosterone	C21H30O4	Lipid metabolism
26	HMDB01547	346.2142	346.2144	Corticosterone	C21H30O4	Lipid metabolism
27	HMDB00063	362.2093	362.2093	Cortisol	C21H30O5	Lipid metabolism
28	HMDB00631	449.3131	449.3141	Deoxycholic acid glycine conjugate	C26H43NO5	Lipid metabolism
29	HMDB00573 /HMDB03231	282.2564	282.2559	Elaidic acid/Vaccenic acid	C18H34O2	Lipid metabolism
30	HMDB00628 /HMDB00234	288.2078	288.2089	Epitestosterone/Testosterone	C19H28O2	Lipid metabolism

31	HMDB00086	257.1026	257.1028	Glycerophosphocholine	C8H20NO6P	Lipid metabolism
32	HMDB00138	465.3096	465.3090	Glycocholic acid	C26H43NO6	Lipid metabolism
33	HMDB02259	270.2557	270.2559	Heptadecanoic acid	C17H34O2	Lipid metabolism
34	HMDB00666	130.0989	130.0994	Heptanoic acid	C7H14O2	Lipid metabolism
35	HMDB00689	116.0833	116.0837	Isocaproic acid	C6H12O2	Lipid metabolism
36	HMDB00673	280.2397	280.2402	Linoleic acid	C18H32O2	Lipid metabolism
37	HMDB00806	228.2088	228.2089	Myristic acid	C14H28O2	Lipid metabolism
38	HMDB00593	785.5943	785.5935	PC(18:1/18:1)	C44H84NO8P	Lipid metabolism
39	HMDB00847	158.1300	158.1307	Pelargonic acid	C9H18O2	Lipid metabolism
40	HMDB00253	316.2395	316.2402	Pregnenolone	C21H32O2	Lipid metabolism
41	HMDB01830	314.2239	314.2246	Progesterone	C21H30O2	Lipid metabolism
42	HMDB00792	202.1194	202.1205	Sebacic acid	C10H18O4	Lipid metabolism
43	HMDB00933	228.1335	228.1362	Traumatic acid	C12H20O4	Lipid metabolism
44	HMDB03231	282.2551	282.2559	Vaccenic acid	C18H34O2	Lipid metabolism
45	HMDB01877	144.1146	144.1150	Valproic acid	C8H16O2	Lipid metabolism
46	HMDB00876	384.3387	384.3392	Vitamin D3	C27H44O	Lipid metabolism
47	HMDB00182 /HMDB03405	146.1048	146.1055	L-lysine/D-lysine	C6H14N2O2	Lysinuric protein intolerance
48	HMDB01923	230.0937	230.0943	Naproxen	C14H14O3	Naproxen action pathway
49	HMDB06344	264.1103	264.1110	Alpha-N-phenylacetyl-L-glutamine	C13H16N2O4	Phenylacetate Metabolism

50	HMDB00159	165.0784	165.1891	L-Phenylalanine	C9H11NO2	Phenylalanine metabolism
51	HMDB00097	103.0994	104.1075	Choline	C5H14NO	Phosphatidylcholine biosynthesis
52	HMDB00157	136.0380	136.0385	Hypoxanthine	C5H4N4O	Purine metabolism
53	HMDB00289	168.0278	168.0283	Uric acid	C5H4N4O3	Purine metabolism
54	HMDB00926	79.0421	79.0422	Pyridine	C5H5N	Pyridine biosynthesis
55	HMDB00975 /HMDB0005 5	324.1052	342.1162	Trehalose/Cellobiose	C12H22O11	Pyrimidine metabolism
56	HMDB00300	112.0266	112.0273	Uracil	C4H4N2O2	Pyrimidine metabolism
57	HMDB00190 /HMDB0131 1	90.0311	90.0317	L-Lactic acid/D-Lactic acid	C3H6O3	Pyruvate metabolism
58	HMDB00252	299.2818	299.2824	Sphingosine	C18H37NO2	Sphingolipid Metabolism
59	HMDB00374 /HMDB0001 6/HMDB009 20	330.2178	330.2195	17-Hydroxyprogesterone/ Deoxycorticosterone/11a- Hydroxyprogesterone	C21H30O3	Steroid biosynthesis
60	HMDB00929	204.0894	204.0899	L-Tryptophan	C11H12N2O2	Tryptophan metabolism
61	HMDB00197	175.0627	175.0633	Indoleacetic acid	C10H9NO2	Tryptophan metabolism
62	HMDB00183	208.0843	208.0848	L-Kynurenine	C10H12N2O3	Tryptophan metabolism
63	HMDB00158 /HMDB0605 0	181.0735	181.0739	L-Tyrosine/o-Tyrosine	C9H11NO3	Tyrosine metabolism
64	HMDB02302	189.0782	189.0790	3-Indolepropionic acid	HMDB02302	Tryptophan deamination
65	HMDB01924	266.1620	266.1630	Atenolol	C14H22N2O3	Beta1-receptor inhibition
66	HMDB06115	106.0416	106.0419	Benzaldehyde	C7H6O	Oxidoreductase activity

67	HMDB00562	113.0586	113.0589	Creatinine	C4H7N3O	Arginine, proline, glycine and serine Metabolism/creatine catabolism
68	HMDB00453	114.0670	114.0681	Delta-hexanolactone	C6H10O2	Hydroxy acid lactonization
69	HMDB04983	94.0084	94.0089	Dimethyl sulfone	C2H6O2S	Methanethiol metabolism
70	HMDB01888	73.0524	73.0528	N,N-Dimethylformamide	C3H7NO	Tertiary carboxylic acid metabolism
71	HMDB00070	129.0785	129.0790	Pipecolic acid	C6H11NO2	Amino acid metabolism

Table 3. Primary and secondary outcomes between DHEA+ and DHEA - control groups

	DHEA+ group (*n* = 28)	DHEA- control group (*n* = 24)	Effect estimate[$] (95% CI)	*p*-value
Primary outcome				
Clinical pregnancy, *n* (%)	2 (7.1)	3 (12.5)	0.57 (0.10-3.14)	0.652
Secondary outcomes[#]: clinical				
No. of oocytes retrieved, mean (SD)	5.2 (2.9)	4.5 (3.6)	0.7 (-1.4 to 2.9)	0.507
No. of metaphase II oocytes, mean (SD)	4.0 (2.5)	3.5 (2.2)	0.6 (-1.0 to 2.1)	0.459
No. of embryos, mean (SD)	2.7 (2.4)	2.3 (1.8)	0.4 (-1.1 to 1.8)	0.601
Embryos transferred, *n* (%)				
0	5 (23.8)	3 (17.7)	1	
1	9 (42.9)	5 (29.4)	1.03 (0.53 to 2.00)	1
2	7 (33.3)	9 (52.9)	0.78 (0.44 to 1.39)	0.667
Secondary outcomes[#]: biomarkers^				
DHEA-S (µg/ml)	870.73 (96.00-3385.57)	182.45 (47.25-589.19)	4.77 (2.83 to 8.04)	< 0.001
Free Testosterone (pg/ml)	87.68 (19.54-457.86)	25.50 (11.56-202.89)	3.44 (2.12 to 5.59)	< 0.001
Estradiol (x10^5) (pg/ml)	17.2 (6.15-137.00)	14.7 (2.15-32.5)	1.17 (0.67 to 2.02)	0.571
AMH (ng/ml)	1.37 (0.21-12.60)	1.56 (0.39-6.12)	0.88 (0.45 to 1.70)	0.687
IGF1 (ng/ml)	0.23 (0.01-18.84)	0.23 (0.08-14.22)	0.99 (0.37 to 2.63)	0.976

[$] The effect estimate refers to relative risk for clinical pregnancy, absolute mean difference for cycle outcomes and relative mean difference for hormonal biomarkers (ratio of geometric means).

[#] Secondary outcomes were obtained from 24 patients in the DHEA group and 20 patients in the control group who had gone through the IVF treatment.

^ Hormones' concentration in follicular fluid of the leading follicle, are summarized by geometric mean (range).