Transcriptome atlas of *Phalaenopsis equestris*

Anna V. Klepikova[1], Artem S. Kasianov[1], Margarita A. Ezhova[1], Aleksey A. Penin[1,2], Maria D. Logacheva[1,2,3]

[1] Laboratory of Plant Genomics, Institute for Information Transmission Problems of the Russian Academy of Sciences, Moscow, Russia
[2] Faculty of Biology, Lomonosov Moscow State University, Moscow, Russia
[3] Center of Life Sciences, Skolkovo Institute of Science and Technology, Moscow, Russia

Corresponding Author:
Anna V. Klepikova [1]
Bolshoy Karetny per. 19, build. 1, Moscow 127051, Russia
Email address: annklepikova@gmail.com

Abstract

The vast diversity of Orchidaceae together with sophisticated adaptations to pollinators and other unique features make this family an attractive model for evolutionary and functional studies. The sequenced genome of *Phalaenopsis equestris* facilitates Orchidaceae research. Here we present an RNA-seq based transcriptome map of *P. equestris* which covers 19 organs of the plant including leaves, roots, floral organs and shoot apical meristem. We demonstrated the high quality of the data and showed the similarity of *P. equestris* transcriptome map with gene expression atlases of other plants. The transcriptome map can be easily accessed through our database Transcriptome Variation Analysis (TraVA) visualizing gene expression profiles. As an example of the application we analyzed the expression of *Phalaenopsis* "orphan" genes – the ones that do not have recognizable similarity with genes of other plants. We found that about a half of them are not expressed; the ones that are expressed have a predominant expression pattern in reproductive structures.

Introduction

The enormous diversity of orchids traditionally attracts attention of plant biologists. Orchidaceae comprises about 25 thousand of species, which makes it the largest plant taxon (Cai et al., 2015). The diversification of orchids has evolved along with complex pollinator-adapted flower structure (Cozzolino & Widmer, 2005), CAM-photosynthesis and epiphytism (Silvera et al., 2009).

Genome assembly of *Phalaenopsis equestris* (the horse phalaenopsis) (Cai et al., 2015) provided novel opportunities for evolutionary and functional studies of Orchidaceae. Genome assembly was used for the functional studies of transcription factor families (Lin et al., 2016; Valoroso et al., 2019), somatic embryogenesis (Chen et al., 2019), retrotransposon insertions (Hsu et al., 2019), as well as for evolutionary studies of ancient polyploidy (Barrett et al., 2019). However,

transcriptome resources of *P. equestris* remain limited even though the de novo transcriptome
assembly was performed based on RNA sequencing of 11 organs (Niu et al., 2016).
In our study we present a transcriptome map of *P. equestris* consisting of 19 samples in two
biological replicates. High-quality RNA of orchid organs and tissues was sequenced using
Illumina technology resulting in 1 687 M reads. We compared expression characteristics of
P. equestris transcriptome map with gene expression atlases of other plants to provide evidence
of reliability of our data. Transcriptome map of *P. equestris* can be applied in a great variety of
functional studies.
49

Materials & Methods

Growing conditions
Plants were grown in a climate chamber under a 16 h light/8 h dark cycle at 22°C and 50–60%
relative humidity. Samples were collected in two biological replicates; each replicate consists of
at least seven plants. Sample collection was performed within two hours (Zeitgeber time ZT8-10)
to reduce the influence of the circadian cycle.
RNA extraction, library preparation and sequencing
RNA was extracted using the RNeasy mini kit (Qiagen, The Netherlands) following the
manufacturer's protocol. To ensure a high quality of *Phalaenopsis* samples, RNA was analyzed
using capillary electrophoresis on Agilent Bioanalyzer 2100. cDNA libraries for Illumina
sequencing were constructed using the NEBNext Ultra II RNA Library Prep Kit for Illumina
(New England BioLabs, MA, USA) following the manufacturer's protocol in 0.5 of the
recommended volume (due to low RNA quantity in such samples as shoot apical meristem).
cDNA libraries were sequenced with the HiSeq4000 and NextSeq500 (Illumina, CA, USA)
instruments (50 bp and 75 bp single read run).
Read mapping
Read trimming was performed using Trimmomatic version 0.36 (Bolger, Lohse & Usadel, 2014)
in a single read mode and parameters "ILLUMINACLIP:common.adapters.file:2:30:10
LEADING:20 TRAILING:20 SLIDINGWINDOW:4:15 MINLEN:30". For read mapping
genome assembly and annotation of *P. equestris* from PLAZA database (version 4.5) was used.
Trimmed reads mapped on the genome assembly using Spliced Transcripts Alignment to a
Reference (STAR) version 2.4.2 (Dobin et al., 2013) in the "GeneCounts" mode and parameters
"--sjdbOverhang 59 --sjdbGTFfeatureExon exon --sjdbGTFtagExonParentTranscript gene_id" to
obtain counts of uniquely mapped reads on each gene.
Expression characteristics of transcriptome map
Gene read counts obtained with STAR were normalized on library size using size factors, as
described in (Anders & Huber, 2010). A threshold of five or higher normalized read counts in
each biological replicate was used to define expressed genes.
To describe gene expression pattern Shannon entropy values were calculated for expressed in at
least one sample genes (Schug et al., 2005). In order to avoid overrepresentation of certain plant
organs, the samples were grouped using distances on clustering tree: gene expression levels were

averaged if samples had distance (1 - Pearson r^2) less than 0.1. Sample groups are listed in Table
S1.
Data availability
The RNA-seq raw data of transcriptome map were deposited in NCBI Sequence Read Archive
(SRA) under BioProject accession PRJNA667255. The TraVA database can be accessed at
http://travadb.org/browse/Species=Phalaenopsis_equestris/.
87

Results

Transcriptome map construction
Ornamental orchid *P. equestris* comprises three varieties and numerous hybrids of various flower
colors and sizes (Hsu & Chen, 2016). To create transcriptome atlas we chose *P. equestris* var.
blue (orchidee.su) as clonal plants are available for the cultivar which helps to reduce
interindividual variability. We have collected 31 samples covering main plant organs and
developmental stages such as roots, young and mature leaves, floral organs, flower buds, and
meristems. Each sample was collected in two biological replicates, and each replicate was pooled
from at least seven plants. Sample RNA was sequenced on Illumina platform resulting in 29 M -
65 M raw single reads (38 M median) for each sample (for sequencing statistics see Table S2).
After removing low-quality reads and technical sequenced 98.7-99.8% of reads remained (Table
S2).
Reads were mapped on the reference genome of *P. equestris* (Cai et al., 2015) with only one
match allowed (unique mapping); 9.2-89.6% of high-quality reads were successfully mapped
(Table S2). 12 samples showed extremely low percentage of read mapping; unmapped reads
were identified as sequences belonging to Cymbidium mosaic virus (GenBank accession
MK816927) which are known to persist in the majority of *P. equestris* population and affect
mainly mature and senescent tissues (Koh, Lu & Chan, 2014). As the library size of infected
samples was insufficient and can distort the conclusions we excluded samples with a percentage
of mapped read lower than 35% in at least one biological replicate. The remained samples had
37.3-89.6% of uniquely mapped reads with median of 81.6%.
Thus, we constructed a transcriptome map of *P. equestris* covering 19 organs and parts of the
plant. Floral organs (anthers, labellum, inner and outer tepals), leaves at different developmental
stages, axes (inflorescence and pedicel), shoot apical and inflorescence meristems, and root parts
were taken into analysis (for detailed description of samples see Table S3). The biological
replicates showed high consistency (median Pearson $r^2 = 0.99$, Table S4).
Clustering of samples generally reflects plant body plan and groups organs with similar
morphology and physiology (Klepikova & Penin, 2019). Hierarchical clustering of *P. equestris*
samples showed the same pattern (Fig. 1A). Sample clusters were formed by floral organs, leaf
parts, meristems and young leaves, inflorescence internode and root; young and mature anthers
were an outgroup for the other samples, similar to *A. thaliana*, rice, and maize (Nobuta et al.,
2007; Wang et al., 2010; Stelpflug et al., 2016; Klepikova et al., 2016). The distances between
samples on clustering tree were closer than in other species we observed (Klepikova et al., 2016;

Penin et al., 2019), which can be explained by the lack of older tissues in *P. equestris*
transcriptome atlas.

123
Figure 1. Expression characteristics of the P. equestris transcriptome map: (a) Hierarchical
clustering tree of transcriptome map samples; (b) The distribution of genes by the number of
samples where gene is expressed. Only expressed genes with 5 or more normalized read counts
in each biological replicate were considered; (c) The distribution of Shannon entropy of P.
equestris genes.
We compare our samples with publicly available *P. equestris* transcriptomes (Table S5). In
general, the clustering of samples was consistent (Fig. S1), though leaf and column from the
BioProject PRJNA288388 (Niu et al., 2016) form outgroup to all other samples.
Expression characteristics of *P. equestris*
Phalaenopsis genome annotation (PLAZA database, version 4.5) includes 29 431 protein-coding
genes. Among them 14 174 (48%) genes were expressed in all samples (using five reads in each
biological replicate as a threshold), when transcripts of 21 671 (74%) genes were found in at
least one sample. These values are in the range of typical expressed gene numbers across plant
transcriptome maps (Klepikova & Penin, 2019). As in other species, samples demonstrated
similarity in the number of expressed genes, which varied form 15 612 (53%) in shoot apical
meristem to 18 947 (64%) in ovules before pollination (Table S6).
Expression patterns of *P. equestris* genes
The study of gene expression pattern can shed light on the biological function of the gene and
place it among essential for a plant existence ubiquitously expressed genes or precise regulators
of tissue features – sample-specific expressed genes. We used two approaches to define gene
expression patterns. A number of samples where gene is expressed is the simplest method to

characterize expression pattern width, as was shown for *Nicotiana tabacum* (Edwards et al.,
2010) or *Vigna unguiculata* (Yao et al., 2016). The majority of genes (16 486) were expressed in
17 or more samples; the second peak (1 896 genes) of the distribution is formed by genes
expressed in 3 or less samples (Fig. 1B). The main patterns of tissue-specific genes were anthers
(56% of tissue-specific genes), roots (11%), and meristems (both shoot apical and inflorescence
meristem, 8%). The high number of anther-associated genes are known for *A. thaliana*
(Klepikova et al., 2016) and is expected for *P. equestris* as young and mature anthers are the
most distant samples on clustering tree (Fig. 1A).
While useful, such approach depends on an arbitrary threshold which separates expressed and
non-expressed genes and does not take into account the variation of expression level between
samples. To overcome the issue, we used Shannon entropy as a measure of expression pattern
width: low entropy values correspond to tissue-specific genes, while high values mark
ubiquitously expressed genes (Schug et al., 2005). The distribution of Shannon entropy in
P. equestris was significantly skewed to the right revealing major part of wide-expressed genes
(Fig. 1C) similarly to *A. thaliana*, *Solanum licopersicum* and *Zea mays* (Sekhon et al., 2013;
Klepikova et al., 2016; Penin et al., 2019).
Using Shannon entropy value lower than 0.25 we identified 521 tissue-specific genes. As in case
of direct count the majority of genes was associated with anthers or roots (Fig. 2). According to
GO enrichment, genes uniquely expressed in the mature anthers were involved in cell wall
organization, biogenesis and modification and had pectinesterase and enzyme inhibitor activity
(Table S7). Young anthers were characterized by genes encoding products with amine and amino
acid binding activity (Table S8). Root-specific genes (expressed in the sample "Root without
apex") were described by terms "response to chemical stimulus", "response to oxidative stress",
"oxidation reduction", and "heme binding" (Table S9).

Figure 2. The heatmap of tissue-specific genes. Expression levels of each gene in each sample
were normalized on its maximal expression level.

To find genes with opposite behaviour which uniformly expressed across tissues we selected
genes with Shannon entropy 3.55 or higher and calculated coefficient of variance (CV) as a
measure of expression stability. For 899 out of 1 340 genes CV was less than 0.25, indicating
uniform expression in all samples and biological replicates. Stable genes had GO enrichment in
terms associated with vesicles, membranes, RNA processing and localization. The list of GO
categories strongly overlapped with the enrichment of *A. thaliana* uniformly expressed genes
indicating inter-species universality of basic biological processes (Table S10).

P. equestris **Transcriptome Variation Database**
We aimed to make our transcriptome data easily accessible and ready to use, so we uploaded
P. equestris transcriptomes into our database Transcriptome Variation Analysis (TraVA,
http://travadb.org/browse/Species=Phalaenopsis_equestris/). TraVA interface demonstrates a
color chart of gene expression profiles in a single- or multiple-gene view. A user can prefer to
show or hide expression values in a chart and choose between several types of read count
normalization (Fig. 3).

Figure 3. Database view.
The application of the TraVA database to the characterization of orchid genes
Graphical interface of TraVA facilitate gene expression patterns analysis and comparison and
can be widely used in *P. equestris* functional studies. Orchids are a large and highly diverse plant
family whose species adapted to a number of ecological niches (typical terrestrial plants,
epiphytes, non-photosynthetic plants). These adaptations reflect in their genome – for example,
P. equestris which has sophisticatedly differentiated perianth the number of *AP3* orthologs is
higher compared to *Apostasia schenzhenica*, the basal orchid species with undifferentiated
perianth. Vice versa, *P. equestris* which is an epiphyte and does not develop typical terrestrial
roots, lacks *AGL12* and several genes of the *ANR1* clade, in contrast to *A. schenzhenica*. This
stresses the importance of the study of lineage-specific genes and gene families. For the genes
that do not have orthologs in model species the analysis of the expression profiles is the first step
towards functional characterization.
We identified 181 (160 after filtering of the proteins that had X on more than 50% of length)
P. equestris proteins that do not have significant similarity to any *Arabidopsis* protein (e-value
cut-off = 10). Out of them 118 share similarity with the proteins of *A. schenzhenica* and are thus
presumably orchid-specific while 42 have no hits and thus emerged after the divergence of
Apostasioideae and Epidendroideae. The survey of the expression profiles showed that 93 of

them are not expressed in any of the samples of the map (Fig. S2). Among the ones which are
expressed most are expressed at very low levels. Higher expression levels are associated with
reproductive structures, in particular, anthers (Fig. 4).

Figure 4. The heatmap of P. equestris-specific genes. Expression levels of each gene in each
sample were normalized on its maximal expression level for the color key. The numbers on the
figure represent normalized gene read count averaged over biological replicates.
Among vegetative structures the most distinct is root (root apex) where three genes –
PEQU_39433, *PEQU_02900*, *PEQU_33696* – have the highest expression levels. *Phalaenopsis*
roots are unique (compared to most other plants, including *A. schenzhenica*, but not to other
epiphytic orchids) in many respects – in particular, they are photosynthetic and develop a special

structure called velamen. Velamen is a tissue of epidermal origin that consists of several layers
of dead cells which help to absorb water and protect photosynthetic tissues of the root from the
UV damage. Notably, *PEQU_39433* and *PEQU_33696* do not have homologs in
A. schenzhenica. *PEQU_02900* has a marginal similarity (34%) with *A. schenzhenica* protein
encoded by Ash001570 gene.
The topic of orphan genes – the ones that lack detectable homologues in other lineages - is
widely discussed, in particular in application to plants (Arendsee, Li & Wurtele, 2014). While a
part of the orphan genes might represent the artifacts of the annotation, others have a function
(for example, *A. thaliana* orphan gene *QQS* which acts in starch metabolism (Li et al., 2009).
The functional analysis of orphan genes however lags behind the typical genes; they are
overlooked in the annotations based on the homology; they are also usually expressed at lower
levels and in a narrower range of tissues (reviewed in Schlötterer, 2015). The study of expression
levels and patterns of a potential orphan gene is a first step towards its characterization – the
detectable level of expression is an evidence of that the ORF is indeed a gene, not an annotation
artifact.
Notably that orphan genes in the well-characterized animal objects (Drosophila, primates) have
expression patterns biased towards male reproductive structures (Begun et al., 2007; Xie et al.,
2012). According to the "out-of-testis" hypothesis (Kaessmann, 2010) this is mediated by the
unique epigenetic state of the chromatin during male gametogenesis. We observed the same bias
in *Phalaenopsis*; the growing availability of plant transcriptome maps will enable to find out if
this is universal for plants.

Conclusions

In this study we present a transcriptome map of orchid *Phalaenopsis equestris* covering 19
organs at various stages of the development. We identified 521 tissue-specific genes the majority
of which expressed in anthers, roots (11%), and meristems. The uniformly expressed genes were
associated with the similar processes as in *Arabidopsis thaliana*, i.e. vesicles, membranes, RNA
processing and localization. In order to improve reuse of the data we integrated transcriptome
map in our database TraVA and demonstrated its usability in the study of *P. equestris* orphan
genes.
245

References

Anders S, Huber W. 2010. Differential expression analysis for sequence count data. *Genome*
Biology 11:R106. DOI: 10.1186/gb-2010-11-10-r106.
Arendsee ZW, Li L, Wurtele ES. 2014. Coming of age: orphan genes in plants. *Trends in Plant*
Science 19:698–708. DOI: 10.1016/j.tplants.2014.07.003.
Barrett CF, McKain MR, Sinn BT, Ge X-J, Zhang Y, Antonelli A, Bacon CD. 2019. Ancient
Polyploidy and Genome Evolution in Palms. *Genome Biology and Evolution* 11:1501–
1511. DOI: 10.1093/gbe/evz092.

Begun DJ, Lindfors HA, Kern AD, Jones CD. 2007. Evidence for *de Novo* Evolution of Testis-
Expressed Genes in the *Drosophila yakuba* / *Drosophila erecta* Clade. *Genetics* 176:1131–
1137. DOI: 10.1534/genetics.106.069245.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics (Oxford, England)* 30:2114–2120. DOI:
10.1093/bioinformatics/btu170.

Cai J, Liu X, Vanneste K, Proost S, Tsai W-C, Liu K-W, Chen L-J, He Y, Xu Q, Bian C, Zheng
Z, Sun F, Liu W, Hsiao Y-Y, Pan Z-J, Hsu C-C, Yang Y-P, Hsu Y-C, Chuang Y-C,
Dievart A, Dufayard J-F, Xu X, Wang J-Y, Wang J, Xiao X-J, Zhao X-M, Du R, Zhang G-
Q, Wang M, Su Y-Y, Xie G-C, Liu G-H, Li L-Q, Huang L-Q, Luo Y-B, Chen H-H, Van de
Peer Y, Liu Z-J. 2015. The genome sequence of the orchid *Phalaenopsis equestris*. *Nature*
Genetics 47:65–72. DOI: 10.1038/ng.3149.

Chen J-C, Tong C-G, Lin H-Y, Fang S-C. 2019. *Phalaenopsis LEAFY COTYLEDON1*-Induced
Somatic Embryonic Structures Are Morphologically Distinct From Protocorm-Like
Bodies. *Frontiers in Plant Science* 10:1594. DOI: 10.3389/fpls.2019.01594.

Cozzolino S, Widmer A. 2005. Orchid diversity: an evolutionary consequence of deception?
Trends in Ecology & Evolution 20:487–494. DOI: 10.1016/j.tree.2005.06.004.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras
TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics (Oxford, England)*
29:15–21. DOI: 10.1093/bioinformatics/bts635.

Edwards KD, Bombarely A, Story GW, Allen F, Mueller LA, Coates SA, Jones L. 2010.
TobEA: an atlas of tobacco gene expression from seed to senescence. *BMC Genomics*
11:142. DOI: 10.1186/1471-2164-11-142.

Hsu C-C, Chen W-H. 2016. The breeding achievements from *Phalaenopsis equestris*. In:
Malayan Orchid Review. WORLD SCIENTIFIC, 41–47. DOI:
10.1142/9789814749954_0007.

Hsu C-C, Su C-J, Jeng M-F, Chen W-H, Chen H-H. 2019. A *HORT1* Retrotransposon Insertion
in the *PeMYB11* Promoter Causes Harlequin/Black Flowers in *Phalaenopsis* Orchids.
Plant Physiology 180:1535–1548. DOI: 10.1104/pp.19.00205.

Kaessmann H. 2010. Origins, evolution, and phenotypic impact of new genes. *Genome Research*
20:1313–1326. DOI: 10.1101/gr.101386.109.

Klepikova AV, Kasianov AS, Gerasimov ES, Logacheva MD, Penin AA. 2016. A High
Resolution Map of the *Arabidopsis thaliana* Developmental Transcriptome Based on
RNA-seq Profiling. *The Plant Journal*. DOI: 10.1111/tpj.13312.

Klepikova AV, Penin AA. 2019. Gene Expression Maps in Plants: Current State and Prospects.
Plants 8:309. DOI: 10.3390/plants8090309.

Koh KW, Lu H-C, Chan M-T. 2014. Virus resistance in orchids. *Plant Science* 228:26–38. DOI:
10.1016/j.plantsci.2014.04.015.

Li L, Foster CM, Gan Q, Nettleton D, James MG, Myers AM, Wurtele ES. 2009. Identification
of the novel protein QQS as a component of the starch metabolic network in *Arabidopsis*
leaves. *The Plant Journal* 58:485–498. DOI: 10.1111/j.1365-313X.2009.03793.x.
Lin Y-F, Chen Y-Y, Hsiao Y-Y, Shen C-Y, Hsu J-L, Yeh C-M, Mitsuda N, Ohme-Takagi M,
Liu Z-J, Tsai W-C. 2016. Genome-wide identification and characterization of *TCP* genes
involved in ovule development of *Phalaenopsis equestris*. *Journal of Experimental Botany*
67:5051–5066. DOI: 10.1093/jxb/erw273.
Niu S-C, Xu Q, Zhang G-Q, Zhang Y-Q, Tsai W-C, Hsu J-L, Liang C-K, Luo Y-B, Liu Z-J.
2016. De novo transcriptome assembly databases for the butterfly orchid *Phalaenopsis*
equestris. *Scientific Data* 3:160083. DOI: 10.1038/sdata.2016.83.
Nobuta K, Venu RC, Lu C, Beló A, Vemaraju K, Kulkarni K, Wang W, Pillay M, Green PJ,
Wang G, Meyers BC. 2007. An expression atlas of rice mRNAs and small RNAs. *Nature*
Biotechnology 25:473–477. DOI: 10.1038/nbt1291.
Penin AA, Klepikova AV, Kasianov AS, Gerasimov ES, Logacheva MD. 2019. Comparative
Analysis of Developmental Transcriptome Maps of *Arabidopsis thaliana* and *Solanum*
lycopersicum. *Genes* 10:50. DOI: 10.3390/genes10010050.
Schlötterer C. 2015. Genes from scratch – the evolutionary fate of de novo genes. *Trends in*
Genetics 31:215–219. DOI: 10.1016/j.tig.2015.02.007.
Schug J, Schuller W-P, Kappen C, Salbaum JM, Bucan M, Stoeckert CJ. 2005. Promoter
features related to tissue specificity as measured by Shannon entropy. *Genome Biology*
6:R33. DOI: 10.1186/gb-2005-6-4-r33.
Sekhon RS, Briskine R, Hirsch CN, Myers CL, Springer NM, Buell CR, de Leon N, Kaeppler
SM. 2013. Maize Gene Atlas Developed by RNA Sequencing and Comparative Evaluation
of Transcriptomes Based on RNA Sequencing and Microarrays. *PLoS ONE* 8:e61005.
DOI: 10.1371/journal.pone.0061005.
Silvera K, Santiago LS, Cushman JC, Winter K. 2009. Crassulacean Acid Metabolism and
Epiphytism Linked to Adaptive Radiations in the Orchidaceae. *Plant Physiology*
149:1838–1847. DOI: 10.1104/pp.108.132555.
Stelpflug SC, Sekhon RS, Vaillancourt B, Hirsch CN, Buell CR, de Leon N, Kaeppler SM. 2016.
An Expanded Maize Gene Expression Atlas based on RNA Sequencing and its Use to
Explore Root Development. *The Plant Genome* 9:0. DOI:
10.3835/plantgenome2015.04.0025.
Valoroso MC, Sobral R, Saccone G, Salvemini M, Costa MMR, Aceto S. 2019. Evolutionary
Conservation of the Orchid MYB Transcription Factors DIV, RAD, and DRIF. *Frontiers*
in Plant Science 10:1359. DOI: 10.3389/fpls.2019.01359.
Wang L, Xie W, Chen Y, Tang W, Yang J, Ye R, Liu L, Lin Y, Xu C, Xiao J, Zhang Q. 2010. A
dynamic gene expression atlas covering the entire life cycle of rice. *The Plant Journal*
61:752–766. DOI: 10.1111/j.1365-313X.2009.04100.x.

Xie C, Zhang YE, Chen J-Y, Liu C-J, Zhou W-Z, Li Y, Zhang M, Zhang R, Wei L, Li C-Y.
2012. Hominoid-Specific De Novo Protein-Coding Genes Originating from Long Non-
Coding RNAs. *PLoS Genetics* 8:e1002942. DOI: 10.1371/journal.pgen.1002942.
Yao S, Jiang C, Huang Z, Torres-Jerez I, Chang J, Zhang H, Udvardi M, Liu R, Verdier J. 2016.
The *Vigna unguiculata* Gene Expression Atlas (VuGEA) from *de novo* assembly and
quantification of RNA-seq data provides insights into seed maturation mechanisms. *The*
Plant Journal 88:318–327. DOI: 10.1111/tpj.13279.