Cross-species analysis between the maize smut fungi *Ustilago maydis* and *Sporisorium reilianum* highlights the role of transcriptional plasticity of effector orthologs for virulence and disease

Weiliang Zuo[1], Jasper RL Depotter[1,*], Deepak K Gupta[2,3,4,*], Marco Thines[2,3,4], Gunther Doehlemann[1]

[1]Institute for Plant Sciences and Cluster of Excellence on Plant Sciences (CEPLAS), University of Cologne, Zuelpicher Str. 47a, 50674 Cologne, Germany; [2]Department for Biological Sciences, Institute of Ecology, Evolution and Diversity, Goethe University Frankfurt am Main, Frankfurt am Main, Germany; [3]Senckenberg Biodiversity and Climate Research Centre, Frankfurt am Main, Germany; [4]Integrative Fungal Research Cluster (IPF), Frankfurt am Main, Germany.

*: D. G and J.D contributed equally to this work

Author for correspondence: Gunther Doehlemann; email: g.doehlemann@uni-koeln.de

Summary

- The constitution and regulation of effector repertoires determines and shapes the outcome of the interaction with the host. *Ustilago maydis* and *Sporisorium reilianum* are two closely related smut fungi, which both infect maize, but cause distinct disease symptoms. Understanding how effector orthologs are regulated in these two pathogens can therefore provide insights to pathogen evolution and host adaption.

- We tracked the infection progress of *U. maydis* and *S. reilianum* in maize leaves, characterized two distinct infection stages for cross species RNA-sequencing analysis and identified 207 out of 335 one-to-one effector orthologs being differentially regulated during host colonization, while transcriptional plasticity of the effector orthologs correlated with the distinct disease development strategies.

- By using CRISPR-Cas9 mediated gene conversion, we identified two differentially expressed effector orthologs with conserved function between two pathogens. Thus, differential expression of functionally conserved genes contributes to species specific adaptation and symptom development. Interestingly, another differentially expressed orthogroup (*UMAG_05318/Sr1007)* showed diverged protein function during speciation, providing a possible case for neofunctionalization.

- Collectively, we showed the diversification of effector genes in related pathogens can be caused both by plasticity on the transcriptional level, as well as through functional diversification of the encoded effector proteins.

Key words: Fungal effectors, cross-species RNA-seq, differentially regulated orthologs, orthogroup, CRISPR-Cas9 gene conversion, *Sporisorium reilianum, Ustilago maydis*

Introduction

During symbiosis, microbes secret effector proteins to facilitate a compatible interaction with their hosts. Effectors hold various functions, such as the suppression of host immunity and host metabolism manipulation to promote host infection. Co-evolution of pathogens with their hosts is driven by an arms race causing selection pressure on perceived effector genes against their recognition by the host immune system (Jones & Dangl, 2006).

Polymorphisms of effectors are diverse, including presence/absence variation, amino acid substitution, epigenetic modification and transcriptional plasticity, which together determine the outcome of the plant-pathogen interaction (Gijzen et al., 2014; Toruño et al., 2016; Franceschetti et al., 2017; Torres et al., 2020). Until now, many effector studies have focused on genome sequencing and effector prediction and variation amongst phylogenetically related pathogens (Raffaele & Kamoun, 2012; Sánchez-Vallet et al., 2018); functional elucidation of single effector proteins (Selin et al., 2016); as well as transcriptional changes during host infection in one-to-one host-pathogen interactions (Lanver et al., 2018). However, information available on the transcriptional regulation of effector orthologs across related pathogen species is scarce.

Smut fungi are a group of fungal plant pathogens consisting of more than 1,500 species, which infect mainly grasses including major cereals such as maize, sorghum, wheat, barley and sugar cane (Begerow et al., 2004). A characteristic feature of many smut fungi is their dimorphic growth, where distinct growth states are observed: saprophytic growth as yeast, and filamentous hyphae which initiate the biotrophic interaction with the host (Kämper et al., 2006a). Smuts often infect their host through the root or coleoptile (Martinez et al., 2002; Laurie et al., 2012), from where the fungi proliferate without evident symptoms during the early vegetative growth stage of the host. When plants switch into the reproductive phase, fungal hyphae residing in floral structure may produce massive black/ brown teliospores (the notorious "smut" phenotype). This type of infection style is also found for the smut fungus *Sporisorium reilianum*, which infects either sorghum (*S. reilianum f. sp. reilianum*) or maize (*S. reilianum f. sp. zeae*) and causes head smut disease. Closely related to this is *Ustilago maydis*, which causes common maize smut disease (Kämper et al., 2006a; Schirawski et al., 2010). While *U. maydis* has the ability to cause disease symptoms on all the aerial maize tissues, *S. reilianum* spreads systemically in the plant while it stays close to the vascular bundles and grows up the main stem axis to reach the cob primordia. *Ustilago. maydis* causes locally restricted plant tumors within a short time of less than 2 weeks under laboratory conditions. Comparative genome analysis of *U. maydis* with *S. reilianum* revealed that the two maize smuts share high genome similarity with regard to gene number and synteny (Schirawski et al., 2010). Furthermore, previous studies suggested that, similar to *U. maydis*, *S. reilianum* can efficiently infect maize leaves to spread systemically and cause head smut disease in floral organs. In recent years, several virulence effectors were identified using seedling infection approaches (Schirawski et al., 2010; Ghareeb et al., 2015;

Schweizer *et al.*, 2018). Together, this renders the two closely related species a well-suited model to
study regulation and function of orthologous genes.

In *U. maydis*, the expression of effector genes is precisely regulated via a network of
hierarchical transcriptional factors (TF), which modulates activity of the transcription of effectors
genes at distinct stages of infection (Skibbe *et al.*, 2010; Lanver *et al.*, 2018). The *b* mating type locus
encoded genes, *bE/bW*, form a heterodimer TF, which not only regulates the expression of 38 effectors,
but also trigger the expression of several TFs including a C2H2 zinc finger TF Rbf1 (regulator of b-
filament) (Heimel *et al.*, 2010b,a), which then turn on the TF Hdp2 (homeodomain transcription factor
2) and Biz1 (*b*-dependent zinc finger protein), together activating the expression of effectors important
for initialing the biotrophic interaction (Flor-Parra *et al.*, 2006; Heimel *et al.*, 2010b). At later
infection stages, the forkhead TF Fox1 controls effectors necessary for full virulence and inhibition of
the host defense after establishment of biotrophic interaction (Zahiri *et al.*, 2010) and the WOPR TF
Ros1 (Tollot *et al.*, 2016) regulate effector gene expression related to fungal sporogenesis in mature
tumors. A comprehensive set of RNA-seq data covering the whole biotrophic growth phase of *U.*
maydis identified distinct expression patterns of effectors and a novel virulence related TF Nlt1 (no
leaf tumors1) (Lanver *et al.*, 2018). The expression of effectors in *U. maydis* was also found to be
regulated in an organ-specific manner (Skibbe *et al.*, 2010), and a more recent study using RNA-
sequencing from laser capture micro-dissected cells revealed a host cell-type specific expression
pattern for several effector genes (Matei *et al.*, 2018).

Reverse genetics studies identified various *U. maydis* effectors holding distinct virulence
functions, and some of these effectors have been functionally characterized on the molecular level
(Lanver *et al.*, 2017). Examples are the apoplastic effectors Pit2 and Pep1, which inhibit the maize
apoplastic Papain-like cysteine proteases (PLCPs) (Mueller *et al.*, 2013; Misas Villamil *et al.*, 2019)
and peroxidase activities (Hemetsberger *et al.*, 2012), respectively. The effectors Cmu1 (Djamei *et al.*,
2011) and Tin2 (Tanaka *et al.*, 2014) are translocated into the host cytoplasm to manipulate host SA
and lignin metabolism, respectively. The leaf-specific effector See1 re-activates host DNA synthesis to
directly promote tumor formation in bundle sheath cells (Redkar *et al.*, 2015a; Matei *et al.*, 2018). In
addition, the secreted repetitive effector protein Rsp3 coats the fungal cell wall and protects it against
a host anti-fungal protein (Ma *et al.*, 2018). In contrast, only one *S. reilianum* specific effector
SAD1suppresses the apical dominance of infected maize to increase branch number has been
functional characterized in *S. reilianum* (Ghareeb *et al.*, 2015), and several species-specific and cluster
19A effectors were identified to contribute virulence (Ghareeb *et al.*, 2019). While these studies
focused on the effectors' function on the protein level, it remains unknown if and how transcriptional
regulation of effectors contributes to virulence and infection style of the pathogen.

Comparative genomic analysis from phylogenetically related plant pathogens suggesting the
effector repertories of smut fungi can be classified as core and accessory effector based on the
conservation level (Schuster *et al.*, 2018; Beckerson *et al.*, 2019; Depotter & Doehlemann, 2020;

Depotter *et al.*, 2020). Core effectors are conserved in all phylogenetically related pathogen and may
be involved in conserved biological process vital for infection, leading to stabilization of such
effectors during speciation (Hemetsberger *et al.*, 2015; Irieda *et al.*, 2019; Thines, 2019).
Complementary, accessory effectors are less conserved and only found in few or individual pathogen
species and these were supposed to have more subtle/specific functions in virulence. Previous studies
suggested that effector orthologs between *U. maydis* and *S. reilianum* are more functionally conserved
compared to *Ustilago hordei* and/or *Melanopsichium pennsylvanicum* (Sharma *et al.*, 2014; Redkar *et*
al., 2015b; Stirnberg & Djamei, 2016).
Clustered regularly interspaced short palindromic repeats (CRISPR)-Cas9 was first identified
in *Streptococcus pyogenes* as part of the bacterial immune system against bacteriophage infection
(Barrangou *et al.*, 2007), and has been widely adapted as genome editing tool in various organisms
including filamentous fungi and oomycetes (Schuster & Kahmann, 2019) . The Cas9 endonuclease
generates DNA double-strand break in a specific target that is recognized by the sgRNA (single guide
RNA). This double strand break can be repaired by non-homologous end-joining (NHEJ) which
eventually results in a gene knock-out mutant. Alternatively, homology direct repair (HDR) with a
donor template can be used for the generation of knock-in or gene conversion mutants. In *U. maydis*,
the CRISPR-Cas9 is achieved by expressing the codon-optimized Cas9 protein and sgRNA in an
autonomous replication plasmid (Schuster *et al.*, 2016, 2018), which shows high efficiency to generate
a selection marker free genome modification mutant. The off-target effects are further reduced by
application of high fidelity variant Cas9HF1 (Zuo *et al.*, 2020).
In this study, we examined the growth of *U. maydis* and *S. reilianum* during leaf infection for
recording similar and divergent biotrophic interaction with maize. We investigated the regulation of
orthologous genes with an emphasis on effector genes.
138
Materials and Methods
Strains, growth conditions and plant infection
The mating compatible isolates of *U. maydis* FB1, FB2 and *S. reilianum* SRZ1, SRZ2 were used for
RNA-seq, and the solo-pathogenic *U. maydis* strain SG200 and its respective knock-out and ortholog
conversion mutants were used for virulence tests. All strains were grown in YEPS light liquid medium
with 200 rpm shaking or potato dextrose agar (PD, Difco) plate at 28°C. *Escherichia coli* strain Top10
was used for cloning purpose and grown in dYT liquid medium or YT agar plate with supplement of
corresponding antibiotics. Maize variety Early Golden Bantam was used for infection. The plants were
grown in a controlled condition of 16 h light at 28 °C and 8 h dark at 22°C. 7 days-old seedlings were
infected with a mixture of compatible *U. maydis*/*S. reilianum* isolates or SG200, and disease
symptoms were scored at 12 days after infection.
150
Staining and microscopy

For microscopy purposes, 0.1% tween-20 was added in the inoculum. To visualize the appressorium,

leaves were stained 20 hours post infection with calcofluor solution (100 µg/ml) for 1 min and briefly

rinsed with water (Lanver *et al.*, 2014). To visualize the fungal growth inside the infected leaves,

WGA-AF488 (Wheat Germ Agglutinin, Alexa Fluor 488) and Propidium Iodide were used for

staining the fungal and plant cell wall, respectively as previously described (Doehlemann *et al.*,

2009a). The microscopy was done on a Nikon Eclipse Ti Inverted Microscope with the Nikon NIS-

ELEMENTS software (Düsseldorf, Germany) and the photos were taken by HAMAMATSU camera.

Confocal microscopy was done by Leica TCS SP8 confocal laser scanning microscope (Leica, Wetzlar,

Germany) and the following filters were used excitation 458 nm and emission 470-490 nm for WGA-

AF488 and excitation 561 nm and emission 590-603 nm for Propidium Iodide.

RNA preparation and RNA-seq

For RNA preparation, 2-cm long of 3^{rd} leaf sections from more than 15 individual plants were

collected for each sample. The compatible haploid *U. maydis* and *S. reilianum* cells from cultures with

OD_{600} around 0.8 were spun down and mixed in a 1:1 ratio as axenic culture (AC) control. The plant

tissues and cell pellets were ground into fine powder with liquid nitrogen and the RNAs were prepared

using TRizol (Thermo Fisher, Waltham, USA) according to the manufacture's protocol and followed

by DnaseI digestion (Thermo Fisher, Waltham, USA). The RNA libraries were prepared using an

Illumina TruSeq Stranded mRNA kit (Illumina, San Diego, USA), and paired-end sequencing was

performed on the a HiSeq4000 platform to produce 2×75 bp long reads at the Cologne Center for

Genomics (Cologne, Germany). Three independent biological replicates have been sampled and used

for RNA-seq analysis.

Data analysis

For the gene expression analysis of *U. maydis* and *S. reilianum* individually, reads of three biological

replicates were filtered using the Trinity software (v2.9.1) option trimmomatic under the standard

settings (Grabherr *et al.*, 2013). They were mapped to a reference assembly using Bowtie 2 (v2.3.5.1)

with the first 15 nucleotides on the 5'-end of the reads being trimmed (Langmead & Salzberg, 2012).

The reference genome was either the genome assembly of *U. maydis* (Kämper *et al.*, 2006b) or *S.

reilianum* (Schirawski *et al.*, 2010) combined with that of *Z. mays* B73 version 3 (Schnable *et al.*,

2009). Reads were counted to the *U. maydis* and *Z. mays* loci using the R package Rsubread (v1.34.7)

(Liao *et al.*, 2019). edgeR package v3.26.8 was used for statistical analysis of differential gene

expression and pairwise comparison was conducted using Generalized linear models (glm) (Robinson

et al., 2009). Genes with log2 fold change >1 and $p<0.05$ were considered as differentially regulated

between timepoints, and genes with log2 fold change <1, $p<0.05$ or log2 fold change >1 but $p>0.05$

were considered as not significantly differenr in the induced pattern analysis. The induced patterns of

effectors were defined based on the differentially regulation relationships between AC, 2 dpi and 4 dpi

samples with following criterias: pattern1 are genes induced *in planta* (expression levels at 2 dpi and 4

dpi were significantly higher than AC samples) but the 4 dpi expression level is higher than 2 dpi;

genes classified as pattern 2 were induced *in planta*, but the expression levels were similar between 2

and 4 dpi; pattern 3 genes were only induced at 2 dpi and had dropped again at 4 dpi to AC level;

pattern 4 genes were significantly induced *in planta* at both timepoints, but the expression level of 2

dpi is higher than 4 dpi; pattern 5 genes are only induced at 4 dpi.

A phylogenetic tree was constructed including the smut species *U. hordei* (Laurie *et al.*, 2012), *M.*

pennsylvanicum (Sharma *et al.*, 2014), *S. scitamineum* (Taniguti *et al.*, 2015), *S. reilianum* (Schirawski

et al., 2010) and *U. maydis* (Kämper *et al.*, 2006a). *Moesziomyces antarcticus* was used as an

outgroup (Morita *et al.*, 2013). A phylogenetic tree was constructed based on 1643 Benchmarking

Universal Single-Copy Orthologs (BUSCOs) from the database "basidiomycota_odb10" that were

present in single copy in all members investigated in the phylogenetic reconstruction (Seppey *et al.*,

2019). For every gene, orthologs were aligned using MAFFT (v7.464) option "--auto" (Katoh &

Standley, 2013). These aligned gene sequences were then concatenated for every species and used for

tree construction using RAxML (v8.2.11) with the substitution model "GTRGAMMA" and 100

bootstraps (Stamatakis, 2014).

For comparison of ortholog expression, high quality paired end transcriptomic reads were mapped

using Bowtie2 with default parameters over all orthologous genes of *Ustilago maydis* and *Sporisorium*

reilianum as calculated using OrthoMCL. Average coverage of mapping of each gene was calculated

from the sorted BAM file by summing the coverage of each base of a gene and then dividing this by

the length of that gene. The relative expression value of a gene was calculated by using the following

formula : $Relative\ Expression\ of\ Gene\ (X) = \dfrac{\frac{\sum_1^n Coverage\,(X) \times (N-n)}{n}}{(M-m)/(N-n)}$, Where n= Length of the

(X) , N = Total number of bases in all genes ; m= Number of bases mapped to gene (X) ; M=

Number of bases mapped to all genes. *Student t-test* and/or one sample *t-test* were used for

significance test with Benjamini-Hochberg *p* value correction for multiple comparison. Orthologs with

fold change >3 and adjust *p*<0.05 were considered as differentially regulated ortholog between *U.*

maydis and *S. reilianum*.

Gene ontology and GO enrichment analysis

The gene ontology classification and overrepresentation analysis were done by using the *U. maydis*

annotation data from PANTHER (pantherdb.org). The overrepresentation was conducted by using

Fisher's exact test with Bonferroni correction.

Quantitative Real-Time PCR

For biomass quantification, the DNA of infected leaves from three biological replicates was prepared

by Buffer A (0.1M Tris-HCl, 0.05M EDTA, 0.5M NaCl, 1.5% SDS), then further purified by

MasterPure Complete DNA and RNA Purification Kit Bulk Reagents (Epicentre, Madison, USA). 100

ng of DNA was used for qPCR, and $2^{-\Delta Ct}$ was calculated to determine the ratio between fungal

peptidylprolyl isomerase (*ppi*) and maize *GAPDH*.

For gene expression, the total RNA was reverse-transcribed with Oligo(dT) primer by using

RevertAid First Strand cDNA Synthesis Kit (Thermo Scientific, Waltham, USA), and the expression

level of each effectors were determined by $2^{-\Delta Ct}$ ratio between effector genes and *ppi* genes from

respective species. The qPCR was conducted in CFX96 Real-Time PCR Detection System (Bio-Rad,

Hercules, USA) with GoTaq qPCR mix (Promega, Madison, USA). The primers designed in this study

were listed in Table. **S1**.

Gene conversion in *U. maydis* by CRISPR and disease scoring

CRISPR-Cas9 mediated genome editing was used to generate the effector ortholog conversion

mutants in *U. maydis* solo-pathogenic strain SG200. An sgRNA was designed to target the coding

region or the promoter region of the effector to generate the double strand break, which was then

repaired by homology-directed repair using a donor plasmid as template. The design of sgRNA and

cloning was done as previously described (Zuo *et al.*, 2020). To construct donor plasmid, 1 kb

fragments flanking the effector coding region or promoter regions and the respective *S. reilianum*

ortholog open reading frame or promoters were amplified by Phusion DNA polymerase (NEB,

Ipswich, USA) with primers (Sigma Aldrich, St. Louis, US) listed in Table. **S1** and cloned into

pAGM1311 vector by Gibson assembly (New England Biolabs, Ipswich, USA). The CRISPR plasmid

and circular donor plasmid were co-transformed into SG200 protoplast and mutants were singled out

on PD plate with 2 µg/ml carboxin for CRISPR plasmid first then transfer to PD plate without

antibiotic to get rid of CRISPR plasmid. The resulting conversion mutants have no antibiotic

resistance. Correct integration of the recombinant DNA was confirmed by southern blot (not shown).

The conversion mutants and corresponding effector deletion mutants (Schilling *et al.*, 2014) were used

for infection. Disease scoring was done as previous described at 12 dpi. Disease indexes 9, 7, 5, 3, 1

and 0 were assigned to dead, heavy tumor, tumor, small tumor, chlorosis and normal symptom,

respectively. The number of diseased plants were multiplied by the corresponding disease index, and

the sum was divided by the total number of plants used for infection to give an average disease index.

Student *t*-test was used for significance test of disease index from three biological replications.

Results

Maize leaf infection of *S. reilianum* and *U. maydis*

For a comparative transcriptomics approach between *S. reilianum* and *U. maydis*, we first tracked

infection and fungal growth inside the maize leaves to record the milestone events during infection

and define appropriate timepoints for analysis (Fig. **1**). 18-24 hours post infection, the compatible

sporidia cells of both pathogens, mated on the leaf surface, and dikaryotic hyphae formed appressoria

to penetrate the leaf surface (Fig. **1a**). At 2 days post infection (dpi), all infected leaves showed

chlorosis as a first visible indication of successful infection (Fig. **S1a**). Microscopic analysis of WGA-

AF488 stained hyphae showed similar colonization in the leaf vascular tissue for both smut fungi (Fig.

1b). At 3 dpi, the earliest microscopic difference was observed between *S. reilianum* and *U. maydis*: in

alignment to our previous report (Matei *et al.*, 2018), host bundle sheath cells underwent *de novo* cell

division in *U. maydis* infected leaves, indicating the initiation of tumorigenesis. In contrast, *S.*

reilianum further accumulated in the leaf vein and showed aggregation as a string of round cells (Fig.

1c). At 4 dpi, the first visible small tumor structures can be observed on *U. maydis* infected leaves,

while *S. reilianum* infected leaves showed individual events of necrosis (Fig. **S1a**) and more

frequently cell aggregation (Fig. **1d**, Fig. **S1b**). In parallel to the microscopic observation, fungal

biomass in the infected leaves was quantified for both pathogens. The relative amount of both fungal

pathogens increased similarly during infection until 3 dpi. Interestingly, at 4 dpi *S. reilianum* showed a

significantly higher abundance compared to *U. maydis,* as the relative amount of *S. reilianum* doubled

while the relative amount of *U. maydis* was maintained in comparison to 3 dpi (Fig. **1e**).

Expression profiling of *U. maydis* and *S. reilianum* during maize leaf colonization

Based on our microscopic and biomass analysis, we decided to collect samples at 2 dpi to represent an

infection stage where both *U. maydis* and *S. reilianum* successfully establish the biotrophic interaction

with the host but do not show detectable differences in microscopical growth and biomass

accumulation. Samples were also taken at 4 dpi when *U. maydis* tumorigenesis is initiated, while *S.*

reilianum exhibits extensive tissue colonization without inducing morphological changes in the leaf.

As control samples, cell pellets from axenic culture (AC) of both pathogens were used for RNA

extraction. All samples have been generated in three independent biological replicates. In total, more

than 670M paired-end reads were generated, of which over 93.7% of reads were uniquely mapped to

either *U. maydis*, *S. reilianum* or the maize genome, respectively. The changes of fungal transcripts in

the RNA-seq samples confirmed the biomass quantification results by qPCR. At 2 dpi, around 2% of

fungal transcripts were mapped from both pathogens, and at 4 dpi, *S. reilianum* reads increased to 12.5%

in the infected samples, which is significantly higher compared to *U. maydis* infected tissues (7.5%)

(Fig. **1f**). Principal component analysis (PCA) analysis of *U. maydis* and *S. reilianum* samples showed

that three biological replications from different conditions form distinct clusters, and variations

between different conditions were larger than the variation of biological replications within the same

condition (Fig. **2a**). To further analyze the RNA-seq data, we normalized reads counts and

investigated the differentially expressed genes (DEGs) by edgeR (Robinson *et al.*, 2009). After

filtering the low- or not-expressed genes between samples, 6537 out of 6765 *U. maydis* genes and

6456 out of 6673 *S. reilianum* genes were used to identify DEGs. Compared to axenic culture,

dramatic changes in gene expression were observed during seedling infection. Around 19.3-24.2%

genes were up-regulated (log2 fold change >1, $p<0.05$) at 2 dpi and 4 dpi in both pathogens, while

12.1-19.1% genes were down-regulated (Fig. **2b**). We detected 435 out of 467 effector genes from *U.*
maydis and 454 out of 489 from *S. reilianum* were expressed. As expected, most effector genes were
specifically expressed in the biotrophic growth stage (311 in *U. maydis* and 307 in *S. reilianum,*
respectively) and showed higher degrees of transcriptional induction compared to non-effector genes
(Fig. **2c, d**), which reflects the explicit role of effectors in host infection. Three biological replicates of
quantitative RT-PCR were conducted to confirmed the expression of several effector genes including
infection markers such as Pit2 and Pep1 from our RNA-seq data (Fig. **S2a, b**).
We clustered effector genes into all five possible expression patterns between three conditions (axenic
culture, 2 dpi, 4 dpi) (Fig. **3a**). This classified groups of effector genes (i) showing constitutively
increasing induction (pattern 1), (ii) induced *in-planta* on a stable level (pattern 2), (iii) being only
transiently expressed (pattern3), (iv) dropping at 4 dpi (pattern 4), (v) or only being induced at 4 dpi
(pattern 5). In *U. maydis* and *S. reilianum*, 227 effectors from each species were highly expressed
across the whole biotrophic growth phases (pattern 1, 2 and 4), while only 14 *U. maydis* effectors and
8 *S. reilianum* effectors were transient induced at 2 dpi (pattern 3) and similar number of effectors
were only required in the 4 dpi (23 from *U. maydis* vs. 25 from *S. reilianum* as pattern 5) (Fig. **3b**).
The expression patterns are found to reflect previously observed virulence function of known effector
genes. For example, the core effector *UmPep1*(*UMAG_01987*) (Doehlemann *et al.*, 2009b;
Hemetsberger *et al.*, 2015) was stably expressed and clustered as pattern 2, while genes coding for the
known virulence factors *UmPit2* (*UMAG_01375*) (Misas Villamil *et al.*, 2019),
UmCmu1(*UMAG_05731*) (Djamei *et al.*, 2011), *UmTin2* (*UMAG_05302*) (Tanaka *et al.*, 2014) and
UmRsp3 (*UMAG_03274*) (Ma *et al.*, 2018) group to pattern 1 cluster. Interestingly, *S. reilianum*
orthologs of these characterized effectors displayed different induction patterns, keeping a steady level
between 2 and 4 dpi. In particular, we identified 100 out 467 of *U. maydis* effectors being induced in
pattern 1, which is significantly higher compared to *S. reilianum* (43/488), whereas 184 out of 488 *S.*
reilianum effector genes were showing stable expressing levels, which is higher as compared to *U.*
maydis (136/467) (Fig. **3b**). This enrichment of different induction patterns of effector genes
suggested different requirements in disease for the two species.
326

Differential regulation of effector orthologues in *U. maydis* and *S. reilianum*
To better understand how *U. maydis* and *S. reilianum* orchestrate their respective effector repertoires
with respect to their differential disease development, we directly compared the relative expression
levels of effector orthologs. We used orthoMCL to identify one-to-one ortholog pairs between *U.*
maydis and *S. reilianum* and obtained 6005 one-to-one ortholog pairs including 335 effectors. This
means that for avoiding ambiguous results due to functional redundancy or paralog compensation,
one-to-many or many-to-many ortholog pairs were not included in this analysis. Relative expression
was corrected for putative differences in length between the orthologs. PCA analysis based on the
relative expression of 6005 ortholog pairs confirmed that plant associated samples form distinct

clusters on time and species levels. In contrast, the axenic cultured samples of the different species

have minor differences and clustered together (Fig. **4a**). Pearson correlation analysis displayed similar

results and suggested the difference between samples increased at 4 dpi compared to 2 dpi, as the

correlation decreased (Fig. S**3a**).

Next, we compared the relative ortholog expression levels to identify differentially expressed

orthologs (DEOs) in the two smut fungi. In total, we found 769 DEOs with comparatively higher

expression in *U. maydis* and 737 DEOs being more highly expressed in *S. reilianum*, respectively (Fig.

4b). The largest number of DEOs (835) was detected from axenic cultured samples (Fig. **4b**), and

around 30 and 50% of these DEOs from *U. maydis* and *S. reilianum* were only confined in this stage

(Fig. S**3b)**. The DEO number dropped to 323 at 2 dpi when the pathogens had switched from *in-vitro*

yeast growth to the initial filamentous, biotrophic growth in the host tissue. At 4 dpi the number

increased to 747, which likely reflects the significant differences in disease development between the

two pathogens, i.e. initiation of tumorigenesis by *U. maydis* versus extended proliferation of *S.*

reilianum for systemic spreading. We also detected more *U. maydis*-induced DEOs during infection,

particularly at 2 dpi, when 227 DEOs were *U. maydis*-induced but only 97 DEOs for *S. reilianum* (Fig.

4b). However, we could not detect overrepresented gene ontology terms, except for 4 dpi, where an

enrichment of cellular lipid catabolic process (GO:0044242) was detected in the higher expressed *U.*

maydis DEOs. The DEOs were involved in different biological processes related to pathogen growth

(Fig. **4c**), suggesting that during the yeast and biotrophic growth no particular cellular or metabolic

pathway was selected to promote cell growth in either *U. maydis* or *S. reilianum*.

From the 335 one-to-one effector orthologs, more than 60% were differentially regulated, including

100 *U. maydis*- and 102 *S. reilianum*-highly expressed effectors, from which 32 and 35 showed

consistently enhanced expression during the two biotrophic time points (Fig. **4d**). The number of

effector DEOs also increased from 99 at 2 dpi to 170 at 4 dpi as disease developed. Previously

identified pathogenesis associated TFs might be responsible in the regulation of effector DEOs,

particularly at 4 dpi, since differential expression was found for virulence related TFs *Rbf1, Biz1,*

*Hdp2, Fo*x1 and *Nlt1* (Fig. **S4**). TF *Rbf1* was highly induced in *S. reilianum*, especially at 4 dpi, the

transcript levels were 17-fold higher than those in *U. maydis*, which may lead to the high abundance of

Biz1 in *S. reilianum* (Fig. **S4**). On the other hand, the expression level of TFs *Hdp2*, *Fox1* and *Nlt1*

were significantly higher in *U. maydis* at 4 dpi, which could indicate that these TFs play a role in the

regulation of effectors highly expressed in *U. maydis* during tumorigenesis (Fig. **S4**). Besides these,

the TF *ROS1* showed similar expression level in both species, which may reflect its dedicated function

in sporogenesis, a process which needs to be initiated in both pathogens for completing the pathogenic

like cycle (Fig. **S4**). The expression of *ROS1* is dramatically increased since 6 dpi in *U. maydis* that is

corresponding to the initiation of sporogenesis (Tollot *et al.*, 2016), which may explain why we did

not observe its differential expression between two species. Cluster 19A (Fig. **S5a**) is the largest

effector cluster, which is a crucial determinant of virulence in both two pathogens (Brefort et al., 2014;

Ghareeb et al., 2019). We observed that 10 out of 14 effectors residing in cluster 19A were *U. maydis*-

enhanced, whereas only two were highly expressed in *S. reilianum,* including the neofunctionalized

Tin2 (Tanaka *et al.*, 2019) (Fig. **S5b**).

In a previous study, Schuster et al. analyzed 12 smut related basidiomycete and suggested that effector

repertoires of smut fungi comprise sets of core and accessory effectors (Schuster *et al.*, 2018). Based

on this information, we classified the 335 one-to one effector pairs into plant smut core effectors,

which were present in all 4 monocot smut fungi, including *Ustilago hordei* and *Sporisorium*

scitamineum, and accessory effectors, which present in *U. maydis* and *S. reilianum* but not in all other

plant smut fungi (Fig. **4e**). *Melanopsychium pennsylvanicum* was included in this analysis due to its

close phylogenetic relationship to monocot smut pathogens, which suggests a recent host jump event

(Sharma et al. 2014). In total, we identified 191 core effectors and 144 accessory effectors. At 2 dpi,

40 core effectors and 31 accessory effectors were differentially expressed between *U. maydis* and *S.*

reilianum, and these numbers were significantly increased to 67 and 61 at 4 dpi, respectively (Fig. **4f**).

We did not find accessory effectors subjected to more intensive transcription regulation compared to

core effectors (Fig. **4f**), suggesting that differential regulation of both core and accessory effectors is

needed for the distinct pathogenic development in the two pathogens.

Functional conservation and neofunctionalization of differentially expressed effector orthologs

Several *U. maydis* effectors have been functionally characterized, which provides a reference to link

the ortholog expression with their biological functions. The conserved core effector Pep1

(Hemetsberger et al., 2012, 2015; Sharma et al. 2019), as well as the virulence factors Pit2 (Misas

Villamil *et al.*, 2019) and Cmu1 (Djamei *et al.*, 2011) were highly induced during biotrophic growth

(Table. **S2**) and showed similar expression level between *U. maydis and S. reilianum*, which is

consistent with their crucial roles in maize immunity inhibition (Fig. **5a**). The effectors Rsp3 (Ma *et*

al., 2018) and ApB73 (Stirnberg & Djamei, 2016) had been validated in previous studies for their

conserved function between two smut fungi by ortholog complement assay, and also these genes show

comparable expression levels (Fig. **5a**). Also the See1 effector, which in *U. maydis* has a virulence

function specifically for leaf tumor-formation (Redkar *et al.*, 2015a), is expressed in *S. reilianum* both

at 2 and 4 dpi (Fig. **5a**). Another effector DEO which caught our attention is Tin2. SrTin2

neofunctionlized during speciation between *U. maydis* and *S. reilianum* (Tanaka *et al.*, 2019)**.** We

found a significantly enhanced expression of *SrTin2* compared to *UmTin2* (Fig. **5a**), in line with the

neofunctionalisation, as the transcriptional level divergence could be involved to promote the

functional divergence.

From a set of 20 organ-specific effectors, we found 9 effector DEOs and 8 of them were leaf-specific

effectors, from which 4 were validated as virulence factors from a deletion mutant screen (Schilling *et*

al., 2014) (Fig. **5a**). Orthogroups *UMAG_11060/Sr14941*, *UMAG_05306/Sr10059* and

UMAG_05318/Sr10075 were particularly interesting as they were induced in both pathogens (Table.

S2), however, the transcripts levels of the orthologs in *U. maydis* were at least 4.8 folds (for
orthogroup *UMAG_11060/Sr14941* at 2 dpi) to more than 50 times (orthogroup
UMAG_05318/Sr10075 at 4 dpi) higher compared to their *S. reilianum* orthologs (Table. S2). These
effectors are specifically required for *U. maydis* induced leaf tumors (Schilling *et al.*, 2014). Thus,
their low expression in *S. reilianum* might suggest that these genes are of lesser importance for the
basal establishment of biotrophic growth. We assessed the protein function of these DEOs to test if
functional diversification occurred and is associated with differentially expression, as in the case of
Tin2. We converted the open reading frame of the candidate effectors in *U. maydis* to their respective
S. reilianum orthologs. For the generation of *in situ* seamless mutants we used a CRISPR-Cas9
approach that allowed marker-free selection of gene-replacement mutants in *U. maydis* (Fig. **5b**). In
each case, an sgRNA was designed to target the effector gene to induce a double strand break, which
was then repaired by HDR with a donor plasmid containing the *S. reilianum* effector ortholog, flanked
by 1kb homologs arms (Fig. **5b**). This CRISPR-Cas9 mediated gene conversion eliminated potential
ectopic effects and allowed expression of the orthologs under the native *in-situ* cis- or trans- regulatory
elements.
In maize infection assays, conversion of UMAG_05306 and UMAG_11060 to their *S. reilianum*
orthologs did not affect the virulence compared to the *U. maydis* progenitor strain SG200, while the
deletion of these two effectors resulted in significantly reduced virulence (Fig. **5d**). However,
conversion of UMAG_05318 into its *S. reilianum* ortholog Sr10075 significantly reduced virulence to
the similar level of the deletion mutant, which indicates that Sr10075 might have a different function,
or lost its activity during speciation (Fig. **5d**). In *S. reilianum*, Sr10075 has a paralog Sr10079, which
has only low sequence identity to UMAG_05318 and therefore was not detected in our one-to-one
ortholog analysis (Fig. **S6a, b**). Deletion of Sr10079 was previously shown to reduce the virulence in *S.
reilianum* (Ghareeb *et al.*, 2019). Conversion of UMAG_05318 into Sr10079 resembled virulence
defect of the UMAG_05318 deletion mutant. Thus, neither Sr10075, nor Sr10079 can functionally
replace   UMAG_05138,   suggesting   the   functional   divergence   of   the   orthogroup
UMAG_05318/Sr10075 - Sr10079 during speciation, which identifies them as an interesting objective
to test for possible neofunctionalization.
To further confirm and investigate how promoters regulate the expression of DEOs, we used the
CRIPSR-Cas9 mediated conversion approach to replace the promoter of the *U. maydis* induced DEOs
UMAG_11060 and *UMAG_05306* with their respective *S. reilianum* counterparts in SG200 (Fig. **S7a**),
and quantitative RT-PCR was used to detect the chimeric gene expression. Infected samples at the
time points when the differential expression levels were detected in RNA-seq (2 dpi for *UMAG_11060*
and 4 dpi for *UMAG_05306*, Fig. **5a**) were tested. Expressing *UMAG_11060* under Pro^{sr14941} showed
lower expression at 2 dpi (Fig. **S7b**), which was similar to our RNA-seq analysis, indicating the cis-
regulation in the promoter region determined their differential expression. However, Pro^{Sr10059}
surprisingly drove a similar expression level of *UMAG_05306* at 4 dpi in *U. maydis* genetic

background, suggesting a trans regulator or distal enhancer, which is variable between two species

control the differential expression (Fig. **S7b**).

Together, the results of the gene conversion approach showed that functional diversification of

effector orthologs can be caused on two levels: (i) the differential expression of orthologous genes

which encode functionally conserved proteins (as found for orthogroups UMAG_05306/Sr10059 and

UMAG_11060/Sr14941), (ii) the functional divergency of effectors on the protein level, which further

fine-tuned on transcription levels for functional adaption (iii) the regulations of DEO is complicated,

involving both cis- and trans-regulatory elements.

Discussion

In this study we applied a cross-species transcriptome comparison between *U. maydis* and *S. reilianum*

to investigate transcriptomic changes of effector orthologs during maize leaf infection.

A crucial challenge in cross-species RNA-seq is to determine the right timing for sampling to exclude

that the observed differences result rather from the different disease progression states than from a

differential regulation as such. Sucher *et al*. (2020) collected samples from the edge of disease lesions,

which were smaller than 25 mm in diameter, as indication of comparable infection stage to study the

different host responses to *Sclerotinia sclerotiorum* for quantitative disease resistance (Sucher *et al.*,

2020). Time resolved microscopes were applied in *Fusarium virguliforme* to facilitate RNA-seq to

study its transcription plasticity on different hosts (Baetsen-Young *et al.*, 2020). In our experiment, we

conducted a series of microscopy combined with qPCR-based biomass quantification, and selected

two biotrophic interaction phases representing a comparable (2 dpi) and a distinct colonization stage (4

dpi) for RNA-seq.

When comparing the induction profiles of effectors in *U. maydis* and *S. reilianum* respectively, we

found a significantly higher number of *S. reilianum* effectors showing stable expression between 2 and

4 dpi, while in *U. maydis* more effectors displayed an increasing expression over time. We further

focused on the expression regulations of one-to-one effector orthogroups. Overall, 207 out of 335 one-

to-one effector orthologs were differentially expressed between *U. maydis* and *S. reilianum* during

biotrophic interaction, and the number of effector DEOs increased at 4 dpi. The difference in effector

induction patterns and differential regulation of effector DEOs are consistent with the appearance of

different symptoms. At 4 dpi, *U. maydis* triggers local host cell proliferation to produce tumors, which

will require the high transcript abundance of effector genes related to tumorigenesis; in contrast, *S.*

reilianum sustaining stable expression levels of effectors likely reflects maintenance of the biotrophic

interaction in already colonized tissue and continuation of systemic proliferation through the leaf veins

with inducing structural changes in the host tissue

Not only effector genes, but also pathogenesis related transcription factors were found to be

differentially regulated, which in turn is probably playing a crucial role in the control of effector DEO

expression. TFs Rbf1 and Biz1 were highly expressed in *S. reilianum*, which may be responsible for

the expression of *S. reilianum* enriched effector DEOs. However, the abundance of TF Hdp2

transcripts, which is activated by Rbf1 (Heimel *et al.*, 2010b), was more highly expressed in *U. maydis*,

indicating the complicated regulation of TF cascade. The promoter conversion of $Pro^{Sr10059}$ showed

comparable expression with Pro^{UMAG_05306} in *U. maydis* (Supplementary Figure S7), indicating the

potential trans effects such as transcription factors or distal enhancers might be driving differential

regulation while for orthogroup UMAG_11060/sr14941, cis-regulatory elements in promoter region

likely determines the expression pattern. Until now, only few direct target genes of TFs were

identified by CHIP-seq in smut fungi (Tollot *et al.*, 2016). Elucidation of the binding motif of these

pathogenesis related TFs and combined with the promoter analysis of effector DEOs, which were

obtained in this study could be a starting point for a more detailed study on the specific roles of TFs in

the orchestration of effector expression with respect to the adaptation to different infection styles in

pathogen speciation.

A major conclusion which can be drawn from the comparative transcriptome analysis is that the

expression levels of effector orthologs between *U. maydis* and *S. reilianum* are related to their function:

effectors with general virulence functions showed similar expression levels between both smut fungi,

whereas effectors adapted to execute specific processes such as the leaf-tumor specific effectors of *U.*

maydis showed significant differences in their transcriptional regulation. UMAG_05306 and

UMAG_11060, whose disruption resulted in reduced virulence in *U. maydis* and were functionally

conserved with respect to the corresponding *S. reilianum* orthologs. This implies that the functional

diversification of effectors can be driven on the transcriptional level and is not necessarily linked to a

changed protein function. However, UMAG_05318 provides another example for combination of

transcriptional plasticity and functional diversification of the protein level. While the orthogroups

UMAG_05318/Sr10075 or Sr10079 and Umtin2/Srtin2 were expressed during leaf infection both in *U.*

maydis and *S. reilianum*, they also show similar induction pattern during biotrophic interaction.

However, when comparing the expression levels of the orthologs, UMAG_05318 is more than 50

times higher than its *S. reilianum* ortholog, while Srtin2 has a 10 times higher expression compared to

its respective ortholog in *U. maydis*. Until now, the molecular mechanisms of these differentially

regulated organ specific effectors are not elucidated in *U. maydis*, and whether they also contribute to

virulence in *S. reilianum* is largely unknown. Deletion of Sr10075 alone did not seem affect virulence,

while a knock-out Sr10079 reduced *S. reilianum* virulence (Ghareeb *et al.*, 2019). The orthologs of

UMAG_05318 were found in *S. scitamineum* and *S. reilianum f. sp. reilianum*, but not in *U. hordei*

and *M. pennsylvanicum*. Elucidation of the function of this ortholog group and its role in smut fungi

evolution and speciation will be our next aim. Interestingly, both effectors showing functional

divergence are located in effector cluster 19A, which is the largest virulence cluster in both pathogens

(Brefort *et al.*, 2014; Ghareeb *et al.*, 2019) and several *U. maydis* highly expressed effector DEOs

were also identified in this cluster from our study. However, the impact of 19A deletion on *S.*

reilianum virulence was monitored by the head smut phenotype in the maize inflorescence (Ghareeb *et*

al., 2019). Thus, it could be possible that these effectors were highly expressed in a different time/spatial pattern to contribute to the systemic spreading of *S. reilianum*. On the other hand, it would be tempting to study if higher expression of cluster 19A effectors could promote the adaptation of *S. reilianum* to leaf infection and eventually even trigger the formation of leaf tumor by this pathogen. Therefore, to elucidate the molecular functions of these DEOs and the identification of their host targets will be instrumental to gain new insights of how evolution of effector orthologs promote the emergence of new pathogens.

With only a limited set of virulence DEOs can be tested in gene conversion experiments, this study identified three effector DEOs as promising candidates for effectors with virulence function in speciation. Despite these findings, one should not neglect the importance of effectors that evolved new functions without fundamental changes in expression patterns. For this, the Pit2 effector is a potential example: SrPit2 showed a similar expression level compared to UmPit2. However, in a previous study we found that SrPit2 only partially complemented a *U. maydis* pit2 knock out mutant, which was linked with a reduced ability of SrPit2 to inhibit SA-associated PLCPs (Misas Villamil *et al.*, 2019). Furthermore, a recent apoplastic proteome analysis suggested that composition of PLCPs is different between maize leaf and root (Schulze Hüynck *et al.*, 2019). Given that in natural conditions, *S. reilianum* infects maize through the root while *U. maydis* infects aerial tissues, one could speculate that Srpit2 adapted to be a more efficient inhibit of root PLCPs, while UmPit2 adapted to inhibit PLCPs in the maize leaf. This reflects that effector function can specialize to different hosts of tissues. In addition to this mechanism of gradual effector specialization, our approach to compare the transcriptional levels of orthologs across two different pathogen species documented that both the transcriptional plasticity and functional divergency of effector proteins contribute to pathogen speciation. Including more phylogenetically related smut fungi infecting different hosts, such as *S. reilianum f. sp. reilianum* for sorghum *and U. hordei* for barley in future research will provide a more comprehensive map of the emergence and adaptation of pathogenicity effectors during host adaptation and specialization. Besides the plastic regulation of effector orthogroups, one should not ignore the role of species-specific effectors in virulence, which is not discussed in our study. Studies on how pathogens organize their effector warehouse in temporal and spatial manner to shape the pathogenesis, and how these effectors were adapted during speciation will shed a light on the evolution and mechanistic basis of plant-pathogen interactions.

Acknowledgments

This work was funded by the European Research Council under the European Union's Horizon 2020 research and innovation program (consolidator grant conVIRgens, ID 771035), as well as funding by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany´s Excellence Strategy- EXC-2048/1- Project ID: 390686111. Weiliang Zuo and Jasper Depotter are supported by the Research Fellowship Programme for Postdoctoral Researchers of the Alexander von

558 Humboldt Foundation. MT and DKG received support by the LOEWE initiative of the government of
559 Hesse, in the framework of the LOEWE Centre for Translational Biodiversity Genomics (TBG) and
560 the LOEWE Cluster for Integrative Fungal Research (IPF).
561

562 **Author contributions**: W.Z. and G.D. designed the research; W.Z. performed the data analysis,
563 molecular experiment and virulence assay; J.D. and D.K.G. mapped the RNA-seq data; J.D. conducted
564 phylogenetic analysis; D.K.G. and M.T. provided ortholog expression normalization; W.Z. and G.D.
565 wrote the paper with contributions from the other authors.
566

567 **Data availability**
568 RNA sequencing data has been submitted to NCBI Genbank and are available under the following
569 accession: BioProject ID PRJNA674747
570

572 **References**
573
Baetsen-Young A, Wai CM, VanBuren R, Day B. **2020**. Fusarium virguliforme transcriptional plasticity is
revealed by host colonization of maize versus soybean. *Plant Cell* **32**: 336–351.

Barrangou R, Fremaux C, Deveau H, Richards M, Boyaval P, Moineau S, Romero DA, Horvath P. **2007**.
CRISPR provides acquired resistance against viruses in prokaryotes. *Science*.

Beckerson WC, Rodríguez De La Vega RC, Hartmann FE, Duhamel M, Giraud T, Perlin MH. **2019**.
Cause and effectors: Whole-genome comparisons reveal shared but rapidly evolving effector sets among host-
specific plant-castrating fungi. *mBio*.

Begerow D, Göker M, Lutz M, Stoll M. **2004**. On the evolution of smut fungi on their hosts. *Frontiers in*
basidiomycote mycology.

Brefort T, Tanaka S, Neidig N, Doehlemann G, Vincon V, Kahmann R. **2014**. Characterization of the
Largest Effector Gene Cluster of Ustilago maydis. *PLoS Pathogens*.

Depotter JRL, Doehlemann G. **2020**. Target the core: durable plant resistance against filamentous plant
pathogens through effector recognition. *Pest Management Science* **76**: 426–431.

Depotter JRL, Zuo W, Hansen M, Zhang B, Xu M, Doehlemann G. **2020**. Effectors with Different Gears:
Divergence of Ustilago maydis Effector Genes Is Associated with Their Temporal Expression Pattern during
Plant Infection. *Journal of Fungi* **7**: 16.

Djamei A, Schipper K, Rabe F, Ghosh A, Vincon V, Kahnt J, Osorio S, Tohge T, Fernie AR, Feussner I, *et*
al. **2011**. Metabolic priming by a secreted fungal effector. *Nature*.

Doehlemann G, Van Der Linde K, Aßmann D, Schwammbach D, Hof A, Mohanty A, Jackson D,
Kahmann R. **2009a**. Pep1, a secreted effector protein of Ustilago maydis, is required for successful invasion of
plant cells. *PLoS Pathogens*.

Doehlemann G, Van Der Linde K, Aßmann D, Schwammbach D, Hof A, Mohanty A, Jackson D,
Kahmann R. **2009b**. Pep1, a secreted effector protein of Ustilago maydis, is required for successful invasion of
plant cells. *PLoS Pathogens* **5**.

Flor-Parra I, Vranes M, Kämper J, Pérez-Martín J. **2006**. Biz1, a zinc finger protein required for plant
invasion by Ustilago maydis, regulates the levels of a mitotic cyclin. *Plant Cell* **18**: 2369–2387.

Franceschetti M, Maqbool A, Jiménez-Dalmaroni MJ, Pennington HG, Kamoun S, Banfield MJ. **2017**. Effectors of Filamentous Plant Pathogens: Commonalities amid Diversity. *Microbiology and Molecular Biology Reviews*.

Ghareeb H, Drechsler F, Löfke C, Teichmann T, Schirawski J. **2015**. SUPPRESSOR OF APICAL DOMINANCE1 of Sporisorium reilianum modulates inflorescence branching architecture in maize and arabidopsis. *Plant Physiology*.

Ghareeb H, Zhao Y, Schirawski J. **2019**. Sporisorium reilianum possesses a pool of effector proteins that modulate virulence on maize. *Molecular Plant Pathology*.

Gijzen M, Ishmael C, Shrestha SD. **2014**. Epigenetic control of effectors in plant pathogens. *Frontiers in Plant Science*.

Grabherr MG., Brian J. Haas, Moran Yassour Joshua Z. Levin, Dawn A. Thompson, Ido Amit, Xian Adiconis, Lin Fan, Raktima Raychowdhury, Qiandong Zeng, Zehua Chen, Evan Mauceli, Nir Hacohen, Andreas Gnirke, Nicholas Rhind, Federica di Palma, Bruce W. N, Friedman and AR. **2013**. Trinity: reconstructing a full-length transcriptome without a genome from RNA-Seq data. *Nature Biotechnology* **29**: 644–652.

Heimel K, Scherer M, Schuler D, Kämper J. **2010a**. The Ustilago maydis Clp1 protein orchestrates pheromone and b-dependent signaling pathways to coordinate the cell cycle and pathogenic development. *Plant Cell*.

Heimel K, Scherer M, Vranes M, Wahl R, Pothiratana C, Schuler D, Vincon V, Finkernagel F, Flor-Parra I, Kämper J. **2010b**. The transcription factor rbf1 is the master regulator for b-mating type controlled pathogenic development in Ustilago maydis. *PLoS Pathogens*.

Hemetsberger C, Herrberger C, Zechmann B, Hillmer M, Doehlemann G. **2012**. The Ustilago maydis effector Pep1 suppresses plant immunity by inhibition of host peroxidase activity. *PLoS Pathogens* **8**.

Hemetsberger C, Mueller AN, Matei A, Herrberger C, Hensel G, Kumlehn J, Mishra B, Sharma R, Thines M, Hückelhoven R, *et al.* 2015. The fungal core effector Pep1 is conserved across smuts of dicots and monocots. *New Phytologist*.

Irieda H, Inoue Y, Mori M, Yamada K, Oshikawa Y, Saitoh H, Uemura A, Terauchi R, Kitakura S, Kosaka A, *et al.* 2019. Conserved fungal effector suppresses PAMP-triggered immunity by targeting plant immune kinases. *Proceedings of the National Academy of Sciences of the United States of America*.

Jones JDG, Dangl JL. **2006**. The plant immune system. *Nature* **444**: 323–329.

Kämper J, Kahmann R, Bölker M, Ma LJ, Brefort T, Saville BJ, Banuett F, Kronstad JW, Gold SE, Müller O, *et al.* 2006a. Insights from the genome of the biotrophic fungal plant pathogen Ustilago maydis. *Nature*.

Kämper J, Kahmann R, Bölker M, Ma LJ, Brefort T, Saville BJ, Banuett F, Kronstad JW, Gold SE, Müller O, *et al.* 2006b. Insights from the genome of the biotrophic fungal plant pathogen Ustilago maydis. *Nature* **444**: 97–101.

Katoh K, Standley DM. **2013**. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* **30**: 772–780.

Langmead B, Salzberg SL. **2012**. Fast gapped-read alignment with Bowtie 2. *Nature Methods* **9**: 357–359.

Lanver D, Berndt P, Tollot M, Naik V, Vranes M, Warmann T, Münch K, Rössel N, Kahmann R. **2014**. Plant Surface Cues Prime Ustilago maydis for Biotrophic Development. *PLoS Pathogens*.

Lanver D, Müller AN, Happel P, Schweizer G, Haas FB, Franitza M, Pellegrin C, Reissmann S, Altmüller J, Rensing SA, *et al.* 2018. The biotrophic development of ustilago maydis studied by RNA-seq analysis. *Plant Cell*.

Lanver D, Tollot M, Schweizer G, Lo Presti L, Reissmann S, Ma LS, Schuster M, Tanaka S, Liang L, Ludwig N, *et al.* 2017. Ustilago maydis effectors and their impact on virulence. *Nature Reviews Microbiology*.

Laurie JD, Ali S, Linning R, Mannhaupt G, Wong P, Güldener U, Münsterkötter M, Moore R, Kahmann
R, Bakkeren G, *et al.* **2012**. Genome comparison of barley and maize smut fungi reveals targeted loss of RNA
silencing components and species-specific presence of transposable elements. *Plant Cell*.

Liao Y, Smyth GK, Shi W. **2019**. The R package Rsubread is easier, faster, cheaper and better for alignment
and quantification of RNA sequencing reads. *Nucleic Acids Research* **47**.

Ma LS, Wang L, Trippel C, Mendoza-Mendoza A, Ullmann S, Moretti M, Carsten A, Kahnt J, Reissmann
S, Zechmann B, *et al.* **2018**. The Ustilago maydis repetitive effector Rsp3 blocks the antifungal activity of
mannose-binding maize proteins. *Nature Communications*.

Martinez C, Roux C, Jauneau A, Dargent R. **2002**. The biological cycle of Sporisorium reilianum f.sp. zeae:
An overview using microscopy. *Mycologia* **94**: 505–514.

Matei A, Ernst C, Günl M, Thiele B, Altmüller J, Walbot V, Usadel B, Doehlemann G. **2018**. How to make
a tumour: cell type specific dissection of Ustilago maydis-induced tumour development in maize leaves. *New*
Phytologist.

Misas Villamil JC, Mueller AN, Demir F, Meyer U, Ökmen B, Schulze Hüynck J, Breuer M, Dauben H,
Win J, Huesgen PF, *et al.* **2019**. A fungal substrate mimicking molecule suppresses plant immunity via an inter-
kingdom conserved motif. *Nature Communications* **10**.

Morita T, Koike H, Koyama Y, Hagiwara H, Ito E, Fukuoka T, Imura T, Machida M, Kitamoto D. **2013**.
Genome sequence of the basidiomycetous yeast Pseudozyma antarctica T-34, a producer of the glycolipid
biosurfactants mannosylerythritol lipids. *Genome Announcements*.

Mueller AN, Ziemann S, Treitschke S, Aßmann D, Doehlemann G. **2013**. Compatibility in the Ustilago
maydis-Maize Interaction Requires Inhibition of Host Cysteine Proteases by the Fungal Effector Pit2. *PLoS*
Pathogens **9**.

Raffaele S, Kamoun S. **2012**. Genome evolution in filamentous plant pathogens: Why bigger can be better.
Nature Reviews Microbiology **10**: 417–430.

Redkar A, Hoser R, Schilling L, Zechmann B, Krzymowska M, Walbot V, Doehlemann G. **2015a**. A
secreted effector protein of Ustilago maydis guides maize leaf cells to form tumors. *Plant Cell*.

Redkar A, Villajuana-Bonequi M, Doehlemann G. **2015b**. Conservation of the Ustilago maydis effector see1
in related smuts. *Plant Signaling and Behavior*.

Robinson MD, McCarthy DJ, Smyth GK. **2009**. edgeR: A Bioconductor package for differential expression
analysis of digital gene expression data. *Bioinformatics*.

Sánchez-Vallet A, Fouché S, Fudal I, Hartmann FE, Soyer JL, Tellier A, Croll D. **2018**. The genome
biology of effector gene evolution in filamentous plant pathogens. *Annual Review of Phytopathology* **56**: 21–40.

Schilling L, Matei A, Redkar A, Walbot V, Doehlemann G. **2014**. Virulence of the maize smut Ustilago
maydis is shaped by organ-specific effectors. *Molecular Plant Pathology*.

Schirawski J, Mannhaupt G, Münch K, Brefort T, Schipper K, Doehlemann G, Di Stasio M, Rössel N,
Mendoza-Mendoza A, Pester D, *et al.* **2010**. Pathogenicity determinants in smut fungi revealed by genome
comparison. *Science*.

Schnable PS, Ware D, Fulton RS, Stein JC, Wei F, Pasternak S, Liang C, Zhang J, Fulton L, Graves TA,
et al. **2009**. The B73 maize genome: Complexity, diversity, and dynamics. *Science* **326**.

Schulze Hüynck J, Kaschani F, van der Linde K, Ziemann S, Müller AN, Colby T, Kaiser M, Misas
Villamil JC, Doehlemann G. **2019**. Proteases underground: Analysis of the maize root apoplast identifies organ
specific papain-like cysteine protease activity. *Frontiers in Plant Science*.

Schuster M, Kahmann R. **2019**. CRISPR-Cas9 genome editing approaches in filamentous fungi and oomycetes.
Fungal Genetics and Biology.

Schuster M, Schweizer G, Kahmann R. **2018**. Comparative analyses of secreted proteins in plant pathogenic

smut fungi and related basidiomycetes. *Fungal Genetics and Biology*.

Schuster M, Schweizer G, Reissmann S, Kahmann R. **2016**. Genome editing in Ustilago maydis using the
CRISPR-Cas system. *Fungal Genetics and Biology*.

Schweizer G, Münch K, Mannhaupt G, Schirawski J, Kahmann R, Dutheil JY. **2018**. Positively Selected
Effector Genes and Their Contribution to Virulence in the Smut Fungus Sporisorium reilianum. *Genome Biology*
and Evolution.

Selin C, de Kievit TR, Belmonte MF, Fernando WGD. **2016**. Elucidating the role of effectors in plant-fungal
interactions: Progress and challenges. *Frontiers in Microbiology* **7**: 1–21.

Seppey M, Manni M, Zdobnov EM. **2019**. BUSCO: Assessing genome assembly and annotation completeness.
In: Methods in Molecular Biology.

Sharma R, Mishra B, Runge F, Thines M. **2014**. Gene Loss rather than gene gain is associated with a host
jump from monocots to dicots in the smut fungus melanopsichium pennsylvanicum. *Genome Biology and*
Evolution.

Skibbe DS, Doehlemann G, Fernandes J, Walbot V. **2010**. Maize tumors caused by ustilago maydis require
organ-specific genes in host and pathogen. *Science*.

Stamatakis A. **2014**. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies.
Bioinformatics **30**: 1312–1313.

Stirnberg A, Djamei A. **2016**. Characterization of ApB73, a virulence factor important for colonization of Zea
mays by the smut Ustilago maydis. *Molecular Plant Pathology*.

Sucher J, Mbengue M, Dresen A, Barascud M, Didelon M, Barbacci A, Raffaele S. **2020**.
Phylotranscriptomics of the pentapetalae reveals frequent regulatory variation in plant local responses to the
fungal pathogen sclerotinia sclerotiorum. *Plant Cell* **32**: 1820–1844.

Tanaka S, Brefort T, Neidig N, Djamei A, Kahnt J, Vermerris W, Koenig S, Feussner K, Feussner I,
Kahmann R. **2014**. A secreted Ustilago maydis effector promotes virulence by targeting anthocyanin
biosynthesis in maize. *eLife*.

Tanaka S, Schweizer G, Rössel N, Fukada F, Thines M, Kahmann R. **2019**. Neofunctionalization of the
secreted Tin2 effector in the fungal pathogen Ustilago maydis. *Nature Microbiology*.

Taniguti LM, Schaker PDC, Benevenuto J, Peters LP, Carvalho G, Palhares A, Quecine MC, Nunes FRS,
Kmit MCP, Wai A, *et al*. 2015. Complete genome sequence of Sporisorium scitamineum and biotrophic
interaction transcriptome with sugarcane. *PLoS ONE*.

Thines M. **2019**. An evolutionary framework for host shifts – jumping ships for survival. *New Phytologist* **224**:
605–617.

Tollot M, Assmann D, Becker C, Altmüller J, Dutheil JY, Wegner CE, Kahmann R. **2016**. The WOPR
Protein Ros1 Is a Master Regulator of Sporogenesis and Late Effector Gene Expression in the Maize Pathogen
Ustilago maydis. *PLoS Pathogens* **12**.

Torres DE, Oggenfuss U, Croll D, Seidl MF. **2020**. Genome evolution in fungal plant pathogens: looking
beyond the two-speed genome model. *Fungal Biology Reviews*.

Toruño TY, Stergiopoulos I, Coaker G. **2016**. Plant-Pathogen Effectors: Cellular Probes Interfering with Plant
Defenses in Spatial and Temporal Manners. *Annual Review of Phytopathology* **54**: 419–441.

Zahiri A, Heimel K, Wahl R, Rath M, Kämper J. **2010**. The Ustilago maydis forkhead transcription factor
Fox1 is involved in the regulation of genes required for the attenuation of plant defenses during pathogenic
development. *Molecular Plant-Microbe Interactions*.

Zuo W, Depotter JR, Doehlemann G. **2020**. Cas9HF1 enhanced specificity in Ustilago maydis. *Fungal*
Biology **124**: 228–234.

	Figure Legends

	Fig.1. Comparison of *U. maydis* and *S. reilianum* growth during leaf infection. a,
	calcofluor white staining of fungal hyphae and appressorium at 20hpi. **b-d**, WGA-AF488-
	Propidium Iodide co-staining to show the fungal biotrophic growth inside the plant cell.
	Fungal cells were stained with WGA-AF488 (green), the vascular cell wall was stained with
	Propidium Iodide (red). White arrows indicate the *de-novo* divided bundle sheath cells
	triggered by *U. maydis* infection. White circles indicate the aggregation of *S. reilianum* cells
	inside the leaf vein. **e**, the relative biomass quantification by qPCR shows the fungal growth
	during infection, and a significant higher abundance of *S. reilianum* was detected at 4 dpi. **f**,
	the relative fungal reads from RNA-seq, which was the ratio between reads uniquely mapped
	to fungi and total reads uniquely mapped (fungi and maize). A significantly higher number of
	mapped reads was detected from *S. reilianum* at 4 dpi. "**", Experiments have been
	performed in three independent biological replicates. *p<0.05*. *Student t-test* was used for
	significance test.

	Fig.2. Overview of *U. maydis* and *S. reilianum* transcriptome, respectively. a, PCA
	analysis of *U. maydis* and *S. reilianum* RNA-seq samples. **b,** expression data from three
	different growth conditions. The number of differentially expressed genes (log2 fold
	change>1, p>0.05) from pairwise comparison were indicated in the triangles. Red and green
	triangles represent up-regulated and down-regulated genes between 2 dpi vs. axenic culture
	(AC), 4 dpi vs. AC and 4 dpi vs. 2 dpi, respectively. **c,** boxplot shows transcriptional changes
	of effector and non-effector genes during biotrophic growth. Effector genes were dramatically
	up-regulated compared to the non-effector genes. **d,** heatmap shows the overview of the
	expression changes of effector genes. "n" shows the number of effectors detected expressing
	in RNA-seq sample and the total number of effectors in the genome.

	Fig.3. The induced pattern shows the expression changes of effectors. a, the induced
	effectors are clustered into 5 patterns based on the expression changes between axenic culture
	(AC), 2 dpi and 4 dpi conditions. The gray lines represent individual gene, and blue and
	orange lines are the mean of all genes in each cluster. "a", "b" and "c" were used to indicate
	the significance levels between each condition. Genes with log2 fold change>1 and p<0.05
	were considered significantly induced in pairwise comparison. **b,** the number of induced
	genes of each pattern. The *chi-square test* was used for statistic test for the significance. "**",
	p<0.01.

	Fig.4. Orthologs were differentially regulated between *U. maydis* and *S. reilianum*. a,
	PCA analysis based on the relative ortholog expression of 6005 one-to-one ortholog pairs
	from *U. maydis* and *S. reilianum*. **b,** the number of DEOs detected in different conditions. The
	fold change>3 and p<0.05 were used to identify DEO. **c,** the GO classification of DEOs in
	three different conditions. **d,** the Venn diagram shows the number of differentially expressed
	effector orthologs during biotrophic growth. **e-f,** the differentially regulation of effectors
	clustered by the conservation between 5 smut fungi. **e,** the phylogenetic tree of 5 smut fungi,
	including *U. maydis*, *S. reilianum*, *Ustilago hordei*, *S. scitamineum* and *M. pennsylvanicum*. **f,**
	the number of effector DEOs in each category. *Chi-square test* was used for significance test.
	"**", *p<0.01*, "***", *p<0.001*.

	Fig.5. Effector orthologs between *U. maydis* and *S. reilianum* are functional conserved. a,
	a heatmap shows the expression difference and protein sequence identity of effectors ortholog
	pairs which have been knocked out and subjected to virulence tests in *U. maydis*. "¶", Pit2
	only conserved in functional PID motif which failed to be detected during ortholog

identification and "%ID" was not showed. **b,** the scheme demonstrated the process of
CRISPR-Cas9 mediated gene conversion. The sgRNA was designed to target the coding
sequence of *U. maydis* effector, and the donor plasmid contains *S. reilianum* ortholog coding
sequence flanked by 1kb homologous region. **c-d,** Disease scoring results of gene conversion
mutants and corresponding *U. maydis* knock out mutant. **c,** Converted UMAG_05306 and
UMAG_11060 to the *S. reilianum* orthologs did not affect the virulence in SG200. **d,**
Converted UMAG_05318 to its *S. reilianum* ortholog Sr10075 or Sr10079, the paralog of
Sr10075, compromise SG200 virulence, and show reduced virulence as deletion mutant.
Experiments have been performed in three independent biological replicates. n indicates the
total number of plants tested in three replicates. The significance as calculated by *student t-*
test with disease index "*", $p<0.05$, "**", $p<0.01$.
798
799
800 **Supplementary Captions**
801
Supplementary Fig.1 The symptom of *U. maydis* and *S. reilianum* infected leaves. a, The
photos shows the symptom development from 2 dpi to 4 dpi. The dashed circle shows the first
visible tumor from *U. maydis* infected leaf while the dashed rectangle shows the necrosis spot
caused by *S. reilianum* infection. The back *of U. maydis* infected leaf from 4 dpi is shown to
better visualize the tumor. **b,** Confocal microscopy shows the details of the *S. reilianum*
aggregation inside the leaf vascular at 4 dpi after WGA-AF488- Propidium Iodide staining.
The fungal cells aggregate inside the vascular tissue. Scale bar=100 µm.
809
Supplementary Fig.2 qPCR confirms the expression of effector genes from *U. maydis*
and *S. reilianum*. a, Expression levels of *U. maydis* effector genes. **b,** Expression levels of *S.*
reilianum effector genes. Infection markers such as *UmPep1* and *UmPit2* and their
corresponding orthologs were detected, as well as organ-specific effector See1 and two
differentially expressed orthogroups *UMAG_05306/Sr10059* and *UMAG_11060/Sr14941*.
The relative expression levels of effectors were normalized by respective reference
peptidylprolyl isomerase (*ppi*) genes by $2^{-\Delta Ct}$ method. Error bar indicate standard deviation
from three biological replications.
818
Supplementary Fig.3 Differential regulation of orthologs between *U. maydis* and *S.*
***reilianum* from different samples. a,** Pearson correlation analysis between *U. maydis* and *S.*
reilianum samples based on relative ortholog expression from 6005 one-to-one orthologs. The
r values indicate a high to low correlation between *U. maydis* and *S. reilianum* samples:
AC>2 dpi>4 dpi. **b,** Venn diagram shows the number of differentially expressed ortholog in *U.*
maydis and *S. reilianum* from different conditions. About 30% and 50% of detected DEOs
were only detected from axenic culture.
826
Supplementary Fig.4 Expression of pathogenic development related transcription
factors between *U. maydis* and *S. reilianum* during biotrophic growth from RNA-seq
data. The transcription factors *Rbf1* and *Biz1* were highly expressed in *S. reilianum* while
transcription factors *Hdp2*, *Fox1* and *Nlt1* were enhanced expressed in *U. maydis*. The
transcription factor *Ros1* shows similar expression between these two smut fungi. *Student t-*
test was used for significance test. "**", $p <0.01$.
833
Supplementary Fig.5 The distribution of effector DEOs on chromosomes. **a,** the
distribution of DEOs in 22 chromosomes based on the *U. maydis* genome organization. The
blue bar shows *U. maydis* highly expressed DEOs, and the yellow bar shows the *S. reilianum*
highly expressed DEOs. 2 dpi DEOs and 4 dpi DEOs were plotted above and below the

chromosome, respectively. The red rectangle shows effector cluster 19A. **b,** the diagram
shows the overview of synteny of effectors in cluster 19A. The effector genes are showed in
green arrows; non-effector genes are shown in dark gray. The synteny of effector genes are in
grey shade, the outline color of the shade indicates the DEO. The *U. maydis* highly expressed
effector ortholog pairs are outlined in blue, the *S. reilianum* highly expressed genes are
outlined in yellow.

Supplementary Fig.6 Sequence comparison of UMAG_05318, Sr10075 and Sr10079. a,
the maximum likelihood phylogenetic tree of UMAG_05318, Sr10075 and Sr10079 based on
the amino acid sequences. Sr10079 is distant from UMAG_05318 and Sr10075. **b,** the amino
acid sequence alignment of UMAG_05318, Sr10075 and Sr10079. The identical amino acids
were shaded in grey.

Supplementary Fig. 7 The regulation of *S. reilianum* promoters in *U. maydis* by
CRIPSR-Cas9 mediated promoter conversion. a, Schematic model of the method used for
CRISPR-Cas9 mediated promoter conversion. **b,** qPCR detection of the corresponding
effectors expression under different orthologous *S. reilianum* promoters. At 4 dpi, $\text{Pro}^{\text{sr10059}}$
regulates gene expression to the similar level as $\text{Pro}^{\text{UMAG\_05306}}$ in *U. maydis*, while at 2 dpi,
$\text{Pro}^{\text{sr14941}}$ drove gene expression was significantly lower compared to $\text{Pro}^{\text{UMAG\_11060}}$. Error bar
indicates standard deviation from three biological replications.

Supplement Table.1 Oligo sequences used.
Supplement Table.2 Log2 CPM (count per million) of fungal reads.
Supplement Table.3 Relative ortholog expression data of 6005 one-to-one orthologs
between *U. maydis* and S.*reilianum*.

Figure 1

Figure 2

Figure 3

(a)

(b)

Figure 4

Figure 5