Full title: Conjunctive spatial and idiothetic codes are topographically organized in the

GABAergic cells of the lateral septum

Short title: Spatial and self-motion coding in the lateral septum

Authors: Suzanne van der Veldt[1], Guillaume Etter[1], Coralie-Anne Mosser[1], Frédéric

Manseau[1], Sylvain Williams[1*]

[1] McGill University & Douglas Mental Health University Institute, Montreal, Canada

6875, Boul. LaSalle, Verdun (Quebec), H4H 1R3, Phone: 514 761-6131, Fax: 514 888-4067

*Corresponding author: sylvain.williams@mcgill.ca

Author contributions

SV and SW designed the study. SV performed surgeries, behavioral recording experiments,

immunohistological experiments, and calcium data analysis. GE wrote code for MI

calculation and Bayesian decoding and implanted mice for CA1 recordings. CAM implanted

and recorded four mice that were used in the current study. FM performed and analyzed

patch-clamp in vitro electrophysiology. SV and SW wrote the manuscript with inputs from all

authors.

Abstract

The hippocampal spatial code's relevance for downstream neuronal populations – particularly its major subcortical output the lateral septum (LS) - is still poorly understood. Here, using calcium imaging combined with unbiased analytical methods, we functionally characterized and compared the spatial tuning of LS GABAergic cells to those of dorsal CA3 and CA1 cells. We identified a significant number of LS cells that are modulated by place, speed, acceleration, and direction, as well as conjunctions of these properties, directly comparable to hippocampal CA1 and CA3 spatially modulated cells. Interestingly, Bayesian decoding of position based on LS spatial cells reflected the animal's location as accurately as decoding using the activity of hippocampal pyramidal cells. A portion of LS cells showed stable spatial codes over the course of multiple days, potentially reflecting long-term episodic memory. The distributions of cells exhibiting these properties formed gradients along the anterior-posterior and dorsal-ventral axes of the LS, directly reflecting the topographical organization of hippocampal inputs to the LS. Finally, we show using trans-synaptic tracing that LS neurons receiving CA3 and CA1 excitatory input send projections to the hypothalamus and medial septum, regions that are not targeted directly by principal cells of the dorsal hippocampus. Together, our findings demonstrate that the LS accurately and robustly represents spatial, directional as well as self-motion information and is uniquely positioned to relay this information from the hippocampus to its downstream regions, thus occupying a key position within a distributed spatial memory network.

Abbreviations

LH   lateral hypothalamus

LS   lateral septum

51 MS   medial septum

52 VTA   ventral tegmental area

Introduction

The lateral septum (LS) is an anatomically complex structure uniquely positioned

within a broader distributed spatial memory network. The LS consists almost exclusively of

highly interconnected inhibitory GABAergic cells [1,2] that are densely innervated by the

principal cells of the hippocampus [3,4], in contrast to the medial septum (MS), which

receives inputs solely from the inhibitory neurons of the hippocampus [5]. Previous research

has implicated the LS in a wide variety of behaviors, including spatial and working memory

[6–10], regulation of feeding [11–14], anxiety [15–18], locomotion [19,20], and social

behaviors [21–24].

To date, functional characterizations of the LS have often focused on the spatial

coding properties of these neurons, primarily due to the extensive excitatory input from

positionally-tuned hippocampal pyramidal neurons, commonly referred to as 'place cells'

[4,24–26]. Yet, this body of work has not yet yielded a consensus. Several studies have

reported place-like cells in the LS, but the number, information content, and stability vary

widely from study to study [20,27–31], while others described a lack of canonical place cells

altogether [4]. As previous studies often depended on variable criteria of what defines a place

cell, estimates range from 5.3% [4] to 56.0% [20] of LS cells classified as spatially-

modulated. Strikingly, even estimates by the same authors on data acquired from the same

subjects engaging the same task ranged between 26.5% [32] to 56.0% [20]. When reported,

LS place cells are typically described as of lesser quality than classic hippocampal place

cells, with lower within-session stability and larger place fields [30,31]. Adding to this

complexity, the LS is a large structure that is cytoarchitecturally subdivided into dorsal,

intermediate, and ventral subregions that spread across its anterior-posterior axis [5,26,33].

This suggests that, in addition to subjective criterias, the striking differences in LS spatial

coding characteristics between studies may be in part a product of variations in recording

locations within the region. In addition to positional information, LS neurons may also

encode direction and self-motion information, including acceleration and velocity [20,30,31].

Beyond their spatial tuning properties, hippocampal place cells are also noted for the

rapid reorganization of the place fields across time (on the order of days) and navigational

contexts, a phenomenon termed remapping [34–38]. If the LS spatial code is directly

inherited from its hippocampal inputs, one would expect comparable dynamics over a similar

timescale. While the LS, similar to the hippocampus, has been implicated in behaviors that

may require a stable representation of space over time, including context-reward [39,40],

contextual fear conditioning [16,41–44], spatial learning and memory [32,45,46] remapping

dynamics in the LS remain largely understudied. Thus, whether and how the LS spatial code

changes across time is crucial to clarifying its relationship to the upstream hippocampal

spatial code, with important implications for its involvement in this broader spatial memory

network and how it supports behaviors across longer timescales.

To resolve these outstanding questions, we functionally characterize hippocampal

connectivity of LS, and asked how information rich LS GABAergic neurons are in relation to

their main pyramidal inputs in dorsal CA3 and CA1. To this end, we used head-mounted

miniaturized microscopes to record ~2000 LS neurons across the anterior-posterior and

dorsal-ventral axes of the LS, and compared their firing characteristics to pyramidal cells

recorded in CA1 and CA3, as animals navigated linear track and open field environments.

Results demonstrate that the LS accurately and robustly represents spatial, speed, and

directional information in an anatomically-organized fashion. While the information of LS

neurons is generally comparable to CA3 and CA1, our tracing studies suggest that LS

information is sent to downstream brain regions such as the MS and hypothalamus that are

not directly connected by the principal cells of the dorsal hippocampus.

Results

LS GABAergic neurons exhibit stable spatial activity during alternation on a linear

track

Previous electrophysiological studies of the LS disagree on both the proportion of place

encoding cells in this region and their spatial content. Cells have been previously described to

be of lower quality than the classically described hippocampal place cells [20,30,32] or to be

virtually absent [4].

First, to assess the distribution of excitatory pyramidal inputs from each hippocampal

subregion to the GABAergic cells of the LS, we injected a monosynaptic retrograde rabies

tracer in dorsal LS (Fig 1A, S1 Fig, n = 5 mice), utilizing the monosynaptically restricted,

Cre-dependent retrograde tracing features of this technique. We observed strong labelling of

pyramidal neurons of the hippocampus in areas CA1 and CA3 (Fig 1B, S1 Fig). To

investigate how the spatial coding properties of the LS compare to these main dorsal

hippocampal inputs, we expressed the calcium (Ca^{2+}) indicator GCaMP6f in GABAergic

cells of the LS or in the pyramidal cells of the dorsal CA1 (dCA1) and dorsal CA3 (dCA3) of

mice (Fig 1C, S2 Fig, S3 Fig). We recorded Ca^{2+} activity from 1082 LS GABAergic neurons

(n = 15 mice), 1251 dCA1 pyramidal cells (n = 5 mice) and 464 cells from dCA3 (n = 6

mice) from animals alternating on a linear track. We subsequently filtered and binarized the

recorded calcium traces (S4A Fig) [47] and excluded periods of immobility ($< 5cm.s^{-1}$).

Using an information-theoretic framework [48–50], we computed the mutual information

(MI) between calcium activity and spatial location for each cell and expressed the results in

bits (S4B Fig). As we observed a significant correlation between probability of a cell to be

active (P(A)) and MI (S4B Fig), we computed the MI value for 30 bootstrap samples (50%

data sampling) and compared those to 30 circularly shuffled surrogates, which allowed us to

compute significance of the MI value independently of P(A) (S4C-E Fig) [47]. We found that

LS GABAergic activity was significantly spatially modulated ($p < .01$, Fig 1D, S4H Fig).

Indeed, both mutual information and within-session stability was higher for spatially

modulated LS cells compared to non-modulated cells (Mann-Whitney test, $p < .0001$, S4E-F

Fig). Using this approach, a total of 43.90% of LS GABAergic cells qualified as spatially

modulated (475/1082 cells), with LS cells similarly distributed along the linear track as place

cells recorded from dorsal CA1 and dorsal CA3 (Fig 1E). In contrast to previous reports

[4,20,32], we observed a significant difference in the proportion of spatially modulated cells

across recording regions (LS: 43.57 ± 4.46 %; dCA1: 29.60 ± 2.99 %; dCA3: 27.01 ± 7.07

%; Kruskal-Wallis, H(3) = 6.212, $p = .0448$; Dunn's multiple comparisons test: LS-dCA1: p

= 0.1757; LS-dCA3: $p = 0.1101$; Fig 1F). We found that within-session stability of spatially

modulated cells (expressed as a correlation where 0 means no stability while 1 means

perfectly stable) was significantly different between LS (0.44 ± .0165) and dCA3 (0.73 ±

0.26; Kruskal-Wallis, H(3) = 80.48, $p < .0001$; Dunn's multiple comparisons test, $p < .0001$),

and between dCA1 and dCA3 ($p < .0001$; Fig 1G). Even though LS GABAergic cells are not

fast spiking (S5 Fig, S6 Fig) [51], P(A) in LS ($0.02005 ± 4.692 \times 10^4$) is higher than both

dCA1 ($0.0094 ± 2.544 \times 10^4$, Kruskal-Wallis, H(3) = 502.0, $p < .0001$; Dunn's multiple

comparisons test, $p > .0001$) and dCA3 ($0.009225 ± 3.759 \times 10^4$, $p < .0001$, S6A,B).

Therefore, we assessed the mutual information in bits carried in each binarized event, and

observed that the LS ($5.624 \times 10^5 ± 1.151 \times 10^6$ bits/binarized event) carries significantly less

information per binarized event than CA1 ($1.430 \times 10^4 ± 1.613 \times 10^6$ bits/binarized event;

Kruskal-Wallis, H(3) = 1288, $p < .0001$; Dunn's multiple comparisons test, $p < .0001$); and

CA3 ($1.130 \times 10^4 \pm 2.507 \times 10^6$; bits/binarized event; $p < .0001$, Fig 1H). To confirm that

the observed differences in proportion of spatially modulated cells and within-session

stability were not due to differences in P(A), we used an activity cut-off of P(A) > .001 and

confirmed our previous results (S6D Fig).

In contrast to their omnidirectional firing properties in open field environments,

hippocampal neurons rapidly develop strong directional selectivity on linear tracks [52–54].

To test whether the LS place code developed comparable directional selectivity, we ranked

spatially modulated cells as a function of the difference between MI for left and right

traversals (S7A-D Fig), and found LS cells tuned to the absolute location regardless of

direction (7.67%, 83/1082 cells), while other cells were selective to the left- (15.52%,

168/1082 cells), or right-trajectories (22.08%, 239/1082 cells).

Fig 1: LS shares spatial coding characteristics with its main hippocampal inputs CA1 and CA3

during goal directed navigation in a 1D environment.

(A) Retrograde rabies tracing injection site, blue = DAPI staining, green = TVA.oG coupled to an

enhanced green fluorescent protein (eGFP), red = Rabies coupled to mCherry, orange: starter cells

expressing both TVA.oG.eGFP and Rabies.mCherry. (B) Retrograde labelling in the dorsal

hippocampus, coronal section. Bottom left: example of mCherry-positive CA1 pyramidal cell, bottom

right: mCherry-positive CA3 pyramidal cells, blue = DAPI staining, red = Rabies coupled to

mCherry. (For additional images see S1 Fig). (C) Top: Diagram of one-photon calcium recording

setup in LS, CA1 and CA3 in freely behaving mice, with GCaMP6f expression restricted to

GABAergic cells in LS, and restricted to pyramidal cells in dorsal CA1 and CA3, middle: histological

verification of implantation site (see also S2 Fig and S3 Fig) and bottom: extracted calcium transients

for LS, CA1 and CA3. (D) Top: Linear track paradigm, with sucrose rewards on either end. Bottom:

Probability of an example cell to be active given the location on the linear track (red) with 95% upper

and lower percentile (grey). Examples shown are from two different LS mice, and a representative

spatially modulated cell from one CA1 and CA3 animal each. (E) Activity of cells sorted along

location in the maze for each region (blue, low; yellow, high). (F) Percent of significantly spatially

modulated cells for each animal for each recording region. LS: N = 15 mice. CA1: N = 5 mice, N = 6

mice. (G) Within-session stability of spatially modulated cells in each region active (LS: n = 475

cells, CA1: n = 336 cells, CA3: n = 143 cells). (H) MI per binarized event for all cells (LS: n = 1030

cells, n = 15 mice. CA1: n = 1251 cells, n = 5 mice, CA3: n = 464 cells, n = 6 mice). *, $p < .05$, **, p

$< .01$, ****, $p < .0001$. Test used in F-H: Kruskal-Wallis with Dunn's multiple comparisons test.

Previous studies have used decoding methods to predict behavioral variables from

calcium imaging data recorded in the hippocampus [34,38,55], yielding insights into the

amount of information encoded by a neuronal assembly. Here, we asked whether we could

reliably estimate the mouse location solely from LS neuronal activity patterns (Fig 2). Using

30 bootstrap samples of 60 cells (Fig 2B,C), decoding location using LS neuronal activity

significantly outperformed a decoder that was trained using shuffled data (paired t-test, t(6) =

13.56, $p < .0001$, Fig 2D), even when decreasing the number of neurons used (S8A,B Fig).

Similarly, a decoder trained using CA1 or CA3 data also yielded significantly less error than

one using shuffled surrogates (CA1: t(4) = 7.558, $p = .0016$; CA3: t(3) = 3.233, $p = .0481$,

Fig 2D, S8C,D Fig). While using a small bootstrap sample size allowed a fair comparison

between recording regions, it also induced higher error rates. To ensure the proper

functioning of our decoder, we confirmed that the decoding error decreased significantly

when increasing bootstrapped sample size (S8C,D Fig).

In order to compare decoding accuracy for LS, CA1 and CA3, a decoding score was

computed (Fig 2E, see methods), and we found that LS significantly outperformed CA3 (one-

way ANOVA, F(2,13) = 6.277, $p = .0124$; Holm-Sidak's multiple comparisons test $p < .05$),

but not CA1. Overall, spatially modulated cells recorded in LS were significantly more active

than those recorded in the hippocampus (H(3) = 155.1, $p < .0001$, S5A,B Fig), which could

be a contributing factor to the overall higher decoding accuracy using LS cells. To confirm

that the differences observed in decoding error are not due to the inclusion of a large number

of pseudo-silent cells in the dCA1 or dCA3 bootstrapped samples, we confirmed our findings

using an activity cut-off of $P(A) > .001$ (S8E-F). Similarly, to confirm that the observed

differences in decoding error are not due to the use of a temporal smoothing window, we

replicated our results using a decoder that omitted temporal filtering (S8G-H).

Fig 2: Location can be reliably decoded from lateral septum GABAergic cells.

(A) Top: Posterior probabilities for each frame estimated from binarized calcium activity, bottom:

corresponding actual (green) and decoded (red) location. (B) Schematic representation of

bootstrapping approach. (C) Method for computing mean Euclidean distance decoding error (in cm)

for each bootstrapped estimate. (D) Decoding error for actual vs shuffled dataset (30 bootstrapped

samples of 60 cells, LS: N = 7 mice, CA1: N = 5 mice, CA3: LS = 4 mice). (E) Mean decoding score

for each region. *, $p < .05$, **, $p < .01$, ****, $p < .0001$, Test used in D, paired t-test, E, one-way

ANOVA, Holm-Sidak's multiple comparisons test.

LS GABAergic neurons exhibit stable spatial activity during free exploration in a 2D

environment

In order to assess spatial coding during free exploration in a two-dimensional environment,

we recorded Ca^{2+} activity of 1899 GABAergic neurons in 28 mice implanted in distinct

subregions of the LS, while the animals were freely exploring a novel open field. Out of 1899

recorded cells, 37.80% (718/1899 cells, n = 28 mice) of LS cells displayed significant spatial

information ($p < .01$; Fig 3A-B, S9 Fig), as compared to 31.33% (323/1031 cells, n = 6 mice)

in CA1 and 25.84% (138/534 cells, n = 8 mice) CA3 cells. Despite the LS' known

involvement in feeding-related behaviors [11–14], we did not observe any increased number

of cells with firing fields around food zones, nor did we observe overrepresentation of objects

or walls (S10 Fig). To assess reward-modulation in the context of increased task demand, we

assessed spatial firing of LS GABAergic neurons recorded while animals performed a

delayed non-matching to place task in a T-maze, but did not observe any significant reward

code compared to shuffled surrogates (S11 Fig).

With theta-rhythmic cells in the LS having higher firing rates than theta-rhythm

independent cells [56], and the suggestion that theta-rhythmic cells could receive direct

hippocampal inputs (though see [57,58]), we asked whether LS cells with higher firing rates

were also more likely to be spatially modulated. Indeed, spatially-modulated cells were

significantly more active than non-modulated cells (Mann-Whitney test, U = 342219, $p <$

.0001; S9G-H Fig) and displayed a higher bursting index, defined as the probability of a cell

being active, given it was already in an active state, or $P(A_t|A_{t-1})$, than non spatially

modulated cells (Mann-Whitney test, U = 368276, $p < .0001$; S9H Fig).

We compared the place field properties of cells recorded in LS to those of dorsal CA1

and CA3 (Fig 3). For each animal, we computed the portion of significantly positionally

modulated cells per region, which was not significantly different across LS (36.71 ± 2.16 %),

dCA1 (38.32 ± 9.36 %), and dCA3 (28.39 ± 6.73 %; one-way ANOVA, F(2,39) = 1.105, $p =$

.3414; Fig 3B). We computed the information for each binarized event, and observed that the

LS (1.020 x 10^4 ± 7.56 x 10^7 bits/binarized event) carries significantly less information per

binarized event than CA1 (1.200 x 10^4 ± 1.067 x 10^6 bits/binarized event; Kruskal-Wallis,

H(3) = 264.6, $p < .0001$; Dunn's multiple comparisons test, $p < .0001$) and CA3 (1.199 x 10^4

± 1.605 x 10^6 bits/binarized event; $p < .0001$). As expected on the basis of prior results

[4,28,30,31], within-session stability of LS cells (0.31 ± 0.011) was significantly lower than

that of spatially modulated cells recorded in CA1 (0.44 ± 0.017; Kruskal-Wallis, H(3) =

89.52, $p < .0001$; Dunn's multiple comparisons test, $p < .0001$) and CA3 (0.56 ± 0.027; $p <$

.0001; Fig 3D), and place fields were more dispersed (LS: 14.47 ± 0.13 cm; CA1: 12.00 ±

0.207 cm; CA3: 10.61 ± 0.35 cm; Kruskal-Wallis, H(3) = 155.7, $p < .0001$, Fig 3E). Similar

to recordings in a 1D environment, we observed large differences in P(A) between spatially

modulated cells recorded from LS and those from dorsal CA1 and CA3 (S12A Fig).

Therefore, we also assessed for each region the proportion of spatially modulated cells, split-

half stability and mean-dispersion using an activity cut-off of P(A) > .001, and confirmed our

results (S12B Fig).

When decoding the location of the animals using bootstrap samples of 80 cells from

each region (Fig 3F), decoding error was significantly lower in LS (20.16 ± 1.48 cm)

compared to shuffled surrogates (26.03 ± 0.84 cm; two-way RM ANOVA, F(1,13) = 89.09, p

< .0001 for main effect of shuffle), even when decreasing the number of neurons used to only

40 neurons (S13A,B). Omission of temporal filtering recapitulated these results (S13E Fig).

The same held true for decoders trained using CA1 or CA3 data respectively (S13C,D).

Strikingly, a decoder trained on data from LS does not perform significantly worse than a

decoder trained on CA1 or CA3 data (F(2,13) = 2.186, $p = 0.2384$ for main effect of region;

Fig 3H). Similarly to recordings on the linear track, we used an activity cut-off of P(A) > .001

to confirm that the differences observed in decoding error between regions are not due to the

inclusion of a large number of pseudo-silent cells in the bootstrapped sample (S13F Fig).

Fig 3: LS shares spatial coding characteristics with its main hippocampal inputs CA1 and CA3

during free exploration in a 2D environment.

(A) Example tuning maps of spatially modulated cells recorded from LS, dorsal CA1 and dorsal CA3

in a 45 x 45 cm open field (3 x 3 cm bins). (B) Proportion of spatial cells per animal (LS: n = 28 mice;

dCA1: n = 6 mice; dCA3: n = 8 mice). (C) MI (bits) per binarized event for all cells recorded from

each region (LS: n = 1899 cells from n = 28 mice; dCA1: n = 1031 cells, n = 6 mice; dCA3: n = 534

cells, n = 8 mice). (D) Within session stability for spatial cells (LS: n = 734 spatial cells from n = 28

mice; dCA1: n = 424 spatial cells, n = 6 mice; dCA3: n = 138 spatial cells, n = 7 mice). (E) Left:

mean dispersion computation, right: mean dispersion for all spatial cells recorded from each region.

(F) Method for computing the mean decoding error. (G) Bootstrapping approach using 80 randomly

selected cells, for 30 bootstrapped samples. (H) Mean decoding error for LS (n = 8 mice), CA1 (n = 4

mice) and CA3 (n = 4 mice). *1, $p < .05$, **, $p < .01$, ****, $p < .0001$, Test used in B, one-way

ANOVA, C-E, Kruskal-Wallis, Dunn's multiple comparison test, H, two-way ANOVA, Sidak's

multiple comparisons test.

A subset of LS cells has stable spatial representations up to 8 days, similar to CA1

To test whether the LS and the hippocampus exhibit comparable evolutions of their

spatial code over time, we recorded from animals implanted in LS and dorsal CA1 in a novel

open field over both short (3 days) and longer timescales (8 days, Fig 4, S14 Fig). We

assessed the stability of the spatial map across sessions, and observed a subset of LS spatially

modulated cells that were stable over time (Fig 4B). Correlating the tuning maps of aligned

LS cell pairs over days lead to a significant increase in mean pairwise correlation value over

days (two-way ANOVA, $F(2, 808) = 3.259$, $p = 0.0389$, main effect of time) with LS cells

being statistically more stable than shuffled comparisons ($F(2,808) = 277.0$, $p < .0001$, main

effect of shuffling, Fig 4C, S14C Fig). We observed the opposite pattern for pyramidal cells

recorded from dorsal CA1, with a significant decrease in mean pairwise correlation value

over days (two-way ANOVA, $F(2,766) = 33.47$, $p < 0.0001$ main effect of time, $F(1,766) =$

319.8, $p < 0.0001$ main effect of shuffle, Fig 4C). For cells recorded in the LS, we observed

the strongest within-session stability on day 3 (Kruskal-Wallis, $H(4) = 19.16$, $p = 0.0003$;

S14E Fig), as well as an increase in spatial information (Kruskal-Wallis, $H(4) = 13.27$, $p =$

0.0041; S14B Fig). There was no significant increase in the proportion of stable cell pairs

over subsequent days (one-way ANOVA, $F(3) = .400$, $p = .9537$, S14D Fig), or the

proportion of spatially modulated cells per animal for each days (one-way ANOVA, $F(3,16)$

$= 1.618, p = .2247$, S14E Fig).

We next assessed the significant tuning map correlations on subsequent day-pairs for

those cells that were significantly spatially modulated on day 1 and were subsequently found

on all recording days (Fig 4D), as well as those that passed our criteria for spatial modulation

on all days (Fig 4E). We then compared the place field correlation for different day pairs

($F(5,363) = 2.625$, $p = .0239$, interaction effect), and we found that on shorter timescales,

tuning maps for LS cells (0.316 ± 0.0507) are significantly less correlated than CA1 ($0.497 \pm$

0.0411; Fisher's LSD day 1-2, $p = 0.0153$, Fig 4F). On the other hand, the correlation for day

pairs 3-8 is higher for the LS (0.369 ± 0.0513) compared to dorsal CA1 (0.174 ± 0.0669;

Fisher's LSD, $p = 0.0098$; Fig 4F). Together, this suggests that a subset of LS spatial cells

encodes spatial information over longer periods of time.

Fig 4: LS place code remains similar to CA1 over longer periods of time.

(A) Experimental setup (top) with a representative example of an animal implanted in the LS, and

aligned spatial footprints of cells recorded over days (bottom). (B) Tuning maps for two sets of stable

cells recorded over all days, with each row being one aligned cell. Within-session correlation

indicated in blue. Tuning map correlation indicated at the bottom in red. (C) Significant tuning map

correlation for aligned cell-pairs (black) versus shuffled pairs (red) for progressive days for LS (day 1-

2, n = 157 cells; day 2-3, n = 122 cells; day 3-8, n = 129 cells; n = 5 mice) and dorsal CA1 (day 1-2, n

= 165 cells; day 2-3, n = 122 cells; day 3-8, n = 99 cells ; n = 3 mice). (D) Significant tuning map

correlations for all cells found on all days for LS (n = 84) and CA1 (n = 86). (E) Matrix of mean

tuning map correlation for all aligned cells that were significantly spatially modulated on each day,

for LS (n = 23-37 cells, n = 5 mice) and CA1 (n = 23 - 41 cells, n = 3 mice). (F) Mean tuning map

correlation for data shown in E. *, $p < .05$, **, $p < .01$, ****, $p < .0001$. Test used in C, two-way

ANOVA, with Sidak's multiple comparisons test, F, two-way ANOVA, with Fisher's LSD post-hoc

test.

LS represents direction, velocity, and acceleration information

In CA1, place cells have been found to carry directional [59–61] and speed-related

information [61–64]. Previous studies found that subsets of LS neurons show some degree of

modulation by direction of travel [31] as well as velocity and acceleration [20] through

correlative measures in spatial alternation or T-maze navigation tasks, where the relationship

between cell activity patterns and movement-associated variables may be confounded with

task-dependent variables. Here, we again employed an information-theoretic approach to

compute the MI between calcium activity and each of these self-motion correlates in a subset

of mice recorded during a 15 min free foraging task in the open field. We found that 28.13%

of LS cells are significantly modulated by head-direction (346/1230 cells, n =19 mice, Fig

5A, S15 Fig), 18.70% by velocity (230/1230 cells, Fig 5C) and 24.63% by acceleration

(303/1230 cells, Fig 5E). We assessed the stability of LS directional and self-motion tuning

over short (3 days) and longer timescales (8 days, S16 Fig), and we observed a subset of LS

directionally modulated cells that were stable over time (S16A Fig). Mean tuning vectors

correlation of aligned LS cell pairs significantly increased over days (two-way ANOVA,

$F(2,774) = 3.682$, $p = 0.0256$, main effect of time) with LS cells being statistically more

stable than shuffled surrogates ($F(1,774) = 54,42$, $p < .0001$, main effect of shuffling, S16C

Fig). For velocity or acceleration tuning, we did not observe such stable tuning over time

(two-way ANOVA, velocity: $F(1,506) = 2.609$, $p = .1069$, acceleration: $F(1,320) = .1986$, p

$= .6561$, main effect of shuffle, S16D,E).

We compared the proportion of LS cells significantly modulated by each of these

variables with the proportion of those cell-types found in dCA1 and dCA3. The proportion of

directionally modulated cells was significantly different across regions (LS: 27.89 ± 2.16,

dCA1: 46.11 ± 4.98, dCA3: 28.92 ± 4.52; Kruskal-Wallis, $H(3) = 6.586$, $p = .0371$, Fig 5B)

with a higher proportion of directionally modulated cells in CA1 as compared to LS (Dunn's

multiple comparisons test, $p = .0340$; Fig 5B). The proportion of velocity encoding cells was

not significantly different between LS (20.43 ± 2.598), CA1 (24.38 ± 7.73), or CA3 (20.04 ±

3.62; Kruskal-Wallis, H(3) = 0.1016, *p* = 0.9504, Fig 5D). Similarly, no difference was found

in the proportion of acceleration encoding cells between LS (25.11 ± 2.34), CA1 (24.14 ±

4.13), or CA3 (22.60 ± 2.98, Kruskal-Wallis, H(3) = 0.3337, *p* = 0.8463, Fig 5F). We found a

proportion of cells that were significantly tuned to more than one of these variables in both

LS as well as dCA1 and dCA3 (Fig 5G, S15A-D Fig, S17 Fig), with 22.03% of LS cells

(271/1230 cells) encoding more than one modality (Fig 5H). Together, this conjunctive

coding for location, directionality, and velocity indicates that LS cells fire in response to

more complex navigational features, similarly to the hippocampus.

Fig 5: LS GABAergic cells encode direction, speed, and acceleration.

(A) Left: Activity of an example LS neuron during free exploration in an open field, top: head

direction (HD, red), middle: binarized activity (yellow), bottom: raw calcium activity (blue). Right:

Corresponding polar plot indicating the probability of the cell being active as a function of the

animals' HD with (black, p(active | direction), red lines indicate 95% upper and lower percentile. Blue

line indicates the normalized time spent for each direction. MI calculated using 40 bins of 9°. S15 Fig.

(B) Proportion of significantly directionally modulated cells in LS, dCA1 and dCA3. (C) Left:

Activity of an example LS neuron during free exploration modulated by velocity (top, red), middle:

binarized activity (yellow), bottom: raw calcium activity (blue). Right: Tuning curve for velocity for

the example cell (red) with 95% upper and lower percentile in grey. MI calculated using 20 bins. (D)

Proportion of significantly speed modulated cells in LS, dCA1 and dCA3. (E) Left: Activity of an

example LS neuron during free exploration modulated by acceleration (top, red), middle: binarized

activity (yellow), bottom: raw calcium activity (blue). Right: Tuning curve for acceleration for the

example cell (red) with 95% upper and lower percentile in grey. MI calculated using 20 bins. (F)

Comparison of proportions of significantly modulated cells for each region. (G) Example of an LS

cell that is both significantly head direction modulated (left), as well as spatially modulated (right).

(H) Left: Proportion of cells that are significantly modulated by only one modality (grey), two

modalities (yellow), three (red) or all 4 of the investigated variables (black). The number above the

bars indicates the absolute number of cells found to be modulated in the total population (n = 1230

cells, n = 19 mice). Right: absolute proportion of cells modulated by any combination of variables.

Test used in B, D, F: Kruskal-Wallis test with Dunn's multiple comparisons test. *, $p < .05$. D =

direction, A = acceleration, S = spatial coding, V = velocity.

To understand the function of this downstream copy of the hippocampal spatial code,

we assessed the downstream targets of the dCA1/dCA3 to LS projection. First, we used a

Cre-dependent, anterograde tracing approach to confirm LS projections to the hypothalamus

and ventral tegmental area in the mouse (S18 Fig), as it was initially described in the rat

[3,5,25]. Next, we leveraged the anterograde transsynaptic properties of Cre-expressing

AAV1 viral vector [65] and assessed the targets of the dorsal CA1-lateral septum (dCA1-LS)

as well as dorsal CA3-LS (dCA3-LS) projections (Fig 6). For either dCA1 or dCA3

injections (Fig 6A, S19 Fig), we observed a direct pathway from the principal cells of the

hippocampus, via the LS, leading to dense innervation of the medial septum (Fig 6B) and the

hypothalamic regions (Fig 6C). We observed that dCA1-LS preferentially targets the LH, and

dCA3-LS targets the hypothalamus more broadly as well as the nuclei of the medial zone.

Whether the GABAergic cells of the lateral septum form functional synapses at the level of

the medial septum/diagonal band complex is debated [5,26,66–68]. Thus, in order to

determine whether the projections observed at the level of the MS with both our anterograde

tracing approaches constituted passing fibers or synaptic connections, we used an AAV-Flex-

Synaptag [69], a Cre-dependent viral construct to express mRuby in the cytosol and eGFP

bound to synaptophysin, a protein primarily expressed in the presynaptic endings of cells

participating in synaptic transmission (Fig 6D). For this, we used a CaMKIIα-Cre mouse line,

expressing Cre only in CAMKII-positive cells, a marker that is abundant in the GABAergic

cells of the LS but absent in the MS, thereby preventing any unspecific labelling. After

expressing AAV-Flex-Synaptag in the intermediate LS, we observed significant synaptophysin-bound projections in the MS (Fig 6E), confirming the existence of a synaptic interface between the LS and the MS. Together, our work shows that LS receives direct inputs from dorsal CA1 and CA3 and in turn projects to regions that are not directly receiving inputs from the hippocampus itself (Fig 6F).

Fig 6: LS cells receiving hippocampal inputs project directly to hypothalamus and medial septum.

(A) Diagram of dual-viral injection strategy for anterograde transsynaptic tracing. The diagram is based on dorsal CA1 targeted injection, dCA3 is also used for injections, together with coronal sections showing primary injection sites in dorsal CA1 (top) or dorsal CA3 (bottom). Red = tdTom expression, blue = DAPI, with zoomed images showing tdTom-positive cell bodies are predominantly located in the pyramidal layer (bottom). For additional images of spread of injection, see S19 Fig. (B) eYFP-positive cell bodies at the anterior dorsal LS and fibers at the level of the medial septum following dCA1 injection (left) and dCA3 injection (right). Inset left: eYFP-positive fibers at the level of MS. (C) eYFP positive axons are seen bilaterally at the level of the lateral hypothalamus following dCA1 injection (left) and dCA3 injection (right). (D) Injection strategy for Cre-dependent AAV. Synaptag mediated tracing in the LS. (E) Coronal section of dorsal LS, with synaptophysin-bound eYFP at the MS. Top right: Zoomed images showing transduction at the injection site. Bottom right: eYFP-positive fibers. (F) Schematic with proposed connections of the LS within the hippocampal network. Scale bars: B, left and right: 800 μm. C, left and right: 500 μm. DM = dorsomedial hypothalamic nucleus, LH = lateral hypothalamus.

LS-cells coding for space, direction and self-motion are non-uniformly distributed

To map out the monosynaptic connections between hippocampal and LS neurons, we injected the anterograde transsynaptic Cre-expressing AAV1 viral vector [65], in either dCA1 or dCA3 (S19 Fig, S20A Fig). We subsequently injected a Cre-dependent eYFP expressing

virus in the LS to visualize cells that receive direct hippocampal input (S20B Fig).

Importantly, this approach allowed us to quantify the cells along the LS anterior-posterior and

dorsal-ventral axis receiving direct hippocampal inputs, and assess their distribution density

within the anterior regions of the LS (Bregma +1.0 to 0.0 anterior-posterior). Because we can

only observe secondary labelled cell-bodies at locations where there was infusion of the

secondary, Cre-dependent virus, we used a relatively large injection volume (total 800 nL,

targeted at the intermediate LS). Both dorsal dCA1 and dCA3 injections resulted in a

majority of labelled LS cell bodies in dorsal regions compared to intermediate and ventral LS

(S20D-F Fig). Hence, LS neurons receiving direct inputs from dorsal CA1 and CA3 appear to

be localized at more dorsal-posterior regions as compared to more ventral-anterior regions of

LS.

Next, we examined whether spatially modulated cells recorded in LS had spatial

properties that were dependent on the localization of their recording. To measure this, we

classified the location of GRIN lens implants along the dorsal-ventral and anterior-posterior

axis (Fig 7A). Post-mortem histological verification of GRIN lens implant combined with

within-animal analysis of the location of recorded cells enabled us to approximate the

location of each cell (Fig 7B). While we observed no differences in cell activity

characteristics, such as bursting index or activity probability along the anterior-posterior axis,

medial-lateral or dorsal-ventral axis (S21 Fig), we found a pronounced increase in the portion

of stable spatially modulated cells (within-session stability > .5) along the anterior-posterior

axis (linear regression, $R^2 = 0.1071$, $p = 0.0022$, the proportion of stable cells per 0.2 mm bin

per animal, n = 24 mice, Fig 7C). We observed a similar gradient along the dorsal-ventral

axis, with a larger proportion of stable spatial cells found at the more dorsal regions of the

LS, although this trend failed to reach significance (linear regression, $R^2 = 0.1399$, $p = .0718$,

n = 24 mice, Fig 7C). Taken together, these data strongly suggest that regions of the LS that

receive stronger innervation from the dorsal hippocampus have a larger proportion of cells

that reliably encode space. This suggests that this information could be directly inherited

from the hippocampus, which should be tested using targeted inactivation of the hippocampal

pyramidal inputs to LS.

We also assessed the topographical distribution of direction and self-motion tuned

cells within the LS. We observed a relatively strong dorsal-ventral gradient in the distribution

of stable (within-session stability > .5) directionally modulated cells (linear regression, R^2 =

0.3837, p = 0.0061, n = 17 mice, Fig 7C), with most directionally modulated cells located at

the more dorsal regions of the LS, but we did not observe such a gradient along the anterior-

posterior axis (linear regression, R^2 = 0.0544, p = .0615, n = 24 mice, Fig 7C). Velocity

modulated cells were primarily found at the ventral pole of the LS (linear regression, R^2 =

0.1836, p = 0.0367, n = 24 mice), but no such gradient was found along the anterior-posterior

axis (linear regression, R^2 = .00348 p = .0874, n = 24 mice). Acceleration-modulated cells

were found evenly distributed throughout the LS (DV: R^2 = .0133, p = .2928; AP: R^2 =

.01332, p = .2929, n = 24 mice).

Fig 7: LS-cells coding for space, direction and self-motion correlates are non-uniformly

distributed along the dorsal-ventral and anterior-posterior axis.

(A) Strategy to record from different anterior-posterior levels in LS (left) and implantation sites

covered. (B) Strategy to approximate cell location. Background: Maximal projection of representative

recording, red: outline of approximated GRIN lens position, white: distance from the centre of the

GRIN lens to the cells of interest. (C) Top: For .2 mm bins along dorsal-ventral axis, proportion of

stable (within-session stability > .5) spatially modulated cells (n = 24 mice), directionally modulated

cells (n = 17 mice), velocity cells (n = 24 mice) and acceleration cells (n = 24 mice) respectively.

Bottom: Same as top for anterior-posterior axis. Each red dot represents the average per animal per

bin along the anterior-posterior axis (left) and along the dorsal-ventral axis. Black lines indicate 95%

confidence intervals. *, $p < .05$, **, $p < .01$, Test used in C: linear regression.

Discussion

While spatial coding has been extensively characterized in the hippocampal

formation, how downstream regions integrate this information has only recently begun to

receive attention. The LS has re-emerged as a critical region implicated in a space encoding

network, although electrophysiological recordings of the LS in rats have led to disparate

estimates of the quantity and quality of place cells in the region [4,20,28,32]: LS place cells

recorded on a circular track were described to be almost absent [4], but were found to be

abundant in reward-seeking tasks [20,32]. Using a large calcium imaging dataset and

unbiased information metric and decoding approaches, we found that 37.80% of GABAergic

LS cells robustly code for space during free exploration in an open field and 43.90% of cells

during linear track alternation. One of the major features of the linear track is the induction of

directional place fields in LS, which may underlie some of the differences we observed in 1D

versus 2D spatial coding characteristics.

In addition to spatial information, we found that the LS also reliably encodes velocity,

acceleration, and directional information, suggesting that the LS encodes more complex

navigational features than previously thought. Neuronal representations in the hippocampus

have been found to change over time, with place code ensembles changing rapidly: an

estimated ~40% [34] to ~75-85% [38] of CA1 neurons were found to remap over days. The

stability of the LS spatial code has received little attention to date. In the current study, we

found a subpopulation of cells in the posterior LS that display stable place fields over eight

days. The functional role of such stability remains to be elucidated. One possibility is that this

ensemble may mediate the stable encoding of contexts over longer durations of time, which

could account for the critical importance of the LS for a number of spatial behaviors,

including contextual fear conditioning [16,41–44], context-reward associations [39,40] and

radial-maze navigation tasks [45,46].

Similar to the change in properties of hippocampal place cells along the dorsal-ventral

axis [70,71], previous work has hinted at a dorsal to ventral organization of the LS place

code: electrophysiology studies recorded a slightly larger proportion of LS place cells in the

more dorsal regions [30]. A complementary anatomical microstructure has been previously

described for the rate-independent phase code in the LS, where the strength of the phase code

increased as a function of recording depth along the dorsal-ventral axis [4]. Here we find that

lateral septal spatially modulated cells are arranged along an anterior-posterior gradient

similar to the gradient of hippocampal inputs into the LS – a finding that helps to reconcile

the variation in previously reported estimates of spatial cells in the LS, which ranged between

5.3% of cells recorded along the dorsal-ventral axis [4] up to 56% recorded at the most dorsal

level of the LS [20]. This pattern suggests that the inhibitory neurons of the LS may inherit a

place code from the hippocampus, with subregions receiving dorsal hippocampal inputs being

most similar to the classical hippocampal place cells. Interestingly, this is not the only case of

GABAergic neurons inheriting some spatial properties from hippocampal pyramidal cells:

interneurons within CA1 have been shown to exhibit significant spatial modulation and

comparable information as pyramidal place cells, although with a higher firing rate and

greater spatial dispersion [72]. It should be considered that, although the hippocampus seems

a likely source of the positional information encoded by the LS, this information could also

arise from incoming projections from any other region projecting to LS, such as the

entorhinal cortex [5,25] which could account for part of head direction and velocity tuning.

In addition to spatial modulation of LS cells, we observed a significant number of

velocity and acceleration modulated cells in the LS. These cells have been previously

described by others on the basis of correlative measures, although estimated proportions vary

widely: for velocity tuning in reward-seeking tasks, estimates range from almost absent [4] to

almost 60% of cells [20]. For acceleration, approximately 45% of LS cells were found to

show some degree of correlation, with almost 30% of recorded cells correlated with both

speed and acceleration [20]. Here, using information metrics and a free exploration task in

which firing characteristics are not confounded with task parameters, we find that 18.70% of

cells were modulated by velocity and 24.63% by acceleration. Cells reliably coding for both

velocity and acceleration were relatively rare (2%). Velocity and acceleration cells were

distributed dissimilarly in the LS, with velocity cells being more abundant towards the ventral

pole, whereas acceleration modulated cells were found to be distributed equally. This raises

the question whether velocity and acceleration information are inherited from different areas

upstream of the LS. Interestingly, cells coding for direction were more abundant towards the

dorsal portion of the LS, but distributed equally along the anterior-posterior axis. Previous

studies have identified classical head direction cells in multiple, well-described circuits,

including the postsubiculum [73,74], anterodorsal thalamic nucleus [75], mammillary bodies

[76], entorhinal cortex [77], retrosplenial cortex [78,79] and parasubiculum [80,81]. A likely

source of directional information to the LS is the entorhinal cortex [82]. Our work supports

the perspective that LS neurons combine a wide range of modalities and may form a complex

representation of context over longer timescales than previously reported. In addition to

space, velocity, and reward information, hippocampal pyramidal cells have been shown to

encode variables such as time [83,84], odor [85,86] and sound frequency [87]. Whether this

information is also relayed to the LS remains to be elucidated. Despite this

interconnectedness with the LH and previous reports of LS place cells being skewed towards

reward in a spatial navigation task [32], we did not observe any overrepresentation of LS

spatially modulated cells around food zones during free exploration or around the reward

zone in a T-maze non-match-to-place paradigm.

The question arises of what the function could be of having a region downstream of

the hippocampus that expresses such seemingly similar coding characteristics. Previous work

estimated the convergence of hippocampal efferents onto LS neurons to be 20 to 800 times

more dense than to any of its cortical targets [4]. Understanding how individual LS neurons

integrate the thousands of synaptic inputs they receive from these hippocampal pyramidal

neurons will be critical to understanding how the hippocampal map is processed downstream.

Through the process of synaptic integration, a target neuron can fire an action potential upon

receiving sufficient temporally coincidental excitatory input on its dendrites. The activation

of a LS neuron could thus require multiple hippocampal pyramidal cells to spike

simultaneously or in close temporal proximity. Hippocampal pyramidal cells will fire

concomitantly when their place fields are in overlapping regions or in close vicinity of

another. Due to the high interconnectivity of LS GABAergic neurons [88,89], the activation

of one LS cell could subsequently lead to the inhibition of other neurons in the nucleus

[24,90], thereby reducing noise. Thus, through this process of coincidence detection and

recurrent inhibition, the spatial map could converge to be represented by a much lower

number of cells. As such, the LS could accurately convey hippocampal information to

downstream regions using a much more distributed code, and thereby support more effective

information transmission and associative learning [91,92].

Finally, our tracing work shows LS cells receiving hippocampal inputs from principal

cells in dorsal CA1 and CA3 in turn project directly to the LH and the MS. Although the

existence of an LS-MS projection has long been debated [5,26,66,68,93], we observed both

dense innervation at the level of the MS after transduction of dCA1-input and dCA3-input

receiving LS cells, as well as synaptic densities between the LS and MS. This suggests that

principal cells of the hippocampus send indirect projections to the MS through LS neurons.

This circuit is complementary to the well-described projections of hippocampal GABAergic

interneurons directly to the MS which may play a key role in theta rhythm generation [94,95].

The MS plays a role in generating and propagating theta rhythms throughout the hippocampal

formation [96–98], which in turn organizes hippocampal place cell activity [99], as well as a

behavioral role in the initiation and velocity of locomotion [100,101]. One hypothesis is that

the information on locomotion-related information reported in specific MS neurons [101]

may originate from the LS. Moreover, we observed a direct projection from CA1 and CA3,

via the LS, to the hypothalamus. This pathway was previously shown by previously using

non-specific monosynaptic anterograde tracing [3], with similar unilateral projection patterns

for CA1 and bilaterally for CA3 as described here. Here we show for the first time that the

same LS neurons that receive inputs from dorsal CA1 and CA3 project directly to the

hypothalamus. The hypothalamus is a highly connected region known for its role in

regulating feeding behaviors [102], arousal and motivation [103], and more recently learning

and memory [104–107]. Additionally, the LH is crucial for the control of locomotion [108–

111] and thought to mediate motivational and goal-directed processes underlying feeding

[12,102,112,113].

Together, our findings show that GABAergic cells of the LS may provide the

hypothalamus and MS with information about location, direction, and speed, and therefore

constitute a core node within a distributed spatial memory network. Within this network, the

LS may be necessary for the translation of spatial information to goal-directed or

environmentally appropriate actions necessary for survival.

Methods

Ethics statement

All procedures were approved by the McGill University and Douglas Hospital Research

Centre Animal Use and Care Committee and in accordance with the Canadian Council on

Animal Care.

Animals

Naive mice (8-16 weeks old) were housed individually on a 12-hour light/dark cycle at 22°C

and 40% humidity with food and water ad libitum. All experiments were carried out during

the light portion of the light/dark cycle. Both male and female mice were used for this study.

Virus injections

Mice, 8-16 weeks old, were anesthetized with isoflurane (5% induction, 0.5-2% maintenance)

and placed in a stereotaxic frame (David Kopf Instruments). Body temperature was

maintained with a heating pad, and eyes were hydrated with gel (Optixcare). Carprofen (10

ml/kg) was administered subcutaneously. All viral injections were performed using a glass

pipette connected to a Nanoject III (Drummond) injector at a flow rate of 1 nL/second. All

stereotaxic coordinates are taken from Bregma (in mm). After surgery, animals were

continuously monitored until recovery. For three days after surgery, mice were provided with

a soft diet supplemented with Carprofen for pain management (MediGel CPF, ~5mg/kg/day).

Virus-mediated expression of genetically encoded calcium indicator

Adeno-associated virus (AAV) of serotype 9 containing the genetically encoded calcium

indicator (GECI) 6 fast (GCaMP6f) under a flex promoter was obtained from the University

of Pennsylvania Vector Core (AAV2/9.Syn.Flex.GCaMP6f.WPRE.SV40, CS0641, Penn

Vector Core via Addgene). Lateral septum viral injections were targeted at the following

coordinates: Anterior LS, AP: 0.86; ML: 0.35; DV: -3.0; Intermediate LS, AP: 0.38; ML:

0.35; DV: -2.7, Posterior LS, AP: 0.10; ML: 0.35; DV: -2.5, with DV + or - .2 to adjust for

more dorsal or ventral placement of injection. Lateral septum injections were done in VGAT-

IRES-Cre transgenic mice (Slc32a1$^{tm2(cre)Lowl}$/J, JAX stock #016962). For hippocampal

principal cell imaging, C57Bl/6J mice (JAX stock #000664) were injected with a viral vector

specifically targeting GCaMP6f expression to CaMKII-positive cells

(AAV5.CamKII.GCaMP6f.WPRE.SV40, Penn Vector Core via Addgene, CS1024).

GCaMP6f injections were targeted at the following coordinates: CA1, AP: -1.8; ML: 1.5;

DV: -1.5; CA3, AP: -2.1; ML: 2.3; DV: -2.2. Three of CA3 animals were Grik4-Cre

transgenic mice (C57BL/6-Tg(Grik4-cre)G32-4Stl/J, JAX stock #006474), restricting

GCaMP6f expression to the pyramidal cells of CA3 specifically.

Retrograde tracing

Retrograde tracing studies were done using a Cre-dependent retrograde Rabies tracing

approach. Briefly, 100 nL of helper virus (AAVdj.hSyn.Flex.TVA.P2A.eGFP.2A.oG,

NeuroPhotonics Center, Laval, Canada; [114,115]) was injected in the LSd (coordinates as

above) of VGAT-IRES-Cre transgenic mice (n = 5 animals, Slc32a1$^{tm2(cre)Lowl}$/J, JAX stock

#016962). Cre specificity was tested by the NeuroPhotonics center. After 4 weeks of

incubation time to allow for the complete expression of the helper virus, 200 nL of

inactivated EnvA pseudotyped rabies virus coupled to mCherry was injected (RABV-EnvA-

deltaG-mCherry, NeuroPhotonics Center) at the same coordinates as the helper virus

injection, and brains were collected 7 days later.

Anterograde transsynaptic tracing

To distinguish synaptic targets from passing projection fibers of dCA1 and dCA3 within the

LS, C57Bl/6J mice (JAX stock #000664) received a bilateral injection of transsynaptic Cre-

inducing anterograde virus (AAV1-hSyn-Cre-WPRE-hGH, Penn Vector Core via Addgene,

V24784) mixed with a Cre-dependent tdTom-expressing reporter virus to visualize injection

sites (AAV2-Ef1a-flex-tdTomato, NeuroPhotonics Center, AAV602) in a 3:1 ratio, 80 nL

total injection volume per hemisphere (n = 5 mice per group). Injections were targeted at the

following coordinates: CA1: AP: -1.86; ML: 1.5; DV: -1.5; CA3: AP: -2.1; ML: 2.3; DV: -

2.2. Following 1 week of incubation time, animals received a bilateral injection of a Cre-

dependent anterograde vector in LS expressing enhanced yellow fluorescent protein (400 nL,

bilateral, AP: 0.5; ML: 0.35; DV: -2.6, AAVdj-DIO-Cheta-eYFP, NeuroPhotonics Center,

AAV938). Brains were collected and processed three weeks post-injection.

Synaptag tracing

To verify presence of synaptic connections at the level of the LS-MS projection, AAVdj-

hSyn-flex-mRuby2-syp-eGFP construct (lot AAV799, NeuroPhotonics Center; Oh et al.,

2014) was injected in the LS of CaMKIIα-Cre mice (B6.Cg-Tg(Camk2a-cre)T29-1Stl/J, JAX

stock #005359), with 100 nL total injection volume injected at 1 nL/second (unilateral, AP:

0.5; ML: 0.35; DV: -2.7). After 4 weeks of incubation time, animals were perfused, and

brains were processed as described below.

GRIN lens implant and baseplate

Two to four weeks post GCaMP6f injection, a 0.5 mm diameter gradient refractive index

(GRIN, Inscopix) lens was implanted above LS, dorsal CA1 or CA3, or a 1.8 mm GRIN

(Edmund Optics) lens was implanted above dorsal CA1. For .5 mm LS lens implant, the lens

was glued to an aluminium baseplate prior to the surgery, which provided two main

advantages: 1) it allowed us to implant the baseplate/lens assembly in one surgery (instead of

separate surgeries) and 2) during the surgery the miniscope (V3, miniscope.org) was used to

image for fluorescence while lowering the lens. The mouse was anesthetized with isoflurane,

and the skull was cleared. A ~.6 mm diameter hole was drilled in the skull at the level of the

injection site. An anchor screw was placed above the contralateral cerebellum to stabilize the

implant. The baseplate and GRIN combination were attached to the miniscope and together

secured to the stereotaxic frame. After making a leading track using a .5 mm diameter needle,

the implant was lowered in place. While lowering the GRIN lens into the tissue, an increase

in fluorescent signal indicates that the injection site has been reached. Both the GRIN lens

and baseplate were permanently secured using C&B-Metabond (Patterson Dental). A plastic

cap was used to protect the GRIN lens from scratching during the recovery period and in

between recording sessions. The animal was given 4 - 6 weeks of recovery time before

experiments started. LS implants were targeted at the following coordinates: Anterior LS,

AP: 0.86; ML: 0.35; DV: -2.9; Intermediate LS, AP: 0.38; ML: 0.35; DV: -2.65; Posterior

LS, AP: 0.10; ML: 0.35; DV: -2.45; with DV + or - .2 to adjust for dorsal, ventral or

intermediate targets; CA3 implants were targeted at AP: -2.1; ML: 2.2; DV: -2.2.

For 1.8 mm GRIN lens implantation, a ~2 mm diameter cranial window was prepared.

Again, an anchor screw was used to secure the implant in place. After removing the dura, a

portion of the cortex above the injection site was gently aspirated using a vacuum pump

without applying pressure on the underlying hippocampal tissue. The 1.8 mm GRIN lens was

lowered at the following coordinates: AP: -1.8; ML: 1.5; DV: -1.8. The GRIN lens was

permanently attached to the skull using Metabond, and Kwik-Sil (World Precision

Instruments) silicone adhesive was placed on the GRIN to protect it. On average 4 weeks

post-implant, the silicone cap was removed and CA1 was imaged using a miniscope mounted

with an aluminium base plate while the mouse was under light anesthesia (~0.5 % isoflurane)

to allow the visualization of cell activity. When a satisfying field of view was found (large

number of neurons, visible landmarks), the baseplate was cemented above the GRIN lens and

a protective cap was used to protect the GRIN from scratches. Imaging would start

approximately 1 week after baseplating.

Miniscope recordings

The animals were habituated by being gently handled for ~5 min per day for 3 days. Animals

were then water or chow-scheduled (2h access per day). Miniscope recordings performed at

30 Hz for 15 minutes, with just one recording session per day to minimize photobleaching.

For open field recordings, animals were freely foraging for randomly placed 10% sucrose or

10% sweetened condensed milk in water rewards.

Recording environments consisted of a 45 x 45 cm dark grey open field with visual

cues, or a 49 x 49 cm white plexiglass open field, which was placed in different recording

chambers. For linear track recordings, rewards (30 ul of 10% sucrose water) were placed at

each end of the 100 cm linear track, or a 130 cm linear track, and the mouse had to consume

one reward before getting the next one delivered. To ensure that all animals received the

same amount of exposure, for analysis of spatial and self-motion modulation, only recordings

were included from the first time the animal spent time in the open field.

For analysis of modulation by food zones or objects, animals were placed again in the,

now familiar, open field on day 1, 3 and 5. On day 2 and 4, food zones or objects were added

in opposing corners of the familiar open field without changing any of the existing visual

cues. Food zones or objects were visible to the animal. Time spent in the region of interest

was computed by taking all timestamps in which animals had their nose within less than 2 cm

from the edge of the food zone or the object.

Non-match to place paradigm: Mice were water scheduled (2 h per day) and trained in

an automatized T-maze (MazeEngineers) to a non-match to place task. Briefly, each trial was

divided into two distinct phases: sample and test. In the initial sample phase, a randomly

selected arm was blocked, which forced mice to explore the opposite arm where the animal

received a 10% sucrose water reward. After consuming the reward, animals would return to

the start box for the test phase, during which both arms could be explored. In the test phase,

only the previously unexplored arm is baited, so that mice have to alternate locations between

sample and test phases. Mice were subjected to 10 trials (sample + test) per day, for 10

consecutive days, and calcium activity in LS was recorded on all days. Data provided in S11

Fig are from Day 10 of training.

Miniscope and behavior video acquisition

Miniscopes (V3 and V4, miniscope.org) were assembled using open-source plans as

described previously (miniscope.org) [116,117]. Imaging data were acquired using a CMOS

imaging sensor (Aptina, MT9V032) and multiplexed through a lightweight coaxial cable.

Data was acquired using a data acquisition (DAQ) box connected via a USB host controller

(Cypress, CYUSB3013). Animal behavior was recorded using a webcam mounted above the

environment. Calcium and behavioral data were recorded using Miniscope custom

acquisition software. The DAQ simultaneously acquired behavioral and cellular imaging

streams at 30 Hz as uncompressed .avi files (1000 frames per file), and all recorded frames

were timestamped in order to perform the subsequent alignment.

Histology

After completion of all experiments, animals were deeply anesthetized with

ketamine/xylazine/acepromazide (100, 16, 3 mg/kg, respectively, intraperitoneal injection).

Mice were then transcardially perfused with 4% paraformaldehyde in PBS (PFA). Brains

were extracted and postfixed in PFA at 4 °C for a minimum of 48 hours. Brains were

sectioned at 50 μm using a vibratome and cryoprotected in a solution of 30% ethylene glycol,

30% glycerol and 40% PBS until used.

Immunohistochemistry

Sections were washed 3 x 20 min in PGT (0.45% Gelatin and 0.25% Triton in 1x PBS) at

room temperature. Next, primary antibodies (1:1000 goat anti-GFP from Novus Biologicals

or 1:10 000 rabbit anti-RFP from VWR (Rockland) were incubated with PGT overnight at

4°C. Following 10, 20, and 40 min washes in PGT, sections were incubated with secondary

antibodies (1:1000 donkey anti-goat coupled to A488 or 1:1000 donkey anti-rabbit coupled to

A555, both from Molecular Probes) in PGT at room temperature for 1 hour. Sections were

subsequently washed for 10 and 20 minutes in PGT and 30 minutes in PBS. Sections were

then mounted on glass slides and permanently coverslipped with Fluoromount that contained

DAPI.

In-vitro **patch clamp electrophysiology**

Mice were deeply anesthetized and acute brain slices were obtained following the protective

recovery method [118]. Briefly, mice were transcardially perfused with N-methyl-D-

glutamine (NMDG-based) solution containing (in mM): 93 NMDG, 93 HCl, 2.5 KCl, 1.2

NaH2PO4, 30 NaHCO3, 20 HEPES, 25 glucose, 5 sodium ascorbate, 2 thiourea, 3 sodium

pyruvate, 10 $MgSO_4$ and 0.5 $CaCl_2$, (pH 7.4, oxygenated with carbogen). The brain was then

quickly extracted and coronal slices (300 μm) were cut using a vibrating microtome (Leica-

VT1000S). Slices were incubated for 10–12 min in 32°C NMDG solution before being

transferred to a holding chamber filled with artificial cerebrospinal fluid (aCSF), kept at room

temperature and containing (in mM): 124 NaCl, 24 NaHCO$_3$, 2.5 KCl, 1.2 NaH$_2$PO$_4$, 5

HEPES, 12.5 glucose, 2 MgSO$_4$, and 2 CaCl$_2$, (pH 7.4, oxygenated with carbogen). For

electrophysiology, slices were transferred to a submerged recording chamber perfused with

aCSF (3 ml /min flow rate, 30 °C). Patch-clamp recordings were obtained from LS

GABAergic (VGAT-positive) neurons expressing the fluorescent marker GFP and visualized

using a 40x water immersion objective on an upright Olympus microscope. Recordings were

performed and analysed using an Axon Multiclamp 700B amplifier and the Clampfit10

software (Molecular Devices). The intrapipette solution contained (in mM) 126 K-gluconate,

4 KCl, 10 HEPES, 4 Mg2ATP, 0.3 Na$_2$GTP, and 10 PO-Creatine, adjusted to pH 7.25 with

KOH (272 mosm). Pipette resistance was 4-6MΩ.

Calcium imaging analysis

Calcium imaging videos were analyzed using the Miniscope Analysis pipeline

(https://github.com/etterguillaume/MiniscopeAnalysis) as described previously [47]. Non-

rigid motion correction was applied using NoRMCorre [119], and videos were spatially

downsampled (3x) before concatenation. Calcium traces were extracted using CNMFe [120]

using the following parameters: gSig = 3 pixels (width of gaussian kernel), gSiz = 20 pixels

(approximate neuron diameter), background_model = 'ring', spatial_algorithm = 'hals',

min_corr = 0.8 (minimum pixel correlation threshold), min_PNR = 8 (minimum peak-to-

noise ratio threshold). After extraction, cells and traces were visually inspected, and cells

with low signal to noise ratio were removed. Raw calcium traces were filtered to remove

high-frequency fluctuations and binarized (normalized amplitude > 2 sd and the first-order

derivative > 0) [47].

Mouse position and head orientation was tracked using a DeepLabCut [121,122]

model trained on several mouse pose markers including head centroid (for position) and nose

tip (used to compute head direction angle, see below). Velocity was extracted by dividing

Δd/Δt where d is the distance and t is time, and subsequently smoothed using a gaussian filter

with sigma = 33 ms to remove movement artefacts. Acceleration was computed by

differentiating (Matlab function *diff*) velocity. Head direction was computed as the angle

between the vertical axis and the line formed by the head centroid and nose tip. Location data

was interpolated to calcium imaging sampling frequency using linear interpolation. Head

directions were interpolated using the same method, but direction data was first unwrapped,

interpolated and converted back into radians.

Activity and bursting index

Following the binarization of raw calcium traces, we compute the probability of a neuron to

be active P(A) using the following formula:

$$P(A) = \frac{time\ active}{total\ time}$$

The bursting index was computed as the probability of a cell being active, given it was

already in an active state, or P(Active, t | Active, t-1). The activity index is computed as the

probability of a cell becoming active when it is in an inactive state, or P(Active, t | Inactive, t-

1).

Spatial modulation

Before processing, input variables were binned (Position in open field: 17 x 17 bins of 3 x 3

cm; Position in linear track: 34 bins of 3 cm; velocity, min = 2.5 cm/s, max = 30 cm/s, 20

bins; acceleration: min = -2 cm/s^2, max = +2 cm/s^2, 20 bins; head direction: 40 bins of 9°).

Velocity was smoothed over thirty frames to reduce discontinuities. For head direction and

positional information, all frames were velocity was < 5cm/s were excluded. The probability

of being active P(A) informs on the activity rate of a neuron. Next, we compute the

probability of spending time in a given bin *i* (spatial, velocity, acceleration or head direction

bin).

$$P(S_i) = \frac{time\ in\ bin\ i}{total\ time}$$

836   Next, we compute the joint probability:

$$P(S_i \cap A) = \frac{time\ active\ while\ in\ bin\ i}{total\ time}$$

837   And the probability that a cell is active given the animal is in a bin:

$$P(A \mid S_i) = \frac{time\ active\ while\ in\ bin\ i}{time\ in\ state}$$

After plotting the binarized activity in space, we perform n = 1000 random circular

permutations. Circular permutations are used to remove the temporal relationship between

neuronal activity and behavior, but it preserves the temporal structure of calcium transients

and therefore leads to more conservative results (as opposed to complete randomization of

every data point, which inflates the significance value of results). Because shuffled surrogates

were not systematically normally distributed, we used a non-parametric approach where the

p-value corresponds to the number of datapoints from the shuffled distribution that are

greater than the actual data for each bin, divided by the number of permutations [123]. A

threshold of $p < .01$ is used to determine the significant data points that make up a significant

place field. In our case, for a cell to be considered spatially modulated, at least one of the

spatial bins must contain activity that is significantly non-random. To compute within-session

stability, the halfway point in the recording was determined by taking the timestamps where

half of the total number of binarized events for the session was reached. Tuning maps were

computed for both the first and second half of the session using the methodology described

above. A Gaussian smoothing filter ($\sigma = 3$) was used to smooth the tuning maps before

computing a correlation between the two. A within-session correlation value of $>.5$ was

considered a stable field.

Calculating mutual information content

Mutual information is used to describe the amount of information about one variable (spatial

location, head direction, velocity or acceleration) through the observation of neuronal

activity, and was calculated using the following formula:

$$MI = \sum_{i=1}^{M} \sum_{j=1}^{2} P(S_i \cap A_j) \times log2(\frac{P(S_i \cap A_j)}{P(S_i) \times P(A_j)})$$

Where M is the total number of possible behavioral states, P($S_i \cap A_j$) is the joint probability of

the animal being in bin i concurrently with activity level j. As we are using binarized activity,

j can only be active or inactive. MI was calculated using 30 bootstrapped surrogates using 50

percent of the data (randomly chosen data points) with replacement, to allow for computation

of mean MI and SEM. Next, the trace was circularly shuffled in order to calculate a shuffled

MI and SEM. Cells were then sorted along with the magnitude of the difference between

mean MI and mean shuffled MI, and a two-way ANOVA with a $p < .01$ threshold was used

to determine whether the difference between the two numbers was significant in order for the

cell to be deemed significantly spatially modulated. In case the assumptions for parametric

ANOVA are not met (normal distribution, variance homogeneity), the significance value was

computed by taking the sum of the shuffled MI values of greater magnitude than actual MI

values, divided by the number of shuffles.

Bayesian decoding

A Bayesian decoder was used to evaluate how well LS neural activity estimated the animal's

location as compared to neural activity recorded from dorsal CA1 and CA3. We used

decoding methods specifically adapted for calcium imaging data with binarized calcium

transients [47]. Using only epochs with velocity > 5 cm/s, a training dataset was generated

using 50% of the data. The remaining 50% of the data was used for testing. Decoding

efficiency (as measured by a decoding error and a decoding agreement score as outlined

below) was calculated using 30 bootstrapped surrogates using randomly chosen data points

with replacement, to allow for computation of mean MI and SEM. In order to allow for fair

comparison between recordings with different numbers of cells recorded, this approach was

done using either 40, 60, 80 or 100 randomly chosen cells with replacement for each

bootstrapped sample. The posterior probability density function, which is the probability that

an animal is in a particular bin S_i , given the neural activity A is calculated using the

following equation:

$$P(S_i \mid A) = \frac{P(A \mid S_i) \times P(S_i)}{P(A)}$$

888   P(S|A) is the posterior probability distribution of states given neuronal activity. No prior

889   assumption is made about the location of the mouse on the linear track, P(S) is kept uniform

890   to attribute equal probability for each location. Every neuron is assumed to be independent of

891   each other. To construct the tuning curves from multiple neurons, we use the following

892   equation:

$$P(S \mid A) = \prod_{k=1}^{N} \frac{P(A_k \mid S) \times P(S)}{P(A_k)}$$

893   With *P(S/A)* a vector of *a posteriori* behavioral states and *N* corresponding to the number of

894   neurons used. For every neuron *k* the tuning curves are constructed, and corresponding

895   posterior location probability can be derived.

896   To reconstruct the position of the mouse, we will consider the location associated with

897   the maximum a posteriori:

$$\hat{y} = arg\,max\,exp\left[\sum_{k=1}^{N} log\left(1 + \frac{P(A_k \mid S) \times P(S)}{P(A_k)}\right) - 1\right]$$

898   With the estimated state among all possible states S. We used a temporal filtering window in

899   the open field of 1.5 seconds and in the linear track of 0.5 seconds to remove erratic jumps in

900   the decoded position of the mouse. Decoding accuracy was measured as 1) decoding error

901   and 2) decoding agreement. In one-dimensional space, decoding error was assessed as

$$decoding\ error\ =\ |\ decoded\ position\ -\ actual\ position\ |$$

902   In two-dimensional space, decoding error was assessed as

$$decoding\ error\ =\ (\sqrt{actual\ position\ -\ decoded\ position})^2$$

904   Decoding score was computed as

$$decoding\ score\ =\ \frac{mean\ (shuffled\ decoding\ error)\ -\ mean\ (decoding\ error)}{length\ of\ track}$$

905   Decoding agreement was defined as the portion of time where the exact location or head

906   direction bin was successfully decoded:

$$decoding\ agreement\ =\ \frac{time\ points\ successfully\ decoded}{total\ time}$$

Tracking cells across sessions

Neurons were tracked over multiple days using a probabilistic method as previously

described [124]. Briefly, spatial footprints were aligned using non-rigid alignment

(transformation smoothness = 2) to correct for brain tissue displacements or miniscope

placement. After alignment, we considered candidate sets of cells to be the same neuron if

their maximal distance was < 12 um, and the spatial correlation between sets of cells was >

.65. Next, we visually identified candidate cell pairs across sessions and manually deleted

those cell pairs that were erroneously selected (< 1% of cases). We assessed the stability of

the spatial representation in LS using simple field correlation, which was assessed by

correlating smoothed place fields of cell pairs. To generate the null hypothesis for place

fields' displacements between pairs of days, we used the place fields tuning maps but shuffled

cell identities over days (30 shuffles).

Materials availability

This study did not generate new unique reagents.

Data and Code availability

All data used for main and supplemental figures are publicly available at [insert DOI from

Gnode].   Code   for   cell   extraction   is   available   at

https://github.com/etterguillaume/MiniscopeAnalysis.   Code   for   behavior   extraction   is

available at https://github.com/etterguillaume/PIMPN. Code for extraction of tuning curves

and decoding were adapted from [47] and can be downloaded at the following address:

https://github.com/etterguillaume/CaImDecoding.   All   other   code   is   available   at

https://github.com/suzannevdveldt/spatialcodingLS.

Statistics

Statistics Statistical analyses were performed using Matlab (Mathworks) and GraphPad Prism

version 6.00 (GraphPad Software, La Jolla, California USA). All data are presented as mean

± standard error of the mean (SEM) and statistical test details are described in the

corresponding results. All t-tests are two-tailed. Normality distribution of each group was

assessed using the Shapiro-Wilk normality test and parametric tests were used only when

distributions were found normal, using Student's t-tests, one-way ANOVA or two-way

ANOVA. For non-normal distributions, non-parametric tests Mann-Whitney or Kruskal-

Wallis tests were performed and described where applicable. Significant main effects or

interactions were followed up with appropriate post hoc testing using Bonferroni corrections

were applicable. $p < 0.05$ was considered statistically significant, or $p < .01$ and described

where applicable *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$, ****, $p < 0.0001$.

Funding

This work was supported by funding from the Canadian Institutes for Health Research

(CIHR), Fonds de la Recherche en Santé du Québec (FRSQ), and the Natural Sciences and

Engineering Research Council of Canada (NSERC). SVDV is supported by a Vanier Canada

Graduate Scholarship and the Richard H. Tomlinson Doctoral Fellowship. C-AM is

supported by a FRQS fellowship.

Acknowledgments

We thank Fernanda Sosa for help with behavioral recording experiments, training of the

animals, immunohistological experiments and quantifications. Alexandra T. Keinath for

valuable comments on a previous version of the manuscript, and Ke Cui for help with colony

maintenance and mice perfusions. We also thank Daniel Aharoni (UCLA) for guidance in

using the UCLA miniscope and Bruno Rivard for help building the miniscopes. The present

study used the services of the Molecular and Cellular Microscopy Platform at the DHRC.

Competing interests

The authors declare no competing interests.

Supplemental figures

S1 Fig: LS receives direct inputs from the pyramidal cells of the hippocampus.

(A) Strategy for retrograde rabies tracing in LS. (B) Mean proportion of cells found in the

hippocampus for each 300μm coronal section; two-way RM ANOVA, F(24, 126) = 2.738, p

= .002, interaction effect, n = 5 mice. (C) Coronal section of ventral hippocampal region,

showing retrogradely labelled cell bodies in subiculum, CA1 and CA3. (D) Retrogradely

labelled cell bodies in the *fasciola cinereum* (FC). (E) Retrogradely labelled cells in the

ventral tegmental area (VTA). (F) Bilateral labelling of the hypothalamic region. (G) Cell

bodies in the periaqueductal grey. (H) Total number of starter cells for each LS subregion,

ANOVA, $F_{(2,12)}$ = 24.61, p > .0001. Scale bars: C, left: 500 µm, right: 300 µm. D,200 µm.

E, 500 µm, F, 500 µm and G, 400 µm. *, p < .05, **, p < .01, ***, p < .001, ns = not

significant. Test used in B, one-way ANOVA. Sub = subiculum, FC = fasciola cinereum, Aq

= aqueduct, mtg = mammillotegmental tract, VTA = ventral tegmental area, MM = medial

mammillary nucleus, PH = posterior hypothalamic area, 3V = third ventricle, LH = lateral

hypothalamus, f = fornix, PAG = periaqueductal grey.

S2 Fig: Histological verification of CA1 implant location.

(A) Examples for CA1 animals included in 1D spatial navigation task, with Cre-dependent

GCaMP6f (green) in a CaMKIIα-Cre mouse with DAPI counterstaining (blue). (B)

Corresponding spatial footprints (SFPs) from extracted cells, color coded for minimum to

maximum mutual information value. (C) Example of approach to compute the correlation

between mutual information and position on the X-axis, linear regression R^2 = 0.005682, p =

2678. (D) Summary of all R^2 values for n = 5 mice.

S3 Fig: Histological verification of CA3 implant location.

(A) Coronal section of dorsal hippocampal region, showing location of GRIN lens implant.

Left: Animals implanted trans-hippocampally. Right: Animals implanted with an

extrahippocampal approach. (B) Coronal sections from anterior to progressively more

posterior regions of the hippocampus, showing extent of damage for a 500 um lens implant.

(C) Comparison between mutual information and probability being active for spatially

modulated cells recorded in animals with trans-hippocampal versus extra-hippocampal

implants (MI: unpaired t-test, t(64) = 0.0367, p = 0.9708; P(A): unpaired t-test, t(51) = .3908,

p = 6875; extra-hipp: n = 22 cells from 2 animals, trans-hipp: n = 44 cells from 3 animals).

S4 Fig: GABAergic cells in the lateral septum are significantly spatially modulated in a

1D environment.

(A) Example of raw calcium fluorescence trace (top), and binarized events (green, bottom).

Trace is divided into early and late recording epochs using 50% of total calcium (using the

area under the curve). Inset: zoomed calcium trace with binarized traces. (B) Linear

correlation between MI and probability being active (linear regression, R^2 = 0.5006, p <

.0001, n = 1030 cells from n = 15 mice). (C) Method for computing the MI and 95%

confidence interval from bootstrapped samples (left) and 30x shuffled surrogates (right). (D)

MI computed from actual traces (black) and shuffled traces (red), sorted by the magnitude of

the difference between these values, two-way ANOVA, F(1029,59740) = 126.3, p < .0001 for

interaction effect. Left inset: zoomed version first 20 cells. Right inset: zoomed version for 20

not significant cells. (E) Scatterplot for significance level for spatial modulation and

probability being active (linear regression, R^2 = .0145, p = 0.007, n = 1030 cells, n = 15

mice). (F) Group averages of actual (grey) versus shuffled (red) MI for spatially modulated

LS cells recorded on linear track (Mann-Whitney test, U = 116394, p < .0001). (G) LS group

averages of spatial cells (grey) versus non-spatial cells (red) for split half stability (Mann-

Whitney test, U = 17743, p < .0011). (H) Examples of significantly spatially modulated cells

representative for each rank, as ranked according to panel C. *, p < .05, ****, p < .0001.

1016

S5 Fig: Intrinsic properties and firing frequency in GABAergic neurons of the lateral

septum and parvalbumin-positive (PV) fast-spiking interneurons of the hippocampus.

(A) VGAT-cre mice were injected with AAVdj-Flex-GFP in LS and patch-clamp whole cell

recordings were performed from fluorescent neurons of the dorsal LS. (B) Representative

photos showing the localization of Cre-dependent fluorescence and VGAT-positive neurons

two weeks after bilateral micro-injections at LS coordinates. Scale bar: 200, 200, 10 μm,

(asterisk marks the location of a patched cell). (C) Current-clamp traces from an GFP positive

LS neuron (shown in b) characterized using depolarizing current injection steps (0-200pA).

(D) Sample traces showing hyperpolarizing and depolarizing responses from a fast-spiking

(PV) hippocampal interneuron. (E) Plot of mean firing frequencies in response to injected

currents of increasing supra-threshold amplitudes in hippocampal PV interneurons (open

circles) and LS VGAT neurons (solid squares). Current injections were 600 ms square pulses.

Firing-frequencies are plotted from threshold current (t) to t + 220 pA (two-way ANOVA

with Bonferroni's multiple comparison test, $F_{(11, 154)} = 21.74$, $p < .0001$ for interaction

effect). *, $p < .05$, ***, $p < .001$, ****, $p < .0001$.

1032

S6 Fig: High proportion of spatially modulated cells in LS is not caused by differences

in probability being active.

(A) Probability being active for all cells recorded for each group (Kruskal-Wallis, $H(3) =$

502.0, $p < .0001$; LS: n = 1030 cells, n = 15 mice. CA1: n = 1251 cells, n = 5 mice, CA3: n =

464 cells, n = 6 mice). (B) Histogram of probabilities being active for all groups (C)

Histogram of MI values for all cells recorded in LS, CA1 and CA3. (D) Using a cut-off of

P(A) > .001 to exclude cells of low activity levels, comparison of proportion of spatially

modulated cells for each region (left, Kruskal-Wallis, $H(3) = 59.57$, $p = .0509$, LS: n = 15

mice. CA1: n = 5 mice, CA3: n = 6 mice) and average split half stability for spatially

modulated cells (right, Kruskal-Wallis, H(3) = 80.48, p < 0.0001, LS: n = 475 cells, n = 15

mice, CA1: n = 363 cells, n = 5 mice, CA3: n = 142 cells, n = 5 mice) . ****, p < .0001.

1044

S7 Fig: Lateral septum spatially modulated cells are significantly directionally

modulated.

(A) Mouse location on day 5 of linear track training, divided in left runs (orange) and right

runs (light blue) with corresponding raw calcium activity (middle, dark blue) and derived

binary trace (bottom, yellow). (B) Probability of cell being active for left (orange) versus

right trajectories (blue), and corresponding mutual information calculated separately for left

and right runs. (C) Corresponding locations where binarized activity was detected (orange for

left trajectories, blue for right trajectories and black for all trajectories). (D) MI values for

right versus left trajectories ranked by magnitude of the difference between the two. Dotted

line p < .01 significance level used to assess spatial modulation of cells, two-way ANOVA,

F(489,28420) = 496.9, p < .0001 for interaction effect. (E) Proportion of spatially modulated

cells in LS that are right-selective (blue, 22.08%), left-selective (orange, 15.52%), spatially

modulated but not directionally modulated (red, 7.67%) and non-spatially modulated (grey,

54,71%, left), CA1 (middle) and CA3 (right).

1059

S8 Fig: Decoding using fewer cells still leads to significant differences between

septohippocampal regions and shuffled surrogates in linear track .

(A) Mean decoding error for location decoding on the linear track using LS cells computed

for 30 bootstrap samples of 40, 60, 80, or 100 cells (black, each dot represents mean of an

animal) compared to a decoded location using shuffled tuning maps (red, each dot represents

mean of an animal), two-way RM ANOVA, F(3,19) = 571.4, p = 0.0024 for main effect of

shuffling, F(3,19) = 6.948, p = 0.0024 for main effect of number of included cells. (B) Left:

same as A, for mean decoding agreement, two-way RM ANOVA, $F(3,19) = 1059$, $p < .0001$

for main effect of shuffling, $F(3,19) = 10.06$, $p = 0.0003$ for main effect of number of

included cells, Right: Method for computing the mean decoding agreement for each

bootstrapped estimate. (C) Same as A, for cells recorded from dorsal CA1. In addition to 30

bootstrapped samples of 40, 60, 80 and 100 cells, this panel includes mean decoding error

using all recorded cells, two-way RM ANOVA, $F(1,20) = 304.3$, $p < .0001$ for main effect of

shuffling, $F(4,20) = 2.647$, $p = 0.0637$ for main effect of number of included cells. (D) Same

as A, for cells recorded from dorsal CA3, two-way RM ANOVA, $F(1,9) = 42.83$, $p = .0001$

for main effect of shuffling, $F(3,9) = 1.480$, $p = 0.2845$ for main effect of number of included

cells. (E) Using only cells with $P(A) > .001$, decoding error for actual versus shuffled dataset

using 60 cells (paired t-tests, LS: $t(5) = .10$, $p = 0.001$, n = 6 mice; CA1: $t(4) = 6.732$, $p =$

.0025, n = 5 mice; CA3: $t(3) = 2.966$, $p = .0592$, n = 4 mice). (F) Using only cells with $P(A)$

$> .001$, mean decoding score for each region (one-way ANOVA, $F(2,12) = 5.765$, $p = 0.0167$.

(G) Same as E, without using a temporal smoothing window (paired t-tests, LS: $t(6) = 10.08$,

$p < 0.0001$, n = 7 mice; CA1: $t(4) = 7.897$, $p = .0014$, n = 5 mice; CA3: $t(3) = 2.614$, $p =$

.0794, n = 4 mice). (H) Same as F, without using a temporal smoothing window (ANOVA,

$F(2,13) = 5.462$, $p = .0190$). *, $p < .05$, **, $p < .01$, ***, $p < .001$, ****, $p < .0001$.

S9 Fig: GABAergic cells in the lateral septum are significantly spatially modulated during free exploration in a 2D environment.

(A) Probability P(active | location) of an example cell to be active in a bin (3 cm) of the open

field (45 or 49 cm size). (B) Significance computed from n = 1000 circular permutations, $p <$

.01 is considered significant. (C) Within session stability is computed by correlation of the

first half (left) and the second half (right) of the recording. (D) MI computed from actual

traces (red) and shuffled traces (black), sorted by the magnitude of the difference between

these values (one-way ANOVA, F(1898, 110142) = 65.50, $p < .0001$ for interaction effect).

Left inset: zoomed version of first 20 cells. Right inset: zoomed version for 20 not significant

cells. (E) Cells are ranked according to the difference between their bootstrapped mean

mutual information (MI) value and 30 circularly shuffled surrogates. Ranks are color-coded

for clarity, n = 1899 cells from N = 28 mice. (F) Examples of significantly spatially

modulated cells with corresponding mean MI value (top) and split within-session stability

(bottom) representative for each rank. (G) Computation of activity index (probability inactive

to active) and bursting index (probability active to active) . (H) Left: Activity index for

spatial vs. non-spatial cells (Mann-Whitney test, U = 342219, $p < .0001$). Right: Bursting

index for spatial versus non-spatial cells (Mann-Whitney test, U = 368276, $p < .0001$).

1102

S10 Fig: LS spatially modulated cells do not anchor to foodzones or objects.

(A) Experimental set-up, with animals freely exploring the same open field for 5 consecutive

days, with two food zones and two objects added in opposing corners on day 2 and 4

respectively. Representative trajectories are shown in grey. Green circles and stars represent

locations of food and objects. (B) Representative example of the average number of cells

active per 3 cm bin for an example mouse (baseline 1: n = 132 cells; food zones: n = 110

cells; baseline 2: n = 118 cells; objects: n = 118 cells; baseline 3: n = 72 cells). (C) For the

same animal, red dots are centroids of spatially modulated cells. (D) Centroids for each

significantly spatially modulated cell of all animals included in analysis (baseline 1: n = 730

cells, n = 10 mice; food zones: n = 559 cells, n = 10 mice; baseline 2: n = 749 cells; n = 10

mice; objects: n = 407 cells, n = 8 mice; baseline 3: n = 471 cells; n = 8 mice). (E) Time spent

in food zone as a percentage of total time spent in session for all animals (n = 10 mice). (F)

Number of visits to food zone (n = 10 mice). (G) Likelihood of cell being active within the

food zone as compared to a shuffle for all cells recorded (Wilcoxon matched-pairs signed

rank test, W = -3490, *p* = .6497, n = 559 cells, n = 10 mice). (H) Time spent in object zone as

a percentage of total time spent in session for all animals (n = 8 mice). (I) Number of visits to

objects (n = 8 mice). (J) Likelihood of a cell being active closely around the objects as

compared to a shuffle (Wilcoxon matched-pairs signed rank test, W = 3789, p = .4251, n =

407 cells, n = 8 mice).

1122

S11 Fig: Significantly spatially modulated cells are not centered around the reward zone

in a non-match to place task

(A) Schematic of T-maze non-match to place task, consisting of a first forced run, followed

by a free run in which the previously non-visited arm is rewarded. (B) Example trajectory of

well-trained animal, with Probability (active | location) of three example cells using 5 cm

bins. (C) For two example animals, red dots are centroids of spatially modulated cells. (D)

Time spent in food zone as a percentage of total time spent in session for all animals (n = 6

mice). (E) Number of visits to reward zone (n = 6 mice). (F) Likelihood of a cell being active

within the reward zone as compared to a shuffle for all cells recorded (Wilcoxon matched-

pairs signed rank test, W = -4783, *p* = 0.2358; n = 365 cells from n = 6 mice, day 10 of

training).

1134

S12 Fig: Differences in spatial coding not due to inclusion of pseudo-silent cells

(A) Probability of being active for spatially modulated cells (Kruskal-Wallis, H(3) = 295.0, *p*

< .0001; Dunn's multiple comparisons test; LS: n = 718 spatial cells from n = 28 mice;

dCA1: n = 323 spatial cells, n = 6 mice; dCA3: n = 138 spatial cells, n = 7 mice). (B) Left:

Using an activity cut-off of P(A) > .001, proportion of spatial cells per animal (one way

ANOVA, F(2,39) = 0.9849, *p* = 0.3826; Tukey's multiple comparisons test; LS: n = 1889

cells from n = 28 mice; dCA1: n = 1017 cells, n = 6 mice; dCA3: n = 521 cells, n = 8 mice).

Middle: Within session stability for spatial cells (Kruskal-Wallis, $H(3) = 89.52$, $p < .0001$;

Dunn's multiple comparisons test; LS: n = 718 spatial cells from n = 28 mice; dCA1: n = 323

spatial cells, n = 6 mice; dCA3: n = 138 spatial cells, n = 7 mice). Right: Mean dispersion for

spatial cells (Kruskal-Wallis, $H(3) = 155.5$, $p < .0001$; Dunn's multiple comparisons test; LS:

n = 718 spatial cells from n = 28 mice; dCA1: n = 323 spatial cells, n = 6 mice; dCA3: n =

138 spatial cells, n = 7 mice). *, $p < .05$, **, $p < .01$, ***, $p < .001$, ****, $p < .0001$.

1148

S13 Fig: Decoding using fewer cells still leads to significant differences between

septohippocampal regions and shuffled surrogates in 2D environment.

(A) Mean decoding error for location decoding in the open field using LS cells computed for

30 bootstrapped samples of 40, 60, 80, and 100 cells (black, each dot represents mean of an

animal) compared to a decoded location using shuffled tuning maps (red, each dot represents

mean of an animal; two-way RM ANOVA, $F(1,33) = 174.9$, $p < 0.0001$ for main effect of

shuffling, $F(3,33) = 0.2041$, $p = 0.893$ for main effect of number of included cells. (B) Left:

same as A, for mean decoding agreement (two-way RM ANOVA, $F(1,33) = 132.8$, $p < .0001$

for main effect of shuffling, $F(3,33) = 1.132$, $p = .3505$ for main effect of number of included

cells). Right: Method for computing the mean decoding agreement for each bootstrap

estimate. (C) Same as A, for cells recorded from dorsal CA1. In addition to 30 bootstrap

samples of 40, 60, 80 and 100 cells, panel includes mean decoding error using all recorded

cells (two-way RM ANOVA, $F(1,17) = 649.0$, $p < .0001$ for main effect of shuffling, $F(4,17)$

$= .7948$, $p = .5447$ for main effect of number of included cells) (D) Same as A, for cells

recorded from dorsal CA3 (two-way RM ANOVA, $F(1,11) = 44.13$, $p < .0001$ for main effect

of shuffling, $F(3,11) = 17.99$, $p = .9078$ for main effect of number of included cells). (E)

Effect of temporal filtering on decoding error in the open field for LS (two-way RM

ANOVA, $F(1,25) = 123.7$, $p < .0001$ for main effect of shuffling, $F(4,25) = 0.1169$, $p = .9753$

for main effect of number of included cells. (F) Comparison of mean decoding error using

P(A) > .001 activity cut-off for cell selection versus without such cut off using 80 cells (LS: n

= 8 mice, CA1: n = 4 mice, CA3: n = 4 mice; two-way RM ANOVA, $F(1,12) = 1.374$, $p =$

2.639 for main effect of activity cut-off, $F(2,12) = 1.023$, $p = .3889$, for main effect of

structure). *, $p < .05$, **, $p < .01$, ***, $p < .001$, ****, $p < .0001$, Test used in A-F, two-way

RM ANOVA.

1173

S14 Fig: LS place code over days

(A) Within session stability for each spatially modulated cell recorded cells in LS (Kruskal-

Wallis, H(4)= 8.921, $p = 0.0304$; Dunn's multiple comparisons test; day 1, n = 181; day 2, n

= 172; day 3, n = 178; day 8, n = 209, n = 5 mice). (B) Average MI for each day for all

recorded cells (Kruskal-Wallis, H(4)= 13.27, $p = 0.0041$; Dunn's multiple comparisons test;

day 1, n = 562; day 2, n = 600; day 3, n = 477; day 8, n = 492, n = 5 mice). (C) Proportion of

cells for tuning map correlation (Friedman test, $\chi2(2) = 0.400$, $p = .9537$, significant cell

pairs only, day1-2, n = 158; day 2-3, n = 122, day 3-8, n = 129). (D) Proportion of stable cells

(tuning map correlation > .3) for each progressive day correlation (one-way ANOVA, F(3,16)

= 1.457, $p = 0.2637$; significant cell pairs only, day 1-2, n = 158; day 2-3, n = 122, day 3-8, n

= 129). (E) Proportion of spatially modulated cells per day (day 1, n = 562; day 2, n = 600;

day 3, n = 477; day 8, n = 492, n = 5 mice). *, $p < .05$, **, $p < .01$

1186

S15 Fig: Direction encoding by lateral septum cells.

(A) Examples of significantly modulated cells. The polar plot indicates the probability of the

cell being active as a function of the animal's head direction. lack lines, p(active | bin); red

lines indicate 95% upper and lower percentile; blue lines indicate the normalized time spent

in each direction. MI calculated using 9° bins. (B) Same as A, for non-modulated cells. (C)

Trajectories (grey) with binarized activity superimposed, color-coded from beginning to end

of the recording for representative cells shown in A. (D) Same as C, but for example, cells

shown in B. (E) Examples of significantly directionally modulated cells with corresponding

mean MI value (top) and within-session stability (bottom) representative for each rank.

1196

S16 Fig: Stability of direction and self-motion encoding over short and longer

timescales

(A) Tuning plots for a stable directionally modulated cell over days, using a similar set up as

described as Fig 4. Tuning map correlation indicated at the bottom in red. (B) Significant

tuning map correlation for aligned cell-pairs (black) versus shuffled pairs (red) for

progressive days for LS (day 1-2, n = 161 cells; day 2-3, n = 119 cells; day 3-8, n = 110 cells;

n = 5 mice) and dorsal CA1 (day 1-2, n = 149 cells; day 2-3, n = 102 cells; day 3-8, n = 90

cells; n = 3 mice). (C) Significant direction map correlations for cells found on all days for

LS (n = 29 cells) and CA1 (n = 24 cells). (D) Same as B, for velocity tuning in LS (day 1-2, n

= 110 cells; day 2-3, n = 73 cells; day 3-8, n = 73 cells; n = 5 mice) and dorsal CA1 (day 1-2,

n = 109 cells; day 2-3, n = 72 cells; day 3-8, n = 61 cells; n = 3 mice). (E) Same as B, for

acceleration tuning in LS (day 1-2, n = 73 cells; day 2-3, n = 47 cells; day 3-8, n = 43 cells; n

= 5 mice) and dorsal CA1 (day 1-2, n = 45 cells; day 2-3, n = 40 cells; day 3-8, n = 35 cells; n

= 3 mice) ****, $p < .0001$. Test used in B, D, E two-way ANOVA, with Sidak's multiple

comparisons test

1212

S17 Fig: Direction, speed and acceleration encoding in dorsal hippocampus.

(A) Left: Distribution of MI (left) and p-values (right) for direction for dCA1, dCA3 and LS.

Right: MI (bits) per binarized event, for all cells recorded from each region (Kruskal-Wallis,

H(3) = 351.2, $p < .0001$; Dunn's multiple comparisons test; LS: n = 1230 cells, n = 19

animals, CA1: n = 677 cells, n = 4 animals, CA3: n = 546 cells, n = 7 animals). (B) Same as

A, for velocity (Kruskal-Wallis, H(3) = 325.3, $p < .0001$; Dunn's multiple comparisons test)

(C) Same as B, for acceleration (Kruskal-Wallis, H(3) = 262.2, $p < .0001$; Dunn's multiple

comparisons test). (D) Left: Proportion of cells that are significantly modulated by only one

modality (grey), two modalities (yellow), three (red) or all 4 of the investigated variables

(black) for dCA1 (n = 677 cells, n = 4 mice). Right: absolute proportion of cells modulated

by any combination of variables. E) Same as D, for dCA3 (n = 546 cells, N = 7 mice). *, $p <$

.05, ****, $p < .0001$

1225

S18 Fig: LS GABAergic neurons project to the medial septum, hypothalamus and

ventral tegmental area.

(A) Injection of anterograde, Cre-dependent eYFP (green) viral tracing in VGAT-Cre mouse

LS. (B) Injection site in intermediate/dorsal LS (LSd) with some cell bodies labelled in

intermediate LS (LSi; Green, eYFP; blue, DAPI counterstaining). (C) eYFP-positive fibers in

the medial septum, (D) hypothalamus and (E) ventral tegmental area. (F) Coronal

hippocampal section shows no anterograde labelling of the hippocampal formation, either

dorsal or (G) ventral. Scale bars: B, 500 μm, C, 500 μm, D, 500 μm, E, 500 μm and F, 500

μm, G) 700 μm. DB = diagonal band, DG = dentate gyrus, LSd = dorsal lateral septum, LSi =

intermediate lateral septum, LSv = ventral lateral septum, MS = medial septum, S =

subiculum, 3V = third ventricle.

1237

1238   **S19 Fig: Extent of primary injection sites in dorsal CA1 and CA3.**

1239   (A) Coronal sections showing expression of tdTom (red) in dorsal CA1 along anterior to

1240   posterior axis. (B) Same as A, for primary injections in dorsal CA3. Scale bars: 500 μm for

1241   all sections.

1242

S20 Fig: Anterograde transsynaptic tracing shows that CA1 and CA3 project

preferentially to dorsal LS.

(A) Primary AAV1 injections in CA1 and CA3, using the same injection strategy as

described in Fig 6A. (B) Coronal section showing expression pattern at different anterior-

posterior levels of the LS, with (red) tdTOM positive CA3 projections to LS and eYFP-

positive second-order transduction in LS (left). Bottom schematic: an overview of the

approximate bregma level of coronal slices shown. Bottom right: zoomed versions showing

tdTom-positive hippocampal afferents, eYFP-positive LS cell bodies, and merge. (C) Total

eYFP-positive cells counted at the level of LS following transsynaptic tracer injection in

dCA1 versus dCA3 (dCA1-LS, N = 3 mice; dCA3-LS, N = 5 mice). (D) Total eYFP-positive

cells along the dorsal-ventral and anterior-posterior axis observed in LS for dorsal CA3

injection and (E) for dorsal CA1 injection. Scale bars: B, top: 500 μm, bottom: 500 μm. C, all

overview images, 500 μm, all zoomed images, 50 μm. ns = not significant.

1256

S21 Fig: No changes in cell activity along dorsal-ventral or anterior-posterior axis of the

LS.

(A) Probability of being active for each cell along the anterior-posterior axis (left), medial-

lateral axis (middle), and dorsal-ventral axis (right). All cells were recorded during free

exploration in the open field (linear regression, AP: R^2 = 2.014 x 10^4, p = 0.5611; ML: R^2 =

1.012 x 10^4, p = 0.6803; DV: R^2 = 2.717 x 10^4, p = 0.6803; n = 1679 cells, n = 24 mice). (B)

Same as A, but for activity index as described in S5 Fig (linear regression, AP: R^2 = 2.691 x

10^4, p = 0.5017; ML: R^2 = 3.287 x 10^5, p = 0.8144; DV: R^2 = 3.254 x 10^4, p = 0.4601; n =

1679 cells, n = 24 mice). (C) Same as for A, but for bursting index (linear regression, AP: R^2

$= 7.085 \times 10^4$, $p = 0.2757$; ML: $R^2 = 7.457 \times 10^5$, $p = 0.7237$; DV: $R^2 = 3.731 \times 10^3$, $p =$

0.0123; n = 1679 cells, n = 24 mice).

1268

References

1. Zhao C, Eisinger B, Gammie S. Characterization of GABAergic neurons in the mouse

lateral septum: a double fluorescence in situ hybridization and immunohistochemical

study using tyramide signal amplification. PLoS One. 2013;8: 1–24.

doi:10.1371/journal.pone.0073750

2. Lin W, McKinney K, Liu L, Lakhlani S, Jennes L. Distribution of vesicular glutamate

transporter-2 messenger ribonucleic acid and protein in the septum-hypothalamus of

the rat. Endocrinology. 2003;144: 662–670.

3. Risold PY, Swanson LW. Structural evidence for functional domains in the rat

hippocampus. Science. 1996;272: 1484–1486. doi:10.1126/science.272.5267.1484

4. Tingley D, Buzsáki G. Transformation of a spatial map across the hippocampal-lateral

septal circuit. Neuron. 2018;98: 1229-1242.e5. doi:10.1016/j.neuron.2018.04.028

5. Swanson LW, Cowan WM. The connections of the septal region in the rat. J Comp

Neur. 1979;186: 621–656. doi:10.1038/nature19789

6. Brito GNO, Thomas GJ. T-maze alternation, response patterning, and septo-

hippocampal circuitry in rats. Behav Brain Res. 1981;3: 319–340. doi:10.1016/0166-

4328(81)90003-6

7. Rawlins JNP, Olton DS. The septo-hippocampal system and cognitive mapping. Behav

Brain Res. 1982;5: 331–358. doi:10.1016/0166-4328(82)90039-0

8. Stanton ME, Thomas GJ, Brito GNO. Posterodorsal septal lesions impair performance

on both shift and stay working memory tasks. Behav Neurosci. 1984;98: 405–415.

doi:10.1037/0735-7044.98.3.405

9.   Thomas GJ. Comparison of effects of small lesions in posterodorsal septum on

spontaneous and rerun correction (contingently reinforced) alternation in rats. J Comp

Physiol Psychol. 1979;93: 685–694. doi:10.1037/h0077608

10.   Thomas GJ, Brito GN. Recovery of delayed alternation in rats after lesions in medial

frontal cortex and septum. J Comp Physiol Psychol. 1980;94: 808–818.

doi:10.1037/h0077834

11.   Calderwood MT, Tseng A, Glenn Stanley B. Lateral septum mu opioid receptors in

stimulation of feeding. Brain Res. 2020;1734: 146648.

doi:10.1016/j.brainres.2020.146648

12.   Carus-Cadavieco M, Gorbati M, Ye L, Bender F, van der Veldt S, Kosse C, et al.

Gamma oscillations organize top-down signalling to hypothalamus and enable food

seeking. Nature. 2017;542: 232–236. doi:10.1038/nature21066

13.   Sweeney P, Yang Y. An inhibitory septum to lateral hypothalamus circuit that

suppresses feeding. J Neurosci. 2016;36: 11185–11195.

doi:10.1523/JNEUROSCI.2042-16.2016

14.   Xu Y, Lu Y, Cassidy RM, Mangieri LR, Zhu C, Huang X, et al. Identification of a

neurocircuit underlying regulation of feeding by stress-related emotional responses.

Nat Commun. 2019;10. doi:10.1038/s41467-019-11399-z

15.   Anthony TE, Dee N, Bernard A, Lerchner W, Heintz N, Anderson DJ. Control of

stress-induced persistent anxiety by an extra-amygdala septohypothalamic circuit. Cell.

2014;156: 522–536. doi:10.1016/j.cell.2013.12.040

16.   Besnard A, Gao Y, Kim MT, Twarkowski H, Reed AK, Langberg T, et al.

Dorsolateral septum somatostatin interneurons gate mobility to calibrate context-

specific behavioral responses. Nat Neurosci. 2019;22: 436–446. doi:10.1038/s41593-

018-0330-y

17.   Sparks P, Ledoux J. The septal complex as seen through the context of fear. In: the

behavioral neuroscience of the septal region. Springer-Verlag; 2000.

18.   Trent NL, Menard JL. The ventral hippocampus and the lateral septum work in tandem

to regulate rats' open-arm exploration in the elevated plus-maze. Physiol Behav.

2010;101: 141–152. doi:10.1016/j.physbeh.2010.04.035

19.   Bender F, Gorbati M, Cadavieco MC, Denisova N, Gao X, Holman C, et al. Theta

oscillations regulate the speed of locomotion via a hippocampus to lateral septum

pathway. Nat Commun. 2015;6: 8521. doi:10.1038/ncomms9521

20.   Wirtshafter HS, Wilson MA. Locomotor and hippocampal processing converge in the

lateral septum. Curr Biol. 2019;29: 3177-3192.e3. doi:10.1016/j.cub.2019.07.089

21.   Bredewold R, Schiavo JK, van der Hart M, Verreij M, Veenema AH. Dynamic

changes in extracellular release of GABA and glutamate in the lateral septum during

social play behavior in juvenile rats: Implications for sex-specific regulation of social

play behavior. Neuroscience. 2015;307: 117–127.

doi:10.1016/j.neuroscience.2015.08.052

22.   Chee S-SA, Menard JL, Dringenberg HC. The lateral septum as a regulator of

hippocampal theta oscillations and defensive behavior in rats. J Neurophysiol.

2015;113: 1831–1841. doi:10.1152/jn.00806.2014

23.   Clemens AM, Wang H, Brecht M. The lateral septum mediates kinship behavior in the

rat. Nat Commun. 2020;11: 1–11. doi:10.1038/s41467-020-16489-x

24.   Leroy F, Park J, Asok A, Brann DH, Meira T, Boyle LM, et al. A circuit from

hippocampal CA2 to lateral septum disinhibits social aggression. Nature. 2018;564:

213–218.

25.   Risold PY, Swanson LW. Connections of the rat lateral septal complex. Brain Res

Rev. 1997;24: 115–195. doi:10.1016/S0165-0173(97)00009-X

26.   Swanson LW, Cowan WM. An autoradiographic study of the organization of the

1342   efferent connections of the hippocampal formation in the rat. J Comp Neur. 1977;172:

1343   49–84. doi:10.1002/cne.901720104

1344   27.   Kita T, Nishijo H, Eifuku S, Terasawa K, Ono T. Place and contingency differential

1345   responses of monkey septal neurons during conditional place-object discrimination. J

1346   Neurosci. 1995;15: 1683–1703.

1347   28.   Leutgeb S, Mizumori S. Context-specific spatial representations by lateral septal cells.

1348   Neuroscience. 2002;112: 655–663. doi:10.1016/S0306-4522(02)00101-X

1349   29.   Nishijo H, Kita T, Tamura R, Uwano T, Terasawa K. Septal neuronal responses related

1350   to spatial representation in monkeys. Hippocampus. 1997;464: 460–464.

1351   30.   Takamura Y, Tamura R, Zhou TL, Kobayashi T, Tran AH, Eifuku S, et al. Spatial

1352   firing properties of lateral septal neurons. Hippocampus. 2006;16: 635–644.

1353   doi:10.1002/hipo

1354   31.   Zhou TL, Tamura R, Kuriwaki J, Ono T. Comparison of medial and lateral septal

1355   neuron activity during performance of spatial tasks in rats. 1999;234: 220–234.

1356   32.   Wirtshafter HS, Wilson MA. Differences in reward biased spatial representations in

1357   the lateral septum and hippocampus. Elife. 2020;9: 1–20. doi:10.7554/eLife.55252

1358   33.   Risold PY, Swanson LW. Chemoarchitecture of the rat lateral septal nucleus. Brain

1359   Res Rev. 1997;24: 91–113. doi:10.1016/S0165-0173(97)00008-8

1360   34.   Gonzalez WG, Zhang H, Harutyunyan A, Lois C. Persistence of neuronal

1361   representations through time and damage in the hippocampus. Science. 2019;365:

1362   821–825. doi:10.1126/science.aav9199

1363   35.   Howard MW, Kahana MJ. A distributed representation of temporal context. J Math

1364   Psychol. 2002;46: 269–299. doi:10.1006/jmps.2001.1388

36.   Mankin EA, Diehl GW, Sparks FT, Leutgeb S, Leutgeb JK. Hippocampal CA2

activity patterns change over time to a larger extent than between spatial contexts.

Neuron. 2015;85: 190–202. doi:10.1016/j.neuron.2014.12.001

37.   Rubin A, Geva N, Sheintuch L, Ziv Y. Hippocampal ensemble dynamics timestamp

events in long-term memory. Elife. 2015;4: 1–16. doi:10.7554/eLife.12247

38.   Ziv Y, Burns LD, Cocker ED, Hamel EO, Ghosh KK, Kitch LJ, et al. Long-term

dynamics of CA1 hippocampal place codes. Nat Neurosci. 2013;16: 264–266.

doi:10.1038/nn.3329

39.   Luo AH, Tahsili-Fahadan P, Wise RA, Lupica CR, Aston-Jones G. Linking context

with reward: a functional circuit from hippocampal CA3 to ventral tegmental area.

Science. 2011;333: 353–358.

40.   McGlinchey EM, Aston-Jones G. Dorsal hippocampus drives context-induced cocaine

seeking via inputs to lateral septum. Neuropsychopharmacology. 2018;43: 987–1000.

doi:10.1038/npp.2017.144

41.   Butler CW, Wilson YM, Gunnersen JM, Murphy M. Tracking the fear memory

engram: Discrete populations of neurons within amygdala, hypothalamus, and lateral

septum are specifically activated by auditory fear conditioning. Learn Mem. 2015;22:

370–384. doi:10.1101/lm.037663.114

42.   Garcia R, Vouimba RM, Jaffard R. Contextual conditioned fear blocks the induction

but not the maintenance of lateral septal LTP in behaving mice. J Neurophysiol.

1997;78: 76–81. doi:10.1152/jn.1997.78.1.76

43.   Reis DG, Scopinho AA, Guimarães FS, Corrêa F, Resstel LBM. Involvement of the

lateral septal area in the expression of fear conditioning to context. Learn Mem.

2010;17: 134–138. doi:10.1101/lm.1534710

44.   Vetere G, Kenney JW, Tran LM, Xia F, Steadman PE, Parkinson J, et al.

Chemogenetic interrogation of a brain-wide fear memory network in mice. Neuron.

1391   2017;94: 363-374.e4. doi:10.1016/j.neuron.2017.03.037

1392 45. Leutgeb S, Mizumori S. Excitotoxic septal lesions result in spatial memory deficits and

1393   altered flexibility of hippocampal single-unit representations. J Neurosci. 1999;19:

1394   6661–72. Available: http://www.ncbi.nlm.nih.gov/pubmed/10414995

1395 46. M'Harzi M, Jarrard LE. Effects of medial and lateral septal lesions on acquisition of a

1396   place and cue radial maze task. Behav Brain Res. 1992;49: 159–165.

1397   doi:10.1016/S0166-4328(05)80160-3

1398 47. Etter G, Manseau F, Williams S. A probabilistic framework for decoding behavior

1399   from in vivo calcium imaging data. Front Neural Circuits. 2020;14: 1–16.

1400   doi:10.3389/fncir.2020.00019

1401 48. Shannon CE. A mathematical theory of communication. Bell Syst Tech J. 1948;27:

1402   379–423. doi:10.1145/584091.584093

1403 49. Skaggs WE, McNaughton BL, Gothard KM. An information-theoretic approach to

1404   deciphering the hippocampal code. Advances in Neural Information Processing

1405   Systems 5 eds Hanson, SJ, Giles, CL & Cowan, JD. 1993. pp. 1030–1037.

1406   doi:10.1109/EUMC.2007.4405137

1407 50. Souza BC., Pavão R, Belchior H, Tort A. On information metrics for spatial coding.

1408   Neuroscience. 2018;375: 62–73. doi:10.1016/j.neuroscience.2018.01.066

1409 51. Gallagher JP, Zheng F, Hasuo H, Shinnick-Gallagher P. Activities of neurons within

1410   the rat dorsolateral septal nucleus (DLSN). Prog Neurobiol. 1995;45: 373–395.

1411   doi:10.1016/0301-0082(95)98600-A

1412 52. Markus EJ, Qin YL, Leonard B, Skaggs WE, McNaughton BL, Barnes CA.

1413   Interactions between location and task affect the spatial and directional firing of

1414   hippocampal neurons. J Neurosci. 1995;15: 7079–7094. doi:10.1523/JNEUROSCI.15-

1415   11-07079.1995

53.   McNaughton BL, Barnes CA, O'Keefe J. The contributions of position, direction and

velocity to single unit activity in the hippocampus of freely-moving rats. Exp Brain

Res. 1983;52.

54.   Navratilova Z, Hoang LT, Daniela Schwindel C, Tatsuno M, McNaughton BL.

Experience-dependent firing rate remapping generates directional selectivity in

hippocampal place cells. Front Neural Circuits. 2012;6: 1–14.

doi:10.3389/fncir.2012.00006

55.   Mau W, Sullivan DW, Kinsky NR, Hasselmo ME, Howard MW, Eichenbaum H. The

same hippocampal CA1 population simultaneously codes temporal information over

multiple timescales. Curr Biol. 2018;28: 1499-1508.e4. doi:10.1016/j.cub.2018.03.051

56.   Pedemonte M, Barrenechea C, Nuñez A, Gambini JP, García-Austt E. Membrane and

circuit properties of lateral septum neurons: Relationships with hippocampal rhythms.

Brain Res. 1998;800: 145–153. doi:10.1016/S0006-8993(98)00517-4

57.   Twery MJ, Gallagher JP. Somatostatin depresses GABA receptor-mediated inhibition

in the rat dorsolateral septal nucleus. Brain Res. 1990;519: 277–286.

doi:10.1016/0006-8993(90)90089-T

58.   Twery MJ, Phelan KD, Gallagher JP. Spontaneous bursting and non-bursting activity

in morphologically identified neurons of the rat dorsolateral septal nucleus, in vitro.

Neuroscience. 1992;46: 669–679. doi:10.1016/0306-4522(92)90153-S

59.   Acharya L, Aghajan ZM, Vuong C, Moore JJ, Mehta MR. Causal influence of visual

cues on hippocampal directional selectivity. Cell. 2016;164: 197–207.

doi:10.1016/j.cell.2015.12.015

60.   Leutgeb S, Ragozzino KE, Mizumori S. Convergence of head direction and place

information in the CA1 region of hippocampus. Neuroscience. 2000;100: 11–19.

doi:10.1016/S0306-4522(00)00258-X

61.   Muller RU, Bostock E, Taube JS, Kubie J. On the directional firing properties of

hippocampal place cells. J Neurosci. 1994;14: 7235–7251. doi:10.1523/jneurosci.14-

12-07235.1994

62.   Czurko A, Hirase H, Csicsvari J, Buzsáki G. Sustained activation of hippocampal

pyramidal cells by "space clamping" in a running wheel. Eur J Neurosci. 1999;11:

344–352. doi:10.1046/j.1460-9568.1999.00446.x

63.   Góis ZHTD, Tort ABL. Characterizing speed cells in the rat hippocampus. Cell Rep.

2018;25: 1872–1884. doi:10.1016/j.celrep.2018.10.054

64.   Iwase M, Kitanishi T, Mizuseki K. Cell type, sub-region, and layer-specific speed

representation in the hippocampal–entorhinal circuit. Sci Rep. 2020;10: 1–23.

doi:10.1038/s41598-020-58194-1

65.   Zingg B, Chou X lin, Zhang Z gang, Mesik L, Liang F, Tao HW, et al. AAV-mediated

anterograde transsynaptic tagging: mapping corticocollicular input-defined neural

pathways for defense behaviors. Neuron. 2017;93: 33–47.

doi:10.1016/j.neuron.2016.11.045

66.   Leranth C, Deller T, Buzsáki G. Intraseptal connections redefined: lack of a lateral

septum to medial septum path. Brain Res. 1992;583: 1–11. doi:10.1016/S0006-

8993(10)80004-6

67.   Staiger JF, Nürnberger F. The efferent connections of the lateral septal nucleus in the

guinea pig: projections to the diencephalon and brainstem. Cell Tissue Res. 1991;264:

391–413. doi:10.1007/BF00319031

68.   Witter MP, Daelmans HE, Jorritsma-Byham B, Staiger JF, Wouterlood FG. Restricted

origin and distribution of projections from the lateral to the medial septal complex in

rat and guinea pig. Neurosci Lett. 1992;148: 164–168. Available:

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cit

ation&list_uids=1284441

69.   Beier KT, Steinberg EE, Deloach KE, Kremer EJ, Malenka RC, Luo L, et al. Circuit

architecture of VTA dopamine neurons revealed by systematic input-output mapping.

Cell. 2015;162: 622–634. doi:10.1016/j.cell.2015.07.015

70.   Jung MW, Wiener SI, McNaughton BL. Comparison of spatial firing characteristics of

units in dorsal and ventral hippocampus of the rat. J Neurosci. 1994;14: 7347–7356.

doi:10.1523/jneurosci.14-12-07347.1994

71.   Kjelstrup KB, Solstad T, Brun VH, Hafting T, Leutgeb S, Witter MP, et al. Finite scale

of spatial representation in the hippocampus. Science. 2008;321: 140–143.

doi:10.1126/science.1157086

72.   Maurer AP, Cowen SL, Burke SN, Barnes CA, McNaughton BL. Phase precession in

hippocampal interneurons showing strong functional coupling to individual pyramidal

cells. J Neurosci. 2006;26: 13485–13492. doi:10.1523/JNEUROSCI.2882-06.2006

73.   Taube J, Muller R, Ranck J. Head-direction cells recorded from the postsubiculum in

freely moving rats. I. Description and quantitative analysis. J Neurosci. 1990;10: 420–

435. doi:10.1523/jneurosci.10-02-00420.1990

74.   Taube JS, Muller RU, Ranck JB. Head-direction cells recorded from the

postsubiculum in freely moving rats. II. Effects of environmental manipulations. J

Neurosci. 1990;10: 436–447. doi:10.1523/jneurosci.10-02-00436.1990

75.   Taube JS. Head direction cells recorded in the anterior thalamic nuclei of freely

moving rats. J Neurosci. 1995;15: 70–86. doi:10.1523/jneurosci.15-01-00070.1995

76.   Stackman RW, Taube JS. Firing properties of rat lateral mammillary single units: Head

direction, head pitch, and angular head velocity. J Neurosci. 1998;18: 9020–9037.

doi:10.1523/jneurosci.18-21-09020.1998

77.   Sargolini F, Fyhn M, Hafting T, McNaughton BL, Witter MP, Moser MB, et al.

Conjunctive representation of position, direction, and velocity in entorhinal cortex.

Science. 2006;312: 758–762. doi:10.1126/science.1125572

78.   Chen LL, Lin LH, Green EJ, Barnes CA, McNaughton BL. Head-direction cells in the

rat posterior cortex. I. Anatomical distribution and behavioral modulation. Exp Brain

Res. 1994;101: 8–23. doi:10.1007/BF00243212

79.   Sharp PE, Blair HT, Cho J. The anatomical and computational basis of the rat head-

direction cell signal. Trends Neurosci. 2001;24: 289–294. doi:10.1016/s0030-

6657(08)70226-9

80.   Boccara CN, Sargolini F, Thoresen VH, Solstad T, Witter MP, Moser EI, et al. Grid

cells in pre-and parasubiculum. Nat Neurosci. 2010;13: 987–994. doi:10.1038/nn.2602

81.   Cacucci F, Lever C, Wills TJ, Burgess N, O'Keefe J. Theta-modulated place-by-

direction cells in the hippocampal formation in the rat. J Neurosci. 2004;24: 8265–

8277. doi:10.1523/JNEUROSCI.2635-04.2004

82.   Oh SW, Harris JA, Ng L, Winslow B, Cain N, Mihalas S, et al. A mesoscale

connectome of the mouse brain. Nature. 2014;508: 207–214. doi:10.1038/nature13186

83.   Eichenbaum H. Time cells in the hippocampus: A new dimension for mapping

memories. Nat Rev Neurosci. 2014;15: 732–744. doi:10.1038/nrn3827

84.   Pastalkova E, Itskov V, Amarasingham A, Buzsáki G. Internally generated cell

assembly sequences in the rat hippocampus. Science. 2008;321: 1322–1328.

doi:10.1126/science.1159775

85.   Deshmukh SS, Bhalla US. Representation of odor habituation and timing in the

hippocampus. J Neurosci. 2003;23: 1903–1915. doi:10.1523/jneurosci.23-05-

01903.2003

86.   MacDonald CJ, Carrow S, Place R, Eichenbaum H. Distinct hippocampal time cell

sequences represent odor memories in immobilized rats. J Neurosci. 2013;33: 14607–

14616. doi:10.1523/JNEUROSCI.1537-13.2013

87. Aronov D, Nevers R, Tank DW. Mapping of a non-spatial dimension by the

hippocampal-entorhinal circuit. Nature. 2017;543: 719–722. doi:10.1038/nature21692

88. Alonso JR, Frotscher M. Organization of the septal region in the rat brain: a Golgi EM

study of lateral septal neurons. J Comp Neurol. 1989;286: 472–487.

89. Staiger JF, Nürnberger F. Pattern of afferents to the lateral septum in the guinea pig.

Cell Tissue Res. 1989;257: 471–490. doi:10.1007/BF00221457

90. Phelan KD, Hasuo H, Twery MJ, Gallagher JP. Projection neurons in the rat

dorsolateral septal nucleus possess recurrent axon collaterals. Neurosci Lett. 1989;97:

259–265. doi:10.1016/0304-3940(89)90607-1

91. Isely G, Hillar CJ, Sommer FT. Deciphering subsampled data: Adaptive compressive

sampling as a principle of brain communication. Adv Neural Inf Process Syst 23 24th

Annu Conf Neural Inf Process Syst 2010, NIPS 2010. 2010; 1–9.

92. Palm G, Sommer FT. Associative data storage and retrieval in neural networks.

Domany E., van Hemmen JL, Schulten K, editors. Models of Neural Networks III.

Physics of Neural Networks. Springer, New York, NY.; 2006.

93. Staiger JF, Nürnberger F. The efferent connections of the lateral septal nucleus in the

guinea pig: intrinsic connectivity of the septum and projections to other telencephalic

areas. Cell Tissue Res. 1991;264: 415–426. doi:10.1007/BF00319032

94. Tóth K, Borhegyi Z, Freund TF. Postsynaptic targets of GABAergic hippocampal

neurons in the medial septum-diagonal band of broca complex. J Neurosci. 1993;13:

3712–3724. doi:10.1523/jneurosci.1413-04.2004

95. Manseau F, Goutagny R, Danik M, Williams S. The hippocamposeptal pathway

generates rhythmic firing of GABAergic neurons in the medial septum and diagonal

bands: an investigation using a complete septohippocampal preparation in vitro. J

Neurosci. 2008;28: 4096–4107. doi:10.1523/JNEUROSCI.0247-08.2008

96.   Leão RN, Targino ZH, Colom L V., Fisahn A. Interconnection and synchronization of

neuronal populations in the mouse medial septum/diagonal band of Broca. J

Neurophysiol. 2015;113: 971–980. doi:10.1152/jn.00367.2014

97.   Patel J, Schomburg EW, Bere A, Fujisawa S. Local generation and propagation of

ripples along the septotemporal axis of the hippocampus. 2013;33: 17029–17041.

doi:10.1523/JNEUROSCI.2036-13.2013

98.   Robinson J, Manseau F, Ducharme G, Vigneault E, El Mestikawy S, Williams S.

Optogenetic activation of septal glutamatergic neurons drive hippocampal theta

rhythms. 2016;36: 3016–3023. doi:10.1523/JNEUROSCI.2141-15.2016

99.   Buzsáki G. Theta oscillations in the hippocampus. Neuron. 2002;33: 325–340.

doi:10.1016/S0896-6273(02)00586-X

100. Fuhrmann F, Justus D, Sosulina L, Kaneko H, Beutel T, Friedrichs D, et al.

Locomotion, theta oscillations, and the speed-correlated firing of hippocampal neurons

are controlled by a medial septal glutamatergic circuit. Neuron. 2015;86: 1253–1264.

doi:10.1016/j.neuron.2015.05.001

101. Justus D, Dalügge D, Bothe S, Fuhrmann F, Hannes C, Kaneko H, et al. Glutamatergic

synaptic integration of locomotion speed via septoentorhinal projections. Nat

Neurosci. 2017;20: 16–19. doi:10.1038/nn.4447

102. Stuber GD, Wise RA. Lateral hypothalamic circuits for feeding and reward. Nat

Neurosci. 2016;19: 198–205. doi:10.1016/j.physbeh.2017.03.040

103. Burdakov D, Peleg-Raibstein D. The hypothalamus as a primary coordinator of

memory updating. Physiol Behav. 2020;223: 112988.

doi:10.1016/j.physbeh.2020.112988

104. Aitta-Aho T, Pappa E, Burdakov D, Apergis-Schoute J. Cellular activation of

hypothalamic hypocretin/orexin neurons facilitates short-term spatial memory in mice.

Neurobiol Learn Mem. 2016;136: 183–188. doi:10.1016/j.nlm.2016.10.005

105.   Izawa S, Chowdhury S, Miyazaki T, Mukai Y, Ono D, Inoue R, et al. REM sleep-

active MCH neurons are involved in forgetting hippocampus-dependent memories.

Science. 2019;365: 1308–1313. doi:10.1126/science.aax9238

106.   Kosse C, Burdakov D. Natural hypothalamic circuit dynamics underlying object

memorization. Nat Commun. 2019;10: 1–8. doi:10.1038/s41467-019-10484-7

107.   Petrovich GD. Lateral hypothalamus as a motivation-cognition interface in the control

of feeding behavior. Front Syst Neurosci. 2018;12: 1–7. doi:10.3389/fnsys.2018.00014

108.   Gladfelter WE, Brobeck JR. Decreased spontaneous locomotor activity in the rat

induced by hypothalamic lesions. Am J Physiol. 1962;203: 811–817.

doi:10.1152/ajplegacy.1962.203.5.811

109.   Grillner S, Wallén P, Saitoh K, Kozlov A, Robertson B. Neural bases of goal-directed

locomotion in vertebrates-An overview. Brain Res Rev. 2008;57: 2–12.

doi:10.1016/j.brainresrev.2007.06.027

110.   Kosse C, Schöne C, Bracey E, Burdakov D. Orexin-driven GAD65 network of the

lateral hypothalamus sets physical activity in mice. Proc Natl Acad Sci U S A.

2017;114: 4525–4530. doi:10.1073/pnas.1619700114

111.   Sinnamon HM. Preoptic and hypothalamic neurons and the initiation of locomotion in

the anesthetized rat. Prog Neurobiol. 1993;41: 323–344. doi:10.1016/0301-

0082(93)90003-B

112.   Mahler S V., Moorman DE, Smith RJ, James MH, Aston-Jones G. Motivational

activation: A unifying hypothesis of orexin/hypocretin function. Nat Neurosci.

2014;17: 1298–1303. doi:10.1038/nn.3810

113.   Mogenson GJ, Jones DL, Yim CY. From motivation to action: Functional interface

1591 between the limbic system and the motor system. Prog Neurobiol. 1980;14: 69–97.

1592 doi:10.1016/0301-0082(80)90018-0

1593 114. Callaway EM. Transneuronal circuit tracing with neurotropic viruses. Curr Opin

1594 Neurobiol. 2008;18: 617–623. doi:10.1016/j.conb.2009.03.007

1595 115. Wickersham IR, Finke S, Conzelmann KK, Callaway EM. Retrograde neuronal tracing

1596 with a deletion-mutant rabies virus. Nat Methods. 2007;4: 47–49.

1597 doi:10.1038/nmeth999

1598 116. Cai DJ, Aharoni D, Shuman T, Shobe J, Biane J, Song W, et al. A shared neural

1599 ensemble links distinct contextual memories encoded close in time. Nature. 2016;534:

1600 115–118. doi:10.1038/nature17955

1601 117. Ghosh KK, Burns LD, Cocker ED, Nimmerjahn A, Ziv Y, Gamal A El, et al.

1602 Miniaturized integration of a fluorescence microscope. Nat Methods. 2011;8: 871–878.

1603 doi:10.1038/nmeth.1694

1604 118. Ting JT, Daigle TL, Chen Q, Feng G. Acute brain slice methods for adult and aging

1605 animals: Application of targeted patch clamp analysis and optogenetics. Methods Mol

1606 Biol. 2014;1183: 221–242. doi:10.1007/978-1-4939-1096-0_14

1607 119. Pnevmatikakis EA, Giovannucci A. NoRMCorre: An online algorithm for piecewise

1608 rigid motion correction of calcium imaging data. J Neurosci Methods. 2017;291: 83–

1609 94. doi:10.1016/j.jneumeth.2017.07.031

1610 120. Zhou P, Resendez SL, Rodriguez-Romaguera J, Jimenez JC, Neufeld SQ, Stuber GD,

1611 et al. Efficient and accurate extraction of in vivo calcium signals from

1612 microendoscopic video data. 2017. doi:http://arxiv.org/abs/1605.07266

1613 121. Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, et al.

1614 DeepLabCut: markerless pose estimation of user-defined body parts with deep

1615 learning. Nat Neurosci. 2018;21: 1281–1289. doi:10.1038/s41593-018-0209-y

1616   122.   Nath T, Mathis A, Chen AC, Patel A, Bethge M, Mathis MW. Using DeepLabCut for

1617   3D markerless pose estimation across species and behaviors. Nat Protoc. 2019;14:

1618   2152–2176. doi:10.1038/s41596-019-0176-0

1619   123.   Cohen MX. Analyzing neural time series data: Theory and practice. Cambridge, MA:

1620   The MIT Press; 2014.

1621   124.   Sheintuch L, Rubin A, Brande-Eilat N, Geva N, Sadeh N, Pinchasof O, et al. Tracking

1622   the same neurons across multiple days in Ca2+ imaging data. Cell Rep. 2017;21:

1623   1102–1115. doi:10.1016/j.celrep.2017.10.013

1624

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A *LS*
head direction
360°
0°
binarized activity
2 sd
calcium activity
10 sec

90
0.2
180   0
270
Direction (°)

B
% Directional cells
80
60
40
20
*
LS CA1 CA3

C *LS*
velocity
25 cm/s
0 cm/s
binarized activity
2 sd
calcium activity
10 sec

P(active | speed)
0.8
0.4
0.0
0   15   30
Velocity (cm/s)

D
Velocity cells (%)
60
40
20
ns
LS CA1 CA3

E *LS*
acceleration
1 cm/s²
0
-1 cm/s²
binarized activity
2 sd
calcium activity
10 sec

P(active | acceleration)
0.4
0.2
0.0
-1.5   0   2.0
Acceleration (cm.s⁻²)

F
Acceleration cells (%)
60
40
20
ns
LS CA1 CA3

G
90
0.1
180   0
270
MI = 0.0272

End
Start
spatial MI = 0.0595

H
D = Direction   S = Spatial   A = Acceleration   V = Velocity

proportion of cells
0.12
0.08
0.04
0.00

168 55 82 55   105 29 20 56 38 23   58 39 15 9   31
S V A D   D+S D+A D+V S+A S+V A+V   D+S+A D+S+V S+A+V A+D+V S+V+A+D

2.52%
9.84%
36.34%
22.03%
29.27%
Total=1230

4 modalities   1 modality
3 modalities   Not modulated
2 modalities

Figure 6

Figure 7

A

+0.86
anterior LS

+0.38
intermediate LS

+0.02
posterior LS

GRIN lens location

B

medial

anterior

+ 220 μm

+ 50 μm

500 μm

C

Dorsal-Ventral axis

spatial

p = 0.0718

Stable spatial cells per bin (%)

Dorsal - Ventral axis
(distance from Bregma)

direction

R² = 0.3837
**

Stable directional cells per bin (%)

Dorsal - Ventral axis
(distance from Bregma)

velocity

R² = 0.1836
*

Velocity cells per bin (%)

Dorsal - Ventral axis
(distance from Bregma)

acceleration

ns

Acceleration cells per bin (%)

Dorsal - Ventral axis
(distance from Bregma)

Anterior-Posterior axis

+0.86

+0.02

R² = 0.1071
**

Stable spatial cells per bin (%)

Anterior - Posterior axis
(distance from Bregma)

p = 0.0615

Stable directional cells per bin (%)

Anterior - Posterior axis
(distance from Bregma)

p = 0.0874

Velocity cells per bin (%)

Anterior - Posterior axis
(distance from Bregma)

ns

Acceleration cells per bin (%)

Anterior - Posterior axis
(distance from Bregma)

S1 Fig

S2 Fig

S3 Fig

A

trans-hippocampal implant   extra-hippocampal implant

B

anterior – posterior axis

C

S4 Fig

A

Start   early | late   event half-point   End

90 sec

B

R² = .5006
****

MI (bits)
P(a)

C

n = 30 bootstrap samples

Epoch included in MI computation

a₁
a₂
a₃
a₄
a₅
⋮
aₙ

s₁
s₂
s₃
s₄
s₅
⋮
sₙ

D

Shuffled MI
Actual MI

MI (bits)
Cell rank

P < 0.01   ns

E

LS

p value
P(a)

F

MI (bits)
Actual   Shuffle
****

G

within-session stability
Spatial   ns
****

H

spatially modulated   non-spatially modulated

rank 10   rank 50   rank 100   rank 500

P(active | location)
Position on track (cm)

p < .01   *p* > .01

S5 Fig

S6 Fig

A

B

C

D

P(A) > .001

S7 Fig

A

left runs   right runs   *LS*

location (cm)

100
0

calcium activity

binarized activity

60 sec

B

LS

left MI = 0.2305
right MI = 0.0112

P(active | location)

1.0
0.5
0.0

0   50   100
Position on track (cm)

C

left trajectories   right trajectories   all trajectories

lap number

1
21

1
22

1
21

0   50   100
position on track (cm)

0   50   100
position on track (cm)

0   50   100
position on track (cm)

D

LS

Right - selective   Left - selective

MI (bits)

0.3
0.2
0.1
0.0

0   200   400

MI right trajectories
MI left trajectories

E

right selective   spatial, non - directional
left selective   non spatial

LS   *CA1*   *CA3*

Total=1030   Total=1251   Total=463

S8 Fig

S9 Fig

S10 Fig

A
GCaMP6f
LS
Baseline 1 — day 1
Foodzones — day 2
Baseline 2 — day 3
Objects — day 4
Baseline 3 — day 5

B example animal
3 cm bins
avg. N cells active
0 — 5

C
Baseline 1   Foodzones   Baseline 2   Objects   Baseline 3
• place cell centroid

D population
Baseline 1   Foodzones   Baseline 2   Objects   Baseline 3

E Time in zone (%)

F number of visits
Foodzones

G likelihood — ns
actual   shuffle

H Time in zone (%)

I number of visits
Objects

J likelihood — ns
actual   shuffle

Non Match To Place

S12 Fig

P(A) > .001

A

B

S13 Fig

S14 Fig

S15 Fig

A significantly modulated (p < .01)

MI = 0.0281   MI = 0.0272   MI = 0.0312

B non-modulated (p > .01)

MI = 0.0081   MI = 0.0071   MI = 0.0068

C

D

E

significantly modulated (p < .01)   non-modulated (p > .01)

rank 10   rank 50   rank 100   rank 500   rank 1000

Cell 28, ID48 MI: 0.0272 Stability: 0.92	Cell 3, ID48 MI: 0.0171 Stability: 0.17	Cell 79, ID51 MI: 0.0173 Stability: 0.87	Cell 40, ID786 MI: 0.0083 Stability: 0.73	Cell 9, ID134 MI: 0.0078 Stability: 0.49
Cell 37, ID25 MI: 0.0234 Stability: 0.69	Cell 21, ID133 MI: 0.0174 Stability: 0.78	Cell 68, ID47 MI: 0.0122 Stability: 0.64	Cell 73, ID24 MI: 0.0097 Stability: 0.50	Cell 65, ID24 MI: 0.0085 Stability: -0.07
Cell 79, ID8 MI: 0.0281 Stability: 0.46	Cell 33, ID8 MI: 0.0179 Stability: 0.69	Cell 44, ID8 MI: 0.0175 Stability: 0.89	Cell 57, ID47 MI: 0.0097 Stability: 0.56	Cell 77, ID48 MI: 0.0077 Stability: 0.59

P < .01   P> .01

S16 Fig

A

Tuning map correlation: 0.929   0.7816   0.5975

B

Direction

LS

CA1

C

Direction

LS

CA1

D

Velocity

LS

CA1

E

Acceleration

LS

CA1

S18 Fig

S19 Fig

S20 Fig

A

CA1 injected

CA1
Dapi tdTom

CA3 injected

Dapi tdTom CA3

B *Lateral Septum*

anterior | intermediate | posterior

LSd
LSi
LSv

LSd
LSi
Ld

LSd

anterior | intermediate | posterior

LSd
LSi

LSd
LSi

+0.98 0.60 0.10 LS

tdTom eYFP merged

C

transfected cells

ns

dCA1-LS dCA3-LS

D

dCA3-LS

bregma

LSd LSi LSv

Transfected cells

dCA3-LS

transfected cells

LSd
LSi
LSv

bregma

E

dCA1-LS

LSd LSi LSv

Transfected cells

S21 Fig