Full title: Conjunctive spatial and idiothetic codes are topographically organized in the

GABAergic cells of the lateral septum

Authors: Suzanne van der Veldt[1], Guillaume Etter[1], Fernanda Sosa[1], Coralie-Anne Mosser[1],

Sylvain Williams[1*]

[1] McGill University & Douglas Mental Health University Institute, Montreal, Canada

6875, Boul. LaSalle, Verdun (Quebec), H4H 1R3, Phone: 514 761-6131, Fax: 514 888-4067

*Corresponding author: sylvain.williams@mcgill.ca

Author contributions

SV and SW designed the study. SV performed surgeries, behavioral recording experiments,

immunohistological experiments, and calcium data analysis. GE wrote code for MI

calculation and Bayesian decoding and implanted mice for CA1 recordings. FS assisted with

behavioral recording experiments, immunohistological experiments and performed

quantifications. CAM implanted two mice that were used in the current study. SV and SW

wrote the manuscript with inputs from all authors.

Abstract

The hippocampal spatial code's relevance for downstream neuronal populations – particularly

its main subcortical output, the lateral septum (LS) - is still poorly understood. Here, using

calcium imaging combined with unbiased analytical methods, we functionally characterized

and compared the spatial tuning of LS GABAergic cells to those of dorsal CA3 and CA1

cells. We identified a significant number of LS cells that are modulated by place, speed,

acceleration, and direction, as well as conjunctions of these properties, directly comparable to

hippocampal CA1 and CA3 place cells. Interestingly, Bayesian decoding of position based on LS place cells reflected the animal's location as accurately as decoding using the activity of hippocampal pyramidal cells. A portion of LS place cells showed stable fields over the course of multiple days, potentially reflecting long-term episodic memory. The distributions of cells exhibiting these properties formed gradients along the anterior-posterior and dorsal-ventral axes of the LS, directly reflecting the organization of hippocampal inputs to the LS. Finally, we show using trans-synaptic tracing that LS neurons receiving CA3 and CA1 excitatory input send projections to the hypothalamus and medial septum, regions that are not targeted directly by principal cells of the dorsal hippocampus. Together, our findings demonstrate that the LS accurately and robustly represents spatial and idiothetic information and is uniquely positioned to relay this information from the hippocampus to its downstream regions, thus occupying a key position within a distributed spatial memory network.

Introduction

The lateral septum (LS) is an anatomically complex structure uniquely positioned within a broader distributed spatial memory network. The LS consists almost exclusively of highly interconnected inhibitory GABAergic cells (Zhao et al., 2013) that are densely innervated by the principal cells of the hippocampus (Risold and Swanson, 1996; Tingley and Buzsáki, 2018). Previous research has implicated the LS in a wide variety of behaviors, including spatial and working memory (Brito and Thomas, 1981; Rawlins and Olton, 1982; Stanton et al., 1984; Thomas, 1979; Thomas and Brito, 1980), the regulation of feeding (Calderwood et al., 2020; Carus-Cadavieco et al., 2017; Sweeney and Yang, 2016; Xu et al., 2019), anxiety (Anthony et al., 2014; Besnard et al., 2019; Sparks and Ledoux, 2000; Trent and Menard, 2010), locomotion (Bender et al., 2015; Wirtshafter and Wilson, 2019), and

social behaviors (Bredewold et al., 2015; Chee et al., 2015; Clemens et al., 2020; Leroy et al.,

2018).

To date, functional characterizations of the LS have often focused on the spatial

coding properties of these neurons, primarily due to the extensive excitatory input from

positionally-tuned hippocampal pyramidal neurons, commonly referred to as 'place cells'

(Leroy et al., 2018; Risold and Swanson, 1997; Swanson and Cowan, 1977; Tingley and

Buzsáki, 2018). Yet, this body of work has not yet yielded a consensus. Several studies have

reported place cells in the LS, but the number, information content, and stability vary widely

from study to study (Kita et al., 1995; Leutgeb and Mizumori, 2002; Nishijo et al., 1997;

Takamura et al., 2006; Wirtshafter and Wilson, 2019; Zhou et al., 1999), while others

described a lack of canonical place cells (Tingley and Buzsáki, 2018). As previous studies

often depended on variable criteria of what defines a place cell, estimates range from 5.3%

(Tingley and Buzsáki, 2018) to 56.0% (Wirtshafter and Wilson, 2019) of LS cells classified

as spatially-modulated. Strikingly, even estimates by the same authors on data acquired from

the same subjects engaging the same task ranged between 26.5% (Wirtshafter and Wilson,

2020) to 56.0% (Wirtshafter and Wilson, 2019). When reported, LS place cells are typically

described as of lesser quality than classic hippocampal place cells, with lower within-session

stability and larger place fields (Takamura et al., 2006; Zhou et al., 1999). Adding to this

complexity, the LS is a large structure that is cytoarchitecturally subdivided into dorsal,

intermediate, and ventral subregions spread across its anterior-posterior axis (Risold and

Swanson, 1997; Swanson and Cowan, 1977), suggesting that, in addition to subjective

criterias, the striking differences in LS spatial coding characteristics between studies may be

in part a product of variations in recording locations within the region. In addition to

positional information, LS neurons may also encode idiothetic information, such as

acceleration, velocity, and direction (Takamura et al., 2006; Wirtshafter and Wilson, 2019;

Zhou et al., 1999).

Beyond their spatial tuning properties, hippocampal place cells are also noted for the

rapid reorganization of the place fields across time (on the order of days) and navigational

contexts, a phenomenon termed remapping (Gonzalez et al., 2019; Howard and Kahana,

2002; Mankin et al., 2015; Rubin et al., 2015; Ziv et al., 2013). If the LS spatial code is

directly inherited from its hippocampal inputs, one would expect comparable dynamics over

a similar timescale. In contrast, the LS has been implicated in behaviors that may require a

stable representation of space over time, including context-reward (Luo et al., 2011a;

McGlinchey and Aston-Jones, 2018), contextual fear conditioning (Besnard et al., 2019;

Butler et al., 2015; Garcia et al., 1997; Reis et al., 2010; Vetere et al., 2017) and spatial

learning and memory (Leutgeb and Mizumori, 1999; M'Harzi and Jarrard, 1992; Wirtshafter

and Wilson, 2020). Thus, whether and how the LS spatial code changes across time is crucial

to clarifying its relationship to the upstream hippocampal spatial code, with important

implications for its involvement in this broader spatial memory network and how it supports

behaviors across longer timescales.

To resolve these outstanding questions, we functionally characterize hippocampal

connectivity of LS, and asked how information rich LS neurons are in relation to their main

hippocampal inputs dorsal CA3 and CA1. To this end, we used head-mounted miniaturized

microscopes to record ~2000 LS neurons across the anterior-posterior and dorsal-ventral axes

of the LS, and compared their firing characteristics to pyramidal cells recorded in CA1 and

CA3, as animals navigated linear track and open field environments. Results demonstrate that

the LS accurately and robustly represents spatial, speed, and directional information in an

anatomically-organized fashion. While the information of LS neurons is generally

comparable to CA3 and CA1, our tracing studies suggest that LS information is sent to

downstream brain regions such as the medial septum and hypothalamus that are not directly

connected by the principal cells of the dorsal hippocampus.

Results

LS GABAergic neurons exhibit stable spatial activity during alternation in linear track

Previous electrophysiological studies of the lateral septum disagree on both the proportion of

place encoding cells in this region and their spatial content. Cells have been previously

described to be of lower quality than the classically described hippocampal place cells

(Takamura et al., 2006; Wirtshafter et al., 2019; 2020) or to be virtually absent (Tingley &

Buzsaki, 2018).

First, to assess the distribution of excitatory pyramidal inputs from each hippocampal

subregion to the GABAergic cells of the LS, we injected a monosynaptic retrograde rabies

tracer in dorsal LS (Figure 1A, Supplemental figure 1, n = 5 mice). We observed strong

labeling of pyramidal neurons of the hippocampus in area CA1 and CA3 (Figure 1B,

Supplemental figure 1). To investigate how the spatial coding properties of the LS compare

to these main dorsal hippocampal inputs, we expressed the calcium (Ca^{2+}) indicator

GCaMP6f in the GABAergic cells of the LS or in the pyramidal cells of the dorsal CA1

(dCA1) and dorsal CA3 (dCA3) of mice (Figure 1C). We recorded Ca^{2+} activity of 1082 LS

neurons (N = 15 mice), 1148 dCA1 pyramidal cells (N = 5 mice) and 144 cells from dCA3

(N = 2 mice) from animals alternating on a linear track. We subsequently filtered and

binarized the recorded calcium traces (Supplemental figure 2A, Etter et al., 2020) and

excluded periods of immobility ($< 5cm.s^{-1}$). Using an information-theoretic framework

(Shannon, 1948; Skaggs et al., 1993; Souza et al., 2018), we computed the mutual

information (MI) between calcium activity and spatial location for each cell and expressed

the results in bits (30 bootstrap samples, 50% data sampling, Supplemental figure 2B-C, Etter

et al., 2020). We found that LS GABAergic activity was significantly spatially modulated in

comparison to circularly shuffled surrogates ($p < .01$, Figure 1D, Supplemental figure 2C, H).

Indeed, both mutual information and within-session stability was higher for spatially

modulated LS cells than for non-modulated (Wilcoxon test, $p < .0001$, Supplemental figure

1E,F). Using this approach, a total of 43.90% of LS GABAergic cells qualified as place cells

(475/1082 cells), as compared to 31.05% in CA1 (336/1148 cells) and 32.63% in CA3 (47

cells/144 cells), with LS cells similarly distributed along the linear track as classical

hippocampal place cells (Figure 1E). In contrast to previous reports (Tingley & Buzsaki,

2018; Wirtshafter & Wilson, 2019, 2020), we thus observed a significantly higher proportion

of place cells in LS as compared to CA1 and CA3 (Kruskall-Wallis, $H(3) = 6.102$, $p = .0298$,

Figure 1F). We found that even though within-session stability of place cells was

significantly different between LS, dCA1, and dCA3, stability in the LS was not significantly

different than in dorsal CA1 (Kruskall-Wallis, $H(3) = 8.922$, $p = .0116$, Figure 1G).

In contrast to their omnidirectional firing properties in open field environments,

hippocampal neurons rapidly develop strong directional selectivity in linear tracks (Markus et

al., 1995; McNaughton et al., 1983; Navratilova et al., 2012). To test whether the LS place

code developed comparable directional selectivity, we ranked place cells as a function of the

difference between left and right MI (Supplemental figure 3A-D), we found LS cells tuned to

the absolute location regardless of direction (83/1082, 7.67%), while other cells were

selective to the left- (168/1082, 15.52%), or right-trajectories (239/1082, 22.08%).

Previous studies have used decoding methods to predict behavioral variables from

calcium imaging data recorded in the hippocampus (Gonzalez et al., 2019; Mau et al., 2018;

Ziv et al., 2013), yielding insights into the amount of information encoded by a neuronal

assembly. Here, we asked whether we could reliably estimate the mouse location solely from

LS neuronal activity patterns. To assess decoding accuracy and make sure our Bayesian

decoder worked well, we computed the mean decoding error and mean decoding accuracy

(Figure 1H-J, see methods) for CA1. Indeed, using a bootstrapped sample of 60 cells, a

decoder using CA1 calcium data significantly outperformed a decoder that was trained using

shuffled data (paired t-test, t(4) = 7.558, p = .0016, Figure 1K). Decoding location using LS

neuronal activity significantly outperformed a decoder that was trained using shuffled data

(paired t-test, t(6) = 13.56, p < .0001, Figure 1K), even when decreasing the number of

neurons used, for both decoding error (two-way RM ANOVA, F(3,19) = 571.4, p = 0.0024

for main effect of shuffle, F(3,19) = 6.948, p = 0.0024 for main effect of number of included

cells; Supplemental figure 4A) and decoding agreement (two-way RM ANOVA, F(3,19) =

1059, p < .0001 for main effect of shuffle, F(3,19) = 10.06, p = 0.0003 for main effect of

number of included cells; Supplemental figure 3A). A decoder trained using CA3 data was

not significantly different from shuffle (t(1) = 2.597, p = .2305), which is likely due to the

low number of animals included in the analysis. In order to compare decoding accuracy for

LS, CA1 and CA3, a decoding score was computed (Figure 1L, see methods), and we found

that LS significantly outperforms CA3 (one-way ANOVA, F(2,11) = 7.662, p = .0082).

Overall, place cells recorded in LS were significantly more active than those recorded in the

hippocampus (H(3) = 155.1, p < .0001, Supplemental figure 2G), which could contribute to

the overall higher decoding accuracy using LS cells.

LS GABAergic neurons exhibit stable spatial activity during free exploration in a 2D

environment

In order to assess spatial coding during free exploration in a two-dimensional environment,

we recorded Ca2+ activity of 1889 neurons in 28 mice implanted in distinct subregions of the

LS, while the animals were freely exploring a novel open field. Out of 1889 recorded cells,

38.01% (718/1889, N = 28) of LS cells displayed significant spatial information ($p < .01$;

Figure 2A-B, Supplemental figure 5), as compared to 31.33% (323/1031, N = 6) in CA1 and

42,15% (86/ 204, N = 5) CA3 cells. Despite the LS's known involvement in feeding-related

behaviors (Calderwood et al., 2020; Carus-Cadavieco et al., 2017; Sweeney and Yang, 2016;

Xu et al., 2019) , we did not observe any increased number of place cells with fields around

food zones, nor did we observe overrepresentation of objects or walls (Supplemental figure

6).

With theta-rhythmic cells in the LS having higher firing rates than theta-rhythm

independent cells (Pedemonte et al., 1998), and the suggestion that theta-rhythmic cells could

receive direct hippocampal inputs (though see Twery et al., 1990, 1992), we asked whether

LS cells with higher firing rates were also more likely to be spatially modulated. Indeed,

spatially-modulated cells were significantly more active than non-modulated cells (Mann-

Whitney test, U = 342219, $p < .0001$; Supplemental figure 5G-H) and displayed a higher

bursting index, defined as the probability of a cell being active, given it was already in an

active state, or p(a|a), than non spatially modulated cells (Mann-Whitney test, U = 36876, $p <$

.0001; Supplemental figure 5H).

We compared the place field properties of cells recorded in LS to those of the place

cells we recorded in CA1 and CA3 (Figure 2). For each animal, we computed the portion of

significantly positionally modulated cells per region, which was not significantly different

across LS, dCA1, and dCA3 (one-way ANOVA, $F_{(2,34)} = 0.499$, $p = .6110$, Figure 2B). As

expected on the basis or prior results (Leutgeb and Mizumori, 2002; Takamura et al., 2006;

Tingley and Buzsáki, 2018; Zhou et al., 1999), the within-session stability of LS cells was

significantly lower than that of place cells recorded in CA1 and CA3 (Kruskal-Wallis, $H(3) =$

49.44, $p < .0001$; Figure 2C). Conversely, the activity of LS neurons was significantly higher

than those of dCA1 and dCA3 principal cells (Kruskall-Wallis, $H(3) = 56.76$, $p < .0001$;

Figure 2D), and place fields were more dispersed (Kruskall-Wallis, H(3) = 93.21, $p < .0001$,

Figure 2E).

When decoding the location of the animals using a bootstrapped sample of 100 cells

from each region (Figure 2G), the decoder trained on data from LS performs significantly

better than a decoder trained on shuffled data (two-way RM ANOVA, F(1,9) = 76.15, $p <$

.0001 for main effect of shuffle), even when decreasing the number of neurons used to only

40 neurons, for both decoding error (two-way repeated-measures (RM) ANOVA, F(3,19) =

571.4, $p = 0.0024$ for main effect of shuffle, F(3,19) = 6.948, $p = 0.0024$ for main effect of

number of included cells; Supplemental figure 4B left) and decoding agreement (two-way

RM ANOVA, F(3,19) = 1059, $p < .0001$ for main effect of shuffle, F(3,19) = 10.06, $p =$

0.0003 for main effect of number of included cells; Supplemental figure 4B right). The same

held true for decoders trained using CA1 or CA3 data respectively (two-way RM ANOVA,

CA1: F(1,15) = 444.1, $p < .0001$ for main effect of shuffle, F(3,15) = 0.0706, $p = 0.97$ for

number of included cells; CA3: F(1,5) = 239.1, $p < .0001$ for main effect of shuffle, F(3,5) =

1.129, $p = 0.42$ for main effect of region, Supplemental figure 4B). Strikingly, a decoder

trained on data from LS does not perform significantly worse than a decoder trained on CA1

or CA3 data (F(2,9) = 0.3579, $p = 0.71$ for main effect of region; Figure 2H).

A subset of LS cells has stable spatial representations up to 8 days, similar to CA1

To test whether the LS and the hippocampus exhibit comparable evolutions of their

spatial code over time, we recorded from CA1 and LS neurons in a novel open field over both

short (3 days) and longer timescales (8 days, Figure 3A). We assessed the stability of the

spatial map across sessions, and we observed a subset of LS place cells that were stable over

time (Figure 3B). Correlating the tuning maps of aligned LS cell pairs over days lead to a

significant increase in correlation value over days (two-way ANOVA, F(2, 1470) = 8.129, p

= 0.0003, main effect of time) with LS cells being statistically more stable than shuffled

comparisons (F(2,1470) = 197.3, $p < .0001$, main effect of shuffle, Figure 3C). There was no

significant increase in the proportion of place cells per animal over days (one-way ANOVA,

F(3,16) = 1.618, $p = .2247$, Figure 3D). We observed the strongest within-session stability on

day 3 (Kruskal-Wallis, H(4) = 19.16, $p = 0.0003$; Figure 3E), as well as a steady increase in

spatial information (Kruskal-Wallis, H(4) = 13.27, $p = 0.0041$; Figure 3F). We computed the

mean stability for each place cell pair recorded over multiple days (Figure 3G), and when

comparing the mean tuning map correlation between day 1 and 8 for LS and CA1 place cells,

we found no significant difference in their stability (unpaired t-test, t(68) = 1.475, $p = 0.1449$,

Figure 3H). Together, this suggests that a subset of LS place cells codes for space over longer

periods of time, similarly to what has been described for CA1 (Gonzalez et al., 2019; Ziv et

al., 2013).

LS integrates direction, velocity, and acceleration information

In CA1, place cells have been found to carry directional (Acharya et al., 2016; Leutgeb et al.,

2000; Muller et al., 1994) and speed-related information (Czurko et al., 1999; McNaughton et

al., 1983; Iwase et al., 2020; Góis and Tort, 2018). Previous studies found that subsets of LS

neurons show some degree of modulation by direction of travel (Zhou et al., 1999) as well as

velocity and acceleration (Wirtshafter and Wilson, 2019) through correlative measures in

spatial alternation or T-maze navigation tasks, where the relationship between cell activity

patterns and movement-associated variables may be confounded with task-dependent

variables. Here, we again employed an information-theoretic approach to compute the MI

between calcium activity and each of these self-motion correlates in a subset of mice

recorded during a 15-minute free foraging task in the open field. We found that 28.13% of LS

cells are significantly modulated by direction (346/1230, N =19 mice, Figure 4A,

Supplemental figure 7), 18.70% by velocity (230/1230, Figure 4C) and 24.63% by

acceleration (303/1230, Figure 4E). We compared the proportion of LS cells significantly

modulated by each of these variables with the hippocampus, and observed no significant

differences in the proportion of velocity (Kruskall-Wallis, $H(3) = 0.255$, $p = 0.88$) or

acceleration (Kruskall-Wallis, $H(3) = 2.262$, $p = 0.32$) modulated cells. The proportion of

directionally modulated cells was significantly different across regions (Kruskall-Wallis,

$H(3) = 9.898$, $p = .0071$). We found a proportion of cells that were significantly tuned to

more than one of these variables in both LS and hippocampus (Figure 4G, Supplemental

figure 8), with 22.03 % of LS cells (271/1230) encoding more than one modality (Figure 4H).

Together, this conjunctive coding for location, directionality, and velocity parameters

indicates that LS cells fire in response to more complex navigational features, similarly to the

hippocampus.

To understand the function of this downstream copy of the hippocampal spatial code,

we assessed the downstream targets of hippocampal to LS projection. To this end, we

transfected an anterograde, transsynaptic Cre-expressing AAV1 viral vector (Zingg et al.,

2017) and assessed the second-order targets of the dorsal CA1-lateral septum (dCA1-LS) and

dorsal CA3-LS (dCA3-LS) projections (Figure 5). For either dCA1 or dCA3 injections, we

observed a direct disynaptic pathway from principal cells of the hippocampus to the medial

septum (Figure 5A-B) and the hypothalamic regions (Figure 5C). We observed that dCA1-LS

preferentially targets the LH, and dCA3-LS targets the hypothalamus more broadly as well as

the nuclei of the medial zone. In order to determine whether the projections observed at the

level of the LS with both our anterograde tracing approaches constituted passing fibers or

synaptic connections, we used an AAV-Flex-Synaptag (Beier et al., 2015), a Cre-dependent

viral construct to express mRuby in the cytosol and eGFP bound to synaptophysin, a protein

primarily expressed in the presynaptic endings of cells participating in synaptic transmission

(Figure 5D). We transfected cells at the level of the dorsal LS and observed significant

synaptophysin-bound projections in the medial septum (Figure 5E), confirming the existence

of direct, functional projections to medial septum neurons. Together, our work shows that LS

receives direct inputs from dorsal hippocampus and in turn projects to regions that are not

directly receiving inputs from the hippocampus itself (Figure 5F).

LS-cells coding for spatial and idiothetic information are non-uniformly distributed

To map out the monosynaptic connections between hippocampal and LS neurons, we

transfected an anterograde transsynaptic Cre-expressing AAV1 viral vector (Zingg et al.,

2017), in either the dCA1 or dCA3 (Supplemental figure 10A). We subsequently injected a

Cre-dependent eYFP expressing virus in the LS to visualize cells that receive direct

hippocampal input (Supplemental figure 10B). Importantly, this approach allowed us to

quantify the cells along the LS anterior-posterior and dorsal-ventral axis receiving direct

hippocampal inputs, and assess the density of their distribution within the anterior regions of

the LS (Bregma +1.0 to 0.0 anterior-posterior). Both dorsal dCA1 and dCA3 injections

resulted in the strongest transfection of LS cell bodies in dorsal regions compared to

intermediate and ventral LS (Supplemental figure 10D-F). Hence, the density of hippocampal

innervation to the LS is a gradient along the dorsal-ventral and anterior-posterior axis, with

dorsal-posterior LS neurons receiving more substantial dorsal hippocampal inputs.

Next, we sought whether LS place cells had spatial properties that were dependent on

the localization of their recording. To measure this, we classified the location of GRIN lens

implants along the dorsal-ventral and anterior-posterior axis (Figure 6A). Post-mortem

histological verification of GRIN lens implant combined with within-animal analysis of the

location of recorded cells enabled us to approximate the location of each recorded cell

(Figure 6B). While we observed no differences in cell activity characteristics, such as

bursting index or activity probability along the anterior-posterior axis, medial-lateral or

dorsal-ventral axis (Supplemental figure 11), we found a pronounced increase in the portion

of stable place cells (within-session stability > .5) along the anterior-posterior axis (linear

regression, $R^2 = 0.1272$, $p = 0.013$, the proportion of stable cells per 0.2 mm bin per animal, n

= 24 mice, Figure 6C). On the other hand, we observed a similar gradient along the dorsal-

ventral axis, with a larger proportion of stable place cells found at the more dorsal regions of

the LS, although this trend failed to reach significance (linear regression, $R^2 = 0.129$, $p =$

.0848, n = 24 mice, Figure 6C). Taken together, these data strongly suggest that regions of

the LS that receive stronger innervation from the dorsal hippocampus have a larger

proportion of cells that reliably encode space; thus it is likely that this information is directly

inherited from the hippocampus.

We also assessed the topographical distribution of self-motion tuned cells within the

LS. We observed a pronounced dorsal-ventral gradient in the distribution of stable (within-

session stability > .5) directionally modulated cells (linear regression, $R^2 = 0.3837$ $p =$

0.0061, n = 17 mice, Figure 6C), with most directionally modulated cells located at the more

dorsal regions of the LS, but we did not observe such a gradient along the anterior-posterior

axis (linear regression, $R^2 = 0.014$, $p = .3542$, n = 24 mice, Figure 6C). Velocity modulated

cells were mostly found at the ventral pole of the LS (linear regression, $R^2 = 0.1836$, $p =$

0.0367, n = 24), but no such gradient was found along the anterior-posterior axis (linear

regression, $R^2 = .0075$, $p = .4481$, $n = 24$). Acceleration-modulated cells were found evenly

distributed throughout the LS (DV: $R^2 = .0146$, $p = .2928$; AP: $R^2 = .00055$, $p = .7981$, n = 24

mice).

Discussion

While spatial coding has been extensively characterized in the hippocampal

formation, how downstream regions integrate this information has only recently begun to

receive attention. The LS has re-emerged as a critical region implicated in a space encoding

network, although electrophysiological recordings of the LS in rats have led to disparate

estimates of the quantity and quality of place cells in the region (Leutgeb and Mizumori,

2002; Wirtshafter and Wilson, 2019, 2020; Tingley and Buzsáki, 2018): LS place cells

recorded on a circular track were described to be almost absent (Tingley and Buzsáki, 2018),

but were found to be abundant in reward-seeking tasks (Wirtshafter and Wilson, 2019, 2020).

Using calcium imaging and unbiased information metric and decoding approaches, we found

that 38.01% of GABAergic LS cells robustly code for space during free exploration in an

open field and 45.29% of cells during linear track alternation. In addition to spatial

information, we found that the LS also reliably encodes velocity, acceleration, and directional

information, suggesting that the LS encodes more complex navigational features than

previously thought.

Neuronal representations in the hippocampus have been found to change over time,

with place code ensembles changing rapidly: an estimated ~40% (Gonzalez et al., 2019) to

~75-85% (Ziv et al., 2013) of CA1 neurons were found to remap over days. In the posterior

LS, we found a subpopulation of cells that display stable place fields over eight days,

comparable to CA1 cells recorded under the same conditions. The functional role of such

stability remains to be elucidated. One possibility is that this ensemble may mediate the

stable encoding of contexts over longer durations of time, which could account for the critical

importance of the LS for a number of spatial behaviors, including contextual fear

conditioning (Besnard et al., 2019; Butler et al., 2015; Garcia et al., 1997; Reis et al., 2010;

Vetere et al., 2017), context-reward associations (Luo et al., 2011b; McGlinchey and Aston-

Jones, 2018) and radial-maze navigation tasks (Leutgeb and Mizumori, 1999; M'Harzi and

Jarrard, 1992). Previous work has hinted at a dorsal to ventral organization of the LS place

code: electrophysiology studies recorded a slightly larger proportion of LS place cells in the

more dorsal regions (Takamura et al., 2006). A complementary anatomical microstructure has

been previously described for the rate-independent phase code in the LS, where the strength

of the phase code increased as a function of recording depth along the dorsal-ventral axis

(Buzsáki and Tingley, 2018). Here we find that lateral septal place cells are arranged along an

anterior-posterior gradient similar to the gradient of hippocampal inputs into the LS – a

finding that helps to reconcile the variation in previously reported estimates of LS place cell

prevalence, which ranged between 5.3% of cells recorded along the dorsal-ventral axis

(Tingley and Buzsáki, 2018) up to 56% recorded at the most dorsal level of the LS

(Wirtshafter and Wilson, 2019). This pattern suggests that the inhibitory neurons of the LS

may inherit a place code from the hippocampus, with subregions receiving dorsal

hippocampal inputs being most similar to the classical hippocampal place cells. Despite this

interconnectedness with the LH and previous reports of LS place cells being skewed towards

reward in a spatial navigation task (Wirtshafter and Wilson, 2020), we did not observe any

increased LS place cell overrepresentation around food zones during free exploration.

In addition to place cells, we observed significant proportions of velocity and

acceleration modulated cells in the lateral septum. These cells have been previously described

by others using correlative measures, although estimated proportions vary widely: for

velocity tuning in reward-seeking tasks, estimates range from almost absent (Buzsáki and

Tingley, 2018) to almost 60% of cells (Wirtshafter and Wilson, 2019). For acceleration,

approximately 45% of LS cells were found to show some degree of correlation, with almost

30% of recorded cells correlated with both speed and acceleration (Wirtshafter and Wilson,

2019). Here, using information metrics and a free exploration task in which firing

characteristics are not confounded with task parameters, we find that 18.70% of cells were

modulated by velocity and 24.63% by acceleration. Cells reliably coding for both velocity

and acceleration were found to be relatively sparse (2%). Velocity and acceleration cells were

distributed dissimilarly in the LS, with velocity cells being more abundant towards the ventral

pole, whereas acceleration modulated cells were found to be distributed equally. This raises

the question whether velocity and acceleration information are inherited from different areas

upstream of the LS. Interestingly, cells coding for direction were more abundant towards the

dorsal portion of the LS, but distributed equally along the anterior-posterior axis. Previous

studies have identified classical head direction cells in multiple, well-described circuits,

including the postsubiculum (Taube et al., 1990), anterodorsal thalamic nucleus (Taube,

1995), mammillary bodies (Stackman and Taube, 1998), and entorhinal cortex (Sargolini et

al., 2006). A probable source of directional information to the LS is the entorhinal cortex

(Allen Brain Connectivity Atlas, Oh et al., 2014). Our work supports the perspective that LS

neurons combine a wide range of modalities and may form a complex representation of

context over longer timescales than previously reported. In addition to space, velocity, and

reward information, hippocampal pyramidal cells have been shown to encode variables such

as time (Eichenbaum, 2014; Pastalkova et al., 2008), odor (Deshmukh and Bhalla, 2003;

MacDonald et al., 2013) and sound frequency (Aronov et al., 2017). Whether this information

is also relayed to the LS remains to be elucidated.

The question arises of what the function of could be having a region downstream of

the hippocampus that expresses such seemingly similar coding characteristics. Previous work

estimated the convergence of hippocampal efferents onto LS neurons to be 20 to 800 times

more dense than to any of its cortical targets (Tingley and Buzsáki, 2018). Understanding

how individual LS neurons integrate the thousands of synaptic inputs they receive from these

hippocampal pyramidal neurons will be critical to understanding how the hippocampal map is

processed downstream. Through the process of synaptic integration, a receiving neuron can

fire an action potential upon receiving sufficient synchronized excitatory input on its

dendrites, leading to rapid depolarization of the cell. The activation of a LS neuron could thus

require multiple hippocampal pyramidal cells to spike simultaneously or in close temporal

proximity. Hippocampal pyramidal cells will fire concomitantly when their place fields are in

overlapping regions or in close vicinity of another. Due to the high interconnectivity of LS

GABAergic neurons (Alonso and Frotscher, 1989; Staiger and Nürnberger, 1989), the

activation of one LS cell could subsequently lead to the inhibition of other neurons in the

nucleus (Leroy et al., 2018; Phelan et al., 1989), thereby reducing noise. Thus, through this

process of coincidence detection and recurrent inhibition, the spatial map could be

compressed to be represented by a much lower number of cells. As such, the LS could

accurately convey hippocampal information to downstream regions using a much sparser

signal, thereby allowing for increased metabolic efficiency (Olshausen and Field, 2004) and

more effective associative learning (Isely et al., 2010; Palm and Sommer, 2006; Zetsche).

Indeed, our tracing work shows LS cells receiving hippocampal inputs from principal

cells in turn project directly to the LH and the MS. Although the existence of an LS-MS

projection has long been debated (Leranth et al., 1992; Staiger and Nürnberger, 1991;

Swanson and Cowan, 1977, 1979; Witter et al., 1992), we observed labeling at the level of

the MS after transfection of dCA1-input and dCA3-input receiving LS cells providing strong

support that principle cells of the hippocampus send indirect projections to the MS through

LS neurons. This circuit is complementary to the well-described projections of hippocampal

GABAergic interneurons directly to the MS which may play a key role in theta rhythm

generation (Toth et al., 1993; Manseau et al., 2008). The MS plays a role in generating and

propagating theta rhythms throughout the hippocampal formation (Leão et al., 2015; Patel et

al., 2013; Robinson et al., 2016), which in turn organizes hippocampal place cell activity

(Buzsáki, 2002), as well as a behavioral role in the initiation and velocity of locomotion

(Fuhrmann et al., 2015; Justus et al., 2017). It may be that the information on locomotion-

related information reported in specific MS neurons (Justus et al., 2017) may originate from

the LS. Moreover, we observed a direct projection from the LS to the hypothalamus, a highly

connected region traditionally known as necessary for the regulation of feeding (Stuber and

Wise, 2016), arousal and motivation (Burdakov and Peleg-Raibstein, 2020), and, more

recently learning and memory (Aitta-Aho et al., 2016; Izawa et al., 2019; Kosse and

Burdakov, 2019; Petrovich, 2018). The LH is crucial for the control of locomotion

(Gladfelter and Brobeck, 1962; Grillner et al., 2008; Kosse et al., 2017; Sinnamon, 1993) and

thought to mediate motivational and goal-directed processes underlying feeding (Carus-

Cadavieco et al., 2017; Mahler et al., 2014; Mogenson et al., 1980; Stuber and Wise, 2016).

In summary, we have shown that the GABAergic cells of the LS may provide the

hypothalamus and medial septum with information about location, direction, and speed, and

therefore constitute a core node within a distributed spatial memory network. Within this

network, the LS may be necessary for the translation of spatial information to goal-directed

or environmentally appropriate actions necessary for survival.

Methods

Animals

All procedures were approved by the McGill University and Douglas Hospital Research

Centre Animal Use and Care Committee and in accordance with the Canadian Council on

Animal Care. Naive mice were housed individually on a 12-hour light/dark cycle at 22°C and

40% humidity with food and water ad libitum. All experiments were carried out during the

light portion of the light/dark cycle. Both male and female mice were used for this study.

Virus injections

Mice, 8-16 weeks old, were anesthetized with isoflurane (5% induction, 0.5-2% maintenance) and placed in a stereotaxic frame (David Kopf Instruments). Body temperature was maintained with a heating pad, and eyes were hydrated with gel (Optixcare). Carprofen (10 ml/kg) was administered subcutaneously. All viral injections were performed using a glass pipette connected to a Nanoject III (Drummond) injector at a flow rate of 1 nL/second. All stereotaxic coordinates are taken from Bregma (in mm). After surgery, animals were continuously monitored until recovery. For three days after surgery, mice were provided with a soft diet supplemented with Carprofen for pain management (MediGel CPF, ~5mg/kg/day).

Virus-mediated expression of genetically encoded calcium indicator

Adeno-associated virus (AAV) of serotype 9 containing the genetically encoded calcium indicator (GECI) 6 fast (GCaMP6f) under a flex promoter was obtained from the University of Pennsylvania Vector Core (AAV2/9.Syn.Flex.GCaMP6f.WPRE.SV40, CS0641, Penn Vector Core via Addgene). Lateral septum viral injections were targeted at the following coordinates: Anterior LS, AP: 0.86; ML: 0.35; DV: -3.0; Intermediate LS, AP: 0.38; ML: 0.35; DV: -2.7, Posterior LS, AP: 0.10; ML: 0.35; DV: -2.5, with DV + or - .2 to adjust for more dorsal or ventral placement of injection. Lateral septum injections were done in VGAT-IRES-Cre transgenic mice (Slc32a1$^{tm2(cre)Lowl}$/J, JAX stock #016962). For hippocampal principal cell imaging, C57Bl/6J mice (JAX stock #000664) were injected with a viral vector specifically targeting GCaMP6f expression to CaMKII-positive cells (AAV5.CamKII.GCaMP6f.WPRE.SV40, Penn Vector Corde via Addgene, CS1024). GCaMP6f injections were targeted at the following coordinates: CA1, AP: -1.8; ML: 1.5; DV: -1.5; CA3, AP: -2.1; ML: 2.3; DV: -2.2.

Retrograde tracing

Retrograde tracing studies were done using a Cre-dependent retrograde Rabies tracing

approach. Briefly, 100 nL of helper virus (AAVdj.hSyn.Flex.TVA.P2A.eGFP.2A.oG,

NeuroPhotonics Center, Laval, Canada; Callaway, 2008; Wickersham et al., 2007) was

injected in the LSd (coordinates as above) of VGAT-IRES-Cre transgenic mice (n = 5

animals, Slc32a1$^{tm2(cre)Lowl}$/J, JAX stock #016962). Cre specificity was tested by the

NeuroPhotonics center. After 4 weeks of incubation time to allow for the complete

expression of the helper virus, 200 nL of inactivated EnvA pseudotyped rabies virus coupled

to mCherry was injected (RABV-EnvA-deltaG-mCherry, NeuroPhotonics Center) at the

same coordinates as the helper virus injection, and brains were collected 7 days later.

Anterograde transsynaptic tracing

To distinguish synaptic targets from passing projection fibers of dCA1 and dCA3 within the

LS, C57Bl/6J mice (JAX stock #000664) received a bilateral injection of transsynaptic Cre-

inducing anterograde virus (AAV1-hSyn-Cre-WPRE-hGH, Penn Vector Core via Addgene,

V24784) mixed with a Cre-dependent tdTom-expressing reporter virus to visualize injection

sites (AAV2-Ef1a-flex-tdTomato, NeuroPhotonics Center, AAV602) in a 3:1 ratio, 80 nL

total injection volume per hemisphere (n = 5 mice per group). Injections were targeted at the

following coordinates: CA1: AP: -1.86; ML: 1.5; DV: -1.5; CA3: AP: -2.1; ML: 2.3; DV: -

2.2. Following 1 week of incubation time, animals received a bilateral injection of a Cre-

dependent anterograde vector in LS expressing enhanced yellow fluorescent protein (400 nL,

bilateral, AP: 0.5; ML: 0.35; DV: -2.6, AAVdj-DIO-Cheta-eYFP, NeuroPhotonics Center,

AAV938). Brains were collected and processed three weeks post-injection.

Synaptag tracing

To verify presence of synaptic connections at the level of the LS-MS projection, AAVdj-

hSyn-flex-mRuby2-syp-eGFP construct (lot AAV799, NeuroPhotonics Center; Oh et al.,

2014) was injected in the LS of CaMKIIα-Cre mice (B6.Cg-Tg(Camk2a-cre)T29-1Stl/J, JAX

stock #005359), with 100 nL total injection volume injected at 1 nL/second. After 4 weeks of

incubation time, animals were perfused, and brains were processed as described below.

GRIN lens implant and baseplate

Two to four weeks post GCaMP6f injection, a 0.5 mm diameter gradient refractive index

(GRIN, Inscopix) lens was implanted above LS, dorsal CA1 or CA3, or a 1.8 mm GRIN

(Edmund Optics) lens was implanted above dorsal CA1. For .5 mm lens implant, the lens was

glued to an aluminium baseplate prior to the surgery, which provided two main advantages:

1) it allowed us to implant the baseplate/lens assembly in one surgery (instead of separate

surgeries) and 2) during the surgery the miniscope (V3, miniscope.org) was used to image for

fluorescence while lowering the lens. The mouse was anesthetized with isoflurane, and the

skull was cleared. A ~.6 mm diameter hole was drilled in the skull at the level of the injection

site. An anchor screw was placed above the contralateral cerebellum to stabilize the implant.

The baseplate and GRIN combination were attached to the miniscope and together secured to

the stereotaxic frame. After making a leading track using a .5 mm diameter needle, the

implant was lowered in place. While lowering the GRIN lens into the tissue, and an increase

in fluorescent signal indicates that the injection site has been reached. Both the GRIN lens

and baseplate were permanently secured using C&B-Metabond (Patterson Dental). A plastic

cap was used to protect the GRIN lens from scratching during the recovery period and in

between recording sessions. The animal was given 4 - 6 weeks of recovery time before

experiments started. LS implants were targeted at the following coordinates: Anterior LS,

AP: 0.86; ML: 0.35; DV: -2.9; Intermediate LS, AP: 0.38; ML: 0.35; DV: -2.65; Posterior

LS, AP: 0.10; ML: 0.35; DV: -2.45; with DV + or - .2 to adjust for dorsal, ventral or

intermediate targets; CA3 implants were targeted at AP: -2.1; ML: 2.2; DV: -2.2.

For 1.8 GRIN lens implantation, a ~2 mm diameter cranial window was prepared.

Again, an anchor screw was used to secure the implant in place. After removing the dura, a

portion of the cortex above the injection site was gently aspirated using a vacuum pump

without applying pressure on the underlying hippocampal tissue. The 1.8 mm GRIN lens was

lowered at the following coordinates: AP: -1.8; ML: 1.5; DV: -1.8. The GRIN lens was

permanently attached to the skull using Metabond, and Kwik-Sil (World Precision

Instruments) silicone adhesive was placed on the GRIN to protect it. On average 4 weeks

post-implant, the silicone cap was removed and CA1 was imaged using a miniscope mounted

with an aluminium base plate while the mouse was under light anesthesia (~0.5 % isoflurane)

to allow the visualization of cell activity. When a satisfying field of view was found (large

number of neurons, visible landmarks), the baseplate was cemented above the GRIN lens and

a protective cap was used to protect the GRIN from scratches. Imaging would start

approximately 1 week after baseplating.

Miniscope recordings

The animals were habituated by being gently handled for ~5 min per day for 3 days. Animals

were then water or chow-scheduled (2h access per day). Miniscope recordings performed at

30 Hz for 15 minutes every day, with just one recording session per day to minimize

photobleaching. For open field recordings, animals were freely foraging for randomly placed

10% sucrose or 10% sweetened condensed milk in water rewards.

Recording environments consisted of a 45 x 45 cm dark grey open field with visual

cues, or a 49 x 49 cm white plexiglass open field, which was placed in different recording

chambers. For linear track recordings, rewards (30 ul of 10% sucrose water) were placed at

each end of the 100 cm linear track, or a 130 cm linear track, and the mouse had to consume

one reward before getting the next one delivered.

Miniscope and behavior video acquisition

Miniscopes (V3, miniscope.org) were assembled using open-source plans as described

previously (miniscope.org, Cai et al., 2016; Ghosh et al., 2011). Imaging data were acquired

using a CMOS imaging sensor (Aptina, MT9V032) and multiplexed through a lightweight

coaxial cable. Data was acquired using a data acquisition (DAQ) box connected via a USB

host controller (Cypress, CYUSB3013). Animal behavior was recorded using a webcam

mounted above the environment. Calcium and behavioral data were recorded using

Miniscope custom acquisition software (miniscope.org). The DAQ simultaneously acquired

behavioral and cellular imaging streams at 30 Hz as uncompressed .avi files (1000 frames per

file), and all recorded frames were timestamped in order to perform the subsequent

alignment.

Histology

After completion of all experiments, animals were deeply anesthetized with

ketamine/xylazine/acepromazide (100, 16, 3 mg/kg, respectively, intraperitoneal injection).

Mice were then transcardially perfused with 4% paraformaldehyde in PBS (PFA). Brains

were extracted and postfixed in PFA at 4 °C for a minimum of 48 hours. Brains were

sectioned at 50 μm using a vibratome and cryoprotected in a solution of 30% ethylene glycol,

30% glycerol and 40% PBS until used.

Immunohistochemistry

Sections were washed 3 x 20 min in PGT (0.45% Gelatin and 0.25% Triton in 1x PBS) at

room temperature. Next, primary antibodies (1:1000 goat anti-GFP from Novus Biologicals

or 1:10 000 rabbit anti-RFP from VWR (Rockland) were incubated with PGT overnight at

4°C. Following 10, 20, and 40 min washes in PGT, sections were incubated with secondary

antibodies (1:1000 donkey anti-goat coupled to A488 or 1:1000 donkey anti-rabbit coupled to

A555, both from Molecular Probes) in PGT at room temperature for 1 hour. Sections were

subsequently washed for 10 and 20 minutes in PGT and 30 minutes in PBS. Sections were

then mounted on glass slides and permanently coverslipped with Fluoromount that contained

DAPI.

Calcium imaging analysis

Calcium imaging videos were analyzed using the Miniscope Analysis pipeline

(https://github.com/etterguillaume/MiniscopeAnalysis) as described previously (Etter et al.,

2020; Pnevmatikakis and Giovannucci, 2017). Non-rigid motion correction was applied using

NoRMCorre (Pnevmatikakis and Giovannucci, 2017), and videos were spatially

downsampled (3x) before concatenation. Calcium traces were extracted using CNMFe (Zhou

et al., 2017) using the following parameters: gSig = 3 pixels (width of gaussian kernel), gSiz

= 20 pixels (approximate neuron diameter), background_model = 'ring', spatial_algorithm =

'hals', min_corr = 0.8 (minimum pixel correlation threshold), min_PNR = 8 (minimum peak-

to-noise ratio threshold). After extraction, cells and traces were visually inspected, and cells

with low signal to noise ratio were removed. Raw calcium traces were filtered to remove

high-frequency fluctuations and binarized (normalized amplitude > 2 sd and the first-order

derivative > 0, Etter et al., 2020).

Position data were inferred offline following the recording, using the red LED on the

CMOS of the miniscope. A subset of behavioral videos were analyzed using DeepLabcut

through Python Integrated Miniscope Pipeline Notebook (PIMPN, available through

https://github.com/etterguillaume/PIMPN) to extract head direction.

Activity and bursting index

Following the binarization of raw calcium traces, we compute the probability of a neuron to

be active P(A) using the following formula:

$$P(A) = \frac{time\ active}{total\ time}$$

The bursting index was computed as the probability of a cell being active, given it was

already in an active state, or P(A|A). The activity index is computed as the probability of a

cell becoming active when it is in an inactive state, or P(A|I).

Spatial modulation

Space was binned (Position in open field: 17 x 17 bins of 3 cm; Position in linear track: 34

bins of 3 cm; velocity, min = 2.5 cm/s, max = 30 cm/s, 20 bins; acceleration: min = -2 cm/s^2,

max = +2 cm/s^2, 20 bins; head direction: 40 bins of 9°). Velocity was smoothed over thirty

frames to reduce discontinuities. For head direction and positional information, all frames

where velocity was < 5cm/s were excluded. The probability of being active P(A) informs on

the activity rate of a neuron. Next, we compute the probability of spending time in a given

bin *i* (spatial, velocity, acceleration or head direction bin).

$$P(S_i) = \frac{time\ in\ bin\ i}{total\ time}$$

616   Next, we compute the joint probability:

$$P(S_i \cap A) = \frac{time\ active\ while\ in\ bin\ i}{total\ time}$$

617   And the probability that a cell is active given the animal is in a bin:

$$P(A \mid S_i) = \frac{time\ active\ while\ in\ bin\ i}{time\ in\ state}$$

618   After plotting the binarized activity in space, we perform n = 1000 random circular

619 permutations. Circular permutations are used to remove the temporal relationship between

620 neuronal activity and behavior, but it preserves the temporal structure of calcium transients

621 and therefore leads to more conservative results (as opposed to complete randomization of

622 every data point, which inflates the significance value of results). In this case, the p-value

623 corresponds to the number of datapoints from the shuffled distribution that are greater than

624 the actual data for each bin, divided by the number of permutations (Cohen, 2014). A

625 threshold of $p < .01$ is used to determine the significant data points that make up a significant

626 place field. In our case, for a cell to be considered a place cell, at least one of the spatial bins

627 must contain activity that is significantly non-random. To compute within-session stability,

628 the halfway point in the recording was determined by taking the timestamps where half of the

629 total number of binarized events for the session was reached. Tuning maps were computed

630 for both the first and second half of the session using the methodology described above. A

631 Gaussian smoothing filter ($\sigma = 3$) was used to smooth the tuning maps before computing a

632 correlation between the two. A within-session correlation value of $>.5$ was considered a

633 stable field.

635 **Calculating mutual information content**

636 Mutual information is used to describe the amount of information about one variable (spatial

637 location, head direction, velocity or acceleration) through the observation of neuronal

638 activity, and was calculated using the following formula:

639     $$MI = \sum_{i=1}^{M} \sum_{j=1}^{2} P(S_i \cap A_j) \times log2(\frac{P(S_i \cap A_j)}{P(S_i) \times P(A_j)})$$

Where *M* is the total number of possible behavioral states, $P(S_i \cap A_j)$ is the joint probability of

the animal being in bin *i* concurrently with activity level *j*. As we are using binarized activity,

j can only be active or inactive. MI was calculated using 30 bootstrapped surrogates using 50

percent of the data (randomly chosen data points) with replacement, to allow for computation

of mean MI and SEM. Next, the trace was circularly shuffled in order to calculate a shuffled

MI and SEM. Cells were then sorted along with the magnitude of the difference between

mean MI and mean shuffled MI, and a two-way ANOVA with a $p < .01$ threshold was used

to determine whether the difference between the two numbers was significant in order for the

cell to be deemed a significant place cell. In case the assumptions for parametric ANOVA are

not met, the significance value was computed by taking the sum of the shuffled MI values of

greater magnitude than actual MI values, divided by the number of shuffles.

Bayesian decoding

A Bayesian decoder was used to evaluate how well LS neural activity estimated the animal's

location. We used decoding methods specifically adapted for calcium imaging data with

binarized calcium transients (Etter et al., 2020). The training dataset was generated using

50% of the data, randomly chosen, using 30 bootstrap samples including 40, 60, 80 or 100

randomly chosen cells (Supplemental figure 4). The posterior probability density function,

which is the probability that an animal is in a particular bin S_i, given the neural activity A is

calculated using the following equation:

$$P(S_i \mid A) = \frac{P\left(A \mid S_i\right) \times P\left(S_i\right)}{P\left(A\right)}$$

660   P(S|A) is the posterior probability distribution of states given neuronal activity. No prior

661   assumption is made about the location of the mouse on the linear track, P(S) is kept uniform

662   to attribute equal probability for each location. Every neuron is assumed to be independent of

663　each other. To construct the tuning curves from multiple neurons, we use the following

664　equation:

$$P\,(S\,|\,A)\;=\;\prod_{k\,=\,1}^{N}\frac{P(A_k\,|\,S)\;\times\;P\,(S)}{P\,(A_k)}$$

665　With *P(S/A)* a vector of *a posteriori* behavioral states and *N* corresponding to the number of

666　neurons used. For every neuron *k* the tuning curves are constructed, and corresponding

667　posterior location probability can be derived.

668　　　　To reconstruct the position of the mouse, we will consider the location associated with

669　the maximum a posteriori:

$$\hat{y}=\;arg\,max\,exp\left[\sum_{k\,=1}^{N}\;log\left(1+\frac{P(A_k\,|\,S)\;\times\;P(S)}{P(A_k)}\right)\;-\;1\right]$$

670　With the estimated state among all possible states S. We used a temporal filtering window in

671　the open field of 1.5 seconds and in the linear track of 0.5 seconds to remove erratic jumps in

672　the decoded position of the mouse. Decoding accuracy was measured as 1) decoding error

673　and 2) decoding agreement. In one-dimensional space, decoding error was assessed as

$$decoding\;error\;=\;|\,decoded\;position\,-\,actual\;position\,|$$

674　In two-dimensional space, decoding error was assessed as

$$decoding\;error\;=\;\left(\sqrt{actual\;position\,-\,decoded\;position}\right)^2$$

676　Decoding score was computed as

$$decoding\;score\;=\;\frac{mean\;(shuffled\;decoding\;error)\,-\,mean\;(decoding\;error)}{length\;of\;track}$$

677　Decoding agreement was defined as the portion of time where the exact location or head

678　direction bin was successfully decoded:

$$decoding\;agreement\;=\;\frac{time\;points\;successfully\;decoded}{total\;time}$$

680 **Tracking cells across sessions**

681 Neurons were tracked over multiple days using a probabilistic method as previously

682 described (Sheintuch et al., 2017). Briefly, spatial footprints were aligned using non-rigid

683 alignment (transformation smoothness = 2) to correct for displacements of the brain or

684 miniscope placement. After alignment, we considered candidate sets of cells to be the same

685 neuron if their maximal distance was < 12 um, and the spatial correlation between sets of

686 cells was > .65. Next, we visually identified candidate cell pairs across sessions and manually

687 deleted those cell pairs that were erroneously selected (< 1% of cases). We assessed the

688 stability of the spatial representation in LS using simple field correlation, which was assessed

689 by correlating smoothed place fields of cell pairs. To generate the null hypothesis for place

690 fields' displacements between pairs of days, we used the place fields tuning maps but shuffled

691 cells identities over days (30 shuffles).

692

693 **Materials availability**

694 This study did not generate new unique reagents.

695

696 **Data and Code availability**

697 All data will be made publicly available. Code for cell extraction is available at

698 https://github.com/etterguillaume/MiniscopeAnalysis. Code for behavior extraction is

699 available at https://github.com/etterguillaume/PIMPN. Code for extraction of tuning curves

700 and decoding were adapted from Etter et al. (2020) and can be downloaded at the following

701 address: https://github.com/etterguillaume/CaImDecoding. All other code is available from

702 the corresponding authors upon request.

703

704 **Statistics**

Statistics Statistical analyses were performed using Matlab (Mathworks) and GraphPad Prism

version 6.00 (GraphPad Software, La Jolla, California USA). All data are presented as mean

± standard error of the mean (SEM) and statistical test details are described in the

corresponding results. All t-tests are two-tailed. Normality distribution of each group was

assessed using the Shapiro-Wilk normality test and parametric tests were used only when

distributions were found normal, using Student's t-tests, one-way ANOVA or two-way

ANOVA. For non-normal distributions, non-parametric tests Mann-Whitney or Kruskal-

Wallis tests were performed and described where applicable. Significant main effects or

interactions were followed up with appropriate post hoc testing using Bonferroni corrections

were applicable. $p < 0.05$ was considered statistically significant, or $p < .01$ and described

where applicable *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$, ****, $p < 0.0001$.

Funding

This work was supported by funding from the Canadian Institutes for Health Research

(CIHR), Fonds de la Recherche en Santé du Québec (FRSQ), and the Natural Sciences and

Engineering Research Council of Canada (NSERC). SVDV is supported by a Vanier Canada

Graduate Scholarship and the Richard H. Tomlinson Doctoral Fellowship.

Acknowledgments

We thank Alexandra T. Keinath for valuable comments on a previous version of the

manuscript. We thank Ke Cui for help with colony maintenance and mice perfusions, and

members of the Williams lab for constructive feedback throughout the work. We also thank

Daniel Aharoni (UCLA) for guidance in using the UCLA miniscope and Bruno Rivard for

building the miniscopes. The present study used the services of the Molecular and Cellular

Microscopy Platform at the DHRC.

Competing interests

The authors declare no competing interests.

References

Aitta-Aho, T., Pappa, E., Burdakov, D., and Apergis-Schoute, J. (2016). Cellular activation of
    hypothalamic hypocretin/orexin neurons facilitates short-term spatial memory in
    mice. Neurobiol. Learn. Mem. *136*, 183–188.

Alonso, J.R., and Frotscher, M. (1989). Organization of the septal region in the rat brain: a
    Golgi EM study of lateral septal neurons. J. Comp. Neurol. *286*, 472–487.

Anthony, T.E., Dee, N., Bernard, A., Lerchner, W., Heintz, N., and Anderson, D.J. (2014).
    Control of stress-induced persistent anxiety by an extra-amygdala septohypothalamic
    circuit. Cell *156*, 522–536.

Aronov, D., Nevers, R., and Tank, D.W. (2017). Mapping of a non-spatial dimension by the
    hippocampal-entorhinal circuit. Nature *543*, 719–722.

Beier, K.T., Steinberg, E.E., Deloach, K.E., Kremer, E.J., Malenka, R.C., Luo, L., Beier,
    K.T., Steinberg, E.E., Deloach, K.E., Xie, S., et al. (2015). Circuit Architecture of
    VTA Dopamine Neurons Revealed by Systematic Input-Output Mapping Article
    Circuit Architecture of VTA Dopamine Neurons Revealed by Systematic Input-
    Output Mapping. Cell *162*, 622–634.

Bender, F., Gorbati, M., Cadavieco, M.C., Denisova, N., Gao, X., Holman, C., Korotkova, T.,
    and Ponomarenko, A. (2015). Theta oscillations regulate the speed of locomotion via
    a hippocampus to lateral septum pathway. Nat. Commun. *6*, 8521.

Besnard, A., Gao, Y., Kim, M.T., Twarkowski, H., Reed, A.K., Langberg, T., Feng, W., Xu,
    X., Saur, D., Zweifel, L.S., et al. (2019). Dorsolateral septum somatostatin

755   interneurons gate mobility to calibrate context-specific behavioral responses. Nat.

756   Neurosci. *22*, 436–446.

757   Bredewold, R., Schiavo, J.K., van der Hart, M., Verreij, M., and Veenema, A.H. (2015).

758   Dynamic changes in extracellular release of GABA and glutamate in the lateral

759   septum during social play behavior in juvenile rats: Implications for sex-specific

760   regulation of social play behavior. Neuroscience *307*, 117–127.

761   Brito, G.N.O., and Thomas, G.J. (1981). T-maze alternation, response patterning, and septo-

762   hippocampal circuitry in rats. Behav. Brain Res. *3*, 319–340.

763   Burdakov, D., and Peleg-Raibstein, D. (2020). The hypothalamus as a primary coordinator of

764   memory updating. Physiol. Behav. *223*, 112988.

765   Butler, C.W., Wilson, Y.M., Gunnersen, J.M., and Murphy, M. (2015). Tracking the fear

766   memory engram: Discrete populations of neurons within amygdala, hypothalamus,

767   and lateral septum are specifically activated by auditory fear conditioning. Learn.

768   Mem. *22*, 370–384.

769   Buzsáki, G. (2002). Theta oscillations in the hippocampus. Neuron *33*, 325–340.

770   Buzsáki, G., and Tingley, D. (2018). Space and Time: The Hippocampus as a Sequence

771   Generator. Trends Cogn. Sci.

772   Cai, D.J., Aharoni, D., Shuman, T., Shobe, J., Biane, J., Song, W., Wei, B., Veshkini, M., La-

773   Vu, M., Lou, J., et al. (2016). A shared neural ensemble links distinct contextual

774   memories encoded close in time. Nature *534*, 115–118.

775   Calderwood, M.T., Tseng, A., and Glenn Stanley, B. (2020). Lateral septum mu opioid

776   receptors in stimulation of feeding. Brain Res. *1734*, 146648.

777   Callaway, E.M. (2008). Transneuronal circuit tracing with neurotropic viruses. Curr. Opin.

778   Neurobiol. *18*, 617–623.

779   Carus-Cadavieco, M., Gorbati, M., Ye, L., Bender, F., van der Veldt, S., Kosse, C., Börgers,

C., Lee, S.Y., Ramakrishnan, C., Hu, Y., et al. (2017). Gamma oscillations organize
top-down signalling to hypothalamus and enable food seeking. Nature *542*, 232–236.

Chee, S.-S.A., Menard, J.L., and Dringenberg, H.C. (2015). The lateral septum as a regulator
of hippocampal theta oscillations and defensive behavior in rats. J. Neurophysiol. *113*,
1831–1841.

Clemens, A.M., Wang, H., and Brecht, M. (2020). The lateral septum mediates kinship
behavior in the rat. Nat. Commun. *11*, 1–11.

Cohen, M.X. (2014). Analyzing Neural Time Series Data: Theory And Practice. (Cambridge,
MA: The MIT Press).

Deshmukh, S.S., and Bhalla, U.S. (2003). Representation of odor habituation and timing in
the hippocampus. J. Neurosci. *23*, 1903–1915.

Eichenbaum, H. (2014). Time cells in the hippocampus: A new dimension for mapping
memories. Nat. Rev. Neurosci. *15*, 732–744.

Etter, G., Manseau, F., and Williams, S. (2020). A Probabilistic Framework for Decoding
Behavior From in vivo Calcium Imaging Data. Front. Neural Circuits *14*, 1–16.

Fuhrmann, F., Justus, D., Sosulina, L., Kaneko, H., Beutel, T., Friedrichs, D., Schoch, S.,
Schwarz, M., Fuhrmann, M., and Remy, S. (2015). Locomotion, Theta Oscillations,
and the Speed-Correlated Firing of Hippocampal Neurons Are Controlled by a Medial
Septal Glutamatergic Circuit. Neuron *86*, 1253–1264.

Garcia, R., Vouimba, R.M., and Jaffard, R. (1997). Contextual conditioned fear blocks the
induction but not the maintenance of lateral septal LTP in behaving mice. J.
Neurophysiol. *78*, 76–81.

Ghosh, K.K., Burns, L.D., Cocker, E.D., Nimmerjahn, A., Ziv, Y., Gamal, A. El, and
Schnitzer, M.J. (2011). Miniaturized integration of a fluorescence microscope. Nat.
Methods *8*, 871–878.

Gladfelter, W.E., and Brobeck, J.R. (1962). Decreased spontaneous locomotor activity in the

rat induced by hypothalamic lesions. Am. J. Physiol. *203*, 811–817.

Gonzalez, W.G., Zhang, H., Harutyunyan, A., and Lois, C. (2019). Persistence of neuronal

representations through time and damage in the hippocampus. Science (80-.). *365*,

821–825.

Grillner, S., Wallén, P., Saitoh, K., Kozlov, A., and Robertson, B. (2008). Neural bases of

goal-directed locomotion in vertebrates-An overview. Brain Res. Rev. *57*, 2–12.

Howard, M.W., and Kahana, M.J. (2002). A distributed representation of temporal context. J.

Math. Psychol. *46*, 269–299.

Isely, G., Hillar, C.J., and Sommer, F.T. (2010). Deciphering subsampled data: Adaptive

compressive sampling as a principle of brain communication. Adv. Neural Inf.

Process. Syst. 23 24th Annu. Conf. Neural Inf. Process. Syst. 2010, NIPS 2010 1–9.

Izawa, S., Chowdhury, S., Miyazaki, T., Mukai, Y., Ono, D., Inoue, R., Ohmura, Y.,

Mizoguchi, H., Kimura, K., Yoshioka, M., et al. (2019). REM sleep-active MCH

neurons are involved in forgetting hippocampusdependent memories. Science (80-.).

365, 1308–1313.

Justus, D., Dalügge, D., Bothe, S., Fuhrmann, F., Hannes, C., Kaneko, H., Friedrichs, D.,

Sosulina, L., Schwarz, I., Elliott, D.A., et al. (2017). Glutamatergic synaptic

integration of locomotion speed via septoentorhinal projections. Nat. Neurosci. *20*,

16–19.

Kita, T., Nishijo, H., Eifuku, S., Terasawa, K., and Ono, T. (1995). Place and contingency

differential responses of monkey septal neurons during conditional place-object

discrimination. J. Neurosci. *15*, 1683–1703.

Kosse, C., and Burdakov, D. (2019). Natural hypothalamic circuit dynamics underlying

object memorization. Nat. Commun. *10*, 1–8.

Kosse, C., Schöne, C., Bracey, E., and Burdakov, D. (2017). Orexin-driven GAD65 network

of the lateral hypothalamus sets physical activity in mice. Proc. Natl. Acad. Sci. U. S.

A. *114*, 4525–4530.

Leão, R.N., Targino, Z.H., Colom, L. V., and Fisahn, A. (2015). Interconnection and

synchronization of neuronal populations in the mouse medial septum/diagonal band of

Broca. J. Neurophysiol. *113*, 971–980.

Leranth, C., Deller, T., and Buzsáki, G. (1992). Intraseptal connections redefined: lack of a

lateral septum to medial septum path. Brain Res. *583*, 1–11.

Leroy, F., Park, J., Asok, A., Brann, D.H., Meira, T., Boyle, L.M., Eric, W., and Siegelbaum,

S.A. (2018). A circuit from hippocampal CA2 to lateral septum disinhibits social

aggression. Nature *564*, 213–218.

Leutgeb, S., and Mizumori, S. (1999). Excitotoxic septal lesions result in spatial memory

deficits and altered flexibility of hippocampal single-unit representations. J. Neurosci.

19, 6661–6672.

Leutgeb, S., and Mizumori, S. (2002). Context-specific spatial representations by lateral

septal cells. Neuroscience *112*, 655–663.

Luo, A.H., Tahsili-Fahadan, P., Wise, R.A., Lupica, C.R., and Aston-Jones, G. (2011a).

Linking context with reward: a functional circuit from hippocampal CA3 to ventral

tegmental area. Science (80-.). *333*, 353–358.

Luo, A.H., Tahsili-fahadan, P., Wise, R.A., Lupica, C.R., and Aston-jones, G. (2011b).

Supporting Online Material for CA3 to Ventral Tegmental Area.

M'Harzi, M., and Jarrard, L.E. (1992). Effects of medial and lateral septal lesions on

acquisition of a place and cue radial maze task. Behav. Brain Res. *49*, 159–165.

MacDonald, C.J., Carrow, S., Place, R., and Eichenbaum, H. (2013). Distinct hippocampal

time cell sequences represent odor memories in immobilized rats. J. Neurosci. *33*,

14607–14616.

Mahler, S. V., Moorman, D.E., Smith, R.J., James, M.H., and Aston-Jones, G. (2014).

Motivational activation: A unifying hypothesis of orexin/hypocretin function. Nat.

Neurosci. *17*, 1298–1303.

Mankin, E.A., Diehl, G.W., Sparks, F.T., Leutgeb, S., and Leutgeb, J.K. (2015).

Hippocampal CA2 Activity Patterns Change over Time to a Larger Extent than

between Spatial Contexts. Neuron *85*, 190–202.

McGlinchey, E.M., and Aston-Jones, G. (2018). Dorsal hippocampus drives context-induced

cocaine seeking via inputs to Lateral Septum. Neuropsychopharmacology *43*, 987–

1000.

Mogenson, G.J., Jones, D.L., and Yim, C.Y. (1980). From motivation to action: Functional

interface between the limbic system and the motor system. Prog. Neurobiol. *14*, 69–

97.

Nishijo, H., Kita, T., Tamura, R., Uwano, T., and Terasawa, K. (1997). Septal Neuronal

Responses Related to Spatial Representation in Monkeys. Hippocampus *464*, 460–

464.

Oh, S.W., Harris, J.A., Ng, L., Winslow, B., Cain, N., Mihalas, S., Wang, Q., Lau, C., Kuan,

L., Henry, A.M., et al. (2014). A mesoscale connectome of the mouse brain. Nature

508, 207–214.

Olshausen, B.A., and Field, D.J. (2004). Sparse coding of sensory inputs. Curr. Opin.

Neurobiol. *14*, 481–487.

Palm, G., and Sommer, F.T. (2006). Associative Data Storage and Retrieval in Neural

Networks. In Models of Neural Networks III. Physics of Neural Networks, Domany

E., J.L. van Hemmen, and K. Schulten, eds. (Springer, New York, NY.), p.

Pastalkova, E., Itskov, V., Amarasingham, A., and Buzsáki, G. (2008). Internally generated

cell assembly sequences in the rat hippocampus. Science (80-.). *321*, 1322–1328.

Patel, J., Schomburg, E.W., Bere, A., and Fujisawa, S. (2013). Local Generation and

Propagation of Ripples along the Septotemporal Axis of the Hippocampus. *33*,

17029–17041.

Petrovich, G.D. (2018). Lateral hypothalamus as a motivation-cognition interface in the

control of feeding behavior. Front. Syst. Neurosci. *12*, 1–7.

Phelan, K.D., Hasuo, H., Twery, M.J., and Gallagher, J.P. (1989). Projection neurons in the

rat dorsolateral septal nucleus possess recurrent axon collaterals. Neurosci. Lett. *97*,

259–265.

Pnevmatikakis, E.A., and Giovannucci, A. (2017). NoRMCorre: An online algorithm for

piecewise rigid motion correction of calcium imaging data. J. Neurosci. Methods *291*,

83–94.

Rawlins, J.N.P., and Olton, D.S. (1982). The septo-hippocampal system and cognitive

mapping. Behav. Brain Res. *5*, 331–358.

Reis, D.G., Scopinho, A.A., Guimarães, F.S., Corrêa, F., and Resstel, L.B.M. (2010).

Involvement of the lateral septal area in the expression of fear conditioning to context.

Learn. Mem. *17*, 134–138.

Risold, P.Y., and Swanson, L.W. (1996). Structural evidence for functional domains in the rat

hippocampus. Science (80-.). *272*, 1484–1486.

Risold, P.Y., and Swanson, L.W. (1997). Connections of the rat lateral septal complex. Brain

Res. Rev. *24*, 115–195.

Robinson, J., Manseau, F., Ducharme, G., Vigneault, E., El Mestikawy, S., and Williams, S.

(2016). Optogenetic Activation of Septal Glutamatergic Neurons Drive Hippocampal

Theta Rhythms. *36*, 3016–3023.

Rubin, A., Geva, N., Sheintuch, L., and Ziv, Y. (2015). Hippocampal ensemble dynamics

timestamp events in long-term memory. Elife *4*, 1–16.

Sargolini, F., Fyhn, M., Hafting, T., McNaughton, B.L., Witter, M.P., Moser, M.B., and

Moser, E.I. (2006). Conjunctive representation of position, direction, and velocity in

entorhinal cortex. Science (80-.). *312*, 758–762.

Sheintuch, L., Rubin, A., Brande-Eilat, N., Geva, N., Sadeh, N., Pinchasof, O., and Ziv, Y.

(2017). Tracking the Same Neurons across Multiple Days in Ca2+Imaging Data. Cell

Rep. *21*, 1102–1115.

Sinnamon, H.M. (1993). Preoptic and hypothalamic neurons and the initiation of locomotion

in the anesthetized rat. Prog. Neurobiol. *41*, 323–344.

Sparks, P., and Ledoux, J. (2000). The septal complex as seen through the context of fea, In:

the behavioral neuroscience of the septal region (Springer-Verlag).

Stackman, R.W., and Taube, J.S. (1998). Firing properties of rat lateral mammillary single

units: Head direction, head pitch, and angular head velocity. J. Neurosci. *18*, 9020–

9037.

Staiger, J.F., and Nürnberger, F. (1989). Pattern of afferents to the lateral septum in the

guinea pig. Cell Tissue Res. *257*, 471–490.

Staiger, J.F., and Nürnberger, F. (1991). The efferent connections of the lateral septal nucleus

in the guinea pig: intrinsic connectivity of the septum and projections to other

telencephalic areas. Cell Tissue Res. *264*, 415–426.

Stanton, M.E., Thomas, G.J., and Brito, G.N.O. (1984). Posterodorsal septal lesions impair

performance on both shift and stay working memory tasks. Behav. Neurosci. *98*, 405–

415.

Stuber, G.D., and Wise, R.A. (2016). Lateral Hypothalamic Circuits for Feeding and Reward.

Nat. Neurosci. *19*, 198–205.

Swanson, L.W., and Cowan, W.M. (1977). An Autoradiographic Study of the Organization

of the Efferent Connections of the Hippocampal Formation in the Rat. J. Comp. Neur.

172, 49–84.

Swanson, L.W., and Cowan, W.M. (1979). The Connections of the Septal Region in the Rat.

J. Comp. Neur. *186*, 621–656.

Sweeney, P., and Yang, Y. (2016). An Inhibitory Septum to Lateral Hypothalamus Circuit

That Suppresses Feeding. J. Neurosci. *36*, 11185–11195.

Takamura, Y., Tamura, R., Zhou, T.L., Kobayashi, T., Tran, A.H., Eifuku, S., and Ono, T.

(2006). Spatial Firing Properties of Lateral Septal Neurons. *644*, 635–644.

Taube, J.S. (1995). Head direction cells recorded in the anterior thalamic nuclei of freely

moving rats. J. Neurosci. *15*, 70–86.

Taube, J., Muller, R., and Ranck, J. (1990). Head-direction cells recorded from the

postsubiculum in freely moving rats. I. Description and quantitative analysis. J.

Neurosci. *10*, 420–435.

Thomas, G.J. (1979). Comparison of effects of small lesions in posterodorsal septum on

spontaneous and rerun correction (contingently reinforced) alternation in rats. J.

Comp. Physiol. Psychol. *93*, 685–694.

Thomas, G.J., and Brito, G.N. (1980). Recovery of delayed alternation in rats after lesions in

medial frontal cortex and septum. J. Comp. Physiol. Psychol. *94*, 808–818.

Tingley, D., and Buzsáki, G. (2018). Transformation of a Spatial Map across the

Hippocampal-Lateral Septal Circuit. Neuron *98*, 1229-1242.e5.

Trent, N.L., and Menard, J.L. (2010). The ventral hippocampus and the lateral septum work

in tandem to regulate rats' open-arm exploration in the elevated plus-maze. Physiol.

Behav. *101*, 141–152.

Vetere, G., Kenney, J.W., Tran, L.M., Xia, F., Steadman, P.E., Parkinson, J., Josselyn, S.A.,

and Frankland, P.W. (2017). Chemogenetic Interrogation of a Brain-wide Fear

Memory Network in Mice. Neuron *94*, 363-374.e4.

Wickersham, I.R., Finke, S., Conzelmann, K.K., and Callaway, E.M. (2007). Retrograde

neuronal tracing with a deletion-mutant rabies virus. Nat. Methods *4*, 47–49.

Wirtshafter, H.S., and Wilson, M.A. (2019). Locomotor and Hippocampal Processing

Converge in the Lateral Septum. Curr. Biol. *29*, 3177-3192.e3.

Wirtshafter, H.S., and Wilson, M.A. (2020). Differences in reward biased spatial

representations in the lateral septum and hippocampus. Elife *9*, 1–20.

Witter, M.P., Daelmans, H.E., Jorritsma-Byham, B., Staiger, J.F., and Wouterlood, F.G.

(1992). Restricted origin and distribution of projections from the lateral to the medial

septal complex in rat and guinea pig. Neurosci. Lett. *148*, 164–168.

Xu, Y., Lu, Y., Cassidy, R.M., Mangieri, L.R., Zhu, C., Huang, X., Jiang, Z., Justice, N.J.,

Xu, Y., Arenkiel, B.R., et al. (2019). Identification of a neurocircuit underlying

regulation of feeding by stress-related emotional responses. Nat. Commun. *10.*

Zetsche, C. Sparse coding: the link between low level vision and associative memory. In

Parallel Processing in Neural Systems and Computers, R. Eckmiller, G. Hartmann,

and G. Hauske, eds. (Elsevier, Amsterdam, Holland), p.

Zhao, C., Eisinger, B., and Gammie, S. (2013). Characterization of GABAergic Neurons in

the Mouse Lateral Septum: A Double Fluorescence In Situ Hybridization and

Immunohistochemical Study Using Tyramide Signal Amplification. PLoS One *8*, 1–

24.

Zhou, P., Resendez, S.L., Rodriguez-Romaguera, J., Jimenez, J.C., Neufeld, S.Q., Stuber,

G.D., Hen, R., Kheirbek, M.A., Sabatini, B.L., Kass, R.E., et al. (2017). Efficient and

accurate extraction of in vivo calcium signals from microendoscopic video data.

Zhou, T.L., Tamura, R., Kuriwaki, J., and Ono, T. (1999). Comparison of Medial and Lateral

Septal Neuron Activity During Performance of Spatial Tasks in Rats. *234*, 220–234.

Zingg, B., Chou, X. lin, Zhang, Z. gang, Mesik, L., Liang, F., Tao, H.W., and Zhang, L.I.

(2017). AAV-Mediated Anterograde Transsynaptic Tagging: Mapping

Corticocollicular Input-Defined Neural Pathways for Defense Behaviors. Neuron *93*,

33–47.

Ziv, Y., Burns, L.D., Cocker, E.D., Hamel, E.O., Ghosh, K.K., Kitch, L.J., Gamal, A. El, and

Schnitzer, M.J. (2013). Long-term dynamics of CA1 hippocampal place codes. Nat.

Neurosci. *16*, 264–266.

Main figures

Figure 1: LS shares spatial coding characteristics with its main hippocampal inputs

CA1 and CA3 during goal directed navigation in a 1D environment. A) Retrograde rabies

tracing injection site, blue = DAPI staining, green = TVA.oG coupled to an enhanced green

fluorescent protein (eGFP), red = Rabies coupled to mCherry, orange: starter cells expressing

both TVA.oG.eGFP and Rabies.mCherry. B) Retrograde transfection in the dorsal

hippocampus, coronal section. Bottom left: example of transfected CA1 pyramidal cell,

bottom right: transfected CA3 pyramidal cells. (For additional images see Supplemental

figure 1.) C) Top: Diagram of one-photon calcium recording setup in LS, CA1 and CA3 in

freely behaving mice, middle: histological verification of implantation site and bottom:

extracted calcium transients for LS, CA1 and CA3. D) Top: Recoding strategy in these

animals, with animal alternating left and right trajectories to obtain a sucrose reward on a 100

or 130 cm linear track. Bottom: Probability of an example cell to be active given the location

on the linear track (red) with 95% upper and lower percentile (grey). Examples shown are

from two different LS mice, and a representative place cell from one CA1 and CA3 animal

each. F) Proportion of significantly spatially modulated cells for each animal for each

recording region. Total recorded cells: LS: n = 1030 cells, N = 15 mice. CA1: n = 1148 cells,

N = 5 mice, CA3: n = 144 cells, N = 2 mice. G) Split half stability of recorded spatial cells in

each region and right: probability being active (LS: n = 475 cells, CA1: n = 336 cells, CA3: n

= 47 cells). H) Top: Posterior probabilities for each frame estimated from binarized calcium

activity, bottom: corresponding actual (green) and decoded (red) location. I) Schematic

representation of bootstrapping approach. J) Method for computing the Euclidean distance

mean decoding error in cm for each bootstrapped estimate and mean decoding agreement for

each bootstrapped estimate for open field recordings. K) Decoding error for actual vs shuffled

dataset (30 times bootstrapped sample of 60 cells, LS: N = 7 mice, CA1: N =5 mice, CA3: LS

= 2 mice). L) Mean decoding score each region, *, $p < .05$, **, $p < .01$, ****, $p < .0001$, Test

used in F-G, Kruskal-Wallis, K, paired t-test, L, one-way ANOVA.

1018

Figure 2: LS shares spatial coding characteristics with its main hippocampal inputs

CA1 and CA3 during free exploration in a 2D environment. A) Tuning curves of example

place cells recorded from LS, dorsal CA1 and dorsal CA3 in a 45 cm open field (3 cm bins).

B) Proportion of place cells per animal (LS: n = 1899 cells from N = 28 mice; dCA1: n =

1229 cells, N = 6 mice; dCA3: n = 204 cells, N = 4 mice). C) Within session stability for

spatial cells (LS: n = 734 place cells from N = 28 mice; dCA1: n = 424 place cells, N = 6

mice; dCA3: n = 86 place cells, N = 4 mice). D) Probability of being active for spatial cells

(LS: n = 734 place cells from N = 28 mice; dCA1: n = 424 place cells, N = 6 mice; dCA3: n

= 86 place cells, N = 4 mice). E) Left: mean dispersion computation, right: mean dispersion

for spatial cells (LS: n = 734 place cells from N = 28 mice; dCA1: n = 424 place cells, N = 6

mice; dCA3: n = 86 place cells, N = 4 mice). F) Bootstrapping approach using 100 cells. G)

Method for computing the mean decoding error and mean decoding agreement for each

bootstrapped estimate using 100 cells. H) Mean decoding error for LS (N = 6), CA1 (N =4)

and CA3 (N = 2). **, $p < .01$, ***, $p < .001$, ****, $p < .0001$, Test used in B, one-way

ANOVA, C-E, Kruskal-Wallis, Dunn's multiple comparison test. H, two-way ANOVA,

Sidak's multiple comparisons test.

1035

Figure 3: LS place code remains similar to CA1 over longer periods of time. A)

Experimental setup (top) and aligned spatial footprints of cells recorded over days (bottom).

B) Tuning maps for two sets of stable cells recorded over all days, with each row being one

aligned cell. Tuning map correlation indicated at the bottom. C) Tuning map correlation for

all aligned cells (black) versus shuffled cells (red) for progressive days (day 1, n = 562; day

2, n = 600; day 3, n = 477; day 8, n = 492, n = 5 mice). D) Proportion of place cells recorded

for each day. Each dot is an animal (N = 5). E) Within session stability for each place cell for

all recorded cells (day 1, n = 181; day 2, n = 172; day 3, n = 178; day 8, n = 209, N = 5 mice).

F) Average MI for each day for all recorded cells (day 1, n = 562; day 2, n = 600; day 3, n =

477; day 8, n = 492, n = 5 mice). G) Matrix of mean tuning map correlation for all aligned

place cells recorded in LS (N = 3 mice) and CA1 (N = 1 mouse). H) Tuning map correlation

between day 1 and day 8 for all aligned place cells recorded in LS vs CA1 (LS: 42 cells, n = 3

mice, CA1: 28 cells, n = 1 mouse). *, $p < .05$, **, $p < .01$, ****, $p < .0001$. Test used in C,

two-way ANOVA, Sidak's multiple comparisons test, D, one-way ANOVA, E-F, Kruskall

Wallis test, with Dunn's multiple comparison test, H, unpaired t-test.

1051

Figure 4: LS GABAergic cells encode direction, speed, and acceleration. A) Left:

Activity of an example LS neuron during free exploration in an open field, top: head direction

(HD, red), middle: binarized activity (yellow), bottom: raw calcium activity (blue). Right:

Corresponding polar plot indicating the probability of the cell being active as a function of

the animals' HD with (black, p(active | bin), red lines indicate 95% upper and lower

percentile. MI calculated using 9° bins. Supplemental figure 7. B) Proportion of significantly

directionally modulated cells in LS, dCA1 and dCA3. C) Left: Activity of an example LS

neuron during free exploration modulated by velocity (top, red), middle: binarized activity

(yellow), bottom: raw calcium activity (blue). Right: Tuning curve for velocity for the

example cell (red) with 95% upper and lower percentile in grey. MI calculated using 20 bins.

D) Proportion of significantly speed modulated cells in LS, dCA1 and dCA3. E) Left:

Activity of an example LS neuron during free exploration modulated by acceleration (top,

red), middle: binarized activity (yellow), bottom: raw calcium activity (blue). Right: Tuning

curve for acceleration for the example cell (red) with 95% upper and lower percentile in grey.

MI calculated using 20 bins. F) Comparison of proportions of significantly modulated cells

for region. G) Example of an LS cell that is both significantly head direction modulated (left),

as well as spatially modulated (right). H) Left: Proportion of cells that are significantly

modulated by only one modality (grey), two modalities (yellow), three (red) or all 4 of the

investigated variables (black). The number above the bars indicates the absolute number of

cells found to be modulated in the total population (n = 1230 cells, N = 19 mice). Right:

absolute proportion of cells modulated by any combination of variables. Test used in B, D, F:

Kruskal-Wallis test with Dunn's multiple comparisons test. *, $p < .05$. D = direction, A =

acceleration, S = spatial coding, V = velocity.

1075

Figure 5: LS cells receiving hippocampal inputs project directly to hypothalamus and

medial septum. A) Diagram of dual-viral injection strategy for anterograde transsynaptic

tracing. The diagram is based on dorsal CA1 targeted injection, dCA3 is also used for

injections, together with coronal sections showing primary injection site in dorsal CA1 (top)

or dorsal CA3 (bottom). Red = tdTom expression, blue = DAPI, with zoomed images

showing transfected cell bodies are predominantly located in the pyramidal layer (bottom). B)

eYFP-positive cell bodies at the anterior dorsal LS and fibers at the level of the medial

septum following dCA1 injection (left) and dCA3 injection (right). Inset left: eYFP-positive

fibers at the level of MS. C) Second-order transfection is seen bilaterally at the level of the

lateral hypothalamus following dCA1 injection (left) and dCA3 injection (right). D) Injection

strategy for Cre-dependent AAV. Synaptag mediated tracing in the LS. E) Coronal section of

transfection at the level of dorsal LS, with synaptophysin-bound eYFP at the MS. Top right:

Zoomed images showing transfection at the injection site. Bottom right: eYFP-positive fibers.

F) Schematic with proposed connections of the LS within the hippocampal network. Scale

bars: B, left and right: 800 μm. C, left and right: 500 μm/ DM = dorsomedial hypothalamic

nucleus, 3V = third ventricle, LH = lateral hypothalamus.

1092

Figure 6: LS-cells coding for spatial and idiothetic information are non-uniformly

distributed along the dorsal-ventral and anterior-posterior axis. A) Strategy to record

from different anterior-posterior levels in LS (left) and implantation sites covered. B)

Strategy to approximate cell location. Background: Maximal projection of representative

recording, red: outline of approximated GRIN lens position, white: distance from the center

of the GRIN lens to the cells of interest. C) Top: For .2 mm bins along dorsal-ventral axis,

proportion of stable (within-session stability > .5) place cells (N = 24 animals), head direction

cells (N = 17 animals), velocity cells (N = 24 animals) and acceleration cells (N = 24

animals) respectively. Bottom: Same as top for anterior-posterior axis. Each dot represents

the average per animal per bin along the anterior-posterior axis (left) and along the dorsal-

ventral axis. Black lines indicate 95% confidence intervals. *, $p < .05$, **, $p < .01$, Test used

in C, linear regression

1105

1106

1107

Figure 1

Figure 2

A LS — CA1 — CA3

MI: 0.0499 MI: 0.0624 MI: 0.0395 MI: 0.0432 MI: 0.046
Stability: 0.56 Stability: 0.82 Stability: 0.70 Stability: 0.82 Stability: 0.91

MI: 0.0517 MI: 0.0521 MI: 0.0603 MI: 0.0456 MI: 0.0732
Stability: 0.81 Stability: 0.60 Stability: 0.77 Stability: 0.86 Stability: 0.70

P(active|location)

B %Spatial cells — ns — LS dCA1 dCA3

C within-session stability — *** **** — LS dCA1 dCA3

D Prob. being active — **** **** ** — LS dCA1 dCA3

E n binarized events

mean dispersion = mean distance to centroid

10 + 20 + 12 + ... + ... / n

+ centroid

mean place field dispersion (cm) — **** **** — LS dCA1 dCA3

F Recorded cells

● included in decoding (100 cells)

n = 30 bootstrap samples

a1
a2
...
a30

G open field   bin 3 6 9 12...

45 cm   9 cm   15 cm

actual position
decoded position

decoding error = 17.49 cm

actual position   decoded position

3 cm bin

decoding agreement   0   0   1

0.333

H mean decoding error (cm) — *** ** ** — LS CA1 CA3

Shuffle
Actual

Figure 3

A AAV.hSyn. GCaMPF6f

recent — *day 1* *day 2* *day 3* — *remote* *day 8*

45 cm

1 2 3 8

B

day 1 *day 2* *day 3* *day 8*

MI : 0.0707 MI : 0.0971 MI : 0.1223 MI : 0.0601

Stability: 0.78 Stability: 0.94 Stability: 0.88 Stability: 0.77

Tuning map correlation: 0.83 0.86 0.75

MI : 0.0499 MI : 0.0564 MI : 0.0766 MI : 0.0812

Stability: 0.83 Stability: 0.87 Stability: 0.88 Stability: 0.61

Tuning map correlation: 0.89 0.93 0.86

P(active|location) 0 0.05 0.1 0.15

C Tuning map correlation — **** **** **** day 1-2 day 2-3 day 3-8

D % Place cells — ns Day 1 Day 2 Day 3 Day 8

E place cell within session stability — * Day 1 Day 2 Day 3 Day 8

F MI (bits) — ** Day 1 Day 2 Day 3 Day 8

G *LS* *CA1* — Day 1 Day 2 Day 3 Day 8 tuning map correlation 0 0.5 1

H Tuning map correlation Day 1-8 — ns CA1 LS

Figure 4

Figure 5

A

AAV.hSyn.Cre
AAV.flex.tdTom
AAV.flex.eYFP

CA1
CA3
HPC
LS
Cre +
no Cre

CA1 injected

CA1
Dapi   tdTom

CA3 injected

Dapi   tdTom   CA3

B

CA1 injected

LSd

CA3 injected

LSd
LSi
LSv
MS

C

CA1 injected

DM

CA3 injected

DM

D

AAVdj.hSyn.flex.
mRuby2.syp.eGFP

LS
MS
DBB

E

LSd
LSi
LSv
MS

Merged   mRuby2   syp.eYFP

syp.eYFP

F

hypothalamus

CA1
Sub
LS
DG
CA3
hippocampus
MS
θ

glutamate
GABA
ACh

Figure 6

A

+0.86
anterior LS

+0.38
intermediate LS

+0.02
posterior LS

GRIN lens location

B

medial

+ 220 μm

anterior

+ 50 μm

500 μm

C

Dorsal-Ventral axis

Anterior-Posterior axis

+0.86

+0.02

spatial

Stable Place Cells (%)

p = 0.084

dorsal - ventral axis
(distance from Bregma)

Stable Place Cells (%)

**

anterior - posterior axis
(distance from Bregma)

head direction

Proporton Stable HD Cells per Bin (%)

**

R2 = 0.3837

Dorsal - Ventral axis
(distance from Bregma)

Proporton Stable HD Cells per Bin (%)

ns

Anterior - Posterior axis
(distance from Bregma)

velocity

Proporton Velocity Cells per Bin (%)

*

R2 = 0.1836

Dorsal - Ventral axis
(distance from Bregma)

Proporton Velocity Cells per Bin (%)

ns

Anterior - Posterior axis
(distance from Bregma)

acceleration

Proporton Acceleration Cells per Bin (%)

ns

Dorsal - Ventral axis
(distance from Bregma)

Proporton Acceleration Cells per Bin (%)

ns

Anterior - Posterior axis
(distance from Bregma)