A region-based method for causal mediation analysis of DNA methylation data

Qi Yan[1*], Erick Forno[2], Juan C. Celedón[2] and Wei Chen[2,3]

[1]Department of Obstetrics and Gynecology, Columbia University Irving Medical Center,

New York, NY

[2]Division of Pediatric Pulmonary Medicine, UPMC Children's Hospital of Pittsburgh,

University of Pittsburgh, Pittsburgh, PA;

[3]Department of Biostatistics, Graduate School of Public Health, University of Pittsburgh,

Pittsburgh, PA;

[*]Correspondence should be addressed to:

Q.Y. (qy2253@cumc.columbia.edu)

Funding

This research was supported by grant HL138098 from the U.S. National Institutes of

Health (NIH) to Q.Y. J.C.C. was supported by grants HL079966, HL117191, MD011764

and HL150431 from the U.S. NIH. W.C. was supported by grant HL150431 from the U.S.

NIH.

ABSTRACT

Exposure to environmental factors can affect DNA methylation at a CpG site or a genomic region, which can then affect an outcome. In other words, environmental effects on an outcome could be mediated by DNA methylation. To date, single CpG site-based mediation analysis has been employed extensively. More recently, however, there has been considerable interest on studying differentially methylated regions (DMRs), both because DMRs are more likely to have functional effects than single CpG sites and because testing DMRs reduces multiple testing. In this report, we propose a novel causal mediation approach under the counterfactual framework to test the significance of total, direct and indirect effects of predictors on response variable with a methylated region (MR) as the mediator (denoted as MR-Mediation). Functional linear transformation is used to reduce the possible high dimension of the CpG sites in a predefined methylated region and to account for their location information. In our simulation studies, MR-Mediation retained the desired Type I error rates for total, direct and indirect effect tests, for both continuous and binary outcomes. Furthermore, MR-Mediation had better power performance than testing mean methylation level as the mediator in most considered scenarios, especially for indirect effect (i.e., mediated effect) test, which could be more interesting than the other two effect tests. We further illustrate our proposed method by analyzing the methylation mediated effect of exposure to gun violence on total immunoglobulin E (IgE) or atopic asthma among participants in the Epigenetic Variation and Childhood Asthma in Puerto Ricans (EVA-PR) study.

INTRODUCTION

Epigenetic studies are key to understanding regulatory mechanisms of gene expression. Of the available epigenetic markers, DNA methylation is the most stable and widely studied epigenetic modification.[1] DNA methylation refers the addition of a methyl group at the 5′ location of cytosine nucleotides (C), with this modification occurring predominantly at Cs that are immediately followed by a guanine (G) in the 5′ to 3′ direction, denoted CpG. DNA methylation often affects gene transcription and has been associated with complex diseases[2-5] and cancers.[6-10]

DNA methylation analysis has traditionally focused on individual CpG sites. Over the last few years, there has been great interest in developing region-based methods to detect differentially methylated regions (DMRs).[11, 12] Such region-based approach is supported by strong correlation in DNA methylation levels across regions of the genome,[13, 14] as well as by the fact that methylated regions such as CpG islands,[15] CpG island shores,[16] or generic 2-kb regions[17] are more often linked to functionally relevant findings than single CpG sites. Statistically, if there are multiple causative CpG sites that have small individual effects, single-marker testing may have limited power to detect those weak signals. On the other hand, region-based approaches have higher power by combining the effects of multiple CpG sites. Moreover, region-based methods greatly reduce the burden of multiple testing in a genome-wide study.

DMR identification methods could be classified into two categories: 1. Supervised methods that first calculate the p-values for the association between each single CpG site

and the outcome of interest, and then identify the regions with adjacent small p-values,

and 2. Unsupervised methods that predefine the regions without using any outcome

information, and then test the association between methylation level in the genomic

regions and the outcome of interest.

Exposure to environmental factors can affect DNA methylation, which can then affect a

disease or condition of interest. In other words, environmental effects on a disease or

condition could be mediated by DNA methylation (Figure 1). Mediation analysis

investigates the relationship between an exposure and an outcome, while examining how

such exposure and outcome relate to the mediator. Although the Baron and Kenny's "four

steps" method[18] has been the most commonly used approach to mediation analysis, such

approach has been recently extended using a counterfactual framework[19-21] to allow the

presence of an interaction between the exposure and mediator, as well as to decompose

the estimate of total effect (TE) into estimates of direct (DE) and indirect effects (IE). To

date, single CpG site-based mediation analysis has been employed extensively. This

approach has indeed shown that DNA methylation can mediate the effects of

environmental exposure on complex traits,[22-24] complex diseases[25-27] and cancers.[28, 29]

Figure 1. Mediation model with DNA methylation as the mediator.

In this study, we propose a novel causal mediation approach under the counterfactual

framework to test the significance of total, direct, and indirect effects of DNA methylation

in a genomic region on an outcome of interest. In this approach, a group of CpG sites

from a predefined region are utilized as the mediator, and functional linear

transformation[30] is used to reduce the possible high dimensions in the region's CpG sites

and account for their location information. We denote the method as MR-Mediation. To

evaluate the performance of the proposed MR-Mediation statistic, we first conduct

extensive simulation studies to assess Type I error rates and power. We then illustrate

our proposed method by analyzing whether DNA methylation mediates the association

between exposure to gun violence and total immunoglobulin E (IgE) or atopic asthma

among children participating in the Epigenetic Variation and Childhood Asthma in Puerto

Ricans study (EVA-PR).[31]

METHODS

Functional linear transformation for region-based CpG sites

The idea is to linearly combine a few basis functions to fit a curve that can closely

represent the methylation levels of the CpG sites in the region (e.g., an illustrative

example is shown in Figure S1). Thus, the dimension can be reduced and the location

information can be captured by the fitted curve. Specifically, we consider n subjects who

have m CpG sites in a predefined region. The physical locations of the m CpG sites are

normalized to a range of $[0, 1]$ (i.e., $0 \leq l_1 \ldots l_m \leq 1$). Thus, we use $M_i(l_j)$ to denote the

methylation level of a CpG site at j^{th} location from the i^{th} subject. We assume that the

$n \times 1$ vector of the continuous outcome $\mathbf{y}$ follows a linear regression model. For the i^{th}

subject,

$$y_i = \mathbf{Z_i}\boldsymbol{\beta}_0 + \int_0^1 M_i(l)\,dl \times \beta_M + \varepsilon_i \qquad (1)$$

where $\mathbf{Z_i}$ is a $1 \times p$ covariate vector, $\boldsymbol{\beta}_0$ is a $p \times 1$ parameter vector (an intercept and $p -$

1 covariates), $M_i(l)$ is a function of CpG site locations and includes m CpG site values,

β_M is the effect of $M_i(l)$, and ε_i is the random error. By using the ordinary linear

smoother,[32] we let

$$M_i(l) = \big(M_i(l_1), \dots, M_i(l_m)\big)\boldsymbol{\Phi}[\boldsymbol{\Phi}'\boldsymbol{\Phi}]^{-1}\boldsymbol{\phi}(l)$$

where $\big(M_i(l_1), \dots, M_i(l_m)\big)$ is a $1 \times m$ vector for m CpG sites in the region, $\boldsymbol{\phi}(l)$ is a

$K_M \times 1$ vector including K_M basis functions, and $\boldsymbol{\Phi}$ is a $m \times K_M$ matrix containing the

values of $\boldsymbol{\phi}(l)$ at $l_1 \dots l_m$. Thus, Formula (1) can be rewritten as

$$y_i = \mathbf{Z_i}\boldsymbol{\beta}_0 + R_i\beta_M + \varepsilon_i \qquad (2)$$

where $R_i = \big(M_i(l_1), \dots, M_i(l_m)\big)\boldsymbol{\Phi}[\boldsymbol{\Phi}'\boldsymbol{\Phi}]^{-1}\int_0^1 \boldsymbol{\phi}(l)\,dl$ is a scalar. Therefore, R_i can be

viewed as a summary of the methylation level in the region. In this study, we focus on

cubic B-spline basis functions.

Association between the mediator and the exposure variable

To test the proportions of the TE, DE and IE of the exposure on the outcome variables

that are mediated by DNA methylation, we need to first make sure there exists an

association between the mediator and exposure variables. To do so, we consider a linear

regression model for n independent subjects,

$$R_i = \mathbf{Z_i}\boldsymbol{\alpha}_0 + X_i\alpha_X + \varepsilon_{R_i} \qquad (3)$$

where R_i is a scalar representing the regional methylation level for the i^{th} subject, $\mathbf{Z}_i$ is a

$1 \times p$ covariate vector, $\boldsymbol{\alpha}_0$ is a $p \times 1$ parameter matrix (an intercept and $p-1$ covariates),

X_i is a scalar for the exposure variable, α_X is the parameter for exposure, and ε_{R_i} is a

random error. In this study, we first test $H_0: \alpha_X = 0$, and then further test any methylation

mediated effects between the exposure and outcome variables only when $\alpha_X \neq 0$.

Counterfactual approach to the causal mediation model

The total effect (TE) from exposure to outcome can be decomposed into the indirect effect

(IE) mediated through DNA methylation and the direct effect (DE) not mediated by DNA

methylation, which could be mediated by other mechanisms or a direct link between

exposure and outcome. For a continuous outcome, the mediation model for the i^{th}

subject can be written as

$$\boldsymbol{E}(y_i|\mathbf{Z}_i, \mathrm{X}_i, \mathrm{R}_i) = \mathbf{Z}_i\boldsymbol{\beta}_0 + \mathrm{X}_i\beta_X + \mathrm{R}_i\beta_M + \mathrm{X}_i\mathrm{R}_i\beta_{XM} \qquad (4)$$

where $\mathbf{Z}_i\boldsymbol{\beta}_0$ is for the covariates, $\mathrm{X}_i\beta_X$ is for the exposure variable, $\mathrm{R}_i\beta_M$ is for the regional

methylation level, and $\mathrm{X}_i\mathrm{R}_i\beta_{XM}$ is for the interaction between the exposure variable and

the regional methylation level. To use the counterfactual approach to the causal mediation

model, four identifiability assumptions need to be satisfied: 1. no unmeasured

confounding of the exposure and outcome relationship; 2. no unmeasured confounding

of the mediator and outcome relationship; 3. no unmeasured confounding of the exposure

and mediator relationship; 4. the exposure must not cause any known confounder of the

mediator and outcome relationship.[33] If all assumptions hold, the DE, IE and TE for

change in exposure from level x_0 to x_1 are given by

$$DE = \boldsymbol{E}(y_i|\mathbf{Z}_i, x_1, \mathrm{R}_i(x_0)) - \boldsymbol{E}(y_i|\mathbf{Z}_i, x_0, \mathrm{R}_i(x_0))$$

$$= (x_1 - x_0)[\beta_X + (\mathbf{Z}_i\boldsymbol{\alpha}_0 + x_0\alpha_X)\beta_{XM}]$$

$$IE = \boldsymbol{E}\big(y_i|\mathbf{Z}_i, x_1, \mathrm{R}_i(x_1)\big) - \boldsymbol{E}\big(y_i|\mathbf{Z}_i, x_1, \mathrm{R}_i(x_0)\big)$$

$$= (x_1 - x_0)\alpha_X(\beta_M + x_1\beta_{XM})$$

$$TE = \boldsymbol{E}\big(y_i|\mathbf{Z}_i, x_1, \mathrm{R}_i(x_1)\big) - \boldsymbol{E}\big(y_i|\mathbf{Z}_i, x_0, \mathrm{R}_i(x_0)\big)$$

$$= DE + IE$$

Since we only focus on the scenarios with $\alpha_X \neq 0$, conceivably, it can be shown that

$$H_0: DE = 0 \text{ equivalent to } H_0: \beta_X = \beta_{XM} = 0$$

$$H_0: IE = 0 \text{ equivalent to } H_0: \beta_M = \beta_{XM} = 0 \qquad (5)$$

$$H_0: TE = 0 \text{ equivalent to } H_0: \beta_X = \beta_M = \beta_{XM} = 0.$$

Thus, we can test $H_0: DE = 0$ by a *F*-test with degrees of freedom $(2,\ n - p - 3)$, $H_0: IE =$

0 by a *F*-test with degrees of freedom $(2,\ n - p - 3)$, and $H_0: TE = 0$ by a *F*-test with

degrees of freedom $(3,\ n - p - 3)$.

Similarly, when the outcome is binary, the mediation model for the i^{th} subject can be

written as

$$logit\big(P(y_i = 1|\mathbf{Z}_i, \mathrm{X}_i, \mathrm{R}_i)\big) = \mathbf{Z}_i\boldsymbol{\beta}_0 + \mathrm{X}_i\beta_X + \mathrm{R}_i\beta_M + \mathrm{X}_i\mathrm{R}_i\beta_{XM} \qquad (6).$$

If all identifiability assumptions hold, the DE, IE and TE for change in exposure from level

x_0 to x_1 are given by

$$log\{OR^{DE}\} = logit\left(P\big(y_i = 1|\mathbf{Z}_i, x_1, \mathrm{R}_i(x_0)\big)\right) - logit\left(P\big(y_i = 1|\mathbf{Z}_i, x_0, \mathrm{R}_i(x_0)\big)\right)$$

$$log\{OR^{IE}\} = logit\left(P\big(y_i = 1|\mathbf{Z}_i, x_1, \mathrm{R}_i(x_1)\big)\right) - logit\left(P\big(y_i = 1|\mathbf{Z}_i, x_1, \mathrm{R}_i(x_0)\big)\right)$$

$$log\{OR^{TE}\} = logit\left(P\big(y_i = 1|\mathbf{Z}_i, x_1, \mathrm{R}_i(x_1)\big)\right) - logit\left(P\big(y_i = 1|\mathbf{Z}_i, x_0, \mathrm{R}_i(x_0)\big)\right).$$

According to Valeri and VanderWeele (2013),[33] a rare outcome assumption (i.e., low

disease prevalence) is needed so as to have the odds ratio (OR) in the case-control

design equivalent to the OR in the population, which is further approximately equivalent

to the relative risk (RR) in the population. We also adopt the rare outcome assumption in

this study. Then, the DE, IE and TE can be derived as

$$log\{OR^{DE}\} \approx (x_1 - x_0)[\beta_X + (\mathbf{Z}_i\boldsymbol{\alpha}_0 + x_0\alpha_X + \beta_M\sigma_R^2)\beta_{XM}] + \frac{1}{2}(x_1^2 - x_0^2)\beta_{XM}^2\sigma_R^2$$

$$log\{OR^{IE}\} \approx (x_1 - x_0)\alpha_X(\beta_M + x_1\beta_{XM})$$

$$log\{OR^{TE}\} = log\{OR^{DE}\} + log\{OR^{IE}\},$$

where σ_R^2 is the variance of random error from the mediator regression, Formula (3). The

existence of σ_R^2 is due to $exp(\varepsilon_R) \sim lognormal(0, \sigma_R^2)$. The null hypotheses for testing

$DE = 0, IE = 0$ and $TE = 0$ for binary outcome are the same as given for continuous

outcome. We can test $H_0: DE = 0$ by a χ^2-distributed Rao's score statistic with 2 degrees

of freedom, $H_0: IE = 0$ with 2 degrees of freedom, and $H_0: TE = 0$ with 3 degrees of

freedom. Please note that the mediator regression, Formula (3), is run only for controls

to account for the case-control design.[33] In other words, because of the rare outcome

assumption, the exposure distribution in the controls from the case-control design is

approximately equivalent to the exposure distribution in the whole population.

The Epigenetic Variation and Childhood Asthma in Puerto Ricans study (EVA-PR)

EVA-PR is a case-control study of asthma in subjects aged 9 to 20 years. Subject

recruitment and study procedures for EVA-PR have been described elsewhere.[31]

Genome-wide DNA methylation was measured in 488 nasal epithelial samples using the

Infinium HumanMethylation450 BeadChip arrays (Illumina, San Diego, CA). The

preprocessing and quality control (QC) procedures were described in our prior study.[31]

After quality control, CpG sites with an overall mean β-value within [0.1, 0.9] were kept,

leaving 227,901 CpG sites in the final dataset. Methylation M-value was calculated by

$\log_2$(β-value/(1–β-value)). This methylation dataset was used in both the following

simulation studies and in real data analysis.

The child's lifetime exposure to gun violence was treated as the exposure variable in the

following real data analysis. Lifetime exposure to gun violence was derived from the

exposure to violence scale[34-36] and analyzed as a binary variable (having heard gunshots

at least twice vs. no more than once).[37] Atopy was defined as an IgE ≥0.35 IU/mL to ≥1

of five common allergens (Der p 1, Bla g 2, Fel d 1, Can f 1, and Mus m 1). Asthma was

defined as a physician's diagnosis plus at least one episode of wheeze in the previous

year. Atopic asthma was defined as the presence of both atopy and asthma, and thus

treated as the binary outcome variable in the real data analysis; controls were non-

asthmatic subjects. Total plasma IgE was treated as continuous outcome after $\log_{10}$-

transformation.

Simulation studies

We used the CpG sites from 488 nasal epithelial samples in EVA-PR for our simulation

studies. The CpG sites were grouped into corresponding genes, and the kept CpG sites

also needed to be within 5 kb distance from its nearest gene. The histogram plot (Figure

S2) showed that most of the genes included less than 20 CpG sites, although some genes

could have several hundred CpG sites. Thus, we selected different sizes of genes in our

simulation studies. Specifically, we selected (1) 705 genes with 10 CpG sites; (2) 306

genes with 15 CpG sites; (3) 152 genes with 20 CpG sites; and (4) 21 genes with 40 CpG

sites.

Furthermore, we need to choose the basis functions and the number of basis functions

to reduce the dimension of each gene. To simulate the exposure $\mathbf{X}$ and outcome y

variables, we used the same cubic (i.e., $order = 4$) B-spline basis functions for $M_i(l)$. We

considered $K_M = \{4, \dots, 8\}$ for all genes.

For each gene, we simulated one 488×1 vector for binary exposure $\mathbf{X}$ via the model:

$$logit\big(P(\mathbf{X} = 1)\big) = \mathbf{M}\boldsymbol{\alpha}_X^*$$

where $\mathbf{M}$ included 15% consecutive CpG sites covering the longest distance in the gene,

each $\alpha_X^* = 1.2/m_{15\%}$ and $m_{15\%}$ is the number of 15% CpG sites in the gene. The purpose

is to make the exposure and the mediator (i.e., methylation) associated. Then, we can

evaluate the power of testing $\alpha_X = 0$ in Formula (3).

For each gene, we then simulated continuous and binary outcomes $\mathbf{y}$ for 488 samples via

the models:

$$\mathbf{y} = \beta_0 + \mathbf{Z}_1\beta_1 + \mathbf{Z}_2\beta_2 + \mathbf{X}\beta_X + \mathbf{M}\boldsymbol{\beta}_M + \mathbf{X}\mathbf{M}\boldsymbol{\beta}_{XM} + \boldsymbol{\varepsilon} \text{ and}$$

$$logit\big(P(\mathbf{y} = 1)\big) = \beta_{0^*} + \mathbf{Z}_1\beta_1 + \mathbf{Z}_2\beta_2 + \mathbf{X}\beta_X + \mathbf{M}\boldsymbol{\beta}_M + \mathbf{X}\mathbf{M}\boldsymbol{\beta}_{XM}$$

where $\mathbf{Z}_1$, $\mathbf{Z}_2$, $\mathbf{X}$ and $\mathbf{M}$ were the same as described above, $\boldsymbol{\varepsilon}$ was generated from

standard normal distribution, $\beta_0 = 0.1$, $\beta_1 = -0.1$, $\beta_2 = 0.1$ and $\beta_{0^*} = -1$. We considered

(1) all $m_{15\%}$ effective CpG sites in the same direction (100%+) and (2) 50% effective CpG

sites in one direction and the other 50% effective CpG sites in the opposite direction

(50%+/50%-). We then considered eight settings for β_X, $\boldsymbol{\beta}_M$ and $\boldsymbol{\beta}_{XM}$ to simulate **y**, and

these eight settings (Table 1) were used to evaluate Type I error rates and power of

testing DE, IE and TE with the null hypotheses described in Formula (5). As shown in

Table 1, Setting (1) was used to assess Type I error rates of testing DE, IE and TE; Setting

(2) was used to assess Type I error rates of testing DE and power of testing IE and TE;

Setting (3) was used to assess Type I error rates of testing IE and power of testing DE

and TE; and Settings (4) ~ (8) were used to assess power of testing DE, IE and TE.

Please note that Setting (2) may not be the true null hypothesis for DE. Although $\beta_X = 0$,

there could still be implicit **X** effect, because **X** and **M** are associated, and $\boldsymbol{\beta}_M \neq 0$.

Analogously, Setting (3) may not be the true null hypothesis for IE.

Table 1. The eight settings for β_X, $\boldsymbol{\beta}_M$ and $\boldsymbol{\beta}_{XM}$ and their corresponding null hypothesis
(H_0) and alternative hypothesis (H_a) for DE, IE and TE

	Settings*				DE	IE	TE
	Continuous y		Binary y				
	100%+	50%+/50%-	100%+	50%+/50%-			
(1)	$\beta_X = \boldsymbol{\beta}_M = \boldsymbol{\beta}_{XM} = 0$		$\beta_X = \boldsymbol{\beta}_M = \boldsymbol{\beta}_{XM} = 0$		H_0	H_0	H_0
(2)	$\beta_X = \boldsymbol{\beta}_{XM} = 0, \boldsymbol{\beta}_M = 0.2$	$\beta_X = \boldsymbol{\beta}_{XM} = 0, \boldsymbol{\beta}_M = 0.2/-0.2$	$\beta_X = \boldsymbol{\beta}_{XM} = 0, \boldsymbol{\beta}_M = 0.4$	$\beta_X = \boldsymbol{\beta}_{XM} = 0, \boldsymbol{\beta}_M = 0.4/-0.4$	$H_0{}^{\#}$	H_a	H_a
(3)	$\boldsymbol{\beta}_M = \boldsymbol{\beta}_{XM} = 0, \beta_X = 0.2$	$\boldsymbol{\beta}_M = \boldsymbol{\beta}_{XM} = 0, \beta_X = 0.2/-0.2$	$\boldsymbol{\beta}_M = \boldsymbol{\beta}_{XM} = 0, \beta_X = 0.4$	$\boldsymbol{\beta}_M = \boldsymbol{\beta}_{XM} = 0, \beta_X = 0.4/-0.4$	H_a	$H_0{}^{\#}$	H_a
(4)	$\beta_X = \boldsymbol{\beta}_M = 0, \boldsymbol{\beta}_{XM} = 0.2$	$\beta_X = \boldsymbol{\beta}_M = 0, \boldsymbol{\beta}_{XM} = 0.2/-0.2$	$\beta_X = \boldsymbol{\beta}_M = 0, \boldsymbol{\beta}_{XM} = 0.4$	$\beta_X = \boldsymbol{\beta}_M = 0, \boldsymbol{\beta}_{XM} = 0.4/-0.4$	H_a	H_a	H_a
(5)	$\beta_X = \boldsymbol{\beta}_{XM} = 0.2, \boldsymbol{\beta}_M = 0$	$\beta_X = \boldsymbol{\beta}_{XM} = 0.2/-0.2, \boldsymbol{\beta}_M = 0$	$\beta_X = \boldsymbol{\beta}_{XM} = 0.4, \boldsymbol{\beta}_M = 0$	$\beta_X = \boldsymbol{\beta}_{XM} = 0.4/-0.4, \boldsymbol{\beta}_M = 0$	H_a	H_a	H_a
(6)	$\boldsymbol{\beta}_M = \boldsymbol{\beta}_{XM} = 0.2, \beta_X = 0$	$\boldsymbol{\beta}_M = \boldsymbol{\beta}_{XM} = 0.2/-0.2, \beta_X = 0$	$\boldsymbol{\beta}_M = \boldsymbol{\beta}_{XM} = 0.4, \beta_X = 0$	$\boldsymbol{\beta}_M = \boldsymbol{\beta}_{XM} = 0.4/-0.4, \beta_X = 0$	H_a	H_a	H_a
(7)	$\beta_X = \boldsymbol{\beta}_M = 0.2, \boldsymbol{\beta}_{XM} = 0$	$\beta_X = \boldsymbol{\beta}_M = 0.2/-0.2, \boldsymbol{\beta}_{XM} = 0$	$\beta_X = \boldsymbol{\beta}_M = 0.4, \boldsymbol{\beta}_{XM} = 0$	$\beta_X = \boldsymbol{\beta}_M = 0.4/-0.4, \boldsymbol{\beta}_{XM} = 0$	H_a	H_a	H_a
(8)	$\beta_X = \boldsymbol{\beta}_M = \boldsymbol{\beta}_{XM} = 0.2$	$\beta_X = \boldsymbol{\beta}_M = \boldsymbol{\beta}_{XM} = 0.2/-0.2$	$\beta_X = \boldsymbol{\beta}_M = \boldsymbol{\beta}_{XM} = 0.4$	$\beta_X = \boldsymbol{\beta}_M = \boldsymbol{\beta}_{XM} = 0.4/-0.4$	H_a	H_a	H_a

255 * $\boldsymbol{\beta}_M$ and $\boldsymbol{\beta}_{XM}$ are vectors for 15% consecutive CpG sites. For 100%+, each element has the same
256 assigned value. For 50%+/50%-, 50% elements have the same positive value and 50% elements have the
257 same negative value. If the number of CpG sites is odd, one CpG site is assigned to have no effect.
258
Setting (2) may not be the true null hypothesis for DE. Although $\beta_X = 0$, there could still be implicit **X**
effect, because **X** and **M** are associated, and $\boldsymbol{\beta}_M \neq 0$. Analogously, Setting (3) may not be the true null
hypothesis for IE.

For each gene, we simulated 10 sets of **y** for genes with 10 CpG sites (total is

705×10=7,050); 30 sets of **y** for genes with 15 CpG sites (total is 306×30=9,180); 50 sets

of **y** for genes with 20 CpG sites (total is 152×50=7,600); and 400 sets of **y** for genes with

40 CpG sites (total is 21×400=8,400). Thus, we simulated 32,230 datasets for continuous

outcome and another 32,230 datasets for binary outcome. After simulating the exposure

X and outcome **y** variables, we used two approaches to conduct the analyses: (1) basis

function fitted regional methylation level as the mediator (i.e., MR-Mediation) and (2)

mean CpG methylation level as the mediator. We assessed the Type I error rates of

testing DE, IE and TE only when the p-values of α_X were less than 0.05, and we evaluated

the power at the significance level that the p-values of the corresponding effect (i.e., DE,

IE or TE) and α_X were both less than 0.05.

RESULTS

Simulation of the Type I error rate

For both continuous and binary outcomes, both approaches retained the desired Type I

error rates for DE, IE and TE across all considered gene sizes and basis numbers in

Setting (1) (Figures 2, S3-S5 and S8-S10). Since Setting (2) might not be the true null

hypothesis for DE, both MR-Mediation and the mean CpG methylation approach had

inflated Type I error rates for scenarios with 100%+ effective CpG sites (Figure S6 and

S11) and correct Type I error rates for scenarios with 50%+/50%- effective CpG sites

(Figure S13 and S15). Although Setting (3) might not be the true null hypothesis for IE,

both approaches retained the desired Type I error rates for all simulated conditions for

both continuous and binary outcomes (Figures S7, S12, S14 and S16). Thus, both

approaches are valid under their true null hypothesis.

Figure 2. QQ plot of the *p*-values from Setting (1) with (A) continuous outcome and (B) binary outcome. The 1st column shows DE results, the 2nd column shows IE results and the 3rd column shows TE results. Each row shows one gene size (i.e., 10, 15, 20 or 40 CpG sites). Two approaches were compared: (1) MR-Mediation; and (2) Mean CpG methylation approach. A 95% pointwise confidence band (gray area) was computed under the assumption that the *p*-values were drawn independently from a uniform [0, 1] distribution.

Simulation of power

In all considered scenarios, the power of MR-Mediation decreased as the basis number

increased when the gene size was relatively small, and remained the similar power as

the basis number increased when the gene size was large (Figures S17-S28). When the

basis number was 4, MR-Mediation was the most powerful method in all scenarios for

testing IE and more powerful than the mean CpG methylation method in almost all

scenarios for testing TE and DE (Figures 3 and 4). In real data studies, one could be more

interested in testing IE than TE and DE. Thus, MR-Mediation with the basis number equal

to 4 is recommended.

Figure 3. Power comparison for continuous outcome. (A) all positively effective CpG sites (100%+) and (B) 50% positively and 50% negatively effective CpG sites (50%+/50%-). The 1st column shows DE results, the 2nd column shows IE results and the 3rd column shows TE results. Each row shows one gene size (i.e., 10, 15, 20 or 40 CpG sites). Two approaches were compared: (1) MR-Mediation; and (2) Mean CpG methylation approach.

Figure 4. Power comparison for binary outcome. (A) all positively effective CpG sites (100%+) and (B) 50% positively and 50% negatively effective CpG sites (50%+/50%-). The 1st column shows DE results, the 2nd column shows IE results and the 3rd column shows TE results. Each row shows one gene size (i.e., 10, 15, 20 or 40 CpG sites). Two approaches were compared: (1) MR-Mediation; and (2) Mean CpG methylation approach.

Results in EVA-PR

We used the proposed MR-Mediation method to study whether the association between exposure to gun violence and atopic asthma was mediated by DNA methylation, where atopic asthma is a binary outcome. In this study, we considered 5,720 genes including at

least 10 CpG sties in 407 subjects without any missing covariates. Note that subjects who had asthma but no atopy were excluded from this analysis, as atopic asthma is the most common type of asthma in children. This analysis was adjusted for age, sex, annual household income (a measure of socioeconomic status), the top five principal components from genotypic data, methylation batch, and latent factors between exposure to gun violence and methylation, and between atopic asthma and methylation, estimated from *sva*.[38] The mediator regression with four cubic B-spline basis functions, Formula (3), was run only for controls to account for the case-control design and we selected the top 20 genes for the further mediation analysis. We focused on the IE test to see if the association between exposure to gun violence and atopic asthma was mediated by methylation. We found that *CFD* on chromosome 19 was the top gene associated with exposure to gun violence ($P = 1.93 \times 10^{-5}$ and *FDR* = 0.1102, Table S1), although it did not reach the genome-wide significance level, and then it could mediate the effect of exposure to gun violence on atopic asthma risk ($P = 3.09 \times 10^{-2}$). The *CFD* gene has been reported to be associated with asthma.[39] The expression of *CFD* can be reduced by IL-17A, which is required for development of airway hyperresponsiveness (a key intermediate phenotype of asthma). In addition, *TBC1D14* ($P = 7.28 \times 10^{-3}$) on chromosome 4, *FAM120B* ($P = 3.33 \times 10^{-2}$) on chromosome 6, *CRB1* ($P = 1.75 \times 10^{-2}$) on chromosome 10, *LOC339166* ($P = 1.42 \times 10^{-2}$) on chromosome 17, *ZGPAT* ($P = 2.55 \times 10^{-3}$) on chromosome 20, *MED16* ($P = 2.05 \times 10^{-2}$) on chromosome 19, *PLEKHG6* ($P = 1.24 \times 10^{-3}$) on chromosome 12, and *TNNI1* ($P = 8.85 \times 10^{-3}$) on chromosome 1 could also mediate the effect of exposure to gun violence on atopic asthma risk at the nominal level in the top 20 gene list.

We also applied MR-Mediation to study whether the association between exposure to

gun violence and total IgE was mediated by DNA methylation, where total IgE is a

continuous outcome. We also used 5,720 genes including at least 10 CpG sties in 473

subjects without any missing covariates. This analysis was adjusted for age, sex, annual

household income, the top five principal components from genotypic data, methylation

batch, and latent factors between exposure to gun violence and methylation, and between

total IgE and methylation, estimated from *sva*.[38] We selected the top 20 genes by *p*-values

from the association between exposure to gun violence and gene-based methylation

(Formula (3)) and conducted mediation analysis (Formula (4)). Again, we focused on the

IE test to see if the association between exposure to gun violence and total IgE was

mediated by methylation. The results showed that *PIP5K1C* (P = 4.33×10^{-3}) on

chromosome 19 could mediate the effect of exposure to gun violence on total IgE change

at the nominal level in the top 20 gene list (Table S2).

DISCUSSION

In this study, we developed a novel causal mediation approach, MR-Mediation, under the

counterfactual framework to test the significance of DE, IE and TE. We implemented MR-

Mediation in R (http://www.r-project.org) and the R package (https://cran.r-

project.org/web/packages/MRmediation/index.html) is available. The major advantage of

the counterfactual framework to mediation analysis is that it allows the decomposition of

TE into DE and IE, even in models with nonlinearities and interactions. The counterfactual

framework extends the widely used Baron and Kenny mediation approach by allowing an

interaction term between exposure and mediator in the outcome regression.

We focused on testing the significance of DE, IE and TE instead of estimating their effects.

For example, in IE, we test $H_0: \beta_M = \beta_{XM} = 0$. When we reject the null hypothesis, it is

possible that $\beta_M = 0$ and $\beta_{XM} \neq 0$, which could not be known from the proposed test.

Then the estimates of β_M may not be reasonable to be used in calculating IE. The

parameters of covariates even need to be significant to calculate the estimates of DE and

TE. In this approach, a group of CpG sites from a predefined region are utilized as the

mediator, and the functional transformation is used to reduce the possible high dimension

in the region-based CpG sites and account for their location information. The region-

based analysis methods can improve power by combining weak signals and by reducing

the multiple testing penalty. Users can use different ways to define regions of interest in

addition to genes used in this study, such as, CpG islands and CpG shores.

For binary outcomes, we used the rare disease assumption, so that the OR in the case-

control design is equal to the OR in the population, which is approximately equal to the

RR in the population. When the disease is common in the population, the OR does not

approximate the RR. In that case, a log link should be used in the generalized linear

model for binary outcomes instead of a logit link, but the log link often fails to work with a

binomial distribution. The Zou's modified Poisson regression[40] could be applied in this

situation, although this is an approximate approach.

In the simulation studies, we show that MR-Mediation retains the correct Type I error rate

for testing DE, IE and TE for both continuous and binary outcomes. In power comparison,

we are more interested in IE, and MR-Mediation with basis=4 achieves better power than

the mean CpG methylation approach. Our real data results show that MR-Mediation could

help to identify potential methylation mediated effects. The genome-wide data analysis

could be completed within an hour using one CPU.

Since MR-Mediation is a joint significance test, the exposure and regional methylation

relationship and the regional methylation and outcome relationship are tested in two steps.

It's possible that CpG sites associated with the exposure and CpG sites associated with

the outcome are different, which cannot be detected by the region-based mediator

method. A further mediation analysis using single CpG sites may be needed. The single

CpG site mediation function is also available in our R package. In addition, the

summarized methylation level in the region can be viewed as the area under the curve.

However, the areas could be still the same, even if curves are different. In future studies,

we will consider to automatically partition the area into several sub-areas and each sub-

area has a summarized methylation value. In such methods, $\boldsymbol{\alpha}_X$, $\boldsymbol{\beta}_M$ and $\boldsymbol{\beta}_{XM}$ are vectors

instead of scalars as in this study. It's possible that $\boldsymbol{\alpha}_X(\boldsymbol{\beta}_M + x_1\boldsymbol{\beta}_{XM}) = 0$, even if $\boldsymbol{\alpha}_X \neq 0$

and $\boldsymbol{\beta}_M \neq 0$ or $\boldsymbol{\beta}_{XM} \neq 0$. We will need to address this potential problem.

MR-Mediation can be extended to high dimensional predictors, such as multiple SNPs

from a region, so as to investigate DNA methylation-mediated genetic effects on an

outcome. These SNPs can be from the same region as the CpG sites. Such region-based

approach on both SNPs and CpG sites can boost statistical power, because the region-

based approach combines small individual effects and reduce multiple testing. Moreover,

the region-based CpG sites can also be treated as predictors and gene expression can

be treated as a mediator in order to understand the regulatory mechanisms of DNA

methylation on gene expression that further affect diseases or traits.

Disclosure of Potential Conflict of Interest

No potential conflicts of interest were disclosed.

Supplemental Material

Supplemental data for this article can be accessed on the publisher's website.

REFERENCES

1.   Laurent L, Wong E, Li G, Huynh T, Tsirigos A, Ong CT, Low HM, Kin Sung KW, Rigoutsos I, Loring J, et al. Dynamic changes in the human methylome during differentiation. Genome Res 2010; 20:320-31.

2.   Portela A, Esteller M. Epigenetic modifications and human disease. Nat Biotechnol 2010; 28:1057-68.

3.   Melotte V, Lentjes MH, van den Bosch SM, Hellebrekers DM, de Hoon JP, Wouters KA, Daenen KL, Partouns-Hendriks IE, Stessels F, Louwagie J, et al. N-Myc downstream-regulated gene 4 (NDRG4): a candidate tumor suppressor gene and potential biomarker for colorectal cancer. J Natl Cancer Inst 2009; 101:916-27.

4.   Schmidt B, Liebenberg V, Dietrich D, Schlegel T, Kneip C, Seegebarth A, Flemming N, Seemann S, Distler J, Lewin J, et al. SHOX2 DNA methylation is a biomarker for the diagnosis of lung cancer based on bronchial aspirates. BMC Cancer 2010; 10:600.

5.   Jain S, Chen S, Chang KC, Lin YJ, Hu CT, Boldbaatar B, Hamilton JP, Lin SY, Chang TT, Chen SH, et al. Impact of the location of CpG methylation within the GSTP1 gene on its specificity as a DNA marker for hepatocellular carcinoma. PLoS One 2012; 7:e35789.

6.   Lord J, Cruchaga C. The epigenetic landscape of Alzheimer's disease. Nat Neurosci 2014; 17:1138-40.

7.   De Jager PL, Srivastava G, Lunnon K, Burgess J, Schalkwyk LC, Yu L, Eaton ML, Keenan BT, Ernst J, McCabe C, et al. Alzheimer's disease: early alterations in brain DNA methylation at ANK1, BIN1, RHBDF2 and other loci. Nat Neurosci 2014; 17:1156-63.

8.   Lunnon K, Smith R, Hannon E, De Jager PL, Srivastava G, Volta M, Troakes C, Al-Sarraj S, Burrage J, Macdonald R, et al. Methylomic profiling implicates cortical deregulation of ANK1 in Alzheimer's disease. Nat Neurosci 2014; 17:1164-70.

9.   Pidsley R, Viana J, Hannon E, Spiers H, Troakes C, Al-Saraj S, Mechawar N, Turecki G, Schalkwyk LC, Bray NJ, et al. Methylomic profiling of human brain tissue supports a neurodevelopmental origin for schizophrenia. Genome Biol 2014; 15:483.

10.   Jaffe AE, Gao Y, Deep-Soboslay A, Tao R, Hyde TM, Weinberger DR, Kleinman JE. Mapping DNA methylation across development, genotype and schizophrenia in the human frontal cortex. Nat Neurosci 2016; 19:40-7.

11.   Pedersen BS, Schwartz DA, Yang IV, Kechris KJ. Comb-p: software for combining, analyzing, grouping and correcting spatially correlated P-values. Bioinformatics 2012; 28:2986-8.

12.   Butcher LM, Beck S. Probe Lasso: a novel method to rope in differentially methylated regions with 450K DNA methylation data. Methods 2015; 72:21-8.

13.   Eckhardt F, Lewin J, Cortese R, Rakyan VK, Attwood J, Burger M, Burton J, Cox TV, Davies R, Down TA, et al. DNA methylation profiling of human chromosomes 6, 20 and 22. Nat Genet 2006; 38:1378-85.

14.   Irizarry RA, Ladd-Acosta C, Carvalho B, Wu H, Brandenburg SA, Jeddeloh JA, Wen B, Feinberg AP. Comprehensive high-throughput arrays for relative methylation (CHARM). Genome Res 2008; 18:780-90.

15.   Jaenisch R, Bird A. Epigenetic regulation of gene expression: how the genome integrates intrinsic and environmental signals. Nat Genet 2003; 33 Suppl:245-54.

16. Irizarry RA, Ladd-Acosta C, Wen B, Wu Z, Montano C, Onyango P, Cui H, Gabo
K, Rongione M, Webster M, et al. The human colon cancer methylome shows similar
hypo- and hypermethylation at conserved tissue-specific CpG island shores. Nat Genet
2009; 41:178-86.
17. Lister R, Pelizzola M, Dowen RH, Hawkins RD, Hon G, Tonti-Filippini J, Nery JR,
Lee L, Ye Z, Ngo QM, et al. Human DNA methylomes at base resolution show widespread
epigenomic differences. Nature 2009; 462:315-22.
18. Baron RM, Kenny DA. The moderator-mediator variable distinction in social
psychological research: conceptual, strategic, and statistical considerations. Journal of
personality and social psychology 1986; 51:1173-82.
19. Imai K, Keele L, Tingley D. A general approach to causal mediation analysis.
Psychol Methods 2010; 15:309-34.
20. Robins JM, Greenland S. Identifiability and exchangeability for direct and indirect
effects. Epidemiology 1992; 3:143-55.
21. Vanderweele TJ, Vansteelandt S. Odds ratios for mediation analysis for a
dichotomous outcome. Am J Epidemiol 2010; 172:1339-48.
22. Bozack AK, Cardenas A, Quamruzzaman Q, Rahman M, Mostofa G, Christiani DC,
Kile ML. DNA methylation in cord blood as mediator of the association between prenatal
arsenic exposure and gestational age. Epigenetics 2018; 13:923-40.
23. Cardenas A, Lutz SM, Everson TM, Perron P, Bouchard L, Hivert MF. Mediation
by Placental DNA Methylation of the Association of Prenatal Maternal Smoking and Birth
Weight. Am J Epidemiol 2019; 188:1878-86.
24. Barfield R, Shen J, Just AC, Vokonas PS, Schwartz J, Baccarelli AA, VanderWeele
TJ, Lin X. Testing for the indirect effect under the null for genome-wide mediation
analyses. Genet Epidemiol 2017; 41:824-33.
25. Tobi EW, Slieker RC, Luijk R, Dekkers KF, Stein AD, Xu KM, Biobank-based
Integrative Omics Studies C, Slagboom PE, van Zwet EW, Lumey LH, et al. DNA
methylation as a mediator of the association between prenatal adversity and risk factors
for metabolic disease in adulthood. Sci Adv 2018; 4:eaao4364.
26. Yang CF, Karmaus WJJ, Yang CC, Chen ML, Wang IJ. Bisphenol a Exposure,
DNA Methylation, and Asthma in Children. Int J Environ Res Public Health 2020; 17.
27. Neophytou AM, Oh SS, Hu D, Huntsman S, Eng C, Rodriguez-Santana JR, Kumar
R, Balmes JR, Eisen EA, Burchard EG. In utero tobacco smoke exposure, DNA
methylation, and asthma in Latino children. Environ Epidemiol 2019; 3:e048.
28. Wu D, Yang H, Winham SJ, Natanzon Y, Koestler DC, Luo T, Fridley BL, Goode
EL, Zhang Y, Cui Y. Mediation analysis of alcohol consumption, DNA methylation, and
epithelial ovarian cancer. J Hum Genet 2018; 63:339-48.
29. Battram T, Richmond RC, Baglietto L, Haycock PC, Perduca V, Bojesen SE, Gaunt
TR, Hemani G, Guida F, Carreras-Torres R, et al. Appraising the causal relevance of
DNA methylation for risk of lung cancer. Int J Epidemiol 2019; 48:1493-504.
30. Fan R, Wang Y, Mills JL, Wilson AF, Bailey-Wilson JE, Xiong M. Functional linear
models for association analysis of quantitative traits. Genet Epidemiol 2013; 37:726-42.
31. Forno E, Wang T, Qi C, Yan Q, Xu CJ, Boutaoui N, Han YY, Weeks DE, Jiang Y,
Rosser F, et al. DNA methylation in nasal epithelium, atopy, and atopic asthma in children:
a genome-wide study. Lancet Respir Med 2019; 7:336-46.
32. Ramsay JO, Silverman BW. Functional Data Analysis. Springer, New York 1996.

33.   Valeri L, Vanderweele TJ. Mediation analysis allowing for exposure-mediator
interactions and causal interpretation: theoretical assumptions and implementation with
SAS and SPSS macros. Psychol Methods 2013; 18:137-50.
34.   Sternthal MJ, Jun HJ, Earls F, Wright RJ. Community violence and urban
childhood asthma: a multilevel analysis. Eur Respir J 2010; 36:1400-9.
35.   Suglia SF, Ryan L, Wright RJ. Creation of a community violence exposure scale:
accounting for what, who, where, and how often. Journal of traumatic stress 2008;
21:479-86.
36.   Thomson CC, Roberts K, Curran A, Ryan L, Wright RJ. Caretaker-child
concordance for child's exposure to violence in a preadolescent inner-city population.
Arch Pediatr Adolesc Med 2002; 156:818-23.
37.   Ramratnam S, Han Y, Rosas-Salazar C, Forno E, Brehm J, Rosser F, Marsland
A, Colón-Semidey A, Alvarez M, Miller G, et al. Exposure to gun violence and asthma
among children in Puerto Rico. Respir Med 2015; In Press.
38.   Leek JT, Johnson WE, Parker HS, Jaffe AE, Storey JD. The sva package for
removing batch effects and other unwanted variation in high-throughput experiments.
Bioinformatics 2012; 28:882-3.
39.   Mathews JA, Wurmbrand AP, Ribeiro L, Neto FL, Shore SA. Induction of IL-17A
Precedes Development of Airway Hyperresponsiveness during Diet-Induced Obesity and
Correlates with Complement Factor D. Front Immunol 2014; 5:440.
40.   Zou G. A modified poisson regression approach to prospective studies with binary
data. Am J Epidemiol 2004; 159:702-6.