THE CORRELATION OF SUBSTITUTION EFFECTS ACROSS POPULATIONS AND GENERATIONS IN

THE PRESENCE OF NON-ADDITIVE FUNCTIONAL GENE ACTION

A. Legarra*, C.A. Garcia-Baccino*[,†], Y.C.J. Wientjes[‡], , Z.G. Vitezica*

* INRAE/INP, UMR 1388 GenPhySE, 31326, Castanet-Tolosan, France.

[†] Departamento de Producción Animal, Facultad de Agronomía, Universidad de Buenos

Aires, C1417DSQ, Buenos Aires, Argentina.

[‡] Wageningen University & Research, Animal Breeding and Genomics, 6700 AH Wageningen,

The Netherlands.

Keywords: QTL, substitution effects, epistasis, dominance, genetic distance

Corresponding author:

Andres Legarra,

andres.legarra@inrae.fr

+33 561285182

INRAE, UMR1388 GenPhySE

CS 52627

31326 Castanet Tolosan, France

ABSTRACT

Allele substitution effects at quantitative trait loci (QTL) are part of the basis of quantitative

genetics theory and applications such as association analysis and genomic prediction. In

presence of non-additive functional gene action, substitution effects are not constant across

populations. We develop an original approach to model the difference in substitution effects

across populations as first order Taylor series expansion from a "focal" population. This

expansion involves the difference in allele frequencies and second-order statistical effects

(additive by additive and dominance). The change in allele frequencies is a function of

relationships (or genetic distances) across populations. As a result, it is possible to estimate

the correlation of substitution effects across two populations using three elements:

magnitudes of additive, dominance and additive by additive variance; relationships across

populations (similar to Fst indexes); and functions of heterozygosities at the markers.

Similarly, the theory applies as well to distinct generations in a population, in which case the

distance across generations is a function of increase of inbreeding. Using published

estimates of the needed parameters, we estimate the correlation between substitution

effects to be around 0.60 for distinct breeds and higher than 0.9 for generations closer than

5. Simulation results confirmed our derivations although our estimators tended to

underestimate the correlation. Our derivations are useful to understand the difficulty in

predicting across populations, to forecast the possibility of prediction across populations,

and we suggest that they can be useful to disentangle genotype by environment interaction

from genotype by genotype interaction.

                                     INTRODUCTION

One of the aims of quantitative genetics is to provide methods for prediction, for instance

genomic prediction (prediction of livestock breeding values or of crop performance) or

polygenic risk score (risk of a disease in humans). These predictions would ideally work

across a range of populations (different breeds, future generations). Ideally, the prediction

goes through a process of identifying causal genes, estimating their effects in some

population, and transposing these effects to newly genotyped individuals (Lande and

Thompson 1990; Meuwissen *et al.* 2001). These "gene effects" are substitution effects –

loosely speaking, the regression of the own phenotype (for polygenic risk scores) or

expected progeny phenotypes (for estimated breeding values) on gene content at the locus.

Being able to use substitution effects at causal genes across populations and generations is

the ultimate goal of genomic prediction, QTL detection and to some extent of molecular

biology description of gene effects.

There are several obstacles for these aims. Finding and validating causal genes and

understanding their functional mechanism is extremely difficult (Grobet *et al.* 1997; Bonifati

et al. 2003; Rupp *et al.* 2015). In practice, predictions are done using SNP markers using

statistical genetics techniques. Use of markers results in very good predictions within

population but mediocre (at best) predictions across populations, even with very

sophisticated techniques (Hayes *et al.* 2009; Karoui *et al.* 2012; Porto-Neto *et al.* 2015;

MacLeod *et al.* 2016). Indeed, livestock and human genetics empirical results show

decreasing predictive ability with increasing genetic distance across distinct populations or

generations (Liu *et al.* 2016; Martin *et al.* 2019). The lack of perfect linkage disequilibrium

across markers and genes has been claimed to be a reason for this decrease in accuracy.

Adding extra information (more dense maps, biological prior information) should result in a

better choice of markers close to causal genes, and therefore in boosting predictive abilities

across populations (Roos *et al.* 2009; MacLeod *et al.* 2016). However, in practice, the

increase in predictive ability across populations is small at best (MacLeod *et al.* 2016;

Moghaddar *et al.* 2019). This is against results in simulations (Roos *et al.* 2009; MacLeod *et*

al. 2016) – but a problem in simulations is that typically gene effects are assumed to be

biologically additive and therefore constant across populations.

In fact, substitution (statistical) additive gene action is not homogeneous across populations,

even for exactly the same mutation. Examples in livestock genetics include myostatin gene

(Aiello *et al.* 2018) or DGAT1 (Gautier *et al.* 2007). Although part of these differences may be

due to genotype by environment interactions, it is also plausible that this is due to epistasis

or dominance; for instance this is the case in DGAT1 (Streit *et al.* 2011). For instance, in the

5-loci epistatic network in Carlborg et al. (2006) some substitution effects at genes switch

signs depending on genetic background. In Drosophila, estimated (GWAS) substitution

effects switched signs depended on the genetic background (Magwire *et al.* 2010; Mackay

2015).

There is indeed widespread evidence of biological epistasis (Mackay 2014). Whereas

biological epistasis does not impede (on the contrary) large additive variation (Hill *et al.*

2008; Mackay 2014; Mäki-Tanila and Hill 2014), it implies is that substitution effects do vary

across genetic backgrounds. Thus, in the presence of functional dominance and epistasis,

there is no stability of substitution effects across different genetic backgrounds. Even if in all

of these populations, additive variation accounts for most genetic variation, and additive

substitution effects are sizable (Hill *et al.* 2008), substitution effects may differ across

populations.

Recent simulations (Dai *et al.* 2020; Duenk *et al.* 2020) showed that the difference in

substitution effects across populations may be quite large under non-additive biological

gene action and increases with divergence of populations. It is relatively easy to derive

algebraic expressions for substitution effects α assuming specific hypothesis of biological

gene action. For instance, assuming biological additive and dominance effects only (Falconer

and Mackay 1996) results in $\alpha = a + (q - p)d$, whereas assuming additive, dominance and

additive by additive biological gene effects results in $\alpha_1 = a_1 + (q_1 + p_1)d_1 + (p_2 -$

$q_2)[aa]_{12}$ (Fuerst *et al.* 1997). Both of the approaches (simulation or analytical) are limited

because the hypotheses of specific biological gene actions (*e.g.* 2-loci interaction but not 3-

loci interaction) are too restrictive. Classically, these hypotheses are bypassed in quantitative

genetics by working on statistical effects (Falconer and Mackay 1996; Lynch and Walsh 1998;

Mäki-Tanila and Hill 2014).

The key parameter to describe the resemblance of statistical effects across populations is

the correlation of substitution effects across populations. Under certain assumptions

(independence of allele frequencies and substitution effects, appropriate coding; (Wientjes

et al. 2017)), this correlation can be estimated from SNP markers and data of two

populations (Karoui *et al.* 2012; Wientjes *et al.* 2017), and can easily be accommodated into

genomic prediction models (Karoui *et al.* 2012; Xiang *et al.* 2017). In addition to empirical

results, some theory to describe the resemblance of substitution effects across different

populations would be helpful to (1) better understand and quantify the change in *true*

substitution effects (*i.e.* if true genes instead of markers were being used), (2) give upper

bounds for genomic prediction across populations/generations, and (3) allow *a priori*

planning of genomic predictions *i.e.* including or not different subpopulations.

The aim of this work is to develop a theory to predict the extent of change of substitution

effects across space (breeds, lines) and across time (generations), without invoking or

assuming specific modes of biological gene action. We obtain explicit estimators that are

functions of additive, dominance and additive by additive variances, genetic distances across

populations, and distributions of allele frequencies. We check and illustrate our theory using

published results and simulations considering dominance and epistasis (additive by additive

and complementary) from 5-loci interactions.

<div align="center">MATERIALS AND METHODS</div>

Theory

General Theory. Here, we model difference of substitution effects across populations as

Taylor expansions around a "focal" population. Using Kojima's method (Kojima 1959, 1961),

we put additive substitution effects as a function of differences in allele frequencies across

populations, "focal" additive substitution effects, and higher order (dominance, additive by

additive epistasis) statistical effects. From here, we show that the correlation of substitution

effects across populations is a function of their relationship (or genetic distance), the

additive, dominant and additive by additive genetic variances, the average heterozygosities

and average squared heterozygosities. All these parameters can be estimated in real

populations. In the following, we try to stick to Mäki-Tanila and Hill (2014) notation. Many

details are given as Appendix in the Supplementary Material.

Note that our procedure is general – it does not invoke neither any particular mechanism for

epistasis or dominance, nor knowledge of individual QTL effects and locations. We assume

that the population mean is a (possibly complex) function of QTL allele frequencies $\boldsymbol{p}$ and

QTL functional or biological effects. The latter, albeit unknown, are assumed to be constant

across populations - we therefore assume no genotype by environment interactions.

Consider the correlation $r\left(\alpha_i^b, \alpha_i^{b'}\right)$ of substitution effects $\alpha^{(b)}$ and $\alpha^{(b')}$ across respective

populations (breeds, heterotic groups, lines, or generations) b and b'. It is known that, in

presence of dominant and epistatic biological interactions, the value of α depends on the

allele   frequencies   and   biological   effects;   for   instance, $\alpha_1 = a_1 + (q_1 + p_1)d_1 +$

$(p_2 - q_2)[aa]_{12}$ for 2-locus epistasis (Fuerst $et\ al.$ 1997), for $p_1 = 1 - q_1$ and $p_2 = 1 - q_2$

frequencies at loci 1 and 2, respectively. Inspired by this, we want to write the correlation

$r\left(\alpha_i^b, \alpha_i^{b'}\right)$ for a locus i as a function of respective allele frequencies, $\boldsymbol{p}^b$ and $\boldsymbol{p}^{b'}$, but in a

general manner, without defining a particular functional or biological gene action. We use a

first order Taylor series expansion to approximate additive substitution effects in a

population b and in population b', as a function of effects in another "focal" population (that

we call "0") and the differences in allelic frequencies across the three populations. By doing

this, it can be shown that the difference of substitution effects between two populations is

(approximately) a function of the genetic distance of the two populations to the focal

population, and the magnitude of dominance and second order epistatic variances in the

focal population.

To derive additive substitution effects α as function of allelic frequencies $\boldsymbol{p}$, we use Kojima's

definition of statistical effects as first, second… derivatives of the mean of the population (μ)

as a function of p . We don't invoke any explicit function – we just presume that there is

one, in other words, change in allele frequencies of the population implies change in the

total average genotypic value. Using Kojima's method, the additive substitution effect at the

i-th locus is the first derivative (Kojima 1959, 1961):

$$\alpha_i = \frac{1}{2}\frac{\partial \mu}{\partial p_i}$$

180   Higher order effects implying locus *i* are higher order derivatives of μ but we remark that

181   they can equally be obtained as derivatives of α_i. The dominance deviation at the *i*-th locus

182   (that we denote as d^* to distinguish from the biological or functional effect d (Falconer and

183   Mackay 1996)) is:

$$d_i^* = -\frac{1}{4}\frac{\partial^2 \mu}{\partial p_i^2} = -\frac{1}{2}\frac{\partial \alpha_i}{\partial p_i}$$

185   The negative sign comes because the dominance deviation is usually understood as a feature

186   of *heterozygosity,* in other words, it is of opposite sign than the increase of homozygosity in

187   ∂p_i^2. Last, the epistatic pairwise deviation of locus i with j is

$$(\alpha\alpha)_{ij} = \frac{1}{4}\frac{\partial^2 \mu}{\partial p_i \partial p_j} = \frac{1}{2}\frac{\partial \alpha_i}{\partial p_j}$$

This is positive because it is the effect of increasing both p_i and p_j.

Kojima's method shows, therefore, that higher order effects of one locus are derivatives of

lower order effects. With these elements we can make a Taylor order expansion of

α_i around frequencies in the focal population, $\boldsymbol{p_0}$, so that $\boldsymbol{p} = \boldsymbol{p_0} + \boldsymbol{\epsilon}$, such that we create a

function $\alpha_i(\boldsymbol{p})$. A meaningful value for $\boldsymbol{p_0}$ is a "focal" population for which there are

estimates of additive, dominance and second order epistatic variances. In the Appendix, we

show that the Taylor linear approximation of the substitution effects α_i^b and $\alpha_i^{b'}$ for two

populations b and b' is:

$$\alpha_i^b \approx \alpha_i^0 + 2\epsilon_i^b(-d_i^{*0}) + 2\boldsymbol{\epsilon^{(b)}}'(\boldsymbol{\alpha\alpha})_i^0 \qquad (1)$$

$$\alpha_i^{b'} \approx \alpha_i^0 + 2\epsilon_i^{b'}(-d_i^{*0}) + 2\boldsymbol{\epsilon^{(b')}}'(\boldsymbol{\alpha\alpha})_i^0$$

199   where we use differences in allele frequencies from the focal population, $\boldsymbol{\epsilon^{(b)}} = \boldsymbol{p^{(b)}} - \boldsymbol{p_0}$

200   and $\boldsymbol{\epsilon^{(b')}} = \boldsymbol{p^{(b')}} - \boldsymbol{p_0}$, d_i^{*0} is the statistical dominance deviation at the locus i and $(\boldsymbol{\alpha\alpha})_i^0$ is

201   a vector containing epistatic substitution effects of locus i with the rest of loci.

203   From equation 1, the covariance across two populations b and b' of the two substitution

204   effects α_i^b and $\alpha_i^{b'}$ of the locus i is a function of the covariance of $\boldsymbol{\epsilon^{(b)}}$ and $\boldsymbol{\epsilon^{(b')}}$ and is:

$$Cov\left(\alpha_i^{(b)}, \alpha_i^{(b')}\right) \tag{2}$$

$$= Cov\left(\alpha_i^0 + 2\epsilon_i^{(b)}(-d_i^{*0}) + 2\boldsymbol{\epsilon}^{(b)'}(\boldsymbol{\alpha\alpha})_i^0, \alpha_i^0\right.$$

$$\left. + 2\epsilon_i^{(b')}(-d_i^{*0}) + 2\boldsymbol{\epsilon}^{(b')'}(\boldsymbol{\alpha\alpha})_i^0\right)$$

$$= Var(\alpha_i^0) + 4Cov\left(\epsilon_i^{(b)}, \epsilon_i^{(b')}\right)Var(d_i^{*0})$$

$$+ 4tr\left(Cov\left(\boldsymbol{\epsilon}^{(b)}, \boldsymbol{\epsilon}^{(b')'}\right)Var((\boldsymbol{\alpha\alpha})_i^0)\right)$$

Where $Cov\left(\boldsymbol{\epsilon}^{(b)}, \boldsymbol{\epsilon}^{(b')'}\right)$ and $Var((\boldsymbol{\alpha\alpha})_i^0)$ are matrices. Note that we can drop terms

including e.g. $Cov(\alpha^0, (\boldsymbol{\alpha\alpha})_i^0)$ as the different statistical effects (additive dominant,

epistatic) are mutually orthogonal. Also, we assume $\boldsymbol{\epsilon}^{(b)}$ (change in allele frequencies) and

the different effects d_i^{*0} and $(\boldsymbol{\alpha\alpha})_i^0$ to be independent. The equation 2 has different terms

that are briefly presented here; more details are in the Appendix.

The variance of statistical additive, dominant and additive by additive effects are (Mäki-

Tanila and Hill 2014):

$$Var(\alpha_i^0) = \sigma_\alpha^{(0)2} = \frac{\sigma_A^2}{n\overline{H}}$$

$$Var(d^{*0}) = \sigma_d^{(0)2} = \frac{\sigma_D^2}{n\overline{H^2}}$$

$$Var((\alpha\alpha)_{i,j,j>i}^0) = \sigma_{(\alpha\alpha)}^{(0)2} = \frac{\sigma_{AA}^2}{0.5n(n-1)\overline{HH}} \approx 4\frac{\sigma_{AA}^2}{n^2\overline{HH}}$$

$$Var((\boldsymbol{\alpha\alpha})_i^0) = \boldsymbol{I}\otimes\sigma_{(\alpha\alpha)}^{(0)2} \approx 4\boldsymbol{I}\frac{\sigma_{AA}^2}{n^2\overline{HH}}$$

214 for n the number of QTL loci and using functions of heterozygosities:

$$\overline{H} = E\left(2p_i(1-p_i)\right)$$

$$\overline{H^2} = E\big(4p_i(1-p_i)p_i(1-p_i)\big)$$

$$\overline{HH} = E_{i>j}\Big(2p_i(1-p_i)2p_j(1-p_j)\Big) \approx \frac{1}{2}\overline{HH}$$

All variances and effects refer to a focal population (with sub/superindex "0") with allele

frequencies p^0 and effects α^0. Note that we assume HWE and LE. Of course, we don't know

allele frequencies or even numbers of true causal genes, but the first can be prudently

guessed and the later cancel out in the following.

A relationship coefficient across populations b and b' is $\gamma_{b,b'} = 8Cov\Big(p_i^{(b)}, p_i^{(b')}\Big)$; also,

$\gamma_{b,b'} = -4D^2$ where D^2 is the average square of the Euclidean distance (Robertson 1975;

Bonhomme *et al.* 2010; Garcia-Baccino *et al.* 2017). Similarly, we introduce a measure of

similarity $\gamma_{b,b'}^\epsilon$ (more details in the Appendix) such that

$$Cov\Big(\epsilon_i^{(b)}, \epsilon_i^{(b')}\Big) = \frac{1}{8}\gamma_{b,b'}^\epsilon = \frac{1}{8}\big(\gamma_{b,b'} - \gamma_{b,0} - \gamma_{b',0} + \gamma_{0,0}\big)$$

This is the remaining similarity of b and b' once similarity of both to "0" has been discounted

If the focal ("0") population is population b, then $Cov\Big(\epsilon_i^{(b')}, \epsilon_i^{(b')}\Big) = Var\Big(\epsilon_i^{(b')}\Big) =$

$\frac{1}{8}\big(\gamma_{b'} + \gamma_b - 2\gamma_{b,b'}\big)$. In fact, in this case $Cov\Big(\epsilon_i^{(b')}, \epsilon_i^{(b')}\Big) = \frac{1}{8}\big(\gamma_{b'} + \gamma_b - 2\gamma_{b,b'}\big)$ is

proportional to the F_{st} differentiation index across b and b' such that the pairwise

$F_{st} = \dfrac{\frac{1}{8}\big(\gamma_{b'} + \gamma_b - 2\gamma_{b,b'}\big)}{1 - \frac{1}{8}\big(\gamma_{b'} + \gamma_b + 2\gamma_{b,b'}\big)}$. This is obtained from the definition of F_{st} from average

relationships within and across populations (Caballero and Toro 2002; Caballero 2020) as

detailed in the Supplementary Material. The main difference between the F_{st} and our

relationship coefficients γ resides in the fact that F_{st} describes divergence, and is relative to

the set of populations being considered, whereas γ coefficients describe similarity, and are

not relative to any particular set of populations; for instance, they can be defined and

obtained for pairs of populations, to be later combined. Across all loci, $Cov\left(\boldsymbol{\epsilon}^{(b)}, \boldsymbol{\epsilon}^{(b')'}\right) =$

$\boldsymbol{I}\frac{1}{8}\gamma^{\epsilon}_{b,b'}$. Thus, the covariance of changes in allele frequencies is a function of relatedness

across populations or, equivalently, of their differentiation.

The term including $Cov\left(\boldsymbol{\epsilon}^{(b)}(\boldsymbol{\alpha\alpha})^0_i, \boldsymbol{\epsilon}^{(b')}(\boldsymbol{\alpha\alpha})^0_i\right)$ uses the covariance of quadratic forms

(Searle 1971) (eq. 58) and is detailed in the Appendix. It assumes uncorrelated $\boldsymbol{\epsilon}^{(b)}$ and

$(\boldsymbol{\alpha\alpha})^{05}_i$, and normality for both. It is customary in population genetics to assume normality

for small changes in allele frequencies (Lewontin and Krakauer 1973; Bonhomme *et al.*

2010).

Combining expressions above, equation (2) becomes

$$Cov\left(\alpha^b_i, \alpha^{b'}_i\right) = Var(\alpha^0_i) + 4Cov\left(\epsilon^{(b)}_i, \epsilon^{(b')}_i\right)Var(d^{*0}_i) \tag{3}$$

$$+ 4tr\left(Cov\left(\boldsymbol{\epsilon}^{(b)}, \boldsymbol{\epsilon}^{(b')'}\right)Var((\boldsymbol{\alpha\alpha})^0_i)\right)$$

$$= \frac{\sigma^2_A}{n\overline{H}} + 4\frac{\gamma^{\epsilon}_{b,b'}}{8}\frac{\sigma^2_D}{n\overline{H^2}} + 4n\frac{\gamma^{\epsilon}_{b,b'}}{8}4\frac{\sigma^2_{AA}}{n^2\overline{H}\overline{H}}$$

$$= \frac{\sigma^2_A}{n\overline{H}} + \frac{1}{2}\gamma^{\epsilon}_{b,b'}\frac{\sigma^2_D}{n\overline{H^2}} + 2\gamma^{\epsilon}_{b,b'}\frac{\sigma^2_{AA}}{n\overline{H}\overline{H}}$$

$$= \frac{1}{n}\left(\frac{\sigma^2_A}{\overline{H}} + \frac{1}{2}\gamma^{\epsilon}_{b,b'}\frac{\sigma^2_D}{\overline{H^2}} + 2\gamma^{\epsilon}_{b,b'}\frac{\sigma^2_{AA}}{\overline{H}\overline{H}}\right)$$

The quantities involved in Equation 3 are (1) the covariance in change of allelic frequencies $\boldsymbol{\epsilon}$

in b and b' (called $\gamma^{\epsilon}_{b,b'}$), which describes the similarity of populations b and b' using the

focal population as a reference, and (2) the variance of statistical additive, dominant and

additive by additive effects. We have used $4tr\left(Cov\left(\boldsymbol{\epsilon}^{(b)},\boldsymbol{\epsilon}^{(b')'}\right)Var((\boldsymbol{\alpha}\boldsymbol{\alpha})_i^0)\right) =$

$4tr\left(\left(\boldsymbol{I}\frac{1}{8}\gamma_{b,b'}^{\epsilon}\right)\left(4\boldsymbol{I}\frac{\sigma_{AA}^2}{n^2\overline{H}\overline{H}}\right)\right) = 4n\left(\left(\frac{1}{8}\gamma_{b,b'}^{\epsilon}\right)\left(4\frac{\sigma_{AA}^2}{n^2\overline{H}\overline{H}}\right)\right) = 2\gamma_{b,b'}^{\epsilon}\frac{\sigma_{AA}^2}{n\overline{H}\overline{H}}$

Whereas

$$Var(\alpha_i^b) = \frac{1}{n}\left(\frac{\sigma_A^2}{\overline{H}} + \frac{1}{2}\gamma_b^{\epsilon}\frac{\sigma_D^2}{H^2} + 2\gamma_b^{\epsilon}\frac{\sigma_{AA}^2}{\overline{H}\overline{H}}\right)$$

254 From here the correlation of α across populations is (factorizing out $\frac{1}{n}$)

$$r\left(\alpha_i^b,\alpha_i^{b'}\right) = \tag{4}$$

$$\frac{\frac{\sigma_A^2}{\overline{H}} + \frac{1}{2}\gamma_{b,b'}^{\epsilon}\frac{\sigma_D^2}{H^2} + 2\gamma_{b,b'}^{\epsilon}\frac{\sigma_{AA}^2}{\overline{H}\overline{H}}}{\sqrt{\left(\frac{\sigma_A^2}{\overline{H}} + \frac{1}{2}\gamma_b^{\epsilon}\frac{\sigma_D^2}{H^2} + 2\gamma_b^{\epsilon}\frac{\sigma_{AA}^2}{\overline{H}\overline{H}}\right)\left(\frac{\sigma_A^2}{\overline{H}} + \frac{1}{2}\gamma_{b'}^{\epsilon}\frac{\sigma_D^2}{H^2} + 2\gamma_{b'}^{\epsilon}\frac{\sigma_{AA}^2}{\overline{H}\overline{H}}\right)}}$$

256 It can already be seen that $\frac{\sigma_A^2}{\overline{H}}$ is both on the numerator and denominator. If the focal

257 population is population b, then $\gamma_b^{\epsilon} = \gamma_{b,b'}^{\epsilon} = 0$, but $\gamma_{b'}^{\epsilon} = \left(\gamma_{b'} + \gamma_b - 2\gamma_{b,b'}\right)$ (as an aside,

258 this is positive, which results in the correlation $r\left(\alpha_i^b,\alpha_i^{b'}\right)$ being positive), and Equation 4

259 becomes

$$r\left(\alpha_i^b,\alpha_i^{b'}\right) = \frac{\frac{\sigma_A^2}{\overline{H}}}{\sqrt{\frac{\sigma_A^2}{\overline{H}}\left(\frac{\sigma_A^2}{\overline{H}} + \frac{1}{2}\gamma_{b'}^{\epsilon}\frac{\sigma_D^2}{H^2} + 2\gamma_{b'}^{\epsilon}\frac{\sigma_{AA}^2}{\overline{H}\overline{H}}\right)}} = \frac{\sqrt{\frac{\sigma_A^2}{\overline{H}}}}{\sqrt{\frac{\sigma_A^2}{\overline{H}} + \frac{1}{2}\gamma_{b'}^{\epsilon}\frac{\sigma_D^2}{H^2} + 2\gamma_{b'}^{\epsilon}\frac{\sigma_{AA}^2}{\overline{H}\overline{H}}}}$$

261 Factorizing out $\overline{H}$ we arrive to the slightly clearer expression

$$r\left(\alpha_i^b, \alpha_i^{b'}\right) = \frac{\sqrt{\sigma_A^2}}{\sqrt{\sigma_A^2 + \gamma_{b'}^\epsilon \left(\frac{1}{2}\frac{\overline{H}}{\overline{H^2}}\, \sigma_D^2 + 2\frac{\sigma_{AA}^2}{\overline{H}}\right)}} \tag{5}$$

$$= \frac{\sqrt{\sigma_A^2}}{\sqrt{\sigma_A^2 + \left(\gamma_{b'} + \gamma_b - 2\gamma_{b,b'}\right)\left(\frac{1}{2}\frac{\overline{H}}{\overline{H^2}}\, \sigma_D^2 + 2\frac{\sigma_{AA}^2}{\overline{H}}\right)}}$$

which shows well that the correlation is a function of relationships across populations

$\left(\gamma_{b'} + \gamma_b - 2\gamma_{b,b'}\right)$ and weights of additive *vs.* non-additive variances.

Finally, our expression for $r\left(\alpha_i^b, \alpha_i^{b'}\right)$ includes values for the three main genetic variances

(which can be estimated from data), measures of genetic distances (which can be estimated

from data) and "good guesses" of the distribution of QTL effects – the latter can come either

via markers (sequence or chip SNP) or guesses (*i.e.*, U-shaped, flat, etc.).

Thus, the algorithm to estimate *a priori* the correlation of α across populations b and b',

$r\left(\alpha_i^b, \alpha_i^{b'}\right)$, is:

    1. Define a focal population with existing estimates of statistical additive, dominance,

and additive by additive variances (this focal population can actually be population b

or b')

2. Compute average heterozygosity $\overline{H}$ and average squared heterozygosity $\overline{H^2}$ in the

focal population, either from markers or from guesses.

3. Using markers, compute relationships γ within and across the three populations

4. Apply equation 4. If the focal population is population b or b', apply the equivalent

Equation 5.

Covariance across generations within one breed. The definition above of two populations is

general enough that we can consider any two populations, *e.g.* two breeds (Angus and

Hereford), two strains (New Zealand Holstein and US Holstein) or two generations or time

frames (*e.g.* animals born in 2000 vs. animals born in 2005, or animals born in 2000 vs. their

descendants). There is evidence that across-generations genetic correlation decreases with

time to values as low as 0.6 (Tsuruta *et al.* 2004; Haile-Mariam and Pryce 2015). As explained

before, the across-generation genetic correlation does not depend only on the across-breed

correlation of substitution effects, first because allele frequencies and substitution effects

might be correlated due to selection, second because of genotype by environment

interactions. Anyway, part of the across-generation genetic correlation can be accounted for

by our model, based on the evolution of average coancestry in the breed.

We will talk about "generations" but ideas apply to pedigrees with overlapping generations

as well. Consider that we called populations b and b' are animals born at time t_1 and t_2, with

$t_2 > t_1$. Allele frequencies at generation t are (for a single locus) $p_t = \frac{1}{2n_t}\mathbf{1}'\mathbf{z}$ where $\mathbf{z}$ has

gene content in the form {0,1,2}. The variance of p_t is $Var(p_t) = \frac{1}{4}\bar{A}_t 2pq = \frac{pq}{2}\bar{A}_t$ (because

$Var(\mathbf{z}) = \mathbf{A}_t 2pq$), where $\mathbf{A}$ is the pedigree-based relationship matrix and $\mathbf{A}_t$ the submatrix

corresponding to animals born at time t. Considering $\mathbf{A}^{(\gamma)}$ build with metafounders (Legarra

et al. 2015) then $Var(p_t) = \bar{A}_t^{(\gamma)}/8$. Note that if there is a single metafounder then

$\mathbf{A}^{(\gamma)} = \mathbf{A}\left(1 - \frac{\gamma}{2}\right) + \mathbf{1}\mathbf{1}'\gamma$. The covariance between two generations is, similarly,

$Cov(p_{t_1}, p_{t_2}) = \overline{\boldsymbol{A}}^{(\gamma)}_{t_1,t_2}/8$, so here $\overline{\boldsymbol{A}}^{(\gamma)}_{t_1,t_2}$ takes the role of the $\gamma_{b,b'}$ before. Leading to, if we

take t_1 as the focal population,

$$Cov\left(\epsilon_i^{(t_2)}, \epsilon_i^{(t_2)}\right) = \frac{1}{8}\left(\bar{A}^{(\gamma)}_{t_2} + \bar{A}^{(\gamma)}_{t_1} - 2\bar{A}^{(\gamma)}_{t_1,t_2}\right)$$

303   This involves average relationships within and across generations. From here, the

304   correlation of substitution effects across t_1 and t_2 is

$$r\left(\alpha_i^{t_2}, \alpha_i^{t_1}\right) = \frac{\sqrt{\sigma_A^2}}{\sqrt{\sigma_A^2 + \left(\overline{\boldsymbol{A}}_{(\gamma)t_2} + \overline{\boldsymbol{A}}_{(\gamma)t_1} - 2\overline{\boldsymbol{A}}_{(\gamma)t_1,t_2}\right)\left(\frac{1}{2}\frac{\bar{H}}{\bar{H}^2}\sigma_D^2 + 2\frac{\sigma_{AA}^2}{\bar{H}}\right)}} \tag{6}$$

Thus, the correlation of substitution effects decreases when there is large drift: high average

relationships within each generation combined with low relationship across. This is the case

for instance if parents of the next generation are very highly selected without restrictions in

future inbreeding, resulting in a considerable change in allele frequencies over the

generations.

Values of $\left(\overline{\boldsymbol{A}}_{(\gamma)t_2} + \overline{\boldsymbol{A}}_{(\gamma)t_1} - 2\overline{\boldsymbol{A}}_{(\gamma)t_1,t_2}\right)$ can be easily computed, even for large pedigrees,

using Colleau (2002). This can be approximated (see Appendix) as $\overline{\boldsymbol{A}}_{(\gamma)t_2} + \overline{\boldsymbol{A}}_{(\gamma)t_1} -$

$2\overline{\boldsymbol{A}}_{(\gamma)t_1,t_2} \approx (t_2 - t_1)\Delta F_{(\gamma)}$ for small values of $\Delta F_{(\gamma)}$ and steady increase of inbreeding. Note

that $\Delta F_{(\gamma)} = \left(1 - \frac{\gamma}{2}\right)\Delta F$ as $\boldsymbol{A}_{(\gamma)} = \boldsymbol{A}\left(1 - \frac{\gamma}{2}\right) + \mathbf{1}\mathbf{1}'\gamma$. This agrees with classical theory

(Falconer and Mackay 1996): the variance in change of allele frequencies from one

generation to the next is simply ΔF. Typical values of ΔF in livestock (rabbits, pigs, cattle or

sheep) are at most 0.01 per generation (Welsh *et al.* 2010; García-Ruiz *et al.* 2016; E. N.

Fernandez, J.P. Sanchez, R. Martinez, A. Legarra, M. Baselga 2017; Rodríguez-Ramilo *et al.*

2019), although this is potentially changing with genomic selection. Substituting in Equation

6 we obtain the correlation of substitution effects as a function of increase in inbreeding:

$$r\left(\alpha_i^{t_2}, \alpha_i^{t_1}\right) = \frac{\sqrt{\sigma_A^2}}{\sqrt{\sigma_A^2 + \Delta F_{(\gamma)} \left(\frac{1}{2} \frac{\bar{H}}{\bar{H}^2} \sigma_D^2 + 2 \frac{\sigma_{AA}^2}{\bar{H}} \right)}} \tag{7}$$

Examples

Estimates of correlation across populations using literature values. From values of the

literature, the previous expressions, equations (5) and (7) were applied to obtain guesses of

the correlation of substitution effects across different populations and generations within

population. Estimates of statistical non-additive variation are scarce, as are estimates of

distances across populations in terms of covariance of allele frequencies. Table 1 shows two

pairs of populations in pigs (Xiang *et al.* 2016) and cattle (VanRaden *et al.* 2011; Legarra *et*

al. 2015), and the corresponding relationships expressed as $\gamma_{b,b'} = 8Cov(p^b, p^{b'})$. In all

cases, the reference allele has been randomized so $E(p) = 0.5$.

Table 1. Relationships within and across two pairs of populations

		Holstein	Jersey
Cattle	Holstein	0.55	0.48
	Jersey	0.48	0.77
		Landrace	Yorkshire
Pigs	Landrace	0.76	0.26
	Yorkshire	0.26	0.73

Estimates from cattle (VanRaden *et al.* 2011; Legarra *et al.* 2015) and pigs (Xiang *et al.* 2016).

Table 2 includes estimates for additive, dominance and epistatic variation iWe used

estimates for milk yield in Simmenthal cattle (Fuerst and Sölkner 1994) as if it was Holstein,

341 and for litter size in a commercial pig line (Vitezica *et al.* 2018) as if it was Landrace, because

342 we could not find estimates in the same populations.

344 Table 2. Variance component estimates, as ratio from phenotypic variance, from literature

Species	σ_A^2	σ_D^2	σ_{AA}^2
Cattle	0.20	0.09	0.15
Pigs	0.092	0.020	0.016

Estimates for milk yield in cattle (Fuerst and Sölkner 1994) and litter size in pigs (Vitezica *et*

al. 2018).

Then, we assumed three distributions for allele frequencies, following Hill et al. (2008): a

uniform distribution ("Uniform"), a U-shaped distributions with $f(p) \propto p^{-1}(1-p)^{-1}$ with

effective population size of 50 ("Hill"). We also assumed a Beta(0.04, 0.04) distribution,

which is an extreme U-shaped distribution ("Extreme") in which roughly 80% of the QTLs

have minor allele frequency lower than 0.01. This results in respective values of $\overline{H} =$

$(0.333, 0.107, 0.037)$ and $\overline{H^2} = (0.133, 0.018, 0.013)$. Using the values described above, we

estimated the correlation of QTL effects across breeds, $r\left(\alpha_i^b, \alpha_i^{b'}\right)$, using equation (5).

Similarly, the correlation of QTL effects a certain number of generations away, $r\left(\alpha_i^{t_2}, \alpha_i^{t_1}\right)$,

was obtained applying Equation (7). An increase $\Delta F = 0.01$ resulting in $\Delta F_\gamma = 0.01(1 -$

$\gamma/2)$ per generation was assumed, for the same assumed variances and allele frequency

distributions.

Simulations. The objective of the simulation is to verify our results considering different

kinds of non-additive biological gene action, and not to infer the values of the correlation of

substitution effects across populations as this requires realistic presumptions of the forms

and magnitudes of epistatic interactions, something that is largely unknown.

We simulated two divergent populations using QMSim (Sargolzaei and Schenkel 2009). First,

2000 generations of the historical population were generated with an effective population

size of 1500 individuals, followed by a gradual contraction during the following 500

generations to an effective population size of 100 individuals. It followed a gradual

expansion to a population size of 1500 during the last 500 generations. From this common

historical population, two divergent populations were selected in opposite directions for 5

generations of 1000 individuals each. Selection was for a trait with narrow-sense heritability

of 1, as a device to speed up divergence; however, for the purposes of this work we

simulated another, uncorrelated, trait (with a broad-sense heritability of 1). There were 30

chromosomes of 100 cM each and 66,000 segregating loci (60,000 SNPs and 6,000 QTLs) in

the first generation of the historical population. The number of segregating QTLs and SNPs in

the last generation of the historical population was 750 and 13,555, respectively.

To simulate non-additive biological gene action, we considered three scenarios: Complete

Dominance (within locus), 5 locus complementary epistasis and 5 locus additive by additive

epistasis. Complete dominance has the genotypic value of the heterozygote equal to one of

the homozygotes. Additive by additive epistasis can be understood as a pure multiplication

of gene contents. Complementary epistasis implies that a simultaneous homozygote

genotype at all loci results in a genotypic value, and any other combination in another

genotypic value. Explicit functional additive effects were not simulated. These scenarios are

similar to Hill *et al.* (2008) but instead of considering 2 loci we consider 5 (details are given in

the Appendix).

For epistasis, networks are internally epistatic but contribute additively to the total

phenotype. Loci within a network were picked up at random within chromosomes, resulting

in very low LD across loci within network. We considered 5 networks per chromosome

yielding a total of 150 networks. We derived substitution effects α algebraically for the three

different cases (details are given in the Appendix). We used those expressions to evaluate

the true substitution effects, and to obtain the true value $r\left(\alpha^b, \alpha^{b'}\right)$. At the end, we had

two populations of 5000 individuals each (1000 individuals per generation). Using markers

(alternatively, QTL loci) and simulated phenotypes, we estimated (1) variance components

σ_A^2, σ_D^2 and σ_{AA}^2 using G-REML (Vitezica *et al.* 2018) in population 1 (all 5 generations totaling

5000 animals). We then computed, using markers (alternatively, QTL loci), relationships γ

within and across populations and the functions of heterozygosities $\bar{H}$ and $\overline{H^2}$. Then, using

Equation 4 we predicted $r\left(\alpha^b, \alpha^{b'}\right)$ across populations 1 and 2 within generation 5, and we

predicted $r\left(\alpha^{t_1}, \alpha^{t_2}\right)$ within population 1 for $t_1 = 1$ and $t_2 = 2 \dots 5$. The purpose of using,

alternatively to genotypes at markers, genotypes at QTL was to verify or discard that results

were due to omperfect Linkage Disequilibrium.

Data Availability Statement. Appendix associated with this manuscript is in Figshare. Code

and   files   for   simulations   can   be   found   in

http://genoweb.toulouse.inra.fr/~alegarra/simulation_alpha.tar.gz . All other results can be

reproduced using equations and figures from the text.

410   RESULTS

Estimates of correlation across populations using literature values. The results for

reasonable guesses of the distribution of QTL substitution effects are in Table 3. The

correlation of QTL substitution effects across breeds were roughly 0.6 for the "Uniform"

distribution, whereas the "Extreme" distribution resulted in numbers close to 0.4. These

numbers agree reasonably well with scarce estimates of genetic correlation (although,

strictly speaking, not the same thing) in the literature: 0.4 to 0.8 for production traits (Karoui

et al. 2012), 0.3 to 0.5 for milk yield (Legarra *et al.* 2014) and 0.6 for body condition (Porto-

Neto *et al.* 2015). It may be argued that these studies used SNPs instead of true QTLs;

however, Wientjes et al. (Wientjes *et al.* 2017) in a simulation study obtained unbiased

estimates of correlation across QTLs using SNP markers for a true correlation of 0.5.

Table 3: Estimates of correlations of QTL effects across breeds based on values from Tables 1

and 2 and Equation 5, for different distributions of QTL frequencies

Species	Uniform	Hill[1]	Extreme[2]
Cattle	0.60	0.39	0.25
Pigs	0.66	0.46	0.31
Maize	0.57	0.36	0.22

[1] U-shaped distribution with effective population size of 50

[2] Beta(0.04, 0.04) distribution

Concerning decrease of correlations across generations, results (Table 4) show a mild

decrease, and this decrease is more sensitive to the assumed distribution of allelic

frequencies than the previous ones concerning correlation across populations. There are no

explicit estimates of genomic correlation across generations, but drops in the accuracy of

genomic prediction have been reported (Wolc *et al.* 2011; Baloche *et al.* 2014; Liu *et al.*

2016).

435
436 Table 4: Correlation of QTL effects across time, based on values from Table 1 and 2 and

437 Equation 7, for different distributions of QTL frequencies

Species	Distance in generations	Uniform	Hill[1]	Extreme[2]
Cattle	1	0.98	0.95	0.88
	2	0.97	0.90	0.79
	5	0.92	0.80	0.63
	10	0.85	0.67	0.50
Pigs	1	1.00	0.99	0.97
	2	0.99	0.98	0.94
	5	0.98	0.94	0.88
	10	0.96	0.89	0.79

[1] U-shaped distribution with effective population size of 50

[2] Beta(0.04, 0.04) distribution

Simulations. Estimated variance components (expressed as ratios from total phenotypic

variance σ^2) with marker genotypes and with QTL genotypes are shown in Table 5. The part

of non-additive variance increases with the complexity of the system, although this is in part

because we chose on purpose very artificial examples. In general, the two estimates agree,

although with markers we estimate non-zero epistatic additive by additive variance in the

case of complete dominance, where it is zero. This is due to imperfect LD between QTL and

markers and LD across QTLs and also across markers (Campos *et al.* 2019). Indeed, when we

estimated variance components with QTLs, this epistatic variance disappeared as shown in

the bottom three rows of Table 5.

Table 5. Estimated Ratios of additive, dominance and additive by additive variances σ_A^2/σ^2 ,

σ_D^2/σ^2 and σ_{AA}^2/σ^2 (standard errors) in population 1, estimated with markers (three top

rows) and QTLs (three bottom rows)

	σ_A^2/σ^2	σ_D^2/σ^2	σ_{AA}^2/σ^2
Marker-based			
Complete Dominance	0.709 (0.017)	0.130 (0.011)	0.144 (0.031)
Complementary Epistasis	0.440 (0.025)	0.144 (0.016)	0.309 (0.047)
Additive by additive	0.232 (0.025)	0.040 (0.015)	0.587(0.053)
QTL-based			
Complete Dominance	0.676 (0.009)	0.324 (0.009)	0.00001(0.004)
Complementary Epistasis	0.512 (0.019)	0.145 (0.011)	0.248 (0.022)
Additive by additive	0.265 (0.020)	0.009 (0.006)	0.520(0.036)

Estimates of correlation across substitution effects are presented in Table 6 (across

population) and 7 (within population). Our estimates tend to agree with true values, but they

underestimate the true values. Still, they clearly tell apart one scenario with low correlation

(i.e. additive by additive) from one with high correlation (i.e. complete dominance). Using

QTL genotypes for estimation instead of markers yield similar results, so incomplete linkage

disequilibrium does not seem to be the reason.

Table 6. True and estimated $r(\alpha^b, \alpha^{b'})$ across populations 1 and 2, generation 5

	True $r(\alpha^b, \alpha^{b'})$	Estimated $r(\alpha^b, \alpha^{b'})$ with markers	Estimated $r(\alpha^b, \alpha^{b'})$ with QTL
Complete Dominance	0.981	0.945	0.970
Complementary Epistasis	0.933	0.842	0.854
Additive by additive	0.76	0.60	0.532

465   Table 7. True and estimated $r(\alpha^b, \alpha^{b'})$ across generations 1 and 5, population 1

	True $r(\alpha^b, \alpha^{b'})$	Estimated $r(\alpha^b, \alpha^{b'})$ with markers	Estimated $r(\alpha^b, \alpha^{b'})$ with QTL
Complete Dominance	0.993	0.976	0.989
Complementary Epistasis	0.975	0.926	0.943
Additive by additive	0.899	0.777	0.780

469   DISCUSSION

The fact that most genetic variation is statistically additive is well known. However, this does

not imply equality of statistical additive effects across populations, something that is possibly

reflected in the difficulty of predicting across (distinct) populations, either in animal or

human genetics. Only recently the role of non-additive biological gene actions in this

difficulty has started to be recognized (Dai *et al.* 2020; Duenk *et al.* 2020). Here we confirm

and reinforce their findings. These authors presented result of simulations, where the

problem is that the results are scenario specific and do not allow a good appraisal of the

different factors.

In this work, we present a general, formal framework that does not depend on specific

hypothesis on gene action or order of the epistatic interactions. In our derivation, we

approached *high-order functional* epistasis by Taylor expansions, leading to expressions that

involve only *low-order statistical* additive epistasis (actually pairwise). Including one extra

term in our Taylor expansion would include *three-loci statistical* epistasis and so on, but extra

terms would lead to more difficult expressions (covariance of allele frequencies across loci

and populations), and it is expected that the magnitude of the statistical effects is smaller

and smaller with higher orders of interactions (Mäki-Tanila and Hill 2014). We find

reasonable agreement with our simulation and values from literature. However, we have

indeed not thoroughly studied by simulation the quality of our approximation, which would

require considerable thinking on how to simulate epistasis in very general manners.

Three main factors influence the correlation of substitution effects between

populations $r(\alpha^b, \alpha^{b'})$: the genetic similarity of the two populations, the extend of additive,

dominance and additive by additive variance, and the distribution of allele frequencies at

QTL. We consider that showing explicitly these three factors is an achievement, as their role

is clear from previous work in simulated and real data, (Martin *et al.* 2019; Dai *et al.* 2020;

Duenk *et al.* 2020). Now we discuss these three factors.

The distance across populations is conveniently summarized by relationships γ, which

describe covariances of allelic frequencies across populations, and are related to F_{st} indexes.

On ideal populations, these γ relationships depend on the divergence time t but also on the

effective population size N, as shown by the expression $\frac{\gamma_{b,b'}}{8} = Cov(p^b, p^{b'}) \approx 1 -$

$\exp(-t/2N)$ (Bonhomme *et al.* 2010). In equation (4), to estimate $r(\alpha^b, \alpha^{b'})$, relationship

enters into the form $\left(\gamma_{b'} + \gamma_b - 2\gamma_{b,b'}\right)$ which as explained is proportional to the F_{st} index.

If the relationship across populations is very small, $\gamma_{b,b'} = 0$ which results in $\left(\gamma_{b'} + \gamma_b -\right.$

$\left.2\gamma_{b,b'}\right)$ bounded between 0 (all allele frequencies are 0.5) and 4 (alleles are fixed). The more

diverse each of the populations $(\gamma_{b'}, \gamma_b)$, and the more similar one to the other, the higher

$r(\alpha^b, \alpha^{b'})$ will be.

The factors $\frac{1}{2}\frac{\bar{H}}{\overline{H^2}}$ and $\frac{2}{\bar{H}}$ are weighting factors on dominance and additive by additive

variances. If the allele frequencies are modelled using symmetric $Beta(a, a)$ distributions

(see Appendix), these become $\frac{1}{2}\frac{\bar{H}}{\overline{H^2}} = \frac{1}{4\frac{a+1}{a}\left(1+\frac{a+2}{2a+2}\left(\frac{a+3}{2a+3}-2\right)\right)}$ and $\frac{2}{\bar{H}} = 1 + \frac{1}{2a}$. The first is

bounded between 1.5 (for U-shaped distributions) and 1 (for peaked distributions), so that

dominance variation does not play a big role in the difference between substitution effects

across populations unless dominance variation is way much larger that additive variation,

something that seems unlikely based on theory and estimates. However, the second weight,

due to epistasis, is not bounded and is large for small values of a (*e.g.* 13.5 for

Beta(0.04,0.04), which reflects that functional epistasis plays a strong role in additive

variation for U-shaped distributions (Hill *et al.* 2008). This result also reinforces the findings

of Duenk et al. (2020) and Dai et al. (2020) that it is mainly biological epistasis that generates

changes in additive substitution effects across populations. For instance, when one

population drifts, the additive by additive statistical variation enters into additive variation

(Hill *et al.* 2006; Walsh and Lynch 2018).

Thus, $r\left(\alpha_i^b, \alpha_i^{b'}\right)$ will be very low only if (1) populations are distinct, (2) there is *statistical*

epistatic variation and (3) QTL distributions are U-shaped.

In most conceivable cases σ_{AA}^2 and σ_D^2 are of magnitudes roughly similar to additive variances

(Fuerst and Sölkner 1994; Palucci *et al.* 2007; Hill *et al.* 2008; Mäki-Tanila and Hill 2014;

Vitezica *et al.* 2018), and relationships across populations are moderate. Then, for

reasonable assumptions about allele distribution of QTLs, we have shown that the

correlation of substitution effects across populations is typically around 0.6, which basically

agrees with scarce estimates of genetic correlations across populations available in the

literature. Our results seem rather robust to different distributions of allele frequencies.

These values are low enough so that, for practical purposes, predictions across populations

are of little interest, moreover if populations are not closely related as is often the case. This

is of practical relevance, *e.g.* for genomic predictions in livestock improvement, but also in

human genetics *e.g.* for the use of European-based Polygenic Risk Scores in individuals from

other ancestries (Martin *et al.* 2019)

Similar results apply to the same populations across generations, in which case the

correlation of substitution effects across generations goes from 0.99 (next 1-2 generations)

to 0.70 (5-10 generations). This illustrates that, even if genomic predictions would use QTLs

or markers in thight LD with QTLs, there would still be a need for a continuous system of

data collecting and re-estimation of effects.

In the simulations we confirmed that our estimates are reasonably good although slightly

biased downwards. Still, we find that, for the challenging forms of non-additive gene action

that we simulated, the results are quite good. For instance, an explicit functional value was

only included in the case of complete dominance. The additive by additive epistasis is highly

extreme (e.g. substitution effects change sign across spectra of allelic frequency). The other

form of epistasis, complementary epistasis, has genotypic value depending on every locus

being at a particular homozygous genotype and is deemed to be a biological mechanism that

contributes to variance stabilization (Mackay 2014). Complementary epistasis generates

more additive variation that additive by additive epistasis and, as expected, it leads to higher

similarity of substitution effects across populations.

Another result of our theory is that it allows, at least in principle, to assess if genetic

correlation across breeds is due to gene-by-environment or gene-by-gene interaction,

without going to a fully parametric analysis. For instance, Legarra *et al.* (2014) estimated a

genetic correlation of 0.5 for Latxa Cara Rubia - Manech Tête Rousse, two breeds that have

frequent exchanges and are almost identical (in this case, $\Gamma = \begin{pmatrix} \gamma_b & \gamma_{b,b'} \\ \gamma_{b,b'} & \gamma_{b'} \end{pmatrix} =$

$\begin{pmatrix} 0.49 & 0.47 \\ 0.47 & 0.51 \end{pmatrix}$. This would result in a value of $r(\alpha^b, \alpha^{b'})$ much higher than 0.5. A possibility

is that the low value of 0.5 is due to GxE interaction, as the production systems may differ.

This opens new perspectives.

568   CONCLUSIONS

We presented a coherent, approximate theory, that does not invoke any particular

mechanism of gene action, to explain and appraise the change in magnitude of (additive)

QTL substitution effects across populations and generations. The theory gives good

approximate estimates of this correlation, that needs to be otherwise explicitly estimated.

More importantly, the theory shows that the main sources for the change of effects are

relationships across populations, magnitudes of additive and first-order non-additive

variances (dominance and additive by additive), and spectra of allele frequencies. These

findings provide better understanding of the properties of genomic prediction methods and

of quantitative genetics in general.

581   ACKNOWLEDGMENTS

Part of this work was while Y. Wientjes was visiting the GenPhySE unit at INRAE, Toulouse,

financed by the Netherlands Organisation of Scientific Research (NWO). Authors thank INRA

SelGen metaprograms Episel and Epifun. This project has received funding from the

European Unions' Horizon 2020 Research & Innovation programme under grant agreement

N°772787 -SMARTER. We are grateful to the Genotoul Bioinformatics Platform Toulouse

Midi- Pyrenees (Bioinfo Genotoul) for providing computing and storage resources.

REFERENCES

Aiello D., K. Patel, and E. Lasagna, 2018 The myostatin gene: an overview of mechanisms of

action and its relevance to livestock animals. Anim. Genet. 49: 505–519.

https://doi.org/10.1111/age.12696

Baloche G., A. Legarra, G. Sallé, H. Larroque, J. M. Astruc, *et al.*, 2014 Assessment of accuracy

of genomic prediction for French Lacaune dairy sheep. J. Dairy Sci. 97: 1107–1116.

Bonhomme M., C. Chevalet, B. Servin, S. Boitard, J. Abdallah, *et al.*, 2010 Detecting Selection

in Population Trees: The Lewontin and Krakauer Test Extended. Genetics 186: 241–

262. https://doi.org/10.1534/genetics.110.117275

Bonifati V., P. Rizzu, M. J. van Baren, O. Schaap, G. J. Breedveld, *et al.*, 2003 Mutations in the

DJ-1 Gene Associated with Autosomal Recessive Early-Onset Parkinsonism. Science

299: 256–259. https://doi.org/10.1126/science.1077209

Caballero A., and M. A. Toro, 2002 Analysis of genetic diversity for the management of

conserved subdivided populations. Conserv. Genet. 3: 289.

Caballero A., 2020 *Quantitative Genetics*. Cambridge University Press.

Campos G. de los, D. A. Sorensen, and M. A. Toro, 2019 Imperfect Linkage Disequilibrium

Generates Phantom Epistasis (& Perils of Big Data). G3 Genes Genomes Genet.

9: 1429–1436. https://doi.org/10.1534/g3.119.400101

Carlborg Ö., L. Jacobsson, P. Åhgren, P. Siegel, and L. Andersson, 2006 Epistasis and the

release of genetic variation during long-term selection. Nat. Genet. 38: 418–420.

https://doi.org/10.1038/ng1761

Dai Z., N. Long, and W. Huang, 2020 Influence of Genetic Interactions on Polygenic

Prediction. G3 Genes Genomes Genet. 10: 109–115.

https://doi.org/10.1534/g3.119.400812

Duenk P., P. Bijma, M. P. L. Calus, Y. C. J. Wientjes, and J. H. J. van der Werf, 2020 The Impact

of Non-additive Effects on the Genetic Correlation Between Populations. G3 Genes

Genomes Genet. 10: 783–795. https://doi.org/10.1534/g3.119.400663

Fernandez E. N., J.P. Sanchez, R. Martinez, A. Legarra, M. Baselga, 2017 Role of inbreeding

depression, non-inbred dominance deviations and random year-season effect in

genetic trends for prolificacy in rabbit closed lines. J. Anim. Breed. Genet. 6: 441–452.

Falconer D. S., and T. F. C. Mackay, 1996 *Introduction to quantitative genetics*. Longman New

York.

Fuerst C., and J. Sölkner, 1994 Additive and Nonadditive Genetic Variances for Milk Yield,

Fertility, and Lifetime Performance Traits of Dairy Cattle. J. Dairy Sci. 77: 1114–1125.

https://doi.org/10.3168/jds.S0022-0302(94)77047-8

Fuerst C., J. W. James, J. Sölkner, and A. Essl, 1997 Impact of dominance and epistasis on the

genetic make-up of simulated populations under selection: a model development. J.

Anim. Breed. Genet. 114: 163–175. https://doi.org/10.1111/j.1439-

0388.1997.tb00502.x

Garcia-Baccino C. A., A. Legarra, O. F. Christensen, I. Misztal, I. Pocrnic, *et al.*, 2017

Metafounders are related to Fst fixation indices and reduce bias in single-step

genomic evaluations. Genet. Sel. Evol. 49: 34. https://doi.org/10.1186/s12711-017-

0309-2

García-Ruiz A., J. B. Cole, P. M. VanRaden, G. R. Wiggans, F. J. Ruiz-López, *et al.*, 2016

Changes in genetic selection differentials and generation intervals in US Holstein

dairy cattle as a result of genomic selection. Proc. Natl. Acad. Sci. 113: E3995–E4004.

Gautier M., A. Capitan, S. Fritz, A. Eggen, D. Boichard, *et al.*, 2007 Characterization of the

DGAT1 K232A and Variable Number of Tandem Repeat Polymorphisms in French

Dairy Cattle. J. Dairy Sci. 90: 2980–2988. https://doi.org/10.3168/jds.2006-707

Grobet L., L. J. Royo Martin, D. Poncelet, D. Pirottin, B. Brouwers, *et al.*, 1997 A deletion in

the bovine myostatin gene causes the double–muscled phenotype in cattle. Nat.

Genet. 17: 71–74. https://doi.org/10.1038/ng0997-71

Haile-Mariam M., and J. E. Pryce, 2015 Variances and correlations of milk production,

fertility, longevity, and type traits over time in Australian Holstein cattle. J. Dairy Sci.

98: 7364–7379. https://doi.org/10.3168/jds.2015-9537

Hayes B. J., P. J. Bowman, A. C. Chamberlain, K. Verbyla, and M. E. Goddard, 2009 Accuracy

649   of genomic breeding values in multi-breed dairy cattle populations. Genet Sel Evol

650   41: 51.

651   Hill W. G., N. H. Barton, and M. Turelli, 2006 Prediction of effects of genetic drift on variance

652   components under a general model of epistasis. Theor. Popul. Biol. 70: 56–62.

653   https://doi.org/10.1016/j.tpb.2005.10.001

654   Hill W. G., M. E. Goddard, and P. M. Visscher, 2008 Data and Theory Point to Mainly Additive

655   Genetic Variance for Complex Traits. PLOS Genet. 4: e1000008.

656   https://doi.org/10.1371/journal.pgen.1000008

657   Karoui S., M. J. Carabaño, C. Díaz, and A. Legarra, 2012 Joint genomic evaluation of French

658   dairy cattle breeds using multiple-trait models. Genet Sel Evol 44: 39.

659   Kojima K., 1959 Role of epistasis and overdominance in stability of equilibria with selection.

660   Proc. Natl. Acad. Sci. U. S. A. 45: 984–989.

661   Kojima K.-I., 1961 Effects of dominance and size of population on response to mass

662   selection*. Genet. Res. 2: 177–188. https://doi.org/10.1017/S0016672300000689

663   Lande R., and R. Thompson, 1990 Efficiency of marker-assisted selection in the improvement

664   of quantitative traits. Genetics 124: 743–756.

665   Legarra A., G. Baloche, F. Barillet, J.-M. Astruc, C. Soulas, *et al.*, 2014 Within-and across-

666   breed genomic predictions and genomic relationships for Western Pyrenees dairy

667   sheep breeds Latxa, Manech, and Basco-Béarnaise. J. Dairy Sci. 97: 3200–3212.

Legarra A., O. F. Christensen, Z. G. Vitezica, I. Aguilar, and I. Misztal, 2015 Ancestral

Relationships Using Metafounders: Finite Ancestral Populations and Across

Population Relationships. Genetics 200: 455–468.

https://doi.org/10.1534/genetics.115.177014

Lewontin R. C., and J. Krakauer, 1973 Distribution of Gene Frequency as a Test of the Theory

of the Selective Neutrality of Polymorphisms. Genetics 74: 175–195.

Liu Z., H. Alkhoder, F. Reinhardt, and R. Reents, 2016 Accuracy and bias of genomic

prediction for second-generation candidates. Interbull Bull. 50: 17-23.

Lynch M., and B. Walsh, 1998 *Genetics and analysis of quantitative traits*. Sinauer associates.

Mackay T. F. C., 2014 Epistasis and Quantitative Traits: Using Model Organisms to Study

Gene-Gene Interactions. Nat. Rev. Genet. 15: 22–33.

https://doi.org/10.1038/nrg3627

Mackay T. F. C., 2015 Epistasis for Quantitative Traits in Drosophila, pp. 47–70 in *Epistasis:*

Methods and Protocols, Methods in Molecular Biology. edited by Moore J. H.,

Williams S. M. Springer, New York, NY.

MacLeod I. M., P. J. Bowman, C. J. Vander Jagt, M. Haile-Mariam, K. E. Kemper, *et al.*, 2016

Exploiting biological priors and sequence variants enhances QTL discovery and

genomic prediction of complex traits. BMC Genomics 17: 144.

https://doi.org/10.1186/s12864-016-2443-6

Magwire M. M., A. Yamamoto, M. A. Carbone, N. V. Roshina, A. V. Symonenko, *et al.*, 2010

Quantitative and Molecular Genetic Analyses of Mutations Increasing Drosophila Life

Span. PLOS Genet. 6: e1001037. https://doi.org/10.1371/journal.pgen.1001037

Mäki-Tanila A., and W. G. Hill, 2014 Influence of Gene Interaction on Complex Trait Variation

with Multilocus Models. Genetics 198: 355–367.

https://doi.org/10.1534/genetics.114.165282

Martin A. R., M. Kanai, Y. Kamatani, Y. Okada, B. M. Neale, *et al.*, 2019 Clinical use of current

polygenic risk scores may exacerbate health disparities. Nat. Genet. 51: 584–591.

https://doi.org/10.1038/s41588-019-0379-x

Meuwissen T. H. E., B. J. Hayes, and M. E. Goddard, 2001 Prediction of total genetic value

using genome-wide dense marker maps. Genetics 157: 1819–1829.

Moghaddar N., M. Khansefid, J. H. J. van der Werf, S. Bolormaa, N. Duijvesteijn, *et al.*, 2019

Genomic prediction based on selected variants from imputed whole-genome

sequence data in Australian sheep populations. Genet. Sel. Evol. 51: 72.

https://doi.org/10.1186/s12711-019-0514-2

Palucci V., L. R. Schaeffer, F. Miglior, and V. Osborne, 2007 Non-additive genetic effects for

fertility traits in Canadian Holstein cattle (Open Access publication). Genet. Sel. Evol.

39: 181. https://doi.org/10.1186/1297-9686-39-2-181

Porto-Neto L. R., W. Barendse, J. M. Henshall, S. M. McWilliam, S. A. Lehnert, *et al.*, 2015

Genomic correlation: harnessing the benefit of combining two unrelated populations

for genomic selection. Genet. Sel. Evol. 47: 84. https://doi.org/10.1186/s12711-015-

0162-0

Robertson A., 1975 Gene Frequency Distributions as a Test of Selective Neutrality. Genetics

81: 775–785.

Rodríguez-Ramilo S., J. M. Elsen, and A. Legarra, 2019 Inbreeding and effective population

size in French dairy sheep: Comparison between genomic and pedigree estimates. J.

Dairy Sci. 102: 4227–4237.

Roos A. P. W. de, B. J. Hayes, and M. E. Goddard, 2009 Reliability of Genomic Predictions

Across Multiple Populations. Genetics 183: 1545–1553.

https://doi.org/10.1534/genetics.109.104935

Rupp R., P. Senin, J. Sarry, C. Allain, C. Tasca, *et al.*, 2015 A Point Mutation in Suppressor of

Cytokine Signalling 2 (Socs2) Increases the Susceptibility to Inflammation of the

Mammary Gland while Associated with Higher Body Weight and Size and Higher Milk

Production in a Sheep Model. PLOS Genet. 11: e1005629.

https://doi.org/10.1371/journal.pgen.1005629

Searle S., 1971 *Linear models*. New York, NY (EUA). John Wiley.

Streit M., N. Neugebauer, T. H. E. Meuwissen, and J. Bennewitz, 2011 Short communication:

evidence for a major gene by polygene interaction for milk production traits in

German Holstein dairy cattle. J. Dairy Sci. 94: 1597–1600.

https://doi.org/10.3168/jds.2010-3834

Tsuruta S., I. Misztal, T. J. Lawlor, and L. Klei, 2004 Modeling final scores in US Holsteins as a

function of year of classification using a random regression model. Livest. Prod. Sci.

91: 199–207. https://doi.org/10.1016/j.livprodsci.2003.09.016

VanRaden P., K. Olson, G. Wiggans, J. Cole, and M. Tooker, 2011 Genomic inbreeding and

relationships among Holsteins, Jerseys, and Brown Swiss. J. Dairy Sci. 94: 5673–5682.

Vitezica Z. G., A. Reverter, W. Herring, and A. Legarra, 2018 Dominance and epistatic genetic

variances for litter size in pigs using genomic models. Genet. Sel. Evol. 50: 1–8.

Walsh B., and M. Lynch, 2018 *Evolution and Selection of Quantitative Traits*. Oxford

University Press.

Welsh C. S., T. S. Stewart, C. Schwab, and H. D. Blackburn, 2010 Pedigree analysis of 5 swine

breeds in the United States and the implications for genetic conservation. J. Anim.

Sci. 88: 1610–1618. https://doi.org/10.2527/jas.2009-2537

Wientjes Y. C. J., P. Bijma, J. Vandenplas, and M. P. L. Calus, 2017 Multi-population Genomic

Relationships for Estimating Current Genetic Variances Within and Genetic

Correlations Between Populations. Genetics 207: 503–515.

https://doi.org/10.1534/genetics.117.300152

Wolc A., J. Arango, P. Settar, J. E. Fulton, N. P. O'Sullivan, *et al.*, 2011 Persistence of accuracy

of genomic estimated breeding values over generations in layer chickens. Genet Sel

Evol 43: 23.

Xiang T., B. Nielsen, G. Su, A. Legarra, and O. F. Christensen, 2016 Application of single-step

genomic evaluation for crossbred performance in pig. J. Anim. Sci. 94: 936–948.

https://doi.org/10.2527/jas.2015-9930

Xiang T., O. F. Christensen, and A. Legarra, 2017 Technical note: Genomic evaluation for

crossbred performance in a single-step approach with metafounders. J. Anim. Sci. 95:

1472–1480.