**Osteosarcoma tumors maintain intra-tumoral transcriptional heterogeneity during bone**

**and lung colonization**

Sanjana Rajan[1,2*], Emily Franz[1,2*], Camille A. McAloney[2,3], Tatyana A. Vetter[4], Maren Cam[2],

Amy C. Gross[2], Cenny Taslim[2], Meng Wang[2], Matthew V. Cannon[2], Alexander Oles[4], Ryan D.

Roberts[2,6,7#]

INSTITUTIONS:

[1]Molecular, Cellular, and Developmental Biology Program, The Ohio State University,

Columbus, Ohio, USA

[2]Center for Childhood Cancers and Blood Diseases, Abigail Wexner Research Institute at

Nationwide Children's Hospital, Ohio, USA

[3]Department of Veterinary Biosciences, College of Veterinary Medicine, The Ohio State

University, Columbus, Ohio, USA

[4]Center for Gene Therapy, Abigail Wexner Research Institute, Nationwide Children's Hospital,

Columbus, Ohio, USA

[6]Division of Pediatric Hematology, Oncology, and BMT, Nationwide Children's Hospital,

Columbus, Ohio, USA

[7]The Ohio State University James Comprehensive Cancer Center, Columbus, Ohio, USA

*Co-first authors

#Corresponding author

**Email addresses:**

SR: SRajan11@its.jnj.com

EF: Emily.Franz@nationwidechildrens.org

CAM: Camille.McAloney@nationwidechildrens.org

TAV: Tatyana.Vetter@nationwidechildrens.org

MC: Maren.Cam@nationwidechildrens.org

ACG: Amy.Gross@nationwidechildrens.org

CT: Cenny.Taslim@nationwidechildrens.org

MW: Meng.Wang@nationwidechildrens.org

MVC: Matthew.Cannon@nationwidechildrens.org

AO: oles@musc.edu

RDR: Ryan.Roberts@nationwidechildrens.org

**Correspondence:**

Ryan D. Roberts

Division of Pediatric Hematology, Oncology, and BMT

Nationwide Children's Hospital

700 Children's Drive, Columbus, OH 43205

Email: Ryan.Roberts@nationwidechildrens.org

Phone: 614-722-29966; Fax: 614-355-2927

**Abstract**

Background

Tumors are complex tissues containing collections of phenotypically diverse malignant and
nonmalignant cells. We know little of the mechanisms that govern heterogeneity of tumor cells
nor of the role heterogeneity plays in overcoming stresses, such as adaptation to different
microenvironments. Osteosarcoma is an ideal model for studying these mechanisms—it exhibits
widespread inter- and intra-tumoral heterogeneity, predictable patterns of metastasis, and a lack
of clear targetable driver mutations. Understanding the processes that facilitate adaptation to
primary and metastatic microenvironments could inform the development of therapeutic
targeting strategies.

Results

We investigated single-cell RNA-sequencing profiles of 47,977 cells obtained from cell line and
patient-derived xenograft models as cells adapted to growth within primary bone and metastatic
lung environments. Tumor cells maintained phenotypic heterogeneity as they responded to the
selective pressures imposed during bone and lung colonization. Heterogenous subsets of cells
defined by distinct transcriptional profiles were maintained within bone- and lung-colonizing
tumors, despite high-level selection. One prominent heterogenous feature involving glucose
metabolism was clearly validated using immunofluorescence staining. Finally, using concurrent
lineage tracing and single-cell transcriptomics, we found that lung colonization enriches for
multiple clones with distinct transcriptional profiles that are preserved across cellular
generations.

Conclusions

Response to environmental stressors occurs through complex and dynamic phenotypic
adaptations. Heterogeneity is maintained, even in conditions that enforce clonal selection.

These findings likely reflect the influences of developmental processes promoting diversification

of tumor cell subpopulations, which are retained, even in the face of selective pressures.

**Key words**

Osteosarcoma, heterogeneity, transcriptional profile, clonal selection, microenvironment,

metastasis

## Background

Osteosarcoma is the most common malignant tumor of bone in pediatric patients[1]. This aggressive disease occurs in the metaphyseal region of the long bones, coinciding in both anatomic location and developmental timing with periods of rapid linear bone growth[2]. Localized disease has a favorable five-year survival rate of over 70%[3], but survival plummets to less than 20% in the event of metastasis[4]. Unfortunately, survival rates have not changed since the introduction of chemotherapy in the 1980s[5]. Staging and therapeutic assignment remain solely based on the presence or absence of metastasis[6]. One unique feature of osteosarcoma is its extreme tropism for the lung, which is the primary site of metastasis in nearly 90% of patients[7,8]. How these tumors adapt to survive and grow as they migrate from a bone to a lung environment remains poorly understood. While others have reported that osteosarcoma metastasis is a polyclonal process[9], the molecular pathways that mediate these adaptations during tissue colonization—which may involve a coordination of behaviors between cells from distinct clones with distinct phenotypes—remain to be studied at a single-cell level. Such investigation may provide novel insights into tumor biology and reveal vulnerabilities not evident in studies that utilize bulk methods.

Most osteosarcoma patients have disseminated tumor cells at the time of diagnosis[6,10]. Thus, studies focused on the latter steps in the metastatic process, especially lung colonization, are those most likely to produce targets that will translate into impactful clinical interventions. A large body of literature suggests that tumor cells face enormous stresses upon dissemination to the lung—only the fittest cells survive the transition from dissemination to lung colonization, while most cells die[11]. Thus, the experiments shown here focus deliberately on understanding the contributions of clonal selection and transcriptional heterogeneity to the adaptive processes that allow tumors to traverse the tissue colonization bottleneck[11–15], using reductionist systems that are agnostic of early steps in the metastatic cascade. For instance, whether the properties

that mediate fitness within the lung arise from rare, but pre-existing clones that carry an intrinsic

survival advantage or from some active adaptive response to the new environment remains

poorly understood. Such fundamental questions have largely remained unanswered due to

technical limitations. In this study, we overcome this gap by combining single-cell

transcriptomics with lineage tracing to study transcriptional heterogeneity dynamics and clonal

evolution using linked single cell datasets.

We analyzed single cell transcriptomic libraries of 47,977 osteosarcoma cells obtained from cell

lines and low-passage patient-derived xenograft (PDX) models that were grown as both

orthotopic primary tumors and lung metastases to identify changes in intra-tumoral

transcriptional heterogeneity that occur during tibia and lung colonization. While we suspected

that adaptation to growth within the tibia or lung would select for a rather narrowly-defined

subset of cells with transcriptional phenotypes that endowed cells with increased fitness in these

distinct environments, we instead found that tumors maintain a high degree of transcriptional

heterogeneity while adapting to growth in both bone and lung environments. However, while the

overall transcriptional heterogeneity was maintained, the process of adaptation to these distinct

environments revealed reproducible patterns of biologic sub-specialization within subsets of

cells that were characteristic for each tissue niche. In one intriguing example, populations of

cells expressing genes suggestive of reliance on glycolytic and aerobic metabolism both

emerged during lung colonization. Using immunofluorescence, we validated this metabolic

heterogeneity in both primary and metastatic mouse tumors. Furthermore, we validated the

heterogeneous activation of multiple pathways identified from our preclinical models in single

cell RNA-seq datasets from patient tumors. Finally, we combined lineage tracing with single cell

transcriptomics to study the contribution of clonal selection to the phenotypic shifts that we

observed. Surprisingly, while several clones exhibited clear expansion during lung colonization,

these clones emerged within clusters spread across the entire spectrum of transcriptional

profiles, rather than a particular cluster representing a genome-wide transcriptional state that

endowed tissue-specific fitness. However, isolation of cells from these expanding clones

revealed activation of a more focused set of genes that again suggested the importance of

metabolic flexibility during lung colonization.

Based on these investigations, we conclude that osteosarcoma cells undergo dynamic changes

in their transcriptional phenotype as they colonize different microenvironmental conditions, while

simultaneously maintaining an overall degree of transcriptional heterogeneity in the wake of

clonal selection. Overall, these data highlight the importance of understanding intra-tumor

heterogeneity in the context of tumor cell transcriptional changes or adaptations during tissue

colonization.

**Results**

**Transcriptional heterogeneity in cell line and PDX models of osteosarcoma**

Recent work has shown that osteosarcoma tumors collected directly from patients exhibit

significant transcriptional heterogeneity[16,17]. However, it is unclear if this heterogeneity is

maintained within cell line and PDX models of disease[6]. We investigated phenotypic

heterogeneity within two *in vitro* and *in vivo* models of osteosarcoma by performing single-cell

RNA sequencing (scRNA-seq). We generated transcriptomic libraries from a cell line (OS-17)

and a PDX (NCH-OS-7; see *Table 1,* supplementary *Table S1*, for full characteristics as per

PDX-MI standards) grown in cell culture or as subcutaneous flank tumors (>2,500 cells per

model). To distinguish tumor cells from murine host cells, we separated cells with reads that

mapped to the human genome from those that mapped to mouse. Principal component analysis

identified that the majority of the variation in gene expression of tumor cells was dominated by

cell cycle related genes, a common finding in proliferative malignant tissues[18] (see

supplementary *Figure S1*). Scaling out cell-cycle related genes unmasked underlying

heterogeneity that did not correspond to variation arising solely from cells being in different

states of cell cycle (see supplementary *Figure S1*, **S2**A, B, and *Figure 1*A, B). Using genes that

were differentially upregulated in each cluster relative to the remaining cells in each sample, we

identified pathways enriched in these subsets of cells as well as those pathways differentially

downregulated in each cluster (*Figure 1*C, see supplementary *Figure S3*A). While subsets

within each model shared some enriched gene sets--for example, OS-17 cells in cluster 0 and

cluster 1 showed enrichment for epithelial to mesenchymal transition (EMT) genes--none of the

subsets had completely identical profiles, suggesting that these subsets are indeed

transcriptionally distinct. Interestingly, while EMT transcription factors play an important role in

promotion of osteosarcoma cell invasion and metastasis[19], expression of EMT-related genes

was associated with certain, distinct subsets. Furthermore, individual subsets had differentially

expressed genes that were associated with multiple gene sets. For example, OS-17 cells in

cluster 2 were enriched for E2F targets, G2M checkpoint target genes, and MYC target genes;

while NCH-OS-7 cells in cluster 0 were enriched for genes associated with EMT, glycolysis,

hypoxia, and MYC target genes. This identification of gene sets exclusive to specific clusters of

cells validated that these cell line and PDX models demonstrate intra-tumor phenotypic

heterogeneity.

**Transcriptional signatures associated with colonization of bone and lung**

**microenvironments**

Microenvironment cues play a critical role in shaping transcriptional plasticity[20]. We

hypothesized that transcriptional signatures shared across models upon colonization of tibia or

lung microenvironments will inform characteristics that are microenvironment-specific rather

than tumor-specific. Single cell suspensions of two cell lines and two PDXs were injected intra-

tibially or intra-venously to generate orthotopic models of bone-colonizing and lung-colonizing

lesions (*Figure 2*A schematic). Both the PDXs included in this study (NCH-OS-2, NCH-OS-7)

are early passage (passage 3, passage 5) and were obtained from patient lung metastases.

*Table 1*. **Characteristics of osteosarcoma models used in this study**

| Model | Species | Type | Host | Endpoint | Characteristics |
|-------|---------|------|------|----------|-----------------|
| **OS-17** | Human | Cell Line | Icr-scid | 70 days | Low passage, very well-characterized, reliable metastases |
| **143B** | Human | Cell Line | Icr-scid | 65 days | Widely utilized, very well-characterized, reliable metastases |
| **NCH-OS-2** | Human | PDX | Icr-scid | 200 days | Very low-passage, highly penetrant, from an untreated synchronous metastasis |
| **NCH-OS-7** | Human | PDX | Icr-scid | 184 days | Very low-passage, highly penetrant, from a heavily-pretreated, multiply-relapsed metastasis |

To study the transcriptional profiles associated with tibia and lung colonization, we performed a

differential expression analysis of the pseudo-bulked scRNA-seq data from bone-colonizing

tumors or lung-colonizing lesions relative to their corresponding starting population (cell

culture/flank tumors). Most of the differentially regulated genes were unique to each model, with

only 0.3-0.8% of genes shared across all four models (*Figure 2*B). Using genes that were either

up- or down-regulated in at least three of the four models (2.7-5.6% of genes), we performed

pathway enrichment analysis to identify pathways associated with colonization of tibia or lung

tissues (*Figure 2*C). We verified the shared pathways between these human osteosarcoma

samples and mouse osteosarcoma models (K7M2, F420) to determine shared pathways

between xenograft and syngeneic models (see supplementary *Figure S4*). Genes that were

significantly upregulated upon tibia colonization included those associated with glycolysis,

hypoxia, MYC targets, fatty acid metabolism, and oxidative phosphorylation. None of the

identified downregulated gene sets were statistically significant. Gene sets that were most

significantly upregulated with lung colonization included those associated with TNFα signaling

via NFκB and EMT, whereas those significantly downregulated included MYC targets and

MTORC1 signaling. Overall, we identified common pathways that were differentially regulated

across the four models upon tibia or lung colonization.

**Transcriptional heterogeneity of individual cells within a tumor**

We analyzed single cell transcriptional profiles of the above datasets using our scRNA-seq

bioinformatics workflow to study the role of intra-tumor heterogeneity in tissue colonization

(*Figure 3*A). To estimate the degree of intra-tumor heterogeneity in our models, we computed

intra-tumor heterogeneity scores (ITH scores) from the average Euclidian distance between

each cell and every other cell in the same dataset[18]. In essence, this is a measure of how

similar any particular cell is to the other cells within a sample. Since osteosarcoma can arise

from transformed progenitor cells with osteoblastic differentiation and osteoid production[6], we

compared the heterogeneity scores in osteosarcoma cells relative to human primary osteoblast

cells grown in cell culture. We found that most osteosarcoma samples exhibit somewhat higher

ITH scores than do normal osteoblast cultures (p <0.001, *Figure 3*B), though there remained a

high degree of overlap in the distributions of phenotypic similarity (ITH scores) within the

populations of all three cell cultures. Indeed, cells from osteoblast cultures exhibit several

distinct phenotypes that readily cluster away from each other, consistent with previous reports[21]

(see supplementary *Figure S5*) and similar to that observed in the osteosarcoma samples. We

have included overlap scores[22] in the figure, which gives a more appropriate statistical

assessment of the proposed hypothesis than a p value, given the type of data[23] (i.e., we are not

evaluating whether two populations containing thousands of cells are identical, rather evaluating

the degree of similarity vs difference in their ITH Score distribution).

We hypothesized that adaptation to distinct microenvironmental conditions would create

selective pressures that would cause a narrowing of the transcriptional phenotypes of the tumor

cells as the process would select for cells expressing genes that endow them with an

environment-specific fitness phenotype. Surprisingly, we found that osteosarcoma cells grown in

tibia and lung microenvironments showed no or little decrease in their ITH score distributions

relative to those grown *in vitro* or as flank tumors (**Figure 3**B, see supplementary **Figure S6**A),

though they do show a broadening of the cell-cell similarity distribution, with small populations of

cells emerging that have increasingly distinct phenotypes (higher ITH scores). These results

suggest that osteosarcoma cells are at least as transcriptionally heterogeneous as non-

malignant osteoblast cells and hint that adaptation to new tissues might drive the emergence of

minor subpopulations. At a high level, these data support the concept that these cells retain the

programs of the tissues they derive from, and that phenotypic diversity (as defined by

transcriptional heterogeneity) may be a fundamental property that is retained as they colonize

the tibia and lung microenvironments.

To determine if bone-colonizing tumors and lung-colonizing lesions display similar phenotypes,

we attempted to use an integrative approach across models (see supplementary **Figure S7**).

However, cells predominantly clustered by tumor model of origin, a phenomenon reported in

other cancers including small cell lung cancers[18]. This indicates a high degree of inter-tumor

heterogeneity, a well-known characteristic of osteosarcoma tumors[24]. We then compared bone-

colonizing and lung-colonizing lesions within the same model (OS-17) and characterized the

transcriptional profile of each cluster (see supplementary **Figure S8**, **S2**C-F). Using a cluster

distribution analysis, we calculated the relative percentages of cells in each cluster to compare

cells that colonized the tibia or the lung (see supplementary **Table S2**). We identified a

significant overlap in the transcriptional profiles of cells that colonized the two

microenvironments, while certain cells demonstrated profiles specific to each microenvironment

(*Figure 3*C, see supplementary *Figure S6*B). For instance, both primary and metastatic OS-17

lesions had cells that exhibited the phenotype associated with cluster 0 (47% in tibia and 42% in

lung), however, cells with the phenotype associated with cluster 2 were significantly enriched in

metastatic lesions (1% in tibia and 40% in lung). Using the pathway enrichment analysis

outlined in *Figure 1*, we investigated the transcriptional profiles associated with each cluster

(*Figure 3*D, see supplementary *Figure S3*B, *S6*C). This cluster enrichment-based strategy

allowed us to overcome the challenges presented by inter-tumoral heterogeneity and instead

focus on the characteristics of subsets that show differential fitness within the bone and lung

microenvironments. Pathways enriched in each cluster were unique, with high heterogeneity in

the activation of individual pathways across clusters within the same tumor (see supplementary

*Figure S9 - Figure S18)*. For example, while OS-17 cells in cluster 0 and cluster 3 were

associated with MTORC1 signaling, only cells in cluster 3 were enriched for glycolysis-related

genes. Our results suggested that the maintenance of subpopulations with distinct phenotypes

is reproducible and likely important for the biology of lung colonization. To ensure that these

results were indicative of generalizable properties of osteosarcoma tumors (and not simply an

artifact of our chosen model system), we validated the presence of these intratumoral subsets

defined by glycolysis, hypoxia, epithelial-mesenchymal transition, and TNFa signaling via NFKB

within scRNA-seq datasets generated directly from patient tumors (see supplementary *Figure*

*S19 - Figure S28*). Our results suggest that colonization of lung tissue requires contributions

from tumor cell subsets exhibiting distinct metabolic behaviors.

**Metabolic heterogeneity in osteosarcoma tumors**

To validate our observations suggesting changes in gene expression related to metabolic

heterogeneity at the protein level, we performed immunofluorescence staining for a known

activation marker of glycolysis, glucose transporter 1 (GLUT1, a primary component of the

observed gene signature), in lung lesions from two models (OS-17 and 143B). Sections were

co-stained for vimentin to distinguish osteosarcoma cells from lung parenchyma. We observed

variable staining for GLUT1 protein within individual lung lesions, proving that the observed

heterogeneity is relevant at the protein level and not an artifact resulting from collections of

homogenous, spatially isolated lesions that stained differently from one another (**Figure 4**A, B).

Interestingly, this heterogeneity was evident even in the smallest lesions, suggesting that this

metabolic heterogeneity either represents a fundamental property intrinsic to subpopulations

when exposed to the lung environment or provides some survival and/or growth advantage for

metastatic lung lesions.

To confirm that the heterogeneity observed within this representative pathway results from

activation of gene expression modules within the cells and not simply from stochastic variations

in gene expression, we stained for other members of the glycolysis gene set (hexokinase 2–

HK2, carbonic anhydrase 9–CA9, monocarboxylate transporter 4–MCT4)[25] and a known

upstream regulator of glycolysis, MYC[26] to evaluate the degree of concordance within individual

cells. We measured the intensity of GLUT1 fluorescence of individual pixels, along with

fluorescence of other markers, within areas positive for vimentin. We noted a high correlation in

staining for GLUT1 and the other markers within 143B and OS-17 metastatic lesions (see

supplementary **Figure S29**). Immunofluorescence staining for markers of glycolysis in bone

tumors from these models identified a high degree of correlation for high, low, and intermediate

staining for all of these targets (see supplementary **Figure S29A**, **Figure S30**). The same

staining of these cells in culture indicated that heterogeneous expression is present in culture

(see supplementary **Figure S31**). This staining validated our sequencing-based finding of

heterogeneity in glycolysis activation at the protein level and confirmed that this heterogeneity

was not the result of stochastic gene expression changes.

**Tumor cell clones with diverse transcriptional phenotypes expand during lung**

**colonization**

To evaluate the contributions of clonal selection to phenotypic adaptation and heterogeneity as

osteosarcoma cells colonize tibia and lung microenvironments, we combined expressed-tag

lineage tracing with single cell transcriptomics (**A,** B) Immunofluorescence staining of mouse

lungs bearing OS-17 and 143B lung-colonizing tumors, respectively, for GLUT1 (green; a

marker of glycolysis) and vimentin (red; marker to identify osteosarcoma cells). Magnified

GLUT1 staining is shown for the boxed regions on the whole-section images. Lesion edges are

indicated by white outlines in the magnified regions. Tumors showed a high degree of intra-

tumor variation in GLUT1 staining intensity. C) FeaturePlots for the Glycolysis module score in

patient primary tumor datasets. The module score for Glycolysis was calculated using the

'AddModuleScore' function in Seurat with msigdb HALLMARK_GLYCOLYSIS genes as input

features for expression program.

**Figure 5**A). Lentivirus infection introduced a very high diversity, heritable, and 3' capture

compatible lineage tag library in osteosarcoma cells in culture. After reserving a representative

sample of the uniquely tagged cells, we generated bone and lung lesions. Single cell libraries of

tibia tumors, lung tumors and cells in culture were processed bioinformatically to overlay

phenotype and lineage information for each cell, and cluster transcriptional profiles were

determined (see supplementary **Figure S2**G). At the level of phenotype, we noted that cells in

tibia and lung cluster away from cells in cell culture, suggesting tumor-level adaptation to tibia or

lung microenvironments (**A,** B) Immunofluorescence staining of mouse lungs bearing OS-17

and 143B lung-colonizing tumors, respectively, for GLUT1 (green; a marker of glycolysis) and

vimentin (red; marker to identify osteosarcoma cells). Magnified GLUT1 staining is shown for

the boxed regions on the whole-section images. Lesion edges are indicated by white outlines in

the magnified regions. Tumors showed a high degree of intra-tumor variation in GLUT1 staining

intensity. C) FeaturePlots for the Glycolysis module score in patient primary tumor datasets. The

module score for Glycolysis was calculated using the 'AddModuleScore' function in Seurat with

msigdb HALLMARK_GLYCOLYSIS genes as input features for expression program.

*Figure 5*B). Since the lineage tags are passed from parent to progeny as each cell divides, we

used a frequency distribution analysis for lineage tags in the tibia and lung tumors to determine

if tibia or lung colonization drove enrichment for specific clonal populations. Lineage barcode

enrichment analysis for cells in cell culture identified 85% of cells to have a unique lineage tag,

confirming that our starting population received lineage tags at high diversity. Surprisingly, cells

from orthotopic primary tumors showed very little evidence of clonal selection, suggesting that

bone represents a relatively permissive colonizing environment (*A,* B) Immunofluorescence

staining of mouse lungs bearing OS-17 and 143B lung-colonizing tumors, respectively, for

GLUT1 (green; a marker of glycolysis) and vimentin (red; marker to identify osteosarcoma

cells). Magnified GLUT1 staining is shown for the boxed regions on the whole-section images.

Lesion edges are indicated by white outlines in the magnified regions. Tumors showed a high

degree of intra-tumor variation in GLUT1 staining intensity. C) FeaturePlots for the Glycolysis

module score in patient primary tumor datasets. The module score for Glycolysis was calculated

using the 'AddModuleScore' function in Seurat with msigdb HALLMARK_GLYCOLYSIS genes

as input features for expression program.

*Figure 5*C). In contrast, cells from lung lesions demonstrated a higher level of enrichment for

specific barcode, suggesting either an environment selecting for clones with specific properties

or the activation of programs driving proliferation within a subset of tumor cells (*A,* B)

Immunofluorescence staining of mouse lungs bearing OS-17 and 143B lung-colonizing tumors,

respectively, for GLUT1 (green; a marker of glycolysis) and vimentin (red; marker to identify

osteosarcoma cells). Magnified GLUT1 staining is shown for the boxed regions on the whole-

section images. Lesion edges are indicated by white outlines in the magnified regions. Tumors

showed a high degree of intra-tumor variation in GLUT1 staining intensity. C) FeaturePlots for

the Glycolysis module score in patient primary tumor datasets. The module score for Glycolysis

was calculated using the 'AddModuleScore' function in Seurat with msigdb

HALLMARK_GLYCOLYSIS genes as input features for expression program.

*Figure 5*C). We confirmed that this pattern of clonal selection was reproducible in two

independent replicates (see supplementary *Figure S32*A; supplementary *Table S4* and *Table*

*S5*, for lentivirus distribution in cell culture). Importantly, datasets from OS-17 cells from distinct

biological replicates (different lot of cells, different lot of virus, different litter of mice) overlaid

completely, even without batch correction, suggesting that intra-tumor transcriptional

heterogeneity of these osteosarcoma cells is not random or stochastic and that our decision to

avoid using batch correction techniques is biologically appropriate (see supplementary *Figure*

*S32*B).

We then overlaid lineage information on phenotypic profiles to identify the transcriptional

phenotypes most permissive to clonal expansion in the lung. Interestingly, while we found that

cells within given enriched clonal families clustered together, suggesting that overarching

transcriptional states were passed from parent to progeny, we also found that these expanding

clones emerged from parents exhibiting phenotypes across the transcriptional spectrum (*A,* B)

Immunofluorescence staining of mouse lungs bearing OS-17 and 143B lung-colonizing tumors,

respectively, for GLUT1 (green; a marker of glycolysis) and vimentin (red; marker to identify

osteosarcoma cells). Magnified GLUT1 staining is shown for the boxed regions on the whole-

section images. Lesion edges are indicated by white outlines in the magnified regions. Tumors

showed a high degree of intra-tumor variation in GLUT1 staining intensity. C) FeaturePlots for

the Glycolysis module score in patient primary tumor datasets. The module score for Glycolysis

was calculated using the 'AddModuleScore' function in Seurat with msigdb

HALLMARK_GLYCOLYSIS genes as input features for expression program.

*Figure 5*D, see supplementary *Figure S33*). This suggested that the process leading to clonal

expansion might come from the triggering of specific gene programs rather than the selection of

cells with any particularly fit phenotype. To identify gene programs that might be activated within

these expanding clones, we performed pathway enrichment analysis using the genes

differentially upregulated in the top 10 expanded clones. This analysis showed that genes

associated with hypoxia, glycolysis, EMT and MTORC1 signaling were upregulated within the

expanding clones, suggesting a potential mechanism for the metabolic heterogeneity noted

earlier, whereas differentially downregulated genes were associated with interferon response

pathways (*A,* B) Immunofluorescence staining of mouse lungs bearing OS-17 and 143B lung-

colonizing tumors, respectively, for GLUT1 (green; a marker of glycolysis) and vimentin (red;

marker to identify osteosarcoma cells). Magnified GLUT1 staining is shown for the boxed

regions on the whole-section images. Lesion edges are indicated by white outlines in the

magnified regions. Tumors showed a high degree of intra-tumor variation in GLUT1 staining

intensity. C) FeaturePlots for the Glycolysis module score in patient primary tumor datasets. The

module score for Glycolysis was calculated using the 'AddModuleScore' function in Seurat with

msigdb HALLMARK_GLYCOLYSIS genes as input features for expression program.

*Figure 5*E-F*,* see *Table S3*, for differentially upregulated genes, see supplementary *Figure S3*C

for downregulated pathways).

**Discussion**

These studies provide insight into the roles that intra-tumor heterogeneity and clonal evolution

play as osteosarcoma tumors adapt to bone and lung microenvironments. Using an orthotopic

cell line and PDX models of osteosarcoma, we show that these tumor cells maintain

transcriptional heterogeneity as they respond to changing microenvironment conditions while

also plastically adapting their transcriptional profiles. We identified that tibia and lung tumors

demonstrate phenotypic heterogeneity where subsets of cells upregulate genes associated with

distinct pathways, including glycolysis, hypoxia, EMT, and TNFα signaling via NFκB. We

validated this functional heterogeneity for the identified pathways in publicly available patient

primary tumor datasets. Furthermore, we showed that intratumoral heterogeneity within at least

one metabolically relevant pathway was reproducible using immunofluorescent staining for key

proteins, suggesting heterogeneity in glycolytic metabolism within individual metastatic lesions,

irrespective of lesion size. Finally, by combining lineage tracing and single-cell transcriptomic

analysis, we showed that rapidly expanding clones emerge from cells across the transcriptional

landscape, each of which adopts a growth-associated sub-phenotype, which is at least

associated with the activation of pathways also associated with metabolism. Together, our

studies suggest a tumor-intrinsic mechanism that allows for maintenance of phenotypic

diversity, despite clonal selection, while undergoing adaptive transcriptional changes during lung

colonization (**Figure 6**). These observations implicate several transcriptional programs in the

process of lung metastasis, including several pathways linked to energy metabolism. Most

interestingly, this data suggests potential cooperation between distinct tumor subpopulations,

which could be a driver for the maintenance of heterogeneity and lung tropism. Further study

will be necessary to determine the functional importance of these pathways.

Developmental processes in normal cells have been shown to require and maintain a

fundamental level of phenotypic heterogeneity[27,28]. Congruently, osteoblast cells grown in cell

culture demonstrated a baseline level of transcriptional heterogeneity. In our studies,

osteosarcoma cells maintained an even higher degree of transcriptional heterogeneity, whether

grown in mice or cell culture, and maintained that same degree of heterogeneity when exposed

to selective pressures of distinct tissue environments. While more samples are needed to

capture the true heterogeneity in osteosarcoma, one might speculate that the normal

developmental programs that support normal bone growth are maintained (or even amplified) in

tumor cells and that these cooperative mechanisms could produce a survival advantage in a

transformed malignant state. Indeed, studies published using patient tumor tissues support this

hypothesis[16,17] and analysis of those published data sets identifies heterogeneity in the same

pathways that we identified in our mouse models.

While pseudo-bulk analyses identified shared pathways associated with tibia and lung

colonization across the models, single cell analysis identified that each tumor was composed of

phenotypically distinct subsets associated with unique combinations of activated pathways. In

this study, we validated findings of heterogeneity in glycolytic activation from our sequencing

experiments by identifying correlated staining patterns for four distinct markers of glycolysis

activation, namely GLUT1, HK2, CA9, MCT4[25], along with a known upstream regulator of

GLUT1, MYC[26]. We showed that this heterogeneity in glycolysis activation is intrinsic to

individual metastatic lesions and not an artifact created by spatially isolated, metabolically

distinct lesions. Strong correlation between stains for each of these five markers confirmed that

the observed heterogeneity did not arise from stochastic gene expression changes. These

findings also suggest that a transition to glycolytic metabolism occurs within subsets within a

tumor. Reliance on glycolytic metabolism is a deeply-studied phenomenon in cancer cells,

where rapid energy production can support the abnormally high needs of malignant

transformation[29,30]. Our findings align with those reported in recent single cell sequencing

studies in other cancers, which also showed heterogeneous activation of glycolytic pathways[31].

This may suggest that tumor cells with distinct phenotypes leverage some type of metabolic

cooperation, with each cell specializing to perform a specific set of necessary tasks while relying

on the activity of cells with different phenotypes for needs that they cannot meet alone. Indeed,

many examples of metabolic cooperation between tumor cells and the surrounding stroma have

been documented. Evidence within the remote literature hints that similar processes may be

active within cultures of tumor cells[32,33]. Whatever the biologic advantage provided by this

heterogeneity, these tumors clearly resemble something more like a developing tissue than a

simple clonal proliferation of malignant cells. While tumors have long been described as

complex organs, based on the myriad essential interactions between tumor and host cells within

the microenvironment[34], studies presented here lend support for the idea that interactions

between tumor cells with distinct phenotypes may also be critical for tumor development.

Combining the power of scRNA-seq with lineage tracing allowed us to make inferences about

the roles of clonal selection and transcriptional plasticity in the emergence of traits that facilitate

lung colonization. We showed that lung colonization enriches for multiple clonal families,

wherein clonally related cells tend to have similar transcriptional profiles. In contrast, tumor cells

retained clonal diversity on tibia colonization, suggesting that this environment is relatively

permissive—at least to cells previously growing in culture.

It should be noted that PDXs and patient derived cell lines may not fully recapitulate the entire

spectrum of heterogeneity within a patient tumor. However, studies evaluating the rate of

evolution within PDX models over time suggest that the low passage PDXs utilized in this study

should retain a high degree of fidelity to the original tissue used to generate the models[35,36].

We acknowledge that the experimental metastasis model chosen for these studies produces

results that are agnostic of the multi-step process required to effect metastasis in a patient. This

choice was deliberate and designed to very simply ask whether the capacity to colonize these

different tissues is contained within a few rare clones with a pre-existing phenotype or whether

diverse groups of cells have the capacity to adapt to stresses imposed by the tibia or lung

environment. Our results suggest that both answers have some truth. The drastic shift in

phenotype from cell culture or flank tumors to bone or lung tumors endorses the strong role that

microenvironmental cues play in shaping tumor cell behavior.

However, lung colonization enriched for specific clonal populations that all demonstrated a

similar growth-associated phenotype. Genes differentially upregulated in the top ten enriched

clonal phylogenies overlapped mostly with genes related to hypoxia and glycolysis, a finding

that may inform the interpretation of the staining patterns seen in our gene signature validation

studies. Indeed, activation of pathways associated with hypoxia and glycolysis is closely linked

and may be important in generating the energy and building blocks needed to support the rapid

proliferation of these tumor cells. While our results cannot definitively show whether the

enriched clones composed individual lung-colonized lesions, or whether the same clones

colonized multiple lesions independently, the immunofluorescence stains for glycolysis-related

gene performed earlier suggest that this metabolic heterogeneity is a phenomenon intrinsic to

this process that likely occurs diffusely. Whether this transition represents a developmental step

that occurs within every lesion as it begins to expand rapidly, what changes (likely epigenetic)

constitute the trigger that facilitates clonal expansion, and whether the conservation of

transcriptional profiles in cells that originated from a common ancestor has an underlying

genetic or epigenetic basis all remain points of active study.

While the overall differential expression analysis identified genes associated with TNFα

signaling via NFκB and EMT to be most significantly upregulated upon lung colonization (*Figure*

*2*C), the comparison between the most enriched clones and non-expanded clones identified a

distinct transcriptional program associated with activation of glycolysis and hypoxia related

genes. These data may suggest that lung colonization follows a two-step process—the first

characterized by survival and niche establishment, with a second occurring through rapid

expansion of a subset of cells that adopt a growth-associated phenotype distinct from the

remaining bulk of the lung-colonizing tumor cells.

**Conclusions**

Overall, we show that osteosarcoma tumors maintain a high level of intra-tumoral transcriptional

heterogeneity, even while experiencing broad transcriptional changes as tumors adapt to tibia

and lung microenvironments. Distinct tumors maintain subpopulations of cells characterized by

activation of gene expression pathways that demonstrate similarities across models within each

environment. The process of lung colonization enriches for distinct clonal populations.

Interestingly, while a particular transcriptional signature appears common to rapidly expanding

clonal phylogenies, these arise from groups of cells with diverse genome-wide transcriptional

profiles that are maintained within each phylogeny. Additionally, maintenance of the

range/degree of intra-tumor heterogeneity endorses the role of cell-autonomous mechanisms in

holding up a basal evolutionary advantage associated with maintaining diverse transcriptional

states[37]. How this transcriptional heterogeneity is regulated in osteosarcoma cells, and whether

underlying genomic heterogeneity engenders transcriptional heterogeneity remain unanswered

questions.

**Methods**

**Experimental model - PDXs, Cell lines and murine studies.**

*Patient derived xenografts (PDXs).* OS-17 PDX tissue was obtained from a primary femur

biopsy performed at St. Jude's Children's Research Hospital in Memphis and was a gift from

Peter Houghton[38]. Patient-derived samples, NCH-OS-2 and NCH-OS-7, were obtained from

patients consented under an Institutional Review Board (IRB)-approved protocol IRB11-00478

at Nationwide Children's Hospital (Human Subject Assurance Number 00002860).

*Cell lines.* OS-17 cells were derived from an early passage of the OS-17 PDX (see above).

These cells were grown in RPMI (Corning, 10-040-CV) supplemented with 10% fetal bovine

serum (FBS). 143B cells were obtained from the American Type Culture Collection (ATCC,

CRL-8303) and grown in DMEM (Corning, 10-013-CV) supplemented with 10% FBS. Human

osteoblast cells (hFOB 1.19) were obtained from the American Type Culture Collection (ATCC,

CRL-11372) and grown in DMEM/F12 1:1 (Gibco, 21041-025) supplemented with 0.3 mg/ml

G418 and 10% FBS at 34°C. All cell lines were tested annually for both STR genotyping and

mycoplasma contamination using testing services provided by Genetica.

*Murine Studies.* All animal studies were approved by Nationwide Children's Hospital Institutional

Animal Care and Use Committee (IACUC protocols AR15-00022 and AR14-00045). *Flank*

*tumors.* Cryopreserved viable tissue pieces from OS-17, 143B, NCH-OS-2 and NCH-OS-7 PDX

tumors were placed in the right flank of recipient C.B-17/lcrHsd-Prkdc$^{scid}$ mice (Envigo,

Frederick, MD). These subcutaneous tumors were allowed to grow to 300 mm$^3$ before excision.

These were then prepped for single-cell RNA-seq. *Orthotopic primary tumors.* Single cell

suspensions of $5x10^5$ cells of OS-17, 143B, NCH-OS-2 and NCH-OS-7 were injected intra-

tibially in C.B-17/lcrHsd-Prkdc$^{scid}$ mice as per IACUC guidelines. Primary tumors were excised

once they grew to 800 mm$^3$ and prepped for single-cell RNA-seq. *Experimental metastasis.*

Single cell suspensions of $1x10^6$ cells of OS-17, 143B, NCH-OS-2 and NCH-OS-7 were injected

intra-venously in C.B-17/lcrHsd-Prkdc$^{scid}$ mice. Lungs were harvested once these mice reached

endpoint. Endpoint criteria for euthanasia was defined as weight loss of >10% or a body

condition score (BCS) of <10. For fluorescence immunohistochemistry studies, tumors were

placed in 10% formalin for 24 hours at 4°C, then moved to (phosphate-buffered saline) PBS for

at least one hour at 4°C. Tissues were then placed in tissue cassettes and placed in fresh PBS.

Tissues were embedded in paraffin.

**Fluorescent immunohistochemistry**

Primary antibodies against glucose transporter 1 (Abcam, ab15309, 1:200), hexokinase 2

(Abcam, [3D3] ab104836, 1:10), carbonic anhydrase 9 (Abcam, [2D3] ab107257, 1:100), c-myc

(Invitrogen, [9E10] MA1-980, 1:100), monocarboxylate transporter 4 (Santa Cruz Biotechnology,

[D-1] sc-376140, 1:50) and vimentin (Cell Signaling Technology, [D21H3] 5741, 1:100 and

Abnova, [SRL33] MAB9596, 1:200), were used. Secondary antibodies and counterstain used

were donkey anti-rabbit Alexa Fluor 488 (Invitrogen, A21206, 1:500), donkey anti-mouse Alexa

Fluor 488 (Invitrogen, A21202, 1:500), donkey anti-rabbit Alexa Fluor 568 (Invitrogen, A10042,

1:500), donkey anti-mouse Alexa Fluor 568 (Invitrogen, A10037, 1:500), and DAPI (4, 6-

diamidino-2-phenylindole dihydrochloride; Invitrogen, D1306, 1:500).

Paraffin embedded tissues were cut into 4uM sections and placed on glass slides. The sections

were deparaffinized with xylene and rehydrated. Sections were submerged into a Tris-EDTA

solution (pH 9.0) or a citrate solution (pH 6.0) and heated for antigen retrieval. Sections were

blocked and permeabilized with a solution of PBS + 0.2% triton + 2% bovine serum albumin for

one hour at room temperature. Primary antibodies were diluted in PBS + 0.2% triton + 2%

bovine serum albumin. The blocking solution was removed from slides and primary antibodies

were applied to the sections overnight at 4°C. Sections were washed three times with PBS +

0.2% triton. All secondary antibodies and DAPI were diluted in PBS + 0.2% triton + 2% bovine

serum albumin and added to the samples for one hour at room temperature. Sections were

washed three times with PBS + 0.2% triton, and once a final time with reverse osmosis water.

Samples were mounted in an aqueous mountant (Invitrogen, 00-4958-02). Each tissue section

was co-stained for vimentin and one of the other markers of interest.

Whole-section multichannel images were captured using a Nikon Ti2-E motorized microscope

with a Lumencor SOLA LED light engine (at 50% power), a Hamamatsu ORCA Fusion camera,

and Nikon Plan Apochromat Lambda objectives using Nikon NIS-Elements AR version 5.30

software. Each sample was imaged at 10x magnification with a final 16-bit image resolution of

0.64 µm/pixel. Identical imaging settings were used for all samples within the same staining and

imaging set.

Images were analyzed using NikoNIS-Elements AR software version 5.30 with the General

Analysis 3 module. Semi-automated tumor segmentation was performed based on vimentin

staining, and the segmented regions were manually modified by an unbiased operator as

needed to ensure specific and complete tumor selection. Images were then rescaled to 1.28

µm/pixel and the intensities of all pixels representing tumor tissue were recorded in a frequency

table for each sample. For assessing correlation between non-vimentin markers, image

channels representing each marker of interest from different tissue sections were aligned using

the multimodal image registration tool. Following alignment, an intensity profile line of 2-3 mm in

length was plotted through tumor regions showing diverse signal landscapes, and the intensities

of pixels along the profile line were recorded for each marker. All marker intensities were plotted

by distance along the profile line in GraphPad Prism software version 9.0.0, and the curves

were smoothed using a $2^{nd}$ order polynomial based on 50 neighboring points (25 on each side)

to account for minor mismatches in image channels from different tissue sections. The Pearson

correlation of each pair of markers was then calculated from the smoothed XY curves.

**Single-cell RNA-Seq**

Tumors and lungs harvested from mice were processed using the human tumor dissociation kit

(Miltenyi Biotec, 130-095-929) with a GentleMacs Octo Dissociator with Heaters (Miltenyi

Biotec, 130-096-427). Single cell suspensions in 0.04% BSA-PBS of cell lines, dissociated

tumor and lung tissues were generated and run on the Chromium Single Cell 3'RNA-sequencing

system (10x Genomics) with the Reagent Kit v3.1 (10XGenomics, PN-1000121) according to

the manufacturer's instructions. Briefly, cells were loaded into Chromium Next GEM Chip G

Single Cell Kit (10x Genomics, PN-1000120) with a targeted cell recovery of 5,000 cells per

sample. After performing cDNA purification, amplification, and library construction as instructed,

we sequenced sample libraries on a half lane of HS4000 (Illumina) to yield (after quality control)

about 65,000 paired-end reads per cell. For samples that contained lineage tracing barcodes

0.9 uL (100uM) of Cellecta FSeqRNA-BC14 primer was added into the sample index pcr

reaction.

**Cellular Barcoding**

Barcoded lentivirus libraries were synthesized using CloneTracker XP™ 10M Barcode-3' Library

with Venus-Puro (plasmid) (Cellecta, BCXP10M3VP-P) as described in Wang and McManus

2009 [39] with polyethylenimine (PEI), (Alfa Aesar, 43896) as the transfection reagent. OS-17

cells were infected with in-house prepared virus library in the presence of Polybrene (8 µg/ml)

(MilliporeSigma, TR1003G). The barcoded plasmids contain ~10M 38-bp semi-random

oligonucleotide sequence that is captured on the Chromium Single Cell 3'RNA-sequencing

system. 48 hours post-infection, OS-17 cells were used to generate orthotopic primary and

metastatic tumor models as described above.

**Cellular Barcoding Computational Analysis**

Raw sequencing data was pre-processed to extract the lineage tag (LT) using known flanking

sequences, and the matching cell ID (CID). Lines without high-confidence CID assignment were

removed and redundant reads were removed. The extracted LT barcode reads were matched

against the known Cellecta barcode library. Reads with barcodes that didn't match the Cellecta

library were eliminated. The LT barcode for each matching CID was integrated into the SEURAT

object metadata to allow for further analysis (see below).

**Single-cell RNA-Seq analysis**

Cell Ranger version 3.0.2 (10x Genomics) was used to convert Illumina BCL files to FASTQ

files. These FASTQ files were then de-multiplexed and aligned to the hg19 human reference

genome provided by 10X Genomics, to generate gene-cell matrices. We used the SEURAT R

package[40–42] for quality control, filtering, and analysis of the data. Cells were filtered to remove

doublets (outliers with high count and high genes per cell), low quality cells (outliers with low

count and low genes per cell), and cells with high mitochondrial genes (indicative of cells with

broken membrane). Cells with fewer than 800 expressed genes and genes expressed in fewer

than 5 cells were filtered out. The total numbers of cells in each model after filtering were as

follows: Osteoblasts (cell culture): 5735; OS-17 (cell culture): 3,327; OS-17 (tibia): 4,574; OS-17

(lung): 2,849; 143B (cell culture): 3,354; 143B (tibia): 6,187; 143B (lung): 5,134; NCH-OS-2

(flank): 6,749; NCH-OS-2 (tibia): 4,285; NCH-OS-2 (lung): 1,645; NCH-OS-7 (flank): 1,998;

NCH-OS-7 (tibia): 2,891; NCH-OS-7 (lung): 5,133. In analyses that required comparison across

conditions or models, samples were subset to contain equal numbers of cells and this number is

specified in the figure legends. We transformed and normalized unique molecular identifier

(UMI) counts using the 'NormalizeData' function in SEURAT package version 4.0.2[43] with

default parameters. We mitigated the effects of cell cycle heterogeneity by regressing out

canonical G2/M- and S-Phase genes using the 'ScaleData' function in SEURAT. Principle

component analysis (PCA) was performed using the top 2,000 highly variable genes identified

by the SEURAT function 'FindVariableFeatures' with default parameters. For each dataset, the

first twenty principal components were selected based on the elbow plot for percentage

explained variances, representing ~55.5–60.3% of total variances. The UMAP transformation[44]

was performed on selected principal components using the 'RunUMAP' function. We applied

shared nearest neighbor (SNN) modularity optimization[45] for clustering. We calculated silhouette

scores for different number of clusters (from two clusters to ten clusters) to measure how similar

one cell was to other cells in its own cluster compared with other clusters using the silhouette

function in the R package cluster. We determined the optimum number of clusters for each

sample by maximizing the Silhouette score. Datasets were integrated using four different

approaches (FastMNN[46], SCTransform[47], Harmony[48], merge Seurat objects[49]) to determine the

optimal method. PCA using the expression of cell cycle genes identified that integrating

datasets with 'merge' function in SEURAT or application of Harmony gave consistent results

where cells did not cluster by cell cycle. Marker genes were identified for each cluster relative to

other clusters using the 'FindMarkers' function, returning only positive markers, with 'test.use'

set to DESeq2[50] which uses a negative binomial distribution. Pathway enrichment analysis was

performed on these marker genes using the enricher function in the R package clusterProfiler[51]

with default parameters with msigdb Hallmark gene sets[52]. Pseudo-bulk analysis was performed

by setting identities of each cell to the sample. We identified gene differentially regulated using

the 'FindMarkers' function for both positive and negative markers upon tibia or lung colonization.

Pathway enrichment analysis was performed on these markers as described above. The ITH

score was calculated as described in Stewart et al.[18] Briefly, it was defined as the average

Euclidean distance between each cell to every other cell in each sample, in terms of the

selected principal components.

We downloaded validation datasets from Gene Expression Omnibus (GEO) for seven patient

primary tumor tissues published in a recent study (GEO accession number-GSE152048)[16]. We

used the Seurat R package[40–42] for quality control, filtering, and analysis of these gene

expression matrices. Cells were filtered to remove doublets, low quality cells, and cells with high

mitochondrial genes as described above. Cells with fewer than 800 expressed genes and genes

expressed in fewer than 5 cells were filtered out. We transformed and normalized UMI counts

using the 'NormalizeData' function in SEURAT package version 4.0.2[43] with default parameters.

Principle component analysis (PCA) was performed using the top 2,000 highly variable genes

identified by the SEURAT function 'FindVariableFeatures' with default parameters. Using the

marker gene sets used in the paper[16], we separated tumor cells from patient host cells. For

each sample, the first twenty principal components were selected based on the elbow plot for

percentage explained variances, representing ~51.8–61.5% of total variances. The UMAP

transformation[44] was performed on selected principal components using the 'RunUMAP'

function. Module scores for Glycolysis, Hypoxia, EMT, TNFα signaling via NFκB were calculated

using the 'AddModuleScore' function in SEURAT with corresponding Hallmark msigdb gene

sets as input features for the expression program.

**Statistical analysis**

Statistical analyses were performed using R software environment for statistical computing[53]

and Prism 9 (GraphPad Software, Inc.). The packages used in R software are mentioned

through the text in the methods section. For differential gene expression analysis, a negative

binomial model (DESeq2)[50] was used. For multiple testing, p values were adjusted using

Benjamini Hochberg (BH) correction. For ITH score comparisons, data were subjected to one-

way analysis of variance (ANOVA) followed by Šidák multiple comparisons test.

**Abbreviations**

**PDX:** patient-derived xenograft

**ITH:** intra-tumor heterogeneity

**EMT:** epithelial to mesenchymal transition

**MTORC1:** mechanistic target of rapamycin (mTOR) complex 1.

**PI3K:** Phosphoinositide 3-kinase

**AKT:** Protein kinase B

**TGF:** Transforming growth-factor

**TNFα:** tumor necrosis factor α

**NFKB:** Nuclear Factor kappa-light-chain-enhancer of activated B cells

**Declarations**

**Ethics approval and consent to participate**

Patient-derived samples were obtained from patients consented under an Institutional Review

Board (IRB)-approved protocol IRB11-00478 at Nationwide Children's Hospital (Human Subject

Assurance Number 00002860).

All animal studies were approved by Nationwide Children's Hospital Institutional Animal Care

and Use Committee (IACUC protocols AR15-00022 and AR14-00045).

**Consent for publication**

Not applicable

**Availability of data and materials**

The datasets supporting the conclusions of this article are available in the NCBI's Gene

Expression Omnibus (GEO) repository,

[https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE179681].

**Code availability**

The bioinformatics analyses were performed using open-source software, including SEURAT

version 4.0.2[43], clusterProfiler version 3.14.3[51], and Cell Ranger version 3.0.2. All code is

deposited in https://github.com/kidcancerlab/OSHetero2021.

**Competing interests**

The authors declare that they have no competing interests.

**Acknowledgements**

We would like to thank our funding sources. This work was generously supported by funding

provided by NIH/NCI (K08CA201638, RDR), St Baldrick's Foundation Scholar Award (RDR),

Hyundai Hope on Wheels Young Investigator Award (RDR), CancerFree KIDS Foundation

(RDR), Steps for Sarcoma Foundation (RDR), Sarcoma Foundation of America (RDR), a

Pelotonia Fellowship (SR), a Nationwide Children's Director's Strategic Development Fund, and

an NIH CTSA Grant UL1TR002733.

**Authors' contributions**

SR and RDR conceived and designed the study. SR and RDR developed the methodology. MC

and ACG prepared libraries for next-generation sequencing. SR, EF, CT, MW, MVC and AO

performed computation analysis. CAM performed immuno-fluorescence staining. TAV

supported image acquisition and analysis of immunofluorescence experiments. SR and RDR

interpreted all the consolidated data and wrote the manuscript. SR, EF, CAM and RDR edited

the manuscript. SR, EF and RDR revised the manuscript. All authors read and approved the

final manuscript.

**References**

1. Casali, P. G. *et al.* Bone sarcomas: ESMO-PaedCan-EURACAN Clinical Practice
Guidelines for diagnosis, treatment and follow-up. *Annals of Oncology* **29**, iv79–iv95
(2018).

2. Savage, S. A. & Mirabello, L. Using epidemiology and genomics to understand
osteosarcoma etiology. *Sarcoma* **2011**, 14–16 (2011).

3. Lindsey, B. A., Markel, J. E. & Kleinerman, E. S. Osteosarcoma Overview. *Rheumatology*
*and therapy* **4**, 25–43 (2017).

4. Geller, D. S. & Gorlick, R. Osteosarcoma: a review of diagnosis, management, and
treatment strategies. *Clinical advances in hematology & oncology : H&O* **8**, 705–718
(2010).

5. Link, M. P. *et al.* The Effect of Adjuvant Chemotherapy on Relapse-Free Survival in
Patients with Osteosarcoma of the Extremity. *New England Journal of Medicine* **314**,
1600–1606 (1986).

6. Fan, T. M., Roberts, R. D. & Lizardo, M. M. Understanding and Modeling Metastasis
Biology to Improve Therapeutic Strategies for Combating Osteosarcoma Progression.
*Frontiers in Oncology* vol. 10 Preprint at https://doi.org/10.3389/fonc.2020.00013 (2020).

7. Aljubran, A. H., Griffin, A., Pintilie, M. & Blackstein, M. Osteosarcoma in adolescents and
adults: Survival analysis with and without lung metastases. *Annals of Oncology* **20**, 1136–
1141 (2009).

8. Khanna, C. *et al.* Toward a drug development path that targets metastatic progression in
osteosarcoma. *Clinical cancer research : an official journal of the American Association*
*for Cancer Research* **20**, 4200–4209 (2014).

9. Gambera, S. *et al.* Clonal dynamics in osteosarcoma defined by RGB marking. *Nature*
*Communications* **9**, (2018).

10.    OS, B., H, H., G, S., S, S. & O, F. Hematogenous micrometastases in osteosarcoma
patients. *Clinical cancer research□: an official journal of the American Association for*
*Cancer Research* **11**, 4666–4673 (2005).

11.    S, V. & J, M. Origins of metastatic traits. *Cancer cell* **24**, 410–421 (2013).

12.    S, V. & RA, W. Tumor metastasis: molecular insights and evolving paradigms. *Cell* **147**,
275–292 (2011).

13.    S, T. & C, S. Metastasis as an evolutionary process. *Science (New York, N.Y.)* **352**, 169–
175 (2016).

14.    Sethi, N. & Kang, Y. Unravelling the complexity of metastasis — molecular understanding
and targeted therapies. *Nature Reviews Cancer 2011 11:10* **11**, 735–748 (2011).

15.    M, C. *et al.* Circulating tumor cells, disease progression, and survival in metastatic breast
cancer. *The New England journal of medicine* **351**, 781–791 (2004).

16.    Zhou, Y. *et al.* Single-cell RNA landscape of intratumoral heterogeneity and
immunosuppressive microenvironment in advanced osteosarcoma. *Nat Commun* **11**, 1–
17 (2020).

17.    Liu, Y. *et al.* Single-Cell Transcriptomics Reveals the Complexity of the Tumor
Microenvironment of Treatment-Naive Osteosarcoma. *Front Oncol* **0**, 2818 (2021).

18.    Chalishazar, M. D. *et al.* Single-cell analyses reveal increased intratumoral heterogeneity
after the onset of therapy resistance in small-cell lung cancer. *Nature cancer* (2020).

19.    Yu, X., Yustein, J. T. & Xu, J. Research models and mesenchymal/epithelial plasticity of
osteosarcoma. *Cell & Bioscience 2021 11:1* **11**, 1–12 (2021).

20.    Sun, X. & Yu, Q. Intra-tumor heterogeneity of cancer cells and its implications for cancer
treatment. *Acta Pharmacologica Sinica 2015 36:10* **36**, 1219–1227 (2015).

21.    Yoshioka, H. *et al.* Single-Cell RNA-Sequencing Reveals the Breadth of Osteoblast
Heterogeneity. *JBMR Plus* **5**, (2021).

22.    Ridout, M. S. & Linkie, M. Estimating overlap of daily activity patterns from camera trap
data. *Journal of Agricultural, Biological, and Environmental Statistics 2009 14:3* **14**, 322–
337 (2009).

23.    Lin, M., Lucas, H. C. & Shmueli, G. Research Commentary—Too Big to Fail: Large
Samples and the p-Value Problem. *https://doi.org/10.1287/isre.2013.0480* **24**, 906–917
(2013).

24.    Sayles, L. C. *et al.* Genome-informed targeted therapy for osteosarcoma. *Cancer Discov*
**9**, 46–63 (2019).

25.    Kim, M.-J., Kim, D.-H., Jung, W.-H. & Koo, J.-S. Expression of metabolism-related
proteins in triple-negative breast cancer. *International journal of clinical and experimental*
*pathology* **7**, 301–312 (2014).

26.    CV, D., A, L. & P, G. MYC-induced cancer cell energy metabolism and therapeutic
opportunities. *Clinical cancer research□: an official journal of the American Association*
*for Cancer Research* **15**, 6479–6483 (2009).

27.    Altschuler, S. J. & Wu, L. F. Cellular Heterogeneity: Do Differences Make a Difference?
*Cell* vol. 141 559–563 Preprint at https://doi.org/10.1016/j.cell.2010.04.033 (2010).

28.    Chang, H. H., Hemberg, M., Barahona, M., Ingber, D. E. & Huang, S. Transcriptome-wide
noise controls lineage choice in mammalian progenitor cells. *Nature* **453**, 544–547
(2008).

29.    RJ, D., JJ, L., G, H. & CB, T. The biology of cancer: metabolic reprogramming fuels cell
growth and proliferation. *Cell metabolism* **7**, 11–20 (2008).

30.    PP, H. & DM, S. Cancer cell metabolism: Warburg and beyond. *Cell* **134**, 703–707
(2008).

31.    Xiao, Z., Dai, Z. & Locasale, J. W. Metabolic landscape of the tumor microenvironment at
single cell resolution. *Nature Communications 2019 10:1* **10**, 1–12 (2019).

32.    Gupta, S., Roy, A. & Dwarakanath, B. S. Metabolic cooperation and competition in the
tumor microenvironment: Implications for therapy. *Front Oncol* **7**, 68 (2017).

33.    Rattigan, Y. I. *et al.* Lactate is a mediator of metabolic cooperation between stromal
carcinoma associated fibroblasts and glycolytic tumor cells in the tumor
microenvironment. *Exp Cell Res* **318**, 326–335 (2012).

34.    Egeblad, M., Nakasone, E. S. & Werb, Z. Tumors as organs: Complex tissues that
interface with the entire organism. *Dev Cell* **18**, 884–901 (2010).

35.    Rajan, S. *et al.* Structurally complex osteosarcoma genomes exhibit limited heterogeneity
within individual tumors and across evolutionary time. *bioRxiv* 2021.08.30.458268 (2022)
doi:10.1101/2021.08.30.458268.

36.    Sun, H. *et al.* Comprehensive characterization of 536 patient-derived xenograft models
prioritizes candidates for targeted treatment. *Nature Communications 2021 12:1* **12**, 1–20
(2021).

37.    Marusyk, A. & Polyak, K. Tumor heterogeneity: causes and consequences. *Biochimica et*
*biophysica acta* **1805**, 105–117 (2010).

38.    Houghton, P. J. *et al.* The pediatric preclinical testing program: Description of models and
early testing results. *Pediatr Blood Cancer* (2007) doi:10.1002/pbc.21078.

39.    Wang, X. & McManus, M. Lentivirus production. *Journal of visualized experiments□:*
*JoVE* (2009) doi:10.3791/1499.

40.    Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of
single-cell gene expression data. *Nature Biotechnology* **33**, 495–502 (2015).

41.    Haber, A. L. *et al.* A single-cell survey of the small intestinal epithelium. *Nature* **551**, 333–
339 (2017).

42.  Macosko, E. Z. *et al.* Highly parallel genome-wide expression profiling of individual cells
using nanoliter droplets. *Cell* **161**, 1202–1214 (2015).

43.  Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e21
(2019).

44.  McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and
Projection for Dimension Reduction. (2018).

45.  Waltman, L. & van Eck, N. J. A smart local moving algorithm for large-scale modularity-
based community detection. *The European Physical Journal B 2013 86:11* **86**, 1–14
(2013).

46.  Haghverdi, L., Lun, A. T. L., Morgan, M. D. & Marioni, J. C. Batch effects in single-cell
RNA-sequencing data are corrected by matching mutual nearest neighbors. *Nature
Biotechnology 2018 36:5* **36**, 421–427 (2018).

47.  Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-
seq data using regularized negative binomial regression. *Genome Biology 2019 20:1* **20**,
1–15 (2019).

48.  Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with
Harmony. *Nature Methods 2019 16:12* **16**, 1289–1296 (2019).

49.  merge: Merge Two Data Frames. https://rdrr.io/r/base/merge.html.

50.  Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biology 2014 15:12* **15**, 1–21 (2014).

51.  Yu, G., Wang, L. G., Han, Y. & He, Q. Y. ClusterProfiler: An R package for comparing
biological themes among gene clusters. *OMICS A Journal of Integrative Biology* **16**, 284–
287 (2012).

52.  A, L. *et al.* Molecular signatures database (MSigDB) 3.0. *Bioinformatics (Oxford,
England)* **27**, 1739–1740 (2011).

53.  Foundation, T. R. The R Project for Statistical Computing. https://www.r-project.org/.

**Figure legends**

*Figure 1.* **Cell line and PDX models of osteosarcoma display transcriptional**

**heterogeneity.** A) UMAP analysis of OS-17 cells (n = 3327) grown in cell culture. Cell cycle

distribution of cells (n = 3327) is visualized as pie charts. B) UMAP analysis of NCH-OS-7 cells

(n = 1998) grown as subcutaneous flank tumor. Cell cycle distribution of cells is visualized as

pie charts. C) Pathway enrichment analysis for hallmark gene sets associated with genes

upregulated in distinct clusters identified in the two osteosarcoma models. P values were

adjusted for multiple comparisons. Boxes in grey identify non-significant enrichments, whereas

boxes in red identify statistically significant enrichments. MTORC1: mechanistic target of

rapamycin (mTOR) complex 1. PI3K: Phosphoinositide 3-kinase. AKT: Protein kinase B. TGF:

Transforming growth-factor. TNF: tumor necrosis factor. NFKB: Nuclear Factor kappa-light-

chain-enhancer of activated B cells.

***Figure 2.* Osteosarcoma cells adopt distinct transcriptional profiles as they colonize tibia**

**and lung microenvironments.** A) Schematic of study design depicting generation of orthotopic

tibia- and lung-colonizing tumors. Tumors were harvested for scRNA-seq when mice reached

endpoint. B) Venn diagrams showing overlap of differentially expressed genes that are up- or

down- regulated in tibia or lung lesions relative to corresponding starting population of cells (cell

culture or PDX flank tumors). Regions outlined in red identify differentially expressed genes

shared across at least three models. C) Pathway enrichment analysis with adjusted p values for

hallmark gene sets associated with these shared genes.

***Figure 3.* Osteosarcoma cells retain phenotypic heterogeneity despite adaptive changes**

**in response to changing microenvironments.** A) Schematic outlining scRNA-seq

bioinformatics workflow. B) Osteosarcoma cells maintained overall heterogeneity with a high

degree of overlap between conditions. The ridge plot shows ITH scores, which represent the

gene expression "distance" between each tumor cell within a sample and all of the other tumor

cells from that same sample. The overlap statistic describes the total percentage of overlap in

the observed distributions between two samples. C) UMAP analysis for merged tibia- and lung-

colonizing tumor samples in each of the four models. Cells in grey represent remaining cells in

merged sample. Cluster enrichment analysis shows distribution of cells in each cluster in the

two microenvironment conditions (tibia, lung). While some cells in the tibia and lung lesions

adopted shared phenotypes, others adopted distinct phenotypes. D) Metabolic heterogeneity in

glycolysis activation. We used a pathway enrichment analysis for hallmark gene sets using

genes differentially upregulated in each cluster relative to every other cluster within the same

model. P values were adjusted for multiple comparisons. Boxes in grey identify non-significant

pathway enrichments, whereas boxes in red identify statistically significant enrichments. Bar plot

shows percentage of cells identified per cluster in lung lesions. In B-D, samples subset to equal

number of cells to allow inter-model comparison (n=1500 per condition).

*Figure 4.* **Heterogeneity in Glycolysis activation identified within lung-colonizing**

**osteosarcoma lesions.** A, B) Immunofluorescence staining of mouse lungs bearing OS-17 and

143B lung-colonizing tumors, respectively, for GLUT1 (green; a marker of glycolysis) and

vimentin (red; marker to identify osteosarcoma cells). Magnified GLUT1 staining is shown for

the boxed regions on the whole-section images. Lesion edges are indicated by white outlines in

the magnified regions. Tumors showed a high degree of intra-tumor variation in GLUT1 staining

intensity. C) FeaturePlots for the Glycolysis module score in patient primary tumor datasets. The

module score for Glycolysis was calculated using the 'AddModuleScore' function in Seurat with

msigdb HALLMARK_GLYCOLYSIS genes as input features for expression program.

*Figure 5.* **Lung colonization bottleneck enriches for clonal populations with a common**

**transcriptional phenotype.** A) Schematic of experimental set-up. B) UMAP analysis of lineage

tagged OS-17 cells in culture, implanted into tibia, or inoculated to generate lung- colonizing

tumors. These lineage tags are heritable, allowing us to track the transcriptional profiles of

daughter cells originating from the same ancestor. Lung- and tibia-colonizing tumor samples

share many cells with similar phenotypes, though cultured cells are dramatically different (n =

2800 per condition). C) Frequency distribution of clones identified in each of the conditions. A

high level of clonal diversity is maintained in the tibia-colonizing tumors (78% unique clones in

tibia-colonizing tumor compared to 86% unique clones in cell culture), while dominant clones

emerge in the lung (only 33% unique clones). D) Overlay of cells sharing a common ancestor

onto the dimensional reduction plots in the top four enriched clonal families. E) UMAP

visualization of cells belonging to top ten enriched clonal families highlighted in red from the

lung-colonizing tumor sample. F) Pathway enrichment analysis for hallmark gene sets

comparing enriched clones relative to the remaining lung-colonizing tumor cells identifies

hypoxia, glycolysis, and EMT signaling related genes to be significantly upregulated.

*Figure 6.* **Schematic of tumor cell adaptation to changing microenvironments.**

Osteosarcoma cells demonstrate high transcriptional heterogeneity (colors) in cell culture,

orthotopic tibia and lung microenvironments. Tumors that grow in the orthotopic tibia lesions

follow dynamics of neutral evolution where different cancer clones (shapes) co-exist and

propagate simultaneously. The selective pressure of lung colonization results in enrichment of

clonal families with different phenotypes (colors), where cells belonging to the same family

(shape) tend to have similar phenotypes (color).

**Additional Files**

**Additional File 1**

File format: .pdf

Title of the data: Supplemental Figures

Description of the data: Supplementary figures supporting the study can be found in this

document.

**Additional File 2**

File format: .docx

Title of the data: Supplementary Tables

Description of the data: Supplementary tables supporting the study can be found in this

document.

Figure 1

**C**

| OS-17 | | | | NCH-OS-7 | | | | |
|---|---|---|---|---|---|---|---|---|
| 0.00 | 0.00 | 0.00 | 0.00 | 0.12 | 2.93 | 0.00 | 1.23 | ADIPOGENESIS |
| 0.02 | 0.14 | 0.00 | 0.62 | 0.40 | 0.00 | 1.14 | 0.00 | ANDROGEN RESPONSE |
| 1.59 | 0.73 | 0.09 | 0.63 | 0.27 | 0.00 | 0.00 | 1.66 | ANGIOGENESIS |
| 0.26 | 0.82 | 0.00 | 0.01 | 0.00 | 0.00 | 0.00 | 0.00 | APICAL JUNCTION |
| 1.13 | 0.25 | 0.31 | 0.24 | 0.00 | 0.00 | 0.00 | 0.00 | APOPTOSIS |
| 0.09 | 0.09 | 0.74 | 0.09 | 0.81 | 0.00 | 0.00 | 1.60 | CHOLESTEROL HOMEOSTASIS |
| 0.26 | 0.82 | 0.00 | 0.01 | 0.00 | 0.00 | 0.00 | 0.00 | COAGULATION |
| 0.07 | 0.01 | 0.22 | 0.84 | 0.00 | 6.36 | 0.00 | 0.00 | DNA REPAIR |
| 0.44 | 0.01 | 2.11 | 0.63 | 0.00 | 6.75 | 0.76 | 0.00 | E2F TARGETS |
| 16.46 | 7.97 | 0.00 | 1.90 | 9.65 | 0.00 | 0.31 | 5.11 | EPITHELIAL MESENCHYMAL TRANSITION |
| 0.09 | 0.00 | 0.00 | 0.24 | 0.57 | 0.00 | 0.00 | 0.00 | ESTROGEN RESPONSE LATE |
| 0.00 | 0.00 | 0.74 | 0.01 | 0.00 | 2.93 | 0.00 | 0.62 | FATTY ACID METABOLISM |
| 0.02 | 0.02 | 0.93 | 0.01 | 0.00 | 0.68 | 1.80 | 0.00 | G2M CHECKPOINT |
| 0.44 | 0.82 | 0.01 | 0.24 | 6.58 | 0.00 | 0.00 | 0.14 | GLYCOLYSIS |
| 0.00 | 0.73 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | HEDGEHOG SIGNALING |
| 1.59 | 1.31 | 0.22 | 0.01 | 13.20 | 0.00 | 1.45 | 0.54 | HYPOXIA |
| 0.00 | 0.00 | 0.74 | 0.00 | 0.00 | 0.00 | 0.00 | 1.60 | INTERFERON ALPHA RESPONSE |
| 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 1.23 | INTERFERON GAMMA RESPONSE |
| 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 10.42 | 0.00 | MITOTIC SPINDLE |
| 0.00 | 0.82 | 2.43 | 0.77 | 6.58 | 1.80 | 0.55 | 0.00 | MTORC1 SIGNALING |
| 0.09 | 0.63 | 15.34 | 0.63 | 2.91 | 39.39 | 0.65 | 5.11 | MYC TARGETS V1 |
| 0.00 | 0.00 | 0.74 | 0.00 | 0.00 | 1.65 | 0.65 | 0.00 | MYC TARGETS V2 |
| 1.59 | 0.25 | 0.00 | 0.24 | 0.00 | 0.00 | 0.00 | 0.00 | MYOGENESIS |
| 0.02 | 0.06 | 0.63 | 0.24 | 0.00 | 36.82 | 0.00 | 13.98 | OXIDATIVE PHOSPHORYLATION |
| 0.00 | 0.00 | 0.00 | 0.00 | 1.32 | 0.00 | 1.65 | 0.00 | P53 PATHWAY |
| 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.62 | 0.00 | 0.76 | PEROXISOME |
| 0.00 | 0.00 | 0.00 | 0.00 | 0.12 | 0.38 | 1.53 | 0.00 | PI3K AKT MTOR SIGNALING |
| 0.26 | 0.04 | 0.77 | 0.17 | 0.12 | 3.28 | 0.00 | 1.60 | REACTIVE OXYGEN SPECIES PATHWAY |
| 0.00 | 0.00 | 0.00 | 0.00 | 0.12 | 0.00 | 4.29 | 0.00 | TGF BETA SIGNALING |
| 0.00 | 0.00 | 0.00 | 0.00 | 0.27 | 0.00 | 1.45 | 0.00 | TNFA SIGNALING VIA NFKB |
| 0.00 | 0.00 | 0.00 | 0.00 | 3.99 | 0.00 | 4.17 | 0.00 | UNFOLDED PROTEIN RESPONSE |
| 0.90 | 0.25 | 0.00 | 0.01 | 0.12 | 0.00 | 1.65 | 0.00 | UV RESPONSE DN |
| 0.00 | 0.00 | 0.00 | 0.00 | 1.16 | 0.05 | 0.28 | 0.00 | UV RESPONSE UP |

Cluster 0 / Cluster 1 / Cluster 2 / Cluster 3 (OS-17)
Cluster 0 / Cluster 1 / Cluster 2 / Cluster 3 (NCH-OS-7)

p value of GSEA
> 0.05
≤ 0.05

Figure 1 (continued)

**Figure 1. Cell line and PDX models of osteosarcoma display transcriptional heterogeneity.** A) UMAP analysis of OS-17 cells (n = 2800) grown in cell culture. Cell cycle distribution of cells (n = 2800) is visualized as pie charts. B) UMAP analysis of NCH-OS-7 cells (n = 1998) grown as subcutaneous flank tumor. Cell cycle distribution of cells is visualized as pie charts. C) Pathway enrichment analysis for hallmark gene sets associated with genes upregulated in distinct clusters identified in the two osteosarcoma models. P values were adjusted for multiple comparisons. Boxes in grey identify non-significant enrichments, whereas boxes in red identify statistically significant enrichments. MTORC1: mechanistic target of rapamycin (mTOR) complex 1. PI3K: Phosphoinositide 3-kinase. AKT: Protein kinase B. TGF: Transforming growth-factor. TNF: tumor necrosis factor. NFKB: Nuclear Factor kappa-light-chain-enhancer of activated B cells.

Figure 2

**Figure 2. Osteosarcoma cells adopt distinct transcriptional profiles as they colonize tibia and lung microenvironments.** A) Schematic of study design depicting generation of orthotopic tibia- and lung-colonizing tumors. Tumors were harvested for scRNA-seq when mice reached endpoint. B) Venn diagrams showing overlap of differentially expressed genes that are up- or down- regulated in tibia or lung lesions relative to corresponding starting population of cells (cell line or PDX flank tumors). Region outlined in red identifies differentially expressed genes shared across at least three models. C) Pathway enrichment analysis with adjusted p values for hallmark gene sets associated with these shared genes.

Figure 3

**Figure 3. Osteosarcoma cells retain phenotypic heterogeneity despite adaptive changes in response to changing microenvironments.** A) Schematic outlining scRNA-seq bioinformatics workflow. B) Osteosarcoma cells maintained overall heterogeneity with a high degree of overlap between conditions. The ridge plot shows ITH scores, which represent the gene expression "distance" between each tumor cell within a sample and all of the other tumor cells from that same sample. The overlap statistic describes the total percentage of overlap in the observed distributions between two samples. C) UMAP analysis for merged tibia- and lung-colonizing tumor samples in each of the four models. Cells in grey represent remaining cells in merged sample. Cluster enrichment analysis shows distribution of cells in each cluster in the two microenvironment conditions (tibia, lung). While some cells in the tibia and lung lesions adopted shared phenotypes, others adopted distinct phenotypes. D) Metabolic heterogeneity in glycolysis activation. We used a pathway enrichment analysis for hallmark gene sets using genes differentially upregulated in each cluster relative to every other cluster within the same model. P values were adjusted for multiple comparisons. Boxes in grey identify non-significant pathway enrichments, whereas boxes in red identify statistically significant enrichments. Bar plot shows percentage of cells identified per cluster in lung lesions. In B-D, samples subset to equal number of cells to allow inter-model comparison (n=1500 per condition).

# Figure 4

Figure 4 (continued)

**Figure 4. Heterogeneity in glycolysis activation identified within lung-colonizing osteosarcoma lesions.**
A, B) Immunofluorescence staining of mouse lungs bearing OS-17 and 143B lung-colonizing tumors, respectively, for GLUT1 (green; a marker of glycolysis) and vimentin (red; marker to identify osteosarcoma cells). Magnified GLUT1 staining is shown for the boxed regions on the whole-section images. Lesion edges are indicated by white outlines in the magnified regions. Tumors showed a high degree of intra-tumor variation in GLUT1 staining intensity. C) FeaturePlots for a glycolysis module score in patient primary amd metastatic tumor datasets (GEO accession number-GSE152048). The module score for glycolysis was calculated using the 'AddModuleScore' function in Seurat with msigdb HALLMARK_GLYCOLYSIS genes as input features for expression program.

Figure 5

**Figure 5. Lung colonization bottleneck enriches for clonal populations with a common transcriptional phenotype.** A) Schematic of experimental set-up. B) UMAP analysis of lineage tagged OS-17 cells in culture, implanted into tibia, or inoculated to generate lung- colonizing tumors. These lineage tags are heritable, allowing us to track the transcriptional profiles of daughter cells originating from the same ancestor. Lung- and tibia-colonizing tumor samples share many cells with similar phenotypes, though cultured cells are dramatically different (n = 2800 per condition). C) Frequency distribution of clones identified in each of the conditions. A high level of clonal diversity is maintained in the tibia-colonizing tumors (78% unique clones in tibia-colonizing tumor compared to 86% unique clones in cell culture), while dominant clones emerge in the lung (only 33% unique clones). D) Overlay of cells sharing a common ancestor onto the dimensional reduction plots in the top four enriched clonal families. E) UMAP visualization of cells belonging to top ten enriched clonal families highlighted in red from the lung-colonizing tumor sample. F) Pathway enrichment analysis for hallmark gene sets comparing enriched clones relative to the remaining lung-colonizing tumor cells identifies hypoxia, glycolysis, and EMT signaling related genes to be significantly upregulated.

Figure 6

**Figure 6. Schematic of tumor cell adaptation to changing microenvironments.** Osteosarcoma cells demonstrate high transcriptional heterogeneity (colors) in cell culture, orthotopic tibia and lung microenvironments. Tumors that grow in the orthotopic tibia lesions follow dynamics of neutral evolution where different cancer clones (shapes) co-exist and propagate simultaneously. The selective pressure of lung colonization results in enrichment of clonal families with different phenotypes (colors), where cells belonging to the same family (shape) tend to have similar phenotypes (color).