1 **Molecular physiology of pumiliotoxin sequestration in a poison frog**

3 Aurora Alvarez-Buylla[1], Cheyenne Y. Payne[1], Charles Vidoudez[2], Sunia A. Trauger[2] and Lauren

4 A. O'Connell*[1]

6 1 Department of Biology, Stanford University, Stanford, CA 94305, USA

7 2 Harvard Center for Mass Spectrometry, Harvard University, Cambridge, MA 02138, USA

* To whom correspondence should be addressed:

Lauren A. O'Connell
Department of Biology
Stanford University
371 Jane Stanford Way
Stanford, CA 94305
loconnel@stanford.edu
22

ABSTRACT

Poison frogs bioaccumulate alkaloids for chemical defense from their arthropod diet. Although many alkaloids are accumulated without modification, some poison frog species can metabolize pumiliotoxin (PTX **251D**) into the more potent allopumiliotoxin (aPTX **267A**). Despite extensive research characterizing the chemical arsenal of poison frogs, the physiological mechanisms involved in the sequestration and metabolism of individual alkaloids remain unclear. We first performed a feeding experiment with the Dyeing poison frog (*Dendrobates tinctorius*) to ask if this species can metabolize PTX **251D** into aPTX **267A** and what gene expression changes are associated with PTX **251D** exposure in the intestines, liver, and skin. We found that *D. tinctorius* can metabolize PTX **251D** into aPTX **267A**, and that PTX **251D** exposure changed the expression level of genes involved in immune system function and small molecule metabolism and transport. To better understand the functional significance of these changes in gene expression, we then conducted a series of high-throughput screens to determine the molecular targets of PTX **251D** and identify potential proteins responsible for metabolism of PTX **251D** into aPTX **267A**. Although screens of PTX **251D** binding human voltage-gated ion channels and G-protein coupled receptors were inconclusive, we identified human CYP2D6 as a rapid metabolizer of PTX **251D** in a cytochrome P450 screen. Furthermore, a CYP2D6-like gene had increased expression in the intestines of animals fed PTX, suggesting this protein may be involved in PTX metabolism. These results show that individual alkaloids can modify gene expression across tissues, including genes involved in alkaloid metabolism. More broadly, this work suggests that specific alkaloid classes in wild diets may induce physiological changes for targeted accumulation and metabolism.

1. INTRODUCTION

Animals have evolved many ways of avoiding predation to enhance their survival and fitness. One strategy is evolving chemical defenses by carrying small molecules that are toxic or repellant to predators. While some organisms are able to produce their own chemical defenses, many animals accumulate and store compounds found in their environment. In some cases environmentally derived compounds are sequestered unchanged, however small molecule metabolism is also used for storage or diversification of a chemical repertoire. For example, the European cinnabar moth (*Tyria jacobaeae*) larvae convert toxic pyrrolizidine alkaloids obtained from their plant diet into inert forms for storage throughout development [1]. In plants, the expansion of cytochrome P450 genes has led to diversification of defensive alkaloids used to combat herbivores and pathogens [2]. Uncovering general principles of how organisms sequester and metabolize chemical defenses allows for a better understanding of how animals evolve to maximize the use of their available environmental chemicals.

Poison frogs (Family Dendrobatidae) are chemically defended against predators [3–5] using alkaloids sequestered from dietary arthropods [6,7]. Within the poison frog clade, the ability to sequester dietary alkaloids has evolved independently at least three times [8,9]. Notably, chemical defense covaries with the diversification of parental care strategies in this clade [10], where in some species mothers provision chemical defenses to their tadpoles by feeding alkaloid-containing trophic eggs [11–13]. Lab feeding experiments have found that alkaloids accumulate primarily on the skin, although detectable quantities are also found in the liver and intestines [14]. This accumulation can occur within a few days [15,16], and is associated with changes in gene expression and protein abundance across tissues [15,16]. However, previous analyses of gene expression have been limited to comparing frogs with alkaloids to those without alkaloids. [15-17], leaving a major gap in our understanding of how frog physiology changes in response to specific alkaloids rather than overall toxicity. Filling this

knowledge gap is important because poison frogs carry many different alkaloid classes, each of

which could potentially induce specific changes in physiology.

Controlled alkaloid-feeding experiments have been crucial in understanding alkaloid

metabolism in poison frogs [18–20]. Many poison frog alkaloids have been found in the ants and

mites they consume, suggesting most alkaloids are sequestered unchanged [21–24]. Alkaloids

of the pumiliotoxin (PTX) family are found across the poison frogs clade, including *Dendrobates,*

Oophaga, Epipedobates, Ameerega, and *Phyllobates* species, and are thought to be sourced

from formicine ants [21]. Pumiliotoxins have also been found in Malagasy poison frogs (family

Mantellidae), a convergent evolution of anuran chemical defenses found in Madagascar [25].

Along with sequestering pumiliotoxins from ants, laboratory feeding experiments have also

shown that some dendrobatid species metabolize PTX **251D** into allopumiliotoxin (aPTX) **267A**

[19]. After PTX **251D** feeding, both PTX **251D** and its metabolite, aPTX **267A,** were detected in

the skin of *Dendrobates auratus*, but only PTX **251D** was detected in the skin of *Phyllobates*

bicolor and *Epipedobates tricolor* [10]. The authors then suggested that an unidentified enzyme

performs the 7'-hydroxylation of PTX **251D** into aPTX **267A** in some dendrobatid species, but

not all. Whether other poison frog species can metabolize PTX **251D** into aPTX **267A** and the

molecular machinery responsible remains unknown.

Of the poison frog alkaloids that have been tested *in vivo* and *in vitro*, many are

considered to be toxic due to their varied effects on different ion channels in the nervous system

[5]. PTX **251D** is a toxic alkaloid that causes pain, hyperactivity, convulsions and death in mice

and insects [19,26,27]. PTX **251D** likely acts as a cardiac depressor in mice, eventually leading

to cardiac arrest [28,29]. Electrophysiology work with PTX **251D** found that it inhibited both

mammalian and insect voltage-gated sodium and potassium channels [30]. PTX **251D** potently

inhibits insect sodium channels, which may explain its repellent effect on mosquitos [27].

Additionally, the effects of PTX **251D** on mice were reduced by previous administration of

anticonvulsants phenobarbital and carbamazepine, which target calcium and sodium channels,

respectively [19]. Together, these studies provide evidence that PTX **251D** is toxic by way of

inhibition of mammalian and insect ion channels, however finer scale studies of concentration-

dependent responses and other molecular targets, such as G-protein coupled receptors, are

needed to better understand the activity of PTX **251D**. The molecular targets of aPTX **267A** are

poorly characterized, however it induces hyperactivity, convulsions, and death when injected

into mice at 2 mg/kg, a lethal dose five times lower than that of PTX **251D**. This finding suggests

that aPTX **267A** is more toxic than its precursor molecule PTX **251D**, and that the ability to

make and sequester aPTX **267A** may be an adaptive strategy to increase toxicity in some

poison frog species.

This study aimed to better understand the molecular physiology of PTX **251D**

metabolism in poison frogs and further characterize potential human molecular targets. We

conducted an alkaloid feeding study with the Dyeing poison frog (*Dendrobates tinctorius*) to test

whether this species can metabolize PTX **251D** into aPTX **267A** and to explore gene expression

changes associated with PTX **251D** exposure. We predicted metabolic enzymes involved in the

hydroxylation of PTX **251D** into aPTX **267A** may be upregulated in response to their metabolic

target. To functionally identify PTX **251D** target proteins that either metabolize PTX or alter

neuronal function, we conducted a series of high-throughput screens with human cytochrome

P450s, ion channels, and G-protein coupled receptors. This body of work provides the first in

depth examination of PTX-induced changes in poison frog physiology.

2. MATERIALS AND METHODS

2.1 Alkaloid feeding

Lab-reared (non-toxic) *Dendrobates tinctorius* were housed in terraria with live plants, a

water pool, and a shelter. Ten adult females were size-matched, randomly assigned to control

or experimental groups (N=5 per group), and then housed individually. To measure the specific

effects of PTX **251D** compared to a background toxicity, the control group was fed 0.01% DHQ

(Sigma-Aldrich, St. Louis, USA) in a solution of 1% EtOH and the experimental group was fed a

solution of 0.01% DHQ and 0.01% PTX **251D** (PepTech, Burlington, MA, USA) in a solution of

1% EtOH in water. Each frog was fed 15 µL each day for five days by pipetting the solution

directly into the mouth between 10am-12pm. On the afternoon of the fifth day, frogs were

euthanized by cervical transection and the dorsal skin, liver, intestines, and eggs were dissected

into Trizol (Thermo Fisher Scientific, Waltham, USA). All procedures were approved by the

Institutional Animal Care and Use Committee at Stanford University (protocol number #32870).

2.2 RNA extraction and library preparation

RNA extraction followed the Trizol (Thermo Fisher Scientific, Waltham, MA, USA)

protocol outlined in Caty *et al.* 2019 [31] and according to the manufacturer's instructions. After

the first spin, the organic layer was saved for alkaloid extraction (see below). Poly-adenylated

RNA was isolated using the NEXTflex PolyA Bead kit (Bioo Scientific, Austin, USA) following

manufacturer's instructions. RNA quality and lack of ribosomal RNA was confirmed using an

Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, USA). Each RNA sequencing

library was prepared using the NEXTflex Rapid RNAseq kit (Bioo Scientific). Libraries were

quantified with quantitative PCR (NEBnext Library quantification kit, New England Biolabs,

Ipswich, USA) and a Agilent Bioanalyzer High Sensitivity DNA chip, both according to

manufacturer's instructions. All libraries were pooled at equimolar amounts and were

sequenced on four lanes of an Illumina HiSeq 4000 machine to obtain 150 bp paired-end reads.

2.3 Transcriptome assembly and differential expression analysis

We created a reference transcriptome using Trinity [32] and filtered the raw assembly by

removing contigs with BLAST hits belonging to microorganisms and invertebrates in the Swiss-

Prot database [33], as these represent likely parasites, prey items, or other contaminants.

Overlapping contigs were clustered using cd-hit-est [34,35] and contigs that were less than

250bp long were removed from the assembly. We mapped the paired quality-trimmed Illumina

reads to the reference transcriptome using kallisto [36]. Samples were compared across

treatment groups (DHQ vs DHQ+PTX) for the skin, liver, and intestines, as these tissues

contained higher levels of PTX. Differences in gene expression levels were calculated using

DESeq2 [37] [$P<0.05$ false discovery rate (Benjamini–Hochberg FDR), 4-fold change]. Contigs

with significant expression differences were compared to the non-redundant (nr) database using

BLAST with an E-value cutoff of 1e-5. Many contigs did not have a BLAST hit, or aligned to

hypothetical or non-vertebrate proteins. Contigs with annotations of interest were chosen based

on candidates from existing literature. Boxplots were made with R package ggplot2 (R version

3.6.3) using TMM (trimmed mean of M-values) normalized expression. All scripts are detailed in

supplementary materials.

2.4 Alkaloid extraction and detection

To isolate alkaloids, 0.3 mL of 100% EtOH was added to 1mL of organic layer from the

Trizol RNA extraction, inverted 10 times, and stored at room temperature for 2-3 minutes to

precipitate genomic DNA, which was pelleted by centrifugation at 2000g for 5 minutes at 4°C.

Then, 300 µL of supernatant was transferred to a new microfuge tube. Proteins were

precipitated by adding 900 µL of acetone, mixing by inversion for 10-15 seconds, incubating at

room temperature for 10 min, and centrifuging at max speed for 10 min at 4°C. Then, 1 mL of

the supernatant containing alkaloids was moved into a glass vial and stored at -20°C until dried

down completely under a gentle nitrogen gas flow.

Samples were resuspended in 200 µl of methanol:chloroform 1:1 and 1 µM Nicotine-d3

(used as an internal standard). A 10-point standard curve was prepared in the same solution

with DHQ and PTX. A QE+ mass spectrometer coupled to an Ultimate3000 LC (ThermoFisher)

was used for analysis. Five µl of each sample were injected on a Gemini C18 column

(100x2mm, Phenomenex). The mobile phases were A: water and B: acetonitrile, both with 0.1%

formic acid. The gradient was 0% B for 1 min, then increased to 100% B in 14 min, followed by

5 min at 100% B and 3.5 min at 0% B. Data were quantified using accurate mass, using the

standard curve for DHQ and PTX for absolute quantification. aPTX was identified by accurate

mass and MS/MS fragmentation similarity to PTX.

2.5 Alkaloid statistical analyses

R version 3.6.3 was used for all statistical analyses, and all plotting and statistics code is

provided in a supplementary file. There were instances in the LC-MS/MS data where the

molecules of interest (DHQ, PTX **251D**, or aPTX **267A**) were not detected, and these were

converted to zeros prior to statistical analyses and visualization. A generalized linear mixed

model was used (glmmTMB package in R [38]) to test for differences in alkaloid abundance

across tissues and treatment type with the frog as a random effect, using a negative binomial

error distribution and a single zero-inflation parameter applied to all observations. PTX **251D**

and DHQ were analyzed separately. The abundance of aPTX **267A** was approximated using

the area-under-the-curve divided by the internal nicotine standard, as there is no standard for

aPTX **267A,** and therefore exact pmol values could not be calculated. A Wilcoxon rank-sum test

(wilcox.test) was used to compare the aPTX values in the skin between treatment groups and

the Kruskal-Wallis test (kruskal.test) with a post-hoc Dunn test (dunnTest from the FSA package

[61]) was used to compare the aPTX values across tissues. Boxplots used to visualize alkaloid

abundance values were created in R using ggplot.

2.6 Assays for PTX target proteins

Assays for human CYP activity (CYP phenotyping), CYP2D6 metabolite discovery

(metID), ion channel inhibition (CiPA), and G-protein coupled receptor (GPCR) activity

(gpcrMAX) were performed through Eurofins Discovery Services (Eurofins Panlabs Inc, St.

Charles MO, USA). In the CYP phenotyping panel, human recombinant CYP1A2, CYP2B6,

CYP2C8, CYP2C9, CYP2C19, CYP2D6, and CYP3A4 were included in the assay. A

concentration of 1E-07 M of PTX **251D** was tested, and the percent compound remaining was

quantified at 0, 15, 30, 45, and 60 minutes using HPLC-MS/MS. As a positive control, a

reference compound was used for each CYP tested (see supplementary files for results). In the

CYP2D6 metabolism assay, human recombinant CYP2D6 was incubated with an initial

concentration of 10 µM and metabolites were quantified after 90 minutes using LC-MS/MS.

Two replicates were used for the CYP phenotyping panel, and one replicate was used for

CYP2D6 metabolite assay. The CiPA assay was a cell-based QPatch on voltage-gated sodium

channel NaV1.5 (peak and late/agonist), voltage-gated potassium channels Kv4.3/KChIP2,

hERG, KCNQ1/minK, Kir2.1, and voltage-gated calcium channel Cav1.2. The concentrations of

PTX **251D** assayed were 3, 10, and 30 µM, and a reference compound for each channel was

used (see supplementary files for results and individual QPatch parameters). In the gpcrMAX

assay a panel of 165 GPCRs was tested through the DiscoverX PathHunter beta-arrestin

enzyme fragment complementation technology, and a concentration of 10 µM PTX **251D** was

tested for both agonist and antagonist activity along with a reference compound for each GPCR

(see supplementary files for results).

2.7 Identification and comparison of poison frog CYP2D6

The human CYP2D6 protein sequence (uniprot: P10635) was used to identify *D.*

tinctorius homologs by sequence similarity using tblastn. We then obtained the top 8 BLAST

contigs from the TMM expression matrix for each tissue. A pairwise t-test was used to

determine if the mean expression value was significantly different between treatments in each

tissue and p-values were corrected for multiple testing using benjamini-hochberg correction.

The only remaining significant difference was one contig in the gut (TRINITY_DN15846_c0_g2).

Using the largest ORF protein sequence of this gene we used tblastn against BLAST nucleotide

databases made from previously assembled unpublished transcriptomes in five other poison

frog species: *Oophaga sylvatica*, *Ranitomeya imitator*, *Epipedobates tricolor*, *Allobates*

femoralis (unpublished genome), and *Mantella aurantiaca* (Family Mantellidae, an independent

origin of chemical defense sequestration in amphibians [25]). The *D. auratus* and *P. bicolor*

sequences were found using the same method from previously published transcriptomes

[40,41]. The *D. tinctorius* TRINITY_DN15846_c0_g2 protein sequence and largest ORF of the

top BLAST hit from each of these transcriptomes was translated and aligned to the human

CYP2D6 protein sequence to identify amino acid differences in important binding residues [39].

For the *B. bufo* (XP_040266170), *R. temporaria* (XP_040185037), and *X. laevis*

(XP_031756601)   sequences,   the   nucleotide   sequence   of   *D.   tinctorius*

TRINITY_DN15846_c0_g2 was used to blastn against the NR database using the BLAST web

portal, and the protein sequence of the top hit from each species was downloaded and aligned.

Benchling software (Benchling Inc., San Francisco, CA) was used to find the largest ORFs,

translate into amino acid sequences, and create a protein alignment using Clustal Omega with

the human CYP2D6 sequence as a reference. All nucleotide sequences and amino acid

sequences are included in supplementary FASTA files.

3. RESULTS

3.1 The Dyeing poison frog metabolizes PTX to aPTX

We conducted a feeding experiment to determine if the Dyeing poison frog (*Dendrobates*

tinctorius) can metabolize PTX **251D** into aPTX **267A** (Figure 1A). Alkaloids were most

abundant in the skin and liver, followed by the intestines, and only trace amounts were detected

in eggs. DHQ abundance did not differ by treatment group (GLMM treatment, $p = 0.377$),

confirming both groups were fed equal amounts. DHQ abundance differed across tissue types

(GLMM tissue, X^2 (3) = 203.642, p < 2e-16), with the highest levels occurring in the liver and

skin (Figure 1B). PTX **251D** abundance differed by tissue and treatment (GLMM

tissue*treatment, X^2 (3) = 57.265, p < 2e-12), with the highest levels in the liver and skin in the

DHQ+PTX feeding group (Figure 1C). We detected aPTX **267A** in the skin of all individuals in

the DHQ+PTX feeding group at higher levels than the DHQ-fed group (Wilcoxon test, W = 25, p

= 0.012, Figure 1D). The amount of aPTX **267A** differed across tissues (Kruskal-Wallis, X^2 (3) =

13.727, p = 0.003), with greater abundance in the skin than the eggs (post-hoc Dunn test, p =

0.001) and intestines (post-hoc Dunn test, p = 0.035, Figure 1E). These data show *D. tinctorius*

can metabolize PTX **251D** into aPTX **267A** and that some alkaloid metabolism may occur in the

liver and intestines.

Figure 1: Alkaloid sequestration in different tissue types. Boxplots showing abundance of
different compounds in different tissues and treatment types, with DHQ-fed frogs in yellow and
DHQ+PTX-fed frogs in blue. **(A)** Frogs were orally administered either DHQ or DHQ+PTX once
a day for five days. **(B)** DHQ abundance differed by tissue but not treatment group, and was

highest in the liver and skin (GLMM tissue, X^2 (3) = 203.642, p < 2e-16). **(C)** PTX levels differed
by tissue and treatment, and were higher in the liver and skin of the DHQ+PTX fed group
(GLMM tissue:treatment, X^2 (3) = 57.265, p < 2e-12). **(D)** The hydroxylated metabolite aPTX
was found in the DHQ+PTX fed frogs (Wilcoxon test, W = 0, p-value = 0.012, n = 5). **(E)** aPTX
abundance differed across tissues within the DHQ-PTX group (Kruskal-Wallis, X^2 (3) = 13.727, p
= 0.003), and was found primarily in the skin, with some in the liver.
263

3.2 PTX alters gene expression across tissues

To identify genes that may be involved in PTX sequestration and metabolism, we next

quantified gene expression changes across tissues using RNA sequencing. The number of

genes upregulated in response to PTX **251D** feeding were 282 in the intestines, 144 in the liver,

and 197 in the skin. Although hundreds of genes were differentially expressed in each tissue,

most did not have annotations or they aligned with unknown, hypothetical, or non-vertebrate

proteins (Supplementary Excel File). Cytochrome P450 (CYP3A29), an enzyme family well-

known for their involvement in small molecule hydroxylation, was upregulated in the intestines (t

= 4.7, log2FC = 5.72, p-adjusted = 0.0045; Figure 2A). In the liver, vitellogenin 2 (VTG2) was

downregulated in the PTX feeding group (t = 3.8, log2FC = -7.73, p-adjusted = 0.0421, Figure

2B). MHC Class I□ was upregulated in both the liver (t = 5.7, log2FC = 3.49, p-adjusted =

0.0005) and intestines (t = 5.7, log2FC = 5.41, p-adjusted = 0.0001) in the presence of PTX

251D (Figure 2A,B). In the skin, syntaxin 1A (STX1A) was upregulated (t = 4.0, log2FC = 2.58,

p-adjusted = 0.0385) and a solute carrier family 2 protein (SLC2) was downregulated (t = -3.7,

log2FC = 6.73, p-adjusted = 0.0496) in response to PTX **251D** (Figure 2C).

Figure 2: Differentially expressed genes in different tissues. Boxplots show TMM normalized expression levels in the DHQ-fed group (yellow) and DHQ+PTX fed group (blue) for a subset of differentially expressed genes. (A) Differentially expressed genes in the intestines include Cytochrome P450 Family 3 Protein 29 (CYP3A29) and MHC Class I alpha (MHCIα). **(B)** Differentially expressed genes in the liver include vitellogenin 2 (VTG2) and MHCIα. **(C)** Differentially expressed genes in the skin include syntaxin 1A (STX1A) and solute carrier family 2 (SLC2). (FC indicates log2 fold change values, * indicates adjusted p-value < 0.05, *** indicates adjusted p-value < 0.005; *y*-axes of individual plots have different scales).

3.3 Molecular targets of PTX 251D

A common reaction performed by cytochrome P450 (CYP) enzymes is the hydroxylation

of small molecule substrates [42]. To identify candidate cytochrome P450s that may metabolize

PTX **251D**, we screened seven human CYPs for PTX **251D** clearance. Human CYP2D6

showed rapid clearance of PTX **251D**, with most of the initial compound depleted after 15

minutes (Figure 3A), while CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, and CYP3A4

showed minimal clearance of PTX **251D** after 60 minutes. We followed this observation with a

metabolite discovery screen using human CYP2D6. PTX **251D** was completely cleared after 90

minutes, and two metabolites were identified with mass-to-charge ratio (m/z) shifts

corresponding to either one (m/z of M+H = 266) or two (m/z of M+H = 282) hydroxylation events

(Figure 3B). Together, this suggests that CYP2D6 may metabolize PTX **251D** into aPTX **267A**.

To better understand the physiological impacts of PTX **251D** exposure, we also

conducted a screen for activity on human G-protein coupled receptors (GPCRs) and ion-

channels. At a maximum concentration of 30 uM PTX **251D** showed no inhibitory effect on the

ion channels NaV1.5, Kv4.3/KChIP2, hERG, KCNQ1/minK, Cav1.2, or Kir2.1. At a

concentration of 10 uM PTX **251D** showed no significant inhibition or activation of any of the 168

GPCRs tested. These results suggest that at these concentrations PTX **251D** does not have an

effect on the specific human GPCRs or ion channels tested in these panels.

Figure 3: Human CYP2D6 PTX 251D metabolism. (A) Human CYP2D6 rapidly clears PTX
251D in-vitro compared to other human CYPs. **(B)** Human CYP2D6 creates two hydroxylation
products from PTX **251D** corresponding to a single hydroxylation event (m/z = 266), suggestive
of aPTX **267A**, and two hydroxylation events (m/z = 282). Mass-to-charge ratio of the peak is
indicated with "m/z", and "peak area" indicates the area under the curve of each peak
corresponding to the m/z indicated.

3.4 CYP2D6 expression and sequence diversity

Given that human CYP2D6 hydroxylates PTX **251D**, we searched for CYP2D6

homologs in the *D. tinctorius* transcriptome. Out of eight contigs with high sequence similarity to

human CYP2D6, one contig showed high sequence similarity (BLAST e-value = 6.89e-121) and

was upregulated in the intestines of frogs fed PTX **251D** (Welch's two-sample t-test, t = -4.0227,

df = 6.3801, p-value = 0.00611; Figure 4A). We next examined if there were sequence

differences in this homolog across amphibians that relate to PTX metabolism [19]. We identified

homologous genes to this upregulated candidate in other amphibian transcriptomes and

genomes and aligned them to the human CYP2D6 sequence to find conserved amino acid

residues. Of the human CYP2D6 main active site residues [39], Asp301 was fully conserved in

all amphibian protein sequences, while Phe120 and Phe483 were changed to aliphatic residues

isoleucine, valine, or leucine. Glu216 in *D. tinctorius* and most other amphibians was changed to

an alanine, however in *O. sylvatica* and *E. tricolor* was changed into a serine. Overall, we did

not identify a clear pattern in active site mutations that reflect the ability to convert PTX **251D**

into aPTX **267A**, although conversion has not been tested in all species for which sequences

are available.

Figure 4: CYP2D6 expression and sequence conservation. (A) A CYP2D6-like protein was
identified in the *D. tinctorius* transcriptome that was upregulated with PTX feeding. **(B)**
Alignment of CYP2D6-like proteins in poison frogs and other amphibians show that important
human CYP2D6 binding residue Asp301 is conserved, however other active site residues are
changed in frogs. Blank spots indicate residues not present in the alignment of those species
due to shorter protein sequences. The orange lines on the dendrobatid phylogeny indicate
independent origins of chemical defense. Species in black have been tested for ability to

metabolize PTX **251D** into aPTX **267A**, whereas species names in gray have not been tested.
Asterisks (*) indicate species that sequester aPTX **267A** when fed PTX **251D**. Amino acid
residues are colored using the RasMol scheme, which corresponds to amino acid properties.

4. DISCUSSION

Poison frogs acquire alkaloids from their diet for chemical defense, and in some cases

are able to metabolize these compounds into more potent forms. This study shows that when

fed PTX **251D,** *D. tinctorius* is able to create and sequester the more potent aPTX **267A**, and

that the consumption of PTX **251D** changes gene expression across multiple tissues.

Furthermore, human CYP2D6 is able to hydroxylate PTX **251D**, and a similar gene found in

poison frogs and other amphibians show sequence conservation in Asp301, an important

binding residue. These results expand our understanding of PTX, and how general alkaloid

metabolism may be co-opted in poison frog physiology for the creation of stronger chemical

defense from available dietary resources.

Performing a controlled feeding study with DHQ and PTX **251D** allowed us to determine

that *D. tinctorius* can metabolize PTX **251D** into aPTX **267A**. Although previous studies have

documented wild *D. tinctorius* with aPTX on their skin [4,21], this is the first experimental

evidence that this species metabolizes PTX **251D** into aPTX **267A** rather than sequestering

aPTX from their diet. Previous work has shown that *D. auratus*, a species closely related to *D.*

tinctorius, is able to convert PTX **251D** in aPTX **267A** [19]. However aPTX **267A** is not detected

on the skin of *P. bicolor*, a species in the same origin of toxicity, and *E. tricolor,* a species in a

different origin, when fed PTX **251D** in the laboratory [19]. It is possible that the ability to convert

PTX into aPTX or sequester aPTX to the skin has only been acquired in specific poison frog

populations or species. The accumulation of both DHQ and PTX **251D** in the liver, intestines,

and skin, indicates that these tissues play an important role in the sequestration of alkaloids and

echoes previous controlled feeding study results where frogs were only fed DHQ [14]. The liver

and intestines are also important sites of alkaloid metabolism in mammals due to high levels of

Cytochrome P450s [43–45]. Together, these results show that *D. tinctorius* is able to metabolize

PTX **251D** into aPTX **267A** and that the tissue distribution of alkaloids includes the skin, liver,

and intestines.

PTX **251D** feeding resulted in gene expression changes in the intestines, liver, and skin,

suggesting a single alkaloid can influence poison frog physiology. Specifically, the upregulation

of CYP3A29 in response to PTX in the intestines implicates this enzyme in the metabolism of

PTX **251D** into aPTX **267A**, or of PTX into a metabolic byproduct to be later discarded. Although

we originally expected to identify metabolism enzymes in the liver, it is possible the liver instead

acts primarily as a detoxification site. In the dendrobatid *Oophaga sylvatica*, feeding DHQ

compared to a non-alkaloid vehicle control led to a downregulation of CYP3A29 in the intestines

[14]. This suggests that expression of CYP3A29 may be downregulated in response alkaloids

generally, yet upregulated in response to specific alkaloids, such as the addition of PTX **251D**.

The upregulation of MHC class I□ proteins in the intestines and liver in response to PTX **251D**

supports previous findings that frog immune systems respond to alkaloids [31,46]. We also

found that VTG2 (vitellogenin-2) was downregulated in response to PTX **251D**. Although

vitellogenins are typically thought to be egg-yolk proteins, they also play regulatory roles and

protect cells from reactive oxygen species that may arise from alkaloid metabolism [47–49].

Finally, SLC2 (solute carrier family 2) which encodes for the GLUT family of glucose

transporters, was downregulated in the skin with PTX feeding. Alkaloids can be potent inhibitors

of GLUTs in mammalian cell lines [50], and the downregulation of GLUTs in this case may be

due to the presence of concentrated PTX **251D** in the frog skin. Together, these gene

expression changes that are in response to PTX **251D** compared to an alkaloid-fed control

support an argument for physiological "fine-tuning" of gene expression in response to certain

alkaloids.

Human CYP2D6 is able to hydroxylate PTX **251D**, and similar sequences in poison frogs

and other amphibians show varying sequence conservation in important binding residues. This

study provides the first evidence that human CYP2D6 is able to rapidly clear PTX **251D**, and a

gene in *D. tinctorius* with high similarity (BLAST e-value = 6.89e-121) showed an increased

expression in the intestines of frogs fed PTX **251D**. Human CYP2D6 is documented to be

involved in the metabolism of many compounds, including plant alkaloids [51]. Although there is

no direct evidence for the upregulation of CYP2D6 in response to exogenous compounds in

mammals, increased abundance of members of the CYP2D family through gene duplications in

humans and mice are hypothesized to allow species to deal with the presence of dietary or

environmental alkaloids [52]. Furthermore, human CYP2D6 is known to perform hydroxylation

reactions mediated through the Asp301 residue binding to nitrogen in lipophilic substrate

compounds [39]. Crystal structure and site-directed mutagenesis studies have found that

additional residues involved in substrate binding in the active site of human CYP2D6 are

Phe120, Glu216, and Phe483 [39]. Our results find that a hydroxylation product of PTX **251D** is

made by human CYP2D6, however it is not possible with our data to know if the position of

hydroxylation corresponds exactly to aPTX **267A.** Nevertheless, given that Asp301 is conserved

across distant frog species, many species may have the ability to create aPTX **267A,** and only

certain poison frog species have evolved the ability to sequester aPTX **267A** to their skin for

chemical defense. *D. auratus* and *D. tinctorius* are the only tested species known to sequester

aPTX **267A** when fed PTX **251D,** and also the only species found with Ile120 instead of Val120

in their CYP2D6-like proteins. Although isoleucine and valine are similar amino acids this

change may still influence the substrate binding affinity of CYP2D6 or positioning of substrate

hydroxylation. It is also likely that there are additional residues that coordinate specificity for

PTX **251D** that are yet undiscovered. Controlled feeding experiments with PTX **251D** in other

species and detailed *in vitro* biochemistry and crystallography studies would need to confirm

how binding specificity may differ between species that can and cannot make aPTX **267A**.

Together, these data suggest that many species of amphibians may be able to hydroxylate PTX

251D, and that *D. tinctorius* may be modulating the expression of a CYP2D6-like protein to

metabolize PTX **251D** into aPTX **267A** for improved chemical defense**.**

We did not find any effect of PTX **251D** on the six ion channels and 168 GPCRs tested.

For the ion channels, this is probably because the concentrations tested were much lower than

that of previous electrophysiology work on PTX **251D** [30]. Patch clamp experiments with

mammalian and insect voltage-gated sodium and potassium channels found that at 100 µM

PTX **251D** inhibits both types of channels, with the strongest effect being on hKv1.3 [30]. We

tested a range of concentrations of PTX **251D** in the ion channel screen based on the active

concentrations tested in electrophysiology studies of batrachotoxin (BTX) [53], and the lack of

effect seen by PTX **251D** at these concentrations supports findings that PTX **251D** is not as

potent of a toxin as BTX for mammalian ion channels [19,54]. In order to better understand the

effects of different poison frog toxins, including aPTX **267A**, further systematic testing of

potential targets would be required, although synthesis of the compounds found on wild poison

frogs is difficult and has limited research progress in this direction.

In summary, this study provides evidence that *D. tinctorius* can metabolize PTX **251D**

into aPTX **267A** and that PTX **251D** exposure changes gene expression across tissues,

demonstrating that specific alkaloids can change poison frog physiology [14,46]. Following up

on candidate genes with biochemical studies is needed in order to fully characterize the

genetics of alkaloid sequestration and metabolism. In the wild, where chemically defended

dendrobatids carry many different alkaloids, subtle alkaloid differences may induce distinct gene

expression changes. More broadly, modulating gene expression in response to specific

alkaloids may set the stage for local adaptation to environmental resources.

5. ACKNOWLEDGEMENTS

We thank Stephanie Caty and Nora Moskowitz for their comments on early versions of this

manuscript. We also thank the O'Connell Lab for frog colony care and general experimental

advice and encouragement. We acknowledge that this research was conducted at Stanford

University, which is located on the ancestral and unceded land of the Muwekma Ohlone tribe.

6. FUNDING

This work was supported by a National Science Foundation (NSF) [IOS-1822025] grant

to LAO. AAB is supported by a NSF Graduate Research Fellowship (DGE-1656518) and an

HHMI Gilliam Fellowship. LAO is a New York Stem Cell – Robertson Investigator.

6. DATA ACCESSIBILITY

All LC-MS/MS data from the alkaloid analysis, the *D. tinctorius* transcriptome, TMM expression

and count data for different tissues, fasta files with nucleotide and amino acid sequences of

CYP2D6-like proteins, and code is available from the Dryad Digital Repository (pending). All

Illumina fastq files are available on the Sequence Read Archive (pending).

7. AUTHOR CONTRIBUTIONS

AAB and LAO designed the experiment. AAB and CYP carried out the experimental procedures.

CV and SAT quantified alkaloids. AAB analyzed the data. AAB and LAO wrote the manuscript

with contributions from all authors.

8. REFERENCES

1. Naumann C, Hartmann T, Ober D. Evolutionary recruitment of a flavin-dependent monooxygenase for the detoxification of host plant-acquired pyrrolizidine alkaloids in the alkaloid-defended arctiid moth Tyria jacobaeae. Proc Natl Acad Sci U S A. 2002;99: 6085–6090.

2. Mizutani M. Impacts of diversification of cytochrome P450 on plant metabolism. Biol Pharm Bull. 2012;35: 824–832.

3. Daly JW, Brown GB, Mensah-Dwumah M, Myers CW. Classification of skin alkaloids from neotropical poison-dart frogs (dendrobatidae). Toxicon. 1978;16: 163–188.

4. Daly JW, Myers CW, Whittaker N. Further classification of skin alkaloids from neotropical poison frogs (dendrobatidae), with a general survey of toxic/noxious substances in the amphibia. Toxicon. 1987;25: 1023–1095.

5. Daly JW, Spande TF, Garraffo HM. Alkaloids from Amphibian Skin: A Tabulation of Over Eight-Hundred Compounds. J Nat Prod. 2005;68: 1556–1575.

6. Darst CR, Menéndez-Guerrero PA, Coloma LA, Cannatella DC. Evolution of dietary specialization and chemical defense in poison frogs (Dendrobatidae): a comparative analysis. Am Nat. 2004;165: 56–69.

7. Daly JW, Garraffo HM, Spande TF, Jaramillo C, Rand AS. Dietary source for skin alkaloids of poison frogs (Dendrobatidae)? J Chem Ecol. 1994;20: 943–955.

8. Vences M. Convergent evolution of aposematic coloration in Neotropical poison frogs: a molecular phylogenetic perspective. Organisms Diversity & Evolution. 2003. pp. 215–226. doi:10.1078/1439-6092-00076

9. Santos JC, Coloma LA, Cannatella DC. Multiple, recurring origins of aposematism and diet specialization in poison frogs. Proceedings of the National Academy of Sciences. 2003;100: 12792–12797.

10. Carvajal-Castro JD, Vargas-Salinas F, Casas-Cardona S, Rojas B, Santos JC. Aposematism facilitates the diversification of parental care strategies in poison frogs. Sci Rep. 2021;11: 19047.

11. Saporito RA, Russell MW, Richards-Zawacki CL, Dugas MB. Experimental evidence for maternal provisioning of alkaloid defenses in a dendrobatid frog. Toxicon. 2019. doi:10.1016/j.toxicon.2019.02.008 PMID - 30790578

12. Fischer EK, Roland AB, Moskowitz NA, Vidoudez C, Ranaivorazo N, Tapia EE, et al. Mechanisms of Convergent Egg Provisioning in Poison Frogs. Curr Biol. 2019;29: 4145–4151.e3.

13. Stynoski JL, Torres-Mendoza Y, Sasa-Marin M, Saporito RA. Evidence of maternal provisioning of alkaloid-based chemical defenses in the strawberry poison frogOophaga pumilio. Ecology. 2014. pp. 587–593. doi:10.1890/13-0927.1

14. O'Connell LA, Course, LS50: Integrated Science Laboratory, O'Connell JD, Paulo JA,

Trauger SA, Gygi SP, et al. Rapid toxin sequestration impacts poison frog physiology.
bioRxiv. 2020; 2020.05.27.119081.

15. O'Connell LA, LS50: Integrated Science Laboratory Course, O'Connell JD, Paulo JA,
Trauger SA, Gygi SP, et al. Rapid toxin sequestration modifies poison frog physiology. J
Exp Biol. 2021;224. doi:10.1242/jeb.230342

16. Caty SN, Alvarez-Buylla A, Byrd GD, Vidoudez C, Roland AB, Tapia EE, et al. Molecular
physiology of chemical defenses in a poison frog. J Exp Biol. 2019;222.
doi:10.1242/jeb.204149

17. Sanchez E, Rodríguez A, Grau JH, Lötters S, Künzel S, Saporito RA, et al. Transcriptomic
Signatures of Experimental Alkaloid Consumption in a Poison Frog. Genes . 2019;10.
doi:10.3390/genes10100733

18. Daly JW, Secunda SI, Garraffo HM, Spande TF, Wisnieski A, Cover JF. An uptake system
for dietary alkaloids in poison frogs (Dendrobatidae). Toxicon. 1994;32: 657–663.

19. Daly JW, Garraffo HM, Spande TF, Clark VC, Ma J, Ziffer H, et al. Evidence for an
enantioselective pumiliotoxin 7-hydroxylase in dendrobatid poison frogs of the genus
Dendrobates. Proceedings of the National Academy of Sciences. 2003;100: 11092–11097.

20. Saporito RA, Spande TF, Martin Garraffo H, Donnelly MA. Arthropod Alkaloids in Poison
Frogs: A Review of the "Dietary Hypothesis." HETEROCYCLES. 2009. p. 277.
doi:10.3987/rev-08-sr(d)11

21. Saporito RA, Garraffo HM, Donnelly MA, Edwards AL, Longino JT, Daly JW. Formicine
ants: An arthropod source for the pumiliotoxin alkaloids of dendrobatid poison frogs. Proc
Natl Acad Sci U S A. 2004;101: 8045–8050.

22. Saporito RA, Donnelly MA, Norton RA, Garraffo HM, Spande TF, Daly JW. Oribatid mites
as a major dietary source for alkaloids in poison frogs. Proceedings of the National
Academy of Sciences. 2007;104: 8885–8890.

23. Moskowitz NA, Dorritie B, Fay T, Nieves OC, Vidoudez C, Class CRAL 2017 B, et al. Land
use impacts poison frog chemical defenses through changes in leaf litter ant communities.
Neotropical Biodiversity. 2020;6: lxxv–lxxxvii.

24. McGugan JR, Byrd GD, Roland AB, Caty SN, Kabir N, Tapia EE, et al. Ant and Mite
Diversity Drives Toxin Variation in the Little Devil Poison Frog. J Chem Ecol. 2016;42: 537–
551.

25. Daly JW, Garraffo HM, Hall GS, Cover JF Jr. Absence of skin alkaloids in captive-raised
Madagascan mantelline frogs (Mantella) and sequestration of dietary alkaloids. Toxicon.
1997;35: 1131–1135.

26. Bargar TM, Lett RM, Johnson PL, Hunter JE, Chang CP, Pernich DJ, et al. Toxicity of
Pumiliotoxin 251D and Synthetic Analogs to the Cotton Pest Heliothis virescens. Journal of
Agricultural and Food Chemistry. 1995. pp. 1044–1051. doi:10.1021/jf00052a037

27. Weldon PJ, Kramer M, Gordon S, Spande TF, Daly JW. A common pumiliotoxin from
poison frogs exhibits enantioselective toxicity against mosquitoes. Proc Natl Acad Sci U S

A. 2006;103: 17818–17821.

28. Daly JW, McNeal ET, Overman LE, Ellison DH. A new class of cardiotonic agents:
structure-activity correlations for natural and synthetic analogues of the alkaloid A new
class of A new class of cardiotonic agents: structure-activity correlations for natural and
synthetic analogues of the alkaloid pumiliotoxin B (8-hydroxy-8-methyl-6-alkylidene-1-
azabicyclo[4.3.0]nonanes). J Med Chem. 1985;28: 482–486.

29. Daly JW, McNeal E, Gusovsky F, Ito F, Overman LE. Pumiliotoxin alkaloids: relationship of
cardiotonic activity to sodium channel activity and phosphatidylinositol turnover. J Med
Chem. 1988;31: 477–480.

30. Vandendriessche T, Abdel-Mottaleb Y, Maertens C, Cuypers E, Sudau A, Nubbemeyer U,
et al. Modulation of voltage-gated Na+ and K+ channels by pumiliotoxin 251D: A "joint
venture" alkaloid from arthropods and amphibians. Toxicon. 2008;51: 334–344.

31. Caty SN, Alvarez-Buylla A, Byrd GD, Vidoudez C, Roland AB, Tapia EE, et al. Molecular
physiology of chemical defenses in a poison frog. J Exp Biol. 2019;222: jeb204149.

32. Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, et al. De novo
transcript sequence reconstruction from RNA-seq using the Trinity platform for reference
generation and analysis. Nat Protoc. 2013;8: 1494–1512.

33. UniProt Consortium. UniProt: a worldwide hub of protein knowledge. Nucleic Acids Res.
2019;47: D506–D515.

34. Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of protein or
nucleotide sequences. Bioinformatics. 2006. pp. 1658–1659.
doi:10.1093/bioinformatics/btl158

35. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-generation
sequencing data. Bioinformatics. 2012;28: 3150–3152.

36. Bray NL, Pimentel H, Melsted P, Pachter L. Erratum: Near-optimal probabilistic RNA-seq
quantification. Nat Biotechnol. 2016;34: 888.

37. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biol. 2014;15: 550.

38. Brooks M, Mollie, Brooks E, Kristensen K, Koen, J., et al. glmmTMB Balances Speed and
Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. The R
Journal. 2017. p. 378. doi:10.32614/rj-2017-066

39. Rowland P, Blaney FE, Smyth MG, Jones JJ, Leydon VR, Oxbrow AK, et al. Crystal
structure of human cytochrome P450 2D6. J Biol Chem. 2006;281: 7614–7622.

40. Stuckert AMM, Moore E, Coyle KP, Davison I, MacManes MD, Roberts R, et al. Variation in
pigmentation gene expression is associated with distinct aposematic color morphs in the
poison frog Dendrobates auratus. BMC Evol Biol. 2019;19: 85.

41. Márquez R, Linderoth TP, Mejía-Vargas D, Nielsen R, Amézquita A, Kronforst MR.
Divergence, gene flow, and the origin of leapfrog geographic distributions: The history of
colour pattern variation in Phyllobates poison-dart frogs. Mol Ecol. 2020;29: 3702–3719.

42. Ortiz de Montellano PR. Hydrocarbon hydroxylation by cytochrome P450 enzymes. Chem
Rev. 2010;110: 932–948.

43. Grant DM. Detoxification pathways in the liver. J Inherit Metab Dis. 1991;14: 421–430.

44. Robinson T. The Metabolism and Biochemical Actions of Alkaloids in Animals. Bioactive
Natural Products (Part C). 2000. pp. 3–54. doi:10.1016/s1572-5995(00)80022-8

45. Thelen K, Dressman JB. Cytochrome P450-mediated metabolism in the human gut wall. J
Pharm Pharmacol. 2009;61: 541–558.

46. Sanchez E, Rodríguez A, Grau JH, Lötters S, Künzel S, Saporito RA, et al. Transcriptomic
Signatures of Experimental Alkaloid Consumption in a Poison Frog. Genes . 2019;10: 733.

47. Havukainen H, Münch D, Baumann A, Zhong S, Halskau Ø, Krogsgaard M, et al.
Vitellogenin recognizes cell damage through membrane binding and shields living cells
from reactive oxygen species. J Biol Chem. 2013;288: 28369–28381.

48. Nunes FMF, Ihle KE, Mutti NS, Simões ZLP, Amdam GV. The gene vitellogenin affects
microRNA regulation in honey bee (Apis mellifera) fat body and brain. J Exp Biol. 2013;216:
3724–3732.

49. Seehuus S-C, Norberg K, Gimsa U, Krekling T, Amdam GV. Reproductive protein protects
functionally sterile honey bee workers from oxidative stress. Proc Natl Acad Sci U S A.
2006;103: 962–967.

50. Reckzeh ES, Waldmann H. Development of Glucose Transporter (GLUT) Inhibitors.
European J Org Chem. 2020;2020: 2321–2329.

51. Darney K, Lautz LS, Béchaux C, Wiecek W, Testai E, Amzal B, et al. Human variability in
polymorphic CYP2D6 metabolism: Implications for the risk assessment of chemicals in food
and emerging designer drugs. Environ Int. 2021;156: 106760.

52. Ingelman-Sundberg M. Genetic polymorphisms of cytochrome P450 2D6 (CYP2D6):
clinical consequences, evolutionary aspects and functional diversity. The
Pharmacogenomics Journal. 2005. pp. 6–13. doi:10.1038/sj.tpj.6500285

53. Finol-Urdaneta RK, McArthur JR, Goldschen-Ohm MP, Gaudet R, Tikhonov DB, Zhorov
BS, et al. Batrachotoxin acts as a stent to hold open homotetrameric prokaryotic voltage-
gated sodium channels. J Gen Physiol. 2019;151: 186–199.

54. Albuquerque EX, Daly JW, Witkop B. Batrachotoxin: Chemistry and Pharmacology.
Science. 1971. pp. 995–1002. doi:10.1126/science.172.3987.995

61. Ogle DH, Doll JC, Wheeler P, Dinno A (2021). FSA: Fisheries Stock Analysis. R package
version 0.9.1, https://github.com/droglenc/FSA.

Figure 4: CYP2D6 expression and sequence conservation. (A) A CYP2D6-like protein was identified in the *D. tinctorius* transcriptome that was upregulated with PTX feeding. **(B)** Alignment of CYP2D6-like proteins in poison frogs and other amphibians show that important human CYP2D6 binding residue Asp301 is conserved, however other active site residues are changed in frogs. Blank spots indicate residues not present in the alignment of those species due to shorter protein sequences. The orange lines on the dendrobatid phylogeny indicate independent origins of chemical defense. Species in black have been tested for ability to metabolize PTX **251D** into aPTX **267A**, whereas species names in gray have not been tested. Asterisks (*) indicate species that sequester aPTX **267A** when fed PTX **251D**. Amino acid residues are colored using the RasMol scheme, which corresponds to amino acid properties.

Figure 3: Human CYP2D6 PTX 251D metabolism. (A) Human CYP2D6 rapidly clears PTX **251D** in-vitro compared to other human CYPs. **(B)** Human CYP2D6 creates two hydroxylation products from PTX **251D** corresponding to a single hydroxylation event (m/z = 266), suggestive of aPTX **267A**, and two hydroxylation events (m/z = 282). Mass-to-charge ratio of the peak is indicated with "m/z", and "peak area" indicates the area under the curve of each peak corresponding to the m/z indicated.

Figure 2: Differentially expressed genes in different tissues. Boxplots show TMM normalized expression levels in the DHQ-fed group (yellow) and DHQ+PTX fed group (blue) for a subset of differentially expressed genes. (A) Differentially expressed genes in the intestines include Cytochrome P450 Family 3 Protein 29 (CYP3A29) and MHC Class I alpha (MHCI□). **(B)** Differentially expressed genes in the liver include vitellogenin 2 (VTG2) and MHCI□. **(C)** Differentially expressed genes in the skin include syntaxin 1A (STX1A) and solute carrier family 2 (SLC2). (FC indicates log2 fold change values, * indicates adjusted p-value < 0.05, *** indicates adjusted p-value < 0.005; *y*-axes of individual plots have different scales).

Figure 1: Alkaloid sequestration in different tissue types. Boxplots showing abundance of different compounds in different tissues and treatment types, with DHQ-fed frogs in yellow and DHQ+PTX-fed frogs in blue. **(A)** Frogs were orally administered either DHQ or DHQ+PTX once a day for five days. **(B)** DHQ abundance differed by tissue but not treatment group, and was highest in the liver and skin (GLMM tissue, X^2 (3) = 203.642, p < 2e-16). **(C)** PTX levels differed by tissue and treatment, and were higher in the liver and skin of the DHQ+PTX fed group (GLMM tissue:treatment, X^2 (3) = 57.265, p < 2e-12). **(D)** The hydroxylated metabolite aPTX was found in the DHQ+PTX fed frogs (Wilcoxon test, W = 0, p-value = 0.012, n = 5). **(E)** aPTX abundance differed across tissues within the DHQ-PTX group (Kruskal-Wallis, X^2 (3) = 13.727, p = 0.003), and was found primarily in the skin, with some in the liver.