Reciprocal lateral hypothalamic and raphé GABAergic projections promote

wakefulness

Mary Gazea[1,2*a], Szabina Furdan[3*], Péter Sere[3,4], Lukas Oesch[2,3], Benedek Molnár[3,4],

Giuseppe Di Giovanni[5,6], Lief E. Fenno[7], Charu Ramakrishnan[8], Joanna Mattis[7], Karl

Deisseroth[7,9], Susan M. Dymecki[10], Antoine R. Adamantidis[2,3¶] and Magor L. Lőrincz[3,4,5¶]

[1]Centre for Experimental Neurology, Department of Neurology, Inselspital University

Hospital Bern, University of Bern, Bern, Switzerland

[2]Department of Biomedical Research (DBMR), Inselspital University Hospital Bern,

University of Bern, Bern, Switzerland

[3]Department of Physiology, Anatomy, and Neuroscience, University of Szeged, Szeged,

Hungary

[4]Department of Physiology, University of Szeged, Szeged, Hungary.

[5]Neuroscience Division, School of Bioscience, Cardiff University, Cardiff, UK

[6]Department of Physiology and Biochemistry, University of Malta, Malta

[7] Departments of Psychiatry & Behavioral Sciences and Bioengineering, Stanford University,

Stanford, USA

[8] CNC Program, Stanford University, Stanford, USA

[9] Howard Hughes Medical Institute, Stanford University, Stanford, USA

[10]Department of Genetics, Harvard Medical School, Boston, MA, USA.

[a] Present address: Concentris research management GmbH, Fürstenfeldbruck, Germany

*¶equal contribution

Correspondence should be addressed to **Magor L. Lőrincz**: mlorincz@gmail.com or **Antoine**

R. Adamantidis: antoine.adamantidis@dbmr.unibe.ch

Keywords: arousal, brain states, neuromodulation, inhibition, GABA, optogenetics,

disinhibition

33   ABSTRACT

The lateral hypothalamus (LH), together with multiple neuromodulatory systems of the brain,

such as the dorsal raphé nucleus (DR), is implicated in arousal, yet interactions between these

systems are just beginning to be explored. Using a combination of viral tracing, circuit

mapping, electrophysiological recordings from identified neurons and combinatorial

optogenetics in mice, we show that GABAergic neurons in the LH selectively inhibit

GABAergic neurons in the DR resulting in increased firing of a substantial fraction of its

neurons that ultimately promotes arousal. These DR_{GABA} neurons are wake active and project

to multiple brain areas involved in the control of arousal including the LH, where their

specific activation potently influences local network activity leading to arousal from sleep.

Our results show how mutual inhibitory projections between the LH and the DR promote

wakefulness and suggest a complex arousal control by intimate interactions between long-

range connections and local circuit dynamics.

Introduction

Brain-wide neuronal activities are strongly modulated across sleep-wake states (Saper et al., 2010; Weber and Dan, 2016) attributable to the influence of neuromodulatory systems of the brainstem (Jones, 2020), basal forebrain (Fuller et al., 2011; Xu et al., 2015) and lateral hypothalamus (LH) (Bonnavion et al., 2016; Arrigoni et al., 2019). In this context, the LH stands both as an anatomical and a sleep-wake hub.

Various LH neuronal populations are known to modulate sleep-wake states (Saper et al., 2010; Weber and Dan, 2016), energy intake and reward (Bernardis and Bellinger, 1996; Stuber and Wise, 2016). Hypocretins/orexins (Hcrt/Ox) and Vgat-expressing neurons of the LH (LH$_{Vgat}$) control wakefulness (Adamantidis et al., 2007; Carter et al., 2009; Carter et al., 2012; Herrera et al., 2016; Venner et al., 2016). Optogenetic activation of LH$_{Vgat}$ neurons induced rapid arousal through direct modulation of the reticular thalamus and brainstem circuits in the locus coeruleus area (Herrera et al., 2016), concommittent to an inhibition of anterior hypothalamic sleep-promoting circuits (Venner et al., 2019). In contrast, activation of MCH neurons promotes REM, and to a lesser extent NREM, sleep (Verret et al., 2003; Jego et al., 2013; Konadhode et al., 2013; Tsunematsu et al., 2014). Yet, whether and how LH$_{Vgat}$ neurons also modulate the activity of other brainstem neuromodulatory circuits remains unclear.

LH$_{Vgat}$ neurons are reciprocally connected to multiple brainstem nuclei (Bonnavion et al., 2016), including the DR (Weissbourd et al., 2014). The DR is an appealing candidate for influencing sleep-wake states since its major neuromodulator serotonin rapidly influences sensory (Kapoor et al., 2016; Lottem et al., 2016), motor (Liu et al., 2014; Correia et al., 2017; Seo et al., 2019) and cognitive (Miyazaki et al., 2011; Liu et al., 2014; Cohen et al., 2015; Fonseca et al., 2015) functions. Importantly, the DR has been involved in the regulation of various patho-physiological functions including depression, which is often accompanied by

sleep disturbances (Lowry et al., 2008; Steiger and Pawlowski, 2019; Riemann et al., 2020).

The largest fraction of DR neurons are non-serotonergic (Descarries et al., 1982), express the

VGAT marker (Allers and Sharp, 2003) and project to various forebrain targets including the

LH (Bang and Commons, 2012). These DR_{VGAT} neurons control food intake and their activity

increases in response to fasting (Nectow et al., 2017), however, their role in sleep-wake

behavior remains to be examined.

Using combinatorial optogenetics and electrophysiological recordings from identified

neurons, we show that LH_{GABA} neurons selectively inhibit DR_{GABA} neurons resulting in

disinhibition of a part of DR neurons ultimately leading to arousal. We found that DR_{GABA}

neurons are highly active during wakefulness compared to NREM and REM sleep. Selective

activation of their projection to the LH area leads to wakefulness, highlighting the importance

of reciprocal hypothalamus-brainstem long range circuits in the control of brain states and

arousal.

MATERIALS AND METHODS

All experimental procedures were performed according to the European Communities

Council Directives of 1986 (86/609/EEC) and 2003 (2003/65/CE) for animal research and

were approved by the Ethics Committee of the University of Szeged and University of Bern.

80 adult (age > 2 months) male mice were used in this study.

Experimental design

Viral targeting

For *in vitro* recordings AAV1-CAG-ChR2-venus-WPRE-SV40, AAV1-CAGS-flex-ChR2-

tdTomato-WPRE-SV40 viruses, (50-150 nl, Penn Vector Core; titer: $5x10^{12}$ gc/ml) were

injected bilaterally in the LH (AP: -1.4 mm, ML: ±1 mm, DV: -5.0-5.4 mm) of WT, GAD-

GFP (Tamamaki et al., 2003) or Vgat-IRES-Cre (Vong et al., 2011) mice using a nanoinjector

(Nanoliter 2000, WPI) connected to a glass pipette (~20 um diameter) at a rate of 1 nl/s.

For *in vivo* optogenetic activation of LH_{VGAT} projections to the DR, one group of Vgat-IRES-

Cre mice (Vong et al., 2011) received AAV1-CAGGS-FLEX-CHR2-td tomato-SV40 or

AAV2-EF1a-DIO-hChR2(H134R)-eYFP, while the second group received AAV2-EF1a-DIO-

eYFP (control, all virus vectors were packaged at the Vector Core of the University of North

Carolina at Chapel Hill, titres between $3 \cdot 10^{12}$ to $5 \cdot 10^{12}$ gc/ml). For the intersectional

optogenetic activation of DR neuronal subpopulations, Vgat-IRES-Cre (Vong et al., 2011) x

Pet1-Flpe (Jensen et al., 2008) double transgenic mice received AAVdj-EF1a-Foff/Con-

hChR2(H134R)-eYFP (Pet1-OFF/Vgat-ON), AAVdj-EF1a-Fon/Con-hChR2(H134R)-eYFP

(Pet1-ON/Vgat-ON), or AAVdj-EF1a-eYFP (eYFP controls). 150-600 nl of each virus was

injected bilaterally into the LH or into the dorsal raphe nucleus (AP: -4.45, ML: +1.2, DV: -

2.9 to -3.1, at a 30° angle). The virus suspension was infused with a micro-infusion pump

(PHD Ultra, Harvard Apparatus) through a 28 G stainless steel cannula (Plastic One) at a rate

of 0.1 µl per minute.

Surgical procedures

Mice were anesthetized using 1-1.5% (v/v) Isoflurane, placed in a stereotaxic frame (Model

900, David Kopf Instruments, USA) on a heating pad (Supertech, Hungary). For sleep studies

8-10 week old male Vgat-IRES-Cre mice were chronically implanted with an optical fiber

(200 um, 0.39 NA Core Multimode Optical Fiber, FT200EMT, TECS Clad, Thorlabs,

inserted into zirconia ferrules, 1,25 mm OD; 230 µm ID, Precision Fiber product) above the

DR (AP -4.45, ML +1.2, DV -2.9 from Bregma at a 20° mediolateral angle), which were

fixated to the skull with C&B-Metabond (Parkell Inc.). Four stainless steel screw electrodes

were inserted epidurally (two into both frontal cortices, two into parietal cortices) for the

recording of EEG signals and wire loops were sewed to both neck muscles for EMG

recordings. The EEG/EMG electrodes and optic fiber were secured to the skull with dental

cement (Patterson dental). After surgical procedures mice received appropriate post-operative

care (analgesics: Rimadyl (15 mg/kg), anti-inflammatory drug: Dexafort (2mg/kg), antibiotic:

Gentamicin (5mg/kg) and were allowed to recover for two weeks.

In vitro electrophysiology

Mice were deeply anesthetized and perfused through the heart with ice cold cutting solution

containing (in mM) 93 *N*-methyl-d-glucamine, 2.5 KCl, 1.2 NaH_2PO_4, 30 $NaHCO_3$, 20

HEPES, 25 glucose, 5 *N*-acetyl-l-cysteine, 5 Na-ascorbate, 3 Na-pyruvate, 10 $MgSO_4$, and 0.5

$CaCl_2$. The same solution was used to cut coronal brainstem slices containing the DR at 4ºC

and for the initial storage of slices (32-34ºC for 12 minutes) following which the slices were

stored in a solution containing (in mM) 30 NaCl, 2.5 KCl, 1.2 NaH_2PO_4, 1.3 $NaHCO_3$, 20

HEPES, 25 glucose, 5 *N*-acetyl-l-cysteine, 5 Na-ascorbate, 3 Na-pyruvate, 3 $CaCl_2$, and 1.5

$MgSO_4$. For recording, slices were submerged in a chamber perfused with a warmed (34°C)

continuously oxygenated (95% O_2, 5% CO_2) ACSF containing (in mM) 130 NaCl, 3.5 KCl, 1

KH_2PO_4, 24 $NaHCO_3$, 1.5 $MgSO_4$, 3 $CaCl_2$, and 10 glucose. DR neurons were visualized

using standard DIC optics and recorded in whole cell current clamp mode using an EPC9

amplifier (Heka Elektronik). Patch pipettes (tip resistance, 4–5 MΩ) were filled with an

internal solution containing the following (in mM): 126 K-gluconate, 4 KCl, 4 ATP-Mg, 0.3

$GTP-Na_2$, 10 HEPES, 10 kreatin-phosphate, and 8 biocytin, pH 7.25; osmolarity, 275 mOsm.

The liquid junction potential (−13 mV) was corrected off-line. Access and series resistances

were constantly monitored, and data from neurons with a >20% change from the initial value

were discarded. DR_{GABA} neuron somata in GAD-GFP mice (Tamamaki et al., 2003) were

targeted under epifluorescent illumination. Photostimulation was performed through the

microscope objective using a blue LED light source (0.5-0.8 mW/mm^2, Thorlabs).

In vivo electrophysiology and juxtacellular labeling

To record the activity of DR neurons mice were anesthetized using 1-1.5% (v/v) Isoflurane

(Forane), placed in a stereotaxic frame (Model 900, David Kopf Instruments, USA) on a

heating pad (Supertech, Hungary). A small craniotomy (1 mm diameter) was made over the

DR (AP: -4.75, ML: 0.2, DV: -3.1 (mm) from Bregma) leaving the dura mater intact. Glass

electrodes (resistance: 10-25 MOhm) were filled with saline containing 1.5% Biocytin

(Sigma, Hungary). The electrode was lowered in a 15º mediolateral angle to avoid the sagittal

sinus. The activity of DR neurons was monitored using a Multiclamp 700B amplifier

(Molecular Devices) operating in current clamp mode (filtered between 100 Hz and 3 kHz).

At the end of some recordings neurons were filled with Biocytin (Sigma Aldrich) using 0.5-4

nA anodal current pulses of 500 ms duration, 50% duty cycle for 2-10 minutes) as previously

described (Pinault, 1996). For recording GABAergic neurons in the DR, the lateral wings

were targeted, where most neurons are GABAergic (Allers and Sharp, 2003). These neurons

possessed small (≤15 µm diameter) oval somata and mean baseline firing rates above 5 Hz

(Allers and Sharp, 2003).

Data acquisition.

Before the start of the experiment, the animals were habituated to a 3-m-long fiber optic patch

cord with protective tubing (Thorlabs) that was connected to the chronically implanted optical

fiber with a zirconia sleeve (Precision Fiber Products) and an EEG/EMG cable for one week.

Subsequently, a 24-h baseline polysomnographic recording was obtained without any

optogenetic manipulations. EEG and EMG signals were sampled with an AM systems 3500

amplifier at 512 Hz and digitized with a national instruments USB X DAQ device. EEG and

EMG data were recorded with the SleepScore software (View Point). All experiments took

place between 10 - 2 PM (Zeitgeber time 3-7).

For unit recordings in freely moving animals, mice were habituated to optical patch cords and

an electrode board head stage (RHD2132, Intan Technologies) for at least 4 days.

Subsequently a 24-h baseline polysomnographic recording was obtained without optogenetic

manipulations to ensure that the animals recovered a normal sleep-wake cycle. Tetrode

signals were acquired at 20 kHz with an open-source acquisition software (Open Ephys) via a

digitizing head stage (RHD2132) and a multi-channel DAQ board (Open Ephys, Intan

Technologies). Opto-tagging was performed by applying short blue light pulses to the DR (10

times 5ms pulses, every 30 seconds). Light-responsive cells were identified based on their

spiking activity within 3 ms of the application of blue light.

EEG and EMG recordings

Sleep-wake stages were manually scored based on EEG/EMG signal characteristics as

described previously (Jego et al., 2013). Briefly, wakefulness was scored when low-amplitude

desynchronized EEG activity was accompanied by high, tonic muscle activity with phasic

bursts in the EMG. NREM sleep was defined by the presence of synchronized, high amplitude

oscillations in the delta frequency band (0.5 – 4 Hz) and low EMG tone. REM sleep was

scored when the EEG showed pronounced theta oscillations with an almost complete absence

of EMG tone except for short twitches. The 24 h baseline recordings were scored in epochs of

5 s, while state-specific optogenetic stimulation recordings were scored at an epoch duration

of 1 s.

Optogenetic activation during polysomnographic recordings

For *in vivo* optogenetic stimulation experiments, the patch cord was connected to a 473nm

DPSS laser (Laserglow Technologies) via a FC/PC connector (250 um, 30126G2-250,

Thorlabs). The laser output was set to a power output of 10 mW from the tip of the optic fiber.

Trains of 5 ms pulses at 5 or 20Hz were applied to LH_{VGAT} fibers or DR Pet1-OFF/Vgat-ON

neurons and fibers using a Master-9 pulse generator (AM Systems). State-specific optical

activation was performed based on visual inspection of online EEG and EMG signal

parameters. For NREM and REM sleep-specific optogenetic activation, pulse trains were

applied through the patch cord and optic fiber when the animal spent at least 10 seconds in

NREM or REM sleep. The optogenetic stimulation was performed throughout the NREM or

REM episode until the mouse transitioned to another state. Afterwards, the animal was left

undisturbed to cycle between NREM sleep, REM sleep and wakefulness for 2-3 cycles before

the next NREM or REM sleep episode was optically stimulated. A total of 9 NREM and REM

sleep episodes per animal were optogenetically stimulated for each frequency (5 and 20 Hz).

Immunohistochemistry following sleep studies

At the end of the experiments, the mice were deeply anesthetized with an intraperitoneal

injection of pentobarbital (250 mg/kg, Streuli Pharma) and transcardially perfused with 5 ml

cold saline (0.9 % NaCl) followed by 20 ml 4% formaldehyde (Grogg Chemie).

Subsequently, the brains were removed and stored in formaldehyde over night at 4 °C for

post-fixation. The next day, the brains were transferred into phosphate-buffered saline (PBS)

containing 30 % sucrose before freezing. The brains were cut into 30 μm sections with a

cryostat (Hyrax C 25, Zeiss) and collected in PBS with 0.1 % Triton A-100 (Sigma-Aldrich),

PBST.

For immunostaining, the sections were first incubated for 1 h at room temperature in a

blocking solution containing 4 % bovine serum albumin (Sigma-Aldrich) in PBS-T. Then the

tissue was incubated with anti-GFP (1:5000, A0262, Life technologies) and anti-Tph2 (1:400,

ABN60, Merck) for 24 h. Afterwards, the sections were washed for 5 x 5 minutes in PBS-T

and then incubated with DyLight 488 and AlexaFluor 555 secondary antibody (1:500 dilution,

A96947 for eYFP, Life technologies; 1:500, A21428 for Tph2; Life technologies) at room

temperature for 1 h. The sections were washed 2 x 10 minutes in PBS-T and 2-10 minutes in

PBS and then mounted on glass slides. Cover slips were fixed on the slides with Fluoromount

(F4680, Sigma-Aldrich). Photomicrographs were taken using a confocal microscope. For

display, the image brightness and contrast was moderately adjusted in Adobe Photoshop CC.

Immunohistochemistry following in vitro recordings

Following *in vivo* electrophysiological recordings, slices were immersed in 4%

paraformaldehyde (PFA) in 0.1 M phosphate buffer (PB; pH = 7.4) at 4 °C for at least

12 h. After several washes with 0.1 M PB, slices were embedded in 10% gelatin, and

further sectioned into slices of 50-70 μm thickness in the cold PB using a vibratome VT1000S

(Leica Microsystems). Coronal sections were subjected to freeze-thaw over liquid nitrogen

after cryoprotecting in 10% and 20% sucrose. The recorded cells were first visualized with

incubation in Cy3-conjugated streptavidin (Jackson ImmunoResearch) for 2 h, diluted at

1:500 in tris buffer saline (TBS, SigmaAldrich). After examination by epifluorescence

microscopy (Olympus BX60), the sections containing the soma of the labeled neurons were

incubated in 10% normal horse serumin (NHS) in TBS to block non-specific antibody-

binding sites. Free- floating sections were incubated in primary antibodies dissolved in TBS

containing 0,1% Trinon X-100 ((TBS-Tx); Sigma Aldrich) for overnight at room temperature

((RT); 22□°C). The following primary antibodies were used: anti-5HT-Rb (1:1000, Jackson

ImmunoResearch); anti-GABA-Rb (1:1000, Jackson ImmunoResearch). After several washes

in TBS, the immunoreactions were visualized with Dylight649-DARb (1:400, Jackson

ImmunoResearch) secondary antibodies. Finally, the sections were mounted in Vectashield

mounting medium (VectorLaboratories). Images were taken by confocal microscope

(Olympus FV1000).

Immunohistochemistry following in vivo single unit recordings

After experiments the animal was quickly perfused through the heart with 50 ml of PBS

followed by 50 ml 4% PFA in 0.1 M PB. Following perfusion, brains were postfixed

overnight and stored in 0.1 M PB with 0.05% sodium azide at 4°C as a preservative. The

tissue was sectioned in PB at 50–70 μm on a vibratome (Leica VT 1000S). Coronal sections

were permeabilized in 0.3 % TBS-Tx and incubated in Cy3 or Alexa488 conjugated

streptavidin (1:500; Jackson ImmunoResearch) for 2h at RT in TBS. Images were taken by

confocal microscope (Olympus FV1000).

Statistical analysis

Data analysis was performed using Spike2 (Cambridge Electronic Design, UK), OriginPro 8.5

(MicroCal, USA), IgorPro (WaveMetrics, USA), Matlab (MathWorks, USA) and Prism

(GraphPad, USA) software. The data were tested for normality with the Shapiro-Wilk and the

Kolmogorov-Smirnov tests and accordingly analysed using parametric or non-parametric

methods. We used three-way ANOVA to test for frequency (5 versus 20 Hz, within-group

comparisons) and opsin (eYFP controls versus ChR2-expressing mice, between-groups

comparisons) effects in the latencies to transition to another state from NREM or REM sleep

of state-specific optogenetic activation experiments. PSPs evoked by photostimulation were

detected in Vm recordings using a threshold of 2xSD in a $\pm$ 50 ms time window following

photostimulation. Action potentials in extracellular recordings were detected in Spike2 using

a straight forward level threshold. Firing rate histograms were computed for each

photostimulation trial of each neuron. The control period was defined as 0.5 seconds before

photostimulation. All data are displayed as averages +/- SEM.

The firing rate change of single units upon photostimulation was assessed by subtracting the

average spike rate during the 10 seconds preceding the stimulation from the average spike rate

during optogenetic activation. To test for significant modulation, 9999 stimulation onsets

were randomly drawn from each single unit's activity time series and the firing rate change

across these random onsets was calculated. If the actual firing rate difference was larger than

95% or smaller than 5% of the randomly calculated ones, the firing of the neuron was

significantly modulated during this photostimulation. Inhibited or disinhibited neurons

showed a significant decrease or increase, respectively, of their firing rate during more than

50% of all photostimulations. Neurons whose firing rate was significantly modulated during

less than 50% of all optogenetic stimulations are reported as unresponsive.

RESULTS

LH$_{GABA}$ projections to the DR promote wakefulness

A subpopulation of LH_{VGAT} (vesicular GABA and glycine transporter) neurons causally

promotes arousal from NREM sleep (Herrera et al., 2016; Venner et al., 2016; Venner et al.,

2019). Here, we tested the role of LH_{VGAT} projections to the DR in the control of sleep-wake

behaviour. To deliver opsins to LH_{VGAT} neurons, VGAT-Cre mice were stereotactically

injected with AAV2-EF1a-DIO-eYFP (control) or AAV2-EF1a-DIO-hChR2(H134R)-eYFP

(ChR2-EYFP) in the LH for *in vivo* optogenetic activation experiments (see Materials and

Methods; Figure 1A and B). Mice were then implanted with EEG/EMG electrodes for

recording and quantification of sleep-wake states together with an optic fiber above the DR

(see Materials and Methods; Figure 1C and D). State-specific optogenetic activation of

LH_{VGAT} fibers in the DR was triggered manually after at least 10 seconds in stable NREM or

REM sleep. Blue light (473 nm) was delivered to the DR at 5Hz or 20Hz (5ms pulses, see

(Herrera et al., 2016)) in both control (eYFP) and ChR2-expressing animals continuously

until a state transition occurred (Figure 1E). Interestingly, optogenetic activation of LH_{VGAT}

projections to the DR during NREM sleep led to a rapid arousal at both 5 and 20 Hz within <

7s (Figure 1F, P < 0.0001, three-way ANOVA), whereas optical activation of these fibers

failed to significantly change REM sleep duration (Figure 1G). These results extend previous

mechanisms for LH_{VGAT} induced arousal to DR neurons in the brainstem.

304 DR_{GABA} neurons receive monosynaptic inputs from LH_{GABA} neurons

We previously found that LH_{GABA} neurons send long-range projections to various brainstem

nuclei, including the DR (Herrera et al., 2016). In order to determine the downstream

circuitries mediating this rapid arousal response, we mapped LH_{GABA} efferents utilizing

ChR2-assisted circuit mapping to functionally characterize this LH-DR circuit (Figure 2A).

We infected LH neurons irrespective of their neurochemical identity with ChR2 using a viral

construct (AAV1-CAG-ChR2-venus-WPRE-SV40) optimised for in vitro photostimulation of

axons (Petreanu et al., 2009). We found that photostimulation of ChR$_2$ expressing LH

terminals in the DR evoked inhibitory postsynaptic potentials (IPSPs, -7.9±1.03 mV peak

amplitude with a 10,91±2,41 ms decay time) in 7 DR cells (14%) recorded (Figure 2DE),

while EPSPs of 7.58±0.9 mV peak amplitude with a 11,07±2,46 ms decay time were recorded

in 10 (20%) DR cells (Figure 2E). No detectable PSPs could be recorded in the remaining

population (n=34, 66%). All DR neurons exhibiting IPSPs following LH axonal

photostimulation were confirmed to be GABAergic (Figure 2B and E) by post-hoc

immunohistochemistry (n=5) or by recording DR$_{GABA}$ neurons in GAD-GFP mice (n=3, see

Materials and Methods), one identified DR$_{GABA}$ neuron received EPSPs (Figure 2E). The

IPSPs could be abolished by SR95531 (i.e. gabazine, n= 5 out of 5 cells, 100%) but not

NBQX (n=0/5, 0%, Figure 2D), suggesting that these are monosynaptic pathways implicating

GABAa receptors. Note that this result was further confirmed by the recording of similar

IPSPs upon photostimulation of LH axon terminals in the presence of TTX and 4-AP (mean

IPSP amplitude: control: 6.5±0.6 mV, TTX/4-AP: 6.8±0.7 mV, n=6 neurons, Figure 2F and

G). When neurons were held at membrane potentials close to action potential initiation

threshold, photostimulation of LH axons also led to a prominent suppression of firing

(12.66±6.65 Hz vs. 0 Hz, MI -1.0, n=3 neurons) due to the presence of IPSPs (Figure 2D).

This suppression of firing was fully abolished when IPSPs were blocked by gabazine (n=3/3,

Figure 2D). These results show that the LH to DR monosynaptic projections are functional,

can be either inhibitory or excitatory, and that DR$_{GABA}$ neurons are preferentially inhibited by

LH$_{GABA}$ fibers.

333 LH$_{GABA}$ to DR projections can rapidly suppress the activity of DR$_{GABA}$ neurons

To test for functional connections from LH$_{GABA}$ projections to DR neurons, we recorded

extracellular single unit activity from morphologically identified neurons in the DR of

anesthetized VGAT-cre mice infected with AAV1-CAGGS-FLEX-CHR2-tdTOM.-SV40 in

the LH while photo-stimulating ChR2 expressing LH_{GABA} axons in the DR (5 light pulses of

10 ms at 20 Hz, 5 mW; Figure 3A).

When comparing the activity of DR neurons upon LH_{GABA} axonal photostimulation, we found

a rapid suppression (peak suppression: 42.37±5.94%, Figure 3B) of neuronal activity

(baseline firing: 7.72±6.43 Hz, baseline firing rate after photostimulation: 5.37±6.77 Hz,

n=25, $p < 0.001$, Wilcoxon's signed rank test, Figure 3C and D). We identified the

morphology of 8 neurons recorded and filled with the juxtacellular labeling technique. These

neurons were classified as putative DR_{GABA} neurons based on their high baseline firing rate

(≥6 Hz) morphology and location within the DR (see Materials and Methods). These results

show that LH_{GABA} neurons exert an inhibitory action onto putative DR_{GABA} neurons.

We next explored the effect of stimulating LH_{GABA} fibers in the DR in awake, head-restrained

animals while recording the activity of multiple single units in the DR (see Methods).

Comparison of the activity of DR neurons recorded in the presence and absence of LH_{GABA}

axonal photostimulation (5 light flashes of 10 ms at 20 Hz, 5 mW) further confirmed the

suppressive effect in a subset of DR neurons (2/12, 17%), while the activity of the remaining

neurons was increased (10/12, 83%). The overall activity of DR neurons was slowly (~200

ms), but persistently (~1 sec) increased (baseline firing: 7.67±6.87 Hz, baseline firing after

photostimulation: 10.98±8.920 Hz, n=12, $p < 0.001$, Wilcoxon's signed rank test, Figure 3E

and F).

The activity of identified DR_{GABA} neurons is brain state dependent

To target DR_{GABA} neurons in the DR, we made use of the intersectional virus approach (Fenno

et al., 2014). In brief, the INTRSECT system ('intronic recombinase sites enabling

combinatorial targeting') consists of orthogonal Cre and Flp recognitions sites (lox and FRT

sites, respectively) that are inserted into synthetic introns of the ChR2-eYFP open reading

frame (ORF) within an adeno-associated viral construct (AAVdj). The starting direction of the

ORF fragments between the lox and FRT recognition sites defines whether the presence of

Flp and/or Cre recombinases is required to produce a functional opsin (ChR2-eYFP or eYFP,

see Figure 4A and B).

Here, we used this approach to exclude the possibility that GABAergic markers (Vgat) are co-

expressed with serotonergic neurons (Huang et al., 2019; Ren et al., 2019; Okaty et al., 2020).

To this end, Pet1^{Flpe} - Vgat^{Cre} double transgenic mice (Vong et al., 2011; Jensen et al., 2008)

were injected with Cre-ON/Flp-OFF or Cre-ON/Flp-ON intersectional viruses (Figure 4A and

B), which allowed the targeting of ChR2-eYFP to distinct subpopulations of the DR that

expressed Pet1, a transcription factor predominantly found in 5-HT neurons (Hendricks et al.,

1999), and Vgat (Pet1-ON/Vgat-ON) or Vgat only (Pet1-OFF/Vgat-ON). The Pet1-

OFF/Vgat-ON GFP-positive neurons co-localized with Vgat neurons that were co-transfected

with a Cre-dependent mCherry virus (Figure 4C).

To investigate whether the activity of DR$_{GABA}$ neurons is modulated across sleep-wake states,

we performed opto-tag unit recording of putative DR Pet1-OFF/Vgat-ON neurons while

simultaneously conducting EEG and EMG recordings in freely moving Pet1^{Flpe}-Vgat^{Cre} mice

(see Materials and Methods; Figure 4D). These neurons were identified by their immediate

increase in spiking in response to short blue light pulses (10 times 5ms pulses, every 30

seconds) delivered to the DR via an optic fiber. A total of 9 optogenetically identified Pet1-

OFF/Vgat-ON neurons (9 out of 66 cells from 4 animals; Figure 4E and F) were recorded

across the sleep-wake cycle (Figure 4F). The activity of Pet1-OFF/VGAT-ON neurons was

modulated across sleep-wake states (Figure 4G-I, NREM vs Wake/REM: $P < 0.05$, Friedman

non-parametric test). Indeed, their firing activity increased significantly upon arousal as

compared to the 5 seconds before the transition occurred (i.e., when the muscle tone was still

low; Figure 4G, H and J; NREM-to-wake $P < 0.01$; REM-to-wake $P < 0.05$, paired t-tests). In

turn, the firing activity decreased when the mice were at the onset of NREM sleep (Wake-to-

NREM $P < 0.01$, paired t-test), while on the other hand, the transition from NREM to REM

sleep was not marked by a substantial change in firing activity (Figure 4J).

Wake promoting effect of DR$_{GABA}$ neurons

Next, we investigated whether Pet1-OFF/VGAT-ON neurons in the DR were causally

involved in arousal control. For this, we optogenetically activated DR Pet1-OFF/Vgat-ON

neurons during NREM sleep (see methods; Figure 5C). Consistent with our correlative data,

we found that photostimulation of DR Pet1-OFF/VGAT-ON neurons induced a rapid arousal

from NREM sleep when light pulses were delivered at 20Hz as compared to controls ($P <$

0.0001, Three-way ANOVA; Pet1-OFF/VGAT-ON vs Ctrl: $P < 0.0001$, Dunnett's multiple

comparison's post-hoc test), whereas 5Hz stimulation frequency did not decrease the latency

to wakefulness transitions (n = 6 animals per group; Figure 5C). The probability of REM

sleep onset decreased upon optogenetic activation of Pet1-OFF/VGAT-ON neurons during

NREM sleep (Figure 5E), suggesting an implication in REM sleep onset. However, both 5

and 20Hz stimulation frequencies during REM sleep led to an immediate arousal from REM

sleep in Pet1-OFF/Vgat-ON mice (Figure 5D, $P < 0.0001$, three-way ANOVA; Pet1-

OFF/VGAT-ON vs Ctrl: $P < 0.0001$, at both 5 and 20 Hz, Dunnett's multiple comparison's

post-hoc test). As expected, optogenetic activation of Pet1-ON/VGAT-ON neurons (n = 6

animals) did not alter sleep-wake behavior as compared to controls (n = 6 animals; Figure 5D,

E).

DR$_{GABA}$ neurons promote wakefulness through disinhibition in LH

To identify potential sleep-wake centers that may mediate the arousal promotion of DR$_{GABA}$

neurons, we performed a ChR2-assisted functional circuit mapping. Animals were prepared as

described above. We found that DR Pet1-OFF/VGAT-ON strongly innervated major sleep-

wake centers, such as the locus coeruleus (LC), the ventral tegmental area (VTA), the

centromedial thalamic nucleus (CMT), and the LH (Figure 6A and B). To further understand

the reciprocal connection between the DR and the LH, we performed optogenetic state-

specific stimulations of Pet1-OFF/VGAT-ON projecting fibers in the LH. We found that

optogenetic stimulation (20 Hz) of DR Pet1-OFF/VGAT-ON projections to the LH

reproduced the arousal-inducing effect that was observed during somatic optogenetic

activation (Figure 6C, D, NREM sleep: $P < 0.0001$, three-way ANOVA; Ctrl. versus LH

stimulation: $P < 0.001$, Dunnett's multiple comparisons post-hoc test; REM sleep: $P < 0.0001$,

three-way ANOVA; Ctrl. versus LH stimulation: $P < 0.001$, Dunnett's multiple comparisons

post-hoc test). Interestingly, optogenetic activation of Pet1-OFF/VGAT-ON projections in the

LC area, a predominant wake-promoting center, resulted in a similar effect (Ctrl. versus LC

stimulation: NREM sleep: $P < 0.0001$; REM sleep: $P < 0.001$, Dunnett's multiple comparisons

post-hoc test), while optogenetic stimulation of Pet1-OFF/VGAT-ON projections in the CMT

led to an arousal only from NREM sleep, but not from REM sleep (Ctrl. versus CMT

stimulation: NREM sleep: $P < 0.001$; REM sleep: n.s., Dunnett's multiple comparisons post-

hoc test), in line with a previous study highlighting a role of the CMT in NREM sleep

regulation (Gent et al., 2018) (Figure 6C, D).

To further delineate the mechanism by which DR Pet1-OFF/VGAT-ON neurons induce

arousal via the LH, we performed single unit recordings in the LH while simultaneously

recording EEG/EMG and conducting optical stimulations of ChR2-expressing Pet1-

OFF/VGAT-ON neurons using a similar preparation as described above (see Materials and

Methods; Figure 6E). First, we clustered all the cells that we recorded in the LH according to

their sleep-wake discharge profiles; recorded cells separated into 2 clusters, with one cluster

showing maximal activity during wakefulness (55% of cells from 2 animals; Figure 6F, G),

while the second cluster showed maximal activity during both REM sleep and wakefulness

(45% of cells from 2 animals; Figure 6F, G). Cells from both clusters were less active during

NREM sleep (Figure 6F). Interestingly, during optogenetic stimulation of DR Pet1-

OFF/VGAT-ON neurons, cells from Cluster 1 barely changed their firing rate, whereas half of

the cells from the second cluster significantly modulated their firing activity in response to

DR Pet1-OFF/VGAT-ON neuronal activation (Figure 6G). In particular, a major part of

responsive cells from Cluster 2 was disinhibited by the optogenetic activation of DR Pet1-

OFF/VGAT ON neurons, while a few cells were inhibited (Figure 5G and H, $P < 0.05$, three-

way ANOVA). These results show that the activation of DR_{GABA} neurons disinhibits the

majority of LH neurons and promotes wakefulness.

Discussion

In this study, we show that activation of LH_{GABA} axons in the DR promote wakefulness

through a local release of GABA that rapidly suppresses activity of identified DR_{GABA}

neurons but indirectly activates other non-GABAergic DR neurons. In addition, we found that

the activity of DR_{GABA} neurons is state dependent, while activation of DR_{GABA} neurons

disinhibits the majority of LH neurons to promote wakefulness. This wake promoting effect

of optogenetic activation of the LH_{GABA}-DR_{GABA} circuit promotes rapid wakefulness

selectively during NREM sleep, consistent with their activity profile across sleep-wake states

(Hassani et al., 2010; Herrera et al., 2016). These findings identify a LH_{GABA}-DR_{GABA}

feedback loop that is essential for rapid alteration of global brain states and facilitation of

wakefulness.

Using viral tracing and *in vitro* ChR$_2$ assisted circuit mapping we show that a subset of

LH$_{GABA}$ neurons establishes monosynaptic connections to DR neurons. Functional analysis of

this mapping revealed that LH$_{GABA}$ neurons exert a GABA$_A$-mediated inhibitory action on

DR$_{GABA}$ neurons leading to rapid NREM sleep-to-Wake transitions. The precise

neurochemical identity of this subset of DR-projecting LH$_{GABA}$ neurons remains to be

established in light of the numerous sub-populations amongst inhibitory neurons of the LH

(Mickelsen et al., 2017).

Local GABA release evoked by optogenetic activation of LH$_{GABA}$ axons in the DR promotes

wakefulness suppressing the activity of DR$_{GABA}$ neurons. This occurs through a direct

synaptic inhibition of DR$_{GABA}$ neurons mediated by GABA$_A$ receptors and leads to a

prominent suppression of firing in DR$_{GABA}$ neurons *in vivo*. Consistent with these *in*

vivo recordings we show that the activity of identified DR$_{GABA}$ cells is strongly modulated

across NREM sleep-to-wake transitions and optogenetic activation of the LH$_{GABA}$-DR circuit

promotes rapid wakefulness selectively during NREM sleep, but not during REM sleep. This

result is in line with previous studies reporting sleep/wake transitions upon optogenetic

activation of LH$_{GABA}$ fibers in the medial septum, locus coeruleus, reticular thalamic nucleus

and ventrolateral preoptic area (Herrera et al., 2016; Venner et al., 2016; Venner et al., 2019).

Our results demonstrate that LH$_{GABA}$ neurons directly inhibit DR$_{GABA}$ neurons and this

probably results in a disinhibition of various DR neurons. Revealing the exact targets of this

disinhibition awaits further investigation and are hard to predict given the extreme

neurochemical and physiological diversity of the raphe nuclei (Domonkos et al., 2016; Szonyi

et al., 2016; Sos et al., 2017). A phasic increase in 5-HT neuron activity can rapidly and

prominently affect cortical activity (Lottem et al., 2016) and thus may lead to brain state

changes. However, recently the selective tonic activation of DR_{5-HT} neurons has been shown

to promote sleep, while burst stimulation induces wakefulness (Oikonomou et al., 2019). In

light of the present results, it is possible that the phasic activity changes of DR neurons and

their consequences at the cortical level at least partly originate from the LH_{GABA} neuron

activation during sleep to wake transitions.

DR_{GABA} neuronal stimulation also promoted wakefulness, possibly by suppressing or

activating local sleep- or wake-promoting neurons in the LH, respectively. One may speculate

that GABA release from DR_{GABA} synapses could suppress the activity of sleep-promoting

neurons including REM-sleep promoting MCH neurons (Jego et al., 2013; Konadhode et al.,

2013; Tsunematsu et al., 2014), REM active LH_{VGAT} neurons (Hassani et al., 2010; Oesch et

al., 2020), REM-active Lh6x GABA neurons from the ventral zona incerta (Liu et al., 2017).

In contrast, it could inhibit local inhibitory circuits that would result in the activation of the

wake-promoting hypocretins/orexins neurons (Adamantidis et al., 2007), or wake-active

LH_{VGAT} neurons (Hassani et al., 2010; Herrera et al., 2016; Oesch et al., 2020). Whether and

how this dual modulatory activity operates during the sleep-wake cycle awaits further

investigation.

Taken together, a brain area involved in energy balance and arousal can affect the activity of

the DR, an area involved in the control of higher brain functions including reward, patience,

mood and sensory coding suggesting a potential route for interactions between metabolic

states and brain states.

FIGURE LEGENDS

Figure 1. Stimulation of LH$_{GABA}$ fibers in the DR promotes wakefulness. (A) Schematics

of the experimental design: AAV-EF1a-DIO-ChR2-eYFP or AAV-EF1a-DIO-eYFP was

injected in the LH of Vgat-cre mice. Following 4 weeks of expression LH$_{GABA}$ fibers were

photostimulated in the DR with simultaneous polysomnographic recordings for sleep stage

scoring. (B) Example ChR2-eYFP expression in the LH. Fluorescent image overlaid on the

stereotaxic atlas (AP: 4.45). (C) Location of the optical fiber in the DR used for

photostimulation. Merged fluorescent image showing LH$_{GABA}$ fibers expressing ChR2-eYFP

(green) and TPH immunhistochemistry (red) to indicate the location of TPH synthetizing

serotonergic neurons within the DR. (D) location of all implanted optical fibers in 11 mice.

Green lines: Vgat::ChR2-expressing mice (n=6), gray lines: eYFP controls (n=5), Aq:

aqueduct). (E) (Top) Schematic experimental timeline for sleep-stage specific

photostimulation. (Bottom) Representative EEG/EMG traces illustrate behavioural response

to optogenetic stimulation. Note the rapid EEG desynchronization and EMG activation upon

photostimulation indicating a NREM-to-Wake transition. (F) Latency of NREM to wake

transition caused by the photostimulation of LH$_{GABA}$ axons in the DR. (G) Latency of REM

to wake transition caused by the photostimulation of LH$_{GABA}$ axons in the DR. * $P < 0.05$.

Figure 2. LH projections synaptically target DR neurons. (A) Schematic of the

experimental design: AAV-ChR2-Venus injection in the LH of WT mice leads to prominent

ChR$_2$ expression in DR projecting axons. Four weeks after the infection local axons of LH

neurons are photostimulated in the DR with simultaneous whole-cell patch clamp recordings

in identified DR neurons. (B) Fluorescent images (Streptavidin, GABA immunostaining,

merged) of a DR_{GABA} neuron responding to LH axonal photostimulation. (C) Membrane

potential responses of the neuron in (B) to hyperpolarizing and depolarizing current steps

shows fast action potential output with moderate frequency adaptation. (D) The

photostimulation of LH axons leads to monosynaptic IPSPs that are resistant to iGluR

blockade (NBQX), but blocked by SR95531 (i.e. gabazine, GBZ). (E) Distribution of PSP

amplitudes in all the neurons recorded (n=51), green circles indicate identified DR_{GABA}

neurons. (F) Membrane potential responses of a physiologically identified DR_{GABA} neuron.

(G) The photostimulation of LH axons leads to IPSPs in the neuron shown in (F) that persist

following the application of TTX and 4-AP.

Figure 3. LH_{GABA} inputs control DR neuronal activity. (A) Schematic of the experimental

design: AAV-Flex-ChR2-td tomato injection in the LH of VGAT mice leads to prominent

ChR2 expression in DR projecting axons. 4 weeks after the infection local axons of LH_{GABA}

neurons are photostimulated in the DR with simultaneous single unit juxtacellular

recoding/labeling. (B) Example of spontaneous activity of a DR_{GABA} neuron and its response

to local photostimulation of LH_{GABA} axons. (Top) Single-trial raw data aligned on

photostimulation onset (blue horizontal bar, 5 mW, 10 ms pulses at 50 Hz). Middle, Raster

plot in which each tic is a single-unit spike and each row represents a single trial. Red line:

photostimulation onset. (Bottom) PSTH of the unit shown above. (Right) Fluorescent pictures

showing the neuron recorded in (B) filled with Neurobiotin using the juxtacellular filling

technique. Left: streptavidin, middle: VGAT immunostaining, right: merged picture. (C)

Modulation index of all recorded neurons under anaesthesia. (D) Grand average PSTH of all

neurons recorded neurons under anaesthesia. (E) Modulation index of all recorded neurons in

awake animals (inh: inhibited, n.s.: not significant, exc: excited). (F) Grand average PSTH of

all neurons recorded neurons in awake animals.

Figure 4. The activity of optogenetically identified DR$_{GABA}$ neurons are modulated across the sleep-wake cycle. (A) Schematics of intersectional viral strategy: ChR2-Con/off-eYFP was targeted to DR neurons in Pet1^{Flpe}-Vgat^{Cre} double transgenic mice. (B) Schematics of INTRSECT system (Fenno et al., 2014): FRT (grey triangles) and lox sites (light blue triangles) were introduced in intronic sequences within the ChR2 ORF allowing for directional control of these ORFs. In the Con/Foff-ChR2 construct, the presence of Cre recombinase produces a functional ChR2. However, the presence of Flp recombinase would invert the exons, thus yielding an inactivated state of ChR2. In the Con/Fon-ChR2 construct, the presence of both Cre and Flp recombinase is required to produce a functional ChR2. (C) ChR2 expression is restricted to DR$_{GABA}$ (Vgat-mCherry expressing) neurons. (D) Schematics of the experimental design: simultaneous DR Pet1-OFF/Vgat-ON neuron and and EEG recordings in behaving mice. (E) Example recording of a DR Pet1-OFF/Vgat-ON neuron throughout various sleep/wake states. (Top) raw trace voltage recording shows clear action potential firing upon photostimulation (blue ticks). (Bottom) Hypnogram and spike ticks of the neuron showed above (W: wake, R: REM, N: NREM). (F) Averaged spike waveforms of optogenetically identified DR Pet1-OFF/Vgat-ON neurons (n = 9 neurons from n = 4 mice). DR Pet1-OFF/Vgat-ON neurons were identified by their immediate firing in response to pulses of blue light generated by a 453 nm laser (10 pulses of 5ms every 30 sec). (G, H) Raster plot of a DR Pet1-OFF/Vgat-ON neuron during NREM-to-wake transitions (G) and REM-to-wake transitions (H) with spike ticks of the neuron (Top) and PSTH of the neuron across all transitions (Bottom). (I) Firing rates of the 9 DR Pet1-OFF/Vgat-ON neurons across wakefulness, NREM and REM sleep. (J) PSTH (Mean + SEM) of DR Pet1-OFF/Vgat-ON neurons during the different sleep state transitions.

Figure 5. Stimulation of DR$_{GABA}$ neurons promotes wakefulness. (A) Schematic of the

experimental strategy: various intersectional viral constructs carrying the coding sequence of

either eYFP (control) or ChR2 are injected in the DR of Pet1^{Flpe} - VGAT^{Cre} double transgenic

mice. The DR is photostimulated 4 weeks post-infection. (B) Location of the fiberoptic

canulae in all the animals used: left, gray lines: control, middle, green lines: Pet1-

OFF/VGAT-ON, right, blue lines: Pet1-ON/VGAT-ON. (C) Latency of NREM to wake

transition caused by the photostimulation of the DR. (D) Latency of REM to wake transition

caused by the photostimulation of the DR. (E) State transition probabilities at 5 versus 20 Hz

photostimulation for all three viral constructs. * $P < 0.05$.

Figure 6. DR$_{GABA}$ neurons promote arousal through disinhibition in the LH. (A) (Top

left) Schematic of the experimental design: ChR2 is expressed in DR$_{GABA}$ neurons following

a viral injection in Pet1-Flpe x VGAT-Cre mice. Axonal projections of DR$_{GABA}$ neurons

express ChR2 in various DR target areas including the LH (top right), centromedial thalamus

(CMT, bottom left) and locus coeruleus (LC, bottom right). (B) Normalized intensity of the

fluorescent eYFP signal in various projection areas of the DR (PFC: prefrontal cortex, NAc:

nucleus accumbens, HDB: horizontal diagonal band of Brocca, MS: medial septum, LHb:

lateral habenula, RTN: thalamic reticular nucleus, CeA: central amygdala, CA1: hippocampus

CA1 region, SNc: substantia nigra pars compacta). (C) Latency of NREM to wake transition

caused by photostimulation of DR$_{GABA}$ projections in the LH, CMT and LC. (D) Latency of

REM to wake transition caused by photostimulation of DR$_{GABA}$ projection targets (LH, CMT,

LC). (E) Schematic of the experimental design: ChR2 expressing Pet1-OFF/Vgat-ON neurons

are photostimulated in the DR while single units are recorded in the LH of freely moving

mice. (F) Activity of all recorded LH neurons during various sleep-wake stages differentiates

two clusters of neurons: Cluster 1 containing wake active neurons while Cluster 2 neurons do not show significant firing rate increases during the awake state. (G) Pie chart of all Cluster 1 and 2 LH neurons constructed on the basis of their response to the photostimulation of DR_{GABA} neurons. (H) Effect of DR_{GABA} photostimulation on the state-dependent firing of inhibited (left) and disinhibited (right) neurons. For details see main text. * $P < 0.05$.

Acknowledgements

This work has been supported by the Hungarian Scientific Research Fund (grant NF 123831 to MLL), the Hungarian Brain Research Program (grant KTIA_NAP_13-2-2014-0014 to MLL), NIH R01 DA034022 (SMD) and the German Research Foundation (GA 2410/1-1 to MG), the Human Frontier Science Program RGY0076/2012 (AA), Swiss National Science Foundation (grant 31003A_156156 to AA), European Research Council (grant 725850 to AA). MLL is a grantee of the János Bolyai Fellowship.

REFERENCES

Adamantidis AR, Zhang F, Aravanis AM, Deisseroth K, de Lecea L (2007) Neural substrates

of awakening probed with optogenetic control of hypocretin neurons. Nature 450:420-

424.

Allers KA, Sharp T (2003) Neurochemical and anatomical identification of fast- and slow-

firing neurones in the rat dorsal raphe nucleus using juxtacellular labelling methods in

vivo. Neuroscience 122:193-204.

Arrigoni E, Chee MJS, Fuller PM (2019) To eat or to sleep: That is a lateral hypothalamic

question. Neuropharmacology 154:34-49.

Bang SJ, Commons KG (2012) Forebrain GABAergic projections from the dorsal raphe

nucleus identified by using GAD67-GFP knock-in mice. The Journal of comparative

neurology 520:4157-4167.

Bernardis LL, Bellinger LL (1996) The lateral hypothalamic area revisited: ingestive

behavior. Neuroscience and biobehavioral reviews 20:189-287.

Bonnavion P, Mickelsen LE, Fujita A, de Lecea L, Jackson AC (2016) Hubs and spokes of

the lateral hypothalamus: cell types, circuits and behaviour. The Journal of physiology

594:6443-6462.

Carter ME, Adamantidis A, Ohtsu H, Deisseroth K, de Lecea L (2009) Sleep homeostasis

modulates hypocretin-mediated sleep-to-wake transitions. The Journal of neuroscience

29:10939-10949.

Carter ME, Brill J, Bonnavion P, Huguenard JR, Huerta R, de Lecea L (2012) Mechanism for

Hypocretin-mediated sleep-to-wake transitions. Proceedings of the National Academy

of Sciences of the United States of America 109:E2635-2644.

Cohen JY, Amoroso MW, Uchida N (2015) Serotonergic neurons signal reward and
punishment on multiple timescales. eLife 4.

Correia PA, Lottem E, Banerjee D, Machado AS, Carey MR, Mainen ZF (2017) Transient
inhibition and long-term facilitation of locomotion by phasic optogenetic activation of
serotonin neurons. eLife 6.

Descarries L, Watkins KC, Garcia S, Beaudet A (1982) The serotonin neurons in nucleus
raphe dorsalis of adult rat: a light and electron microscope radioautographic study. The
Journal of comparative neurology 207:239-254.

Domonkos A, Nikitidou Ledri L, Laszlovszky T, Cserep C, Borhegyi Z, Papp E, Nyiri G,
Freund TF, Varga V (2016) Divergent in vivo activity of non-serotonergic and
serotonergic VGluT3-neurones in the median raphe region. The Journal of physiology
594:3775-3790.

Fenno LE, Mattis J, Ramakrishnan C, Hyun M, Lee SY, He M, Tucciarone J, Selimbeyoglu
A, Berndt A, Grosenick L, Zalocusky KA, Bernstein H, Swanson H, Perry C, Diester
I, Boyce FM, Bass CE, Neve R, Huang ZJ, Deisseroth K (2014) Targeting cells with
single vectors using multiple-feature Boolean logic. Nature methods 11:763-772.

Fonseca MS, Murakami M, Mainen ZF (2015) Activation of dorsal raphe serotonergic
neurons promotes waiting but is not reinforcing. Current biology : CB 25:306-315.

Fuller PM, Sherman D, Pedersen NP, Saper CB, Lu J (2011) Reassessment of the structural
basis of the ascending arousal system. The Journal of comparative neurology 519:933-
956.

Gent TC, Bandarabadi M, Herrera CG, Adamantidis AR (2018) Thalamic dual control of
sleep and wakefulness. Nature neuroscience 21:974-984.

Hassani OK, Henny P, Lee MG, Jones BE (2010) GABAergic neurons intermingled with

orexin and MCH neurons in the lateral hypothalamus discharge maximally during

sleep. The European journal of neuroscience 32:448-457.

Hendricks T, Francis N, Fyodorov D, Deneris ES (1999) The ETS domain factor Pet-1 is an

early and precise marker of central serotonin neurons and interacts with a conserved

element in serotonergic genes. The Journal of neuroscience 19:10348-10356.

Herrera CG, Cadavieco MC, Jego S, Ponomarenko A, Korotkova T, Adamantidis A (2016)

Hypothalamic feedforward inhibition of thalamocortical network controls arousal and

consciousness. Nature neuroscience 19:290-298.

Huang KW, Ochandarena NE, Philson AC, Hyun M, Birnbaum JE, Cicconet M, Sabatini BL

(2019) Molecular and anatomical organization of the dorsal raphe nucleus. eLife 8.

Jego S, Glasgow SD, Herrera CG, Ekstrand M, Reed SJ, Boyce R, Friedman J, Burdakov D,

Adamantidis AR (2013) Optogenetic identification of a rapid eye movement sleep

modulatory circuit in the hypothalamus. Nature neuroscience 16:1637-1643.

Jensen P, Farago AF, Awatramani RB, Scott MM, Deneris ES, Dymecki SM (2008)

Redefining the serotonergic system by genetic lineage. Nature neuroscience 11:417-

419.

Jones BE (2020) Arousal and sleep circuits. Neuropsychopharmacology : official publication

of the American College of Neuropsychopharmacology 45:6-20.

Kapoor V, Provost AC, Agarwal P, Murthy VN (2016) Activation of raphe nuclei triggers

rapid and distinct effects on parallel olfactory bulb output channels. Nature

neuroscience 19:271-282.

Konadhode RR, Pelluru D, Blanco-Centurion C, Zayachkivsky A, Liu M, Uhde T, Glen WB,

Jr., van den Pol AN, Mulholland PJ, Shiromani PJ (2013) Optogenetic stimulation of

MCH neurons increases sleep. The Journal of neuroscience 33:10257-10263.

Liu K, Kim J, Kim DW, Zhang YS, Bao H, Denaxa M, Lim SA, Kim E, Liu C, Wickersham
IR, Pachnis V, Hattar S, Song J, Brown SP, Blackshaw S (2017) Lhx6-positive
GABA-releasing neurons of the zona incerta promote sleep. Nature 548:582-587.

Liu Z, Zhou J, Li Y, Hu F, Lu Y, Ma M, Feng Q, Zhang JE, Wang D, Zeng J, Bao J, Kim JY,
Chen ZF, El Mestikawy S, Luo M (2014) Dorsal raphe neurons signal reward through
5-HT and glutamate. Neuron 81:1360-1374.

Lottem E, Lorincz ML, Mainen ZF (2016) Optogenetic Activation of Dorsal Raphe Serotonin
Neurons Rapidly Inhibits Spontaneous But Not Odor-Evoked Activity in Olfactory
Cortex. The Journal of neuroscience 36:7-18.

Lowry CA, Hale MW, Evans AK, Heerkens J, Staub DR, Gasser PJ, Shekhar A (2008)
Serotonergic systems, anxiety, and affective disorder: focus on the dorsomedial part of
the dorsal raphe nucleus. Annals of the New York Academy of Sciences 1148:86-94.

Mickelsen LE, Kolling FWt, Chimileski BR, Fujita A, Norris C, Chen K, Nelson CE, Jackson
AC (2017) Neurochemical Heterogeneity Among Lateral Hypothalamic
Hypocretin/Orexin and Melanin-Concentrating Hormone Neurons Identified Through
Single-Cell Gene Expression Analysis. eNeuro 4.

Miyazaki K, Miyazaki KW, Doya K (2011) Activation of dorsal raphe serotonin neurons
underlies waiting for delayed rewards. The Journal of neuroscience 31:469-479.

Nectow AR, Schneeberger M, Zhang H, Field BC, Renier N, Azevedo E, Patel B, Liang Y,
Mitra S, Tessier-Lavigne M, Han MH, Friedman JM (2017) Identification of a
Brainstem Circuit Controlling Feeding. Cell 170:429-442 e411.

Oesch LT, Gazea M, Gent TC, Bandarabadi M, Gutierrez Herrera C, Adamantidis AR (2020)
REM sleep stabilizes hypothalamic representation of feeding behavior. Proceedings of
the National Academy of Sciences of the United States of America 117:19590-19598.

Oikonomou G, Altermatt M, Zhang RW, Coughlin GM, Montz C, Gradinaru V, Prober DA

(2019) The Serotonergic Raphe Promote Sleep in Zebrafish and Mice. Neuron

103:686-701 e688.

Okaty BW, Sturrock N, Escobedo Lozoya Y, Chang Y, Senft RA, Lyon KA, Alekseyenko

OV, Dymecki SM (2020) A single-cell transcriptomic and anatomic atlas of mouse

dorsal raphe Pet1 neurons. eLife 9.

Petreanu L, Mao T, Sternson SM, Svoboda K (2009) The subcellular organization of

neocortical excitatory connections. Nature 457:1142-1145.

Pinault D (1996) A novel single-cell staining procedure performed in vivo under

electrophysiological control: morpho-functional features of juxtacellularly labeled

thalamic cells and other central neurons with biocytin or Neurobiotin. Journal of

neuroscience methods 65:113-136.

Ren J, Isakova A, Friedmann D, Zeng J, Grutzner SM, Pun A, Zhao GQ, Kolluru SS, Wang

R, Lin R, Li P, Li A, Raymond JL, Luo Q, Luo M, Quake SR, Luo L (2019) Single-

cell transcriptomes and whole-brain projections of serotonin neurons in the mouse

dorsal and median raphe nuclei. eLife 8.

Riemann D, Krone LB, Wulff K, Nissen C (2020) Sleep, insomnia, and depression.

Neuropsychopharmacology : official publication of the American College of

Neuropsychopharmacology 45:74-89.

Saper CB, Fuller PM, Pedersen NP, Lu J, Scammell TE (2010) Sleep state switching. Neuron

68:1023-1042.

Seo C, Guru A, Jin M, Ito B, Sleezer BJ, Ho YY, Wang E, Boada C, Krupa NA, Kullakanda

DS, Shen CX, Warden MR (2019) Intense threat switches dorsal raphe serotonin

neurons to a paradoxical operational mode. Science 363:538-542.

Sos KE, Mayer MI, Cserep C, Takacs FS, Szonyi A, Freund TF, Nyiri G (2017) Cellular

architecture and transmitter phenotypes of neurons of the mouse median raphe region.

Brain structure & function 222:287-299.

Steiger A, Pawlowski M (2019) Depression and Sleep. International journal of molecular

sciences 20.

Stuber GD, Wise RA (2016) Lateral hypothalamic circuits for feeding and reward. Nature

neuroscience 19:198-205.

Szonyi A, Mayer MI, Cserep C, Takacs VT, Watanabe M, Freund TF, Nyiri G (2016) The

ascending median raphe projections are mainly glutamatergic in the mouse forebrain.

Brain structure & function 221:735-751.

Tamamaki N, Yanagawa Y, Tomioka R, Miyazaki J, Obata K, Kaneko T (2003) Green

fluorescent protein expression and colocalization with calretinin, parvalbumin, and

somatostatin in the GAD67-GFP knock-in mouse. The Journal of comparative

neurology 467:60-79.

Tsunematsu T, Ueno T, Tabuchi S, Inutsuka A, Tanaka KF, Hasuwa H, Kilduff TS, Terao A,

Yamanaka A (2014) Optogenetic manipulation of activity and temporally controlled

cell-specific ablation reveal a role for MCH neurons in sleep/wake regulation. The

Journal of neuroscience 34:6896-6909.

Venner A, Anaclet C, Broadhurst RY, Saper CB, Fuller PM (2016) A Novel Population of

Wake-Promoting GABAergic Neurons in the Ventral Lateral Hypothalamus. Current

biology : CB 26:2137-2143.

Venner A, De Luca R, Sohn LT, Bandaru SS, Verstegen AMJ, Arrigoni E, Fuller PM (2019)

An Inhibitory Lateral Hypothalamic-Preoptic Circuit Mediates Rapid Arousals from

Sleep. Current biology : CB 29:4155-4168 e4155.

Verret L, Goutagny R, Fort P, Cagnon L, Salvert D, Leger L, Boissard R, Salin P, Peyron C,

769   Luppi PH (2003) A role of melanin-concentrating hormone producing neurons in the

770   central regulation of paradoxical sleep. BMC neuroscience 4:19.

771   Vong L, Ye C, Yang Z, Choi B, Chua S, Jr., Lowell BB (2011) Leptin action on GABAergic

772   neurons prevents obesity and reduces inhibitory tone to POMC neurons. Neuron

773   71:142-154.

774   Weber F, Dan Y (2016) Circuit-based interrogation of sleep control. Nature 538:51-59.

775   Weissbourd B, Ren J, DeLoach KE, Guenthner CJ, Miyamichi K, Luo L (2014) Presynaptic

776   partners of dorsal raphe serotonergic and GABAergic neurons. Neuron 83:645-662.

777   Xu M, Chung S, Zhang S, Zhong P, Ma C, Chang WC, Weissbourd B, Sakai N, Luo L,

778   Nishino S, Dan Y (2015) Basal forebrain circuit for sleep-wake control. Nature

779   neuroscience 18:1641-1647.

A AAV-DIO-ChR2
AAV-DIO-eYFP
473 nm

-1.4mm
-4.5mm
VGAT-IRES-Cre

B ChR2-eYFP

C ChR2-eYFP / Tph2

D Aq

E
State onset → State transition
10 sec | STIM (5 or 20 Hz) |

Vgat::ChR2
20 Hz

NREM | WAKE

F NREM

Transition latency [sec]

5 Hz   20 Hz

G REM

Transition latency [sec]

5 Hz   20 Hz

■ eYFP Control (n=5)
■ Vgat::ChR2 (n=6)

A AAV-ChR2-Venus WT mice

LH

aq

DR

Record/PS DRN

B

C

D

Con

NBQX

GBZ

10 mV

100 ms

E

GABA

PSP amplitude (mV)

F

20 mV

200 ms

G

Con

TTX
4AP

5 mV

A INTRSECT viruses

Pet1-Flpe x Vgat-Cre mice

B

Con/Foff-ChR2

Exon 1 — Exon 2 — Exon 3

↓ +Cre

Exon 1 — Exon 2 — Exon 3
Functional ChR2

↑ +Cre +Flp

Con/Fon-ChR2

Exon 3 — Exon 2 — Exon 1

C Pet1^OFF-Vgat^ON
Vgat::mCherry

D Unit & EEG/EMG recording
Laser 453 nm
Micro-drive

E R N W
10 s

F 100 μm

G R N W
Transition nr
Firing rate (Hz)
Time (s)

H R N W
Transition nr
Firing rate (Hz)
Time (s)

I Firing rate (Hz)
Wake   NREM   REM

J Firing rate (Hz)
N→W
N→R
R→W
W→N
Time (s)

A Pet1^{OFF}-Vgat^{ON}

Pet1-Flpe x Vgat-Cre mice

B

C NREM STIMS

D REM STIMS

Control (eYFP) ■ Pet1^{OFF}-Vgat^{ON} ■

E Record LH

PS DRN

F

G Cluster 1 (n=35)

Cluster 2 (n=28)

■ Unresponsive
■ Inhibited
■ Disinhibited

H Inhibited

Disinhibited

□ Before STIM
■ During STIM

A INTRSECT viruses

PS DRN

Pet1-Flpe x Vgat-Cre mice

B

C NREM-specific stimulations

Transition latency [sec]

$*$

5 Hz   20 Hz

D REM-specific stimulations

Transition latency [sec]

$*$ $*$

5 Hz   20 Hz

E

Transition probability (%)

5 Hz   20 Hz

Control (n=6)
Pet1^{ON}-Vgat^{ON} (n=6)
Pet1^{OFF}-Vgat^{ON} (n=6)

NREM to Wake
NREM to REM