Title: Phylogenomic conflict coincides with rapid morphological innovation

Authors: Caroline Parins-Fukuchi[a,1], Gregory W. Stull[b,c], Stephen A. Smith[d,1]

[a]Department of Geophysical Sciences, University of Chicago, Chicago, IL, USA

[b]Germplasm Bank of Wild Species, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, Yunnan 650201, China

[c]Department of Botany, Smithsonian Institution, Washington, DC 20013, USA

[d]Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, Michigan 48109, USA

[1]To whom correspondence may be addressed. Email: parinsfukuchi@uchicago.edu or eebsmith@umich.edu.

Abstract

Evolutionary biologists have long recognized that rates of phenotypic change vary widely throughout time and across lineages. Researchers have been particularly fascinated with evolutionary 'radiations'--periods during which novel phenotypic characteristics and new species emerge at an unusually high rate. Despite considerable speculation and empirical testing of the geologic, environmental, and ecological contexts coinciding with the radiations of major lineages, our understanding of how population processes during these epochs contribute to emergent macroevolutionary patterns in phenotypic change remains poor. However, the increasing prevalence of phylogenomic datasets over the past decade has revealed that phylogenomic conflict, frequently caused by population-level processes, is often highest at the origin of major lineages. We reconstruct patterns in both rates of morphological change and levels of phylogenomic conflict across six clades to identify whether the large spikes in the rate of phenotypic change that accompany major radiations coincide with elevated conflict. We found that the most conflict-rich regions of the six clades also tended to experience the highest rates of morphological evolution, suggesting that the complex population processes that underlie evolutionary radiations leave signatures at deep phylogenetic scales in both genomic and phenotypic data. Thus, closer examination of the drivers and implications of phylogenomic conflict may permit us to make improved connections between micro- and macroevolution, vastly increasing our understanding of the processes that shape the origin of major lineages across the Tree of Life.

Introduction

Phylogenetic conflict, where gene-trees disagree about species tree resolution, is common across genomes and throughout the tree of life (Smith et al. 2015; Chen et al. 2017; Smith et al. 2020). Conflict is often driven by population processes such as incomplete lineage sorting (ILS), hybridization, and rapid evolution. Regions of high conflict often coincide with the emergence of major clades, such as mammals (e.g., Song et al. 2012), angiosperms (e.g., Wickett et al. 2014), and metazoa (e.g., Nosenko et al. 2013). This suggests that heterogeneity in the population

processes underlying the origins of modern biodiversity has been recorded in particular nodes
across the tree of life (e.g., Smith et al. 2020, Leebens-Mack et al. 2019).
Most large-scale phylogenetic and phylogenomic studies meant to resolve species
relationships have treated conflicting gene regions as an analytical nuisance to be filtered or
accommodated (e.g., Song et al. 2012; Wickett et al. 2014; Leebens-Mack et al. 2019). However,
since conflict might be thought of as the phylogenomic imprint of past population genetic
processes, studying its correlation with other macroevolutionary patterns may shed light on the
microevolutionary processes underlying major transitions across the tree of life. Researchers
have long recognized that rates of morphological change vary across the tree of life, with
pronounced bursts in morphological change interspersed with periods of relative stasis (Simpson
1944, Eldredge and Gould 1972). However, the relationship between rates of morphological
change and genomic evolution is not well-understood. In particular, it may be worthwhile to
examine whether phylogenomic conflict tends to coincide with periods of rapid morphological
change. This may help to identify whether genomic conflict and morphological innovation
detectable over macroevolutionary timescales stem from the same population processes.
Anecdotally, conflict often appears to coincide with important episodes of morphological
differentiation among major lineages. For example, the major differences in life history and body
plan that distinguish mammalian orders emerged rapidly following the K-Pg mass extinction
(O'Leary et al. 2013). The phylogenetic resolution of these early diverging lineages has proved
difficult and attempts at resolution have included the use of both sophisticated methodological
approaches and the generation of large-scale genomic datasets (Morgan et al. 2013; Romiguier et
al. 2013; Teeling et al. 2013). The early avian radiation has experienced similar challenges and is
notable for the establishment of disparate lineages and morphological innovation leading the
extant diversity. Several large-scale studies using massive genomic datasets have revealed
extensive conflict among phylogenetic branches coinciding with the early radiation of crown
Aves (Jarvis et al. 2014; Prum et al. 2015). Since the origin of land plants, there have been
numerous major phases of morphological and ecological innovation, including the initial
radiation of vascular plant body plans in the late Silurian (Kenrick and Crane 1997); the
origination of major lycophyte and euphyllophyte structures (e.g., roots, megaphyllous leaves,
seeds, wood) by the late Devonian or earlier (Gensel et al. 2020); radiations of ferns, seed ferns,
and lycophytes with diverse life histories during the Carboniferous (Stewart and Rothwell 1993);
radiations of various gymnosperm lineages in the Permian and different parts of the Mesozoic
(Beck 1988); and distinct phases of angiosperm radiation from the Cretaceous to the present
(Friis et al. 2011). As with the vertebrate lineages, the origins of many major plant clades show
elevated levels of phylogenomic conflict. Many of these have been the subject of intense analysis
and debate (e.g., Drew et al. 2014; Leebens-Mack et al. 2019; Smith et al. 2020) and recent
sequencing efforts have failed to confidently resolve most of these challenges.
Here we examine whether periods of high genomic conflict coincide with episodes of
rapid morphological evolution across several phylogenetically disparate clades. Many clades
show qualitative evidence that time periods with particularly rampant conflict are often
associated with important phenotypic changes. However, if phylogenomic conflict indeed
represents the remnants of complex population processes that are adaptively important, we
reasoned that there should be a correspondence between areas of high conflict and elevated rates
of phenotypic evolution. To examine whether this is the case, we reconstructed concurrent
patterns in phylogenomic conflict and phenotypic evolution across six major clades of the

eukaryotic tree of life: crown Aves, Mammalia, and four major lineages of vascular plants
(Ericales, Caryophyllales, Polypodiopsida, and Acrogymnospermae).
93
Results
We assembled and analyzed six genomic datasets for plants, mammals, and birds.
Specifically, we examined published phylogenomic datasets including 624 genes of
Caryophyllales from Yang et al. (2018); 387 genes of Ericales from Larson et al. (2020); 1332
gene regions of Polypodiopsida from Shen et al. (2018); 1308 gene regions representing
Acrogymnospermae from Ran et al. (2018); 424 genes of Mammalia from Song et al. (2012);
and 259 gene regions of Aves from Prum et al. (2015). We gathered publicly available
phenotypic matrices corresponding to each of these datasets (see Materials and Methods). For the
Caryophyllales, a suitable morphological matrix is not available, and so we examined the
phylogenetic locations of major changes in the betalain and photosynthetic pathways. Generally,
broad phenotypic matrices that cover the taxa in each of these datasets are rare or non-existent.
Furthermore, overlap between the genomic and phenotypic matrices was such that each needed
to be reduced to ensure comparable edges were being compared. Divergence times continue to
be controversial across the tree of life. To overcome these difficulties, we used one set of
divergence times and one topology for both the morphological and molecular analyses (see
Materials and Methods).
We found that mammals experienced elevated levels of conflict near the K-Pg boundary,
coinciding in time with the origin of most placental orders (Fig. 1). We compared these genomic
patterns to phenotypic rates calculated from two independent datasets: a dataset of mammalian
body masses (Jones et al. 2009), which we interpreted as a proxy for many strongly correlated
aspects of life history and ecology (Calder 1983), and a massive 'phenomic' matrix of >4000
qualitative characters (O'Leary et al. 2013). These analyses revealed that, in addition to
witnessing extensive phylogenomic conflict, the epoch coinciding with the early divergence of
major placental lineages also witnessed a peak in rates of phenotypic evolution. The two
phenotypic datasets displayed remarkably similar patterns, showing that both life history and
anatomical characteristics diversified rapidly during the early divergences of placental lineages.
Unlike the angiosperm examples, but more similar to the fern and gymnosperms examples (albeit
with an order of magnitude higher sampling of characters), the 'phenomic' dataset examined
here sampled characters at all phenotypic levels. Although skeletal and dental characters
comprised the majority of the dataset, characters representing internal and external organs,
musculature, and behavior were also included. It has previously been recognized that the life
history habits displayed by major mammalian lineages were established early during the
radiation of major mammalian clades (Stearns 1983), but our analyses demonstrate that these
changes occurred as rapid bursts during a single temporal episode that also witnessed both rapid
rates of anatomical change and high phylogenomic discordance.
Our genomic analysis revealed rampant conflict among early-diverging avian lineages
immediately following the K-Pg boundary (Fig. 1). We also identified an additional episode of
increased conflict that occurred at the onset of the Miocene. We compared this genomic pattern
to those displayed by two morphological datasets: a large qualitative character matrix (Livezey
and Zusi 2007) and a dataset representing the scores calculated from the first three axes of a
principal component analysis performed on a dataset of linear measurements that corresponded
to body size, relative beak size, and relative wing size, respectively (Pigot 2019). In both
phenotypic datasets, we found evidence for elevated rates of evolution during the epoch

immediately following the K-Pg, coincident with the elevated levels of genomic conflict.
Although qualitatively similar in this regard, we found that each dataset displayed outliers in rate
not shared by the other, with the qualitative character matrix displaying two instances of
extremely high rate coinciding with the K-Pg boundary and the quantitative dataset displaying
two such instances during the late Miocene. The outliers displayed by the qualitative dataset are
consistent with the overall pattern of coincidence between episodes of conflict and rapid
morphological change, differing from the quantitative dataset only in magnitude. However, we
could not identify a clear explanation for the outliers present in the quantitative dataset. In
addition to the congruence between patterns of conflict and phenotypic evolution near the K-Pg,
the quantitative dataset displayed weak evidence for an overall increase in morphological rate
coinciding with the Miocene increase in phylogenomic conflict. Although the qualitative dataset
did not display a Miocene rate increase, this is likely a result of taxon sampling rather than
biological reality, since the genomic data also fail to display the Miocene boost in conflict when
the dataset is reduced to reflect the same taxa as the qualitative data matrix.
Acrogymnospermae (known colloquially as "gymnosperms") show dramatic and
coincident spikes in genomic conflict and rates of morphological evolution, with several of these
spikes occurring near major stratigraphic boundaries marked by mass extinction or, at least,
major ecological upheaval (Fig. 2). The largest spike, for both conflict and innovation, occurs
before the P-T boundary and corresponds to the divergence of Gnetales. The phylogenetic
placement of this lineage has long been contentious, with different molecular datasets (or
analytical methods) resulting in different topologies (e.g., Zhong et al. 2010). Morphologically,
Gnetales is also highly derived, which has led to various hypotheses concerning its placement,
including sister to angiosperms (Crane 1985), given that both lineages share several (presumably
independently derived) traits such as vessels in the wood, net veined leaves (in some Gnetales),
and siphonogamy, as well as flower-like sporangiate structures in Gnetales. Genomic datasets,
however, appear to be reaching a consensus that Pinaceae may indeed be the closest living
relative of this enigmatic order (e.g., Ran et al. 2018; Smith et al. 2020). Our results, which
follow this placement of Gnetales (sister to Pinaceae), suggest that the divergence of the order
involved a coordinated burst in both genomic and morphological evolution; the possible causes
are unclear but deserve further study. Although no whole-genome duplications have been
detected in the stem of Gnetales and polyploidy in gymnosperms is generally considered less
common than in other vascular plant lineages (Leebens-Mack et al. 2019), this phenomenon has
nevertheless been relatively understudied in gymnosperms. It is noteworthy that WGDs have
been detected in two of the three genera comprising Gnetales, *Welwitschia* (Li et al. 2015) and
Ephedra (Ickert-Bond 2020), with the latter including a large number of recently formed
polyploids. Two of the other major (coincident) peaks in gymnosperms correspond to the
divergence of Ginkgo and cycads (ca. 290 Ma) and the divergence of *Gnetum* and *Welwitchia*
(ca. 114 Ma). As with the Gnetales divergence, these two spikes in morphological innovation
include the acquisition of novel states spanning the plant bauplan (including, for example,
changes in stem anatomy, leaf structure, microsporangia and pollen, and megasporangia and
seeds). The monotypic genus *Welwitschia*, in particular, is characterized by highly unique and
bizarre morphology, with continuously growing strap-shaped leaves that sprawl along the desert
floor. The significance of WGD as a possible driver of phenotypic novelty in this genus deserved
further study.
The Ericales also exhibit a high amount of conflict during their early evolution. This
conflict coincides with the rapid evolution of perianth merism, androecial structure (e.g., stamen

number and anther orientation and attachment), and carpel number. Similar variation in floral
morphology is evident in some of the earliest fossil representatives of this order (e.g.,
Schönenberger and Friis 2001; Crepet et al. 2013; Martínez et al. 2016) and likely relates to early
co-evolution with insect pollinators, particularly hymenoptera (Crepet 1996; Nixon and Crepet
1993). This observed genomic conflict and innovation in floral morphology precedes the
evolution of the major families and also correspond to several life history shifts and movements
out of the tropics (Rose et al. 2018) occurring by ~79-90 MYA based on available fossil
evidence (e.g., Crepet et al. 2013) and between ~100-110 based on molecular divergence-time
estimates (Magallón et al. 2015; Rose et al. 2018). These early ecological shifts in the Cretaceous
may have involved adaptation to fire-prone habitats of the mid-latitude Northern Hemisphere,
based on taphonomic information from some of the earliest fossils (charcoalified flowers; Friis et
al. 2011; Crepet et al. 2013).
In Polypodiopsida (=monilophytes, Pryer et al. 2001; or "ferns" more colloquially), as
with gymnosperms, there are major coincident spikes in morphological rate and genomic
conflict. These correspond to the most contentious deep nodes in fern phylogeny. The
monilophytes are a heterogeneous assemblage of ancient lineages—including horsetails
(Equisetiidae), marratioid ferns (Marattiidae), whisk and ophioglossid ferns (Ophioglossidae),
and leptosporangiate ferns (Polypodiidae)—that trace back to the Devonian radiation of vascular
plants ca. 393 to 359 Ma (e.g., Kenrick and Crane 1997; Gensel et al. 2020). Prior to
phylogenetic analyses including molecular data (e.g., Pyrer et al. 2001), the lineages comprising
the monilophytes were not thought to form a clade, but analyses have continued to recover this
relationship with strong support (Pryer et al. 2001; Wickett et al. 2014; Leebens-Mack et al.
2019). However, relationships among the four major constituent lineages remain contentious
(e.g., Shen et al. 2018; Leebens-Mack et al. 2019). Here, we show that the conflict underlying
earliest divergences in monilophyte phylogeny coincides with pronounced phases of rapid
morphological innovation. The two largest bursts (dated here at ca. 290 Ma and ca. 245 Ma)
correspond, respectively, to the divergence of leptosporangiate ferns, and the split of marratioid
and whisk/ophioglossid ferns (topology following Shen et al. 2018). Various aspects of
morphological evolution across these lineages remain poorly known, including the origin(s) of
megaphyll leaves, which show divergent morphologies and range from complex and highly
diverse, to reduced, to entirely absent across the different groups of monilophytes (Harrison and
Morris 2017). Major changes in sporangial structure also characterize (and serve as the namesake
for) the largest extant monilophyte clade, the leptosporangiate ferns (Pryer et al. 2001). Overall,
these pronounced bursts of rapid morphological innovation across the early radiation of
monilophytes appear to correspond to major genomic upheavals. These events render the
resolution of relationships difficult, but also may hold insight into the evolutionary origins of
complex morphological features. It is likely that WGD played an important role in the early
evolution of this group, as WGD events have been identified in all major monilophyte lineages,
with events particularly clustered on the earlier branches of the monilophyte radiation (e.g.,
Leebens-Mack et al. 2019). In addition to other population level processes, WGD provides an
important mechanism whereby major genomic changes/conflict may both signal and underlie
major leaps in morphological novelty.
The Caryophyllales show substantial conflict at the edges preceding the emergence of the
major clades that correspond to Linnean families (Fig 2). Based on available divergence times,
these conflicts occurred between ~59-101 MYA with the most extensive conflict at ~77 MYA. In
particular, there are hotspots of conflict along the backbone of the "core Caryophyllales" and

then several within the Caryophyllaceae and Phytolaccaceae+Nyctaginaceae. Major floral
innovations have been observed along these nodes including changes in the number and
configuration of perianth and androecial parts (e.g., Brockington et al. 2009, 2013); the lineages
stemming from these nodes have been claimed to show the highest diversity in androecial
patterns among all flowering plants (Ronse De Craene 2013; Ronse De Craene and Brockington
2013). Morphological sampling beyond floral characters would likely reveal other major
instances of innovation given the many distinctive phenotypic features documented across this
ecologically diverse clade (e.g., betalain pigments, carnivory, succulence, successive cambia). In
particular, successive cambia and betalains likely evolved in high-conflict areas of the
Caryophyllales backbone (e.g., Carlquist 2010; Brockington et al., 2011; Timoneda et al. 2019),
and both traits show complicated yet intriguing patterns of presence/absence across the
phylogeny. We found that genetic innovations in the betalain pathway, major whole genome
duplications, and transitions in photosynthetic pathways transitions throughout the evolution of
the group experienced an increase in evolutionary rate coinciding with the highest peak in gene-
tree conflict (Fig. 2).
244
Discussion
We demonstrate that several instances of high phylogenomic conflict correspond to
instances of high morphological innovation. This has implications for understanding both
phylogenomic conflict and morphological evolution. First, biological processes, perhaps in
addition to error, have a large impact on the patterns of conflict, even at deep time scales and
when filtering for strongly supported signals. That biological processes can cause conflicting
signals in gene-trees is well known. However, typically, researchers attempt to overcome this
conflict through modeling or methodological developments intended to construct a species tree.
We suggest that conflict itself is important to dissect and not just accommodate in phylogenetic
reconstruction. Importantly, we do not suggest a perfect correlation between instances of
morphological novelty and conflict. There are many processes that might lead to increases in
both of these measures and so it is unlikely that there will be an exact correspondence of rapid
morphological rates and conflict at all nodes across the tree of life. Nevertheless, we suggest that
the events that led to high conflict in the past (e.g., dramatic episodes of population expansion or
contraction during rapid speciation events) had such a large impact that many aspects of these
lineages, including morphological evolution, were significantly affected.
Disentangling the sources of conflict within phylogenomic datasets is fraught with
difficulties. In the datasets examined here, while there may be instances of hybridization and, to
a lesser extent, horizontal gene transfer that are contributing to the patterns of conflict, there is no
published evidence that these phenomena are the source of biological conflict at the nodes that
correspond to morphological innovation. There has been some suggestion of ancient
hybridization early in the Caryophyllales (e.g., Morales-Briones et al. 2019); however, there are
challenges of identifiability that have yet to be overcome. Whole genome duplications (perhaps
in many cases stemming from hybridization, i.e., allopolyploidy events) may also result in
conflicts across the tree of life, especially as they may create errors as a result of misidentified
paralogy. Other systematic errors and background noise may be a source of conflict, especially
when exacerbated by rapid evolution (Smith et al. 2020; Larson et al. 2020). Furthermore,
conflicts may be the result of errors due to uncertainty resulting from rapid evolution. Finally,
incomplete lineage sorting has been recognized as a major source of gene-tree disagreement at
all depths of the tree of life (Maddison and Knowles 2006). Conflicts due to incomplete lineage

sorting will be exacerbated by changes in population processes such as changes in effective
population size from selection, bottlenecks, or other phenomena and/or rapid evolution (Pease
and Hahn 2013). While errors also, undoubtedly, have contributed to some of the conflict we
document, the vast majority of the nodes we highlight have been demonstrated to be difficult to
resolve using multiple data types and analytical techniques (Jarvis et al. 2014; Prum et al. 2015;
Wickett et al. 2015; Walker et al. 2017; Leebens-Mack et al. 2019; Larson et al. 2020).
One notable attribute of several of the clades examined (e.g., Ericales, Caryophyllales) is
that many phenotypic traits show complex patterns of presence/absence across lineages
stemming from regions of high conflict. In most cases, such a distribution of states would be
interpreted as independent gains and losses (i.e., homoplasy). However, it is possible that the
inheritance of phenotypic traits, from a polymorphic ancestral population, might in some cases
follow an ILS-like process (i.e., hemiplasy; Avise and Robinson 2008; Guerrero and Hahn
2018). If the genes underlying phenotypic traits are subject to ILS, it follows that phenotypic
traits themselves are also subject to this process. This might serve as a better explanation than
independent gains/losses for complex distributions of phenotypic states in some clades.
Hypothetically, this would imply that patterns of trait evolution during rapid "adaptive"
radiations in some cases might be partly or largely stochastic--a byproduct of the sorting of
underlying genetic variation. However, combinations of traits inherited through such a process
could subsequently prove adaptive. These possibilities are speculative but deserve further
attention, although demonstrating ILS-driven phenotypic evolution at macroevolutionary scales
would likely be a major challenge.
We suggest that population processes occurring episodically in the past exacerbated
gene-tree conflict and were correlated or causal in generating new morphological innovations. A
link between population processes and phenotypic innovation has long been theorized (Wright
1932). Although the specific relationship between microevolutionary processes and phenotypic
changes observable at a macroevolutionary scale is not well-understood, previous work has
considered how demographic events such as rapid fluctuations in population size and and genic
exchange between diverging populations might work synergistically with subsequent
environmental adaptation to drive the exploration of diverse areas of morphospace by facilitating
the emergence of both new phenotypic features and novel combinations of character states
(Simpson 1944, Marques et al. 2019, Parins-Fukuchi 2020, McGee et al. 2020). Importantly,
these processes are also likely to cause phylogenetic conflict if they persist over many
generations and coincide with lineage divergences. We therefore suggest that phylogenomic
conflict observable at deep timescales may often stem from the same population processes that
facilitated the rapid and dramatic episodes of phenotypic change underpinning the radiation and
morphological diversification of major clades. For example, the rapid speciation events that
occur during evolutionary radiations may both generate conflicting phylogenomic signal due to
population-level processes (e.g. rapid fluctuations in population size and selection) and drive the
emergence of novel morphologies as diverging lineages carve out new ecological niches. In
particular, a large population size will result in a smaller probability that two lineages will
coalesce within a particular time. This should result in high conflict between gene trees. Several
episodes of high conflict observed here (for example, in Ericales) are succeeded by periods of
low conflict. This would suggest population size reduction if ILS can safely be assumed to be the
source of conflict.
In addition to the notable coincidence of nodes with high conflict and rapid
morphological innovation, many of these nodes also appear to correspond with major geologic

events. As has long been observed, major radiations in mammals and crown birds seem closely
tied to the KPg mass extinction; our results here linking conflict to morphological innovation
hint at underlying processes that may have driven these radiations into the wake of KPg
ecological collapse. Gymnosperms and ferns show coincidence spikes near the Permian-Triassic
boundary, another widely studied mass extinction event (Benton and Twitchett 2003). But given
our limited sampling of these particular clades, this association is much more tenuous.
Unfortunately, examining these patterns in greater detail requires a reliance on accurate
divergence times, which are challenging to estimate from molecular phylogenies. A better
integration of fossil and phylogenomic data may offer a clearer understanding of how major
geologic events have influenced coordinated bursts of conflict and phenotypic innovation.
331
Conclusion. This study provides an important link between genomic and morphological
evolution at deep timescales. We suggest that episodic evolutionary and population events leave
signatures of conflict within genomes that may offer important insight on the processes
responsible not only for conflict but also for massive changes in phenotype across disparate
lineages. Moving forward, we suggest that closer examinations using more densely sampled
molecular and morphological datasets representing taxa across the tree of life will help us to
generate a deeper understanding of the micro- and macroevolutionary processes that drive major
organismal changes from the genomic to the phenotypic level.
340
341

Materials and Methods
Genomic datasets. We examine several published phylogenomic datasets including 624 loci of
Caryophyllales from Yang et al. (2018); 387 loci of Ericales from Larson et al. (2020); 1332 loci
of ferns from Shen et al. 2018; 1308 loci of gymnosperms from Ran et al. (2018); 424 loci of
mammals from Song et al. (2012), respectively; and 259 loci of Aves from Prum et al. (2015).
Species trees were all based on those presented in the original publications. Gene trees were re-
estimated using iqtree (v. 1.6.11) with support calculated using the approximate likelihood ratio
test (-alrt 1000) and with the GTR and Γ model for nucleotides (-m GTR+G) and the LG and Γ
model for amino acids (-m LG+G).
351

Morphological datasets. Large morphological datasets for major clades of flowering plants are
surprisingly few, and additionally most available datasets are relatively limited in their sampling
of either taxa and/or characters. For the Caryophyllales, we coded characters representing
genome duplications (from Yang et al. 2018), presence of betalains, betalain pathway related
duplications (from Sheehan et al. 2020), and photosynthetic pathway type (i.e., C3, C4, and
CAM). This matrix is available in the data supplement. For the gymnosperms, we used the
published data from Nixon et al. (1994) where the data are primarily scored at family level. The
taxon circumscriptions applied in Nixon et al. (1994) for their OTUs are monophyletic and
consistent with our current understanding of gymnosperm phylogeny, and thus were applicable
to the clades examined here. Thus multiple exemplars within a clade corresponding to one of
their OTUs will be identical in their states. For the Ericales, we used the published matrix
developed by Chartier et al. (2017) focused on floral characters. For mammals, we used the
O'Leary et al. (2013) published dataset that covers mammal morphology broadly as well as body
size data gathered from the panTHERIA database (Jones et al. 2009). For the ferns, we used the
morphological dataset from Pryer et al. (2001), which was developed for analysis of fern (i.e.,

monilophyte) relationships within the context of vascular plant phylogeny. Finally, for the birds,
we used the morphological matrix compiled by Livezay and Zusi (2007) as well as the principal
component scores calculated from the large dataset of linear measurements presented by Pigot et
al. (2019). Many of the morphological datasets were sampled across more taxa than the genomic
datasets-- in these cases, we pruned the morphological datasets to include only the taxa present in
the genomic data.
373
374 *Phylogenomic analyses.* Conflict analyses were conducted with the program ***bp*** (available from
375 https://github.com/FePhyFoFum/gophy) using species and gene trees. We used the options -tv to
376 print the concordance, conflict, and noninformative values back to the species tree and -scut 80
377 to ignore edges that had less than 80% for the aLRT statistic as calculated from the iqtree
378 analyses.
379
Divergence-time analyses. Although divergence-time estimates were not required for conflict
analyses, they were required for calculating rates of morphological evolution. They are also
useful for visualizing the concurrence of conflict and morphological innovation and placing these
in a geologic context. Therefore, for each dataset, regardless of the source of divergence time
estimates, both the morphological and genomic phylogenies used the same dated tree. The Aves
phylogeny used the published dates from Prum et al. (2015). The final pruned topology
contained 82 taxa matching both the morphological and genomic datasets, from an original 198.
The Caryophyllales dataset uses divergence time estimates from Smith et al. (2017) with a final
topology of 166. For the Ericales dataset, we used the divergence times and topology from
Chartier et al. (2019). For the fern dataset, we used the divergence times and topology from Shen
et al. (2018). For the gymnosperms dataset, no suitable dated phylogeny was available. Instead
we used the topology from Ran et al. (2018) and then constructed a set of calibrations based on
the available fossil data as well as divergence times available in TimeTree (Supplemental Text).
For the mammal dataset, we used the chronogram produced in the course of the original genomic
study (Chen et al. 2017).
395
Morphological analyses. We calculated rates of discrete character evolution by estimating the
number of character transitions using parsimony occurring along each phylogenetic branch and
dividing this estimate of the amount of character divergence along each branch by the amount of
time represented by the corresponding branch in the time-scaled tree. Rates calculated from
discrete-character matrices were therefore presented in units of inferred character transitions per
unit time. For quantitative traits, we calculated the average character divergences along each
branch by estimating ancestral states for each character under a Brownian motion model of
evolution and then representing the amount of change along each edge as the absolute value of
the difference between the trait value inferred for the ancestor and that displayed by the
descendant. Like with the qualitative analyses, we then transformed these raw divergences to
rates by calculating their ratio to the corresponding amount of elapsed time for the corresponding
edge in the timetree.
408
409 *Selection of species tree topology*: We used the same topology for both the morphological and
410 molecular analyses. Conflict and morphological rates were mapped onto the species tree resolved
411 in the study from which the genomic data were sourced.
412

413 **Data availability**
414 Gene trees and dated species trees for each of the datasets examined are available from Figshare.

459 **Figures**

463 *Figure 1.* Co-occurring patterns in gene-tree conflict and morphological rates of evolution across

464 bird and mammal datasets.

472
473 *Figure 2.* Co-occurring patterns in gene-tree conflict and morphological rates of evolution across
474 major plant clades.

References

Avise, J. C., T. J. Robinson. 2008. Hemiplasy: A new term in the lexicon of phylogenetics. *Systematic Biology* 57: 503-507.

Beck, C. B. 1988. Origin and Evolution of Gymnosperms. Cambridge University Press, New York, New York.

Benton, M. J., R. J. Twitchett. 2003. How to kill (almost) all life: the end-Permian extinction event. *Trends in Ecology and Evolution* 7: 358-365.

Brockington, S. F., R. Alexandre, J. Ramdial, M. J. Moore, S. Crawley, A. Dhingra, K. Hilu, D. E. Soltis, P. S. Soltis. 2009. Phylogeny of Caryophyllales sensu lato: Revisiting hypotheses on pollination biology and perianth differentiation in the core Caryophyllales. *International Journal of Plant Sciences* 170: 627-643.

Brockington, S. F., R. H. Walker, B. J. Glover, P. S. Soltis, D. E. Soltis. 2011. Complex pigment evolution in the Caryophyllales. *New Phytologist* 190: 854-864.

Brockington, S. F., P. Dos Santos, B. Glover, L. P. Ronse De Craene. 2013. Androecial evolution in Caryophyllales in light of a paraphyletic Molluginaceae. *American Journal of Botany* 100: 1757-1778.

Calder, W. A. 1983. Ecological scaling: mammals and birds. Annual Review of Ecology and Systematics, 14: 213-230.

Carlquist, S. 2010. Caryophyllales: a key group for understanding wood anatomy character states and their evolution. *Botanical Journal of the Linnean Society* 164: 342-393.

Crane, P. R. 1985. Phylogenetic analysis of seed plants and the origin of angiosperms. *Annals of the Missouri Botanical Garden* 72: 716-793.

Crepet, W. L. 1996. Timing in the evolution of derived floral characters: Upper Cretaceous (Turonian) taxa with tricolpate and tricolpate-derived pollen. *Review of Paleobotany and Palynology* 90: 339-359.

Crepet, W. L., K. C. Nixon, C. P. Daghlian. 2013. Fossil Ericales from the Upper Cretaceous of New Jersey. *International Journal of Plant Sciences* 174: 572-584.

Drew, B. T., B. R. Ruhfel, S. A. Smith, M. J. Moore, B. G. Briggs, M. A. Gitzendanner, P. S. Soltis, D. E. Soltis. 2014. Another look at the root of the angiosperms reveals a familiar tale. *Systematic Biology* 63: 368-382.

Jones, K. E., J. Bielby, M. Cardillo, S. A. Fritz, J. O'Dell, C. D. L. Orme, K. Safi, W. Sechrest, E. H. Boakes, C. Carbone, C. Connolly, M. J. Cutts, J. K. Foster, R. Grenyer, M. Habib, C. A. Plaster, S. A. Price, E. A. Rigby, J. Rist, A. Teacher, O. R. P. Bininda-Emonds, J. L. Gittleman, G. M. Mace, A. Purvis. 2009. PanTHERIA: a species-level database of life history, ecology, and

geography of extant and recently extinct mammals: Ecological Archives E090-184. *Ecology* 90:
2648-2648.

Friis, E. M., P. R. Crane, K. R. Pedersen. 2011. Early flowers and angiosperm evolution.
Cambridge University Press: NY, NY, USA.

Gensel, P. G., I. Glasspool, R. A. Gastaldo, M. Libertin, J. Kvacek. 2020. Back to the
beginnings: The Silurian-Devonian as a time of major innovations in plants and their
communities, pp. 367 to 398 in E. Martinetto et al. (eds.), *Nature through Time*, Springer
Textbooks in Earth Sciences, Geography and Environment, https://doi.org/10.1007/978-3-030-
35058-1_15.

Guerrero, R. F., M. W. Hahn. 2018. Quantifying the risk of hemiplasy in phylogenetic inference.
Proceedings of the National Academy of Sciences, U.S.A. 50: 12787-12792.

Harrison, C. J., J. L. Morris. 2018. The origin and early evolution of vascular plant shoots and
leaves. *Philosophical Transactions of the Royal Society B* 373: 20160496.

Ickert-Bond, S. M., A. Sousa, Y. Min, I. Loera, J. Metzgar, J. Pellicer, O. Hidalgo, I. J. Leitch.
2020. Polyploidy in gymnosperms-Insights into the genomic and evolutionary consequences of
polyploidy in Ephedra. *Molecular Phylogenetics and Evolution* 147: 106786.

Jarvis, E. D., S. Mirarab, A. J. Aberer, B. Li, P. Houde, C. Li, S. Y. Ho, B. C. Faircloth, B.
Nabholz, J. T. Howard, A. Suh. 2014. Whole-genome analyses resolve early branches in the tree
of life of modern birds. *Science* 346:1320-1331.

Kenrick, P., P. R. Crane. 1997. Origin and early diversification of land plants. Smithsonian
Institution Press, Washington D.C.

Larson, D. A., J. F. Walker, O. M. Vargas, S. A. Smith. 2020. A consensus phylogenomic
approach highlights paleopolyploid and rapid radiation in the history of Ericales. *American
Journal of Botany* 107: 773-789.

Leebens-Mack, J. H., M. S. Barker, E. J. Carpenter, *et al.* 2019. One thousand plant
transcriptomes and the phylogenomics of green plants. *Nature* 574: 679–685.

Li, Z., A. E. Baniaga, E. B. Sessa, M. Scascitelli, S. W. Graham, L. H. Rieseberg, M. S. Barker.
2015. Early genome duplications in conifers and other seed plants. *Science advances* 1:
e1501084.

Livezey, B. C., R. L. Zusi. 2007. Higher-order phylogeny of modern birds (Theropoda, Aves:
Neornithes) based on comparative anatomy. II. Analysis and discussion. *Zoological Journal of
the Linnean Society* 149: 1-95.

Magallón, S., S. Gómez-Acevedo, L. L. Sánchez-Reyes, T. Hernández-Hernández. 2015. A
metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity. *New
Phytologist* 207: 437-453.

Marques, D. A., J. I. Meier, O. Seehausen. 2019. A combinatorial view on speciation and adaptive radiation. *Trends in Ecology and Evolution* 34: 531-544.

Martínez, C., T. Y. S. Choo, D. Allevato, K. C. Nixon, W. L. Crepet, R. S. Harbert, C. P. Daghlian. 2016. Rariglanda jerseyensis, a new ericalean fossil flower from the Late Cretaceous of New Jersey. *Botany* 94: 747-758.

McGee, M. D., S. R. Borstein, J. I. Meier, D. A. Marques, S. Mwaiko, A. Taabu, M. A. Kishe, B. O'Meara, R. Bruggmann, L. Excoffier, O. Seehausen. 2020. The ecological and genomic basis of explosive adaptive radiation. *Nature* 586: 75-79.

Morales-Briones, D.F., G. Kadereit, D. T. Tefarikis, M. J. Moore, S. A. Smith, S. F. Brockington, A. Timoneda, W. C. Yim, J. C. Cushman, Y. Yang. 2019. Disentangling Sources of Gene Tree Discordance in Phylotranscriptomic Datasets: A Case Study from Amaranthaceae sl. *Systematic Biology*, https://doi.org/10.1093/sysbio/syaa066.

Morgan, C. C., P. G. Foster, A. E. Webb, D. Pisani, J. O. McInerney, M. J. O'Connell. 2013. Heterogeneous models place the root of the placental mammal phylogeny. *Molecular Biology and Evolution* 30: 2145-2156.

Nixon, K. C., W. L. Crepet. 1993. Late Cretaceous fossil flowers of ericalean affinity. *American Journal of Botany* 80: 616-623.

Nixon, K.C., W. L. Crepet, D. Stevenson, E. M. Friis. 1994. A reevaluation of seed plant phylogeny. *Annals of the Missouri Botanical Garden* 81: 484-533.

Nosenko, T., F. Schreiber, M. Adamski, M. Eitel, J. Hammel, M. Maldonado, W. E. Müller, M. Nickel, B. Schierwater, J. Vacelet. 2013. Deep metazoan phylogeny: when different genes tell different stories. *Molecular Phylogenetics and Evolution* 67: 223-233.

O'Leary, M. A., J. I. Bloch, J. J. Flynn, T. J. Gaudin, A. Giallombardo, N. P. Giannini, S. L. Goldberg, B. P. Kraatz, Z. X. Luo, J. Meng, X. Ni. 2013. The placental mammal ancestor and the post–K-Pg radiation of placentals. *Science* 339: 662-667.

Parins-Fukuchi, C. 2020. Mosaic evolution, preadaptation, and the evolution of evolvability in apes. *Evolution* 74: 297-310.

Pease, J. B., M. W. Hahn. 2013. More accurate phylogenies inferred from low-recombination regions in the presence of incomplete lineage sorting. *Evolution* 67: 2376-2384.

Pigot, A. L., C. Sheard, E. T. Miller, T. P. Bregman, B. G. Freeman, U. Roll, N. Seddon, C. H. Trisos, B. C. Weeks, J. A. Tobias. 2020. Macroevolutionary convergence connects morphological form to ecological function in birds. *Nature Ecology and Evolution* 4: 230-239.

629   Prum, R. O., J. S. Berv, A. Dornburg, D. J. Field, J. P. Townsend, E. M. Lemmon, A. R.
630   Lemmon. 2015. A comprehensive phylogeny of birds (Aves) using targeted next-generation
631   DNA sequencing. *Nature* 526: 569-573.
632
633   Pryer, K. M., H. Schneider, A. R. Smith, R. Cranfill, P. G. Wolfe, J. S. Hunt, S. D. Sipes. 2001.
634   Horsetails and ferns are a monophyletic group and the closest relatives to seed plants. *Nature*
635   409: 618-622.
636
637   Ran, J.-H., T.-T. Shen, M.-M. Wang, X.-Q. Wang. 2018. Phylogenomics resolves the deep
638   phylogeny of seed plants and indicates partial convergent or homoplastic evolution between
639   Gnetales and angiosperms. *Proceedings of the Royal Society B* 285: 20181012.
640
641   Romiguier, J., V. Ranwez, F. Delsuc, N. Galtier, E. J. Douzery. 2013. Less is more in
642   mammalian phylogenomics: AT-rich genes minimize tree conflicts and unravel the root of
643   placental mammals. *Molecular Biology and Evolution* 30: 2134-2144.
644
645   Ronse De Craene, L. P. 2013. Reevaluation of the perianth and androecium in Caryophyllales:
646   implications for flower evolution. *Plant Systematics and Evolution* 299: 1599-1636.
647
648   Ronse De Craene, L. P., S. F. Brockington. 2013. Origin and evolution of petals in angiosperms.
649   *Plant Ecology and Evolution* 146: 5-25.
650
651   Rose, J.P., T. J. Kleist, S. D. Löfstrand, B. T. Drew, J. Schönenberger, K. J. Sytsma. 2018.
652   Phylogeny, historical biogeography, and diversification of angiosperm order Ericales suggest
653   ancient Neotropical and East Asian connections. *Molecular Phylogenetics and Evolution* 122:
654   59-79.
655
656   Schönenberger, J., E. M. Friis. 2001. Fossil flowers of Ericalean affinity from the Late
657   Cretaceous of southern Sweden. *American Journal of Botany* 88: 467-480.
658
659   Shen, H., D. Jin, J.-P. Shu, X.-L. Zhou, M. Lei, R. Wei, H. Shang, H.-J. Wei, R. Zhang, L. Liu,
660   Y.-F. Gu, X.-C. Zhang, Y.-H. Yan. 2018. Large-scale phylogenomic analysis resolves a
661   backbone phylogeny in ferns. *GigaScience* 7: gix116.
662
663   Simpson, G. G.. 1944. Tempo and mode in evolution. Columbia University Press. New York,
664   NY.
665
666   Smith, S. A., M. J. Moore, J. W. Brown, Y. Yang. 2015. Analysis of phylogenomic datasets
667   reveals conflict, concordance, and gene duplications with examples from animals and plants.
668   *BMC Evolutionary Biology* 15: 150.
669
670   Smith, S.A., J. M. Brown, Y. Yang, R. Bruenn, C. P. Drummond, S. F. Brockington, J. F.
671   Walker, J.F., N. Last, N. A. Douglas, M. J. Moore. 2018. Disparity, diversity, and duplications in
672   the Caryophyllales. *New Phytologist* 217: 836-854.
673

674 Smith, S. A., N. Walker-Hale, J. F. Walker, J. W. Brown. 2020. Phylogenetic conflicts,
675 combinability, and deep phylogenomics in plants. *Systematic Biology* 69: 579-592.
676

677 Song, S., L. Liu, S. V. Edwards, S. Wu. 2012. Resolving conflict in eutherian mammal
678 phylogeny using phylogenomics and the multispecies coalescent model. *Proceedings of the*
679 *National Academy of Sciences U. S. A.* 109: 14942-14947.
680

681 Stearns, S. C. 1983. The influence of size and phylogeny on patterns of covariation among life-
682 history traits in the mammals. *Oikos*: 173-187.
683

684 Stewart, W. N., G. W. Rothwell. 1993. Paleobotany and the evolution of plants. Cambridge
685 University Press, New York, NY.
686

687 Teeling, E. C., S. B. Hedges. 2013. Making the impossible possible: rooting the tree of placental
688 mammals. *Molecular Biology and Evolution* 30: 1999-2000.
689

690 Timoneda, A., T. Feng, H. Sheehan, N. Walker-Hale, B. Pucker, S. Lopez-Nieves, R. Guo, S.
691 Brockington. 2019. The evolution of betalain biosynthesis in Caryophyllales. *New Phytologist*
692 224: 71-85.
693

694 Wickett, N. J., S. Mirarab, N. Nguyen, T. Warnow, E. Carpenter, N. Matasci, S. Ayyampalayam,
695 M. S. Barker, J. G. Burleigh, M. A. Gitzendanner, *et al*. 2014. Phylotranscriptomic analysis of
696 the origin and early diversification of land plants. *Proceedings of the National Academy of*
697 *Sciences, U.S.A* 111: E4859-E4868.
698

699 Wright, S. 1932. The roles of mutation, inbreeding, crossbreeding and selection in evolution.
700 Proceedings of the 6th International Congress of Genetics: 356-366.
701

702 Yang, Y., M. J. Moore, S. F. Brockington, J. Mikenas, J. Olivieri, J. F. Walker, S. A. Smith.
703 2018. Improved transcriptome sampling pinpoints 26 ancient and more recent polyploidy events
704 in Caryophyllales, including two allopolyploidy events. *New Phytologist* 217: 855-870.
705

706 Zhong, B., T. Yonezawa, Y. Zhong, M. Hasegawa. 2010. The position of Gnetales among seed
707 plants: overcoming pitfalls of chloroplast phylogenomics. *Molecular Biology and Evolution* 27:
708 2855-2863.
709

711