Complex population structure of the Atlantic puffin revealed by whole genome analyses

Oliver Kersten[1]*, Bastiaan Star[1], Deborah M. Leigh[2], Tycho Anker-Nilssen[3], Hallvard Strøm[4],

Jóhannis Danielsen[5], Sébastien Descamps[4], Kjell E. Erikstad[4,6], Michelle G. Fitzsimmons[7],

Jérôme Fort[8], Erpur S. Hansen[9], Mike P. Harris[10], Martin Irestedt[11], Oddmund Kleven[3], Mark L.

Mallory[12], Kjetill S. Jakobsen[1], Sanne Boessenkool[1]*

[1] Centre for Ecological and Evolutionary Synthesis (CEES), Department of Biosciences, University of Oslo, Blindernveien 31, 0371

Oslo, Norway

[2]WSL Swiss Federal Research Institute, Zürcherstrasse 111, 8903 Birmensdorf, Switzerland

[3] Norwegian Institute for Nature Research (NINA), Høgskoleringen 9, NO-7034 Trondheim, Norway

[4] Norwegian Polar Institute, Fram Centre, Postbox 6606, Langnes, 9296 Tromsø, Norway

[5]Faroe Marine Research Institute (FAMRI), Nóatún 1, FO-100 Tórshavn, Faroe Islands

[6]Centre for Biodiversity Dynamics (CBD), Norwegian University of Science and Technology (NTNU), Trondheim, Norway

[7]Environment and Climate Change Canada, Newfoundland and Labrador, Canada

[8]Littoral, Environment et Sociétés (LIENSs), UMR 7266 CNRS – La Rochelle Université, 17000 La Rochelle, France

[9]South Iceland Nature Research Centre, Ægisgata 2, 900 Vestmannaeyjar, Iceland

[10]UK Centre for Ecology & Hydrology, Penicuik, Midlothian, EH26 0QB, UK

[11]Department for Bioinformatics and Genetics, Swedish Museum of Natural History, Box 50007, 104 05 Stockholm, Sweden

[12]Department of Biology, Acadia University, Wolfville, Nova Scotia, Canada, B4P 2R6

*Email: oliver.kersten@ibv.uio.no, sanne.boessenkool@ibv.uio.no

Abstract

The factors underlying gene flow and genomic population structure in vagile seabirds are

notoriously difficult to understand due to their complex ecology with diverse dispersal barriers and

extensive periods at sea. Yet, such understanding is vital for conservation management of

seabirds that are globally declining at alarming rates. Here, we elucidate the population structure

of the Atlantic puffin (*Fratercula arctica*) by assembling its reference genome and analyzing

genome-wide resequencing data of 72 individuals from 12 colonies. We identify four large,

genetically distinct clusters, observe isolation-by-distance between colonies within these clusters,

and obtain evidence for a secondary contact zone. These observations disagree with the current

taxonomy, and show that a complex set of contemporary biotic factors impede gene flow over

different spatial scales. Our results highlight the power of whole genome data to reveal

unexpected population structure in vagile marine seabirds and its value for seabird taxonomy,

evolution and conservation.

Main

Seabirds are important ecosystem indicators and drivers[e.g. 1–3], and have long had an integral place in human culture and economy[4–6]. Nevertheless, global seabird numbers have deteriorated by an alarming 70% since halfway of the 20th century[e.g. 7,8]. These declines pose a serious threat to marine ecosystems, human society, and culture[e.g. 7,9,10], highlighting the importance of seabird conservation management. Within such management the identification of distinct population units is a fundamental first step towards optimized conservation[11–13]. Defining such units is, however, difficult for many seabirds because their complex ecology promotes dispersal barriers due to e.g. oceanography, foraging behavior, at-sea range, and phenology[14]. Detailed genomic data including thousands of loci provide new possibilities to assess levels of connectivity and gene flow between distinct breeding populations and, thus, help identify relevant conservation units for seabirds[14,15]. Indeed, a few recent publications using reduced genomic representation approaches (e.g. RAD-seq) report fine-scale structure over various spatial scales[16–20]. These studies highlight the great potential of genomic data to disentangle barriers to gene flow that would otherwise remain undetected, but remain nonetheless limited due to incomplete sampling of the genome[21].

The Atlantic puffin (*Fratercula arctica*, Linnaeus, 1789, hereafter "puffin") is an iconic seabird species, prevalent in popular culture[e.g. 22], important for tourism[e.g. 23,24] and inherently valuable for the marine ecosystem[e.g. 1]. Puffins were historically widely harvested for their meat and down[6,25,26] and exploitation remains an important cultural tradition in Iceland and the Faroe Islands[6,23]. Its breeding range stretches from the Arctic coast and islands of European Russia, Norway, Greenland, and Canada, southward to France and the USA[27] (Figure 1a). Puffins have been designated as "vulnerable" to extinction globally and listed as "endangered" in Europe[28]. Notably, the once world's largest puffin colony (Røst, Norway) has experienced complete fledging failure during nine of the last 13 seasons and has lost nearly 80% of its breeding pairs during the last 40 years[28–30]. Similarly, Icelandic and Faroese puffins have experienced low productivity and

negative population growth since 2003[31]. Puffins have been broadly classified into three

taxonomic groups along a latitudinal gradient based on size, with the *smallest* puffins found

around France, Britain, Ireland and southern Norway (*F. a. grabae*), *intermediate* sized puffins

around Norway, Iceland and Canada (*F. a. arctica*) and the *largest* puffins found in the High Arctic,

e.g. Spitsbergen, Greenland and northeastern Canada[32] (*F. a. naumanni*)[33] (Figure 1a).

Nevertheless, this broad classification into three subspecies has been controversial[27,34,35] and the

population structure of puffins remains unresolved at all spatial scales[34]. This knowledge gap

obstructs efforts towards an assessment of dispersal barriers, limits our understanding of cause-

and-effect dynamics between population trends, ecology and the marine ecosystem, and hinders

the development of adapted large-scale conservation actions.

Here, we present the first whole genome analysis of structure, gene flow and taxonomy of

a pelagic, North Atlantic seabird. We generated a *de novo* draft assembly for the puffin and

resequenced 72 individuals across 12 colonies representing the majority of the species' breeding

range (Figure 1a). Our work provides compelling evidence that a complex interplay of ecological

factors contributes to the range-wide genomic population structure of this vagile seabird.

Results

Genome assembly and population sequencing

Based on synteny with the razorbill (*Alca torda*), a total of 13,328 puffin scaffolds were placed into

26 pseudo-chromosomes, leaving 17.06 Mbp (1.4%) unplaced and yielding an assembly of 1.294

Gbp (Table S1, S2). This assembly contains 4,522 of the 4,915 genes (92.0%) of complete protein

coding sequences from the avian set of the OrthoDB v9 database (Table S1). We also assembled

the puffin mitogenome (length of 17,084 bp) with a similar arrangement of genomic elements as

other members within the bird family Alcidae[36,37] (Figure S1, Table S3). For the 72 resequenced

specimens, we analyzed a total of 5.77 billion paired reads, obtaining an average fold-coverage

of 7X (range 3.0 - 10) for the nuclear genome and 591X (5.3 - 1800) for the mitochondrial genome

per specimen (Figure 1a, Table S4). One individual (IOM001) was removed from the dataset due

to a substantially lower number of mapped reads (endogeny) relative to all other samples (Table

S4) resulting in a large proportion of missing sites (Figure S2). Additional filtering produced a final

genotype likelihood dataset of 1,093,765 polymorphic nuclear sites and 192 mitochondrial single-

nucleotide-polymorphisms (SNPs) in 71 birds (36 males and 35 females).

Genomic population structure

Genomic variation across 71 puffin mitogenomes defines 66 polymorphic haplotypes that indicate

a recent global population expansion and show no significant population structure (Figure 1b,

Figure S3, Table S5, S6). In contrast, we inferred four main population clusters using Principal

Component Analysis (PCA) of the nuclear whole-genome dataset (Figure 1c). Puffins from

Spitsbergen are most distinct, while puffins from Bjørnøya are located between Spitsbergen and

a larger, central cluster consisting of populations from Norway, Iceland and the Faroe Islands

(Figure 1c, Figure S4a). Puffins from Canada form their own distinct cluster, as do those from the

Isle of May, southeast Scotland (Figure 1c, Figure S4b). Hierarchical PCA analyses of the cluster

comprising the mainland Norwegian, Icelandic and Faroese colonies reveal further fine-scale

structure separating Norwegian (Hornøya and Røst) and Faroese/Icelandic colonies (Figure S4c).

Model-based clustering (ngsAdmix) agrees with the results from the PCA (Figure 1b). The optimal

model fit for the entire dataset is either K = 2 or K = 4 (Figure S5a), as determined by the method

of Evanno et al.[38]. At K = 2, ngsAdmix separates Spitsbergen from the other colonies, with

Bjørnøya being admixed (following separation along PCA 1), whereas at K = 4, ngsAdmix reflects

the structure of three additional distinct clusters representing Spitsbergen, Canada, the Isle of

May, and a central group with more shared ancestry. The shared ancestry of the central group

remains present in hierarchical admixture analyses excluding Spitsbergen and Bjørnøya

individuals (Figure S5b, S6). We find no fixed alleles and pairwise F_{ST} values between colonies

and genomic clusters are low (<0.01) (Table S6), apart from any comparisons involving the

Spitsbergen population, which show substantially higher F_{ST} values (0.03-0.08).

Phylogenetic reconstructions using individual-based Neighbor-Joining (NJ) and maximum

likelihood (ML) methods (Figure 2a, Figure S7), as well as population-based analyses in Treemix

(Figure 2b), support the distinctiveness of the Spitsbergen, Canada and the Isle of May puffins

with each group forming monophyletic clades with 100% bootstrap support. In contrast, Bjørnøya

forms a paraphyletic clade between Spitsbergen and northern Norway (Figure 2a). The population

clusters identified by the PCA and ngsAdmix at smaller spatial scales are also identified in the

topologies of the NJ and ML trees, sorting individuals predominantly according to geographical

location, although with low bootstrap support (> 80) due to large inter-individual variability (Figure

2a, S6). Allowing a single migration edge in the Treemix phylogeny identifies recent gene flow

from Spitsbergen to Bjørnøya (likelihood = 792.106; Figure S8, S9a). Adding additional migration

edges to the population-based ML tree does not improve the model fit and such edges are

therefore not further interpreted (Figure S8-S10).

Genetic diversity, heterozygosity and inbreeding

Tajima's D does not significantly deviate from neutral expectation per colony (Table S5).

Nucleotide diversity (π) of puffins is significantly different between colonies, with the Spitsbergen

population having significantly lower nucleotide diversity than the global median (Wilcoxon Rank

Sum test, U = 4824, n_{SPI} = 25, n_{Global} = 300, p < 0.05, Table S5). Colonies also differ significantly

in levels of heterozygosity (Kruskal-Wallis Test, p < 0.001; Figure 3a) and inbreeding (Kruskal-

Wallis Test, p < 0.001, Figure 3b), whereby individual inbreeding (F_{RoH}) was approximated based

on Runs of Homozygosity (RoH)[39]. Again, the Spitsbergen colony has significantly lower levels of

heterozygosity (0.00220 - 0.00223) and significantly higher levels of F_{RoH} values (0.161 - 0.172),

compared to the Faroese and Icelandic colonies (Dunn test with Holm correction, p < 0.05). The

Faroese and Icelandic colonies contain the highest levels of heterozygosity and lowest F_{RoH}

values (Figure 3a, 3b, S11) overall. The remaining colonies display intermediate levels (Figure

3a, 3b), although heterozygosity is significantly lower (Figure 3a, S11) and inbreeding is

significantly higher (Figure 3b, S11) on Gull Island and Bjørnøya compared to the Icelandic and

Faroese colonies (Dunn test with Holm correction, p < 0.05). Moreover, Spitsbergen harbors the

most (an average of 718 per individual) and longest RoHs with eight being ≥ 2.3 Mbp long (4.21

± 3.02 % of respective chromosome), whereas none of the RoHs in the remaining colonies were

> 2.15 Mbp long (Figure 3c). The only exception was a 9.65 Mbp long RoH on pseudo-

chromosome 7 (18% of chromosome length) in an Isle of May individual (Figure 3c).

Patterns of gene flow and Isolation-By-Distance (IBD)

We investigated patterns of gene flow and IBD between the colonies using two-dimensional

estimated effective migration surface (EEMS) analyses. Levels of gene flow between the Icelandic

and Faroese colonies and within the Canadian group is high (3-10x higher than the global

average), while intermediate between the Norwegian mainland colonies (around the global

average). In contrast, the Spitsbergen colony is split from the remaining colonies by migration

rates up to 100x lower than the global average (Figure 4a, S12), while additional regions of low

gene flow (2-3x lower than the global average) separate the Isle of May, Canadian, and Bjørnøya

colonies from the rest (Figure 4a, Figure S12). Geographic distance between all puffin colonies

is a poor predictor of pairwise genetic distance, driven by the high Slatkin's linearized F_{ST} values

of the Spitsbergen colony (Figure S13). Nevertheless, geographic distance among a subset of

puffin colonies is significantly correlated with genetic distance using Mantel tests (Figure 4b, S13).

Specifically, by progressively removing the more distant colonies (Spitsbergen, Bjørnøya, Isle of

May, Canada), the fit of a linear IBD model is significantly improved and the proportion of genetic

dissimilarity explained by geographic distance is more than doubled (No SPI = 38.16 %,

SPI/BJO/IOM removed = 79.77 %) (Figure 4b, S13). The isolated colonies are characterized by

high Slatkin's linearized F_{ST} values at relatively small geographic distances (Figure S13).

Admixture on Bjørnøya

We specifically tested for patterns of admixture in Bjørnøya. Significantly negative $f3$ statistics (Z

score < -3) are found for all unique combinations of the phylogeny (Spitsbergen, *X*; Bjørnøya)

(Table S7), indicating an admixed colony on Bjørnøya caused by gene flow between Spitsbergen

and the remaining colonies. Similarly, significantly positive D-statistics (Z score > 3) caused by

an excess of ABBA sites reveal excessive allele sharing between Spitsbergen and Bjørnøya

(Figure S14a). The close association and gene flow from Spitsbergen to Bjørnøya is further

confirmed by D-statistics not being significantly different from 0 for the (((Bjørnøya, Spitsbergen),

H3), Razorbill) topology (Figure S14b).

Discussion

Barriers to gene flow leading to population structure are notoriously difficult to identify and remain largely unknown for most seabirds[14,40]. Using whole genome analyses, we here provide novel insights into the genetic structure of the Atlantic puffin. First, we identify four main puffin population clusters consisting of (1) Spitsbergen (High Arctic), (2) Canada, (3) Isle of May, and (4) multiple colonies in Iceland, the Faroe Islands and Norway. Second, we find that *within* such clusters, genetic differentiation is driven by isolation-by-distance (IBD). Finally, we find evidence for secondary contact between two clusters. These observations show that a complex set of drivers impacts gene flow over different spatial scales (100s -1000s of km) between these clusters and the colonies within. In particular, the interplay between overwintering grounds, philopatry, natal dispersal, geographic distance, and potentially ocean regimes appears to explain the genomic differentiation between puffin colonies.

Mature puffins rarely, if ever, change their colonies, resulting in very high colony fidelity once they start breeding[27]. Immatures, however, have been observed to visit other nearby colonies during the summer and may breed in non-natal colonies[27,41]. Nevertheless, data on natal philopatry remain scarce, but existing evidence shows rates vary greatly (38-92%) between colonies[27,41]. If either breeding or natal philopatry alone drive the puffin population structure, each colony should constitute its own distinct genomic entity and significant relationships between geographic and genomic distance across the puffin's entire breeding range would be observed. Yet, philopatry alone cannot explain the presence of the four large-scale population clusters we observe here. Additional factors must therefore promote the distinctiveness of the four clusters. For instance, the Isle of May birds have a largely separate overwintering distribution mainly in the North Sea (Figure S15)[27,35,42]. Such potential geographical separation during the winter season might limit the likelihood of immatures intermixing between the Isle of May and other colonies. Similarly, distinct overwintering distributions have been found to lead to increased genetic diversification in other philopatric seabird species[14,40,43], such as the thick-billed murre (*Uria*

lomvia)[20] and black-browed albatross (*Thalassarche melanophris*)[44]. The presence of a Canadian

cluster can also be largely explained by their winter distribution around Newfoundland[42,45]. There

is, however, some fragmentary overlap in the overwintering distribution of the Canadian and

Icelandic colonies off southwestern Greenland[42,45], suggesting that barriers to migration of

immatures and gene flow in the western Atlantic may be further enforced by the large geographic

distance. In contrast, the winter distribution from the colonies in Iceland, Norway and the Faroe

Islands overlaps off the coast of southern Greenland (Figure S15)[42]. This shared overwintering

area, combined with the tendency to return to the natal colony and immature visits of nearby (up

to 100s km) colonies during the summer, appears to drive a pattern of IBD among colonies (Figure

3b). Indeed, IBD has previously been recognized as an important driver of genomic structure in

seabirds, for instance in the little auk (*Alle alle*)[46] and band-rumped storm-petrel (*Oceanodroma*

castro)[47]. While these illustrated mechanisms provide reasonable explanations for the observed

dispersal barriers and population structure based on our current knowledge, validation requires

additional evidence, specifically on the winter distribution of immature puffins and natal emigration

rates across colonies covering the entirety of the puffin's breeding range.

High Arctic puffins from Spitsbergen are genetically the most divergent group within our

dataset. They are also characterized by significantly lower levels of genetic diversity, greater

inbreeding coefficients, and longer and more abundant RoHs compared to other colonies. These

observations may either result from a historical bottleneck followed by isolation (e.g. founder

effect), local adaptation to their extreme environment or generally lower effective population sizes.

Population abundance estimates of less than 10,000 breeding pairs on Spitsbergen compared to

500,000 in the West Atlantic, two million on Iceland and more than two million in the boreal East

Atlantic potentially indicate a lower effective population size[27]. The High Arctic puffins exclusively

inhabit harsh, cold-current environments, also in the winter, during which they likely stay in an

area bounded by the East Greenland ice edge, a latitudinal border at 70° N and the front between

the Barents and Greenland Sea (Figure S15). They are also substantially larger than birds from

lower latitudes[27], following Bergmann's[48] or James's[49] rule, as has been observed in other

seabirds[e.g. 50,51]. This matches the clinal size variation of puffins that closely tracks sea

temperatures in their breeding areas[52]. Despite these distinctions, we find that the relatively small

population of puffins on Bjørnøya (< 1,000 pairs[27]), midway between Spitsbergen and mainland

Norway, represents an area of secondary contact between the puffins from the High Arctic and

other puffin colonies. Based on D- and the f3-statistics, the most likely southern sources are

Iceland, the Faroe Islands, Norway, or a combination thereof. Thus, the barriers to gene flow that

keep the Spitsbergen colonies distinct do not prevent formation of a hybrid colony where

individuals from the High Arctic and the cluster composed of mainland Norwegian, Icelandic and

Faroese colonies meet.

The distinct population structure in the nuclear data is not observed in the mitochondrial

genomes, which reveal an abundance of rare alleles and lack of significant population

differentiation. The mitogenomic variation suggests that puffins experienced a recent population

expansion, possibly out of a refugium after the Last Glacial Maximum. Indeed, it has been shown

that mitogenomic variation in seabirds is dominated by historical factors rather than representing

contemporary gene flow[40], and a lack of mitogenomic population structure has been observed in

many marine birds with high philopatry[e.g. 46,53,54]. In contrast to the mitogenomes, the structure in

the nuclear data therefore likely originated after the last glacial period and reflects the influence

of relatively recent barriers to gene flow in a context of historical demography[14,40]. Such results

are relevant for understanding the "seabird paradox", which contrasts the life-history traits of high

philopatry and restricted dispersal in otherwise highly mobile species[55].

Our results have major implications for the conservation management of the Atlantic

puffin. The genetic structure we identify in puffins disagrees with the suggestion of three

subspecies (*F. a. naumanni, F. a. arctica, F. a. grabae*)[33]. Although the genetically distinct

Spitsbergen cluster coincides with the classification of morphologically large puffins in the High

Arctic (*F. a. naumanni*)[27], we observe gene flow from Spitsbergen into Bjørnøya, which has been

considered *F. a. arctica*[27]. Furthermore, the geographic divide between *F. a. grabae* and *F. a.*

arctica lies farther south than previously thought, with the Faroese puffins being genetically closer

to *F. a. arctica* than to *F. a. grabae*. Nonetheless, *F. a. grabae* is currently represented by a single

colony (Isle of May) in our study and the geographical extent of this genomic cluster needs to be

refined by additional sampling, particularly in the western UK, Ireland, and France. Finally, puffins

from the Western Atlantic region (e.g. colonies in Canada) form their own distinct genetic cluster

that is not recognized within the current classification. Our results do not only warrant a revision

of Salomonsen's taxonomic classification of three subspecies[33], but also highlight the need to

acknowledge the four identified clusters as distinct units within the conservation management of

puffins[11–13]. Although puffin colonies *within* clusters are not genetically distinct entities, ecological

independence illustrated by contrasting population dynamics across relatively small spatial scales

(e.g. western Norway[30]) suggests that higher resolution local management units based on

ecological differences should be considered. Nonetheless, the observation that the genetically

distinct clusters are at the outer edges of the puffin's distribution, indicates that the management

and conservation of these distinct clusters must be a priority. Finally, our sampling does not cover

several outskirts of the puffin's distribution, such as the US, northern Canada, Greenland, Ireland,

western UK, France or Russia, and we may therefore still underestimate the true biological and

genetic complexity of this species.

In conclusion, our study shows that a complex interplay of barriers to gene flow drives a

previously unrecognized population diversification in the iconic Atlantic puffin. So far, much of

seabird population genetics research has been based on mitochondrial and microsatellite

data[14,40], which have limited power to characterize contemporary factors that determine

population structure and gene flow[19,56]. High-resolution nuclear data are therefore essential to

help define evolutionary significant population units, disentangle convoluted ecological

relationships, and are particularly important for seabird conservation, which aims to preserve

genetic diversity considering profound global population declines[e.g. 7,8] and the threat of global

warming, which negatively impacts ecosystems worldwide[57].

Methods

Draft reference genome assembly

A *de novo* Atlantic puffin draft genome was generated from the blood of a female Atlantic puffin.

Read data was sequenced on three Illumina HiSeqX lanes using the 10x Genomics Chromium

technology and assembled with the Supernova assembler (v2.1.1)[58] after subsampling to 0.8

billion and 1 billion reads to maximize performance and remain within the computational capacity

of the assembler. We refined the two assemblies through several steps, including merging of

'haplotigs', removal of contaminant sequences, misassembly correction, re-scaffolding using

mapping coverage and linkage information, and gap filling (Table S1a). The most complete and

continuous 800M and 1000M assemblies together with the 3rd best assembly overall were

selected for a second round of refinement (Table S1b) resulting in a total of 72 draft assemblies.

Of these, we kept the four most complete and continuous assemblies for additional gap filling and

polishing, after which the most complete draft genome was selected for downstream analyses

(Table S1c). The puffin mitogenome was confidently identified by blasting (blastn) all scaffolds

shorter than 25 kb against a custom-built database of 135 published mitogenomes of the order

'Charadriiformes' and annotated with the MITOS web server[59] (Figure S1). The remaining nuclear

scaffolds were ordered and concatenated into "pseudo-chromosomes" by mapping them to the

razorbill genome (*Alca torda* - NCBI: bAlcTor1 primary, GCA_008658365.1) and applying 200 N's

as padding between each scaffold. We combined unmapped scaffolds into an "unplaced" pseudo-

chromosome. We assessed order and placement of scaffolds by investigating synteny in

coverage and length between the puffin and razorbill chromosomes (Table S2). Details on the

draft reference genome assembly and refinement can be found in the Supplementary File.

DNA extraction and sequencing

Samples from a total of 72 puffins collected across 12 breeding colonies (Figure 1a) were made

available for the present study by SEAPOP (http://www.seapop.no/en), SEATRACK

(http://www.seapop.no/en/seatrack) and ARCTOX (http://www.arctox.cnrs.fr/en/home - Canadian

colonies). These samples had been collected between 2012-2018 and consisted of blood

preserved in EtOH or lysis buffer, or feathers (Table S4). We extracted DNA using the DNeasy

Blood & Tissue kit (Qiagen) following the manufacturer's protocol for animal blood or the

nail/hair/feathers protocol applying several modifications for improved lysis and DNA yield.

Individuals that had no sexing data associated with them were sexed using PCR amplification of

specific allosome loci and visualization via gel electrophoresis. Genomic libraries were built by

the Norwegian Sequencing Centre and sequenced on an Illumina HiSeq4000. We processed

sequencing reads in PALEOMIX v1.2.14[60] and split the resulting bam files into nuclear and

mitochondrial bam files. Additional details on the DNA extraction, sexing, sequencing and

mapping are listed in the Supplementary File.

Mitogenome analyses

Genotypes across the mitochondrial genome were jointly called with GATK v4.1.4[61] by using the

HaplotypeCaller, *CombineGVCFs* and *GenotypeGVCFs* tool. We filtered genotypes according to

GATKs Best Practices[62] and set genotypes with a read depth less than 3 or a quality less than 15

as missing. Indels and non-biallelic SNPs were removed and only SNPs present in all individuals

were kept for subsequent analyses. The SNP dataset was annotated with snpEff[63] utilizing the

annotation of the newly assembled mitogenome of the Atlantic puffin and converted into a

mitogenome sequence alignment. To serve as an outgroup, we appended four other species of

the family Alcidae, i.e. the Razorbill (*Alca torda*, NCBI: CM018102.1), the Crested Auklet (*Aethia*

cristatella, NCBI: NC_045517.1), the Ancient Murrelet (*Synthliboramphus antiquus*, NCBI:

NC_007978.1) and the Japanese Murrelet (*Synthliboramphus wumizusume*, NCBI:

NC_029328.1), to the alignment. To construct a maximum-likelihood phylogenetic tree, we split

the alignment into seven partitions, i.e. one partition for a concatenated alignment of each of the

three codon positions of the protein coding genes, one partition for the concatenated alignment

of the rRNA regions, one partition for the concatenated alignment of the tRNAs, one partition for

the alignment of the control region, and one partition for the concatenated alignment of the

"intergenic" regions. The best-fitting evolutionary model for each partition was found by

ModelFinder[64] and the tree was built with IQTree v1.6.12[65] using 1000 ultrafast bootstrap

replicates. We used the resulting tree to draw a haplotype genealogy graph with Fitchi[66]. Using

Arlequin v.3.5[67], we calculated haplotype (h), nucleotide diversity (π) and Tajima's D[68] for each

colony, for each genomic cluster defined by the nuclear analysis, and globally. Additionally, an

Ewens–Watterson test[69], Chakraborty's test of population amalgamation[70] and Fu's F_s test[71] were

conducted for each of those groups. To further identify population differentiation, the proportion

of sequence variation (Φ_{ST}) was estimated for all pairs of populations and genomic clusters.

Hierarchical AMOVA tests subsequently determined the significance of *a priori* subdivisions into

colonies and genomic clusters. Calculation of Φ_{ST} and AMOVA tests were also conducted in

Arlequin. Additional details on the mitochondrial analyses are given in the Supplementary File.

Nuclear genome clustering and phylogenetic analyses

The majority of population genomic analyses were based on nuclear genotype likelihoods as

implemented in ANGSD v.0.931[72]. After assessing the quality of the mapped sequencing data in

an ANGSD pre-run, we removed an individual from the Isle of May from the dataset. Genotype

likelihoods for nuclear SNPs covered in all individuals were calculated and filtered in ANGSD.

Accounting for linkage disequilibrium, we further pruned the dataset by only selecting the most

central site within blocks of linked sites ($r^2 > 0.2$) as in Orlando and Librado (2019)[73]. Subsequently,

all variants located on the Z-pseudo-chromosome and "unplaced scaffolds" were excluded from

the analyses yielding a final genotype likelihood panel consisting of 1,093,765 sites. We

investigated genomic population structure with a Principal Component Analysis (PCA) of the

genotype likelihood panel using PCAngsd v0.982[74] and estimated individual ancestry proportions

using a maximum likelihood (ML) approach implemented in ngsAdmix v32[75], with the number of

ancestral populations (K) set from 1 to 10. Additional "hierarchical" PCA and admixture analyses

were conducted for genomic sub-cluster(s) using identical methods.

After adding the razorbill genome as an outgroup to the genotype likelihood panel by

mapping unpublished, raw 10x Genomics sequencing data used for the assembly of the

embargoed razorbill genome to the puffin draft assembly, we built a neighbor-joining (NJ) tree

based on pairwise genetic distance matrices (p-distance) and a sample-based ML phylogenetic

tree in FastMe v2.1.5[76] and Treemix v1.13[77], respectively. For both trees, 100 bootstrap replicates

were generated. To infer patterns of population splitting and mixing, we produced population-

based ML trees including up to ten migration edges. The optimal number of migrations was

selected using a quantitative approach by evaluating the distribution of explained variance, the

log likelihoods, the covariance with an increase in migration edges, and by applying the method

of Evanno[38] and several different linear threshold models. The topology for m_0 and m_{BEST} was

evaluated by generating 100 bootstrap replicates. Additional details on the cluster and

phylogenetic analyses are given in the Supplementary File.

Genetic diversity, heterozygosity and inbreeding

We calculated a set of neutrality tests and population statistics in ANGSD using colony-based

one-dimensional (1D) folded Site-Frequency-Spectra (SFS). For each population, genomic

cluster, and globally, Tajima's D and nucleotide diversity (π) were computed utilizing the per-site

θ estimates. Individual genome-wide heterozygosity was calculated in ANGSD using individual,

folded, 1D SFS. We calculated heterozygosity by dividing the number of polymorphic sites by the

number of total sites present in the SFS.

The proportion of runs of homozygosity (RoH) within each puffin genome was computed

by calculating local estimates of heterozygosity in 100 kbp sliding windows (50 kbp slide) following

the approach in Sánchez-Barreiro et al. (2020)[39]. We defined the 10% quantile of the average

local heterozygosity across all samples as the cutoff for a "low heterozygosity region". RoHs were

declared as all regions with at least two subsequent windows of low heterozygosity (below cutoff)

and their final length was calculated as described in Sánchez-Barreiro et al. (2020)[39]. We

calculated an individual inbreeding coefficient based on the RoH, F_{RoH} , as in Sánchez-Barreiro

et al. (2020)[39] by computing the fraction of the entire genome falling into RoHs, with the entire

genome being the total length of windows scanned. Additional details on these analyses can be

found in the Supplementary File.

Patterns of gene flow and admixture

Assessing potential landscape genetic patterns of IBD within the breeding range of the puffin, the

program EEMS[78] was used to model the association between genetic and geographic data by

visualizing the existing population structure and highlighting regions of higher-than-average and

lower-than-average historic gene flow. We calculated a pairwise genetic distance matrix in

ANGSD by sampling the consensus base (*-doIBS 2 -makeMatrix 1*) at the sites included in the

genotype likelihood set (see *Nuclear cluster and phylogenetic analyses*) for each sample. The

matrix was fed into 10 independent runs of EEMS, each consisting of one MCMC chain of six

million iterations with a two million iteration burn-in, 9999 thinning iterations, and 1000 underlying

demes. Supplementing the results of the EEMS analysis, we conducted a traditional IBD analysis

by determining geographical and genetic distances between the 12 colonies and assessing the

significance of the correlation between the two distance matrices with a Mantel test[79] and a

Multiple Regression on distance Matrix (MRM)[80] analysis. F_{ST} was used as a proxy for genetic

distance and computed for each population pair in ANGSD by applying two-dimensional (2D),

folded SFS. We converted pairwise F_{ST} values to Slatkin's linearized F_{ST}[81]. Great circle distances

between colony coordinates (latitude/longitude) were calculated using the R package marmap[82]

and used as geographic distances. We performed the Mantel test (10,000 permutations) and

MRM analysis with the R package *ecodist*[83]. All analyses for IBD were re-run on subsets of

colonies by progressively removing the Spitsbergen, Bjørnøya, Isle of May, and Canadian

colonies from the geographic and genetic distance matrices.

Additional assessments of gene flow and admixture were conducted by calculating f3-

statistics and multi-population D-statistics (aka ABBA BABA test)[84]. We calculated f3-statistics in

Treemix for each unique combination of ((A,B),C)) of the 12 puffin populations. D-statistics were

calculated in ANGSD (-doAbbababa2) for each combination of ((A,B),C),Outgroup) using the 12

puffin colonies. The outgroup was generated in ANGSD using the 10xGenomics sequencing data

of the razorbill mapped to the puffin reference genome (see *Nuclear cluster and phylogenetic*

analyses). Additional details on the IBD and admixture analyses are given in the Supplementary

File.

References

1. Otero, X. L., De La Peña-Lastra, S., Pérez-Alberti, A., Ferreira, T. O. & Huerta-Diaz, M. A.

Seabird colonies as important global drivers in the nitrogen and phosphorus cycles. *Nat.*

Commun. **9**, 246 (2018).

2. Velarde, E., Anderson, D. W. & Ezcurra, E. Seabird clues to ecosystem health. *Science*

365, 116–117 (2019).

3. Piatt, J. F., Sydeman, W. J. & Wiese, F. Introduction: a modern role for seabirds as

indicators. *Mar. Ecol. Prog. Ser.* **352**, 199–204 (2007).

4. Boersma, P. D., Clark, J. A. & Hillgarth, N. Seabird conservation. in *Biology of Marine Birds*

(eds. Schreiber, E. & Burger, J.) 559–579 (CRC Press Boca Raton, Florida, 2002).

5. Denlinger, L. & Wohl, K. Seabird harvest regimes in the circumpolar nations. *Conservation*

of Arctic Flora and Fauna (CAFF) (2001).

6. Merkel, F. & Barry, T. Seabird harvest in the Arctic. *Conservation of Arctic Flora and Fauna*

(CAFF) (2008).

7. Croxall, J. P. *et al.* Seabird conservation status, threats and priority actions: a global

assessment. *Bird Conserv. Int.* **22**, 1–34 (2012).

8. Paleczny, M., Hammill, E., Karpouzi, V. & Pauly, D. Population trend of the world's

monitored seabirds, 1950-2010. *PLoS One* **10**, e0129342 (2015).

9. Frederiksen, M. Seabirds in the North East Atlantic. Summary of status, trends and

anthropogenic impact. *TemaNord* **587**, 21–24 (2010).

10. Chardine, J. & Mendenhall, V. Human disturbance at Arctic seabird colonies. *Conservation*

of Arctic Flora and Fauna (CAFF) (1998).

11. Allendorf, F. W., Hohenlohe, P. A. & Luikart, G. Genomics and the future of conservation

genetics. *Nat. Rev. Genet.* **11**, 697 (2010).

12. Fraser, D. J. & Bernatchez, L. Adaptive evolutionary conservation: towards a unified

concept for defining conservation units. *Mol. Ecol.* **10**, 2741–2752 (2001).

13. Funk, W. C., McKay, J. K., Hohenlohe, P. A. & Allendorf, F. W. Harnessing genomics for

delineating conservation units. *Trends Ecol. Evol.* **27**, 489–496 (2012).

14. Friesen, V. L. Speciation in seabirds: why are there so many species… and why aren't

there more? *J. Ornithol.* **156**, 27–39 (2015).

15. Taylor, R. S. *et al.* Sympatric population divergence within a highly pelagic seabird species

complex (*Hydrobates spp.*). *J. Avian Biol.* **49**, (2018).

16. Rexer-Huber, K. *et al.* Genomics detects population structure within and between ocean

basins in a circumpolar seabird: The white-chinned petrel. *Mol. Ecol.* **28**, 4552–4572

(2019).

17. Clucas, G. V. *et al.* Comparative population genomics reveals key barriers to dispersal in

Southern Ocean penguins. *Mol. Ecol.* **27**, 4680–4697 (2018).

18. Frugone, M. J. *et al.* More than the eye can see: Genomic insights into the drivers of

genetic differentiation in Royal/Macaroni penguins across the Southern Ocean. *Mol.*

Phylogenet. Evol. **139,** (2019).

19. Cristofari, R. *et al.* Unexpected population fragmentation in an endangered seabird: the

case of the Peruvian diving-petrel. *Sci. Rep.* **9**, 2021 (2019).

20. Tigano, A., Shultz, A. J., Edwards, S. V., Robertson, G. J. & Friesen, V. L. Outlier analyses

to test for local adaptation to breeding grounds in a migratory arctic seabird. *Ecol. Evol.* **7**,

2370–2381 (2017).

21. Lowry, D. B. *et al.* Breaking RAD: an evaluation of the utility of restriction site-associated

DNA sequencing for genome scans of adaptation. *Mol. Ecol. Resour.* **17**, 142–152 (2017).

22. Somvichian-Clausen, A. Behind the stunning photo of a puffin gorging on fish. *National*

Geographic (2017).

23. Huijbens, E. H. & Einarsson, N. Feasting on friends: Whales, puffins, and tourism in

Iceland. in *Tourism Experiences and Animal Consumption* 10–27 (Routledge, 2018).

24. Lund, K. A., Kjartansdóttir, K. & Loftsdóttir, K. 'Puffin love': Performing and creating Arctic

landscapes in Iceland through souvenirs. *Tourist Studies* **18**, 142–158 (2018).

25. Hodgetts, L. M. Animal bones and human society in the late younger stone age of arctic

Norway. (Durham University, 1999).

26. Dove, C. J. & Wickler, S. Identification of bird species used to make a Viking age feather

pillow. *Arctic* **69**, 29–36 (2016).

27. Harris, M. P. & Wanless, S. *The puffin*. (T & AD Poyser, Bloomsbury Publishing, 2011).

28. BirdLife International. Fratercula arctica. *The IUCN Red List of Threatened Species 2017*

(2017)

29. Anker-Nilssen, T. & Aarvak, T. The population ecology of puffins at Røst. Status after the

breeding season 2001. *NINA Oppdragsmelding* **736**, 1–40 (2002).

30. Anker-Nilssen, T. *et al.* Key-site monitoring in Norway 2019, including Svalbard and Jan

Mayen. *SEAPOP Short Report 1-2020* (2020).

31. Lilliendahl, K. *et al.* Recruitment failure of Atlantic puffins *Fratercula arctica* and sandeels

Ammodytes marinus in Vestmannaeyjar Islands. *Náttúrufræðingurinn* **83**, 65–79 (2013).

32. Gaston, A. J. & Provencher, J. F. A Specimen of the High Arctic Subspecies of Atlantic

Puffin, *Fratercula arctica naumanni*, in Canada. *Can. Field-Nat.* **126**, 50–54 (2012).

33. Salomonsen, F. *The Atlantic Alcidae*. vol. 6 (Elanders boktryckeri aktiebolag, 1944).

34. Moen, S. M. Morphologic and genetic variation among breeding colonies of the Atlantic

puffin (*Fratercula arctica*). *Auk* **108**, 755–763 (1991).

35. Harris, M. P. Measurements and weights of British Puffins. *Bird Study* **26**, 179–186 (1979).

36. Kim, J. A., Kang, S.-G., Yang, J. W., Hur, W.-H. & Kil, H.-J. Complete mitochondrial

genome of *Aethia cristatella* (Charadriiformes: Alcidae). *Mitochondrial DNA Part B* **5**, 31–32

(2020).

37. Eo, S. H. & An, J. The complete mitochondrial genome sequence of Japanese murrelet

(Aves: Alcidae) and its phylogenetic position in Charadriiformes. *Mitochondrial DNA A DNA*

Mapp Seq Anal **27**, 4574–4575 (2016).

38. Evanno, G., Regnaut, S. & Goudet, J. Detecting the number of clusters of individuals using

the software STRUCTURE: a simulation study. *Mol. Ecol.* **14**, 2611–2620 (2005).

39. Sánchez-Barreiro, F. *et al.* Historical population declines prompted significant genomic

erosion in the northern and southern white rhinoceros (*Ceratotherium simum*). *bioRxiv*

(2020) doi:10.1101/2020.05.10.086686.

40. Lombal, A. J., O'dwyer, J. E., Friesen, V., Woehler, E. J. & Burridge, C. P. Identifying

mechanisms of genetic differentiation among populations in vagile species: historical

factors dominate genetic differentiation in seabirds. *Biol. Rev. Camb. Philos. Soc.* **95**, 625–

651 (2020).

41. Breton, A. R., Diamond, A. W. & Kress, S. W. Encounter, survival, and movement

probabilities from an Atlantic puffin (*Fratercula arctica*) metapopulation. *Ecological*

Monographs **75**, (2006).

42. Fayet, A. L. *et al.* Ocean-wide drivers of migration strategies and their influence on

population breeding performance in a declining seabird. *Curr. Biol.* **27**, 3871–3878 (2017).

43. Friesen, V. L., Burg, T. M. & McCoy, K. D. Mechanisms of population differentiation in

seabirds. *Mol. Ecol.* **16**, 1765–1785 (2007).

44. Burg, T. M. & Croxall, J. P. Global relationships amongst black-browed and grey-headed

albatrosses: analysis of population structure using mitochondrial DNA and microsatellites.

Mol. Ecol. **10**, 2647–2660 (2001).

45. Lowther, P. E., Diamond, T., Kress, S. W., Robertson, G. J. & Gill, F. Atlantic Puffin

(*Fratercula arctica*). *The Birds of North America Online* **18**, (2002).

46. Wojczulanis-Jakubas, K. *et al.* Weak population genetic differentiation in the most

numerous Arctic seabird, the little auk. *Polar Biol.* **37**, 621–630 (2014).

47. Smith, A. L., Monteiro, L., Hasegawa, O. & Friesen, V. L. Global phylogeography of the

band-rumped storm-petrel (*Oceanodroma castro*; Procellariiformes: Hydrobatidae). *Mol.*

Phylogenet. Evol. **43**, 755–773 (2007).

48. Bergmann, C. Über die Verhältnisse der Wärmeökonomie der Tiere zu ihrer Grösse.

Gottinger Studien **3**, 595–708 (1847).

49. James, F. C. Geographic size variation in birds and its relationship to climate. *Ecology* **51**,

365–390 (1970).

50. Yamamoto, T. *et al.* Geographical variation in body size of a pelagic seabird, the streaked

shearwater *Calonectris leucomelas*. *J. Biogeogr.* **43**, 801–808 (2016).

51. Barrett, R. T., Anker-Nilssen, T. & Krasnov, Y. V. Can Norwegian and Russian razorbills

(*Alca torda*) be identified by their measurements? *Mar. Ornithol.* **25**, 5–8 (1997).

52. Anker-Nilssen, T., Aarvak, T. & Bangjord, G. Mass mortality of Atlantic Puffins *Fratercula*

arctica off Central Norway, spring 2002: causes and consequences. *Atl. Seabirds* **5**, 57–72

(2003).

53. Pearce, R. L. *et al.* Mitochondrial DNA suggests high gene flow in ancient murrelets.

Condor **104**, 84–91 (2002).

54. Thomas, J. E. *et al.* Demographic reconstruction from ancient DNA supports rapid

extinction of the great auk. *eLife* **8**, (2019).

55. Milot, E., Weimerskirch, H. & Bernatchez, L. The seabird paradox: dispersal, genetic

structure and population dynamics in a highly mobile, but philopatric albatross species. *Mol.*

Ecol. **17**, 1658–1673 (2008).

56. Edwards, S. & Bensch, S. Looking forwards or looking backwards in avian

phylogeography? A comment on Zink and Barrowclough 2008. *Mol. Ecol.* **18,** 2930–2936

(2009).

57. IPCC. *Global Warming of 1.5 °C - Summary for Policy Makers*. (2018).

58. Weisenfeld, N. I., Kumar, V., Shah, P., Church, D. M. & Jaffe, D. B. Direct determination of

diploid genome sequences. *Genome Res.* **27**, 757–767 (2017).

59. Bernt, M. *et al.* MITOS: improved de novo metazoan mitochondrial genome annotation.

Mol. Phylogenet. Evol. **69**, 313–319 (2013).

60. Schubert, M. *et al.* Characterization of ancient and modern genomes by SNP detection and

phylogenomic and metagenomic analysis using PALEOMIX. *Nat. Protoc.* **9**, 1056–1082

(2014).

61. McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing

next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

62. Van der Auwera, G. A. *et al.* From FastQ data to high confidence variant calls: the Genome

Analysis Toolkit best practices pipeline. *Curr. Protoc. Bioinformatics* **43**, 11.10.1–33 (2013).

63. Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide

polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118;

iso-2; iso-3. *Fly* **6**, 80–92 (2012).

64. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.

ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**,

587–589 (2017).

65. Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: a fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* **32**,

268–274 (2015).

66. Matschiner, M. Fitchi: haplotype genealogy graphs based on the Fitch algorithm.

Bioinformatics **32**, 1250–1252 (2016).

67. Excoffier, L. & Lischer, H. E. L. Arlequin suite ver 3.5: a new series of programs to perform

population genetics analyses under Linux and Windows. *Mol. Ecol. Resour.* **10**, 564–567

(2010).

68. Tajima, F. Statistical method for testing the neutral mutation hypothesis by DNA

polymorphism. *Genetics* **123**, 585–595 (1989).

69. Watterson, G. A. Heterosis or neutrality? *Genetics* **85**, 789–814 (1977).

70. Chakraborty, R. Mitochondrial DNA polymorphism reveals hidden heterogeneity within

some Asian populations. *Am. J. Hum. Genet.* **47**, 87–94 (1990).

71. Fu, Y. X. Statistical tests of neutrality of mutations against population growth, hitchhiking

and background selection. *Genetics* **147**, 915–925 (1997).

72. Korneliussen, T. S., Albrechtsen, A. & Nielsen, R. ANGSD: Analysis of next generation

sequencing data. *BMC Bioinformatics* **15**, 356 (2014).

73. Orlando, L. & Librado, P. Origin and evolution of deleterious mutations in horses. *Genes*

10, (2019).

74. Meisner, J. & Albrechtsen, A. Inferring population structure and admixture proportions in

low-depth NGS data. *Genetics* **210**, 719–731 (2018).

75. Skotte, L., Korneliussen, T. S. & Albrechtsen, A. Estimating individual admixture

proportions from next generation sequencing data. *Genetics* **195**, 693–702 (2013).

76. Lefort, V., Desper, R. & Gascuel, O. FastME 2.0: A comprehensive, accurate, and fast

distance-based phylogeny inference program. *Mol. Biol. Evol.* **32**, 2798–2800 (2015).

77. Pickrell, J. K. & Pritchard, J. K. Inference of population splits and mixtures from genome-

wide allele frequency data. *PLoS Genet.* **8**, e1002967 (2012).

78. Petkova, D., Novembre, J. & Stephens, M. Visualizing spatial population structure with

estimated effective migration surfaces. *Nat. Genet.* **48**, 94–100 (2016).

79. Mantel, N. The detection of disease clustering and a generalized regression approach.

Cancer Res. **27**, 209–220 (1967).

80. Lichstein, J. W. Multiple regression on distance matrices: a multivariate spatial analysis

tool. *Plant Ecol.* **188**, 117–131 (2007).

81. Slatkin, M. A measure of population subdivision based on microsatellite allele frequencies.

Genetics **139**, 457–462 (1995).

82. Pante, E., Simon-Bouhet, B. & Irisson, J.-O. *marmap - R package*. (2019).

83. Goslee, S. & Urban, D. The ecodist package for dissimilarity-based analysis of ecological

data. *Journal of Statistical Software, Articles* **22**, 1–19 (2007).

84. Soraggi, S., Wiuf, C. & Albrechtsen, A. Powerful inference with the D-statistic on low-

coverage whole-genome data. *G3* **8**, 551–566 (2018).

607
608

Acknowledgements

Financial support was provided by the Nansen Foundation and the Faculty of Mathematical and

Natural Sciences, University of Oslo (UiO). We are grateful to the SEAPOP program

(www.seapop.no/en, Norwegian Research Council grant number 192141), and the SEATRACK

(http://www.seapop.no/en/seatrack) and ARCTOX (https://arctox.cnrs.fr/en/home) projects for

collecting and sharing of samples. In particular, we thank Dave Fifield and Greg Robertson for

the provision of puffin samples from Gull Island. The authors acknowledge support from the

National Genomics Infrastructure in Stockholm funded by the Science for Life Laboratory, the

Knut and Alice Wallenberg Foundation and the Swedish Research Council, and the

SNIC/Uppsala Multidisciplinary Center for Advanced Computational Science for assistance with

massively parallel sequencing (reference genome) and access to the UPPMAX computational

infrastructure. Special thanks goes to the Norwegian Sequencing Centre (UiO) for the genomic

libraries and resequencing of samples analyzed in this study. The razorbill genome data was

made available for this study by Tom Gilbert and the Vertebrate Genome Project. Computation

was performed using the resources and assistance from SIGMA2. Albína Pálsdóttir shared scripts

for ANGSD, and Emiliano Trucchii advised on the manuscript. Pictures of puffins used in the

figures were taken by Annemarie Look.

Author Contributions

SB and BS conceptualized the project. MI performed the DNA extraction for the reference

genome. OK did all other laboratory work. KSJ advised on the sequencing strategy and co-

supervised OK. OK refined the reference genome assembly. OK carried out the population

genomic analyses with input from SB, BS and DML. OK designed the figures with input from SB,

BS and DML. TAN, HS and ESH advised on colony selection and provided ecological context.

TAN, HS, OdK, SD, ESH, JF, MPH, JD, KE, MLM and MGF provided samples. OK wrote the

paper with SB, BS and DML. All authors read and revised the final version of the manuscript.

Competing Interests

The authors declare no competing interests.

Data availability

Raw read data have been deposited in the European Nucleotide Archive (ENA,

www.ebi.ac.uk/ena) under study accession number PRJEB40631 (see Table S4 for individual

sample accession numbers). Nuclear and mitochondrial scaffolds (GCA_905066775.1,

CAJHIB010000001-CAJHIB010013329), as well as pseudo-chromosomes (GCA_905066775.2,

CAJHIB020000001-CAJHIB020000027), have been uploaded to ENA (Project PRJEB40926,

Sample SAMEA7482542).

Code availability

Full code used for the population genomic analyses is available on the first author's GitHub

(https://github.com/OKersten/PuffPopGen).

652
Figure 1: Sampling distribution and genomic structure of 71 Atlantic puffin individuals across 12
colonies throughout the breeding range. a) Map presenting the location of the 12 sampling sites.
Colour shading indicates the breeding range of the species as a whole, as well as the recognized
subspecies. b) Mitochondrial haplotype network constructed using the program Fitchi and a Maximum
Likelihood tree generated in IQTree. It contains 66 unique haplotypes identified by 192 mitogenome-wide
SNPs. Sizes of circles are proportional to haplotype abundance. Color legend is provided in c). Black dots
represent inferred haplotypes that were not found in the present sampling. c) Principal Component
Analysis (PCA) using genotype likelihoods at 1,093,765 polymorphic nuclear sites calculated in ANGSD
to project the 71 individuals onto PC axes 1 and 2. Each circle represents a sample and colours indicate
the different colonies. Percentage indicates proportion of genomic variation explained by each axis. The
colour coding of the colonies is consistently used throughout the manuscript. d) CLUMPAK-averaged
admixture plots of the best K's using the same genotype likelihood panel as in c). Each column
represents a sample and colonies are separated by solid white lines. Optimal K's were determined by the
method of Evanno et al. (2005) (see Figure S5a) and colors indicate the ancestry fraction to the different
clusters.

669
670
Figure 2: Phylogenetic reconstruction of individual and colony relationships from 71 Atlantic
puffin individuals sampled across 12 colonies throughout the species' breeding range. a) An
individual-based neighbor-joining tree constructed using pairwise p-distances calculated from genotype
likelihoods at 1,093,765 polymorphic nuclear sites. Branch lengths and the outgroup were removed for
the zoomed-in section to improve visualization. b) A population-based maximum likelihood Treemix
analysis using allele frequencies at the same 1,093,765 polymorphic nuclear sites as in a). Both trees are
rooted using the razorbill as an outgroup. The tree in b) is visualized with and without the outgroup.
Branch lengths are equivalent to a genetic drift parameter. The heatmap indicates the residual fit of the
tree displaying the standard error of the covariance between populations. In a) and b), the colour coding
of the colonies is consistent with those in Figure 1 and node labels show bootstrap support >80.

681
682
Figure 3: Genome-wide heterozygosity, inbreeding and Runs-of-Homozygosity (RoH) compared
between 12 Atlantic puffin colonies across the species' breeding range. a) Estimates of individual
genome-wide heterozygosity based on the per-sample one-dimensional Site Frequency Spectrum
calculated in ANGSD. b) Individual inbreeding coefficients, F_{RoH}, defined as the fraction of the individual
genomes falling into RoHs of a minimum length of 150 kbp. RoHs were declared as all regions with at
least two subsequent 100 kbp windows harboring a heterozygosity below 1.435663×10^{-3}. c) RoH length
distribution across the 12 colonies only including RoHs longer than 500 kbp. A single 9.65 Mbp long RoH
on pseudo-chromosome 7 in an Isle of May individual required to introduce a break in the y-axis. In a)
and b), black dots indicate individual sample estimates and black lines the median per colony, while in c),
black dots represent single RoHs. Statistical significance of differences in heterozygosity and F_{RoH}
between populations was assessed with a global Kruskal-Wallis test. The results of *post hoc* Dunn tests
with Holm corrections are presented in Figure S11. Different colonies in all three plots are indicated using
the same color code as in Figure 1.

696
697
Figure 4: Estimates of continuous long-distance gene flow and isolation by distance (IBD) across the breeding range of the Atlantic puffin estimated from 71 individuals across 12 colonies. a) Effective migration surfaces inferred by the program EEMS using the average distance between pairs of individuals calculated in ANGSD by sampling the consensus base for each individual at 1,093,765 polymorphic nuclear sites. Darker reds indicate reduced migration across those areas, while darker blues highlight higher migration rates than the global mean. Different colonies are indicated using colors consistent with those in Figure 1. b) Correlation between genetic (Slatkin's linearized F_{ST}) and geographic (great-circle) distance presented after removing the Spitsbergen, Bjørnøya, Isle of May and Canadian individuals. The diagonal line visualizes the result of the Multiple Regression on Distance Matrices (MRM) analysis (slope and y-intercept). The Mantel test between genetic and geographic distance ($r = 0.826$, $p < 0.01$) was significant and 68.18% of variance in Slatkin's linearized F_{ST} was explained by geographic distance (regression coefficient of linear IBD model = 0.83×10^{-6}, $p < 0.01$). A two-dimensional kernel density estimation (kde2d) highlights dense groups of data points, thus substructure in the genomic landscape pattern. Analyses were conducted and results visualized in R using the *ecodist*, *marmap* and *MASS* packages.