Transcription and splicing dynamics during early *Drosophila* development

Pedro Prudêncio [1, 2 §], Rosina Savisaar [1§], Kenny Rebelo [1§$], Rui Gonçalo Martinho[1, 2, 3*] and Maria Carmo-Fonseca[1*]

[1] Instituto de Medicina Molecular João Lobo Antunes, Faculdade de Medicina, Universidade de Lisboa, 1649-028 Lisboa, Portugal.

[2] Algarve Biomedical Center Research Institute (ABC-RI), Universidade do Algarve, 8005-139 Faro, Portugal.

[3] Department of Medical Sciences and Institute for Biomedicine (iBiMED), Universidade de Aveiro, 3810-193 Aveiro, Portugal.

[§]These authors have contributed equally to the work.

[$]Present address: Donnelly Centre, University of Toronto, Toronto, ON M5S 3E1, Canada

[*]Joint corresponding authors: carmo.fonseca@medicina.ulisboa.pt and rgmartinho@ua.pt

RUNNING TITLE: Transcription and splicing dynamics in *Drosophila*

KEYWORDS: splicing kinetics; Pol II pausing; transcription termination; NET-seq; *Drosophila melanogaster* embryo

ABSTRACT

Widespread co-transcriptional splicing has been demonstrated from yeast to human. However, the precise kinetics of splicing relative to transcription remains unclear, with discrepancies between studies that use different experimental approaches. Here, we adapted native elongating transcript sequencing technology (NET-seq) to measure co-transcriptional splicing dynamics during the early developmental stages of *Drosophila melanogaster* embryos. Our results reveal the position of RNA polymerase II (Pol II) when both canonical and recursive splicing occur. We found heterogeneity in splicing dynamics, with some RNAs spliced immediately after intron transcription, whereas for other transcripts no splicing was observed over the first 100 nucleotides of the downstream exon. Introns that show splicing completion before Pol II has reached the end of the downstream exon are necessarily intron-defined. We studied the splicing dynamics of both nascent pre-mRNAs transcribed in the early embryo, which have few and short introns, as well as pre-mRNAs transcribed later in embryonic development, which contain multiple long introns. As expected, we found a relationship between the proportion of spliced reads and intron size. However, intron definition was observed at all intron sizes. We further observed that transcription termination was very efficient for isolated genes but that the presence of an overlapping antisense gene was often associated with transcriptional readthrough. Taken together, our data unravels novel dynamic features of Pol II transcription and splicing in the developing *Drosophila* embryo.

INTRODUCTION

It is by now largely accepted that splicing can happen, and often does happen, whilst transcription is still in progress (Ameur et al. 2011; Khodor et al. 2011; Beyer and Osheim 1988; Brugiolo et al. 2013; Carrillo Oesterreich et al. 2010; Windhager et al. 2012; Nojima et al. 2015; Alpert et al. 2017). It is also known that the processes of splicing and transcription are tightly interlinked. The RNA Polymerase II (Pol II) elongation rate can affect exon inclusion (Fong et al. 2014; Maslon et al. 2019; Aslanzadeh et al. 2018), and many of the proteins involved in splicing associate with the Pol II large subunit C-terminal domain (CTD) (Hsin and Manley 2012; Emili et al. 2002; David et al. 2011; Görnemann et al. 2011; Morris and Greenleaf 2000; Nojima et al. 2018). Conversely, splicing may also affect transcription, with evidence suggesting that Pol II slows down at exons (Alexander et al. 2010; Carrillo Oesterreich et al. 2010; Jonkers et al. 2014; Mayer et al. 2015; Veloso et al. 2014), potentially to allow for splicing to complete.

The exact timing of intron excision relative to Pol II elongation remains more controversial. A nascent RNA sequencing study (Carrillo Oesterreich et al. 2016) showed that in *Saccharomyces cerevisiae*, splicing could occur as soon as the 3' splice site had emerged from the transcription machinery, suggesting that splicing may be completed immediately after intron transcription. Previous attempts to determine the duration of splicing *in vivo* had returned estimates ranging from a few seconds to several minutes (Alpert et al. 2017). The lowest of these estimates, such as the few-second estimate reported in Martin et al. (2013), are consistent with splicing being completed right after the 3' splice site is transcribed.

More recently, two long-read nascent RNA sequencing studies addressed the timing of splicing in metazoa. One reported little immediate splicing in human and *Drosophila* cultured cell lines, with most splicing occurring when Pol II had elongated thousands of nucleotides past the splice junction (Drexler et al. 2020). Another found that splicing in mouse cells occurred immediately after transcription, when Pol II was within 75-300 nucleotides of 3' splice sites (Reimer et al. 2021). These discrepant results probably reflect biases introduced by the distinct methodological approaches used. Indeed, one study (Drexler et al. 2020) relies on metabolic labelling of newly synthesized RNA with 4sU, while the other extracts RNA from chromatin (Reimer et al. 2021).

An alternative approach to analyse nascent pre-mRNA is Native Elongating Transcript Sequencing (NET-seq), a technique that involves isolation of transcription complexes formed by Pol II, the DNA template and the nascent RNA by immunoprecipitation, without crosslinking (Churchman and Weissman 2011; Nojima et al. 2015). Solubilisation of Pol II complexes under native conditions is typically incomplete in metazoan cells (Kimura et al. 1999), but this problem was circumvented by extensive micrococcal nuclease (MNase) digestion of isolated native chromatin (Nojima et al. 2015). Elongation complexes can then be immunoprecipitated using antibodies that specifically recognize the different phosphorylation states of the Pol II CTD. The main advantage of this approach is that it provides global, bidirectional maps of Pol II CTD phosphorylation specific nascent transcripts and coupled RNA processing at single nucleotide resolution. Although initially applied to mammalian cells, and thus termed mNET-seq (Nojima et al. 2015), the technology can be adjusted to any organism, as recently illustrated in plants (Zhu et al. 2018). Here, we have adapted NET-seq to study the dynamic properties of transcription and co-transcriptional splicing during early stages of development in *Drosophila melanogaster* embryos. An advantage of this model system is that compared to human, the *Drosophila* genome is more compact (Graveley et al. 2011) and thus the coverage of NET-seq reads on intragenic regions is higher. Another advantage of the *Drosophila* model is that genes transcribed initially in the embryo have few and short introns (like yeast genes), whereas genes transcribed later contain multiple long introns (more similar to human genes). An additional feature of *Drosophila* is the presence of many genes with exceptionally long introns that are subdivided by a non-canonical mechanism termed recursive splicing (Duff et al. 2015; Joseph et al. 2018; Pai et al. 2018). For all these reasons, *Drosophila* is an attractive model to study co-transcriptional splicing dynamics.

Drosophila early development is characterized by rapid mitotic cycles that lack cytokinesis, resulting in nuclear proliferation in a syncytial cytoplasm (Campos-Ortega 1985; Laver et al. 2015). During these initial mitotic divisions, which impose significant constraints on transcription and splicing (Shermoen and O'Farrell 1991; Rothe et al. 1992; Guilgur et al. 2014; Martinho et al. 2015; Sandler et al. 2018; Kwasnieski et al. 2019), the embryo largely relies on maternally deposited mRNAs. Subsequently, the duration of the cell cycle is progressively expanded, membranes form between the nuclei and segregate them into cells, and the zygotic transcriptome starts to be fully expressed (Campos-

108   Ortega 1985; Yuan et al. 2016; Blythe and Wieschaus 2015). By performing *d*NET-seq in the

109   developing *Drosophila* embryo, we have unravelled novel dynamic features of Pol II transcription and

110   splicing.

RESULTS

Native elongating transcript sequencing in *Drosophila* embryos (*d*NET-seq)

Our first task was to adapt the NET-seq technology for use in *Drosophila* embryos (henceforth referred to as *d*NET-seq). We collected embryos at 2-3 hours after fertilization (referred to as early embryos) and 4-6 hours after fertilization (referred to as late embryos) (**Fig 1A**). Analysis of embryos stained with a fluorescent dye to visualize DNA (**Fig 1B**) revealed that early embryos were predominantly in mitotic cycle 14 (stage 5), when membranes form between the nuclei, which are evenly distributed in a monolayer under the egg surface, segregating them into separate cells (Campos-Ortega 1985; Laver et al. 2015). The majority of late embryos were in the late stage of germ-band extension (stage 10; **Fig 1C**), when the embryo trunk (also known as the germ-band) elongates in the antero-posterior axis and narrows in the dorso-ventral axis (Campos-Ortega 1985; Laver et al. 2015).

In adapting mNET-seq to *Drosophila* embryos, we optimized buffers and washing conditions to purify the chromatin fraction from manually sorted embryos, solubilize the transcription complexes with MNase digestion and immunoprecipitate Pol II with antibodies. We used rabbit polyclonal antibodies raised against synthetic peptides of the YSPTSPS repeat of the CTD of the largest Pol II subunit in *Saccharomyces cerevisiae*, phosphorylated at either S5 (ab5131 Abcam) or S2 (ab5095 Abcam). Both antibodies have been extensively used for chromatin immunoprecipitation experiments in *Drosophila melanogaster* (Boija et al. 2017; Vizcaya-Molina et al. 2018; Akhtar et al. 2019; Dahlberg et al. 2015; Lee et al. 2015; Yan et al. 2014; Arzate-Mejía et al. 2020).

To enable directional sequencing, the 5' hydroxyl (OH) generated by MNase digestion of RNA was first converted to a 5' phosphate by T4 polynucleotide kinase (**Fig 1D**). RNA was then purified from the immunoprecipitated Pol II complexes and size-selected using an RNA purification kit procedure that combines a unique buffer system with a column technology (see the methods section for more detail). RNAs with a size above 60 nucleotides (nt) were used for subsequent ligation of specific adapters to the 5' P and 3' OH ends of each RNA fragment followed by PCR-based preparation of a cDNA library for high-throughput Illumina sequencing (**Fig 1D**). After sequencing, adapter sequences were trimmed and paired-end reads with sequence overlaps were merged into a single read that spans

the full length of the original RNA fragment (dark orange; **Fig 1E**). The resulting single reads were

aligned to the *Drosophila* reference genome. The nucleotide at the 3' end of each RNA fragment was

identified and its genomic position recorded (asterisk; **Fig 1E**).

Two to three NET-seq libraries were independently prepared from early and late embryos using

S5P antibody; three additional libraries were prepared from late embryos using S2P antibody. Each

library was sequenced to a high coverage with a read length of 150 bp (Supplemental **Fig S1A**; see the

methods section for more detail). Experimental reproducibility was demonstrated by strong agreement

of uniquely aligned read density between biological replicates prepared with antibodies raised against

the CTD phosphorylated on either serine 5 (S5P) or serine 2 (S2P) positions (Supplemental **Fig S1 B-**

E). This suggests that both S5P and S2P antibodies recognize the CTD of elongating Pol II in

Drosophila embryos.

Figure 1. Native elongating transcript sequencing in *Drosophila* embryos.
(A) Timeline of *Drosophila* early embryonic development. **(B)** Representative images (stained for DNA) of embryos in mitotic cycle 14 (stage 5) and late germ-band expansion (stage 10). **(C)** The graph depicts the developmental stage of embryos sorted into the "early" and "late" groups. **(D)** Outline of the *d*NET-seq experimental protocol. **(E)** Outline of *d*NET-seq data analysis.

_d_NET-seq captures splicing intermediates and spliceosomal snRNAs.

NET-seq captures not only the final (3' OH end) nucleotide of nascent RNA but also the 3' OH end of

RNAs that associate with the Pol II elongation complex (Nojima et al. 2015, 2018; Schlackow et al.

2017). Notably, in humans, mNET-seq of Pol II phosphorylated on CTD serine 5 detected splicing

intermediates formed by cleavage at the 5' splice site after the first splicing reaction. The presence of

such intermediates manifests as an enrichment of reads whose 3' ends map precisely to the last

nucleotide of an exon. In both early and late *Drosophila* embryos, we indeed observed large peaks of

*d*NET-seq/S5P reads mapping to the last nucleotide of spliced exons, as shown for the *kuk* gene

(asterisk; **Fig 2A**). We also detected *d*NET-seq/S5P peaks at the last nucleotide of introns, as shown for

the *eEF1alpha1* gene (asterisk; **Fig 2B**); enrichment for these reads results from co-

immunoprecipitation of released intron lariats after completion of the splicing reaction (**Fig 2B**). In

addition, we observed reads corresponding to mature snRNAs engaged in co-transcriptional

spliceosome assembly, suggesting that *d*NET-seq was capturing the free 3' OH ends of the snRNAs.

(**Fig 2C**). We found prominent peaks at the end of spliceosomal U1, U2, U4 and U5 snRNAs. As

expected, no peak was detected mapping to the end of the U3 snRNA, which is involved in the

processing of pre-rRNA synthesized by Pol I. Noteworthy, we observed an accumulation of *d*NET-seq

signal at the end of U6 snRNA (Supplemental **Fig S2A**), contrasting with a lack of peak observed in

mammalian cells (Nojima et al. 2018). This is consistent with the finding that most mammalian U6

snRNAs contain a 2',3'-cyclic phosphate terminal group at the 3' end, whereas U6 3' ends in

Drosophila cells consist of either a cyclic 2',3'-phosphate, a 3'-phosphate or a 2',3'-hydroxyl group

(Lund and Dahlberg 1992).

To quantify how many constitutively spliced exons have *d*NET-seq peaks at the end, we applied

an algorithm that finds nucleotides where the NET-seq read density is at least three standard deviations

above the transcript mean in a local region defined by 100 bp upstream and downstream (Churchman

and Weissman 2011; Prudêncio et al. 2020). Upon analysing replicates of *d*NET-seq/S5P and *d*NET-

seq/S2P libraries, we identified over 10,000 exons showing splicing intermediate peaks (**Fig 2D**). As

peaks were more frequently detected on exons of genes with higher read density (Supplemental **Fig**

S2B), we classified the exons into four groups (quartiles) based on the *d*NET-seq read density of the

corresponding gene and restricted the analysis to exons in the fourth quartile, i.e., from genes with the

highest read density. The results show that splicing intermediate peaks are detected in approximately

80% of all constitutively spliced exons. The proportion is similar for pre-MBT genes, i.e, zygotic genes

that start to be transcribed before cycle 14 and remain active thereafter (Chen et al. 2013), and for genes

expressed in late embryos (**Fig 2E**). Similar proportions were found in the S5P and the S2P datasets

(**Fig 2E**). We then used the same methodology and the same set of genes to detect peaks at the last

intronic nucleotide, corresponding to released intron lariats. Such peaks were detected in less than 10%

of introns (**Fig 2F**).

Taken together, these results demonstrate that *d*NET-seq is capable of detecting splicing

intermediates and spliceosomal snRNAs in *Drosophila* embryos, as previously reported in mammalian

cells (Nojima et al. 2015; Schlackow et al. 2017; Nojima et al. 2018).

Figure 2. *d*NET-seq captures splicing intermediates and spliceosomal snRNAs. (A-C) *d*NET-
seq/S5P profiles over the indicated genes in late embryos. The diagrams outline the 3' OH ends
generated by co-transcriptional cleavage at the 5' splice site (green, **A**), the 3' splice site (red, **B**) and
the free 3' OH end of spliceosomal snRNAs (blue, **C**). Arrows indicate the direction of transcription.

Exons are represented by boxes. Thinner boxes represent UTRs. Introns are represented by lines connecting the exons. Asterisks denote the peak at the end of the exon (**A**), the end of the intron (**B**), and the end of the U2 snRNA gene (**C**). (**D**) Comparison (Venn diagrams) of exons with a splicing intermediate peak detected in biological replicates of *d*NET-seq/S5P and *d*NET-seq/S2P libraries. (**E, F**) Frequency of peaks corresponding to splicing intermediates (**E**) and released intron lariats (**F**) in pre-MBT genes and genes expressed in late embryos. Only genes with the highest read density (fourth quartile) were considered.

dNET-seq specifically captures nascent RNA.

To validate that we were detecting nascent transcription in *Drosophila* embryos, we analysed maternal mRNAs that are transcribed during oogenesis and loaded into the egg (**Fig 3A**). As expected, maternal transcripts such as *bicoid* (**Fig 3B**), *nanos* (Supplemental **Fig S3A**), *gurken* (Supplemental **Fig S3B**) and *Rab32* (Supplemental **Fig S3C**) were detected by RNA-seq in embryos collected 2-3 hours after fertilization, but *d*NET-seq signal over these genes was negligible. However, robust *d*NET-seq signal was found at the *pumilio* gene (Supplemental **Fig S3D**). Expression of this gene was considered to be exclusively maternal based on RNA-seq (Lott et al. 2011) and Pol II ChIP-seq data (Chen et al. 2013), but a GRO-seq study detected *pumilio* nascent transcripts in embryos collected 2-2.5 hours post-fertilisation (Saunders et al. 2013). The detection of *pumilio* RNA in early embryos by GRO-seq and *d*NET-seq highlights the sensitivity of these two techniques in capturing low-level nascent transcripts.

The stage at which the embryo switches from relying on maternally deposited mRNAs and proteins to undergoing its own transcription is termed the mid-blastula transition (MBT). However, a few genes (known as pre-MBT genes) become transcriptionally active before MBT (Kwasnieski et al. 2019). Strong *d*NET-seq signal was detected in early embryos over the bodies of pre-MBT genes such as *snail* (**Fig 3C**), *fushi tarazu* (**Supplemental Fig S3E**) and *odd skipped* (**Supplemental Fig S3F**). Altogether, we examined 117 previously identified pre-MBT genes and 35 maternal mRNAs (**Fig 3D and Supplemental Fig S3G**).

A characteristic feature of pre-MBT genes is that they are short in length with few if any introns (De Renzis et al. 2007; Ali-Murthy et al. 2013; Kwasnieski et al. 2019; Heyn et al. 2014). In contrast, most genes expressed in late embryos contain multiple introns, as shown for *Akap200* (**Fig 3E**), *His3.3A* (**Supplemental Fig S3H**) and *twinstar* (*tsr*) (**Supplemental Fig S3I**). Analysis of *d*NET-seq datasets

from late embryos revealed widespread signal on introns and regions downstream of the

polyadenylation site, further confirming that the technique is capturing nascent transcripts.

To conclude, robust *d*NET-seq signal was recovered from zygotically but not from maternally

expressed transcripts. In addition, we detected reads mapping to introns and downstream of the poly-

adenylation site. These findings suggest that *d*NET-seq is indeed specifically targeting the nascent

transcriptome.

dNET-seq reveals transcriptional readthrough associated with the presence of overlapping

antisense genes.

Having confirmed that *d*NET-seq was capturing nascent RNA, we next investigated the distribution of

Pol II density over transcript regions in early and late embryos. *d*NET-seq profiles (**Fig 3C-E and**

Supplemental Fig S3) do not show the characteristic higher read density near the promoter, as

previously described in mammalian cells (Nojima et al. 2015; Mayer et al. 2015). This is most likely

because Pol II typically pauses ~30-60 bp downstream of the transcription start site (TSS) (Kwak et al.

2013) and in our *d*NET-seq approach we enrich for RNAs longer than 60 nt (**Supplemental Fig S1F**);

thus, we only record the position of polymerases that have transcribed at least 60 bp past the TSS.

We then turned our attention to the *d*NET-seq signal around polyadenylation sites (also known

as the transcription end sites or TES). Many genes, such as *Akap200* (**Fig 3E**), *His3.3A* (**Supplemental**

Fig S3H) and *tsr* (**Supplemental Fig S3I**), have considerable levels of *d*NET-seq/S5P signal after the

TES, indicative that in these cases, Pol II continues to transcribe after cleavage and polyadenylation.

We noted that the 3' UTR of these three genes is overlapped by another gene (*gurken, Nepl3,* and *IntS1,*

respectively), which is transcribed in the opposite direction. We therefore asked whether transcriptional

readthrough is related to the presence of an overlapping convergent gene. In order to identify all the

genes that are transcriptionally active in late embryos, we used a strategy adapted from GRO-seq

analysis (Core et al. 2008) that relies on read density in gene desert regions as background reference for

absence of transcription. Very large intergenic regions (gene deserts) were divided into 50kb windows,

and read densities were calculated by dividing read counts in each window by the window length in bp

(**Supplemental Fig S4A**). Genes with *d*NET-seq signal over the gene body (in RPKM) above the 90th

percentile of read density for all intergenic regions analysed were considered to be transcriptionally

active (**Supplemental Fig S4B**). We identified approximately 7 thousand active genes, and similar

results were obtained from *d*NET-seq/S5P and *d*NET-seq/S2P datasets (**Supplemental Fig S4C**). This

set of active genes includes over 85% of the 3500 genes previously identified as actively transcribed

after the mid-blastula transition (MBT) based on ChIP-seq experiments (Chen et al. 2013). Next, we

divided the genes transcribed in late embryos into two groups, depending on whether their 3' UTR was

or was not overlapped by another convergent gene. The metagene analysis shows that when averaged

across all transcribed genes, the *d*NET-seq/S5P signal is abruptly reduced at the TES (**Fig 3G**). This is

expected, as when Pol II reaches this site, the nascent transcript is cleaved and polyadenylated and

therefore there is no RNA to be sequenced. The presence of nascent transcripts after the TES indicates

that Pol II fails to terminate transcription after cleavage and polyadenylation. Unexpectedly, this is

mainly observed on genes that have the 3' UTR overlapped by an antisense gene (**Fig 3G**). Similar

results were observed after metagene analysis of *d*NET-seq/S2P datasets (**Fig 3H**), confirming an

association between transcriptional readthrough and genomic architecture at the TES. We then asked

whether transcriptional readthrough depends on the transcriptional activity of the overlapping antisense

(convergent) gene. Using the methodology described above to identify transcribed genes, we found that

the vast majority (>80%) of overlapping convergent genes were transcriptionally active and only 219

genes were silent. The metagene analysis shown in **Fig 3I** clearly indicates that transcription of the

convergent overlapping gene is not required for inefficient termination at the overlapped TES.

The vast majority of pre-MBT genes do not show *d*NET-seq signal past the TES (**Fig 3C, 3F**

and Supplemental Fig S4D, E). Most likely, in these genes Pol II releases both the nascent RNA and

the template DNA within a few nucleotides from the polyadenylation site. Notably, for the majority of

pre-MBT genes devoid of *d*NET-seq signal past the TES (65), there is no other gene on either strand

within a region of 500 bp downstream of the TES (referred to henceforth as 'isolated genes'), as shown

for *tailless (tll)* (**Supplemental Fig S4D**). Another 26 pre-MBT genes are embedded in larger genes, as

shown for *nullo*, which is located within a long intron of the *CG12541* gene (**Supplemental Fig S4E**).

A smaller group of pre-MBT genes (21) have neighbouring genes located on either strand within a

region of 500 bp downstream of the TES, as shown for *Elba2* (**Supplemental Fig S4F**). We further

identified 5 pre-MBT genes that have the 3' UTR overlapped by an antisense convergent gene, as shown for *spook (spo)* (**Supplemental Fig S4G**). Transcriptional readthrough was observed in these pre-MBT genes (**Supplemental Fig S4G**). Importantly, the proportion of isolated genes is significantly higher among pre-MBT genes (65/117 or ~56%) than among late genes (1251/7233 or ~17% of all transcriptionally active genes for replicate 1 S5P) ($P < 2.2 * 10^{-16}$, one-tailed binomial test). This suggests that pre-MBT genes may be under selection to avoid overlaps with other genes, thereby minimizing transcriptional readthrough. Note, however, that we cannot exclude the possibility that in some cases, a lack of NET-seq signal past the TES could be due to loss of CTD phosphorylation at this position.

In conclusion, *d*NET-seq reveals a strong correlation between inefficient transcription termination at the cleavage and polyadenylation site, and presence of an antisense convergent gene overlapping the TES. The vast majority of genes that start to be transcribed before the mid-blastula transition, when the mitotic cycles are still very short, do not have the TES overlapped by other genes and lack *d*NET-seq signal past the TES. In contrast, many genes that start to be transcribed after the mid-blastula transition, when the mitotic cycles become much longer, show transcriptional readthrough and are overlapped by an antisense convergent gene.

Figure 3. Transcription over gene bodies and beyond the TES.
(A) The diagram illustrates the temporal expression of maternal, pre-MBT, MBT, and post-MBT genes during *Drosophila* embryonic development. *d*NET-seq/S5P and RNA-seq profiles over the maternal gene *bicoid* (**B**), the pre-MBT gene *snail* (**C**), and the post-MBT gene *Akap200* (**E**). Reads that aligned to the positive strand are in blue, and reads that aligned to the negative strand are in red. **(D)** Meta-analysis of mean *d*NET-seq read density around the transcription start site (TSS) in maternal and pre-MBT genes (replicate 1). **(F-I)** Normalized metagene analysis in arbitrary units (A.U.). The *d*NET-seq signal is depicted along the normalized gene length (grey background), as well as 500 bp upstream of the transcription start site (TSS) and 500 bp downstream of the transcription end site (TES). **(F)** *d*NET-seq/S5P signal on pre-MBT genes in early embryos. **(G)** *d*NET-seq/S5P signal on transcriptionally active genes in late embryos; the signal over genes that have the 3' UTR overlapped by an antisense gene is depicted in dark red, while the signal over genes with no other genes within 500 bp is depicted in light red. **(H)** *d*NET-seq/S2P signal on transcriptionally active genes in late embryos. **(I)** *d*NET-seq/S5P signal on transcriptionally active genes in late embryos; the signal over genes that have the 3' UTR overlapped by a transcriptionally active antisense gene is depicted in dark red, while the signal over genes with the 3' UTR overlapped by a transcriptionally inactive antisense gene is depicted in light red.

Analysis of *d*NET-seq read density profiles.

We next focused on the distribution of *d*NET-seq reads on exons and introns. The number of nascent

RNA reads whose 3' ends map at a particular genomic position is proportional to the number of Pol II

molecules at that position. Thus, Pol II pause sites can be detected as local peaks in read density

(Churchman and Weissman 2011; Larson et al. 2014). To search for *d*NET-seq peaks representative of

paused Pol II, we excluded signal resulting from splicing intermediates (i.e., reads that map to the very

last nucleotide of introns and exons were discarded). We analysed both S5P and S2P data. However,

we were particularly interested in exploring the multiple peaks of dNET-seq/S5P signal observed on

both exons and introns (**Fig 4A**), because co-transcriptional splicing has been linked to S5P in humans

(Nojima et al. 2015).

A difficulty when performing the systematic identification of *d*NET-seq peaks is that transcripts

with higher initiation rates will contain more reads and thus peaks are more likely to be detected than

in more lowly transcribed genes. To control for this confound, we developed a peak calling algorithm

that detects regions where the local read density is significantly higher than expected by chance, given

the over-all read density of the transcript (**Fig 4A;** see also the methods section). We emphasize that for

detecting reads corresponding to splicing intermediates (**Fig 2D-F**), we used a peak calling method that

looks for significant single-nucleotide positions (Churchman and Weissman 2011), whereas this new

method detects higher read regions of variable length.

We then aligned exons and introns on the splice sites and calculated the average peak density

at each position. Exons have a higher over-all peak density than introns (mean proportion of nucleotides

in peaks ~0.022/~0.015 for replicate 1/replicate 2 introns and ~0.043/~0.034 for replicate 1/replicate 2

exons; $P < 2.2 * 10^{-16}$ for both replicates; two-tailed Mann-Whitney U-test with the peak densities of

individual introns/exons as data points), suggesting that the elongation rate is decreased over exons (**Fig**

4C). This is consistent with previous reports in mammalian (Mayer et al. 2015; Jonkers et al. 2014) and

Drosophila cells (Kwak et al. 2013). In addition, peak density is sharply increased around the 5' splice

site (**Fig 4B**). This could indicate Pol II pausing associated with splice site recognition. However, we

cannot exclude that misaligned splicing intermediate reads are contributing to the observed increase in

peak density around the 5' splice site.

The profile around the 3' splice site shows a higher peak density region just after the intron-

exon boundary (**Fig 4B**). A progressive increase in average exonic peak density is further observed

starting roughly 60 nt after the 3' splice site (**Fig 4B**), suggesting slower elongation or more frequent

Pol II pausing in this region. Alternatively, such increases in peak density could correspond to regions

where the Pol II CTD more frequently exhibits the phosphoisoform targeted by the antibodies.

However, note that *d*NETseq/Ser5 and *d*NETseq/Ser2 lead to very similar peak density metaprofiles

(**Fig 4B**; **Supplemental Fig S7I-J**).

A further potential caveat is that nucleotide composition varies systematically across exons and

introns. For example, exons tend to have a higher GC content than introns (Zhu et al. 2009). This could

be problematic as NET-seq relies on MNase digestion of DNA and RNA to solubilize chromatin.

MNase digestion of DNA is known to be sequence-biased, with most notably a preference for cleaving

just 5' of an adenine (Dingwall et al. 1981; Hörz and Altenburger 1981; Gaffney et al. 2012). An

analysis of the 5' ends of our reads revealed similar biases for MNase digestion of RNA (**Fig 4D**). This

sequence preference could lead to artefactual variation in read density, with more reads being sampled

from transcripts and transcript regions whose nucleotide composition is more similar to MNase

digestion biases. To verify to what extent our results were affected by this confound, we performed a

simulation to determine the expected distribution of reads based on the digestion bias alone

(Supplementary Methods). We concluded that MNase biases are unlikely to explain either the

enrichment of peaks in exons or the general profile of peak densities past the 3' splice site.

Figure 4. Analysis of *d*NET-seq read density profiles.
(**A**) *d*NET-seq/S5P and RNA-seq profiles over the post-MBT gene *smoke alarm* (*smal*). The read
number is depicted at two magnification levels in two biological replicates. For replicate 1, the line
"Peak Caller 1" shows peaks called using the "large peaks" setting, which is appropriate for detecting
larger regions of putative Pol II pausing. The line "Peak Caller 2" shows peaks called using the "small
peaks" setting, which provides higher spatial resolution and has been used for subsequent analyses.
RNA-seq data for the same regions is also shown. (**B**) Metagene analysis of peak density estimated
from *d*NET-seq/S5P data from late embryos (replicate 1, N=12799). To calculate peak density for each
position, we divided the number of introns that overlap with a peak at that position by the total number
of introns. The last 50 nucleotides of exons, the first 50 nucleotides of introns, the last 25 nucleotides
of introns, and the first 100 nucleotides of exons are shown. Only internal and fully coding exons from
transcriptionally active genes that are at least 100 nucleotides long are shown. Exons shorter than 150
nucleotides contribute to both the exon end and start. Only introns that were at least 50 nt long were
considered. (**C**) Peak density in the exons and introns of transcriptionally active genes (*d*NET-seq/S5P,
replicate 1). Peak density has been defined as the percentage of nucleotides within a given exon or
intron that overlap with a significant peak. (**D**) Sequence logo of nucleotide frequencies within a 10-nt
window around the 5' ends of NET-seq reads. The combined height of the bases at each position is
proportional to the information content. Position 6 corresponds to the 5'-most nucleotide of the read.
Putative internal priming reads, as well as reads mapping to the last nucleotide of exons or introns
(possible splice intermediate and intron lariat reads) were ignored.
402

404 **dNET-seq captures recursive splicing intermediates.**

Having shown that the spliceosome forms a complex with the elongating Pol II in *Drosophila* embryos, we asked when splicing takes place relative to transcription. We first looked at recursive splicing of long introns because this process involves the formation of inherently unstable intermediates that are more likely to be formed soon after the transcription of each intronic splice site (Pai et al. 2018). In recursive splicing, long introns are removed by sequential excision of adjacent sections involving separate splicing reactions, each producing a distinct lariat (Hatton et al. 1998). Recursively spliced intron segments are bounded at one or both ends by recursive sites or ratchet points (Burnette et al. 2005), which correspond to zero nucleotide exons consisting of juxtaposed 3' and 5' splice sites around a central AG|GT motif, where the vertical line represents the splice junction (**Fig 5A**).

To capture recursive splicing intermediates using *d*NET-seq, it is essential to have a good coverage of reads corresponding to nascent transcripts and spanning the splice junctions. The total number of reads resulting from nascent RNA in each *d*NET-seq dataset is depicted in **Supplemental Fig S1G**. By merging the sequencing information of overlapped paired-end reads (**Fig 1E**), we were able to sequence on average ~103 nucleotides per nascent RNA (**Supplemental Fig S1F and S1H**). Focusing on previously identified *Drosophila* ratchet points (Joseph et al. 2018; Duff et al. 2015), we found *d*NET-seq/S5P reads that span the junction between the canonical 5' splice site at the end of the exon and the first ratchet point (RP1) internal to the downstream intron, as shown for the second intron of the *Megalin* gene (**Fig 5B**). Reads spanning the subsequent intronic RPs were also observed (**Fig 5B**). Overall, we detected *d*NET-seq/S5P and *d*NET-seq/S2P spliced reads supporting most of the previously identified recursive splicing events (**Fig 5C and Supplemental Fig S5A**).

Analysis of *d*NET-seq profiles around a RP reveals an enrichment of reads in a region located a few nucleotides downstream of the RP, as shown for RP2 in the first intron of the *Tenascin major* gene (**Fig 5D**). Noteworthy, most of these reads are already spliced to the previous RP (**Fig 5D**). A meta-analysis of *d*NET-seq/S5P reads around 137 RPs confirms that many spliced reads can be observed just downstream of RPs (**Fig 5E, F**). Taken together, these results indicate that recursive splicing can occur soon after the transcription of intronic recursive sites and suggest the presence of paused Pol II downstream of these sites.

Figure 5. *d*NET-seq captures recursive splicing intermediates.
(A) Schematic illustrating recursive splicing. A ratchet point (RP) with juxtaposed acceptor and donor splice site motifs is indicated. **(B)** Visualization of *d*NET-seq/S5P reads that align to the second intron of the *Megalin* (*mgl*) gene. Recursively spliced reads align to exon 2 (green) and the intron after RP1 (dark red). The number of spliced and unspliced reads at each RP in the intron is indicated. **(C)** Venn diagram comparing RPs identified in two *d*NET-seq/S5P biological replicates and in previously reported studies (Joseph et al. 2018; Duff et al. 2015). **(D)** Number of *d*NET-seq/S5P reads that have the 3' end mapped around RP2 in the first intron of *Tenascin major* (*Ten-m*) gene. The top panel depicts all reads, and the bottom panel depicts only reads that have been spliced to RP1. **(E-F)** Meta-analysis with single nucleotide resolution of normalized *d*NET-seq/S5P reads around RPs (n=137) using all reads **(E)** or only reads spliced to the previous RP or exon **(F).**

Splicing takes place as Pol II transcribes past the 3' splice site.

Having established that *d*NET-seq captures recursive splicing, we then asked whether *d*NET-seq reads spanning canonical exon-exon junctions were also detected (**Fig 6A**). To identify splicing events, we considered all internal and fully-coding exons that are at least 100 nt long in actively transcribed genes in early and late embryos. For each splice junction, we counted how many reads had the 3' end mapped to the first 100 nt of the exon. Only exons with at least 10 reads mapping to this region were considered (see the methods section for justification of the threshold). Then, the *d*NET-seq splicing ratio (SR) was calculated by dividing the number of spliced reads by the sum of the number of spliced and unspliced reads (**Fig 6A**). Reads could be counted as spliced or unspliced if their 3' end mapped to within the first 100 nt of the exon and their 5' end reached upstream of the 3' splice site, allowing to check whether the intron was still present. A robust agreement of estimated SR values was observed between biological replicates for pre-MBT genes (**Supplemental Fig S6A**) and genes expressed in late embryos (**Fig 6B**).

SR values showed a bimodal distribution, with peaks at both extremes (SR = 0 and SR = 1; **Fig 6C** and **Supplemental Fig S6B**), indicating that a subset of junctions were always spliced immediately after transcription (SR=1), while others remained unspliced (SR=0). Notably, differences in SR distribution were found between *d*NET-seq/S5P and *d*NET-seq/S2P datasets (**Fig 6C, D and Supplemental Fig S6B, C**). Significantly lower SR values were detected for S2P compared to S5P ($P < 2.2*10^{-16}$ from binomial regression, see Fig 6 legend for statistical details). In particular, junctions that were most frequently spliced (SR values close to 1) were predominantly captured by *d*NET-seq/S5P (**Fig 6C and Supplemental Fig S6B**). This clearly points to a preferential association between co-transcriptional splicing and S5 phosphorylation of Pol II CTD, as previously proposed in human cells (Nojima et al. 2015, 2018). We therefore continued to focus solely on the S5P datasets in the remainder of our splicing analysis.

We observed that only ~5% of splice junctions in pre-MBT genes were devoid of S5P reads spanning ligated exons and thus presented an SR of 0 (~5.10% replicate 1/~4.35% replicate 2), whereas in genes expressed in late embryos this proportion was ~20% (~19.72% replicate 1/~18.55% replicate 2) (**Fig 6E**; two-tailed binomial test for difference between late and pre-MBT, $P \sim 6.141 * 10^{-5}/9.44 * 10^{-4}$ (replicate 1/replicate 2)). However, this difference between pre-MBT genes and genes expressed

in late embryos disappeared once the higher read density of pre-MBT genes had been controlled for

(**Fig 6E**).

Thus, in most cases, *Drosophila* co-transcriptional splicing can occur when Pol II is still

transcribing the downstream exon, implying an intron definition mechanism as previously proposed for

S. cerevisiae (Carrillo Oesterreich et al. 2016). Consistent with this view, many *Drosophila* transcripts

have relatively long exons separated by short introns – a gene architecture suggested to be conducive

to intron definition. Very long introns flanked by short exons have instead been associated with exon

definition (under which the downstream exon needs to be fully transcribed before splicing can take

place) (Keren et al. 2010). A switch to exon definition was proposed once the size of the intron surpasses

~200 nt (Fox-Walsh et al. 2005). It is unclear, however, whether the choice between exon and intron

definition is dependent on the absolute sizes of exons and introns, or rather the ratio of intron to exon

size.

We found no relationship between the *d*NET-seq splicing ratio and exon length (**Supplemental**

Fig S6D, H for replicates 1 and 2; see **Supplementary Table 1** for details on statistical significance

and sample sizes for all of the gene architecture parameters discussed here). Regarding intron size, we

found that introns with SR = 0 (and thus no evidence for intron definition) were, on average, indeed

larger than other introns (**Fig 6F** and **Supplemental Fig S6G**). However, more careful examination

revealed a more complex picture, with a lower proportion of introns with SR = 0 both for introns of

intermediate size (~55-100 nt, which corresponds to ~55% of the introns studied) and for very large

introns (>1000 nt) (**Fig 6F**). These intron sizes may thus be optimal for fast splicing. We also uncovered

a lower exon to intron length ratio for introns with SR = 0 than for others (**Supplemental Fig S6E, I**;

note that the difference was significant for replicate 2 and near-significant for replicate 1).

Taken together, these results suggest that although fast splicing (implying intron definition) is

indeed skewed towards small introns flanked by large exons, there is also frequent and efficient intron

definition for large introns. Our results are inconsistent with a threshold model, where splicing would

systematically switch to exon definition after a given intron size is reached.

We also investigated the relationship between SR and several other gene architecture

parameters. Firstly, as the GC content in exons and introns decreases, the proportion of introns with SR

= 1 increases and the proportion with SR = 0 decreases, showing more efficient immediate splicing (**Fig 6G, H** and **Supplemental Fig S6K, L**). A similar effect is observed as the ratio of the downstream exon GC content to intron GC content increases (**Fig 6I** and **Supplemental Fig S6M**). Thus, the most efficient immediate splicing is observed when the GC content is low in both exons and introns but higher in exons than introns. Secondly, as 3' splice site strength increases, the proportion of introns with SR = 0 decreases (**Fig 6J** and **Supplemental Fig S6N**). We found no significant relationship between SR and 5' splice site strength (**Supplemental Fig S6F, J**). However, we replicated previous observations (Khodor et al. 2012) that transcripts with only a single intron tend to be spliced less efficiently than multi-intron ones, with a higher proportion of SR = 0 introns (**Supplemental Fig S6O, P**). Thirdly, similarly to previous reports (Khodor et al. 2011, 2012; Herzel et al. 2018), we uncovered an effect of exon rank, whereby exons that are more central appear to be spliced more efficiently (**Fig 6K** and **Supplemental Fig S6Q**). Finally, in light of recent results in mouse linking transcriptional readthrough and inefficient splicing (Reimer et al. 2021), we asked whether such a relationship could be detected in our data. We found no significant difference in levels of readthrough between genes with high or low mean SR (**Supplemental Fig S6R, S**).

In conclusion, *d*NET-seq reveals that splicing can occur immediately as Pol II transcribes past the 3' splice site, yet many nascent transcripts remain unspliced. As expected, we found a relationship between the proportion of spliced reads and intron size. However, immediate splicing was observed at all intron sizes.

Figure 6. *d*NET-seq reveals immediate splicing at all intron sizes.
(A) Visualization of *d*NET-seq/S5P reads that align to exon 10 of the *Tao* gene. For **(B-J)**, unless otherwise specified, only introns from transcriptionally active genes where the downstream exon is a fully coding internal exon at least 100 nt long were included. In addition, enough spliced/unspliced reads had to end within the first 100 exonic nucleotides that obtaining a splicing ratio (SR) of 0 or 1 by chance alone was highly unlikely (see the methods section for details). For genes expressed in late embryos, this threshold was 10 reads for both replicates. For pre-MBT genes, it was 14 for replicate 1 and 9 for replicate 2. For S2P data, we used a threshold of 10 to enable better comparison with S5P. **(B)** SR values estimated in two biological replicates of *d*NET-seq/S5P datasets from late embryos

(Spearman correlation, ρ = ~0.734, $P < 2.2 * 10^{-16}$; N = 3708). **(C)** Histogram of SR values for dNET-seq/S5P (N = 5626) and dNET-seq/S2P (N = 6888). To test the significance of the difference between S5P and S2P, a binomial regression with a logit link was performed without filtering by read number (N = 12833 for S5P; N = 13229 for S2P). The number of spliced and unspliced reads was specified as the dependent variable and the status of each data point as S5P or S2P was the sole predictor. The model predicted a splicing ratio of ~0.369/~0.394 for S5P replicate 1/2 and of 0.162/0.269 for S2P replicate 1/2. **(D)** SR values estimated in replicate 1 of dNET-seq/S5P and dNET-seq/S2P datasets from late embryos (Spearman correlation, ρ = ~0.666, $P < 2.2 * 10-16$; N = 4773). **(E)** Proportion of splice junctions in pre-MBT genes and genes expressed in late embryos classified according to their SR values. As many pre-MBT genes are single-intron, last introns were exceptionally included in this analysis. To make the two columns on the right, we used a subset of the genes expressed in late embryos (post-MBT genes) that was as similar as possible to the pre-MBT set in read number. Concretely, to match each pre-MBT gene, we picked the post-MBT gene that had the most similar total count of spliced and unspliced reads, making sure that every post-MBT gene only appeared in the subset once. **(F-K)** Several parameters of gene architecture show a relationship with SR. For the sample sizes and statistical tests used, see **Supplementary Table 1**. Note that in **(F, J-K)**, the bin ranges have been set so that intron numbers would be as equal as possible between bins.

Immediate splicing associates with specific read density profiles.

Analysis of individual dNET-seq profiles around constitutive splice junctions with a high splicing ratio frequently revealed an enrichment of reads downstream of the 3' splice site, coincident with the appearance of spliced reads, as shown for the *cno* gene (**Fig 7A**). In contrast, profiles around a constitutive splice junction with SR = 0 had an accumulation of reads further along the exon, as shown for the *ND-51* gene (**Fig 7B**). A clearly distinct type of profile was observed on skipped exons, on which very few reads were observed, as shown for the *zip* gene (**Fig 7C**).

We next performed a meta-analysis of dNET-seq peak densities over different exonic and intronic regions for introns with differing SR values. A difficulty of analysing the peak density profile around the 3' splice site is that reads mapping to the final nucleotide of the intron may represent intron lariats and thus be non-nascent. We removed reads mapping to this position prior to peak calling. However, through misalignment, intron lariat reads may also map to the few nucleotides around the 3' splice site and thus still affect the final meta-profile. In order to minimize the impact of such misalignment events, we excluded from the meta-analysis all introns with a read mapping to the final intronic nucleotide, as this is expected to also discard the introns most likely to contain misaligned intron lariat reads (**Fig 7D, Supplemental Fig S7A**). When such filtering is not done, a sharp peak is observed around the 3' splice site, notably when SR values are low (**Supplemental Fig S7 B-C**). It

appears that intron lariat reads are primarily captured for introns with low splicing ratios (**Fig 7E and Supplemental Fig S7D**). This contrasts with splicing intermediate reads from the end of the upstream exon, which are associated to higher splicing ratios instead (**Fig 7E and Supplemental Fig S7D**).

The peak density profile downstream of the 3' splice site contains two primary regions of interest (highlighted in **Fig 7D and Supplemental Fig S7 A-C, G-H**). The first is a region of increased peak density located roughly 10-20 nucleotides after the 3' splice site. This peak is independent of the presence of putative intron lariat reads (**Fig 7D, Supplemental Fig S7 A-C**), and is thus likely to correspond to a genuine region of Pol II pausing or slow-down. However, it could also represent a region of preferential Ser5 phosphorylation, with no concomitant pausing. Noteworthy, pausing in this region has also been observed in studies that did not target specific Pol II phosphoisoforms (e.g. (Kwak et al. 2013)). Interestingly, this peak is more prominent when the SR is higher (**Fig 7D, Supplemental Fig S7 A-C, G-H**), suggesting an association between immediate co-transcriptional splicing and increased Pol II pausing at the start of the exon.

The second feature of interest is a rapid increase in peak density ~60 nucleotides into the exon. Contrary to the first region of interest, the peak density in this region is higher when SR is *lower* (**Fig 7D, Supplemental Fig S7 A-C, G-H**). We found no evidence that this increase in peak density could be a result of MNase digestion biases (**Supplementary Methods**). A further hypothesis is that this pattern could be reflecting a gradual slowdown of Pol II as it moves from an AT-rich intronic region to a GC-rich exonic one. The peak is indeed more prominent in introns where the ratio of exonic to intronic GC content is higher (**Supplemental Fig S7 E-F**). However, when SR=1, then even exons where this ratio is high do not display the peak (**Supplemental Fig S7 E-F**). Moreover, introns with higher SR tend to have *higher* exonic to intronic GC ratios (**Fig 6I, Supplemental Fig S6M**). Hence, the co-variation with SR cannot be explained simply through the correlation between SR and GC content patterns. This suggests that there is an effect of the SR beyond any GC content effect. Potentially, this region could correspond to where delayed splicing starts (because these events have low immediate splicing ratios), and could represent splicing-associated Pol II pausing.

Note that we obtained qualitatively similar meta-profiles when considering read rather than peak densities (**Supplemental Fig S7 G-H**). Our results therefore cannot be explained through any

599   biases introduced by the peak calling approach. The meta-profile obtained using *d*NET-seq/S2P data

600   was also qualitatively similar, although we did not divide the introns by SR given the rarity of reads

601   with high SR in S2P data (**Supplemental Fig S7 I-J**). We also attempted a separate meta-analysis of

602   skipped exons. These exons showed no accumulation of Pol II. However, the sample size was too small

603   for meaningful comparisons with included exons.

604   Finally, we asked how prevalent co-transcriptional splicing is in the developing *Drosophila*

embryo. As shown in **Fig 2E**, approximately 80% of exons in pre-MBT genes and genes expressed in

late embryos covered by a high density of NET-seq reads have a peak corresponding to the splicing

intermediate formed after cleavage at the 5' splice site but before exon-exon ligation. The majority of

these exons (~85%) were covered by *d*NET-seq reads that span the junction to the downstream exon

either directly on nascent transcripts or indirectly on splicing intermediates formed by cleavage at the

5' splice site of the downstream exon **(Fig 7F, H)**, confirming that they are co-transcriptionally spliced.

We then focused on those exons for which no splicing intermediate spike was detected by the peak

calling algorithm, as illustrated for the *Doc3* gene **(Fig 7G, I)**. Over 71% of these exons were also

covered by *d*NET-seq reads that span the junction to the downstream exon either directly on nascent

transcripts or indirectly on splicing intermediates formed by cleavage at the 5' splice site of the

downstream exon **(Fig 7I)**, arguing that even exons without a detectable splicing intermediate peak are

co-transcriptionally spliced. Altogether, our *d*NET-seq results support co-transcriptional splicing for

over 95% of the analysed exons.

Figure 7. Immediate splicing associates with higher density of *d*NET-seq signal.
(A-C) *d*NET-seq/S5P profiles surrounding the indicated exons in the post-MBT genes *cno* **(A)**, *ND-51*
(B), and *zip* **(C)**. The top panels depict all reads. The bottom panels depict either the 3' end coordinate
of reads that span the splice junction **(A, B)**, or the RNA-seq profile **(C)**. **(D)** Metagene analysis of peak
density estimated from *d*NET-seq/S5P datasets from late embryos (replicate 1) for different ranges of
SR values. Peak density has been calculated as the proportion of introns that overlap with a peak at any
given position. The last 50 nucleotides of exons, the first 50 nucleotides of introns, the last 25
nucleotides of introns, and the first 100 nucleotides of exons are shown. Only internal and fully coding
exons from transcriptionally active genes that are at least 100 nucleotides long are shown (N = 4783).
In addition, at least 10 spliced/unspliced reads had to end within the first 100 nucleotides of the exon.
(E) The proportion of introns with at least one read whose 3' end maps to the final position of the

upstream exon (putative splicing intermediates) or to the final position of the intron (putative intron lariats) in *d*NET-seq/S5P late replicate 1. **(F, G)** *d*NET-seq/S5P profiles on the indicated regions of the *velo2* and *Doc3* genes. Below, spliced reads are depicted. Asterisks denote the splicing intermediate peaks at the end of exons. **(H, I)** Venn diagrams showing how many junctions with or without a splicing intermediate peak are covered by spliced reads or have a downstream splicing intermediate covered by spliced reads.

DISCUSSION

In this study, we used *d*NET-seq to map Pol II over the bodies of genes that become transcriptionally active in *Drosophila* embryos during the initial stages of development. The use of embryos allowed us to perform an important test to verify that the captured RNA is truly nascent. Indeed, early *Drosophila* embryos contain abundant mRNAs that are transcribed during oogenesis and loaded into the egg. In our analysis of early embryos, these maternal transcripts were readily detected by RNA-seq, but not by *d*NET-seq.

Transcription in the *Drosophila* early embryo begins in mitotic cycle 8 for a few genes (Erickson and Cline 1993; Pritchard and Schubiger 1996) and then the number of active genes gradually increases until cycle 14 (Kwasnieski et al. 2019). Notably, the initial mitotic cycles have a duration of approximately 10 minutes, with cycle 13 taking about 21 minutes and cycle 14 lasting for at least 65 minutes (Ji et al. 2004). Thus, assuming an elongation rate of 2.4-3.0 kb per minute (Fukaya et al. 2017), there are significant constraints on the transcription of genes that are active before the end of cycle 13 (pre-MBT genes). Although pre-MBT genes are on average shorter than genes expressed at later stages of development (Artieri and Fraser 2014; Hoskins et al. 2011), longer genes are nevertheless transcribed before MBT. However, before cycle 14, transcriptional elongation of long genes is either prematurely terminated (Sandler et al. 2018) or aborted (Shermoen and O'Farrell 1991; Kwasnieski et al. 2019). It has also been reported that pre-MBT expression is associated with the generation of DNA damage due to stalling of DNA replication at transcriptionally engaged loci (Blythe and Wieschaus 2015).

Our *d*NET-seq analysis revealed a novel feature of pre-MBT genes that most likely contributes to reduce conflicts between transcription and replication during the short interphases of early embryos. We found that in the majority of pre-MBT genes, no *d*NET-seq signal is detected after the TES, whereas many of the genes that are only expressed after MBT show high *d*NET-seq signal past the TES. We

also observed that most pre-MBT genes are isolated from other transcriptional units, while the TES of

many post-MBT genes is overlapped by another gene transcribed in the opposite direction. Taken

together, these results suggest that *Drosophila* genome architecture may be under selection for efficient

transcription termination of pre-MBT genes. Our observations further reveal an unexpected link

between transcriptional readthrough and the presence of an overlapping convergent gene in the

Drosophila genome. The finding that readthrough is detected even when the convergent overlapping

gene is not transcribed is intriguing. One possibility is the induction of local conformational changes in

chromatin and/or Pol II, Indeed, pioneer transcription factors can potentially regulate gene expression

at later stages of development by inducing significant chromatin conformational changes during early

embryonic development (Blythe and Wieschaus 2016; Schulz et al. 2015). Although such chromatin

remodelling events can potentially influence transcriptional read-through, regardless of convergent

gene transcription, further studies are needed to understand how transcription termination is regulated

during early *Drosophila* development.

A classical view in the splicing field is that for some introns, the 5' and 3' splice sites are

recognized directly by 'intron definition'. For other introns, spliceosome assembly starts with the

recognition of the downstream exon, and only at a later stage a cross-intron complex is formed ('exon

definition') (Robberson et al. 1990; Berget 1995; Sterner et al. 1996; Fox-Walsh et al. 2005). These

alternative models are supported by *in vitro* evidence, as well as by *in vivo* reporter gene experiments

(Robberson et al. 1990; Berget 1995; Fox-Walsh et al. 2005; De Conti et al. 2013). There is also indirect

evidence, for example from analysis of exon-intron size patterns (Berget 1995). However, these models

have never been directly tested *in vivo* at a transcriptome-wide level. It thus remains uncertain how

intron definition and exon definition operate *in vivo* and what determines the choice between them.

Introns that show splicing completion before Pol II has reached the end of the downstream exon

are necessarily intron-defined, as exon definition requires the presence of the 5' splice site of the

downstream intron. Therefore, introns with an SR above 0 represent an experimentally-determined set

of introns that are spliced *via* intron definition at least part of the time, allowing us to test long-held

assumptions in the field. Notably, *in vitro* and reporter gene work suggested that *Drosophila* uses intron

definition for introns smaller than ~200-250 nt and exon definition for larger introns (Fox-Walsh et al.

2005). Other studies have suggested that the crucial factor is not just intron size but the relative size of

the intron compared to the flanking exons (Sterner et al. 1996).

Using dNET-seq, we have found that although there is a relationship between intron size and

the proportion of introns with SR > 0, intron definition can be observed at all intron sizes. Our results

are thus inconsistent with a simple "threshold" model, where splicing would systematically switch to

exon definition at a particular intron size. Rather, we observe an increased proportion of introns with

SR > 0 within an optimal intron size range of ~55-100 nt. The median intron size in our data set is 71

nt. It thus appears that intron definition is best optimized for introns of a "typical" size, and is less

efficient when introns are unusually small or large. Intriguingly, however, for very large introns (>1000

nt), the proportion with SR > 0 increases again, suggesting that specific mechanisms may have evolved

for the intron-defined splicing of very large introns. The tendency for faster splicing kinetics with

average intron sizes has also been reported using metabolic labelling in both *Drosophila* (Pai et al.

2017) and human cells (Windhager et al. 2012).

We propose that the tethering of the upstream exon to Pol II (de Almeida and Carmo-Fonseca

2008) could be crucial for the intron-defined splicing of such large introns. Indeed, introns with SR > 0

where the upstream exon has one or more putative splicing intermediate reads tend to be larger than

ones without splicing intermediate reads (one-tailed Mann-Whitney U-test, W = 779623, P ~ 0.003, N

= 4589; exon size and read number filtering has been applied like for the gene architecture analysis).

In addition, Pai et al. (2017) found that GC poorer introns were spliced faster. We uncovered a

similar relationship with GC content, with the highest proportion of introns with SR > 0 observed when

both the exon and the intron were AT-rich but the exon had a higher GC content than the intron. We

emphasize that the gene architecture parameters that we have investigated are not independent of each-

other. For instance, exon GC content is correlated negatively with exon rank from start (Spearman's ρ

~ -0.098; P ~ 1.196 * 10^{-12}). Further work is thus needed to distinguish between causative and merely

correlative factors. In addition, it must be determined how the GC content effect relates to patterns of

nucleosome density (Gelfman et al. 2013).

Our results indicate that the majority (>80%) of *Drosophila* introns expressed in late embryos

can be spliced through intron definition. . A caveat of our work is that introns that show no evidence

for splicing completion during transcription of the downstream exon could still be spliced *via* intron

definition, if their splicing takes too long for it to be detected during the 100 nt window studied.

Therefore, we do not know whether introns with a SR of 0 are exon-defined, intron-defined but spliced

when Pol II has elongated past the first 100 exonic nucleotides, or a mixture of both. Similarly, introns

with a splicing ratio between 0 and 1 could either always be intron-defined, or they could use either

exon or intron definition depending on the splicing event.

Taken together, our results show that splicing in *Drosophila* embryos can be completed very

rapidly after transcription of the 3' splice site, as previously reported in other cellular models (Carrillo

Oesterreich et al. 2016; Reimer et al. 2021; Martin et al. 2013). Furthermore, our findings are relevant

to controversies regarding Pol II pausing in association with splicing. Indeed, splicing-dependent Pol II

pausing near the 3' splice site was first reported in yeast (Alexander et al. 2010; Chathoth et al. 2014).

Evidence for Pol II pausing at exon boundaries was further detected in mammalian cells using NET-

seq (Nojima et al. 2015; Mayer et al. 2015). Notably, NET-seq density profiles suggestive of Pol II

pausing were observed with S5P antibodies but not with antibodies to unphosphorylated CTD (Sheridan

et al. 2019; Nojima et al. 2015). In contrast, a recent PRO-seq analysis found no splicing-associated Pol

II pausing (Reimer et al. 2021).

Finally, we emphasize that our analysis of nascent *Drosophila* transcripts associated with Pol

II S5P CTD revealed highly heterogeneous splicing dynamics. Indeed, while only a minority of splice

junctions were devoid of reads spanning ligated exons (corresponding to an SR of 0), many introns

remained unspliced in our analysis. Is it relevant for cells whether any particular intron is rapidly

excised after Pol II transcribes the 3' splice site or is left unspliced while the transcript elongates? Is

this a stochastic process? What are the consequences for gene expression of fast versus delayed

splicing? Addressing these questions will likely dependent on further advances in methodologies to

study splicing in time and space at a transcriptome-wide level.

METHODS

Embryo collection

Drosophila melanogaster flies (Oregon R (OrR) strain) were raised at 25°C, in polypropylene

vials containing standard enriched culture medium (cornmeal, molasses, yeast, soya flour, and beetroot

syrup). Three-day-old flies (counting from pupae eclosion) were fattened in culture medium

supplemented with fresh yeast for 2 days. Embryos were collected into apple juice-agar plates

supplemented with fresh yeast using appropriate cages containing approximately 200 flies each. To

avoid female retention of older embryos, three pre-collections of 30min each were made before the first

collection of embryos. To maximize egg laying, and avoid larvae contamination, adult flies were

transferred to clean embryo collection cages every day over five days. For early stage embryos (2-3

hours after egg-laying), adult females were allowed to lay eggs for 1 hour in apple juice-agar plates.

Plates were subsequently collected and embryos were aged at 25°C for 90 min. During the following

30 min, embryos were harvested from the plates, dechorionated in 50% bleach solution for 2 min, and

washed once in Phosphate Buffered Saline supplemented with 0.1% Tween-20 (PBT) and twice in

deionized water. In order to discard older embryos (stage 6 and older), manual staging of collected

embryos was performed with the help of forceps and under a magnifier scope. Embryos were then

resuspended in a solution containing 120mM NaCl and 0.04% Triton, and washed twice with 120mM

NaCl solution. At the end of the 3 hours collection, the solution was removed and embryos were frozen

in liquid nitrogen and stored at -80ºC. For the late stage (4-6 hours), eggs were laid for 2 hours and aged

at 25°C for 3.5 hours. Embryo collection and processing was similar to the early stage embryos, but in

this case, no manual staging was performed.

Embryo DNA staining

For each embryo collection, and after dechorionation, a representative embryo sample was

collected from the total pool and fixed in a scintillation flask, using a solution containing 1 volume of

4% formaldehyde in PBT and 4 volumes of heptane, for 20 min at 100rpm. The lower aqueous phase

solution was subsequently removed, 4ml of methanol was added and embryos were shaken vigorously

during 1 min. Embryos were then collected from the bottom of the scintillation flask, washed twice

with methanol, and frozen at -20ºC in methanol. To rehydrate the embryos, they were washed for 5

min each, with 3:1, 1:1, and 1:3 mix solutions of methanol:PBT. Embryos were subsequently washed

twice in PBT and incubated with 1:5000 Sytox green (Invitrogen), supplemented with 5μg/ml RNase

A (Sigma-Aldrich) in PBT for 15 min. After washing with PBT, embryos were mounted in fluorescence

mounting medium (Dako) and examined in a Zeiss AxioZoom V16 Fluorescence Stereo Microscope

for image acquisition and embryo staging. Images were processed using ImageJ software (NIH).

*d*NET-seq and library preparation

The *d*NET-seq protocol was adapted from mNET-seq (Nojima et al. 2016). Briefly, 300 ul of

frozen embryos was resuspended in 3.5ml of Buffer B1 (15mM HEPES-KOH, pH 7.6; 10mM KCl;

5mM MgCl2; 1mM DTT; 0.1mM EDTA; 0.35M Sucrose; 4μg/ml Pepstatin; 10mM Sodium

Metabisulfite; 0.5mM EGTA supplemented with complete EDTA free protease inhibitor (Roche) and

PhoSTOP (Roche)). Embryos were homogenised in a Dounce homogenizer with 11x strokes using a

tight pestle on ice. The suspension was centrifuged at 7700g for 15 min at 4ºC, the supernatant was

discarded and the white pellet containing the nuclei was resuspended in 500 μl of Buffer B1. The

suspension was again homogenised in the Dounce with 4x strokes and loaded without mixing on the

top of buffer B2 (15mM HEPES-KOH, pH 7.6; 10mM KCl; 5mM MgCl2; 1mM DTT; 0.1mM EDTA;

0.8M Sucrose; 4μg/ml Pepstatin; 10mM Sodium Metabisulfite; 0.5mM EGTA supplemented with

complete EDTA free protease inhibitor (Roche) and PhosSTOP (Roche)). The suspension was

centrifuged at 1310g during 30 min at 4ºC, and the pellet was resuspended with 125 μl of NUN1 buffer

(20mM Tris-HCl (pH 7.9); 75 mM NaCl; 0.5mM EDTA and 50% Glycerol). 1.2ml of Buffer NUN2

(300mM NaCl, 7.5mM MgCl2, 1% NP-40, 1M Urea supplemented with complete EDTA free protease

inhibitor (Roche) and PhoSTOP (Roche) was mixed with the nuclei and incubated on ice during 15 min

preforming short vortex every 3 min. Chromatin was then centrifuged at 10000g for 10 min at 4ºC,

washed with 100 µl of 1x MNase Buffer and incubated in 100µl of MNase reaction mix (1x MNase

buffer and 30 gel unit/µl MNase (New England Biolabs)) during 3 min at 37ºC with mix at 1400rpm.

The reaction was stopped with 10 µl of 250mM EGTA, centrifuged at 10000g for 5 min at 4ºC, and the

supernatant containing the solubilized chromatin was recovered. For the early embryos sample, 2x

300µl of embryos were prepared in parallel and pooled together after the chromatin solubilization.

Immunoprecipitation of Pol II-RNA complexes was performed using 50µl of Protein G Dynabeads

(Thermo Fisher Scientific), pre-incubated over night with 5µg of the correspondent antibody: anti-Pol

II CTD S5P (ab5131 Abcam) or anti-Pol II CTD S2P (ab5095 Abcam) in 100µl NET2 (50mM Tris-

HCl pH 7.4; 150mM NaCl and 0.05% NP-40) and washed 3 times with NET2. Beads were incubated

with the solubilized chromatin in 1ml total volume of NET2 during 1 hour at 4ºC, washed 7 times with

500µl of NET2 and once with 100µl of PNKT (1x PNK buffer and 0.1% tween) before incubation

during 6 min in 50µl of PNK reaction mix (1x PNKT, 1 mM ATP and 0.05 U/ml T4 PNK 3′phosphatase

minus (NEB) in a thermomixer at 37ºC and 1400rpm. After washing the beads with NET2, long RNA

fragments were isolated using Quick-RNA MicroPrep (Zymo research): 300µl of RNA Lysis Buffer in

33% EtOH was mixed to the beads by pipetting. Beads were discarded and the suspension was loaded

into a Zymo spin column that was centrifuged at 10000g during 30sec. The column was washed once

with 400µl RNA prep buffer and twice with 700µl and 400µl RNA wash buffer respectively. RNA was

then eluted in 15µl of DNase/RNase-Free water (zymo) and stored at -80ºC. 100ng of RNA was used

to prepare each library, following the standard protocol of the Truseq small RNA library prep kit

(Illumina). After adapter ligation and reverse transcription, the libraries were PCR amplified using 16

PCR cycles and cDNA libraries were fractionated in the gel between 130 to 300bp. The libraries were

sequenced using PE-150 on the Illumina HiSeq X platform by Novogene Co., Ltd.

  QUANTIFICATION AND STATISTICAL ANALYSIS

***d*NET-seq data processing**

Adapter sequences were removed from all *d*NET-seq paired-end samples using Cutadapt

(version 1.18) (Martin 2011) with the following parameters: -a TGGAATTCTCGGGTGCCAAGG -A

GATCGTCGGACTGTAGAACTCTGAAC -m 10 -e 0.05 --match-read-wildcards -n 1. Paired-end

read merging was performed using bbmerge.sh from BBMap (Bushnell et al. 2017) with the 'xloose'

parameter. Merged reads were then aligned to the *Drosophila* reference genome (*dm6*; Ensembl release

95) (Cunningham et al. 2019) using STAR (version 2.6.0b) (Dobin et al. 2013) with –chimSegmentMin

set to 20. Only uniquely mapped reads were considered, extracted using SAMtools (version 1.7) (Li et

al. 2009) with -q set to 255. Exceptionally, in **Supplemental Fig S2A**, HiSat2 was used, using the same

dm6 genome annotation file for genome indexing and using default parameters, with the   --no-

discordant –no-mixed flags set. PCR internal priming events generated during library preparation were

removed using a custom Python script (Prudêncio et al. 2020) with the following parameters: -a TGG..

-s paired. To obtain single-nucleotide resolution, a custom Python script (Prudêncio et al. 2020) was

used to extract the 5' end nucleotide of read 2 (after trimming) in each sequencing pair, with the

directionality indicated by read 1 (**Fig 1E**).

Publicly available RNA-seq datasets used

Publicly available *Drosophila* embryonic transcriptome sequencing data (Poly(A) RNA-seq),

performed in developmental stages similar to the dNET-seq early and late samples, were used in this

study. RNA-seq datasets corresponding to cycle 14B (Lott et al. 2011) were obtained from the Gene

Expression Omnibus (GEO) (samples GSM618409, GSM618410, GSM618421 and GSM618422 from

dataset GSE25180). RNA-seq datasets from 4-6h old embryos were obtained from modENCODE

project PRJNA75285 (Graveley et al. 2011) (accessions SRR023696, SRR023746, SRR023836,

SRR035220, SRR023669, SRR035405, SRR035406, SRR024014 and SRR023539).

RNA-seq data processing

Adapters were removed from all datasets using Trim Galore (version 0.4.4) (http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/; last accessed 26 April 2020). Datasets from modENCODE and GEO were aligned to the *dm6 Drosophila* reference genome (Ensembl release 95) (Cunningham et al. 2019) using STAR (version 2.6.0b) (Dobin et al. 2013) with --chimSegmentMin set to 20. Stringtie (version 1.3.3b) (Pertea et al. 2015) was used to quantify normalized gene expression as Transcripts Per Kilobase Million (TPM) values with the following parameters: -a 5 -e. In addition, the isoform list was provided together with the -G parameter corresponding to the *dm6* (Ensembl release 95) GTF file. Genes with TPM values above 2 were considered to be expressed.

Selection of genes and isoforms for analysis

Similar to a previous GRO-seq analysis (Core et al. 2008), we used the read density of very large intergenic regions (gene desert regions) to define the reference for absence of transcription. Gene deserts were divided into 50kb windows, and *d*NET-seq read densities were calculated by dividing the read counts in each window by the window length (in bp). Read counts per window were obtained with bedtools genome coverage (version 2.27.1-1-gb87c465) (Quinlan and Hall 2010), and an arbitrary density threshold was defined as the 90th percentile of the read density distribution (**Supplemental Fig S4A**). Transcripts whose gene body *d*NETseq read density exceeded this threshold were considered to be transcriptionally active (**Supplemental Fig S4B**). For all of the analyses performed on late genes, only transcriptionally active genes were considered. In addition, only one isoform per gene was considered for all analysis, selected as the isoform with the highest RPKM value in the RNA-seq dataset for the corresponding developmental stage.

The coordinates of previously identified pre-MBT genes were obtained from Chen et al. (2013) and converted to *dm6* coordinates using the *liftOver* tool (https://genome.ucsc.edu/cgi-

bin/hgLiftOver). The most representative isoform for each gene was manually selected through

visualization of individual profiles.

Splicing intermediate and lariat detection

Exons containing splicing intermediates or introns containing lariats were identified using a

peak finder algorithm (NET_snrPeakFinder) (Churchman and Weissman 2011; Prudêncio et al. 2020)

that detects the presence of a peak in the last nucleotide of an exon (splicing intermediate) or an intron

(lariat), by comparing the accumulation of 3' end reads mapping at that position with the mean read

density of the flanking 200 nucleotides. A peak is called when the read density at the peak is superior

to the mean of this surrounding region plus 3 standard deviations (Churchman and Weissman 2011).

Since gene read density influences peak detection, the exons were divided into quartiles based on the

*d*NET-seq read density of the corresponding gene. Only exons from the highest quartile (i.e. from genes

with the highest read density) were considered in **Fig 2** for *d*NET-seq Late analysis.

Analysis of read density and peak calling

RPKM values for the merged *d*NET-seq datasets were calculated in the following manner:

$$RPKM\ (transcript) = \frac{reads * \ 10^4 * 10^6}{(total\ uniquely\ mapped\ reads) * (gene\ length\ in\ bp)}$$

Where 10^4 normalizes for gene length and 10^6 normalizes for sequencing depth. Queries of gene

3'UTR overlaps (**Figure 3**) between genes were performed with bedtools intersect (version 2.27.1-1-

gb87c465) (Quinlan and Hall 2010).

To detect splicing intermediate and intron lariat peaks, the algorithm from Churchman and

Weissman (2011) was adapted as described in Prudêncio et al. (2020). To be able to call larger regions

as peaks, a custom algorithm was developed and implemented in Python 3.7. Note that like for all of

the custom Python code reported here, there is heavy reliance on *bedtools* v2.29.0 (Quinlan and Hall

2010) for operations on coordinate intervals. In addition, NumPy v1.17.2 (Harris et al. 2020) and SciPy

1.4.0 (SciPy 1.0 Contributors et al. 2020) were used. Our peak caller detects regions where the local read density is significantly higher than expected by chance given the overall read density of the transcript. Almost all of the numerical parameters used by the peak caller can be adjusted. In **Fig 4A**, results obtained with two particular parameterizations are shown. Peak Caller 1 ("large peaks") is adapted to detecting larger peaks and provides results that are more intuitive to a human observer. Peak Caller 2 ("small peaks") provides a finer spatial resolution, and corresponds to the settings used in all of the analyses in this study. The peak caller takes as input a BED file with the 3' ends of reads (single-nucleotide resolution), as well as a GTF file with transcript and exon annotations and a list of transcripts to analyse. It functions by calculating a sliding average of read density within each transcript (window size 5/21 for small/large; only reads mapping to the same strand as the annotated transcript are considered). It then randomly shuffles the positions of the reads within the transcript and recalculates the sliding averages to determine the random expectation. This can be repeated several times (5 in this study) for more robustness. Windows obtained with the true read distribution are called as significant if their read density is higher than the 99[th] percentile of the simulated windows. Note that in this study, we used a setting whereby this threshold is calculated separately for each exon (and its upstream intron, for exons other than the first), by excluding the intron-exon pair of interest and reads overlapping it during the simulation step. This is necessary so that when calling peaks within a given exon (and its upstream intron), the threshold set would not be affected by the reads within that particular exon and its upstream intron. This way, for instance, the calling of a peak in the beginning of the exon is not affected by the calling of a peak in the middle of the same exon (except through potential merging, see below). After the initial peaks are called, they are filtered to remove peaks where more than 90% of the reads come from a single nucleotide (probable PCR duplicates), that are shorter than 5 nucleotides, or that overlap with fewer reads than a specified threshold (10/5 for large/small). Finally, peaks that are within a specified distance of each-other (21/5 nucleotides for large/small) are merged together.

Individual gene profiles

Individual dNET-seq gene profiles were generated by separating reads by strand using SAMtools (version 1.7) (Li et al. 2009). Strand-separated read data was converted to bedGraph format using bedtools genomecov with the -bg flag (version 2.27.1-1-gb87c465) (Quinlan and Hall 2010). Coverage values were normalized per nucleotide accounting for the total number of uniquely aligned reads and with the scale set to reads per 10^8 sequences. The outcome was converted to bigwig files through the bedGraphToBigWig tool (Kent et al. 2010) and uploaded to the UCSC genome browser (James Kent et al. 2002).

Metagene analysis

The read density metagene plots in **Fig 3F-I** and **Fig 5E,F** were created with deepTools (version 3.0.2) (Ramírez et al. 2016). Metagenes with normalized gene size (Figure 3) have bins of 10 bp while all other metagene plots in this study have single nucleotide resolution. Normalized gene and intron lengths (**Fig 3F-I**) were obtained through the scale-regions option. Exon-intron junctions without normalized lengths were obtained using the reference-option set to the 3'SS or the 5'SS. For normalization, we divided the number of reads at each nucleotide (or bin) by the total number of reads in the entire genomic region under analysis. These values were then used to calculate the mean for each nucleotide, and the results were plotted in an arbitrary units (A.U.) ranging from 0 to 1.

The peak or read density metagene plots in **Fig 3**, **4B** and **7D**, and **Supplemental Fig S3G, S7A-G** and **S7E-J** were prepared using custom Python and R scripts (https://github.com/rosinaSav/dNETseq_code). The peak density value represents the proportion of introns/exons that overlap with a peak at that position. Only internal fully coding exons that were at least 100 nucleotides long were included. In addition, the intron just upstream of the exon had to be at least 50 nt long. For **Fig 7D**, **S7A-G** and **S7E-H**, further filtering based on read coverage was performed (see below). Note that exons shorter than 150 nucleotides contribute both to the upstream and downstream exonic proportion of the plot.

Immediate splicing analysis

The immediate splicing analysis was performed solely on the S5P data sets. Only the 117 previously annotated pre-MBT genes (Chen et al. 2013) were analysed for early data sets. Reads were considered as spliced if they contained 'N's in the CIGAR string, in a position corresponding to an annotated intron. Reads that overlapped both the (unspliced) intron and the downstream exon were considered as unspliced. In both cases, only reads where the 3' end was located at least 5 nt downstream of the 3' ss were included, to avoid analysing misaligned reads whose 3' end should have mapped to the end of the upstream exon instead. Spliced reads that had the 5' end mapped to the upstream exon and the 3' end mapped to the intron were considered indicative of recursive splicing if the first nucleotide of the downstream end (indicative of the ratchet point position) matched the second G in the AGGT canonical splicing motif. If the last nucleotide of a read matched the last nucleotide of an exon, it was considered a splicing intermediate read and not representative of nascent RNA. A splicing ratio was calculated by dividing the number of nascent RNA spliced reads by the sum of the number of spliced and unspliced nascent RNA reads, only including reads whose 3' ends mapped to the first 100 nt of the downstream exon. Only fully coding internal exons at least 100 nt long were considered (exceptionally, in **Fig 6E** and **K** and **Supplemental Fig S6O-Q**, the 3' most coding exon was also analysed). Finally, we performed filtering to remove exons where the read coverage was too low to allow for robust estimation of the splicing ratio. The relevant threshold was calculated for each dataset separately. We calculated the total proportion of spliced reads out of all spliced/unspliced reads for the dataset to obtain the expected splicing ratio. We then performed a binomial test to know the probability of sampling only spliced/unspliced reads by chance under the null that the true splicing ratio equalled this expectation. We set the threshold as the lowest number of reads that had to be sampled for the probability to be below 0.01. Through his procedure, the threshold was set at >=10 reads for replicate 1 and 2 of the late data set (no terminal coding exons), at >=11/10 reads for replicate 1/2 of the late data set including terminal coding exons, and at >=14/9 reads for replicate 1/2 of the early data set (terminal coding exons always included).

Gene architecture and nucleotide composition analysis

Gene architecture and nucleotide composition parameters were calculated using custom Python and R scripts based on Ensembl annotations for dm6.18 (Cunningham et al. 2019). Splice site strength scores were calculated using MaxEntScan (Yeo and Burge 2004) with default parameters. Significance testing was performed using Kruskal-Wallis tests with a Bonferroni correction for multiple comparisons, with the correction applied separately for either replicate. For predictors where the corrected *P*-value was <0.05, Dunn's test was performed on the pairwise comparisons using *R* package *dunn.test* (Dinno 2017). The sequence logo in **Fig 4D** was generated using the *seqLogo* package version 1.52.0 in R (Bembom 2019).

Data access

All raw and processed sequencing data generated in this study have been submitted to the NCBI Gene Expression Omnibus (GEO; https://www.ncbi.nlm.nih.gov/geo/) under accession number GSE152585. The Python code used is available at https://github.com/rosinaSav/dNETseq_code and https://github.com/kennyrebelo. The Python code can also be found in the **Supplemental Code** file.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

We thank Takayuki Nojima and Nicholas Proudfoot (University of Oxford, UK) for critical discussion. This work was supported by funding to M.C.-F. (Fundação para a Ciência e Tecnologia, FCT/ Ministério da Ciência, Tecnologia e Ensino Superior - Fundos do Orçamento de Estado (UIDB/50005/2020), and FCT/FEDER/POR Lisboa 2020, Programa Operacional Regional de Lisboa PORTUGAL 2020, grant LISBOA-01-0145-FEDER-016394) and to R.G.M (FCT grant PTDC/BIA-

1021   BID/28441/2017). P.P. was a recipient of a FCT fellowship (SFRH/BD/109689/2015). R.S. was a

1022   recipient of an EMBO Long-Term Fellowship (EMBO ALTF 101-2019). This project has received

1023   funding from the European Union's Horizon 2020 research and innovation programme under grant

1024   agreements No 842695 (Marie Skłodowska-Curie Actions) and No 857119 (RiboMed).

1025

1026   **Author Contributions**

1027

1028   P.P. performed all molecular biology experiments. R.S. performed the majority of splicing kinetics

1029   analyses and developed the peak caller for detecting putative pause sites. K.R. performed additional

1030   bioinformatics analyses. P.P., R.S., R.G.M. and M.C.-F conceived the study and wrote the paper.

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

REFERENCES

Akhtar J, Kreim N, Marini F, Mohana G, Brüne D, Binder H, Roignant J-Y. 2019. Promoter-proximal pausing mediated by the exon junction complex regulates splicing. *Nat Commun* **10**: 521.

Alexander RD, Innocente SA, Barrass JD, Beggs JD. 2010. Splicing-dependent RNA polymerase pausing in yeast. *Molecular Cell* **40**: 582–593.

Ali-Murthy Z, Lott SE, Eisen MB, Kornberg TB. 2013. An Essential Role for Zygotic Expression in the Pre-Cellular Drosophila Embryo. *PLoS Genetics* **9**.

Alpert T, Herzel L, Neugebauer KM. 2017. Perfect timing: splicing and transcription rates in living cells. *WIREs RNA* **8**.

Ameur A, Zaghlool A, Halvardson J, Wetterbom A, Gyllensten U, Cavelier L, Feuk L. 2011. Total RNA sequencing reveals nascent transcription and widespread co-transcriptional splicing in the human brain. *Nature Structural and Molecular Biology* **18**: 1435–1440.

Artieri CG, Fraser HB. 2014. Transcript length mediates developmental timing of gene expression across drosophila. *Molecular Biology and Evolution* **31**: 2879–2889.

Arzate-Mejía RG, Josué Cerecedo-Castillo A, Guerrero G, Furlan-Magaril M, Recillas-Targa F. 2020. In situ dissection of domain boundaries affect genome topology and gene transcription in Drosophila. *Nat Commun* **11**: 894.

Aslanzadeh V, Huang Y, Sanguinetti G, Beggs JD. 2018. Transcription rate strongly affects splicing fidelity and cotranscriptionality in budding yeast. *Genome Research* **28**: 203–213.

Bembom O. 2019. seqLogo: Sequence logos for DNA sequence alignments. R package version 1.52.0.

Berget SM. 1995. Exon Recognition in Vertebrate Splicing. *The Journal of Biological Chemistry* **270**: 2411–2414.

Beyer AL, Osheim YN. 1988. Splice site selection, rate of splicing, and alternative splicing on nascent transcripts. *Genes & Development* **2**: 754–765.

Blythe SA, Wieschaus EF. 2015. *Coordinating Cell Cycle Remodeling with Transcriptional Activation at the Drosophila MBT*. 1st ed. Elsevier Inc.

Blythe SA, Wieschaus EF. 2016. Establishment and maintenance of heritable chromatin structure during early Drosophila embryogenesis. *Elife* **5**.

Boija A, Mahat DB, Zare A, Holmqvist P-H, Philip P, Meyers DJ, Cole PA, Lis JT, Stenberg P, Mannervik M. 2017. CBP Regulates Recruitment and Release of Promoter-Proximal RNA Polymerase II. *Molecular Cell* **68**: 491-503.e5.

Brugiolo M, Herzel L, Neugebauer KM. 2013. Counting on co-transcriptional splicing.
F1000Prime Reports **5**.

Burnette JM, Miyamoto-Sato E, Schaub MA, Conklin J, Lopez AJ. 2005. Subdivision of
large introns in Drosophila by recursive splicing at nonexonic elements. *Genetics* **170**:
661–674.

Bushnell B, Rood J, Singer E. 2017. BBMerge – Accurate paired shotgun read merging via
overlap. *PLoS ONE* **12**.

Campos-Ortega JA. 1985. Genetics of early neurogenesis in Drosophila melanogaster. *Trends
in Neurosciences* **8**: 245–250.

Carrillo Oesterreich F, Herzel L, Straube K, Hujer K, Howard J, Neugebauer KM. 2016.
Splicing of Nascent RNA Coincides with Intron Exit from RNA Polymerase II. *Cell*
165: 372–381.

Carrillo Oesterreich F, Preibisch S, Neugebauer KM. 2010. Global analysis of nascent RNA
reveals transcriptional pausing in terminal exons. *Molecular Cell* **40**: 571–581.

Chathoth KT, Barrass JD, Webb S, Beggs JD. 2014. A Splicing-Dependent Transcriptional
Checkpoint Associated with Prespliceosome Formation. *Molecular Cell* **53**: 779–790.

Chen K, Johnston J, Shao W, Meier S, Staber C, Zeitlinger J. 2013. A global change in RNA
polymerase II pausing during the Drosophila midblastula transition. *eLife* **2013**.

Churchman LS, Weissman JS. 2011. Nascent transcript sequencing visualizes transcription at
nucleotide resolution. *Nature* **469**: 368–373.

Core LJ, Waterfall JJ, Lis JT. 2008. Nascent RNA Sequencing Reveals Widespread Pausing
and Divergent Initiation. *Science* **322**: 1845–1849.

Cunningham F, Achuthan P, Akanni W, Allen J, Amode MR, Armean IM, Bennett R, Bhai J,
Billis K, Boddu S, et al. 2019. Ensembl 2019. *Nucleic Acids Research* **47**: D745–
D751.

Dahlberg O, Shilkova O, Tang M, Holmqvist P-H, Mannervik M. 2015. P-TEFb, the Super
Elongation Complex and Mediator Regulate a Subset of Non-paused Genes during
Early Drosophila Embryo Development ed. G.P. Copenhaver. *PLOS Genetics* **11**:
e1004971.

David CJ, Boyne AR, Millhouse SR, Manley JL. 2011. The RNA polymerase II C-terminal
domain promotes splicing activation through recruitment of a U2AF65-Prp19
complex. *Genes and Development* **25**: 972–982.

de Almeida SF, Carmo-Fonseca M. 2008. The CTD role in cotranscriptional RNA processing
and surveillance. *FEBS Letters* **582**: 1971–1976.

De Conti L, Baralle M, Buratti E. 2013. Exon and intron definition in pre-mRNA splicing.
Wiley interdisciplinary reviews RNA **4**: 49–60.

De Renzis S, Elemento O, Tavazoie S, Wieschaus EF. 2007. Unmasking activation of the zygotic genome using chromosomal deletions in the Drosophila embryo. *PLoS Biology* **5**: 1036–1051.

Dingwall C, Lomonossoff GP, Laskey RA. 1981. High sequence specificity of micrococcal nuclease. *Nucleic Acids Research* **9**: 2659–2674.

Dinno A. 2017. dunn.test: Dunn's Test of Multiple Comparisons Using Rank Sums. R package version 1.3.5.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15–21.

Drexler HL, Choquet K, Churchman LS. 2020. Splicing Kinetics and Coordination Revealed by Direct Nascent RNA Sequencing through Nanopores. *Molecular Cell* **77**: 985-998.e8.

Duff MO, Olson S, Wei X, Garrett SC, Osman A, Bolisetty M, Plocik A, Celniker SE, Graveley BR. 2015. Genome-wide identification of zero nucleotide recursive splicing in Drosophila. *Nature* **521**: 376–379.

Emili A, Shales M, McCracken S, Xie W, Tucker PW, Kobayashi R, Blencowe BJ, Ingles CJ. 2002. Splicing and transcription-associated proteins PSF and p54nrb/NonO bind to the RNA polymerase II CTD. *Rna* **8**: 1102–1111.

Erickson JW, Cline TW. 1993. A bZIP protein, sisterless-a, collaborates with bHLH transcription factors early in Drosophila development to determine sex. *Genes and Development* **7**: 1688–1702.

Fong N, Kim H, Zhou Y, Ji X, Qiu J, Saldi T, Diener K, Jones K, Fu XD, Bentley DL. 2014. Pre-mRNA splicing is facilitated by an optimal RNA polymerase II elongation rate. *Genes and Development* **28**: 2663–2676.

Fox-Walsh KL, Dou Y, Lam BJ, Hung S -p., Baldi PF, Hertel KJ. 2005. The architecture of pre-mRNAs affects mechanisms of splice-site pairing. *Proceedings of the National Academy of Sciences* **102**: 16176–16181.

Fukaya T, Lim B, Levine M. 2017. Rapid Rates of Pol II Elongation in the Drosophila Embryo. *Current Biology* **27**: 1387–1391.

Gaffney DJ, McVicker G, Pai AA, Fondufe-Mittendorf YN, Lewellen N, Michelini K, Widom J, Gilad Y, Pritchard JK. 2012. Controls of Nucleosome Positioning in the Human Genome. *PLoS Genetics* **8**.

Gelfman S, Cohen N, Yearim A, Ast G. 2013. DNA-methylation effect on cotranscriptional splicing is dependent on GC architecture of the exon-intron structure. *Genome Research* **23**: 789–799.

Görnemann J, Barrandon C, Hujer K, Rutz B, Rigaut G, Kotovic KM, Faux C, Neugebauer KM, Séraphin B. 2011. Cotranscriptional spliceosome assembly and splicing are independent of the Prp40p WW domain. *Rna* **17**: 2119–2129.

Graveley BR, Brooks AN, Carlson JW, Duff MO, Landolin JM, Yang L, Artieri CG, Baren MJ Van, Boley N, Booth BW, et al. 2011. The Developmental Transcriptome of Drosophila melanogaster. *Nature* **471**: 473–479.

Guilgur LG, Prudêncio P, Sobral D, Liszekova D, Rosa A, Martinho RG. 2014. Requirement for highly efficient pre-mRNA splicing during Drosophila early embryonic development. *eLife* **3**.

Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, Cournapeau D, Wieser E, Taylor J, Berg S, Smith NJ, et al. 2020. Array programming with NumPy. *Nature* **585**: 357–362.

Hatton AR, Subramaniam V, Lopez AJ. 1998. Generation of alternative Ultrabithorax isoforms and stepwise removal of a large intron by resplicing at exon-exon junctions. *Molecular Cell* **2**: 787–796.

Herzel L, Straube K, Neugebauer KM. 2018. Long-read sequencing of nascent RNA reveals coupling among RNA processing events. 1008–1019.

Heyn P, Kircher M, Dahl A, Kelso J, Tomancak P, Kalinka AT, Neugebauer KM. 2014. The earliest transcribed zygotic genes are short, newly evolved, and different across species. *Cell Reports* **6**: 285–292.

Hörz W, Altenburger W. 1981. Sequence specific cleavage of DNA by micrococcal nuclease. *Nucleic Acids Research* **9**: 2643–2658.

Hoskins RA, Landolin JM, Brown JB, Sandler JE, Takahashi H, Lassmann T, Booth BW, Zhang D, Wan KH, Yang L, et al. 2011. Genome-wide analysis of promoter architecture in Drosophila melanogaster. *Genome Research* **21**: 182–192.

Hsin JP, Manley JL. 2012. The RNA polymerase II CTD coordinates transcription and RNA processing. *Genes and Development* **26**: 2119–2137.

James Kent W, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, Haussler D. 2002. The human genome browser at UCSC. *Genome Research* **12**: 996–1006.

Ji J-Y, Squirrell JM, Schubiger G. 2004. Both Cyclin B levels and DNA-replication checkpoint control the early embryonic mitoses in Drosophila Jun-Yuan. *Development* **131**: 401–411.

Jonkers I, Kwak H, Lis JT. 2014. Genome-wide dynamics of Pol II elongation and its interplay with promoter proximal pausing, chromatin, and exons. *eLife* **2014**.

Joseph B, Kondo S, Lai EC. 2018. Short cryptic exons mediate recursive splicing in Drosophila. *Nature Structural and Molecular Biology* **25**: 365–371.

Kent WJ, Zweig AS, Barber G, Hinrichs AS, Karolchik D. 2010. BigWig and BigBed: Enabling browsing of large distributed datasets. *Bioinformatics* **26**: 2204–2207.

Keren H, Lev-Maor G, Ast G. 2010. Alternative splicing and evolution: Diversification, exon definition and function. *Nature Reviews Genetics* **11**: 345–355.

Khodor YL, Menet JS, Tolan M, Rosbash M. 2012. Cotranscriptional splicing efficiency
differs dramatically between Drosophila and mouse. *RNA* **18**: 2174–2186.

Khodor YL, Rodriguez J, Abruzzi KC, Tang CHA, Marr MT, Rosbash M. 2011. Nascent-seq
indicates widespread cotranscriptional pre-mRNA splicing in Drosophila. *Genes and*
Development **25**: 2502–2512.

Kimura H, Tao Y, Roeder RG, Cook PR. 1999. Quantitation of RNA Polymerase II and Its
Transcription Factors in an HeLa Cell: Little Soluble Holoenzyme but Significant
Amounts of Polymerases Attached to the Nuclear Substructure. *Molecular and*
Cellular Biology **19**: 5383–5392.

Kwak H, Fuda NJ, Core LJ, Lis JT. 2013. Precise maps of RNA polymerase reveal how
promoters direct initiation and pausing. *Science* **339**: 950–953.

Kwasnieski JC, Orr-Weaver TL, Bartel DP. 2019. Early genome activation in Drosophila is
extensive with an initial tendency for aborted transcripts and retained introns. *Genome*
Research **29**: 1188–1197.

Larson MH, Mooney RA, Peters JM, Windgassen T, Nayak D, Gross CA, Block SM,
Greenleaf WJ, Landick R, Weissman JS. 2014. A pause sequence enriched at
translation start sites drives transcription dynamics in vivo. *Science* **344**: 1042–1047.

Laver JD, Marsolais AJ, Smibert CA, Lipshitz HD. 2015. *Regulation and Function of*
Maternal Gene Products During the Maternal-to-Zygotic Transition in Drosophila.
1st ed. Elsevier Inc.

Lee JEA, Mitchell NC, Zaytseva O, Chahal A, Mendis P, Cartier-Michaud A, Parsons LM,
Poortinga G, Levens DL, Hannan RD, et al. 2015. Defective Hfp-dependent
transcriptional repression of dMYC is fundamental to tissue overgrowth in Drosophila
XPB models. *Nat Commun* **6**: 7404.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin
R. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**:
2078–2079.

Lott SE, Villalta JE, Schroth GP, Luo S, Tonkin LA, Eisen MB. 2011. Noncanonical
compensation of zygotic X transcription in early Drosophila melanogaster
development revealed through single-embryo RNA-Seq. *PLoS Biology* **9**.

Lund E, Dahlberg JE. 1992. Cyclic 2 ', 3 "-Phosphates and Nontemplated Nucleotides at the
3" End of Spliceosomal U6 Small Nuclear RNAs. *Science* **255**: 327–330.

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet.journal* **17**: 10–12.

Martin RM, Rino J, Carvalho C, Kirchhausen T, Carmo-Fonseca M. 2013. Live-Cell
Visualization of Pre-mRNA Splicing with Single-Molecule Sensitivity. *Cell Reports*
4: 1144–1155.

Martinho RG, Guilgur LG, Prudêncio P. 2015. How gene expression in fast-proliferating
cells keeps pace. *BioEssays* **37**: 514–524.

Maslon MM, Braunschweig U, Aitken S, Mann AR, Kilanowski F, Hunter CJ, Blencowe BJ,
Kornblihtt AR, Adams IR, Cáceres JF. 2019. A slow transcription rate causes
embryonic lethality and perturbs kinetic coupling of neuronal genes. *The EMBO*
Journal **38**.

Mayer A, Di Iulio J, Maleri S, Eser U, Vierstra J, Reynolds A, Sandstrom R,
Stamatoyannopoulos JA, Churchman LS. 2015. Native elongating transcript
sequencing reveals human transcriptional activity at nucleotide resolution. *Cell* **161**:
541–554.

Morris DP, Greenleaf AL. 2000. The splicing factor, Prp40, binds the phosphorylated
carboxyl-terminal domain of RNA Polymerase II. *Journal of Biological Chemistry*
275: 39935–39943.

Nojima T, Gomes T, Carmo-Fonseca M, Proudfoot NJ. 2016. Mammalian NET-seq analysis
defines nascent RNA profiles and associated RNA processing genome-wide. *Nature*
Protocols **11**: 413–428.

Nojima T, Gomes T, Grosso ARF, Kimura H, Dye MJ, Dhir S, Carmo-Fonseca M, Proudfoot
NJ. 2015. Mammalian NET-seq reveals genome-wide nascent transcription coupled to
RNA processing. *Cell* **161**: 526–540.

Nojima T, Rebelo K, Gomes T, Grosso AR, Proudfoot NJ, Carmo-Fonseca M. 2018. RNA
Polymerase II Phosphorylated on CTD Serine 5 Interacts with the Spliceosome during
Co-transcriptional Splicing. *Molecular Cell* **72**: 369-379.e4.

Pai AA, Henriques T, McCue K, Burkholder A, Adelman K, Burge CB. 2017. The kinetics of
pre-mRNA splicing in the Drosophila genome and the influence of gene architecture.
eLife **6**.

Pai AA, Paggi JM, Yan P, Adelman K, Burge CB. 2018. Numerous recursive sites contribute
to accuracy of splicing in long introns in flies. *PLoS Genetics* **14**.

Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. 2015. StringTie
enables improved reconstruction of a transcriptome from RNA-seq reads. *Nature*
Biotechnology **33**: 290–295.

Pritchard DK, Schubiger G. 1996. Activation of transcription in Drosophila embryos is a
gradual process mediated by the nucleocytoplasmic ratio. *Genes and Development* **10**:
1131–1142.

Prudêncio P, Rebelo K, Grosso AR, Martinho RG, Carmo-fonseca M. 2020. Analysis of
Mammalian Native Elongating Transcript sequencing (mNET-seq)high-throughput
data. *Methods* **178**: 89–95.

Quinlan AR, Hall IM. 2010. BEDTools: A flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**: 841–842.

Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dündar F,
Manke T. 2016. deepTools2: a next generation web server for deep-sequencing data
analysis. *Nucleic acids research* **44**: W160–W165.

Reimer KA, Mimoso CA, Adelman K, Neugebauer KM. 2021. Co-transcriptional splicing
regulates 3′ end cleavage during mammalian erythropoiesis. *Molecular Cell* **81**: 998-
1012.e7.

Robberson BL, Cote GJ, Berget SM. 1990. Exon Definition May Facilitate Splice Site
Selection in RNAs with Multiple Exons. *Molecular and Cellular Biology* **10**: 84–94.

Rothe M, Pehl M, Taubert H, Jäckle H. 1992. Loss of gene function through rapid mitotic
cycles in the Drosophila embryp. *Nature* **359**: 156–159.

Sandler JE, Irizarry J, Stepanik V, Dunipace L, Amrhein H, Stathopoulos A. 2018. A
Developmental Program Truncates Long Transcripts to Temporally Regulate Cell
Signaling. *Developmental Cell* **47**: 773-784.e6.

Saunders A, Core LJ, Sutcliffe C, Lis JT, Ashe HL. 2013. Extensive polymerase pausing
during Drosophila axis patterning enables high-level and pliable transcription. *Genes
and Development* **27**: 1146–1158.

Schlackow M, Nojima T, Gomes T, Dhir A, Carmo-Fonseca M, Proudfoot NJ. 2017.
Distinctive Patterns of Transcription and RNA Processing for Human lincRNAs.
Molecular Cell **65**: 25–38.

Schulz KN, Bondra ER, Moshe A, Villalta JE, Lieb JD, Kaplan T, McKay DJ, Harrison MM.
2015. Zelda is differentially required for chromatin accessibility, transcription factor
binding, and gene expression in the early Drosophila embryo. *Genome Res* **25**: 1715–
1726.

SciPy 1.0 Contributors, Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T,
Cournapeau D, Burovski E, Peterson P, Weckesser W, et al. 2020. SciPy 1.0:
fundamental algorithms for scientific computing in Python. *Nat Methods* **17**: 261–
272.

Sheridan RM, Fong N, D'Alessandro A, Bentley DL. 2019. Widespread Backtracking by
RNA Pol II Is a Major Effector of Gene Activation, 5′ Pause Release, Termination,
and Transcription Elongation Rate. *Molecular Cell* **73**: 107-118.e4.

Shermoen AW, O'Farrell PH. 1991. Progression of the Cell Cycle through Mitosis Leads to
Abortion of Nascent Transcripts. *Cell* **67**: 303–310.

Sterner DA, Carlo T, Berget SM. 1996. Architectural limits on split genes. *Proc Natl Acad
Sci USA* **93**: 15081–15085.

Veloso A, Kirkconnell KS, Magnuson B, Biewen B, Paulsen MT, Wilson TE, Ljungman M.
2014. Rate of elongation by RNA polymerase II is associated with specific gene
features and epigenetic modifications. *Genome Research* **24**: 896–905.

Vizcaya-Molina E, Klein CC, Serras F, Mishra RK, Guigó R, Corominas M. 2018. Damage-
responsive elements in *Drosophila* regeneration. *Genome Res* **28**: 1852–1866.

Windhager L, Bonfert T, Burger K, Ruzsics Z, Krebs S, Kaufmann S, Malterer G, L'Hernault
A, Schilhabel M, Schreiber S, et al. 2012. Ultrashort and progressive 4sU-tagging

reveals key characteristics of RNA processing at nucleotide resolution. *Genome*
Research **22**: 2031–2042.

Yan D, Neumüller RA, Buckner M, Ayers K, Li H, Hu Y, Yang-Zhou D, Pan L, Wang X,
Kelley C, et al. 2014. A Regulatory Network of Drosophila Germline Stem Cell Self-
Renewal. *Developmental Cell* **28**: 459–473.

Yeo G, Burge CB. 2004. Maximum entropy modeling of short sequence motifs with
applications to RNA splicing signals. *Journal of Computational Biology* **11**: 377–394.

Yuan K, Seller CA, Shermoen AW, Farrell PHO. 2016. Timing the Drosophila Mid-Blastula
Transition: A Cell Cycle-Centered View. *Trends in Genetics* **32**: 496–507.

Zhu J, Liu M, Liu X, Dong Z. 2018. RNA polymerase II activity revealed by GRO-seq and
pNET-seq in Arabidopsis. *Nature Plants* **4**: 1112–1123.

Zhu L, Zhang Y, Zhang W, Yang S, Chen J, Tian D. 2009. Patterns of exon-intron
architecture variation of genes in eukaryotic genomes. *BMC Genomics* **10**.

SUPPLEMENTAL FIGURES

A

Dataset	Raw reads	Uniquely Mapped after Merge
Early_2-3h_S5P_rep1	62,891,575	12,959,070
Early_2-3h_S5P_rep2	56,917,697	3,610,607
Late_4-6h_S5P_rep1	65,865,258	21,564,673
Late_4-6h_S5P_rep2	75,909,131	27,375,998
Late_4-6h_S5P_rep3	71,923,876	11,987,034
Late_4-6h_S2P_rep1	77,838,463	38,595,780
Late_4-6h_S2P_rep2	64,862,996	23,137,529
Late_4-6h_S2P_rep3	76,719,886	17,912,504

B

C

D

E

F

G

Dataset	Number of nascent reads	Nascent reads / Uniquely mapped reads (%)
Early_2-3h_S5P_rep1	7,984,411	61.6
Early_2-3h_S5P_rep2	2,430,876	67.3
Late_4-6h_S5P_rep1	13,413,045	62.2
Late_4-6h_S5P_rep2	13,777,880	50.3
Late_4-6h_S5P_rep3	8,505,056	71.0
Late_4-6h_S2P_rep1	23,153,659	60.0
Late_4-6h_S2P_rep2	12,926,707	55.9
Late_4-6h_S2P_rep3	10,356,743	57.8

H

Dataset	min	mean	max	25th percentile	50th percentile	75th percentile
Early_2-3h_S5P_rep1	35	99.6	293	74	81	119
Early_2-3h_S5P_rep2	35	94.3	293	72	81	110
Late_4-6h_S5P_rep1	35	114.2	293	86	108	135
Late_4-6h_S5P_rep2	35	102.0	294	80	87	119
Late_4-6h_S5P_rep3	35	88.4	294	76	81	89
Late_4-6h_S2P_rep1	35	107.0	294	81	96	128
Late_4-6h_S2P_rep2	35	111.1	294	81	102	132
Late_4-6h_S2P_rep3	35	109.0	293	83	105	130

Supplemental Fig. S1. (A) For each *d*NET-seq library prepared, the number of total reads and of uniquely aligned reads is indicated. **(B)** Density of uniquely aligned reads per gene (RPKM in log2 scale) for two *d*NET-seq/S5P biological replicates from early embryos (Pearson's correlation, R = 0.97). **(C)** Density of uniquely aligned reads per gene (RPKM in log2 scale) from late embryos analysed by *d*NET-seq/S5P and *d*NET-seq/S2P (Pearson's correlation, R = 0.95). **(D, E)** Density of uniquely aligned reads per gene (RPKM in log2 scale) for two *d*NET-seq/S5P and *d*NET-seq/S2P biological replicates from late embryos (Pearson's correlation, R = 0.92; R = 0.96). **(F)** Histogram of the read lengths in *d*NET-seq/S5P data from late embryos. **(G)** For each *d*NET-seq library generated, the number of uniquely aligned reads corresponding to nascent transcripts is indicated. **(H)** Length of sequenced nascent RNA (in nucleotides).

Supplemental Fig. S2. (A) *d*NET-seq/S5P profile over the U6 snRNA gene in the late dataset

(replicate 1). **(B)** Frequency of peaks corresponding to splicing intermediates (green) detected by

*d*NET-seq/S5Pon exons of genes expressed in late embryos. Genes were grouped into quartiles

(Q) based on their *d*NET-seqread density.

Supplemental Fig. S3. **(A-D)** *d*NET-seq/S5P and RNA-seq profiles over maternal genes in early embryos: *nanos* (**A**), *gurken* (**B**), *Rab32* (**C**) and *pumilio (pum)* (**D**). **(E-F)** *d*NET-seq/S5P and RNA-seq profiles over pre-MBT genes in early embryos: *fushi tarazu* (**E**) and *odd skipped (odd)* (**F**). (**G**) Meta-analysis of *d*NET-seq/S5P mean read density around the transcription start site (TSS) in maternal and pre-MBT genes (replicate 2). **(H-I)** *d*NET-seq/S5P and RNA-seq profiles over MBT genes in late embryos: *His3.3A* (**H**) and *twinstar (tsr)* (**I**). Reads that aligned to the positive strand are in blue, and reads that aligned to the negative strand are in red. The direction of transcription is indicated by an arrow.

Supplemental Fig. S4. **(A-C)** Identification of transcriptionally active genes based on *d*NET-seq signal along the gene body in late embryos. **(A)** Read density distribution in intergenic regions (gene deserts). The red dashed line represents the 90th percentile of read density in all intergenic regions analysed. **(B)** Read density distribution per gene (RPKM) represented in Log2 scale. The 90th percentile of read density over gene deserts is set as threshold (red dashed line). **(C)** Comparison of read density per active gene (RPKM in log2 scale) between the indicated datasets. Total number of active genes identified in each dataset is indicated (*n*). Venn diagrams show common active genes between datasets. **(D-G)** *d*NET-seq/S5P and RNA-seq profiles over the indicated pre-MBT genes. Reads that aligned to the positive strand are in blue, and reads that aligned to the negative strand are in red. The direction of transcription is indicated by an arrow.

Supplemental Fig . S5. **(A)** Venn diagram comparing RPs identified in two *d*NET-seq/S2P

biological replicates and in previously reported studies (Joseph et al. 2018; Duff et al. 2015).

Supplemental Fig. S6. In all panels, unless otherwise specified, only introns from transcriptionally active genes where the downstream exon is a fully coding internal exon at least 100 nt long were included. In addition, enough spliced/unspliced reads had to end within the first 100 exonic nucleotides that obtaining a splicing ratio (SR) of 0 or 1 by chance alone was highly unlikely (see the Methods section for details). For genes expressed in late embryos, this threshold was 10 reads for both replicates. For pre-MBT genes, it was 14 for replicate 1 and 9 for replicate 2. For S2P, a threshold of 10 was used for better comparability with S5P data. (**A**) SR values estimated in two biological replicates of dNET-seq/S5P datasets for pre-MBT genes in early embryos (Spearman correlation, $\rho = \sim 0.775$, $P \sim 1.348 * 10^{-14}$). (**B**) Distribution of SR values for dNET-seq/S5P (N = 4510) and dNET-seq/S2P (N = 4373). (**C**) SR values estimated in replicate 2 of dNET-seq/S5P and dNET-seq/S2P from late embryos (Spearman correlation, $\rho = \sim 0.737$, $P < 2.2 * 10^{-16}$; N = 3295). (**D-Q**) SR values for dNET-seq/S5P data (replicate 1 and 2) plotted out against several gene architecture parameters. For sample sizes and statistical tests, see **Supplementary Table 1**. In (**D-J, N-Q**), the bin ranges have been set so that intron numbers would be as equal possible between bins. (**R-S**) dNET-seq/S5P read density around the transcription end site (TES) (averaged over a 10-nucleotide sliding window) for genes with different mean splicing ratios in replicate 1 (**R**) and 2 (**S**). There is no significant correlation between SR and the count of reads 101-300 nucleotides past the TES (partial Spearman correlation controlling for the count of reads in the region 500 to 301 nucleotides before the TES as a proxy for gene body rad density; replicate 1: $\rho = \sim 0.013$, $P = \sim 0.519$; N = 2450; replicate 2: $\rho = \sim -0.034$, P = ~ 0.126; N = 2052).

Supplemental Fig. S7. (**A-C**) Metagene analyses of peak density estimated from *d*NET-seq/S5P datasets from late embryos for different SR ranges. (**A**) shows replicate 2, with introns with putative intron lariat reads removed (N = 3623). (**B**) and (**C**) show replicates 1 and 2, respectively, with introns with putative intron lariat reads included (N = 5614 for replicate 1 and N = 4503 for replicate 2). Peak density has been calculated as the proportion of introns that overlap with a peak at any given position. The last 50 nucleotides of exons, the first 50 nucleotides of introns, the last 25 nucleotides of introns, and the first 100 nucleotides of exons are shown. Only internal and fully coding exons from transcriptionally active genes that are at least 100 nucleotides long are shown. In addition, at least 10 spliced/unspliced reads had to end within the first 100 nucleotides of the exon. (**D**) The proportion of introns with at least one read whose 3' end maps to the final position of the upstream exon (putative splicing intermediates) or to the final position of the intron (putative intron lariats) in *d*NET-seq/S5P late replicate 2. (**E-F**) *d*NET-seq/S5P peak density profiles for late embryos, replicate 1 (**E**) and replicate 2 (**F**). The final 50 nucleotides of the intron and the first 100 nucleotides of the downstream exon are shown. For each data point, the GC content over the first 100 exonic nucleotides was divided by the GC content over the last 50 intronic nucleotides, generating the exonic to intronic GC ratio. Introns were split into two classes along the median of this metric. Only introns with an SR of 0 or 1 were plotted. (**G-H**) Similar to 7D and S7A, respectively, but showing read densities (averaged over a 10-nucleotide sliding window) rather than peak densities. (**I-J**) Metagene analysis of peak density estimated from *d*NET-seq/S2P datasets from late embryos, in replicate 1(**I**) and replicate 2 (**J**). The last 50 nucleotides of exons, the first 50 nucleotides of introns, the last 25 nucleotides of introns, and the first 100 nucleotides of exons are shown. Only internal and fully coding exons from transcriptionally active genes that are at least 100 nucleotides long are shown (N = 13229 for replicate 1, N = 11831 for replicate 2). Exons shorter than 150 nucleotides contribute to both the exon end and start. Only introns that were at least 50 nt long were considered.

SUPPLEMENTARY METHODS
*d*NET-seq sequence biases

Previous work has shown that MNase cleaves DNA more frequently at AT than at GC base pairs (Dingwall *et al.*, 1981). More specifically, Hörz and Altenburger (1981) found a preference for sites where a few alternating AT-TA base pairs were surrounded by a more GC-rich region, rather than for sites within longer AT-rich stretches. More recent genome-wide work in humans has come to similar conclusions (Gaffney *et al.*, 2012). *d*NET-seq relies on MNase cleavage of both DNA and RNA to solubilize chromatin. Any nucleotide biases in cleavage may be problematic, as they could mean that the solubilization is more efficient in regions where the nucleotide composition is more in line with such cleavage biases. This could lead to an artefactually increased read density in such regions.

To verify whether there was any evidence for RNA cleavage biases, we determined the nucleotide composition around the locations of the 5' ends of NET-seq reads (**Figure 4D** in main text). A clear preference for cleavage just 5' of an adenine was observed, with additional weaker biases at surrounding sites. Our results are therefore broadly in line with those obtained previously for DNA.

To what extent could these biases affect our results? We generated a simulated version of a *d*NET-seq dataset (S5P late, replicate 1) where the read positions were randomized in such a way as to preserve the nucleotide biases at the 5' ends of reads. Only reads that overlapped transcriptionally active genes were considered. More specifically, for each read, we defined the "starting hexamer" as the hexamer centred on the 5' end of the read (the three nucleotides just 5' of the read and the first three nucleotides of the read). We then randomly picked another occurrence of the same hexamers from within transcriptionally active genes and defined a "simulated read" at that position, preserving the length of the initial read. We then called peaks on the simulated reads similarly to true reads.

The mean proportion of nucleotides within simulated peaks was similar for both exons and introns (~0.017). However, this belies a difference in medians, with a higher peak density in exons than in introns (exon median: ~0.008; intron median: 0; $P < 2.2 * 10^{-16}$; two-tailed Mann-Whitney U-test; **Supplementary Methods Figure 1**). To know if the 5' end nucleotide bias effect could explain the bias towards exons observed in real data, we calculated a normalized peak density for each exon and intron, as $\frac{true\ peak\ density - simulated\ peak\ density}{simulated\ peak\ density}$. Exons and introns with a simulated peak density of 0 were discarded. Normalized peak densities were still higher for exons than for introns ($P < 2.2 * 10^{-16}$; two-tailed Mann-Whitney U-test). Hence, 5' end nucleotide biases may exaggerate but do not explain the bias towards exons in real data.

Supplementary Methods Figure 1: distribution of peak densities for exons and introns. A: true peaks. B: simulated peaks. C: normalized peaks, obtained as $\frac{true\ peak\ density\ -\ simulated\ peak\ density}{simulated\ peak\ density}$. For all three panels, the y axis has been limited to values below 10 for visualization reasons.

It may seem surprising that simulated read density would be higher in exons, given that *Drosophila* exons tend to have a higher GC content than introns (Zhu *et al.*, 2009). Indeed, one would, at first sight, expect the bias to exaggerate intronic peak density, rather than exonic peak density. However, peaks are called based on the locations of the 3' ends of reads, whereas the bias affects the locations of the 5' ends of reads. Therefore, many reads whose 3' ends are in GC-rich exonic sequence are expected to have their 5' ends in AT-rich intronic sequence.

Further examination of the meta-profiles reveals that the simulated peaks do not display the increased peak density in the beginning of the exons (first highlighted region in **Supplementary Methods Figure 2B**), as observed for true data (**Supplementary Methods Figure 2A**). It thus appears that this peak cannot be explained by read 5' end biases. However, the second highlighted region shows increased peak densities both in true and simulated data (**Supplementary Methods Figure 2A-B**). To verify if the increase observed in this region with true data could be explained by read 5' end nucleotide composition biases, we divided the true data into two classes based on whether or not any simulated peaks mapped to within the second highlighted region (**Supplementary Methods Figure 2C-D**). There was no visible difference in true mean peak densities between the two groups within the second highlighted region. This suggests that any nucleotide biases acting at the 5' ends of reads are unlikely to explain this increase in peak densities.

Supplementary Methods Figure 2: average peak density for the last 50 nucleotides of introns and the first 100 nucleotides of exons (concatenated). The orange boxes show the positions of exons. Only exons at least 100 nt long have been included.

Dingwall, C, Lomonossoff, GP, and Laskey, RA (1981). High sequence specificity of micrococcal nuclease. Nucleic Acids Res 9, 2659–2674.

Gaffney, DJ, McVicker, G, Pai, AA, Fondufe-Mittendorf, YN, Lewellen, N, Michelini, K, Widom, J, Gilad, Y, and Pritchard, JK (2012). Controls of Nucleosome Positioning in the Human Genome. PLoS Genet 8, 1–13.

Hörz, W, and Altenburger, W (1981). Sequence specific cleavage of DNA by micrococcal nuclease. Nucleic Acids Res 9, 2643–2658.

Zhu, L, Zhang, Y, Zhang, W, Yang, S, Chen, J, and Tian, D (2009). Patterns of exon-intron architecture variation of genes in eukaryotic genomes. 12, 1–12.