Comparison of SARS-CoV-2 infection in two non-human primate species: rhesus

and cynomolgus macaques

Kinga P. Böszörményi[1], Marieke A. Stammes[2], Zahra C. Fagrouch[1], Gwendoline Kiemenyi-

Kayere[1], Henk Niphuis[1], Daniella Mortier[1], Nikki van Driel[3], Ivonne Nieuwenhuis[2], Ella

Zuiderwijk-Sick[4], Lisette Meijer[2], Petra Mooij[1], Ed J. Remarque[1], Gerrit Koopman[1], Alexis C. R.

Hoste[5], Patricia Sastre[5], Bart L. Haagmans[6], Ronald E. Bontrop[7,8], Jan A.M. Langermans[3,9], Willy

M. Bogers[1], Ernst J. Verschoor[1*], and Babs E. Verstrepen[1]

[1]Department of Virology, Biomedical Primate Research Centre (BPRC), Rijswijk, The

Netherlands

[2]Department of Parasitology, BPRC, Rijswijk, The Netherlands

[3]Animal Science Department, BPRC, Rijswijk, The Netherlands

[4]Alternatives unit, BPRC, Rijswijk, The Netherlands

[5]Eurofins-Inmunología y Genética Aplicada (Eurofins-INGENASA), Madrid, Spain.

[6]Department of Viroscience, Erasmus University Medical Center, Rotterdam, The

Netherlands

[7]Department of Comparative Genetics and Refinement, BPRC, Rijswijk, The Netherlands

[8]Department of Biology, Theoretical Biology and Bioinformatics, Utrecht University, Utrecht,

The Netherlands

[9]Department of Population Health Sciences, Unit Animals in Science and Society, Veterinary

Faculty, Utrecht University, Utrecht, The Netherlands

*Corresponding author:

E-mail: verschoor@bprc.nl (EV)

Abstract

SARS-CoV-2 is a coronavirus that sparked the current COVID-19 pandemic. To stop the

shattering effect of COVID-19, effective and safe vaccines, and antiviral therapies are urgently

needed. To facilitate the preclinical evaluation of intervention approaches, relevant animal

models need to be developed and validated. Rhesus macaques (*Macaca mulatta*) and

cynomolgus macaques (*Macaca fascicularis*) are widely used in biomedical research and serve

as models for SARS-CoV-2 infection. However, differences in study design make it difficult to

compare and understand potential species-related differences. Here, we directly compared

the course of SARS-CoV-2 infection in the two genetically closely-related macaque species.

After inoculation with a low passage SARS-CoV-2 isolate, clinical, virological, and

immunological characteristics were monitored.

Both species showed slightly elevated body temperatures in the first days after exposure while

a decrease in physical activity was only observed in the rhesus macaques and not in

cynomolgus macaques. The virus was quantified in tracheal, nasal, and anal swabs, and in

blood samples by qRT-PCR, and showed high similarity between the two species.

Immunoglobulins were detected by various enzyme-linked immunosorbent assays (ELISAs)

and showed seroconversion in all animals by day 10 post-infection. The cytokine responses

were highly comparable between species and computed tomography (CT) imaging revealed

pulmonary lesions in all animals. Consequently, we concluded that both rhesus and

cynomolgus macaques represent valid models for evaluation of COVID-19 vaccine and

antiviral candidates in a preclinical setting.

Author summary

SARS-CoV-2 infection can have a wide range of symptoms. It can cause asymptomatic or mild

disease, but can also have a severe, potentially deadly outcome. Vaccines and antivirals will

therefore be crucial in fighting the current COVID-19 pandemic. For testing these prophylactic

and therapeutic treatments, and investigating the progression of infection and disease

development, animal models play an essential role. In this study, we compare the course of

SARS-CoV-2 infection in rhesus and cynomolgus macaques. Both species showed moderate

disease symptoms as shown by pulmonary lesions by CT imaging. Shedding of infectious virus

from the respiratory system was also documented. This study provides a detailed description

of the pathogenesis of a low-passage SARS-CoV-2 isolate in two macaque models and suggests

that both species represent an equally good model in research for both COVID-19 prophylactic

and therapeutic treatments.

Introduction

At the end of 2019, the first cases of coronavirus disease 2019 (COVID-19) were described in the Wuhan region, China [1]. Since then, the causative agent of COVID-19, the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) has spread rapidly across the globe, leading the World Health Organization to officially declare COVID-19 a pandemic on 11 March 2020. To halt the devastating impact of this disease and the ongoing pandemic, vaccines and antiviral therapies are desperately needed, as well as fundamental knowledge to understand the mode of infection and its associated pathologies. Hence, enormous efforts are initiated to develop a wide range of COVID-19 vaccines.

Between humans, transmission mainly occurs via aerosolized droplets after sneezing or coughing or via direct contact with contaminated surfaces. Similar to the related SARS-CoV, SARS-CoV-2 enters the body via docking with its spike (S) protein to the angiotensin-converting enzyme-2 (ACE2) receptor protein [2]. ACE2 is abundantly expressed on cells of the respiratory system but also on a variety of other organ tissues, including those of the brain and the gastrointestinal tract [3]. This wide-spread distribution of the ACE2 receptor may explain the complex clinical picture of COVID-19. The spectrum of COVID-19 ranges from asymptomatic or mild disease, via flu-like illness to a severe, potentially deadly disease [4]. The most typical hallmark of severe COVID-19 is pneumonia leading to respiratory failure, which occurs after the onset of dyspnea and hypoxemia [5]. However, COVID-19 patients may also suffer from a variety of other symptoms, including amongst other gastrointestinal symptoms, neurological disorders, coagulopathy, and cardiac injury [5-8].

Animal models to investigate the progression of infection and disease development, and to evaluate prophylactic and therapeutic treatment options, are essential to warrant progress in SARS-CoV-2 fundamental and applied research. Different animal species have shown their

value in SARS-CoV-2 research. Originally developed to study SARS-CoV [9], transgenic mice

that express the ACE2 receptor elucidated various aspects of COVID-19 [10, 11]. Rodent

species, such as the Syrian hamster, turned out to be susceptible to infection with SARS-CoV-

2, and hamsters were subsequently utilized to test a live-attenuated YF17D-vectored SARS-

CoV2 vaccine candidate [12, 13]. Ferrets represent an established animal model to study the

pathology and transmission of respiratory viruses, like influenza viruses [14]. During the

course of SARS-CoV infection, viral replication was documented in the upper and lower

respiratory tract [15]. Transmission of SARS-CoV-2 among ferrets via either air or direct

contact has been shown recently, where virus was shed in nasal washes, saliva, feces, and

urine of infected animals [16, 17]. Notwithstanding the great importance of rodent and ferret

models, also non-human primates (NHPs) will likely play a pivotal role in COVID-19 research.

Like with SARS-CoV, NHPs are susceptible to infection with human SARS-CoV-2. Due to their

close phylogenetic relationship, NHPs and humans share many immunological and

pathological characteristics [18-20]. This makes NHPs suitable for preclinical evaluation of

vaccines and antiviral or immunomodulatory compounds to combat SARS-CoV-2. So far, five

species of NHPs; rhesus and cynomolgus macaques, common marmosets (*Callithrix jacchus*),

African green monkeys (*Chlorocebus sabaeus*), and sacred baboons (*Papio hamadryas*) have

featured in SARS-CoV-2 studies. Common marmosets, a small New World monkey species, are

susceptible to infection with SARS and MERS coronaviruses [21, 22], but appear to be

relatively resistant to infection with SARS-CoV-2 as only low levels of virus replication could

be measured [23, 24]. African green monkeys presented robust virus replication and also

showed evidence of respiratory disease [25, 26], in contrast to baboons where variable levels

of virus replication were measured [24]. The two most widely-used NHP species in biomedical

research are rhesus macaques and cynomolgus macaques. Both macaque species had already

proven their value in research on the related coronaviruses that caused the SARS and MERS epidemics [27, 28], and thus are considered relevant NHP models for preclinical COVID-19 studies. Cynomolgus macaques have been deployed in studies describing aspects of SARS-CoV-2 pathogenesis [23, 29, 30], and have been utilized to evaluate the efficacy of hydroxychloroquine as an antiviral compound [31]. Rhesus macaques have also been applied in COVID-19 pathogenesis studies [22, 24, 32, 33], and to test the efficacy of remdesivir in the treatment of SARS-CoV-2 infection [34]. Additionally, several prototype COVID-19 vaccine candidates have received their first efficacy evaluation in the rhesus macaque model [35-42]. Some research groups [23, 24] shed light on the heterogeneity in SARS-CoV-2 infection and investigated disease progression in different NHP species. Most of these studies were conducted by different research teams, and a controlled comparative approach is lacking thus far.

In other NHP disease models, like those developed for AIDS, TB, and influenza research, the choice of macaque (sub)species can influence the disease outcome considerably [43-47]. The choice of a specific NHP species for research on a new and complex disease, like COVID-19, is therefore not a trivial one and the key question which macaque species is best suited to investigate specific aspects of COVID-19 research needs to be answered. To address this issue, we compared SARS-CoV-2 replication in rhesus and cynomolgus macaque species and monitored signs of COVID-19-like disease symptoms for three weeks after infection. The macaques were infected in parallel with the same virus stock, received completely identical treatment, and the course of infection was followed using the same analyses, including monitoring of lung pathology using computed tomography (CT), and continuous telemetric recording of body temperature and activity of the animals.

Results

Infection of macaques with SARS-CoV-2

After administration of the virus in the upper trachea and nose, levels of viral RNA were detectable in the tracheal and nasal swabs of all monkeys at day 1 pi. Viral RNA remained evident in swab samples for several days. In the tracheal swab sample of rhesus macaque R14002, viral RNA was first time below the detection time at 10 days pi. (Fig 1A, S1 Table). The individual variation of SARS-CoV-2 RNA levels detected in the macaques, regardless of species, was considerable. Peak viral RNA levels in the trachea varied between 1.7×10^4 copies/ml (R15096; day 1 pi) and 1.8×10^8 copies/ml (J16017; day 2 pi). The time frame in which viral genetic material could be detected varied from only one day (R15096; day 1 pi) up to day 10 pi. (animal R14002, RNA in the trachea).

Peak viral loads detected in nasal swabs were generally lower than levels observed in the throat samples and did not exceed 9.5×10^4 copies/ml (R15090; day 1 pi). The high virus loads measured in the first two days post-infection may suggest that some remaining RNA from the original inoculum was still present. However, in all macaques, viral RNA was also isolated from nasal swabs at later time points, showing that SARS-CoV-2 was excreted via the nose, and thus indicative for viral replication. The total viral RNA production over time is shown in Fig 1B. The patterns of viral RNA detection in swabs also varied between individuals. The most outstanding observation was made for cynomolgus macaque J16017 that was positive in the nose at day 1 pi, then had no detectable viral RNA for a period of three days, but later the animal became again positive in the nose swabs for three consecutive days. Other animals (R15096, J16004, J16012, and Ji40805) also became PCR-positive again after a period of one or more days characterized by undetectable levels. In the anal swabs, viral genetic material was rarely detected. Only at a few time points, three macaques tested positive, with a

maximum viral RNA load of 3 x 10^3 copies/ml at day 1 pi., namely in cynomolgus macaque

J16017. One animal tested positive for viral RNA in blood at a single time point; R15080 at day

5 pi. (S1 Table). Notably, no significant differences in viral RNA loads were calculated between

the macaque species (Fig 1B).

Body temperature, activity, clinical symptoms and blood parameters after SARS-CoV-2

infection

Body temperature and activity of each animal was continuously monitored using a Physiotel

Digital telemetric device during the entire study. Upon infection, elevated body temperatures

were measured in both macaque species, which could be correlated to the episodes of viral

replication in the nose and trachea as was evidenced by qRT-PCR. In Fig 2, we show the body

temperature alterations from the baseline during the study. In both groups of animals, the

body temperature was significantly higher during the first two weeks after infection as

compared to later time points (Fig 2). The temperature curves for the individual animals are

depicted in the supplementary data (S1 Fig). The group of cynomolgus macaques showed

elevated body temperature in the first 8 to 10 days following infection. This is in contrast with

the measurements of the rhesus macaques where no substantial rise in temperature was

measured, except for two animals (R14002 and R15090) that showed a sudden peak in body

temperature of 0.7°C at day 8 pi.

The activity curves measured in the 3-weeks observation period for all individuals are

documented in the S2A Fig. The cumulative activity scores in the first 2 weeks pi. were

compared with activity scores in the last week of the study period (S2B Fig). The paired t-test

illustrated a significantly lower activity in all four rhesus macaques during the first period after

infection (p=0.0072), while this difference in cynomolgus macaques was less obvious and only

found in 2 out of 4 macaques.

We applied a clinical scoring list to enumerate clinical symptoms that may be caused by the

SARS-CoV-2 infection (S2 Table). The cumulative clinical scores per week did not exceed 50 (of

a maximum 490 score per week; data not shown), confirming the absence of serious COVID-

19-related symptoms. However, in the second week of infection, cynomolgus macaques

showed more, but still mild, clinical symptoms than rhesus macaques. This was less evident

during the first and third weeks, probably due to outlier clinical scores of individual animals

(Fig 3).

Blood samples were analyzed for changes in cell subsets and in biochemical parameters. These

data were related to a set of normal (standard) values derived from uninfected, healthy

rhesus, and cynomolgus macaques from the same breeding colony. No significant deviations

from the normal values were seen in blood cell subsets of the infected monkeys. C-reactive

protein levels, which are increased in COVID-19 patients with pneumonia [48], were not found

higher in infected macaques. In humans, acute kidney injury has been related to SARS-CoV-2

infection [49, 50], and elevated levels of serum creatinine and blood urea were detected in

10-15% of a cohort of COVID-19 patients [51]. Hence, we measured creatinine and urea levels

in macaque blood samples at days 0, 5, 10, 14, and 22 pi., but did not find evidence of kidney

malfunction in the infected, but otherwise seemingly healthy monkeys. Equally, depending on

the severity of the disease, blood coagulation disorders, like highly elevated D-dimer levels,

have been reported for patients [52, 53], but no elevated D-dimer levels were measured in

either macaque species. Elevated levels of glucose and alanine transferase were measured in

the first week pi. in the blood of most animals, and amylase was increased in one rhesus

macaque, R15080 (S3 Fig).

Detection of lung lesions in macaques after SARS-CoV-2 infection.

Chest CTs of the macaques after infection revealed several manifestations of COVID-19 with

a variable time course and lung involvement (Table 1). The most common lesion types that

were found in both rhesus and cynomolgus macaques were ground glass opacities (GGO),

consolidations, and crazy paving patterns (CCP) (Fig 4).

Table 1. CT scores of lung lesions in SARS-CoV-2-infected macaques

		Animals							
		R14002	R15080	R15090	R15096	J16004	J16012	J16017	Ji408005
Days pi	0	0	0	0	0	0	0	0	0
	2	0	1	2	2	2	2	0	0
	4	0	0	0	2	0	4	0	0
	6	0	0	0	2	0	5	0	4
	8	3	0	2	3	0	5	3	2
	10	2	2	0	2	2	1	3	4
	12	3	3	0	1	3	1	0	0
	14	1	2	0	3	1	0	0	2
	16	0	0	0	4	3	6	2	4
	22	3	0	0	3	3	6	1	2

Lung lesions (max. CT score 2/35) were already seen in CTs early after infection on day 2 in 5

out of 8 monkeys, three rhesus, and two cynomolgus macaques. Thereafter, lung involvement

was seen in most animals and CT scores increased. Around days 8 and 10 pi., lung lesions were

manifest in all animals, and in several macaques the coverage had increased (Table 1). Again,

individual variations in lung pathology were considerable; for instance, rhesus macaque

R15090 showed low CT scores of 2 only at days 2 and 8 pi, while rhesus macaque R15096 had

positive CT scores at all time points after experimental infection, varying from 1 to 4.

Cynomolgus macaque J16012 suffered from serious lung damage as early as day 4 pi (score 5)

and had a positive chest CT at 8 out of 9 time points. In Fig 5, the cumulative CT scores over

the monitoring period are depicted. The increase in cumulative CT score was 0.692 points per

day (95% CI 0.657 to 0.726) and no difference was observed between rhesus and cynomolgus

macaques (P = 0.2708).

Immune response to SARS-CoV-2 infection

Humoral immune responses developed relatively fast after infection (Fig 6 and S4 Fig). Total

immunoglobulin (Ig) in serum was detected by the DR-ELISA. The antibody response directed

to the nucleoprotein (N) of SARS-CoV-2 became evident between day 10 and 12 pi., and

continued to rise thereafter, but with individual variations in total Ig levels and patterns. While

Ig levels in several animals, both rhesus and cynomolgus macaques, continued to rise until day

21 pi., the Ig levels in others already showed first signs of a decline at day 17 pi. The total Ig

pattern seen in the sera of all macaques was reflected by the measurement of IgG directed to

the N protein in the same sera. Notably, the development of IgM titers was barely detected

in the longitudinal serum samples. In sera from only one animal, cynomolgus macaque

J106012, IgM was detectable, beginning at day 12 pi. but titers already started to decline to

baseline levels at the end of the study at day 22 pi.

Cytokine and chemokine measurements

Serum levels of 13 cytokines and chemokines were measured to examine the nature of the

inflammatory response triggered by the infection with SARS-CoV-2. We investigated if any

species differences could be distinguished in the inflammatory response to infection. All

metadata obtained from the individual animals are depicted in S5 Fig. Directly upon infection,

IP-10 and MCP-1 levels peaked in sera of all cynomolgus macaques. The IP-10 peak at day 2

pi. quickly returned to baseline values. In contrast, in rhesus macaques, only a minor rise in

IP-10 levels was observed at the same time interval. In the same time span, but in both

macaque species, the levels of chemokine MCP-1 (CCL2) increased. Levels of other

chemokines, like eotaxin (CCL11), MIP-1α (CCL3), MIP-1β (CCL4), but also the cytokines IL-6

and IFN-γ dropped in the first week after infection and started rising around two weeks pi.

This decline of particular serum cytokine and chemokine levels coincided with the time period

that viral RNA was detectable in nasal and tracheal swabs, indicative for virus production. A

different pattern was seen for RANTES (CCL5) and TNFα proteins; serum levels were high in

the first 2 weeks after infection and then started decreasing to undetectable levels. RANTES

serum concentrations also showed an initial drop very early after infection. This was most

prominent in the cynomolgus macaques where RANTES levels dropped to zero on day 1 pi.

and started to increase at day 6 pi. Post-infection levels of I-TAC (CXCL11), MIG (CXCL9), and

IL-8 (CXCL8) were below detection or were not influenced by the SARS-CoV-2 infection.

Discussion

In humans, COVID-19 was initially regarded as a respiratory disease, but now it is clear that

individuals that succumb to this disease can display a complex array of pathologies that cover

a broad spectrum of symptoms. To sort out the factors leading to the different COVID-19

manifestations, animal models are essential. Due to their similarity to humans, specifically,

non-human primates can play a pivotal role in this type of preclinical research. To best

appreciate the potential of the various macaque species as SARS-CoV-2 infection models, a

thorough characterization of the course of infection is needed. The comparative study

reported in this communication contributes to the knowledge, but more importantly validates

the macaque models that are currently in use in SARS-CoV-2 research.

Unlike in some human patients, we found no evidence for renal involvement or coagulation

disorders in our monkeys. Equally, increasing C-reactive protein (CRP) levels are a marker in

the early diagnosis of pneumonia [48], but different from humans; these levels were not

subject to change in the infected macaques during the 3-weeks monitoring period. A direct

comparison with humans is hampered by the fact that the macaques were infected, but

seemingly healthy, while most published findings in humans were obtained from COVID-19

patients in various stages of disease. Unbiased measurement of the body temperature and

activity of each animal was done using a telemetric device implanted in the abdomen.

Monitoring by telemetry is an important asset as in both macaque species a notable difference

in body temperature was recorded in the first 2 weeks (period of active virus replication) as

compared to the third week. Significant differences in animal activity indicated that SARS-CoV-

2 infection also influenced the well-being of the animals without causing obvious clinical

symptoms. Notably, the elevation in body temperature on the first day of infection, as was

reported by Munster *et al.*, was not seen in our study, possibly pointing to a potential

difference in the sampling method. While we used 24/7 monitoring, in the study described by

Munster *et al.*, temperature was measured only on selected days on anesthetized animals

[54]. However, variation can also be due to a difference between the challenge viruses used,

the methods used for virus inoculation, or the origin and adaptation of the animals used.

Antibody responses were detectable in all animals after infection. Interestingly, no IgM was

detected in 7 out of 8 animals. This result was confirmed by a second, in-house developed

IgM-ELISA, but instead, the SARS-CoV-2 S protein was used as coating antigen. This approach

excludes a technical flaw in the serological assay used. We cannot explain the lack of IgM

response to infection, but similar observations were made in an infection study using African

Green monkeys. Hartman *et al.* did detect IgM, but the titers were very low, not exceeding

$1Log_{10}$ above the assay limit of detection [26]. In contrast, the development of IgG accurately

followed the course of virus infection, as it became first detectable within one week after the

virus had become undetectable in sera by RT-PCR. This was in line with findings of others [25,

26, 29].

In a subset of COVID-19 patients, the acute respiratory syndrome coincides with a 'cytokine

storm' or hypercytokinemia, which eventually can result in multi-organ failure [55]. Patients

with severe COVID-19 symptoms on intensive care units had significantly elevated plasma

levels of proinflammatory cytokines, like IL-2, IL-7, IL-10, GSCF, IP-10, MCP-1, MIP-1α, and TNF-

α [56]. In this study, macaques did not show overt disease symptoms of COVID-19, but certain

cytokines, like IL-6, IFN-γ, MIP-1α, and MIP-1β increased in the plasma of both macaque

species, indicating the involvement of the chemokine system during SARS-CoV-2 infection.

The cytokine profiles after SARS-CoV-2 were highly comparable between species, except for

IP-10 and MCP-1, suggesting differential involvement of monocyte activation between the

two species. The similarity in cytokine response after SARS-CoV-2 infection contrasts with

observations made after infection of macaques with another respiratory virus, pandemic

H1N1 influenza [47]. In that study, macaque species-specific cytokine responses (IL-6, MCP-1,

IL-15, IL-1Ra, MIP-1α, and IL-8) were induced upon infection with pH1N1, highlighting the virus

type-specific reaction of the chemokine system.

Unlike most published studies, we decided not to conduct a necropsy on animals early, 4-5

days, post-infection. At that time point after infection, evidence was found for acute viral

interstitial pneumonia [30, 32, 34, 54]. Instead, we performed CT imaging to visualize lung

pathology induced by SARS-CoV-2. In humans, the sensitivity of CT scanning for lung pathology

is high (positive predictive value of 92%), but the type of lesions found are not COVID-19-

specific, and can also be observed in a number of other infectious and non-infectious diseases

[57, 58]. In this study, we used purpose-bred NHPs with a well-documented health status and

we could compare the scans with a CT obtained just before infection. Therefore, CT imaging

provides a valuable tool to specifically monitor the progression of COVID-19-related lung

pathology during the entire course of the study. Based on the criteria set to determine clinical

severity [59], the macaques in our panel featured moderate disease levels as all eight

individuals show levels of pneumonia. In another study using only cynomolgus macaques and

using CT imaging as well, lesions were found as early as 2 days post-infection in infected

animals [29]. Type-wise, the lung lesions described in that report were comparable to the ones

in this communication, but they tend to be located deeper in the lungs. An explanation for

this difference may be that the method of instillation of the virus is the underlying cause. Finch

et al. administered the virus into each bronchus by direct bilateral primary post-carinal

intrabronchial instillation, whereas we applied the virus relatively high in the trachea, just

below the vocal cords. For similar reasons as reported by the same research team [29], we did

not collect bronchoalveolar lavage (BAL) samples from our animals to avoid unwanted

interventions in the natural infection process caused by SARS-CoV-2. Studies performed at our

institute with respiratory viruses have shown that lung lavages can indeed significantly

influence the infection process (manuscript in preparation). Instead of BAL samples, we

collected tracheal swabs for virological analysis. The viral RNA loads, but also the temporal

pattern of detection in swabs samples were like those observed in BAL samples from other

studies [22, 24, 34, 38]. This demonstrates that tracheal swabs are a good alternative for BAL

sampling. In addition, the collection of tracheal swabs is a less invasive technique that causes

relatively minor discomfort to the animals.

In most SARS-CoV-2 studies in non-human primates, the animals are euthanized shortly after

infection in the first week, or after a period of 3 weeks. The animals from this study were not

euthanized to be able to perform re-infection studies or to monitor them for late clinical signs,

or co-morbidities related to COVID-19.

We conclude that the course of SARS-CoV-2 infection of both macaque species is highly

similar, indicating that they are equally suitable models to test vaccines and antivirals in a

preclinical setting for safety and efficacy. The macaque model for SARS-CoV-2 infection in

humans manifests important virological aspects of this disease in humans. Given their

immunological and physiological resemblance to humans, NHPs likely will continue to play a

pivotal role in research for both COVID-19 prophylactic and therapeutic treatments.

Materials and Methods

Ethics and Biosafety Statement

All housing and animal care procedures took place at the Biomedical Primate Research Centre

(BPRC) in Rijswijk, the Netherlands. The BPRC is accredited by the American Association for

Accreditation of Laboratory Animal Care (AAALAC) International and is compliant with

European directive 2010/63/EU as well as the "Standard for Humane Care and Use of

Laboratory Animals by Foreign Institutions" provided by the Department of Health and Human

Services of the US National Institutes of Health (NIH, identification number A5539-01). Upon

positive advice by the independent ethics committee (DEC-BPRC) the competent authorities

(CCD, Central Committee for Animal Experiments) issued a project license (license

AVD5020020209404). Approval to start was obtained after further assessment of the detailed

study protocol by the institutional animal welfare body (AWB) (in Dutch: Instantie voor

Dierenwelzijn, IvD). All animal handlings were performed within the Department of Animal

Science (ASD) according to Dutch law. ASD is regularly inspected by the responsible national

authority (Nederlandse Voedsel- en Warenautoriteit, NVWA), and the AWB.

Animals

Four Indian-origin rhesus macaques and four cynomolgus macaques were used in this study (S3 Table). All macaques were mature, outbred animals, purpose-bred, and housed at the BPRC. The animals were in good physical health with normal baseline biochemical and hematological values. All were pair-housed with a socially compatible cage-mate in cages of at least 4 m^3 with bedding to allow foraging and were kept on a 12-hour light/dark cycle. The monkeys were offered a daily diet consisting of monkey food pellets (Ssniff, Soest, Germany) supplemented with vegetables and fruit. Enrichment was provided daily in the form of pieces of wood, mirrors, food puzzles, and a variety of other homemade or commercially available enrichment products. Drinking water was available *ad libitum* via an automatic watering system. Animal Care staff provided daily visual health checks before infection, and twice-daily after infection. The animals were monitored for appetite, general behavior, and stool consistency. All possible precautions were taken to ensure the welfare and to avoid any discomfort to the animals. All experimental interventions (intratracheal and intranasal infection, swabs, blood samplings, and CT scans) were performed under anesthesia.

Virus stock

The animals were infected with the SARS-CoV-2 strain BetaCoV/BavPat1/2020. This strain was isolated from a patient who traveled from China to Germany, and an aliquot of a Vero E6 cell culture was made available through the European Virus Archive-Global (EVAg). The viral stock for the infection study was propagated on Vero E6 cells. For this study, a fifth passage virus stock was prepared with a titer of 3.2×10^6 TCID$_{50}$ per ml. The integrity of the virus stock was confirmed by sequence analysis.

Experimental infections and post-exposure study follow-up

Three weeks before the experimental infection, a Physiotel Digital device (DSI Implantable Telemetry, Data Sciences International, Harvard Bioscience, UK) was implanted in the abdominal cavity of each animal. This device allowed the continuous real-time measurement of the body temperature and the animals' activity remotely using telemetry throughout the study.

At day 0, all animals were exposed to a dose of 1×10^6 TCID$_{50}$ of SARS-CoV-2, diluted in 5 ml phosphate buffered saline (PBS). The virus was inoculated via a combination of the intratracheal route (4.5 ml) and intranasal route (0.25 ml per nostril). Virus infection was monitored for 22 days, during which period the animals were checked twice-daily by the animal caretakers and scored for clinical symptoms according to a previously published, adapted scoring system [60] (S2 Table). A numeric score of 35 or more per observation time point was predetermined to serve as an endpoint and justification for euthanasia. Every time an animal was sedated, the body weight was measured. Blood was collected using standard aseptic methods from the femoral vein at regular time points post-infection (pi). In parallel, tracheal, nasal, and anal swabs were collected using Copan FLOQSwabs (MLS, Menen, Belgium). Swabs were placed in 1 ml DMEM, supplemented with 0.5% bovine serum albumin (BSA), fungizone (2.5 µg/ml), penicillin (100 U/ml), and streptomycin (100 µg/ml) and directly transported to the BSL3 lab.

Biochemistry and hematology analysis

Clinical biochemistry was performed using a Vetscan VS2 Chemical analyzer (Zoetis Benelux, Capelle aan de IJssel, The Netherlands) with the use of the Comprehensive Diagnostic profile.

This profile allows testing for alanine aminotransferase, albumin, alkaline phosphatase,

amylase, calcium, creatinine, globulin, glucose, phosphorus, potassium, sodium, total

bilirubin, total protein, and blood urea nitrogen. Hematology was performed using a Vetscan

HM5 Hematology analyser (Zoetis Benelux, Capelle aan de IJssel, The Netherlands). C-reactive

protein and D-dimer levels were measured using Cobas Integra 400 plus analyzer (Roche

Diagnostics Nederland B.V.).

Viral RNA detection

Viral RNA was isolated from plasma and swab sample supernatants using a QIAamp Viral RNA

Mini kit (Qiagen Benelux BV, Venlo, The Netherlands) following the manufacturer's

instructions. Viral RNA was reverse-transcribed to cDNA using a Transcriptor First Strand cDNA

Synthesis kit (Roche Diagnostics BV, Almere, The Netherlands). Viral RNA was quantified by

real-time quantitative RT-PCR specific for the RdRp gene of SARS-CoV-2, as described by

Corman *et al*. [61]. The lower detection limit of the qRT-PCR was 3.6 viral RNA copies per

reaction.

Computed tomography

Computed tomography (CT) data were acquired on several time points until day 22 (D0, 2, 4,

6, 8, 10, 12, 14, 16, 22 post-infection) using a MultiScan Large Field of View Extreme Resolution

Research Imager (LFER) 150 PET-CT (Mediso Medical Imaging Systems Ltd., Budapest,

Hungary). Animals were sedated with ketamine (10 mg/kg ketamine hydrochloride (Alfasan

Nederland BV, Woerden, The Netherlands) combined with medetomidine hydrochloride, 0.05

mg/kg (Sedastart; AST Farma B.V., Oudewater, The Netherlands) to induce sedation and

muscle relaxation, both applied intramuscularly (IM). The monkeys were positioned head first

supine (HFS) with the arms up and fixated in a vacuum pillow. A single CT of the thorax takes

35 seconds by which respiratory motion is inevitable, therefore, to mitigate the impact of

respiratory motion and improve the image quality, respiratory gating was applied. The

respiratory amplitude was detected with a gating pad placed next to the belly button. At the

end of the procedure, when the macaques returned to their home cage, atipamezole

(Sedastop; AST Farma B.V., Oudewater, The Netherlands) was given IM (0.25 mg/kg).

For the final reconstruction, the expiration phases were exclusively used and manually

selected. A semi-quantitative scoring system for chest CT evaluation was used to estimate

SARS-CoV-2-induced lung disease [29, 62]. Quantification of the CTs was performed

independently by two persons based on the sum of the lobar scores. The degree of

involvement in each zone was scored as: 0 for no involvement, 1 for <5%, 2 for 5-24%, 3 for

25-49%, 4 for 50-74% and 5 for >=75% involvement. An additional increase or decrease of 0.5

was used to indicate alterations in CT density of the lesions. By using this scoring system, a

maximum score of 35 could be reached for the combined lobes per time point.

Assessment of cytokine and chemokine protein levels in serum

Cytokine and chemokine concentrations in sera of infected macaques, including IL-1β, IL-6,

CCL11 (Eotaxin), CXCL10 (IP-10), CXCL11 (I-TAC), CCL2 (MCP-1), CXCL9 (MIG), CCL3 (MIP-1α),

CCL4 (MIP-1β), CCL5 (RANTES), CXCL8 (IL-8), TNFα, and IFNγ, were determined using

LEGENDplex™ NHP Chemokine/Cytokine Panel (13-plex) (Biolegend, San Diego, CA, USA)

according to manufacturer's instruction. Samples were measured on an LSRII FACS machine

and analyzed by using company software.

Antibody response

The total antibody response in macaque sera was analyzed in a double recognition enzyme-

linked immunosorbent assay (DR-ELISA) that detects total immunoglobulins in serum and

targeted to the SARS-CoV-2 nucleoprotein (N) protein as described by Hoste *et al.* [63]

(INgezim COVID19 DR; Eurofins-INGENASA, Madrid, Spain). Additionally, two in-house indirect

ELISAs were used to detect monkey immunoglobulin G (IgG) and immunoglobulin M (IgM)

directed to SARS-CoV-2 N protein. Briefly, Corning® 96-Well High-Binding Flat-Bottom

Microplates were coated with N protein and incubated overnight at 4°C in carbonate buffer,

pH 9.6. After washing the wells with PBS pH 7.4/0.05% Tween 20 (PBST), a blocking step was

performed with StabilZyme® SELECT Stabilizer (SurModics, Inc., Eden Prairie, MN, USA) for 1h

at room temperature (RT). The plate was then incubated with serum samples diluted 1:100 in

PBST with 2.5% fetal bovine serum (Gibco; Thermo Fisher Scientific, Inc., Waltham, MA, USA)

for 1h at RT. Two positive, cut-off, and negative controls were added to each plate. For

macaque-IgG detection, the wells were washed as described above and incubated with

Peroxidase-conjugated AffiniPure Goat Anti-Human IgG, Fcγ Fragment Specific (Jackson

ImmunoResearch Laboratories, Inc., PA, USA) diluted 1:25 000 in StabilZyme® HRP Conjugate

Stabilizer (SurModics), supplemented with 0.5M NaCl for 1h at RT. Finally, after a washing

step, the plate was incubated for 15 min with the substrate (TMB-MAX, Neogen Corporation,

Lexington, KY, USA) and the reaction was stopped by addition of 0.5 M sulfuric acid. The

absorbance was measured at 450 nm using a SpectraMax M5 plate reader (Molecular Devices,

LLC., San Jose, CA, USA). To detect macaque-IgM, the same protocol was followed, but the

secondary antibody used was an anti-Monkey IgM (µ-chain specific)-Peroxidase antibody

produced in goat (Sigma-Aldrich, Merck KGaA, MI, USA) at 1:5000 dilution.

Acknowledgements

We would like to thank the Animal Science Department, especially the animal caretakes and

veterinarians for excellent care of the animals. We also thank Cristina Aira (Eurofins Ingenasa,

Madrid) for technical assistance.

References:

1.   Chan JF, Yuan S, Kok KH, To KK, Chu H, Yang J, et al. A familial cluster of pneumonia
associated with the 2019 novel coronavirus indicating person-to-person transmission: a study
of a family cluster. Lancet. 2020;395(10223):514-23. Epub 2020/01/28. doi: 10.1016/S0140-
6736(20)30154-9. PubMed PMID: 31986261; PubMed Central PMCID: PMCPMC7159286.
2.   Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, Erichsen S, et al. SARS-
CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease
Inhibitor. Cell. 2020;181(2):271-80 e8. Epub 2020/03/07. doi: 10.1016/j.cell.2020.02.052.
PubMed PMID: 32142651; PubMed Central PMCID: PMCPMC7102627.
3.   Hamming I, Timens W, Bulthuis ML, Lely AT, Navis G, van Goor H. Tissue distribution of
ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding SARS
pathogenesis. J Pathol. 2004;203(2):631-7. Epub 2004/05/14. doi: 10.1002/path.1570.
PubMed PMID: 15141377; PubMed Central PMCID: PMCPMC7167720.
4.   Verity R, Okell LC, Dorigatti I, Winskill P, Whittaker C, Imai N, et al. Estimates of the
severity of coronavirus disease 2019: a model-based analysis. Lancet Infect Dis.
2020;20(6):669-77. Epub 2020/04/03. doi: 10.1016/S1473-3099(20)30243-7. PubMed PMID:
32240634; PubMed Central PMCID: PMCPMC7158570.
5.   Berlin DA, Gulick RM, Martinez FJ. Severe Covid-19. N Engl J Med. 2020. Epub
2020/05/16. doi: 10.1056/NEJMcp2009575. PubMed PMID: 32412710.
6.   Han C, Duan C, Zhang S, Spiegel B, Shi H, Wang W, et al. Digestive Symptoms in COVID-
19 Patients With Mild Disease Severity: Clinical Presentation, Stool Viral RNA Testing, and
Outcomes. Am J Gastroenterol. 2020;115(6):916-23. Epub 2020/04/18. doi:
10.14309/ajg.0000000000000664. PubMed PMID: 32301761; PubMed Central PMCID:
PMCPMC7172493.
7.   Paterson RW, Brown RL, Benjamin L, Nortley R, Wiethoff S, Bharucha T, et al. The
emerging spectrum of COVID-19 neurology: clinical, radiological and laboratory findings.
Brain. 2020. Epub 2020/07/09. doi: 10.1093/brain/awaa240. PubMed PMID: 32637987;
PubMed Central PMCID: PMCPMC7454352.
8.   Puntmann VO, Carerj ML, Wieters I, Fahim M, Arendt C, Hoffmann J, et al. Outcomes
of Cardiovascular Magnetic Resonance Imaging in Patients Recently Recovered From
Coronavirus Disease 2019 (COVID-19). JAMA Cardiol. 2020. Epub 2020/07/31. doi:
10.1001/jamacardio.2020.3557. PubMed PMID: 32730619; PubMed Central PMCID:
PMCPMC7385689.
9.   Peiris JS, Lai ST, Poon LL, Guan Y, Yam LY, Lim W, et al. Coronavirus as a possible cause
of severe acute respiratory syndrome. Lancet. 2003;361(9366):1319-25. Epub 2003/04/25.
doi: 10.1016/s0140-6736(03)13077-2. PubMed PMID: 12711465; PubMed Central PMCID:
PMCPMC7112372.
10.   McCray PB, Jr., Pewe L, Wohlford-Lenane C, Hickey M, Manzel L, Shi L, et al. Lethal
infection of K18-hACE2 mice infected with severe acute respiratory syndrome coronavirus. J
Virol. 2007;81(2):813-21. Epub 2006/11/03. doi: 10.1128/JVI.02012-06. PubMed PMID:
17079315; PubMed Central PMCID: PMCPMC1797474.
11.   Lutz C, Maher L, Lee C, Kang W. COVID-19 preclinical models: human angiotensin-
converting enzyme 2 transgenic mice. Hum Genomics. 2020;14(1):20. Epub 2020/06/06. doi:

10.1186/s40246-020-00272-6. PubMed PMID: 32498696; PubMed Central PMCID:
PMCPMC7269898.

12. Sanchez Felipe L, Vercruysse T, Sharma S, Ma J, Lemmens V, van Looveren D, et al. A
single-dose live-attenuated YF17D-vectored SARS-CoV2 vaccine candidate. Nature. 2020 *(in*
press).

13. Boudewijns R, Thibaut HJ, Kaptein SJF, Li R, Vergote V, Seldeslachts L, et al. STAT2
signaling as double-edged sword restricting viral dissemination but driving severe pneumonia
in SARS-CoV-2 infected hamsters. bioRxiv. 2020.

14. Enkirch T, von Messling V. Ferret models of viral pathogenesis. Virology. 2015;479-
480:259-70. Epub 2015/03/31. doi: 10.1016/j.virol.2015.03.017. PubMed PMID: 25816764;
PubMed Central PMCID: PMCPMC7111696.

15. Martina BE, Haagmans BL, Kuiken T, Fouchier RA, Rimmelzwaan GF, Van Amerongen
G, et al. Virology: SARS virus infection of cats and ferrets. Nature. 2003;425(6961):915. Epub
2003/10/31. doi: 10.1038/425915a. PubMed PMID: 14586458; PubMed Central PMCID:
PMCPMC7094990.

16. Kim YI, Kim SG, Kim SM, Kim EH, Park SJ, Yu KM, et al. Infection and Rapid Transmission
of SARS-CoV-2 in Ferrets. Cell Host Microbe. 2020. Epub 2020/04/08. doi:
10.1016/j.chom.2020.03.023
10.1016/j.chom.2020.03.023. PubMed PMID: 32259477.

17. Richard M, Kok A, de Meulder D, Bestebroer TM, Lamers MM, Okba NMA, et al. SARS-
CoV-2 is transmitted via contact and via the air between ferrets. Nat Commun.
2020;11(1):3496. Epub 2020/07/10. doi: 10.1038/s41467-020-17367-2. PubMed PMID:
32641684; PubMed Central PMCID: PMCPMC7343828.

18. Matute-Bello G, Frevert CW, Martin TR. Animal models of acute lung injury. Am J
Physiol Lung Cell Mol Physiol. 2008;295(3):L379-99. Epub 2008/07/16. doi:
10.1152/ajplung.00010.2008. PubMed PMID: 18621912; PubMed Central PMCID:
PMCPMC2536793.

19. Estes JD, Wong SW, Brenchley JM. Nonhuman primate models of human viral
infections. Nat Rev Immunol. 2018;18(6):390-404. Epub 2018/03/21. doi: 10.1038/s41577-
018-0005-7. PubMed PMID: 29556017; PubMed Central PMCID: PMCPMC5970954.

20. Cleary SJ, Pitchford SC, Amison RT, Carrington R, Robaina Cabrera CL, Magnen M, et al.
Animal models of mechanisms of SARS-CoV-2 infection and COVID-19 pathology. Br J
Pharmacol. 2020. Epub 2020/05/29. doi: 10.1111/bph.15143. PubMed PMID: 32462701;
PubMed Central PMCID: PMCPMC7283621.

21. Greenough TC, Carville A, Coderre J, Somasundaran M, Sullivan JL, Luzuriaga K, et al.
Pneumonitis and multi-organ system disease in common marmosets (Callithrix jacchus)
infected with the severe acute respiratory syndrome-associated coronavirus. Am J Pathol.
2005;167(2):455-63. Epub 2005/07/29. doi: 10.1016/S0002-9440(10)62989-6. PubMed PMID:
16049331; PubMed Central PMCID: PMCPMC1603565.

22. Yu P, Xu Y, Deng W, Bao L, Huang L, Xu Y, et al. Comparative pathology of rhesus
macaque and common marmoset animal models with Middle East respiratory syndrome
coronavirus. PLoS One. 2017;12(2):e0172093. Epub 2017/02/25. doi:
10.1371/journal.pone.0172093. PubMed PMID: 28234937; PubMed Central PMCID:
PMCPMC5325479.

23. Lu S, Zhao Y, Yu W, Yang Y, Gao J, Wang J, et al. Comparison of nonhuman primates
identified the suitable model for COVID-19. Signal Transduct Target Ther. 2020;5(1):157. Epub

2020/08/21. doi: 10.1038/s41392-020-00269-6. PubMed PMID: 32814760; PubMed Central
PMCID: PMCPMC7434851.

24. Singh A, Singh RS, Sarma P, Batra G, Joshi R, Kaur H, et al. A Comprehensive Review of
Animal Models for Coronaviruses: SARS-CoV-2, SARS-CoV, and MERS-CoV. Virol Sin.
2020;35(3):290-304. Epub 2020/07/02. doi: 10.1007/s12250-020-00252-z. PubMed PMID:
32607866; PubMed Central PMCID: PMCPMC7324485.

25. Woolsey C, Borisevich V, Prasad AN, Agans KN, Deer DJ, Dobias NS, et al. Establishment
of an African green monkey model for COVID-19. bioRxiv. 2020. Epub 2020/06/09. doi:
10.1101/2020.05.17.100289. PubMed PMID: 32511377; PubMed Central PMCID:
PMCPMC7263506.

26. Hartman AL, Nambulli S, McMillen CM, White AG, Tilston-Lunel NL, Albe JR, et al. SARS-
CoV-2 infection of African green monkeys results in mild respiratory disease discernible by
PET/CT imaging and shedding of infectious virus from both respiratory and gastrointestinal
tracts. PLoS Pathog. 2020;16(9):e1008903. Epub 2020/09/19. doi:
10.1371/journal.ppat.1008903. PubMed PMID: 32946524.

27. Gong SR, Bao LL. The battle against SARS and MERS coronaviruses: Reservoirs and
Animal Models. Animal Model Exp Med. 2018;1(2):125-33. Epub 2019/03/21. doi:
10.1002/ame2.12017
10.1002/ame2.12017. eCollection 2018 Jun. PubMed PMID: 30891557.

28. Sutton TC, Subbarao K. Development of animal models against emerging
coronaviruses: From SARS to MERS coronavirus. Virology. 2015;479-480:247-58. Epub
2015/03/21. doi: 10.1016/j.virol.2015.02.030
10.1016/j.virol.2015.02.030. Epub 2015 Mar 16. PubMed PMID: 25791336.

29. Finch CL, Crozier I, Lee JH, Byrum R, Cooper TK, Liang J, et al. Characteristic and
quantifiable COVID-19-like abnormalities in CT- and PET/CT-imaged lungs of SARS-CoV-2-
infected crab-eating macaques (Macaca fascicularis). bioRxiv. 2020. Epub 2020/06/09. doi:
10.1101/2020.05.14.096727. PubMed PMID: 32511338; PubMed Central PMCID:
PMCPMC7241101.

30. Rockx B, Kuiken T, Herfst S, Bestebroer T, Lamers MM, Oude Munnink BB, et al.
Comparative pathogenesis of COVID-19, MERS, and SARS in a nonhuman primate model.
Science. 2020;368(6494):1012-5. Epub 2020/04/19. doi: 10.1126/science.abb7314. PubMed
PMID: 32303590; PubMed Central PMCID: PMCPMC7164679.

31. Maisonnasse P, Guedj J, Contreras V, Behillil S, Solas C, Marlin R, et al.
Hydroxychloroquine use against SARS-CoV-2 infection in non-human primates. Nature. 2020.
Epub 2020/07/23. doi: 10.1038/s41586-020-2558-4. PubMed PMID: 32698191.

32. Chandrashekar A, Liu J, Martinot AJ, McMahan K, Mercado NB, Peter L, et al. SARS-
CoV-2 infection protects against rechallenge in rhesus macaques. Science.
2020;369(6505):812-7. Epub 2020/05/22. doi: 10.1126/science.abc4776. PubMed PMID:
32434946; PubMed Central PMCID: PMCPMC7243369.

33. Deng W, Bao L, Gao H, Xiang Z, Qu Y, Song Z, et al. Ocular conjunctival inoculation of
SARS-CoV-2 can cause mild COVID-19 in rhesus macaques. Nat Commun. 2020;11(1):4400.
Epub 2020/09/04. doi: 10.1038/s41467-020-18149-6. PubMed PMID: 32879306.

34. Williamson BN, Feldmann F, Schwarz B, Meade-White K, Porter DP, Schulz J, et al.
Clinical benefit of remdesivir in rhesus macaques infected with SARS-CoV-2. Nature.
2020;585(7824):273-6. Epub 2020/06/10. doi: 10.1038/s41586-020-2423-5. PubMed PMID:
32516797; PubMed Central PMCID: PMCPMC7486271.

35.   van Doremalen N, Lambe T, Spencer A, Belij-Rammerstorfer S, Purushotham JN, Port
JR, et al. ChAdOx1 nCoV-19 vaccine prevents SARS-CoV-2 pneumonia in rhesus macaques.
Nature. 2020. Epub 2020/07/31. doi: 10.1038/s41586-020-2608-y. PubMed PMID: 32731258.
36.   Gao Q, Bao L, Mao H, Wang L, Xu K, Yang M, et al. Development of an inactivated
vaccine candidate for SARS-CoV-2. Science. 2020;369(6499):77-81. Epub 2020/05/08. doi:
10.1126/science.abc1932.   PubMed   PMID: 32376603;   PubMed   Central   PMCID:
PMCPMC7202686.
37.   Erasmus JH, Khandhar AP, O'Connor MA, Walls AC, Hemann EA, Murapa P, et al. An
Alphavirus-derived replicon RNA vaccine induces SARS-CoV-2 neutralizing antibody and T cell
responses in mice and nonhuman primates. Sci Transl Med. 2020;12(555). Epub 2020/07/22.
doi: 10.1126/scitranslmed.abc9396. PubMed PMID: 32690628; PubMed Central PMCID:
PMCPMC7402629.
38.   Mercado NB, Zahn R, Wegmann F, Loos C, Chandrashekar A, Yu J, et al. Single-shot
Ad26 vaccine protects against SARS-CoV-2 in rhesus macaques. Nature. 2020. Epub
2020/07/31. doi: 10.1038/s41586-020-2607-z. PubMed PMID: 32731257.
39.   Yu J, Tostanoski LH, Peter L, Mercado NB, McMahan K, Mahrokhian SH, et al. DNA
vaccine protection against SARS-CoV-2 in rhesus macaques. Science. 2020;369(6505):806-11.
Epub 2020/05/22. doi: 10.1126/science.abc6284. PubMed PMID: 32434945; PubMed Central
PMCID: PMCPMC7243363.
40.   Patel A, Walters J, Reuschel EL, Schultheis K, Parzych E, Gary EN, et al. Intradermal-
delivered DNA vaccine provides anamnestic protection in a rhesus macaque SARS-CoV-2
challenge model. bioRxiv. 2020.
41.   Corbett KS, Flynn B, Foulds KE, Francica JR, Boyoglu-Barnum S, Werner AP, et al.
Evaluation of the mRNA-1273 Vaccine against SARS-CoV-2 in Nonhuman Primates. N Engl J
Med. 2020. Epub 2020/07/30. doi: 10.1056/NEJMoa2024671. PubMed PMID: 32722908;
PubMed Central PMCID: PMCPMC7449230.
42.   Yang J, Wang W, Chen Z, Lu S, Yang F, Bi Z, et al. A vaccine targeting the RBD of the S
protein of SARS-CoV-2 induces protective immunity. Nature. 2020. Epub 2020/07/30. doi:
10.1038/s41586-020-2599-8. PubMed PMID: 32726802.
43.   Hatziioannou T, Ambrose Z, Chung NP, Piatak M, Jr., Yuan F, Trubey CM, et al. A
macaque model of HIV-1 infection. Proc Natl Acad Sci U S A. 2009;106(11):4425-9. Epub
2009/03/04. doi: 10.1073/pnas.0812587106. PubMed PMID: 19255423; PubMed Central
PMCID: PMCPMC2657417.
44.   Zhou Y, Bao R, Haigwood NL, Persidsky Y, Ho WZ. SIV infection of rhesus macaques of
Chinese origin: a suitable model for HIV infection in humans. Retrovirology. 2013;10:89. Epub
2013/08/21. doi: 10.1186/1742-4690-10-89. PubMed PMID: 23947613; PubMed Central
PMCID: PMCPMC3765527.
45.   Foreman TW, Mehra S, Lackner AA, Kaushal D. Translational Research in the
Nonhuman Primate Model of Tuberculosis. ILAR J. 2017;58(2):151-9. Epub 2017/06/03. doi:
10.1093/ilar/ilx015. PubMed PMID: 28575319; PubMed Central PMCID: PMCPMC6279141.
46.   Cardona PJ, Williams A. Experimental animal modelling for TB vaccine development.
Int J Infect Dis. 2017;56:268-73. Epub 2017/02/07. doi: 10.1016/j.ijid.2017.01.030. PubMed
PMID: 28163168.
47.   Mooij P, Koopman G, Mortier D, van Heteren M, Oostermeijer H, Fagrouch Z, et al.
Pandemic Swine-Origin H1N1 Influenza Virus Replicates to Higher Levels and Induces More
Fever and Acute Inflammatory Cytokines in Cynomolgus versus Rhesus Monkeys and Can
Replicate in Common Marmosets. PLoS One. 2015;10(5):e0126132. Epub 2015/05/07. doi:

10.1371/journal.pone.0126132. PubMed PMID: 25946071; PubMed Central PMCID:
PMCPMC4422689.

48.   Wang L. C-reactive protein levels in the early stage of COVID-19. Med Mal Infect.
2020;50(4):332-4. Epub 2020/04/04. doi: 10.1016/j.medmal.2020.03.007. PubMed PMID:
32243911; PubMed Central PMCID: PMCPMC7146693.

49.   Ronco C, Reis T. Kidney involvement in COVID-19 and rationale for extracorporeal
therapies. Nat Rev Nephrol. 2020;16(6):308-10. Epub 2020/04/11. doi: 10.1038/s41581-020-
0284-7. PubMed PMID: 32273593; PubMed Central PMCID: PMCPMC7144544.

50.   Ronco C, Reis T, Husain-Syed F. Management of acute kidney injury in patients with
COVID-19. Lancet Respir Med. 2020;8(7):738-42. Epub 2020/05/18. doi: 10.1016/S2213-
2600(20)30229-0. PubMed PMID: 32416769; PubMed Central PMCID: PMCPMC7255232.

51.   Cheng Y, Luo R, Wang K, Zhang M, Wang Z, Dong L, et al. Kidney disease is associated
with in-hospital death of patients with COVID-19. Kidney Int. 2020;97(5):829-38. Epub
2020/04/06. doi: 10.1016/j.kint.2020.03.005. PubMed PMID: 32247631; PubMed Central
PMCID: PMCPMC7110296.

52.   Spiezia L, Boscolo A, Poletto F, Cerruti L, Tiberio I, Campello E, et al. COVID-19-Related
Severe Hypercoagulability in Patients Admitted to Intensive Care Unit for Acute Respiratory
Failure. Thromb Haemost. 2020;120(6):998-1000. Epub 2020/04/22. doi: 10.1055/s-0040-
1710018. PubMed PMID: 32316063; PubMed Central PMCID: PMCPMC7295272.

53.   Connors JM, Levy JH. COVID-19 and its implications for thrombosis and
anticoagulation. Blood. 2020;135(23):2033-40. Epub 2020/04/28. doi:
10.1182/blood.2020006000. PubMed PMID: 32339221; PubMed Central PMCID:
PMCPMC7273827.

54.   Munster VJ, Feldmann F, Williamson BN, van Doremalen N, Perez-Perez L, Schulz J, et
al. Respiratory disease in rhesus macaques inoculated with SARS-CoV-2. Nature.
2020;585(7824):268-72. Epub 2020/05/13. doi: 10.1038/s41586-020-2324-7. PubMed PMID:
32396922; PubMed Central PMCID: PMCPMC7486227.

55.   Coperchini F, Chiovato L, Croce L, Magri F, Rotondi M. The cytokine storm in COVID-
19: An overview of the involvement of the chemokine/chemokine-receptor system. Cytokine
Growth Factor Rev. 2020;53:25-32. Epub 2020/05/25. doi: 10.1016/j.cytogfr.2020.05.003.
PubMed PMID: 32446778; PubMed Central PMCID: PMCPMC7211650.

56.   Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected
with 2019 novel coronavirus in Wuhan, China. Lancet. 2020;395(10223):497-506. Epub
2020/01/28. doi: 10.1016/S0140-6736(20)30183-5. PubMed PMID: 31986264; PubMed
Central PMCID: PMCPMC7159299.

57.   Hope MD, Raptis CA, Shah A, Hammer MM, Henry TS, six s. A role for CT in COVID-19?
What data really tell us so far. Lancet. 2020;395(10231):1189-90. Epub 2020/04/01. doi:
10.1016/S0140-6736(20)30728-5. PubMed PMID: 32224299; PubMed Central PMCID:
PMCPMC7195087.

58.   Simpson S, Kay FU, Abbara S, Bhalla S, Chung JH, Chung M, et al. Radiological Society
of North America Expert Consensus Statement on Reporting Chest CT Findings Related to
COVID-19. Endorsed by the Society of Thoracic Radiology, the American College of Radiology,
and RSNA - Secondary Publication. J Thorac Imaging. 2020;35(4):219-27. Epub 2020/04/24.
doi: 10.1097/RTI.0000000000000524. PubMed PMID: 32324653; PubMed Central PMCID:
PMCPMC7255403.

59.   Fan L, Liu S. CT and COVID-19: Chinese experience and recommendations concerning
detection, staging and follow-up. Eur Radiol. 2020;30(9):5214-6. Epub 2020/05/08. doi:

10.1007/s00330-020-06898-3. PubMed PMID: 32377812; PubMed Central PMCID:
PMCPMC7202679.
60. Brining DL, Mattoon JS, Kercher L, LaCasse RA, Safronetz D, Feldmann H, et al. Thoracic
radiography as a refinement methodology for the study of H1N1 influenza in cynomologus
macaques (Macaca fascicularis). Comp Med. 2010;60(5):389-95. Epub 2011/01/26. PubMed
PMID: 21262125; PubMed Central PMCID: PMCPMC2958208.
61. Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DK, et al. Detection of
2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. Euro Surveill. 2020;25(3). Epub
2020/01/30. doi: 10.2807/1560-7917.ES.2020.25.3.2000045. PubMed PMID: 31992387;
PubMed Central PMCID: PMCPMC6988269.
62. Pan F, Ye T, Sun P, Gui S, Liang B, Li L, et al. Time Course of Lung Changes at Chest CT
during Recovery from Coronavirus Disease 2019 (COVID-19). Radiology. 2020;295(3):715-21.
Epub 2020/02/14. doi: 10.1148/radiol.2020200370. PubMed PMID: 32053470; PubMed
Central PMCID: PMCPMC7233367.
63. Hoste ACR, Venteo A, Fresco-Taboada A, Tapia I, Monedero A, Lopez L, et al. Two
serological approaches for detection of antibodies to SARS-CoV-2 in different scenarios: a
screening tool and a point-of-care test. Diagn Microbiol Infect Dis. 2020;98(4):115167. Epub
2020/09/06. doi: 10.1016/j.diagmicrobio.2020.115167. PubMed PMID: 32890908; PubMed
Central PMCID: PMCPMC7417941.
740
741

Fig 1. Virus load in swab samples. (A) Viral RNA quantification in tracheal and nasal swabs of

rhesus and cynomolgus macaques by qRT-PCR. The limit of quantification (154 RNA

copies/ml) is indicated by the dotted horizontal line. (B) Total virus loads in throat and nose

samples of macaques throughout the study. Horizontal bars represent geometric means. The

sum of the viral copies was calculated rather than area under the curve (AUC), as AUC

interpolates for time points when virus loads were not determined.

The different colors used for each animal as shown in the legend of 1A are used to denote

the same individual in all figures of this manuscript. The group of rhesus macaques is

indicated by yellow to red colors; cynomolgus macaques by green to blue. In each graph

rhesus macaques are depicted left and cynomolgus macaques right.

Fig 2. Body temperature during the study. The body temperature was measured by

telemetry throughout the study. The daily average body temperature of rhesus and

cynomolgus macaques was calculated and the deviations from baseline body temperature

(in °C) are depicted. Measurements done during biotechnical handlings were omitted from

the calculations.

Fig 3. Cumulative clinical scores. The cumulative clinical scores were calculated per week

and per individual animal (day 1-7, 8-14 and 15-21). Horizontal bars represent medians.

764
765 **Fig 4. Types of lung lesions detected via CT scans in SARS-CoV-2-infected macaques. (**A)

766 Ground glass opacities (GGO), (B) consolidations, and (C) crazy paving patterns (CCP).

Fig 5. Cumulative CT scores. Cumulative CT scores for each animal were calculated based on

the CT scores depicted in Table 1.

Fig 6. Development of SARS-CoV-2 antibody response in rhesus and cynomolgus macaques. The humoral immune response was determined using (A) DR-ELISA measuring the total antibody response, (B) an IgG-specific IgG assay, and (C) an IgM-specific IgM serological test. Results are shown as S/P: sample to positive control ratio:

$$S/P = \frac{test\ sample - mean\ negative\ control}{mean\ positive\ control - mean\ negative\ control} \text{ and } Units = \frac{test\ sample\ at\ x\ dpi}{test\ sample\ at\ 0\ dpi}.$$