Viscophobic turning regulates microalgae transport in viscosity gradients

Michael R. Stehnach[1,†], Nicolas Waisbord[1,†], Derek M. Walkama[1,2] and Jeffrey S. Guasto[1,*]

[1]*Department of Mechanical Engineering, Tufts University,*
200 College Avenue, Medford, MA 02155, USA
[2]*Department of Physics and Astronomy, Tufts University,*
574 Boston Avenue, Medford, Massachusetts 02155, USA
[†]*These authors contributed equally.* [*]*Corresponding author.*

Gradients in fluid viscosity characterize microbiomes ranging from mucus layers on marine organisms[1] and human viscera[2,3] to biofilms[4]. While such environments are widely recognized for their protective effects against pathogens and their ability to influence cell motility[2,5], the physical mechanisms controlling cell transport in viscosity gradients remain elusive[6–8], primarily due to a lack of quantitative observations. Through microfluidic experiments with a model biflagellated microalga (*Chlamydomonas reinhardtii*), we show that cells accumulate in high viscosity regions of weak gradients as expected, stemming from their locally reduced swimming speed. However, this expectation is subverted in strong viscosity gradients, where a novel viscophobic turning motility – consistent with a flagellar thrust imbalance[9,10] – reorients the swimmers down the gradient and causes striking accumulation in low viscosity zones. Corroborated by Langevin simulations and a three-point force model of cell propulsion, our results illustrate how the competition between viscophobic turning and viscous slowdown ultimately dictates the fate of population scale microbial transport in viscosity gradients.

Microbial transport is driven by external cues[5,13] and plays a crucial role in regulating the function of ecosystems ranging from cell colonies[4] to the Earth's biome[14]. Some cues are directly linked to a microorganism's metabolic activity, including gradients in chemical concentration[3] and light intensity[15] that guide cells toward regions of optimal nutrient uptake and photosynthetic productivity, respectively. Alternatively, external cues can serve as proxies to facilitate microbial navigation toward favorable conditions within a biome: Gravitational fields guide marine microbes within the water column[16] and magnetic fields serve as a compass for some cells in sedimentary environments[17]. Directed motility in response to an external cue is termed a "-taxis", which cells predominantly achieve through either (*i*) a response to a sensory receptor signal via an internal pathway (e.g. chemotaxis[18], phototaxis[15]) or (*ii*) a passive mechanical response to external forces or torques acting upon a cell or its organelles (e.g. gravitaxis[19], magnetotaxis[17]). Beyond these canonical examples, swimming cells are widely known to respond to gradients in fluid viscosity[5]. Viscous mucus layers serve as the first line of defense against motile pathogenic bacteria in many organisms[1], including the human body[2,20], and they sustain endangered coral ecosystems by trapping and distributing microorganisms and nutrient particles[1]. However, little is known about the physical mechanisms regulating cell swimming in viscosity gradients nor their implications for population scale transport. Despite its recognized importance for cell migration, no clear paradigm has emerged for viscotactic motility.

Contrary to the widely studied effects of elevated homogeneous viscosity on cell motility[11,21], experimental observations of swimming cell transport in inhomogeneous viscous environments are severely lacking and largely inconclusive. In viscosity gradients of unspecified strength, both positive[6,7] and negative[8] (i.e. viscophobic) viscotaxis have been observed toward high and low viscosity regions, respectively. These qualitative observations were based predominantly on microbial density and ignored the potential for purely mechanical explanations through cell swimming kinematics. In contrast, recent theoretical and numerical work[9,10,22] explored the microhydrodynamics of swimming in a viscosity gradient, which showed that tuning the propulsion strategy or body asymmetry could produce swimming either up or down the gradient. While these early experiments and deterministic simulations have provided important insights, a wholistic understanding of viscotactic motility has been hampered by the dearth of quantitative experimental evidence and assimilation with the established stochasticity of active matter transport[12].

In this Letter, we provide the first direct evidence of viscophobic motility for the model, biflagellated, microalga *Chlamydomonas reinhardtii*[23,24] (Fig. 1a), which is achieved through a comprehensive, experimental description of its transport in a prescribed, microfluidic viscosity gradient (Fig. 1b). Our experiments show that as the magnitude of the gradient is increased, cell accumulation surprisingly shifts from the high viscosity region (Fig. 1c,d) to the low viscosity region (Fig. 1e). Analysis of cell trajectories and complementary Langevin simulations reveals that the transition results from the competition between viscous accumulation due to reduced swimming speed[12,25] and a novel viscophobic reorientation mechanism. The latter is consistent with a physical, hydrodynamic phenomenon[10], which results in an ensemble drift of swimming *C. rein-*

Fig. 1: **Viscosity gradient strength regulates dispersal of swimming microalgae. a**, Biflagellated swimming microalga, *Chlamydomonas reinhardtii*. Scale bar, 5 μm. **b**, Schematic of three-inlet microfluidic device for viscosity gradient generation (100 μm deep). Viscous media (PEO in minimal M1 media), *C. reinhardtii* suspension, and M1 media are initially flow-stratified (Extended Data Fig. 1 and Methods), and the viscosity gradient is formed by polymer diffusion upon stopping the flow (Extended Data Fig. 4). **c-e**, Multi-exposure images (0.5 s duration; intensity increases with time; 60 min post flow) of *C. reinhardtii* show a homogeneous distribution in uniform viscosity (**c**; $\nabla \eta = 0$ cP·μm^{-1}, control), accumulation in high viscosity zone for moderate viscosity gradient (**d**; $\nabla \eta = 3.4 \times 10^{-3}$ cP·μm^{-1}), and accumulation in low viscosity zone for large gradient (**e**; $\nabla \eta = 7.2 \times 10^{-3}$ cP·μm^{-1}). Colorbar is measured viscosity profile from **g**. Scale bar, 100 μm. **f**, Ensemble-averaged cell swimming speed, $\langle V \rangle$, in bulk viscous media, η (Supplementary Section 3.1). Error bars are mean standard deviation (N = 2-3) and black line is a power-law fit[11] ($\propto \eta^{-0.93}$). **g**, Microfluidic viscosity profiles measured by microrheology (see Methods), $\eta(x)$, characterized by the viscosity gradient magnitude, $\nabla \eta$. Shaded area is standard deviation (30-90 min post flow). **h-j**, Measured profiles of local mean cell swimming speed (**h**), cell density (**i**), and flux magnitude (**j**), corresponding to **g**. Shaded regions indicate standard error (N = 3-4). Dotted lines are predictions for random walk swimmers (Supplementary Section 4.1), where deviation of experiments in **j** indicates biased transport[12].

hardtii down the viscosity gradient. Viscophobic reorientation dynamics bear striking similarities to a range of other taxes[13,16,17] and add to the rich diversity of microbial transport mechanisms that regulate important macroscopic ecological processes[1–4].

To explain the shift of migratory behavior observed in *C. reinhardtii*, microfluidic devices were designed to precisely control the viscosity gradient over length scales relevant to the swimming cells (Fig. 1b). Briefly, three solutions, including a Newtonian viscous polymer solution (polyethylene oxide, PEO), a dilute suspension of *C. reinhardtii*, and cell media, are stratified by continuous injection into the microchannel test section (Extended Data Fig. 1 and Methods). Upon stopping the flow, the polymer diffuses across the strata for 30 min to form a smooth viscosity gradient (Extended Data

Fig. 4); thus providing ample time for swimming cells to disperse across the channel. Observations are performed between 30 and 90 min, during which time the viscosity profile, $\eta(x)$, quantified by microrheology, is quasi-steady (Fig. 1g and Methods). In the absence of a viscosity gradient, their random walk motility yields a uniform cell density, ρ, across the channel ($\nabla \eta = 0$ cP·μm^{-1}; Fig. 1i).

For a directionally unbiased random walk, cell density is well known to scale inversely with local swimming speed[12,25,26], $\rho(x) \propto 1/V(x)$. Accordingly, imposing a moderate viscosity gradient ($\nabla \eta \leq 3.4 \times 10^{-3}$ cP·μm^{-1}) results in locally reduced swimming speed (Fig. 1h, solid lines) and accumulation of cells (Fig. 1i, solid lines) in the high viscosity region. The swimming speed distribution quantitatively agrees with predictions (Fig.

Fig. 2: **Coupling of swimming kinematics to viscosity gradient induces viscophobic turning. a,b,** Measured cell trajectories for control (**a**; $\nabla\eta = 0$ cP·μm^{-1}) and maximum viscosity gradient (**b**; $\nabla\eta = 7.2 \times 10^{-3}$ cP·μm^{-1}). Conditionally sampled from $400 \leq x \leq 600$ μm and perpendicular to viscosity gradient ($|\theta| = \pi/2 \pm 5°$, dashed lines). Thick colored lines are mean trajectories, and black curve in **b** is analytical prediction (Supplementary Section 4.3). Background qualitatively indicates $\nabla\eta$ and t^* is characteristic swimming time in channel ($t^* = 500$ μm/$\langle V(x)\rangle$). Scale bar, 175 μm. **c,** Angular cell velocity exhibits orientation dependence in a viscosity gradient ($\nabla\eta = 7.2 \times 10^{-3}$ cP·μm^{-1}, orange) compared to control ($\nabla\eta = 0$ cP·μm^{-1}, black). Dashed lines are sinusoidal fits, and shaded region is standard error. **d,** Viscophobic turning rate amplitude, ω_{visc}, (**c**) increases with normalized viscosity gradient, $\nabla\eta/\eta_0$. Black line is linear fit, horizontal error bars stem from measured viscosity profile, and vertical error bars are from fit in **c**. Inset: Potential mechanism for orientation-dependent viscophobic turning via propulsive force imbalance between flagella (Supplementary Section 4.2). Background and arrow indicate increasing viscosity gradient.

1h, dotted lines) based on the measured swimming speed of *C. reinhardtii* in homogeneous conditions[11] (Fig. 1f) and the independently measured viscosity profile (Fig. 1g and Supplementary Section 4.1). However, this theoretical slowdown accumulation largely over-predicts the observed positive viscotactic behavior (Fig. 1i, dotted line). Stronger gradients reveal a counterintuitive cell accumulation in the low viscosity region (Fig. 1i, $\nabla\eta = 7.2 \times 10^{-3}$ cP·μm^{-1}), establishing a second, viscophobic regime. Moreover, the local cell flux magnitude, $\rho(x)V(x)$, is expected to be spatially uniform across the channel for an unbiased random walk, but rather is progressively skewed toward low viscosity (Fig. 1j and Supplementary Section 4.1). Taken together, these results suggest the presence of a novel underlying viscophobic motility. To unravel these population scale observations and determine the nature of the observed behavior – whether a sensory response or a passive physical mechanism – we examine the kinematics of single cell motility in the viscosity gradients.

Cells swimming in a viscosity gradient exhibit directional dependence on their rotational kinematics and ultimately reorient down the viscosity gradient. In the absence of a viscosity gradient, cell trajectories conditionally sampled from the channel center and oriented perpendicular to the gradient ($\theta = \pm\pi/2$) generally maintain their initial swimming direction (Fig. 2a). Conversely, in the largest viscosity gradient investigated, the trajectories continuously bend toward the low viscosity region ($\theta = 0$; Fig. 2b). Due to the stochastic nature of microbial motility, for example stemming from noise in flagellar actuation and thermal fluctuations[27,28] (persistence time, $\tau_{\text{p}} = 1/2D_{\text{r}} = 7.2$ s; rotational diffusion coeffi-

cient, D_{r}; see Methods), a large number of observations are necessary to unveil the underlying deterministic dynamics. Considering all swimming directions, the cells' instantaneous angular velocity displays a sinusoidal dependence on orientation, which is absent from control experiments (Fig. 2c). We hypothesize that viscophobic turning in *C. reinhardtii* results from a slight viscosity difference experienced by its two flagella, causing an imbalance of their drag-based thrust forces[29] (Fig. 2d, inset).

A three-point force swimmer model (Supplementary Section 4.2) quantitatively predicts the sinusoidal form of the observed cell rotation rate (Fig. 2c, dashed curves) and captures the mean swimming trajectory of the cells (Fig. 2b, black curve). The angular cell velocity is well fit by the model prediction (Fig. 2c, dashed curves), $\omega(\theta) = -\omega_{\text{visc}}\sin(\theta)$, and quantifies the turning rate amplitude, ω_{visc}. The amplitude grows continuously with the normalized viscosity gradient, $\nabla\eta/\eta_0$, where η_0 is the viscosity in the channel center, and the largest measured gradient yields a turning time, $\tau = 1/\omega_{\text{visc}} \approx 14$ s (Fig. 2d). Furthermore, the linear dependence[10] on $\nabla\eta/\eta_0$ (Fig. 2d, black line) stems from the growth of the turning rate with the gradient $\nabla\eta$, which is tempered by enhanced rotational drag through increasing η_0. A fit of the model prediction (Fig. 2d, black line; Supplementary Section 5.2), $\omega_{\text{visc}} = (V^*/2)\nabla\eta/\eta_0$, gives $V^* \approx 66$ μm/s, which is comparable to the swimming speed of *C. reinhardtii* in the tested gradients. Viscophobic reorientation of *C. reinhardtii* is thus a deterministic and omnipresent phenomenon, in line with the present hydrodynamic model and other recent squirmer and point force models[9,10]. However, viscotaxis exhibits

surprising similarities to a range of established taxes, including some with internal sensory pathways.

While viscophobic turning is rooted in swimmer hydrodynamics, the strikingly simple orientation dynamics comprise a broader, universal class of motilities. The time evolution of the measured cell orientation distribution, $p(\theta|t)$, illustrates the condensation of the cell swimming direction down the viscosity gradient (Fig. 3a, top and Extended Data Fig. 8). Similar to most microbes, the random walk nature of *C. reinhardtii* motility stems from noise in their orientation[27], modeled as

$$\dot{\theta} = -\omega_{\mathrm{visc}} \sin(\theta) + \sqrt{2D_r}\xi(t), \qquad (1)$$

where ξ is Gaussian noise. A Langevin simulation (see Methods) accurately reproduces the time evolution of cell orientation (Fig. 3a, bottom and Extended Data Fig. 8), which forms a well-defined peak after several seconds (Fig. 3b). Our observations of viscotactic turning are thus consistent with deterministic responses that underlie most eukaryotic systems[30], as opposed to the stochastic navigation of bacterial chemotaxis[18]. Importantly, the form of the dynamical equation is identical to that of other passive, physical taxes – including gravitaxis[16,19] and magnetotaxis[17] – which are governed by a sinusoidal, paramagnetic-like torque in response to an external field. However, viscophobic reorientation is distinctive in that self-propulsion is fundamental to the cell response, akin to chemotactic[3,18] and phototactic[15] sensing.

Beyond the observed universal orientation dynamics, viscophobic motility also exhibits a log-sensing response to the viscosity gradient. The condensation of cell orientation compares favorably to the known steady-state distribution[31] for equation (1), $p(\theta) \propto \exp[\omega_{\mathrm{visc}}/D_r \cos\theta]$. The directional bias is ultimately responsible for population-scale migration via a mean viscotactic drift velocity, $V\langle\cos\theta\rangle$. Borrowing from paramagnetism theory[17], the mean orientation is

$$\langle\cos(\theta)\rangle = \coth\left(\frac{\omega_{\mathrm{visc}}}{D_r}\right) - \frac{D_r}{\omega_{\mathrm{visc}}}, \qquad (2)$$

which scales as $\langle\cos(\theta)\rangle \propto \nabla(\ln\eta)$ in the limit of small $\omega_{\mathrm{visc}}/D_r$. Thus, the orientation exhibits log-sensitivity to the viscosity gradient, which is a prominent feature of chemosensing in bacteria[32,33] as well as some synthetic active matter systems[34]. The resulting viscophobic drift of *C. reinhardtii* toward low viscosity zones competes with accumulation in high viscosity due to locally reduced swimming speed. Understanding the interplay between these effects is crucial for determining the ultimate fate of the cell population.

To elucidate the relative contributions of viscophobic turning versus viscous slowdown to cell transport, we employ a simplified, two-dimensional Langevin model. The simulated cell orientational dynamics are set by equation (1), and the translational dynamics are given by $\dot{x} = V(x)\cos(\theta)$ and $\dot{y} = V(x)\sin(\theta)$ (Supplemen-

Fig. 3: **Simulations illustrate the competition between viscophobic drift and viscous slowdown driving cell accumulation.** **a**, Time evolution of conditional probability density of cell orientation ($\nabla\eta = 7.2 \times 10^{-3}$ cP·μm^{-1}) for experiment (top) and Langevin simulation (bottom), showing condensation of cell alignment down the viscosity gradient (Extended Data Fig. 8). Experimental cell tracks selected from $400 \leq x \leq 600$ μm. **b**, Orientation distribution at $t = 6$ s from experiments and Langevin simulations (**a**) agree with a Fokker-Planck model (Supplementary Section 5.3). **c**, Kullback-Leibler divergence (colormap; see Methods) of cell density between Langevin simulations, ρ, and theory without viscophobic turning, ρ_0 (Supplementary Section 4.1). Squares are best match density profile simulations with experiments (circles; Fig. 2d and Extended Data Fig. 7). Horizontal and vertical error bars propagated from Fig. 1h and Fig. 2d, respectively. Black curve is empirical scaling law (Supplementary Section 5.2), arrow indicates increasing $\nabla\eta$, and diamonds correspond to experimental conditions (Fig. 1).

tary Section 5.1). The essential features of the viscosity dependence are captured by a spatially varying (linear) cell swimming speed profile, $V(x)$, which is parameterized by the speed difference across the channel normalized by the maximal speed in the low viscosity

zone, $\Delta V/V_{\max}$. Resulting Langevin simulations that best match the experimental cell density profiles (Extended Data Fig. 7) faithfully reproduce the measured viscophobic turning rate (Fig. 3c, squares). The simulations also agree well with an empirical scaling law (Fig. 3c, black curve; Supplementary Section 5.2) based on the viscosity dependent swimming speed (Fig. 1f) and turning rate (Fig. 2d).

While viscophobic reorientation (Fig. 2d) and viscous slowdown (Fig. 1h) are intrinsically linked in experiments through the viscosity profile, simulations allow us to untangle these effects by independently varying ω_{visc} and $\Delta V/V_{\max}$. Density profiles from Langevin simulations, $\rho(x)$, and from theory based solely on the viscous slowdown, $\rho_0(x)$, are compared using the Kullback-Leibler divergence, $D_{\mathrm{KL}}(\rho|\rho_0)$, which provides an "entropic" measure of the impact of viscophobic turning on cell transport[35] (Fig. 3c, colormap and Methods). The sharp increase in D_{KL} with increasing viscophobic reorientation rate, ω_{visc}, is indicative of the progressive skewing of the cell density toward the low viscosity region (Extended Data Fig. 7). For a fixed ω_{visc}, the viscous slowdown causes cell accumulation on the viscous side, yet viscophobic motility moderates this trend more efficiently as the swimming speed contrast is increased. In real systems, the simultaneous increase of ω_{visc} and $\Delta V/V_{\max}$ (Fig. 3c, black arrow) indicates that viscophobicity dominates for large viscosity gradients. The growing influence of viscophobic motility with increasing gradient presents an ecologically valuable asset for the escape or repulsion of cells from viscous environments[1,2,5]. We anticipate that both viscophobic turning rate and swimming speed contrast – and thus, cell accumulation – will have a rich dependency on organism flagellation, inviting comparisons with diverse microbes.

Our work provides the first direct evidence of a fundamentally new viscotactic response of swimming microorganisms, where a novel interplay between self propulsion and a viscosity gradient reorients biflagellated microalgae toward low viscosity regions. Viscophobic turning shares universal dynamics that underpin a host of microbial responses to their physical environments, including magnetotaxis and gravitaxis[16,31]. Yet, the degree of alignment of *C. reinhardtii* to the viscosity gradient exhibits log-sensitivity, which has curious parallels to the sensory response of chemotactic microbes[33] and synthetic systems[34]. While swimming cells are generally expected to accumulate in viscous regions due to reduced swimming speed, our experimental and numerical results demonstrate that the orientational viscophobic bias overcomes this viscous slowdown in strong gradients and provides an escape to regions of lower viscosity. Complementary studies at the level of single flagella will be required to fully elucidate the potential for mechanosensing and the possible impact of viscosity gradients on flagellar beat patterns and coordination[11,27,28]. The discovery of viscophobic turning has the potential to refine our understanding of cell motility in complex microbiological systems, where the impact of viscosity gradients on cell motility have been widely overlooked, ranging from biofilms[4] to the protective mucus layers[2,3] to endangered coral ecosystems[1]. The experimental and analytical approaches established in this work will pave the way for understanding the manifestation of viscophobic motility in a range of important microorganisms and in complex fluid rheologies.

[1] Wild, C. *et al.* Coral mucus functions as an energy carrier and particle trap in the reef ecosystem. *Nature* **428**, 66–70 (2004).

[2] Wheeler, K. M. *et al.* Mucin glycans attenuate the virulence of *Pseudomonas aeruginosa* in infection. *Nat. Microbiol.* **4**, 2146–2154 (2019).

[3] Suarez, S. S. & Pacey, A. A. Sperm transport in the female reproductive tract. *Hum. Reprod. Update* **12**, 23–37 (2006).

[4] Hall-Stoodley, L., Costerton, J. W. & Stoodley, P. Bacterial biofilms: from the natural environment to infectious diseases. *Nat. Rev. Microbiol.* **2**, 95–108 (2004).

[5] Swidsinski, A. *et al.* Viscosity Gradient Within the Mucus Layer Determines the Mucosal Barrier Function and the Spatial Organization of the Intestinal Microbiota. *Inflamm. Bowel Dis.* **13**, 963–970 (2007).

[6] Petrino, M. G. & Doetsch, R. N. 'Viscotaxis', a New Behavioural Response of *Leptospira interrogans* (*biflexa*) Strain b16. *J. Gen. Microbiol.* **109**, 113–117 (1978).

[7] Daniels, M. J., Longland, J. M. & Gilbart, J. Aspects of Motility and Chemotaxis in Spiroplasmas. *J. Gen. Microbiol.* **118**, 429–436 (1980).

[8] Sherman, M. Y., Timkina, E. & Glagolev, A. Viscosity taxis in *Escherichia coli*. *FEMS Microbiol. Lett.* **13**, 137–140 (1982).

[9] Liebchen, B., Monderkamp, P., Ten Hagen, B. & Löwen, H. Viscotaxis: Microswimmer Navigation in Viscosity Gradients. *Phys. Rev. Lett.* **120**, 208002 (2018).

[10] Datt, C. & Elfring, G. J. Active Particles in Viscosity Gradients. *Phys. Rev. Lett.* **123**, 158006 (2019).

[11] Qin, B., Gopinath, A., Yang, J., Gollub, J. P. & Arratia, P. E. Flagellar Kinematics and Swimming of Algal Cells in Viscoelastic Fluids. *Sci. Rep.* **5**, 9190 (2015).

[12] Schnitzer, M. J. Theory of continuum random walks and application to chemotaxis. *Phys. Rev. E* **48**, 2553–2568 (1993).

[13] Stocker, R. Marine Microbes See a Sea of Gradients. *Science* **338**, 628–633 (2012).

[14] Cavicchioli, R. *et al.* Scientists' warning to humanity: microorganisms and climate change. *Nat. Rev. Microbiol.* **17**, 569–586 (2019).

[15] Jékely, G. *et al.* Mechanism of phototaxis in marine zooplankton. *Nature* **456**, 395–399 (2008).

[16] Sengupta, A., Carrara, F. & Stocker, R. Phytoplankton can actively diversify their migration strategy in response to turbulent cues. *Nature* **543**, 555–558 (2017).

[17] Blakemore, R. P. Magnetotactic bacteria. *Science* **190**, 377–379 (1975).

Berg, H. C. & Brown, D. A. Chemotaxis in *Escherichia coli* analysed by Three-dimensional Tracking. *Nature* **239**, 500–504 (1972).

Durham, W. M., Kessler, J. O. & Stocker, R. Disruption of Vertical Motility by Shear Triggers Formation of Thin Phytoplankton Layers. *Science* **323**, 1067–1070 (2009).

Cone, R. A. Barrier properties of mucus. *Adv. Drug Deliv. Rev.* **61**, 75–85 (2009).

Martinez, V. A. *et al.* Flagellated bacterial motility in polymer solutions. *Proc. Natl Acad. Sci. USA* **111**, 17771–17776 (2014).

Shoele, K. & Eastham, P. S. Effects of nonuniform viscosity on ciliary locomotion. *Phys. Rev. Fluids* **3**, 043101 (2018).

Harris, E. H. *The Chlamydomonas Sourcebook: Introduction to Chlamydomonas and its laboratory use*, vol. 1 (Academic Press, 2009).

Goldstein, R. E. Green Algae as Model Organisms for Biological Fluid Dynamics. *Annu. Rev. Fluid Mech.* **47**, 343–375 (2015).

Frangipane, G. *et al.* Dynamic density shaping of photokinetic *E. coli*. *Elife* **7**, e36608 (2018).

Cates, M. E. Diffusive transport without detailed balance in motile bacteria: does microbiology need statistical physics? *Reports Prog. Phys.* **75**, 042601 (2012).

Polin, M., Tuval, I., Drescher, K., Gollub, J. P. & Goldstein, R. E. *Chlamydomonas* Swims with Two "Gears" in a Eukaryotic Version of Run-and-Tumble Locomotion. *Science* **325**, 487–490 (2009).

Geyer, V. F., Jülicher, F., Howard, J. & Friedrich, B. M. Cell-body rocking is a dominant mechanism for flagellar synchronization in a swimming alga. *Proc. Natl Acad. Sci. USA* **110**, 18058–18063 (2013).

Lauga, E. & Powers, T. R. The hydrodynamics of swimming microorganisms. *Reports Prog. Phys.* **72**, 096601 (2009).

Waisbord, N. & Guasto, J. S. Peculiar polygonal paths. *Nat. Phys.* **14**, 1161–1162 (2018).

Rupprecht, J. F., Waisbord, N., Ybert, C., Cottin-Bizonne, C. & Bocquet, L. Velocity Condensation for Magnetotactic Bacteria. *Phys. Rev. Lett.* **116**, 168101 (2016).

Block, S. M., Segall, J. E. & Berg, H. C. Adaptation kinetics in bacterial chemotaxis. *J. Bacteriol.* **154**, 312 – 323 (1983).

Saragosti, J. *et al.* Directional persistence of chemotactic bacteria in a traveling concentration wave. *Proc. Natl Acad. Sci. USA* **108**, 16235–16240 (2011).

Theurkauff, I., Cottin-Bizonne, C., Palacci, J., Ybert, C. & Bocquet, L. Dynamic Clustering in Active Colloidal Suspensions with Chemical Signaling. *Phys. Rev. Lett.* **108**, 268303 (2012).

Dechant, A. & Sasa, S.-i. Fluctuation–response inequality out of equilibrium. *Proc. Natl Acad. Sci. USA* **117**, 6430 – 6436 (2020).

Methods

Cell culturing and media preparation. *Chlamydomonas reinhardtii* (CC-1690 wild type mt+ (WT); www.chlamycollection.org) were grown in 100 mL of minimal media (M1)[36] at 22°C using gentle aeration and synchronized on a 14 hr on and 10 hr off light cycle. Cells were subcultured (10 mL) every three days at mid-exponential phase. Viscous media was prepared with polyethylene oxide (PEO; molecular weight, 400,000 g/mol; Sigma-Aldrich) in M1 (Extended Data Table 1) at PEO concentrations of 0%, 0.05%, 0.1%, 0.25%, 0.5%, and 0.82% (w/v), which remain Newtonian (overlap concentration, $c^* \approx 3\%$ (w/v))[37,38]. PEO solutions were mixed for 2 hrs at 40 rpm (multi-purpose tube rotator; Fisherbrand), and 0.2% (w/v) bovine serum albumen was added to all solutions to prevent sticking to microchannel walls.

Microfluidics. Microfluidic devices were made using standard soft lithography techniques[39]. Polydimethylsiloxane (PDMS; Dow Corning SYLGARD 184) channels were cast on photoresist (Microchem) molds, fabricated via photolithography, and plasma bonded to standard glass slides. Circular observation channels (diameter, 5 mm; height, 250 μm) were used for bulk cell motility and microrheology measurements. Viscosity gradient generation channels (Fig. 1b, Extended Data Fig. 1) were designed with three inlets (width, 0.5 mm) carrying a viscous PEO solution, cell suspension, and M1 media, respectively. The three solutions were flow stratified for a minimum of 2 min using two syringe pumps (Harvard Apparatus), where flow rates were adjusted to maintain a 4:1:4 ratio of the stream widths (Supplementary Section 1). Upon halting the flow, a monotonic viscosity profile, $\eta(x)$, slowly evolves over 90 min (Extended Data Fig. 4) due to the relatively small PEO diffusivity[40]. The thin central band of *C. reinhardtii*, bracketed by fresh M1, ensured no chemotactic bias was introduced[21] (Supplementary Section 3.2).

Microrheology and viscosity profile measurement. The viscosity of bulk Newtonian PEO solutions was measured through well-developed microrheology approaches[41] by seeding with dilute (0.5 μL/mL) tracer particles (radius, R = 0.25 μm; 2% solid; Life Technologies). Tracers were imaged using epifluorescence microscopy (10× 0.5NA objective; Nikon Ti-e) and recorded (Blackfly S, FLIR Systems) at 4 fps for 25 s. Mean square displacements (MSD) of 400-1,000 particles, $\langle \Delta x^2 \rangle$, were measured over time, Δt, via particle tracking. A linear fit (Extended Data Fig. 2) of the MSD, $\langle \Delta x^2 \rangle = 2D\Delta t$, yields the diffusivity, D, which was used to determine the viscosity from the Stokes-Einstein relation, $D = k_B T/6\pi\eta R$ (Boltzmann constant, k_B; absolute temperature, T). Measured bulk viscosities (Extended Data Table 1) compared well with previously reported results[37]. Spatially-resolved viscosity profiles, $\eta(x)$, generated by the viscosity gradient assay (Fig. 1g), were measured using an identical approach to bulk experiments, but particle tracks were partitioned into 11 equally spaced bins across the channel width (Extended Data Fig. 3). Time evolution of the viscosity profile was evaluated by acquiring video of fluorescent tracer motion (10× 0.2NA objective) for 250 frames (63 s long) every 10 minutes for 90 min (Zyla, Andor Technology). The local viscosity was measured from the MSD (Extended Data Fig. 3) in each spatially resolved bin over the 90 min experiment. Transient variations in viscosity rapidly decreased after 30 min (Extended Data Fig. 4). Thus, we analyze our viscotaxis assay results from 30-90 min after halting the flow, and we report the average viscosity profile over this time window (Fig. 1g). Viscosity gradient, $\nabla\eta$, is calculated from a linear fit to the viscosity profile in the middle third of the microchannel. The viscosity gradient for each viscotaxis experiment is the average gradient within the 30-90 min interval (Extended Data Table 2). See Supplementary Section 2 for details.

Cell imaging and tracking. Imaging for viscotaxis experiments with *C. reinhardtii* was performed using phase-contrast microscopy (10× 0.2NA objective; Nikon Ti-e). To ensure that phototactic effects did not bias cell motility measurements[23], experiments were performed in a darkened room equipped with black-out shades, a light shield was constructed to block stray light, and the microscope's transmitted light source was equipped with a 665 nm longpass filter (Thorlabs, Inc.), which was outside of the spectral detection range of *C. reinhardtii*[42]. Control experiments in the absence of viscosity gradients (Extended Data Fig. 5) confirmed that no phototactic bias was present. Upon halting the flow and initializing the viscosity gradient formation, videos were recorded (Zyla, Andor Technology) for 1000 frames each at 30 fps, every 5 min for the first 90 min. Dilute cell suspensions ($\approx$ 66 cells per frame) ensured negligible cell-cell interactions[43] and high fidelity cell tracking. Cells were tracked using a custom predictive particle tracking code written in MATLAB (MathWorks). Briefly, cell centroids were identified by intensity thresholding[44,45] and their trajectories were reconstructed via distance minimization of quadratic projections of cell motion in comparison to new candidate cell locations[46]. Three to four replicate experiments (3,454-7,730 cell trajectories each) were performed for each viscosity gradient. Spatial cell density and swimming speed profiles were measured by partitioning the channel into 11 equally spaced bins ($\approx$ 90 μm wide; Fig 1h,i). To quantify the cell turning rate, ω, as a function of orientation (Fig. 2), trajectories were smoothed with a Gaussian filter (1 s standard deviation) to remove high frequency fluctuations, stemming from the natural helical trajectories of *C. reinhardtii*[47]. Occasional stuck cells and mis-trackings were removed by requiring a minimum ratio of track contour length to radius of gyration ≥ 4.

Rotational cell diffusion. Rotational diffusion of *C. reinhardtii* was measured by recording cells (2× 0.1NA objective; Blackfly S, FLIR Systems) for 10 min at 10 fps and 15 fps in bulk 0.996 cP and 4.48 cP media, respectively. Cell tracks with a minimum duration of 15 s

were filtered as described above, and a minimum mean cell swimming speed of 54.2 μm/s and 15.0 μm/s was required for the 0.996 cP and 4.48 cP media, respectively. The MSDs (Extended Data Fig. 6) were fitted (Extended Data Fig. 6, solid lines) using the analytical result[48,49], $\langle \Delta r^2 \rangle = 0.5 V^2 / D_{\mathrm{r}}^2 (2 D_{\mathrm{r}} t + \exp(-2 D_{\mathrm{r}} t) - 1)$. For measured mean swimming speeds $V = 74.7$ μm/s and 22.7 μm/s, the resulting rotational diffusion coefficients were $D_{\mathrm{r}} = 0.070$ s^{-1} and 0.068 s^{-1} for 0.996 cP and 4.48 cP, respectively.

Langevin simulation parameters. Langevin simulations (Fig. 3 and Extended Data Fig. 7-8) were performed by numerically integrating the equations of motion for cell position and orientation using a fourth-order Runge-Kutta method with a time step of 0.001 s (Supplementary Section 5.1). Swimmers are uniformly seeded across the domain with random orientation, including: 500 swimmers for Fig. 3c and Extended Data Fig. 7 and 100,000 for Fig. 3a,b and Extended Data Fig. 8, which were integrated for 1000 s and 6 s, respectively. Based on measured cell motility parameters (Fig. 1-2), viscophobic rotation rate and viscous slow-down were independently varied via ω_{visc} and $\Delta V / V_{\mathrm{max}}$, respectively. Typical, constant values were chosen for all simulations including $V_{\mathrm{max}} = 100$ μm/s and $D_{\mathrm{r}} = 0.069$ s^{-1} (Fig. 3 and Extended Data Figs. 7-8). Boundary interactions were treated with simple specular reflections. 900 total simulations spanning $\Delta V / V_{\mathrm{max}}$ and ω_{visc} were performed for Fig. 3c. For a fixed $\Delta V / V_{\mathrm{max}}$ (corresponding to a $\nabla \eta$), the minimum root mean square (RMS) difference between experimental (Fig. 1i, solid lines) and simulation (Extended Data Fig. 7a) density profiles determined the best match ω_{visc} from simulations (Fig. 3c, squares), which compared well with measured values (Fig. 3c, circles).

Kullback-Leibler divergence. Langevin simulation results were compared to a theoretical prediction for the cell density profile, $\rho_0(x)$, without viscophobic turning (Supplementary Section 4.1). For $\omega_{\mathrm{visc}} = 0$, swimmers are predicted to accumulate in the regions of slow swimming speed (Extended Data Fig. 7b, black dashed curve; $\Delta V / V_{\mathrm{max}} = 0.8$ corresponding to $\nabla \eta = 7.2 \times 10^{-3}$ cP·μm^{-1}). For $\omega_{\mathrm{visc}} > 0$, Langevin simulation

density profiles, $\rho(x)$, departed from theory (Extended Data Fig. 7b) by shifting to high swimming speed zones. The effect of viscophobic turning on the density profile departure was captured by the Kullback–Leibler (KL) divergence[35], or relative entropy,

$$D_{\mathrm{KL}}(\rho \| \rho_0) = \int_{x=0}^{W} \rho(x) \log \left(\frac{\rho(x)}{\rho_0(x)} \right) dx,$$

where $D_{\mathrm{KL}} \geq 0$, and $D_{\mathrm{KL}} = 0$ indicates two identical distributions. D_{KL} was computed for all 900 Langevin simulations investigated (Fig. 3c, colormap).

Reporting Summary. Further information on research design is available in the Nature Research Reporting Summary linked to this Article.

Data availability. The data generated during the current study are available from the corresponding author upon reasonable request.

Code availability. The algorithms and simulation codes are described in the Methods and Supplementary Information.

Acknowledgements. We thank G.J. Elfring for helpful discussions and R.J. Henshaw for comments on the manuscript. This work was funded by NSF Awards CAREER-1554095 and CBET-1701392.

Author contributions. M.R.S, N.W., and J.S.G. designed research; M.R.S. and N.W. performed experiments; M.R.S., N.W., and D.M.W. analyzed experimental data; N.W., D.M.W., M.R.S, and J.S.G. contributed to simulations and theoretical analyses; M.R.S., N.W., D.M.W., and J.S.G. wrote the paper.

Competing interests. The authors declare no competing interests.

Additional information. Correspondence and requests for materials should be addressed to Jeffrey.Guasto@tufts.edu.

---

[36] Sager, R. & Granick, S. Nutritional studies with *Chlamydomonas reinhardi*. *Ann. N. Y. Acad. Sci.* **56**, 831–838 (1953).

[37] Ebagninin, K. W., Benchabane, A. & Bekkour, K. Rheological characterization of poly(ethylene oxide) solutions of different molecular weights. *J. Colloid Interface Sci.* **336**, 360–367 (2009).

[38] Grigorescu, G. & Kulicke, W.-M. *Prediction of Viscoelastic Properties and Shear Stability of Polymers in Solution*, 1–40 (Springer, Berlin, Heidelberg, 2000).

[39] Xia, Y. & Whitesides, G. M. SOFT LITHOGRAPHY. *Annu. Rev. Mater. Sci.* **28**, 153–184 (1998).

[40] Devanand, K. & Selser, J. C. Polyethylene oxide does not necessarily aggregate in water. *Nature* **343**, 739–741 (1990).

[41] Waigh, T. A. Microrheology of complex fluids. *Reports Prog. Phys.* **68**, 685–742 (2005).

[42] Kreis, C. T., Le Blay, M., Linne, C., Makowski, M. M. & Bäumchen, O. Adhesion of *Chlamydomonas* microalgae to surfaces is switchable by light. *Nat. Phys.* **14**, 45–49 (2018).

[43] Rafaï, S., Jibuti, L. & Peyla, P. Effective Viscosity of Microswimmer Suspensions. *Phys. Rev. Lett.* **104**, 098102 (2010).

[44] Cheezum, M. K., Walker, W. F. & Guilford, W. H. Quantitative Comparison of Algorithms for Tracking Single

Fluorescent Particles. *Biophys. J.* **81**, 2378–2388 (2001).

Crocker, J. C. & Grier, D. G. Methods of Digital Video Microscopy for Colloidal Studies. *J. Colloid Interface Sci.* **179**, 298–310 (1996).

Ouellette, N. T., Xu, H. & Bodenschatz, E. A quantitative study of three-dimensional Lagrangian particle tracking algorithms. *Exp. Fluids* **40**, 301–313 (2006).

Crenshaw, H. C. A New Look at Locomotion in Microorganisms: Rotating and Translating. *Am. Zool.* **36**, 608–618 (1996).

Howse, J. R. *et al.* Self-Motile Colloidal Particles: From Directed Propulsion to Random Walk. *Phys. Rev. Lett.* **99**, 48102 (2007).

Rusconi, R., Guasto, J. S. & Stocker, R. Bacterial transport suppressed by fluid shear. *Nat. Phys.* **10**, 212–217 (2014).

Korson, L., Drost-Hansen, W. & Millero, F. J. Viscosity of water at various temperatures. *J. Phys. Chem.* **73**, 34–39 (1969).

Extended Data Fig. 1: **Microfluidic viscosity gradient generation.** Schematic of the junction and test section of a microfluidic viscosity gradient generation device, where inlets containing a viscous solution of PEO polymer in M1 media ($\mathcal{S}_1$; left), *C. reinhardtii* cell suspension in M1 ($\mathcal{S}_2$; center), and M1 media ($\mathcal{S}_3$; right) meet. The flow rates Q_1, Q_2, and Q_3 are adjusted for solutions having viscosities η_1, η_2, and η_3 to maintain the initial stratification of these streams in the main channel of $w_1 = w_3 = 4W/9$ and $w_2 = W/9$. Background is qualitative and illustrates initial stratification of viscous fluid (blue), cell suspension, and media.

Extended Data Fig. 2: **Mean square displacements of tracer particles for viscosity measurements in bulk fluid.** Microrheology quantifies the viscosity of PEO solutions in M1 media at various concentrations. The slopes of linear fits (solid lines) to the mean square displacements (MSDs; markers) yield the tracer particle diffusion coefficients, which are used to determine the viscosity of the solution via the Stokes-Einstein relation (Supplementary Section 2.1). Measured viscosities are listed in Extended Data Table 1.

Extended Data Fig. 3: **Spatially resolved microrheology quantifies the viscosity profile in a microfluidic device.** **a**, The mean square displacement (MSD) of tracer particles in different locations across the microfluidic viscosity gradient device at time, t = 10 min ($\nabla\eta = 3.4 \times 10^{-3}$ cP·μm^{-1}; see also Extended Data Fig. 4). Solid lines are linear fits to the MSDs. **b**, The slopes of the tracer particle MSDs in **a** are used to determine the spatially resolved viscosity profile, $\eta(x)$, across 11 bins of the microchannel width using Stokes-Einstein relation (Supplementary Section 2.2). Shaded region (smaller than the markers) represents the propagation of uncertainty from the error of the fit in **a**.

Extended Data Fig. 4: **Time evolution of the viscosity profile.** Transient diffusion of the PEO molecules results in a viscosity profile that evolves from a near step function to a smooth monotonic profile. Swimming cell assays are analyzed in the time range $30 \leq t \leq 90$ min, when the gradient evolves more slowly and the profile is approximately linear in the center of the channel. The slope of the viscosity profile is quantified by fitting a line to the five central points in the channel. The reported viscosity gradient, $\nabla \eta$, is the average slope of the viscosity profiles from 30-90 min (Extended Data Table 2).

Extended Data Fig. 5: **Control experiments show negligible chemotaxis and chemokinetic effects for *C. reinhardtii* in PEO gradients. a,b**, Viscotaxis assays reveal the local, measured cell density (**a**) and mean swimming speed (**b**) remain spatially uniform for small viscosity gradients. Base PEO concentrations of 0.05% in $\mathcal{S}_1$ ($\nabla\eta = 0.46 \times 10^{-3}$ cP $\cdot\mu$m^{-1}; N=3; right) are comparable to the control experiment with no viscosity gradient ($\nabla\eta = 0$ cP $\cdot\mu$m^{-1}; N=4; left, repeated from Fig. 1h,i). For these control experiments, the base PEO concentration (0.05%) is nearly the same order of magnitude as the maximum gradient (0.82%), yet has no appreciable effect on the local viscosity. Thus, the lack of discernible cell accumulation and the lack of statistically significant spatial variations in swimming speed indicate that PEO likely does not act as a chemoattractant nor does it have any chemokinetic effect on *C. reinhardtii*. The shaded areas represent the standard error.

Extended Data Fig. 6: **Measurement of mean square displacement (MSD) of *C. reinhardtii* in uniform viscosity to determine rotational diffusion coefficient.** Mean square displacements of *C. reinhardtii* were measured in 0.996 cP ($\approx$ 1,800 trajectories) and 4.48 cP ($\approx$ 10,700 trajectories) bulk viscosities. The resulting MSDs were fitted using an analytical result for a persistent random walk[48,49], $\langle \Delta r^2 \rangle = 0.5 V^2/D_\mathrm{r}^2 (2D_\mathrm{r}t + \exp(-2D_\mathrm{r}t) - 1)$, where the ensemble averaged cell swimming speeds, V, are known independently (74.7 μm/s and 22.7 μm/s, respectively), and the rotational diffusion coefficient, D_r, is the lone fitting parameter. The resulting rotational diffusion coefficients were D_r = 0.070 s^{-1} and 0.068 s^{-1}. As a sensitivity analysis, varying the fit window from 10 s to 20 s resulted in D_r = 0.071 $\pm$ 0.006 s^{-1} and 0.068 $\pm$ 0.001 s^{-1} for 0.996 cP and 4.48 cP, respectively, where the error is the standard deviation of the resulting rotational diffusion coefficients.

Extended Data Fig. 7: **Cell density distribution comparison of Langevin simulations and experiments. a**, Cell density profiles, $\rho(x)$, from experiments (repeated from Fig. 1i) compare favorably to Langevin simulations of cell motility with corresponding spatially varying swimming speed (parameterized by $\Delta V/V_{\max}$) and best match turning rate amplitude, ω_{visc}, based on the minimum root-mean-square (RMS) difference between the experiments and simulations (Fig. 3c, squares). **b**, Langevin simulations enable independent exploration of the parameter $\Delta V/V_{\max}$ and ω_{visc}. Cell densities profiles, $\rho(x)$, are shown for fixed $\Delta V/V_{\max}$ and a range of turning rate amplitudes (solid lines). The resulting density profiles are compared to a theoretical distribution, ρ_0, for cells with spatially varying swimming speed in the absence of viscophobic turning (black dashed curve; Supplementary Section 4.1).

Extended Data Fig. 8: **Time evolution of swimming cell orientation distribution in a viscosity gradient. a-c**, In the absence of a viscosity gradient or viscophobic turning, an initially random uniform distribution of cell swimming orientations remains random, as illustrated across experiments, Langevin simulations, and a Fokker-Plank model, respectively. **d-f**, In contrast, in the presence of a viscosity gradient or orientation dependent viscophobic turning rate, the orientation of cells readily condenses in the down-gradient direction. Experimentally measured cell swimming trajectories were sub-sampled from the central portion (400 μm $\leq x \leq$ 600 μm) of the microfluidic gradient generating channel with a uniform orientation distribution $p(\theta|t)$ at $t = 0$ for $\nabla\eta = 0$ cP·μm^{-1} (**a**; control, $\approx$ 1,170 trajectories) and for $\nabla\eta = 7.2 \times 10^{-3}$ cP·μm^{-1} (**d**; $\approx$ 1,330 trajectories). Peaks in **a** at $\approx$ 6 s are due to cell collisions with microchannel walls. Langevin simulations in the absence (**b**) and presence (**e**) of viscophobic turning ($\omega_{\text{visc}} = 0.07$ s^{-1}, $\Delta V/V_{\text{max}} = 0.776$) show quantitative agreement with experiments having matching conditions (Extended Data Fig. 7 and Fig. 3a,b) in **a** and **d**, respectively. A one dimensional Fokker-Planck model of the cell swimming orientation distribution (Supplementary Section 5.3) likewise captures the time evolution of the conditional probability density for the control (**c**) and corresponding maximum viscosity gradient conditions (**f**; Fig. 3b).

PEO Concentration (w/v)	Viscosity (cP) [Ebagninin et al.[37]]	Viscosity (cP) [current study]
0.0%	1*	0.996 ± 0.0052
0.05%	1.09†	-
0.1%	1.36†	1.38 ± 0.0071
0.125%	1.51†	-
0.25%	2	2.23 ± 0.0121
0.375%	3.44†	-
0.5%	5	4.48 ± 0.0195
0.82%	9.01†	8.96 ± 0.0264
1.0%	12	-

Extended Data Table 1: **Viscosity of bulk PEO solutions.** The dynamic viscosities of aqueous PEO solutions of various concentrations reported in the literature[37] compared to the present study in M1 media (measured by microrheology; $\approx 22°$C). The * indicates the known viscosity value at $20°$C[50] and the † indicates a quadratic interpolation of Ebagninin et al.[37]. Experimental uncertainty represents the propagation of error from the calculation of the diffusion coefficient.

PEO concentration in inlet $\mathcal{S}_1$ (w/v)	Measured viscosity gradient, $\nabla\eta$ (cP $\cdot\mu$m^{-1})
0.0%	0
0.05%	0.46×10^{-3} †
0.1%	$0.79\times10^{-3} \pm 0.75 \times10^{-4}$
0.25%	$2.2\times10^{-3} \pm 3.41 \times10^{-4}$
0.5%	$3.4\times10^{-3} \pm 9.61 \times10^{-4}$
0.82%	$7.2\times10^{-3} \pm 1.9 \times10^{-3}$

Extended Data Table 2: **Measured viscosity gradients for viscotaxis experiments.** Viscosity gradients, $\nabla\eta$, for each of the tested gradient conditions were generated by controlling the bulk PEO concentration/viscosity in the inlet, $\mathcal{S}_1$ (Extended Data Fig. 1). Viscosity gradients were measured by fitting a line to the measured viscosity profiles in the central third of the channel. The slopes of the fits were then averaged to determine the mean $\nabla\eta$ over the experiment time window of 30-90 min. The † indicates interpolation by quadratic fit. Uncertainty represents the standard deviation of the fitted viscosity profile slopes from 30-90 min (Extended Data Fig. 4).

Supplementary Information:
Viscophobic turning regulates microalgae transport in viscosity gradients

Michael R. Stehnach,[1,][*] Nicolas Waisbord,[1,][*] Derek M. Walkama,[1,2] and Jeffrey S. Guasto[1,][†]

[1]*Department of Mechanical Engineering, Tufts University,*
200 College Avenue, Medford, Massachusetts 02155, USA
[2]*Department of Physics and Astronomy, Tufts University,*
574 Boston Avenue, Medford, Massachusetts 02155, USA

CONTENTS

1. MICROFLUIDIC VISCOSITY GRADIENT ASSAY

Monotonic viscosity profiles of various gradient magnitude were generated by prescribing the viscosity of the solutions in each of the three inlet channels (Fig. 1b and Extended Data Fig. 1): (*i*) The leftmost inlet contained a viscous polyethylene oxide (PEO) solution (molecular weight (MW), 400,000 g/mol; Sigma-Aldrich) based in minimal (M1) media (solution, $\mathcal{S}_1$; Extended Data Table 1) [1]. (*ii*) The center channel contained a dilute *C. reinhardtii* suspension in M1 media (solution, $\mathcal{S}_2$). (*iii*) The rightmost channel contained M1 media (solution, $\mathcal{S}_3$). In the case of viscosity profile measurements, the *C. reinhardtii* suspension in $\mathcal{S}_2$ was replaced by M1 media and all three solutions were seeded with a dilute suspension of tracer particles (see §2). To initiate the viscosity gradient, two syringe pumps (one for $\mathcal{S}_1$ and one for $\mathcal{S}_2$ and $\mathcal{S}_3$; Harvard Apparatus) were flowed at a high rate of speed for 2 min to establish stratification between the three viscous solutions in the main test channel (Extended Data Fig. 1). At the start of the assay,

---

[*] These authors contributed equally
[†] Corresponding author

both syringe pumps were abruptly stopped, and videos of cell or tracer particle motion were captured periodically over the course of the experiment. Upon halting the flow, the initially step-wise concentration profile (and consequently viscosity profile, $\eta(x)$) of PEO slowly evolves over the course of 90 min into a smooth concentration/viscosity profile (Extended Data Fig. 4) due to the small diffusivity ($D \approx 2 \times 10^{-7}$ cm^2/s) of the relatively large PEO molecules [2].

We chose the target widths of the initial fluid bands for $\mathcal{S}_1$, $\mathcal{S}_2$, and $\mathcal{S}_3$ in the test section to be $w_1 = 4W/9$, $w_2 = W/9$, and $w_3 = 4W/9$, respectively (Extended Data Fig. 1). The thin central band of *C. reinhardtii* in M1 ($\mathcal{S}_2$) ensures that no bias is introduced in the assay due to the starting position of the cells. Furthermore, bracketing the cell suspension by solutions based on fresh M1 media (i.e. $\mathcal{S}_1$ and $\mathcal{S}_3$) ensured that there were no extraneous chemical gradients that might bias motility, for example due to the consumption of resources by the cells [3]. Owing to the differential viscosities, η_i, of the PEO and M1 solutions, initial flow rates, Q_i, were adjusted to maintain the symmetry of the desired band widths, w_i, established above for the main channel for $\mathcal{S}_1$, $\mathcal{S}_2$, and $\mathcal{S}_3$. From the predicted hydraulic flow resistance of each solution in the main channel [4], the desired band widths are related to the respective flow rates and viscosities of the solutions approximately by $w_i \eta_i / Q_i$ = constant. Due to the dilute nature of the cell suspensions, we assume that the viscosities of $\mathcal{S}_2$ and $\mathcal{S}_3$ are the same, $\eta_2 = \eta_3$, as the M1 media alone [5]. Additionally for viscosity profile measurements, we assume that the addition of dilute tracer particles (see §2.1) does not augment the viscosities of the respective solutions. Solutions $\mathcal{S}_2$ and $\mathcal{S}_3$ were flowed through the device using glass syringes (Hamilton) with various barrel radii, r_i ($r_3 > r_2$), and controlled by a single syringe pump resulting in relative flow rates $Q_3 = 17$ μL/min and $Q_2 = Q_3 (r_2/r_3)^2 = 4.25$ μL/min. This ensured achieving the prescribed flow stream widths. The viscous PEO solution, $\mathcal{S}_1$, was injected using a second, separate syringe pump and syringe with an initial flow rate $Q_1 = Q_3 \eta_3 / \eta_1$. Phase contrast microscopy enabled visualization of the stratified fluid interfaces, and the flow rate Q_1 was fine tuned to ensure that the initial, desired symmetry of the stratified solutions ($w_1 = w_3$) was maintained for all experiments. The final adjusted flow rates were $Q_1 \approx 17$, 14, 11, 7, 5 and 3.5 μL/min for PEO solutions with concentrations 0%, 0.05%, 0.1%, 0.25%, 0.5%, and 0.82% (w/v), respectively (Extended Data Table 1). Finally, during control experiments, a negligible change in swimming speed (Extended Data Fig. 5b, left) compared to the bulk measurements (Fig. 1f) verified that this assay had no negative effects on the motility of the robust *C. reinhardtii* cells.

2. MICROSCALE VISCOSITY MEASUREMENTS

Directly quantifying spatially varying viscosity on sub-millimeter scales was necessary to understand the viscotactic transport of swimming cells in viscosity gradients. To measure the viscosity of both the spatially varying viscotaxis experiments and bulk PEO solutions, we adapted techniques from well-developed microrheology approaches [6, 7].

2.1. Bulk media

For uniform, bulk fluids, we measure the Brownian thermal motion of micron-sized tracer particles suspended in viscous fluids of interest [8], which we then relate to the viscosity, η. For simple, Newtonian fluids, the viscosity is quantified by measuring the mean square displacements (MSD), $\langle \Delta x^2 \rangle$, of microparticles over time, Δt, through particle tracking. We focus on particle displacements, Δx, perpendicular to the direction of the flow in the microfluidic devices to circumvent any minor tracer particle drift due to residual flows in the channels during the assay. The diffusion coefficient, D, of the tracer particles is first measured from a linear fit to the MSD over Δt:

$$\langle \Delta x^2 \rangle = 2D\Delta t \tag{S1}$$

Applying the Stokes-Einstein relation, the viscosity, η, is inferred from the measured diffusion coefficient (equation (S1)) [6]:

$$D = \frac{k_{\mathrm{B}} T}{6\pi \eta R} \tag{S2}$$

where k_{B} is the Boltzmann constant, $T = 295$ K is the absolute ambient temperature, and R is the radius of the tracer particles.

To quantify the media viscosity, epifluorescence microscopy (Nikon Ti-e) captured the diffusive motion of fluorescent tracer particles. Viscous media – M1 media with PEO concentrations of 0.1%, 0.25%, 0.5%, and 0.82% (w/v) – was seeded with 0.5 μL/mL of fluorescent tracer particles ($R = 0.25$ μm radius; 2% solid; carboxylated FluoroSpheres, Life Technologies) and placed in a microfluidic observation chamber, where the inlet and outlet were sealed with wax. Using a 10× magnification objective (0.5NA) with an additional 1.5× magnification multiplier lens, videos of

the particles were recorded (Blackfly S, FLIR Systems) at a frame rate of 4 fps for 25 s. Particle tracking (see Methods) was used to measure the $\langle \Delta x^2 \rangle$ of 400-1,000 tracer particles. With increasing concentrations of PEO, the diffusion coefficient (equation (S1)) decreases (Extended Data Fig. 2), revealing an increase in bulk viscosity. The experimentally measured bulk viscosities are reported in Extended Data Table 1, and compare well with previously reported results [9]. Furthermore, PEO suspensions are known to remain Newtonian under these concentration conditions ($c < 1\%$ w/v), which are well below the transition to semi-dilute solutions, $c^* \approx 3\%$ (w/v) [9].

2.2. Viscosity profile

Using the same microfluidic device and conditions as described in §1 and in the main text for the cell motility assays (Fig. 1b), the spatially varying viscosity profile in the channel test section was quantified, in the absence of *C. reinhardtii*, by seeding the media with the fluorescent tracer particles. Following the viscosity gradient assay procedure along with the bulk viscosity measurement procedure above, epifluorescence microscopy captured the diffusive motion of the tracer particles within the viscosity gradient. Viscosity profiles were quantified using a 10× magnification objective (0.2NA) with an additional 1.5× magnification multiplier lens and 10 videos were recorded at 4 fps for 250 frames (63 s long) every 10 minutes for 90 min (Zyla, Andor Technology). For the control experiments, in which solution $\mathcal{S}_1$ contained no PEO, videos were recorded with the same parameters as the bulk media at 4 fps for 100 frames every 3.5 minutes for 10 minutes (Blackfly S, FLIR Systems). By spatially resolving the seeded tracer particles using particle tracking, the channel was partitioned into 11 equally spaced bins, where the $\langle \Delta x^2 \rangle$ was calculated over time, Δt, for each bin (Extended Data Fig. 3). Following this method over the course of the experiment, the viscosity profile's evolution over time was realized (Extended Data Fig. 4). After about 30 min, the strong transient variations in the viscosity profile decrease. Thus, we report and analyze our results 30 to 90 minutes after ceasing the initial stratifying flow, when the viscosity profile varies weakly in time. While the slope of the viscosity profile changes slightly from 30 min to 90 min, the motility of *C. reinhardtii* enables the cells to rapidly sample the channel (effective diffusion coefficient, $D \approx 0.68 \times 10^{-3}$ cm^2/s [10]), resulting in quasi-steady cell transport. This substantial difference in time evolution of the PEO concentration and cell concentration justifies the use of MW 400,000 g/mol PEO molecules for the generation of the viscosity profile. The average viscosity profile, over the 30-90 min, was calculated by averaging the local measured viscosity for each bin over time (Fig. 1g).

From each measured viscosity profile, $\eta(x)$, the viscosity gradient, $\nabla \eta$, was calculated from a linear fit to the middle third of the viscosity profile in the center of the microfluidic channel. The viscosity gradient reported for each experiment throughout the main text is the average gradient within the 30-90 min interval (Extended Data Table 2).

3. CONTROL EXPERIMENTS

3.1. Bulk viscosity swimming speed

C. reinhardtii is a biflagellate (Fig. 1a) with a slightly prolate body (typical diameter of 10 μm) [11]. These cells have two 10-12 μm long anterior flagella (≈ 0.2-0.3 μm diameter) [12, 13], and swim with a breast stroke pattern (≈ 60 Hz beat frequency) [11]. In order to determine the effects of spatially inhomogeneous viscous media on the transport properties of *C. reinhardtii*, we first examined the effect of enhanced bulk viscosity on the motility of these cells by quantifying their ensemble average swimming speed, $\langle V \rangle$. The cells were gently resuspended in PEO solutions of various concentration (and thus viscosities) prepared in M1 media (Extended Data Table 1). To minimize dilution of the viscous media in the resuspension process, the cell suspension was washed three times: (i) 1 mL of cell suspension was pipetted into a 1.5 mL microcentrifuge tube and centrifuged (Eppendorf) at 2,800 rcf for 30 seconds to form a soft pellet, (ii) the supernatant (≈ 0.9 mL) was removed and replaced with an equal amount of viscous media, and (iii) the entire process was repeated two more times. The cells were pipetted into a PDMS microfluidic observation chamber and the inlet and outlet were sealed with wax to eliminate residual flow. Care was taken to ensure that the cells were not damaged during the pipetting and resuspension steps, and the robustness of motility was confirmed both visually and through quantitative video microscopy analysis (Fig. 1f).

Using video microscopy and particle tracking, the ensemble mean swimming speed, $\langle V \rangle$, of the cell trajectories over multiple videos were measured. For each bulk viscosity, 2-3 videos were recorded using the same specifications as for the viscotaxis experiments (see Methods). For bulk viscosities of $\eta = 0.996$ cP, 1.51 cP, 2.23 cP, 3.44 cP, 4.48 cP, and 8.96 cP (Extended Data Table 1), the ensemble averaged swimming speeds of *C. reinhardtii* were $\langle V \rangle = $ 92.2 μm/s, 85.4 μm/s, 51.2 μm/s, 30.3 μm/s, 21.8 μm/s, and 14.5 μm/s, respectively (Fig. 1f). We observed a power law decrease of swimming speed with increasing viscosity of slope −0.934 (Fig. 1f, black line), which is consistent with prior observations [14] and constant thrust flagellar propulsion, $\langle V \rangle \sim \eta^{-1}$. Error bars in Fig. 1f represent the mean

standard deviation across $N = 3$ videos (or the range in the case of $N = 2$ for 0.996 cP and 8.96 cP) for each bulk viscosity (each with $\sim 2,700$ trajectories).

3.2. Potential effects of chemotaxis and chemokinesis

Many microorganisms are capable of sensing, responding to, and metabolizing small polymers in suspension, which may introduce biases into experimental measurements [3, 15, 16]. Chemotaxis can cause cell accumulation in a chemical gradient [17] and chemokinesis can enhance cell swimming speeds in elevated chemical concentrations [18]. Through control experiments, performed at low concentrations of PEO in M1 media, we demonstrate that neither chemotaxis nor chemokinesis affect the motility of $C.\ reinhardtii$ in our system. The base PEO concentration (0.05% for solution $\mathcal{S}_1$; Extended Data Fig. 1) was selected such that the viscosity was within 10% of the M1 media (corresponding to $\nabla\eta = 0.46 \times 10^{-3}$ cP·μm^{-1}), while differing from the maximum concentration by approximately one order of magnitude. In repeating these assays with this smaller viscosity gradient, no cell accumulation was discernible in the cell density profile (Extended Data Fig. 5a, right) compared to the no gradient control experiment ($\nabla\eta = 0$ cP·μm^{-1}; Extended Data Fig. 5a, left). This observation indicates that chemotaxis effects are likely negligible in this system. Furthermore, no spatial variations in swimming speed were discernible for the lower viscosity gradient ($\nabla\eta = 0.46 \times 10^{-3}$ cP·μm^{-1}; Extended Data Fig. 5b, right) compared to the control. We note that a lower mean swimming speed was observed for $\nabla\eta = 0.46 \times 10^{-3}$ cP·μm^{-1}, but this is consistent with the slightly elevated viscosity, the natural variations in population swimming speed, and the experimental uncertainty. Thus, we likewise conclude that there are no discernible chemokinesis effects by PEO on $C.\ reinhardtii$ that affect the outcomes of our study.

3.3. Potential effects of transient viscosity gradients on viscotaxis measurements

Viscosity gradients were generated in the microfluidic device by the diffusion of PEO polymer (see §1), which is an inherently unsteady process. This effect is apparent in viscosity profile measurements (Extended Data Fig. 4), and its effect on viscotaxis experiments was minimized by restricting our analysis to the 30-90 min window, post gradient formation (i.e. after the strongest transient regime subsides). As a test of the sensitivity of our results, we examined the variation of the local cell density, ρ, and swimming speed, V, profiles between 30-60 min and 60-90 min for the maximum viscosity gradient tested ($\nabla\eta = 7.2 \times 10^{-3}$ cP·μm^{-1}; Fig. S1). For the cell density and swimming speed, we observe a mean percent difference across the 30-60 min and 60-90 min time intervals (relative to the 30-90 min interval) of less than 2.33% and 6.08%, respectively. This variation is well within acceptable tolerances and consistent with the uncertainty in our experiments.

3.4. Potential effects of diffusiophoresis

Concentration gradients of small dissolved molecules can cause a drift of suspended colloidal particles due to the Fickian solute flux, a process known as diffusiophoresis [19]. Through the control experiments below, we definitively confirm that diffusiophoretic effects, due to the established concentration gradients of PEO, have a negligible effect on the transport of $C.\ reinhardtii$ in our viscotaxis experiments. In order to isolate the potential effects of diffusiophoresis, $C.\ reinhardtii$ suspensions were rendered immotile while preserving the cell shape by fixing the cells in a 4% (v/v) solution of 37% formaldehyde (Sigma-Aldrich). Diffusiophoretic drift is known to scale with the solute concentration gradient. Therefore, control viscotaxis experiments were performed with fixed $C.\ reinhardtii$ cells at the maximum tested viscosity (i.e. concentration) gradient of $\nabla\eta = 7.2 \times 10^{-3}$ cP ·μm^{-1} under identical conditions and for the same 90 min duration as the experiments shown in the main text (Fig. S2a). The maximum displacement of the tracked non-motile cells in the viscosity (concentration) gradient direction was just $\Delta x \approx 10\ \mu$m (≈ 1 cell diameter or $\approx 1\%$ of the channel width; Fig. S2b) during the 90 min experiment. The observed displacements are consistent with Brownian diffusion and are many orders of magnitude smaller than the effect of cell motility in the same time frame. This result illustrates unequivocally that diffusiophoresis does not affect the observed viscotactic/viscophobic transport of $C.\ reinhardtii$ described in the main text (Fig. 1).

4. THEORETICAL CONSIDERATIONS

4.1. Cell density based on measured bulk swimming speed

When cells are free from directional bias (e.g. viscophobic turning), local cell swimming speed, $V(x)$ is widely recognized to be inversely proportional to cell density [20], $\rho_0(x)$, and in a one dimensional system:

$$\rho_0(x) = \frac{\rho_0(x_0)V(x_0)}{V(x)},\qquad(S3)$$

where x_0 is an arbitrary reference location in the domain $0 \leq x \leq W$. In the experiments shown in this paper, and generally in situations where speed $V(x)$ is heterogeneous, we consider these predicted distributions $\rho_0(x)$ for unbiased random walks as the reference distributions because any deviation signals a bias in the motility. Here, we combine independent experimental measurements of cell swimming speed in bulk viscous media, locally measured viscosity profiles, and theory to predict these reference distributions representing the expected cell density profiles in our system if there was no viscophobic turning. The resulting cell density, speed, and flux profiles are presented in the main text (Fig. 1h-j, dashed curves). We show that the swimming speed of *C. reinhardtii* in bulk viscous media decays as a power law (Fig. 1f) with increasing viscosity, η:

$$V(\eta) = a\eta^b,$$

where a least squares fit gives $b = -0.934$ and $a = 103$, and V and η have units of $\mu m/s$ and cP, respectively. Combining this result with the known spatial variation of the viscosity, $\eta(x)$, in our microfluidic gradient generating device – measured independently from microrheology experiments (Fig. 1g and Extended Data Fig. 3-4) – we obtain an expression for the swimming speed profile of the cells in a quasi-steady viscosity gradient:

$$V_0(x) = a\eta(x)^b.\qquad(S4)$$

This empirical prediction (Fig. 1h, dashed lines) agrees well with the local mean swimming speed profile of *C. reinhardtii* (Fig. 1h, solid lines), measured *in situ* for the viscotaxis experiments, and gives a high degree of confidence in the reliability of the predicted density profile below.

Next, we recognize that locally, the product of the local density and speed (equation (S3)) should be constant throughout the channel under these conditions: $\rho_0(x)V_0(x) = k$. Combining equations (S4) and (S3) with this flux condition, we obtain the following expression for the predicted cell density profile:

$$\rho_0(x) = \frac{k}{a}\eta(x)^{-b}.$$

Because ρ_0 is a probability density function, the normalization constant is defined by $k = a/\int_{x=0}^{W}\eta(x)^{-b}dx$, without any additional fitting parameters, and the expression for the predicted density profile (Fig. 1i, dashed lines) is

$$\rho_0(x) = \frac{\eta(x)^{-b}}{\int_{x=0}^{W}\eta(x)^{-b}dx}.$$

Following from the cell density, the predicted flux magnitude (i.e. a constant value; Fig. 1j, dashed lines) in the viscosity gradient is

$$\rho_0(x)V_0(x) = \frac{a}{\int_{x=0}^{W}\eta(x)^{-b}dx}.$$

As described in the main text, ρ_0 over-predicts the slope of the measured cell density profile, and the measured cell flux progressively deviates from the predicted constant value with increasing viscosity gradient. The strong agreement of the measured, $V(x)$, and predicted, $V_0(x)$, velocity profiles combined with the deviations of the cell density and flux are indicative of a new phenomenon that depends on the magnitude of the viscosity gradient, namely the viscophobic reorientation described in the main text.

4.2. Three-point force swimmer model

In the low Reynolds number regime, biflagellated swimming microalgae, such as *C. reinhardtii*, are commonly modeled using well-established three point force models [21, 22]. In particular, the thrust from the dual flagella are

approximated by one point force each, while the drag from the body is approximated by a third point force or the Stokes drag on a spheroidal body. Here, we extend this idea for a biflagellated microswimmer in a viscosity gradient to corroborate our experimental observations (Fig. 2-3). The model considers a three point force model swimmer in a two-dimensional domain having an inhomogeneous viscosity landscape, $\eta(\mathbf{r})$. In the analysis that follows, we approximate the drag and thrust forces as steady, which is equivalent to averaging over the flagellar beat cycle. This approximation is justified because the beat frequency of the flagella (~ 50 Hz) is at least two orders of magnitude faster than other relevant time scales, for example including the observed viscophobic turning rate, the effective rotational diffusion, and the travel time across the gradient. Furthermore, we assume the swimming dynamics do not affect the viscosity gradient (i.e. through mixing), which is consistent with other recent models [23–25]. In the analysis developed below, we illustrate that the two primary effects regulating the trajectory of a three-point force model swimmer in a viscosity gradient are (i) the orientation-dependent propulsive force imbalance between the two flagella and (ii) the coupling of rotational and translational motion of the cell body [23].

4.2.1. Asymmetric flagellar propulsive force

For drag-based thrust in the Stokes flow regime, the effective propulsive force of each flagellum is proportional to the local viscosity, as well as the velocity and conformation of the flagellum. Thus, the force generated by a flagellum, i, is described as

$$\mathbf{F}_i = \mathbf{k}_i \eta(\mathbf{r}_i),$$

where $\mathbf{r}_i$ is the location of the center of thrust. The geometry and kinematics of the flagellum are encompassed in $\mathbf{k}_i$, which in the most general case is not necessarily aligned with the body axis. The center of thrust for a given flagellum can be expressed as

$$\mathbf{r}_i = \mathbf{r} + \mathbf{d}_i = \mathbf{r} + \left(d_{\parallel,i} \hat{\mathbf{p}} + d_{\perp,i} \hat{\mathbf{n}} \right),$$

where $\mathbf{d}_i$ is the relative location from the body center (i.e. center of drag), $\mathbf{r}$, and the cell body axis and its normal are $\hat{\mathbf{p}}$ and $\hat{\mathbf{n}}$, respectively. For linear viscosity gradients – or small separations of the center of thrust from the body center, compared to the viscosity gradient – the flagellar thrust is

$$\mathbf{F}_i = \mathbf{k}_i \left(\eta(\mathbf{r}) + \nabla\eta(\mathbf{r}) \cdot \mathbf{d}_i \right).$$

To focus on the effect of the viscosity gradient on the viscophobic turning behavior of the cells, we make the following simplifying assumptions in line with other recent simulations in viscosity gradients [25]: (i) The thrusting of the two flagella is symmetric and independent of the local viscosity, $\mathbf{k} = \mathbf{k}_i$. ($ii$) The resultant thrusting is in the direction of the cell body, $\mathbf{k} = k\hat{\mathbf{p}}$. ($iii$) The centers of thrust of the two flagella are located lateral to the center of drag of the swimmer's body, where $\mathbf{d}_i = \pm d\hat{\mathbf{n}}$ and $i = 1, 2$. The flagellar thrust forces become

$$\mathbf{F}_i = k \left(\eta \pm \nabla\eta \cdot d\hat{\mathbf{n}} \right) \hat{\mathbf{p}}, \tag{S5}$$

which also generate a torque about the center of the cell body

$$\boldsymbol{\Gamma}_i = \mathbf{d}_i \times \mathbf{F}_i = \pm k d^2 \left(\frac{\eta}{d} \hat{\mathbf{z}} \pm \nabla\eta \times \hat{\mathbf{p}} \right). \tag{S6}$$

Thus, the forces and torques generated by the two flagella are not in general equal in the presence of a viscosity gradient.

4.2.2. Rotational and translational coupling of swimmer body dynamics

The bodies of unicellular swimming microalgae – including *C. reinhardtii* used in the current work – are spherical to slightly prolate spheroids. Here, we model the dynamics of the swimmer body as a sphere of radius, R. Even for such highly symmetric bodies, a unique feature of their kinematics in a low Reynolds number viscosity gradient is the coupling of rotational and translational motion [23], which we quote below. The translational and rotational Stokes drag for a spherical body in a viscosity gradient are

$$\mathbf{F}_{\mathrm{b}} = -6\pi\eta R\mathbf{V} + 2\pi R^3 \nabla\eta \times \boldsymbol{\omega}, \tag{S7}$$

$$\boldsymbol{\Gamma}_{\mathrm{b}} = -8\pi\eta R^3 \boldsymbol{\omega} - 2\pi R^3 \nabla\eta \times \mathbf{V}, \tag{S8}$$

where $\mathbf{V}$ is the velocity (i.e. cell swimming speed), and $\boldsymbol{\omega}$ is the angular rotation rate. Physically, if a sphere were to be dragged through a viscosity gradient at a velocity $\mathbf{V}$ in the direction perpendicular to the gradient, it would rotate at a speed $\boldsymbol{\omega}$, but not if the force is applied in the viscosity gradient direction. This result is due to the differential viscous traction forces on either side of the body when moving perpendicular to the gradient, which ultimately cause it to rotate. In a similar fashion, if a sphere is rotated by an external torque, it will translate in the direction perpendicular to the gradient.

4.2.3. Rotation rate of a swimmer in a viscosity gradient

Incorporating the flagellar thrust and cell body drag into a force and torque balance on the swimmer, we establish an expression for the rotation rate due to self-propulsion in a viscosity gradient, which we then compare to our experimental results in the main text. A torque balance on the three point force swimmer yields an expression in terms of the rotation rate:

$$\mathbf{0} = \boldsymbol{\Gamma}_1 + \boldsymbol{\Gamma}_2 + \boldsymbol{\Gamma}_{\mathrm{b}} = 2kd^2 \nabla\eta \times \hat{\mathbf{p}} - 8\pi\eta R^3 \boldsymbol{\omega} - 2\pi R^3 \nabla\eta \times \mathbf{V}, \tag{S9}$$

where the first and last terms on the right hand side represent the competition between viscophobic turning due to asymmetric flagellar thrust and viscophilic turning due to translational/rotational cell body coupling, respectively. Likewise, the force balance gives:

$$\mathbf{0} = \mathbf{F}_1 + \mathbf{F}_2 + \mathbf{F}_{\mathrm{b}} = 2k\eta\hat{\mathbf{p}} - 6\pi\eta R\mathbf{V} + 2\pi R^3 \nabla\eta \times \boldsymbol{\omega}. \tag{S10}$$

Recognizing that the rotation rate (equation (S9)) must be proportional to the viscosity gradient $|\boldsymbol{\omega}| \propto |\nabla\eta|$, the final term of the force balance (equation (S10)) is second order in $|\nabla\eta|$. Thus, to first order in the viscosity gradient, swimmers largely move along their body axis with velocity

$$\mathbf{V} = V\hat{\mathbf{p}} = \frac{k}{3\pi R}\hat{\mathbf{p}}.$$

Inserting this result into equation (S9) and eliminating the flagellar thrust coefficient, k, the rotation rate of the swimmer, $\boldsymbol{\omega} = \omega\hat{\mathbf{z}}$, in a viscosity gradient becomes

$$\boldsymbol{\omega} = \omega\hat{\mathbf{z}} = \frac{3d^2 - R^2}{4R^2} \frac{\nabla\eta \times \mathbf{V}}{\eta}. \tag{S11}$$

The prefactor depends solely on cell morphology and dictates whether swimmers are viscophobic or viscophilic. If the center of thrust of the flagella is significantly large compared to the cell radius, $d/R > 1/\sqrt{3} \approx 0.577$, then cells will behave in a viscophobic manner and turn away from high viscosity zones. This condition is consistent with the typical geometry of *C. reinhardtii* and our experimental observations of cell turning (Fig. 2-3). Conservatively, for a typical cell radius [13], $R \approx 5$ μm, flagellar length, $\ell \approx 10$ μm, and flagellar center of thrust, $d \approx \ell/2$, *C. reinhardtii* are generally expected to exhibit viscophobic behavior, $d/R \approx 1 > 0.577$. Furthermore, the prefactor in equation (S11) reduces to $\approx 1/2$, which is similar in magnitude to recent squirmer models [25].

For a one-dimensional viscosity profile, $\eta(x)$, and a viscosity dependent swimming velocity, $\mathbf{V} = V\hat{\mathbf{p}}$, the viscophobic rotation rate is

$$\omega = \frac{3d^2 - R^2}{4R^2} \frac{\partial_{\mathrm{x}}\eta}{\eta} V \sin\theta = -\omega_{\mathrm{visc}} \sin\theta, \tag{S12}$$

where ω_{visc} is the viscophobic turning rate amplitude, θ is the swimming direction, $\hat{\mathbf{p}}$. The geometrical prefactor above depends upon the particular microhydrodynamics of the cell motility, which is beyond the scope of this work. However, this analysis allows us to make quantitative predictions on the scaling of the swimmer turning rate amplitude, which is linearly proportional to the swimming speed, $\omega_{\mathrm{visc}} \propto V$, as might be expected for low Reynolds number locomotion. Importantly, we also find that the turning rate scales with $\omega_{\mathrm{visc}} \propto |\nabla\eta|/\eta$, which agrees with our experiments (Fig. 2) and is also consistent with recent predictions for squirmer models [25].

4.3. Viscophobic turning trajectories

Experimentally measured trajectories of swimming *C. reinhardtii* were demonstrated to turn away from regions of high viscosity and toward regions of low viscosity in the presence of a viscosity gradient (Fig. 2a,b; 65 and 77 tacks

shown, respectively). In particular, cells starting with initial orientation $\theta_0 = \pm\pi/2$ in the center of the maximum viscosity gradient ($\nabla\eta = 7.2 \times 10^{-3}$ cP·μm^{-1}), reveal smoothly curved trajectories stemming from viscophobic turning (Fig. 2b). Here, we examine the equations of motion for viscophobic turning and develop an analytical expression for the expected cell trajectories (Fig. 2b, black curve), which show strong agreement with experiments. The equations of motion for a cell with local swimming speed, V, become

$$\dot{x} = V\cos\theta, \tag{S13}$$
$$\dot{y} = V\sin\theta, \tag{S14}$$
$$\dot{\theta} = -\omega_{\mathrm{visc}}\sin\theta, \tag{S15}$$

where θ is the cell orientation with $\theta = 0$ being in the direction of decreasing viscosity, and ω_{visc} is the viscophobic turning rate amplitude (Fig. 2c,d). Substituting $u = \tan(\theta/2)$, we write

$$\dot{u} = \frac{du}{d\theta}\frac{d\theta}{dt} = -\frac{1}{2}\sec^2\left(\frac{\theta}{2}\right)\omega_{\mathrm{visc}}\sin\theta = -\omega_{\mathrm{visc}}\tan\left(\frac{\theta}{2}\right) = -\omega_{\mathrm{visc}}u,$$

where the over dot indicates a time derivative. Integrating the above expression yields $u(t) = u_0 e^{-\omega_{\mathrm{visc}}t}$ or $\tan(\theta/2) = \tan(\theta_0/2)e^{-\omega_{\mathrm{visc}}t}$. This allows us to rewrite the equations of motion as

$$\dot{x} = V\frac{1-u^2}{1+u^2} = V\frac{1-u_0^2 e^{-2\omega_{\mathrm{visc}}t}}{1+u_0^2 e^{-2\omega_{\mathrm{visc}}t}},$$
$$\dot{y} = V\frac{2u}{1+u^2} = V\frac{2u_0 e^{-\omega_{\mathrm{visc}}t}}{1+u_0^2 e^{-2\omega_{\mathrm{visc}}t}},$$
$$\dot{\theta} = -\omega_{\mathrm{visc}}\frac{2u}{1+u^2} = -\omega_{\mathrm{visc}}\frac{2u_0 e^{-\omega_{\mathrm{visc}}t}}{1+u_0^2 e^{-2\omega_{\mathrm{visc}}t}}.$$

Integrating $\dot{x}$ and $\dot{y}$ gives a parametric expression for the cell trajectory in $x(t)$ and $y(t)$ as

$$x(t) = \int_0^t V\frac{1-u_0^2 e^{-2\omega_{\mathrm{visc}}t'}}{1+u_0^2 e^{-2\omega_{\mathrm{visc}}t'}}dt' = V\left[\frac{\ln\left(u_0^2 + e^{2\omega_{\mathrm{visc}}t}\right)}{\omega_{\mathrm{visc}}} - t - \frac{\ln\left(u_0^2+1\right)}{\omega_{\mathrm{visc}}}\right],$$
$$y(t) = \int_0^t V\frac{2u_0 e^{-\omega_{\mathrm{visc}}t'}}{1+u_0^2 e^{-2\omega_{\mathrm{visc}}t'}}dt' = \frac{2V}{\omega_{\mathrm{visc}}}\left[\arctan\left(\frac{e^{\omega_{\mathrm{visc}}t}}{u_0}\right) - \arctan\left(\frac{1}{u_0}\right)\right].$$

Finally, if we consider initial orientations of the cells in the direction perpendicular to the gradient, $\theta_0 = \pm\pi/2$ and $u_0 = \tan(\theta_0/2) = \pm1$, the trajectory becomes:

$$x(t) = \frac{V}{\omega_{\mathrm{visc}}}\left[\ln\left(1+e^{2\omega_{\mathrm{visc}}t}\right) - \omega_{\mathrm{visc}}t - \ln 2\right], \tag{S16}$$
$$y(t) = 2\frac{V}{\omega_{\mathrm{visc}}}\left[\arctan\left(\pm e^{\omega_{\mathrm{visc}}t}\right) \mp \pi/4\right], \tag{S17}$$

which is shown in Fig. 2b (solid black curve). While this relatively simple analysis omits the stochasticity of cell motility and spatial variations in swimming speed, the model largely captures the shape of the observed cell trajectories in the viscosity gradient.

5. STOCHASTIC DYNAMICS OF VISCOPHOBIC TURNING MOTILITY

5.1. Langevin equations of motion

Microswimmers such as *C. reinhardtii* are subject to stochasticity in their motility stemming from both thermal motion and noise in their flagellar propulsion. In particular, noise in the cell orientation dominates the effects on self-transport compared to translation [26, 27]. We model these effects as a persistent random walk by the addition of a noise term in the orientation dynamics and obtain the Langevin equations of motion

$$\dot{x} = V(x)\cos(\theta), \tag{S18}$$
$$\dot{y} = V(x)\sin(\theta), \tag{S19}$$
$$\dot{\theta} = -\omega_{\mathrm{visc}}\sin(\theta) + \sqrt{2D_{\mathrm{r}}}\xi(t), \tag{S20}$$

where D_r is the effective rotational diffusion coefficient of the cells, and $\xi(t)$ is a Gaussian noise term. The rotational diffusion coefficient is taken to be a constant, as our experiments indicate that it does not vary significantly over the range of viscosities explored in this work (Extended Data Fig. 6 and Methods). Here, we allow the swimming speed, $V(x)$, to vary spatially, which captures the main effect of viscous slow-down on cell translational motion. For simplicity, we take the cell speed profile to be linear across the channel $V(x) = (V_{max} - \Delta V) + \Delta V x/W$, where V_{max} and V_{min} are the cell speed at $x = W$ and $x = 0$, respectively, and $\Delta V = V_{max} - V_{min}$. We also define the dimensionless variables

$$X = \frac{x}{W}, \qquad Y = \frac{y}{W}, \qquad T = \frac{tV_{max}}{W},$$

which lead to the following dimensionless coefficients

$$\tilde{D}_r = \frac{D_r W}{V_{max}}, \qquad \tilde{V}(X) = \left(1 - \frac{\Delta V}{V_{max}}\right) + \frac{\Delta V}{V_{max}} X, \qquad \tilde{\omega}_{visc} = \omega_{visc} \frac{W}{V_{max}}.$$

Finally, substituting into the Langevin equations of motion, we obtain

$$\dot{X} = \tilde{V}(X)\cos(\theta), \tag{S21}$$

$$\dot{Y} = \tilde{V}(X)\sin(\theta), \tag{S22}$$

$$\dot{\theta} = -\tilde{\omega}_{visc}\sin(\theta) + \sqrt{2\tilde{D}_r}\xi(T), \tag{S23}$$

where the over dot now indicates derivatives with respect to dimensionless time, T. Thus, the Langevin equation dynamics are ultimately dictated by the three dimensionless parameters $\tilde{D}_r$, $\Delta V/V_{max}$, and $\tilde{\omega}_{visc}$. Due to the weak variation of rotational diffusivity, $\tilde{D}_r$ was held constant. Simulations were performed as described below and as a function of $\Delta V/V_{max}$ and $\tilde{\omega}_{visc}$, where the results are shown in below (Extended Data Fig. 7-8) and in the main text (Fig. 3).

5.2. Semi-empirical relation for turning rate and swimming speed contrast

Langevin simulations facilitate the arbitrary exploration of the parameter space of ω_{visc} and $\Delta V/V_{max}$ (Fig. 3c). However, experimentally, both parameters directly depend upon the viscosity gradient (Figs. 1f and 2d), which is formed by two fluids of high and low viscosity, initially injected into the test channel. During the experiment time, the extrema of the channel evolve to corresponding viscosities η_{max} and η_{min} (Fig. 1g), whose ratio parametrically controls the trajectory of experiments in the parameters space (Fig. 3c). Here, we develop a semi-empirical relationship for the evolution of ω_{visc} with $\Delta V/V_{max}$ based on this control parameter η_{max}/η_{min}. The swimming speed contrast is evaluated from the power-law relationship (Fig. 1f) of the swimming speed with viscosity, $V = a\eta^{-b}$:

$$\frac{\Delta V}{V_{max}} = 1 - \frac{V_{min}}{V_{max}} = 1 - \left(\frac{\eta_{max}}{\eta_{min}}\right)^b. \tag{S24}$$

Similarly, the observed linear relationship for viscophobic rotation rate (Fig. 2d; equation S12) can be written in terms of the approximate viscosity gradient $\nabla\eta \approx (\eta_{max} - \eta_{min})/W$ and mean viscosity $\eta_0 \approx \beta(\eta_{max} + \eta_{min})/2$:

$$\omega_{visc} \approx \frac{1}{2}V^* \frac{\nabla\eta}{\eta_0} \approx V^* \frac{\beta}{W}\left(1 - \frac{2}{1 + \eta_{max}/\eta_{min}}\right), \tag{S25}$$

where $V^* = 66\ \mu m/s$ is based on the fit in Fig. 2d and $\beta = 1.7 \pm 0.1$ accounts for the observed magnitude of the gradient in the central region of the channel relative to the slightly sigmoidal shape in our experiments (Fig. 1g). Thus, equations (S24) and (S25) form a parametric set of equations in η_{max}/η_{min}, which well represent the experimental observations and matching Langevin simulations (Fig. 3c, black curve).

5.3. Fokker-Planck model of viscophobic reorientation dynamics

To complement the experiments and Langevin simulations discussed above, we also employ a simplified one-dimensional Fokker-Planck (FP) model. The FP model takes the form of a rotational advection diffusion equation to describe the joint probability density of orientation and time, $p(\theta, t)$. The swimming cells are subjected to an

orientation dependent rotation rate due to viscophobic turning motility, and they experience random orientational fluctuations parameterized by rotational diffusion:

$$\frac{\partial p}{\partial T} = \tilde{D}_\text{r} \frac{\partial^2 p}{\partial \theta^2} + \tilde{\omega}_\text{visc} \frac{\partial}{\partial \theta} \left(p \sin \theta \right), \tag{S26}$$

where $\tilde{D}_\text{r}$ and $\tilde{\omega}_\text{visc}$ are constants, as above. The equation is solved using a second-order finite difference scheme for various $\tilde{\omega}_\text{visc}$, corresponding to experimental conditions that include both control experiments ($\nabla \eta = 0$ cP·μm^{-1}; Extended Data Fig. 8c) and maximum viscosity gradient conditions ($\nabla \eta = 7.2 \times 10^{-3}$ cP·μm^{-1}; Extended Data Fig. 8f). As with the Langevin simulations, the turning rate amplitude and rotational diffusion parameters are taken directly from experimental measurements, and thus, there are no fitting parameters. The result is viewed as a conditional probability density, $p(\theta|t)$, that illustrates the time evolution of the cell orientation distribution (Extended Data Fig. 8). Physically, the cell rotation rate in the viscosity gradient drives a condensation, or narrowing, of the orientation distribution toward the low viscosity direction ($\theta = 0$). This effect is in competition with random cell reorientation, which tends to widen the distribution. The results of the FP theory were compared to both experiments (Extended Data Fig. 8a,d) and Langevin simulations (Extended Data Fig. 8b,e), which remain accurate (Fig. 3b) when the cells are isolated far from the channel walls.

6. SUPPLEMENTARY FIGURES

Fig. S1. **Comparison of measured local cell density and swimming speed for various assay time intervals. a,** The distribution of *C. reinhardtii* cell density in the maximum viscosity gradient tested $(\nabla\eta = 7.2 \times 10^{-3}$ cP $\cdot\mu\text{m}^{-1})$ between 30-60 min and 60-90 min intervals remain relatively constant, with a mean percent difference of 1.84% and 2.33%, respectively, when compared to the full analysis time interval of 30-90 min. **b,** The distribution of the cell's local mean swimming speed exhibits similarly small deviations for the 30-60 min and 60-90 min time intervals, having a mean percent error of 5.08% and 6.08%, respectively, when compared to the full analysis time interval of 30-90 min.

Fig. S2. **Control experiments exhibit no discernible diffusiophoretic drift of non-motile *C. reinhardtii*. a,** Visco-taxis control experiments using (fixed) non-motile cells in the maximum viscosity gradient tested ($\nabla \eta = 7.2 \times 10^{-3}$ cP $\cdot \mu m^{-1}$) show no significant observable displacement over the course of a 90 min experiment under identical conditions to Fig. 1h,i (Scale bar, 100 μm). **b,** Maximum cell displacements in the viscosity gradient direction from **a** are minute (on the order of ≈ 1 cell diameter or 1% of the channel width) and consistent with Brownian diffusion of the non-motile cells.

[1] Sager, R. & Granick, S. Nutritional studies with *Chlamydomonas reinhardi*. *Ann. N. Y. Acad. Sci.* **56**, 831–838 (1953).

[2] Devanand, K. & Selser, J. C. Polyethylene oxide does not necessarily aggregate in water. *Nature* **343**, 739–741 (1990).

[3] Martinez, V. A. *et al.* Flagellated bacterial motility in polymer solutions. *Proc. Natl Acad. Sci. USA* **111**, 17771–17776 (2014).

[4] Kirby, B. J. *Micro-and nanoscale fluid mechanics: transport in microfluidic devices* (Cambridge University Press, 2010).

[5] Rafaï, S., Jibuti, L. & Peyla, P. Effective Viscosity of Microswimmer Suspensions. *Phys. Rev. Lett.* **104**, 098102 (2010).

[6] Gardel, M. L., Valentine, M. T. & Weitz, D. A. Microrheology. In Breuer, K. S. (ed.) *Microscale Diagnostic Tech.*, 1–49 (Springer, Berlin, Heidelberg, 2005).

[7] Liu, J. *et al.* Microrheology Probes Length Scale Dependent Rheology. *Phys. Rev. Lett.* **96**, 118104 (2006).

[8] Waigh, T. A. Microrheology of complex fluids. *Reports Prog. Phys.* **68**, 685–742 (2005).

[9] Ebagninin, K. W., Benchabane, A. & Bekkour, K. Rheological characterization of poly(ethylene oxide) solutions of different molecular weights. *J. Colloid Interface Sci.* **336**, 360–367 (2009).

[10] Polin, M., Tuval, I., Drescher, K., Gollub, J. P. & Goldstein, R. E. *Chlamydomonas* Swims with Two "Gears" in a Eukaryotic Version of Run-and-Tumble Locomotion. *Science* **325**, 487–490 (2009).

[11] Goldstein, R. E. Green Algae as Model Organisms for Biological Fluid Dynamics. *Annu. Rev. Fluid Mech.* **47**, 343–375 (2015).

[12] Harris, E. H. *Chlamydomonas* as a model organism. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **52**, 363–406 (2001).

[13] Harris, E. H. *The Chlamydomonas Sourcebook: Introduction to Chlamydomonas and its laboratory use*, vol. 1 (Academic Press, 2009).

[14] Qin, B., Gopinath, A., Yang, J., Gollub, J. P. & Arratia, P. E. Flagellar Kinematics and Swimming of Algal Cells in Viscoelastic Fluids. *Sci. Rep.* **5**, 9190 (2015).

[15] Berg, H. C. & Turner, L. Movement of microorganisms in viscous environments. *Nature* **278**, 349–351 (1979).

[16] Schneider, W. R. & Doetsch, R. N. Effect of Viscosity on Bacterial Motility. *J. Bacteriol.* **117**, 696–701 (1974).

[17] Berg, H. C. & Brown, D. A. Chemotaxis in *Escherichia coli* analysed by Three-dimensional Tracking. *Nature* **239**, 500–504 (1972).

[18] Garren, M. *et al.* A bacterial pathogen uses dimethylsulfoniopropionate as a cue to target heat-stressed corals. *ISME J.* **8**, 999–1007 (2014).

[19] Moran, J. L. & Posner, J. D. Phoretic Self-Propulsion. *Annu. Rev. Fluid Mech.* **49**, 511–540 (2017).

[20] Schnitzer, M. J. Theory of continuum random walks and application to chemotaxis. *Phys. Rev. E* **48**, 2553–2568 (1993).

[21] Drescher, K., Goldstein, R. E., Michel, N., Polin, M. & Tuval, I. Direct Measurement of the Flow Field around Swimming Microorganisms. *Phys. Rev. Lett.* **105**, 168101 (2010).

[22] Bennett, R. R. & Golestanian, R. A steering mechanism for phototaxis in *Chlamydomonas*. *J. R. Soc. Interface* **12**, 20141164 (2015).

[23] Oppenheimer, N., Navardi, S. & Stone, H. A. Motion of a hot particle in viscous fluids. *Phys. Rev. Fluids* **1**, 014001 (2016).

[24] Liebchen, B., Monderkamp, P., Ten Hagen, B. & Löwen, H. Viscotaxis: Microswimmer Navigation in Viscosity Gradients. *Phys. Rev. Lett.* **120**, 208002 (2018).

[25] Datt, C. & Elfring, G. J. Active Particles in Viscosity Gradients. *Phys. Rev. Lett.* **123**, 158006 (2019).

[26] Howse, J. R. *et al.* Self-Motile Colloidal Particles: From Directed Propulsion to Random Walk. *Phys. Rev. Lett.* **99**, 48102 (2007).

[27] Rusconi, R., Guasto, J. S. & Stocker, R. Bacterial transport suppressed by fluid shear. *Nat. Phys.* **10**, 212–217 (2014).