Title: Mass spectrometry and CLIP-seq analysis reveal BCL11b interactions with RNA processing pathways.

Haitham Sobhy[1]*, Marco De Rovere[1]*, Amina Ait-Ammar[1,2]*, Clementine Wallet[1], Fadoua Daouad[1], Carine Van Lint[2&], Christian Schwartz[1&] and Olivier Rohr[1&]

[1] University of Strasbourg, EA 7292, DHPI, IUT Louis Pasteur, France

[2] Université Libre de Bruxelles, ULB, Gosselies, Belgium

* HS, MDR and AAA can be considered as equal contributors

& CVL, CS and OR can be considered as equal contributors

Correspondence should be addressed to Haitham Sobhy (haithamsobhy@gmail.com) and Olivier ROHR (olivier.rohr@unistra.fr)

Conflict of interest: Not known

Author contribution: HS conceived and designed bioinformatics analysis; HS, MDR and AAA conceived and performed experiments; CW and FD provided scientific and logistic supports; CVL, CS and OR conceived the experiments, supervised the work and supported with resources. HS drafted the manuscript, all authors agreed for the final version of the manuscript.

Data deposited on SRA database with ID PRJNA661202

Abstract: 168 words

Text: 4076 words (4832 words with legend)

Figures: 4

Abstract (168 words)

Although BCL11b (B-cell lymphoma/leukemia 11B, CTIP2) is a well-known transcription repressor and tumor suppressor, its functions and cellular pathways are largely unknown. Here, we show that BCL11b interacts with RNA splicing/processing and nonsense-mediated decay (NMD) proteins, including FUS, SMN1, UPF1 and Drosha. Mass spectrometry analysis (LC-MS/MS) shows that BCL11b interacts with histones, polymerases, and chromatin remodeling (CHD, SWI/SNF, and topoisomerase) proteins. BCL11b-bound RNAs were UV cross-linked and sequenced (CLIP-seq) showing that BCL11b binds to coding and noncoding RNAs (ncRNAs). Surprisingly, RNA transcripts and proteins produced by the same genes like FUS, ESWR1, CHD and Tubulin, were found bound to BCL11b. Deeper analysis of the CLIP-seq data further suggested that BCL11b binds to nonsense mediated RNA decay and retained introns transcripts. Our study is the first genome-wide study of BCL11b-protein and BCL11b-RNA interactants. Our results suggest that the functions of BCL11b are not restricted to the regulation of gene transcription. BCL11b may also control physiologic and physiopathologic pathways by direct bindings to protein complexes, coding RNA and non-coding RNA.

Importance

First genome-wide BCL11b-proteins interactomics

First genome-wide BCL11b-RNA interactomics

BCL11b interacts with RNA processing and RNA splicing proteins

BCL11b interacts with neurodegenerative genes and sarcoma genes

Keywords

Transcription factor BCL, zinc figure family, RNA processing, RNA splicing, gene

expression regulation, T-cell development, neuron cell development,

Introductions

The transcription regulator B-cell lymphoma/leukemia 11B (Bcl11b, also known as COUP-TF-interacting protein 2, CTIP2) plays crucial roles in the epigenetic regulation of gene transcription and specifically in the control of the elongation process by interacting with an inactive form of the p-TEFb complex (7SK RNA, CDK9, cyclin T1, and HEXIM1) (Cherrier et al., 2013). BCL11b, which harbors six zinc finger domains (ZnFs) of C2H2-type, shares >60% homology with the human BCL11a, as well as the mouse, chicken, and Xenopus homologues. BCL11 proteins were isolated from T-cell lines derived from patients with T-cell leukemia, suggesting the role of BCL11 proteins in blood cell development and lymphomagenesis, reviewed in (Fu et al., 2017; Lennon et al., 2016, 2017). Beside, BCL11b was deleted in γ-ray induced mouse lymphomas, suggesting a role as a tumor suppressor and modulator of radiation-induced DNA damages, reviewed in (Kominami, 2012).

Studying the function of BCL11b by knocking out the gene (loss-of-function) is challenging (Longabaugh et al., 2017). The mice that lack the two alleles of BCL11b die shortly after birth exhibiting defects in multiple tissues, including immune system, central nervous system (CNS), skin, hair cells in cochlea, teeth, and thymus among other organs (Kominami, 2012) , which suggests the importance of BCL11b during the development. As consequences of absence of the BCL11b-deficient cells, studying the functions and pathways triggered by BCL11b is challenging.

In this study, we used co-immunoprecipitation of proteins followed by mass spectrometry (LC-MS/MS, or MS for short) as well as immunoprecipitation of cross-linked RNA (CLIP-seq), to identify the proteins and RNAs bound to BCL11b respectively. Our results reveal that the partners of BCL11b are involved in cellular pathways dedicated to neuron development, the control of the cell cycle, RNA splicing and neurodegenerative diseases.

Results

Protein interactomics and pathways of BCL11b

To characterize the protein interactants and pathways triggered by BCL11b, we used HEK cells overexpressing Flag-BCL11b. After proteins pull-down using anti-flag antibodies, we performed quantitative LC-MS/MS. After statistical analysis, we identified 629 BCL11b protein partners (supplementary method file S1 and supplementary dataset S1). Among the interacting proteins, we can confirm the previously described interactions between BCL11b and the P-TEFb complex (including CDK9 and CycT1) (Cherrier et al., 2013), HDACs (Marban et al., 2007), HMGA1 (Eilebrecht et al., 2014), HP1 proteins (Rohr et al., 2003), KU proteins (Shadrina et al., 2020), and DCAF1 (Forouzanfar et al., 2019) as a validation of our experimental approach (Figure 1A).

BCL11b interacts with proteins dedicated to genetic and epigenetic regulations

Performing gene ontology (GO) analysis (figure 1B and supplementary data S2), we observed that BCL11b interacts with proteins and complexes that are involved in cellular genetics, DNA replication and regulations of mitosis (Figure 1B), including proteins that are involved in the regulation of gene transcription (CCNT1, CDK9, SMARCAs and SMARCCs), and epigenetic regulators, including HDACs, CHDs, EHMTs, KMT2A (Figure 1B). Indeed, BCL11b binds to more than 20% of the total protein components of H2AX, NuRD and SNW1 complexes, among other proteins involved in chromatin organization, and DNA recombination (Figure 1C). Additionally, we observed BCL11b is associated with proteins involved in DNA repair and cellular response to stimuli (such as BRCA2, MCMs, PARP1, and SMCs proteins). In agreement to our results, BCL11b was previously shown to interact with Ku70 and Ku80 proteins, which have an important role in the non-homologous end-joining DNA repair pathway (Shadrina et al., 2020).

BCL11b interacts with structural proteins and the nuclear pore complex

Among our novel findings, we observed BCL11b interacts with structural proteins involved in cytoskeleton, and motor activity, such as multiple isoforms of tubulin (supplementary data S1 and S2). Another significant GO term is the nucleocytoplasmic shuttling pathways, which include transport of the RNA or ribonucleoprotein complexes, such as NUP133, NUP153, NUP93 and NUP214. Noteworthy, nucleoporins could have molecular roles beyond the nucleocytoplasmic shuttling, such as regulations of gene transcription and chromatin organization (Sun et al., 2019).

BCL11b interacts with proteins involved in RNA translation and RNA splicing

The role of BCL11b during RNA processing have not been described by any previous research. We found that multiple proteins bound to BCL11b are involved in RNA processing pathways such as translation of RNA and ribosome assembly, such as 40S and 60S ribosomal subunits (figure 1C). Moreover, BCL11b interacts with RNA-binding proteins, including RNA splicing and ribonucleoproteins (RNPs), such as spliceosomes. Among these complexes, the survival motor neuron (SMN) protein complex, which is made by binding of SMN protein with other proteins so-called GEMIN and Sm proteins (Matera and Wang, 2014). It is known that SMN complex assembles in nucleus before it is translocalized to the cytoplasm. SMN complex is required for assembly of RNP complexes, which are the building block of spliceosomes. Other RNPs that are involved in post-transcriptional RNA regulation, include small nuclear RNP (snRNPs), heterogeneous nuclear RNP (hnRNP), serine/arginine (SR)-rich proteins, and small nuclear RNA (snRNAs). BCL11b interacts to 22 different hnRNPs or snRNPs, and about 17% of proteins that constitute the spliceosome (figure 1B-1D).

On the other hand, FUS is another crucial splicing factor, which is involved in regulations of RNA splicing, transport of RNA, and regulation of gene expression as response to DNA damage

(Chen et al., 2019). FUS can accumulate near the transcription start sites (TSS) and binds to

transcription initiation factors to inhibit phosphorylation of the RNA polymerase 2 C-terminal

domain (POLR2-CTD) (Chen et al., 2019). We performed co-IP and immunoblot to confirm

that BCL11b interacts with SMN1 and FUS, which are among the hallmarks of RNA splicing

processes (Figure 1E and 1F). Our IP-MS and IP-immunoblot confirm that BCL11b is able to

interact with SMN1 and FUS proteins, and therefore BCL11b could have a role in regulation

of RNA splicing and translation of RNA.

BCL11b interacts with the Drosha and the DGCR8 protein complexes

FUS is a member of the Drosha and the DGCR8 complexes, which are able to regulate

gene expression at transcriptional and post-transcriptional levels (Pong and Gullerova, 2018).

On the other hand, Ago is the key protein in RNA interference (RNAi) pathway. Ago2 can bind

to Dicer or TRBP proteins forming TRBP containing complex. Drosha and TRBP complexes

can regulate translation of mRNA, and degradation of mRNA of certain transcripts (Pong and

Gullerova, 2018). Here, by IP-MS experiment, we found that BCL11b is associated with 15 of

the 20 proteins of the Drosha complex and 10 proteins of the 11 proteins of the DGCR8 complex

(Figure 1D, and 1E), in addition to 20 proteins of TRBP complex, including Ago2. To confirm

these results we performed co-IP experiment, which confirms the association of BCL11b with

FUS and Drosha (Figure 1F, 1G and 1H) and further suggest that BCL11b may coordinate with

FUS and Drosha to regulate gene expression at transcriptional and post-transcriptional levels

and to control mRNA translation and stability.

BCL11b binds to protein from the nonsense-mediated mRNA decay (NMD) pathway

NMD pathway is crucial to decay mRNA transcripts with premature stop codons

(Kurosaki et al., 2019). These transcripts lead to gain- or loss-of-function of some genes, which

could be deleterious during the development. UPF1 is the hallmark of the NMD pathway.

Among the significant IP-MS hits, we observed interactions between BCL11b and proteins

involved in the NMD pathway. Out of 120 proteins identified in this specific GO molecular

pathway, 61 were found bound to BCL11b (figure 1C). Co-IP experiments confirmed that

BCL11b associates with UPF1, which is the hallmark of NMD pathway, and thereby this

confirms that BCL11b interacts with the protein complex dedicated to NMD (Figure 1I). Again,

these results show that BCL11b is involved in the regulation of gene expression at the post-

transcriptional level (Figure 1F-1H).

BCL11b is associated with cellular RNA

Although BCL11b has six C2H2 ZnF domains, to our knowledge the RNA regulation

activities has not been described for any of the two BCL11 proteins, However, BCL11b can

bind the 7SK ncRNA to regulate transcription elongation (Cherrier et al., 2013; Eilebrecht et

al., 2014). Our MS proteomics data show that BCL11b binds to RNA-binding proteins in

complexes dedicated to RNA processing and RNA splicing. To further study the RNA-binding

activity and obtain genome-wide landscape of BCL11b-bound RNAs, we cross-linked the

BCL11b protein to cellular RNAs using UV radiation before immunoprecipitation and

sequencing (CLIP-seq). These experiments have been performed by targeting the endogenous

BCL11b in microglial cells, and an overexpressed Flag-BCL11b in HEK293 cells. We analyzed

the results by two approaches, as the following:

First, number of CLIP reads mapped to a gene feature were counted and the genes that

are enriched over the control samples were selected (see method in Suppl. file S1), which lead

to a dataset of 740 different genes (figure 2, 3A, S1-S5 and Suppl. data S3). We found that

about 96% of the BCL11b-bound RNAs are mapped to protein-coding genes (Table S1).

BCL11b tends to bind to transcripts of long genes. Among the genes that have high numbers of

CLIP reads, we found that about 20% are over 100kb in length, whereas the average length of

these genes is 61kb, which is two-fold the genomics length (~30kb). The longest gene is Arf-

GAP with GTPase, ANK repeat and PH domain-containing protein 1 (AGAP1, NCBI GeneID:

116987), which is 637,753 bp in length. In agreement with previous studies focusing on

BCL11b control of cellular gene expression (Kominami, 2012; Wakabayashi et al., 2003), we

found that the GO terms of BCL11b-bound RNA include genes from neuronal development

and neurogenesis of multiple parts of brain (e.g. midbrain, hindbrain and substantia nigra).

BCL11b binds to RNA of proteins that have been described to regulate proliferation,

regeneration and migration of neuronal and non-neuronal cells, including glial cells and

Schwann cells (details are found in figures 3B and S6, Suppl. data 4). However, BCL11b was

also found associated with RNA from genes involved in the development of the immune

system, hemopoiesis, blood vessels, and hair follicles. Finally, we found RNA coding for actin

and tubulin associated with BCL11b as well as RNA coding for proteins of the major and the

minor spliceosome (Figure 1E, Table S2, Suppl. data 5).

As a second analysis approach, we divide the whole genome into 10kb regions (bins)

and we count the number of reads corresponding to each bin (see method in Suppl. file S1). For

additional confirmation, we performed peak calling to cluster the RNAs into bins. We then

selected the regions that are enriched over the control. The results show that over 40% of CLIP

reads are mapped within 0-50kb down-stream of the transcription start sites (TSS) (Figure S7,

Suppl. data S6, S7). Noting that some these RNAs mapped to long genes, we concluded that

about half of the CLIP reads are within the genes and at least one third could be located in the

intergenic regions. Aligning the sequences, we found two conserved motifs (CUCRGCCU and

UCCCAGCW) in BCL11b-bound RNA regions (figure 3C). Gene ontology analysis for the

genes localized in these regions suggested their involvement into structural process,

extracellular matrix organization, and response to stress (Suppl. data S6, S7). The gene products

are involved in angiogenesis, p53 and inflammation pathways, cytoskeletal regulation and

multiple diseases such as cancers, Huntington's disease and Alzheimer (Figure S8, Suppl. data

S6, S7).

Taken together, our results demonstrate that half of the BCL11b-binding RNAs are

transcribed from long genes and mapped to the genic regions further suggesting an impact of

BCL11b on protein expression or function at post-transcriptional levels.

BCL11b binds to RNA transcripts and proteins product of the same gene

We found BCL11b associated with 137 RNA transcripts and proteins encoded by the

same gene (Figure 3A, and Suppl. data S8). Among them, we found EWSR1, FUS, SPFQ,

UPF1, tubulins, CHDs and hnRNPs transcripts and proteins. Interestingly, GO analysis reveals

that these proteins are involved in RNA processing, NMD pathway and translation (Figure 3D,

Suppl. data S8, S9). Interestingly, tubulin was reported to use a unique mechanism to auto-

regulate its expression level within the cell to ensure a stable production of the protein in cells,

and over-expression of tubulin gene does not lead to increase its protein expression level (Gasic

and Mitchison, 2019). The mechanism of tubulin auto-regulation is largely unknown. However,

it thought that tubulins α and β could polymerize through an unknown mediator, which then

bind to the nascent tubulin peptide (Gasic and Mitchison, 2019). By unknown mechanism, the

mediator can bind to ribosomes and/or ribonucleases to terminate the translation. We found that

BCL11b interacts with multiple isoforms of tubulin proteins, tubulin RNAs, as well as

ribonucleases, and ribosomes (Figure 3E). In addition, we have noted that BCL11b binds to

multiple members of the same protein family, such as tubulins, actins, RNPs, or ribosomes.

Again, these results may support the involvement of BCL11b in the regulation of protein

expressions at post-transcriptional levels.

BCL11b binds differentially transcripts of the same genes

To further study the association of BCL11b with cellular RNA, we counted the number

of reads per exon for CLIP reads that mapped to genes. We found that more than 75% of the

CLIP reads were mapped to exons (Figure 3F). Having a deeper look to the CLIP-seq reads on

genome browsers, we found that when reads were mapped to intronic regions of genes such as

EWSR1, FUS, and SRRM2, the locations of the reads corresponded to the coordinates of RNA

isoforms that do not code for proteins (we refer to them here as INCP), such as retained introns

(IR), nonsense mediated decay (NMD), non-stop decay or processed transcript, (Figure 2,

Suppl. Figure S2-5). This result is consistent with our results demonstrating that BCL11b

interacts with proteins involved in RNA splicing and NMD pathways, including FUS, SMN1,

RNPs, Drosha, and UPF1. These results show that BCL11b can regulate gene expression at

post-transcriptional levels (Figure 1F-1I).

To strengthen this finding, we collected the transcripts (or RNA isoforms) of 23 genes

that encode for isoforms coding for proteins (ICPs), and isoforms that do not code for proteins

(INCPs), such as NMD and IR. We then counted the number of reads that correspond to each

isoform. We found that BCL11b binds to different isoforms of the same gene in differential

manner (Figure 3G, Suppl. data S10). For example, TOP3A gene encodes for 21 RNA isoforms,

similarly, ESWR1, Drosha, SIRT7 and RAD52 genes encode for 20-21 RNA isoforms. The

fold increase to control of CLIP-seq for TOP3A ranges from 1.1 to 2.7 folds, whereas, the

binding to RNA of RAD52 is up to 7-folds increase (log2 fold change ranges from 0.8 to 2.8

folds) (Figure 3G). The same concept applies to SMN1 (10 isoforms), COL27A1 (8 isoforms)

and RAD51 (11 isoforms). We counted the CLIP-seq reads per each transcript (or isoform) and

then normalized to the length of the transcripts. We found that isoforms coding for protein

(ICPs) have 2 folds CLIP reads than INCPs (Figure 3H). The Spearman correlation rank (R)

between the CLIP reads and transcripts length for ICPs is 0.51 and 0.44 for INCP, which

suggest that BCL11b tends to bind to long isoforms of ICPs and less degree INCPs. Together

these results suggest that BCL11b preferentially binds to long RNA isoforms that code for

proteins.

We still noticed some CLIP reads that correspond to intronic regions within ICPs, but

the CLIP reads are mapped to the coordinates of exons in other INCP isoforms. Noteworthy,

NMD or IR transcripts introduce additional exons instead of introns. To explain this concept,

we highlighted the genomics introns in pink colors, see (Figure 2 and Figures S2-S5). Although these intron regions are usually removed in case of ICP transcripts (thick dark bars in e panel in Figure 2,), we found that part of these regions corresponded to exons in INCPs (light blue bars in e panel in Figure 2). Computationally, we counted number of reads mapped to exons and introns, we normalized them to the number of exons or introns as shown in Figure 3I and Suppl. data S10. If BCL11b binds to exons regions of RNA, we expect to find that most of the CLIP reads are biased to exons coordinates. However, in Figure 3I, we observed that most of the reads are mapped to introns. In case of INCPs, the number of reads were almost equally distributed throughout the exons and introns. Statistically, we calculated the Spearman correlation rank (R) between number of reads and number of exons, which show that number of exons correlate with number of CLIP reads, in case of ICPs, but not in case of INCPs (R of INCPs=0.2, and R of ICPs=0.46). This result suggests that INCPs with few numbers of exons have multiple CLIP reads. The DNA (cytosine-5-)-methyltransferase 1 (DNMT1) transcripts are good examples that support this observation. Number of reads mapped to IR transcripts are 2-3 fold higher than the number of reads mapped to INCPs transcripts. Indeed, the 5 exons IR transcripts (ENST00000591764, with 589 reads) have 3-folds more reads than the 9 exons protein coding transcripts (ENST00000588952, with 196 reads) (Figure 3J, Suppl. data S10).

Taken together, Our CLIP-seq results confirm that BCL11b binds to complexes with ncRNA and ribonucleoproteins (Figure 2 and S2-S5). The results support our proteomics findings that BCL11b interacts with RNA processing and RNA splicing proteins, such as FUS, SMN1, Drosha complex and NMD complex (Figure 1). Our results further suggest that BCL11b may be able to differentiate between the types of RNA isoforms to contribute to the selection of the one to be translated. Together, BCL11b could interact with other proteins, such as FUS or UPF1 to regulate mRNA translation of selected genes.

Involvement of BCL11b- associated factors in diseases

Described as a tumor suppressor gene, BCL11b expression level was reported to be

modulated in T-Cell Lymphoma (Fu et al., 2017; Kominami, 2012). Previous reports studying

Ewing Sarcoma (Orth et al., 2020; Wiles et al., 2013), spinal muscular atrophy (SMA) (Doktor

et al., 2016; Maeda et al., 2014), and Parkinson's disease (PD) (Scherzer et al., 2007) showed

differential expression of BCL11b. In addition, since BCL11b was reported to be involved in

development, it is speculated that BCL11b could have roles in neurodegenerative diseases

(Lennon et al., 2017), particularly in motor neuron amyotrophic lateral sclerosis (ALS) disease

(Lennon et al., 2016), and Huntington's disease (Ahmed et al., 2015). However, the role of

BCL11b in these disease was not clear, because of the lack of studies related to protein and

RNA interactions of BCL11b nor its functional complexes, it was challenging to understand

the exact role of BCL11b in these diseases.

As we discussed above, using IP-MS, and CLIP-seq experiments, we detected

interactions between BCL11b and multiple proteins that are involved in diseases such as:

EWSR1, which is known to contribute to Ewing's sarcoma (Wiles et al., 2013); SMN1, which

is involved in SMA (Maeda et al., 2014); FUS, TDP-43 and TAF15 proteins that are involved

in ALS (Yamazaki et al., 2012); SRRM2, which is a key factor causing Parkinson's disease

(Shehadeh et al., 2010) and 7SK, MALAT1, NEAT1 and TUG1 ncRNAs, SFPQ protein, RNPs

and spliceosomes proteins that are known to contribute to cancer, as well as the motor neuron

SMA and ALS diseases (Bhat et al., 2020; Briese et al., 2018; Wu and Kuo, 2020). Noteworthy,

FUS associates with SMN, RNPs and RNA-binding proteins to form a large complex (De Santis

et al., 2017; Yamazaki et al., 2012). We found that one third of these proteins were found bound

to BCL11b by LC-MS, which suggest the ability of BCL11b to form complexes with FUS or

SMN1 (table S2, S3). On the other hand, DNA damage pathway may regulate RNA splicing

process by selecting specific RNA isoforms to be translated (Shkreta and Chabot, 2015). FUS,

Drosha and Ago2 are examples of regulators that are recruited in response to DNA damage to

regulate gene expression of certain protein isoforms (Chen et al., 2019; Pong and Gullerova,

2018).

Our CLIP-seq and RNA IP results confirmed the binding of BCL11b to the RNAs

coding for FUS, SFPQ and SRRM2 genes and specifically to exon and exon-intron junctions

regions (Figure 2, 4B, S1-S3, S9 and S10, table S4, Suppl. data S10). Noteworthy, we could

not observe significant increase of the gene expression of RNA encoded from the same regions

(using RT-qPCR) after over-expression of BCL11b (Figure 4C). These results suggest that

BCL1b could contribute to isoform selection by binding to RNA isoforms, without impacting

the global level of the transcripts.

Discussion and Conclusion

This study describes for the first time BCL11b-associated proteome and RNAome. To

our knowledge this is the first time to report a protein to bind to RNA transcript and protein

encoded from the same gene, such as FUS, Figure 1F, 1G, 3 and 4B. Our results suggest that

BCL11b is involved in multiple pathways and functions. In addition to its function as

transcription regulator, BCL11b binds to proteins involved in cytoskeleton and development,

as well as ribonucleic complexes such as ribosomes and RNPs. We confirmed that BCL11b

binds to proteins involved in RNA processing, such as RNA splicing and NMD pathway, in

addition to the ncRNA that are involved in RNA splicing. Our results show that BCL11b binds

to DNA damage repair proteins, to chromatin remodeling and histone-modifying proteins

(epigenetic modifications). One major GO term found is the assembly and disassembly of the

complexes. We speculate that one of the major roles of BCL11b is to assemble various

complexes to perform a function.

There are number of regulators found in close association to chromatin as checkpoint

safeguard. As response to cell cycle checkpoints, stimuli, or DNA damage, the safeguard factors

regulate the expression level of some specific genes. Here, it emerges the importance of a

regulator that can coordinate between replication, transcription and translation (Shkreta and

Chabot, 2015). First, we demonstrate that BCL11b binds to RNA processing proteins (such as

FUS, Drosha, Ago2, UPF1, and RNA splicing factors, Figure 1F-1I) revealing a role in the

regulation of gene expression in response to DNA damage. Then, we highlighted the potential

role of BCL11b in autoregulation of some specific gene translation such as tubulin (Figure 3E).

We found that BCL11b preferentially binds to RNA isoforms coding for proteins

(Figure 3F). We also observed multiple MS hits that correspond to 40S and 60S ribosome

complexes (Figure 1C). On the other hand, BCL11b was found associated with isoforms that

do not code for proteins (INCPs), such as NMD and IR. Generally, translation is regulated by

selecting the desired mRNA for being translated, whereas, the RNA decay pathways, such as

Drosha, UPF1 (NMD pathway), and Ago (RNA interference) are able to decay or silence the

undesirable RNA isoforms (Tuck et al., 2020; Wang and Aifantis, 2020). Here, we showed

based on reliable methods (IP, LC-MS, CLIP-seq, and RIP-PCR) that BCL11b contributes to

the selection of RNA isoforms. Selecting the isoform (so-called isoform switching) is a crucial

process through which cells select a specific RNA isoform during the development or immune

response. Some isoforms might be needed at specific steps during development, cell cycle or

response to stimuli. Therefore, malfunction of RNA splicing or processing has a great impact

on cell proliferation and cancer prognosis (Wang and Aifantis, 2020).

In conclusion, our analysis of BCL11b-protein and BCL11b-RNA interactions provide

the first global overview of BCL11b partners. The results suggest that BCL11b may bind to

RNA processing proteins to regulate translation and/or select the desirable RNA isoform. One

can speculate that depletion of BCL11b could lead to isoform switching and translation of

wrong isoform of the protein (misfolded or short) but this needs to be formally demonstrated.

Although the exact role of BCL11b in diseases deserve to be studied by future research, our

results offer potential genome-wide RNA and proteins datasets, which will help future efforts

to understand the molecular basis of cancer and neurodegenerative diseases.

Conflict of interest: no conflict of interest is known

Human or animal experiments: no human or animal work have been performed

Funding:

This work was supported by the French agency for research on AIDS and viral hepatitis

(ANRS); SIDACTION; the European Union's Horizon 2020 research and innovation

programme under grant agreement No 691119-EU4HIVCURE-H2020-MSCA-RISE-2015;

The Belgian Fund for Scientific Research (FRS-FNRS, Belgium), the "Fondation Roi

Baudouin", the Walloon Region (Fonds de Maturation) and the university of Brussels (Action

de Recherche Concertée (ARC) grant). The laboratory of CVL is part of the ULB-Cancer

Research Centre (U-CRC). AAA is a fellow of the Wallonie-Bruxelles International program

and of the Marie Skłodowska Curie COFUND action. HS is a fellow of ANRS. MDR is a

fellow of University of Strasbourg. CVL is Directeur de Recherches of the F.R.S-FNRS,

Belgium.

 Figure Legends

Figure 1. BCL11b binds to multiple complexes and pathways as deciphered by MS results. A)

These proteins were previously shown to interact with BCL11b and we can detect them by IP-

MS method. The gene names of BCL11b-bound ncRNAs are written in blue. B) GO terms of

BCL11b-interacting proteins (BIPs), X-axis show percent of BIPs to the total number of

proteins identified in this category. C) Four representative GO terms with the list of proteins

that are detected to bind to BCL11b. D) The complexes that constitute of BIPs, X-axis show

percent of BIPs to the total number of proteins in the complex; the total number of proteins in

the complex is written in dark blue outside the bars. E) Four representative complexes with the

name of proteins. The BIPs are found in rectangles, BCL11b-interacting ncRNAs are in blue;

whereas the non-BIPs are in hexagonal red shapes. The arrows refer to protein-protein

interactions. F-I) western blot images show co-IP using antibodies directed to anti-BCL11b,

anti-FUS, and anti-Drosha to detect BCL11b, FUS, SMN1, and UPF1. The antibodies (anti-)

used to probe western blots are listed at left of the figure. The figures were constructed from

Supplementary data files S1 and S2.

Figure 2. Representative CLIP-seq results of three protein-coding genes (EWSR1, FUS, and

SRRM2), and non-protein coding gene (NEAT1 ncRNA). The olive green bars refer to the

NCBI RefSeq genes. X-axis represents the coordinates of the gene and the chromosome

number. The Y-axis represents the number of CLIP reads in each condition. a) Over-expressed

BCL11b in HEK cells, b) endogenous BCL11b in microglial cells, whereas, c and d are their

control experiments, respectively. e) The black bars represent the GENCODE transcripts of the

stated genes, the sky blue transcripts refer to the protein non-coding transcripts that harbor

significant CLIP reads and their coordinates correspond to intron of RefSeq gene, these intronic

regions are highlighted with light rose colored boxes. The light blue horizontal boxes refer to

introns without significant BCL11b CLIP reads. For additional figures and CLIP-seq reads

alignments, see Figure S1-S5.

Figure 3. CLIP-seq results of BCL11b-bound RNA. A) Number of BCL11b-interacting

proteins, from IP-MS experiment, that passed our cut-off (so-called MS hits); number of genes,

which have CLIP reads passed our cut-off (so-called CLIP hits); 137 genes can bind to BCL11b

in form of RNA or in form of protein. B) GO terms of the genes that passed CLIP-seq cut-offs.

C) The predicted BCL11b-binding motif. D) GO terms of 137 shared genes (BCL11b binds to

RNA and protein product). E) Schematic diagram of the hypothesis regarding tubulin

autoregulation mechanism. BCL11b binds to multiple tubulin RNA, in brown box, in addition

BCL11b binds to multiple ribosomes (oranges box), tubulins (blue box) and ribonucleases (red

box). F) Violin shape represents the number of CLIP reads mapped to exons and introns of the

significant genes. G) BCL11b binds differentially to the selected 23 genes. Number next to the

names are number of transcripts. H) Numbers of CLIP reads normalized to number of exons

for isoforms coding for proteins (ICPs) or isoforms do not code for proteins (INCPs). The table

shows the Spearman's rank correlation coefficient (R) between numbers of CLIP reads in case

of ICPs or INCPs from one side and length or number of exons from the other side. I) Number

of CLIP reads mapped to exons and those mapped to introns in case of ICPs or INCPs. J)

Number of CLIP reads mapped to exons of DNMT1 gene are counted and normalized to

number of exons. The table shows the average number of CLIP reads and number of exons

within protein-coding transcripts and non-protein-coding transcripts. The blue colors refer to

ICPs, whereas red, green and orange colors refer to nonsense mediated decay, processed

transcript, and retained intron transcripts, respectively. The figures 3G-3J were constructed

from Supplementary data files S10.

Figure 4. A) The table shows the contribution of BCL11b and other genes in the Parkinson's

disease (PD), SMA, ALS and Ewing sarcoma diseases. B) Results of RNA IP-qPCR (RIP-PCR)

of three genes, the fold change of RNA bound to endogenous BCL11b to the input control. For

the location of the amplicons on the gene, please refer to figure S3, and for PCR primers, Table

S4. C) Gene expression of the amplicons used in G using overexpressed BCL11b plasmid and

pCDNA3 vector. For raw Ct values presented in (B) and (C), see Supplementary data S11. D)

The graphical abstract shows the major proteins pathways, complexes and based on GO terms

obtained after IP-MS and CLIP-seq experiments.

References

Ahmed, I., Sbodio, J.I., Harraz, M.M., Tyagi, R., Grima, J.C., Albacarys, L.K., Hubbi, M.E.,
Xu, R., Kim, S., Paul, B.D.*, et al.* (2015). Huntington's disease: Neural dysfunction linked to
inositol polyphosphate multikinase. Proceedings of the National Academy of Sciences *112*,
9751-9756.
Bhat, A.A., Younes, S.N., Raza, S.S., Zarif, L., Nisar, S., Ahmed, I., Mir, R., Kumar, S.,
Sharawat, S.K., Hashem, S.*, et al.* (2020). Role of non-coding RNA networks in leukemia
progression, metastasis and drug resistance. Molecular Cancer *19*, 57.
Briese, M., Saal-Bauernschubert, L., Ji, C., Moradi, M., Ghanawi, H., Uhl, M., Appenzeller,
S., Backofen, R., and Sendtner, M. (2018). hnRNP R and its main interactor, the noncoding
RNA 7SK, coregulate the axonal transcriptome of motoneurons. Proc Natl Acad Sci U S A
115, E2859-E2868.
Chen, C., Ding, X., Akram, N., Xue, S., and Luo, S.-Z. (2019). Fused in Sarcoma: Properties,
Self-Assembly and Correlation with Neurodegenerative Diseases. Molecules *24*, 1622.
Cherrier, T., Le Douce, V., Eilebrecht, S., Riclet, R., Marban, C., Dequiedt, F., Goumon, Y.,
Paillart, J.C., Mericskay, M., Parlakian, A.*, et al.* (2013). CTIP2 is a negative regulator of P-
TEFb. Proc Natl Acad Sci U S A *110*, 12655-12660.
De Santis, R., Santini, L., Colantoni, A., Peruzzi, G., de Turris, V., Alfano, V., Bozzoni, I.,
and Rosa, A. (2017). FUS Mutant Human Motoneurons Display Altered Transcriptome and
microRNA Pathways with Implications for ALS Pathogenesis. Stem Cell Reports *9*, 1450-
1462.
Doktor, T.K., Hua, Y., Andersen, H.S., Brøner, S., Liu, Y.H., Wieckowska, A., Dembic, M.,
Bruun, G.H., Krainer, A.R., and Andresen, B.S. (2016). RNA-sequencing of a mouse-model
of spinal muscular atrophy reveals tissue-wide changes in splicing of U12-dependent introns.
Nucleic Acids Research *45*, 395-416.
Eilebrecht, S., Le Douce, V., Riclet, R., Targat, B., Hallay, H., Van Driessche, B., Schwartz,
C., Robette, G., Van Lint, C., Rohr, O.*, et al.* (2014). HMGA1 recruits CTIP2-repressed P-
TEFb to the HIV-1 and cellular target promoters. Nucleic Acids Res *42*, 4962-4971.
Forouzanfar, F., Ali, S., Wallet, C., De Rovere, M., Ducloy, C., El Mekdad, H., El
Maassarani, M., Ait-Ammar, A., Van Assche, J., Boutant, E.*, et al.* (2019). HIV-1 Vpr
mediates the depletion of the cellular repressor CTIP2 to counteract viral gene silencing. Sci
Rep *9*, 13154.
Fu, W., Yi, S., Qiu, L., Sun, J., Tu, P., and Wang, Y. (2017). BCL11B-Mediated Epigenetic
Repression Is a Crucial Target for Histone Deacetylase Inhibitors in Cutaneous T-Cell
Lymphoma. J Invest Dermatol *137*, 1523-1532.
Gasic, I., and Mitchison, T.J. (2019). Autoregulation and repair in microtubule homeostasis.
Current Opinion in Cell Biology *56*, 80-87.

Kominami, R. (2012). Role of the transcription factor Bcl11b in development and
lymphomagenesis. Proc Jpn Acad Ser B Phys Biol Sci *88*, 72-87.
Kurosaki, T., Popp, M.W., and Maquat, L.E. (2019). Quality and quantity control of gene
expression by nonsense-mediated mRNA decay. Nature Reviews Molecular Cell Biology *20*,
406-420.
Lennon, M.J., Jones, S.P., Lovelace, M.D., Guillemin, G.J., and Brew, B.J. (2016). Bcl11b: A
New Piece to the Complex Puzzle of Amyotrophic Lateral Sclerosis Neuropathogenesis?
Neurotoxicity Research *29*, 201-207.
Lennon, M.J., Jones, S.P., Lovelace, M.D., Guillemin, G.J., and Brew, B.J. (2017). Bcl11b—
A Critical Neurodevelopmental Transcription Factor—Roles in Health and Disease. Frontiers
in Cellular Neuroscience *11*.
Longabaugh, W.J.R., Zeng, W., Zhang, J.A., Hosokawa, H., Jansen, C.S., Li, L., Romero-
Wolf, M., Liu, P., Kueh, H.Y., Mortazavi, A.*, et al.* (2017). Bcl11b and combinatorial
resolution of cell fate in the T-cell gene regulatory network. Proceedings of the National
Academy of Sciences *114*, 5800-5807.
Maeda, M., Harris, A.W., Kingham, B.F., Lumpkin, C.J., Opdenaker, L.M., McCahan, S.M.,
Wang, W., and Butchbach, M.E.R. (2014). Transcriptome profiling of spinal muscular
atrophy motor neurons derived from mouse embryonic stem cells. PLoS One *9*, e106818.
Marban, C., Suzanne, S., Dequiedt, F., de Walque, S., Redel, L., Van Lint, C., Aunis, D., and
Rohr, O. (2007). Recruitment of chromatin-modifying enzymes by CTIP2 promotes HIV-1
transcriptional silencing. EMBO J *26*, 412-423.
Matera, A.G., and Wang, Z. (2014). A day in the life of the spliceosome. Nature Reviews
Molecular Cell Biology *15*, 108-121.
Orth, M.F., Hölting, T.L.B., Dallmayer, M., Wehweck, F.S., Paul, T., Musa, J., Baldauf,
M.C., Surdez, D., Delattre, O., Knott, M.M.L.*, et al.* (2020). High Specificity of BCL11B and
GLG1 for EWSR1-FLI1 and EWSR1-ERG Positive Ewing Sarcoma. Cancers *12*, 644.
Pong, S.K., and Gullerova, M. (2018). Noncanonical functions of microRNA pathway
enzymes – Drosha, DGCR8, Dicer and Ago proteins. FEBS Letters *592*, 2973-2986.
Rohr, O., Lecestre, D., Chasserot-Golaz, S., Marban, C., Avram, D., Aunis, D., Leid, M., and
Schaeffer, E. (2003). Recruitment of Tat to heterochromatin protein HP1 via interaction with
CTIP2 inhibits human immunodeficiency virus type 1 replication in microglial cells. J Virol
77, 5415-5427.
Scherzer, C.R., Eklund, A.C., Morse, L.J., Liao, Z., Locascio, J.J., Fefer, D., Schwarzschild,
M.A., Schlossmacher, M.G., Hauser, M.A., Vance, J.M.*, et al.* (2007). Molecular markers of
early Parkinson's disease based on gene expression in blood. Proceedings of the National
Academy of Sciences *104*, 955-960.
Shadrina, O., Garanina, I., Korolev, S., Zatsepin, T., Van Assche, J., Daouad, F., Wallet, C.,
Rohr, O., and Gottikh, M. (2020). Analysis of RNA binding properties of human Ku protein
reveals its interactions with 7SK snRNA and protein components of 7SK snRNP complex.
Biochimie *171-172*, 110-123.
Shehadeh, L.A., Yu, K., Wang, L., Guevara, A., Singer, C., Vance, J., and Papapetropoulos,
S. (2010). SRRM2, a potential blood biomarker revealing high alternative splicing in
Parkinson's disease. PLoS One *5*, e9104.
Shkreta, L., and Chabot, B. (2015). The RNA Splicing Response to DNA Damage.
Biomolecules *5*, 2935-2977.
Sun, J., Shi, Y., and Yildirim, E. (2019). The Nuclear Pore Complex in Cell Type-Specific
Chromatin Structure and Gene Regulation. Trends in Genetics *35*, 579-588.
Tuck, A.C., Rankova, A., Arpat, A.B., Liechti, L.A., Hess, D., Iesmantavicius, V., Castelo-
Szekely, V., Gatfield, D., and Bühler, M. (2020). Mammalian RNA Decay Pathways Are
Highly Specialized and Widely Linked to Translation. Molecular Cell *77*, 1222-1236.e1213.
Wakabayashi, Y., Watanabe, H., Inoue, J., Takeda, N., Sakata, J., Mishima, Y., Hitomi, J.,
Yamamoto, T., Utsuyama, M., Niwa, O.*, et al.* (2003). Bcl11b is required for differentiation
and survival of alphabeta T lymphocytes. Nat Immunol *4*, 533-539.
Wang, E., and Aifantis, I. (2020). RNA Splicing and Cancer. Trends in Cancer.

Wiles, E.T., Lui-Sargent, B., Bell, R., and Lessnick, S.L. (2013). BCL11B Is Up-Regulated
by EWS/FLI and Contributes to the Transformed Phenotype in Ewing Sarcoma. PLOS ONE
8, e59369.
Wu, Y.-Y., and Kuo, H.-C. (2020). Functional roles and networks of non-coding RNAs in the
pathogenesis of neurodegenerative diseases. Journal of Biomedical Science *27*, 49.
Yamazaki, T., Chen, S., Yu, Y., Yan, B., Haertlein, Tyler C., Carrasco, Monica A., Tapia,
Juan C., Zhai, B., Das, R., Lalancette-Hebert, M.*, et al.* (2012). FUS-SMN Protein
Interactions Link the Motor Neuron Diseases ALS and SMA. Cell Reports *2*, 799-806.

A)

	BCL11b interacts with	PD	SMA	ALS	Ewing's sarcoma
BCL11b		Down-regulated	Down-regulated		Up-regulated
SRRM2	RNA	Up-regulated	-	-	-
SMN1	Protein	-	Down-regulated	-	-
NEAT1	RNA	-	Up-regulated	-	-
EWSR1	RNA and protein	-	-	Contribute in	Contribute in
FUS	RNA and protein	-	-	Mutated	-
RNPs	RNA, protein & ncRNA	-	Defects	Defects	-
TDP43	RNA and protein	-	-	Mutated	-
TAF15	RNA and protein	-	-	Contribute in	-
VCP	RNA	-	-	Contribute in	-

B)

C)

D)