Temporal and trophic partitioning promote coexistence between mesocarnivores in a Mediterranean landscape

Iago Ferreiro-Arias[1]*, Jorge Isla[1], Pedro Jordano[1], Ana Benítez-López[1]

[1]Department of Integrative Ecology, Estación Biológica de Doñana, Consejo Superior de Investigaciones Científicas, Avda. Américo Vespucio, 26, 41092, Sevilla, Spain.

*Corresponding author. email: iago.ferreiro.arias@gmail.com

Iago Ferreiro-Arias - ORCID: 0000-0003-4178-5783

Pedro Jordano - ORCID: 0000-0003-2142-9116

Jorge Isla - ORCID: 0000-0002-2307-9730

Ana Benítez-López - ORCID: 0000-0002-6432-1837

Abstract

1. The partition of the ecological niche can enhance the coexistence of predators due to differences in how they exploit three main resources: food, space, and time, the latter being an axis that often remains unexplored.

2. We studied niche segregation in a Mediterranean mesocarnivore community in sites where the top predator, the Iberian lynx, is absent, addressing simultaneously two niche axes: the temporal and the trophic axes.

3. Temporal overlap between prey and predator and between potential competitors was analysed by comparing daily activity patterns of predators and prey species through data obtained by using 24 camera-traps deployed in three different sites of the Doñana National Park. We further examined the trophic dimension by compiling and analysing diet information in the study area and exploring trophic preferences between potential competitors.

4. We found an overall temporal segregation between trophic generalist species while those species which showed higher values of temporal overlap differ largely in their trophic preferences. Furthermore, we observed an overall high degree of overlap between the activity patterns of predators and their putative prey (the common genet vs small mammals and the red fox vs European rabbit).

5. Our study suggests that coexistence of the different species that compose the mesocarnivore assemblage in Mediterranean landscapes can be facilitated by the marked temporal segregation between generalist species, but also by differences in trophic preferences by specialist species when their activity patterns overlap with other

23   mesocarnivores. Our findings reinforce that idea that the coexistence mechanisms underlying

24   community structure are multidimensional.

Keywords

26   Diel activity, Doñana, *Genetta genetta, Herpestes ichneumon,* interspecific competition,

27   *Meles meles,* niche differentiation, *Vulpes vulpes.*

Introduction

29   Understanding the mechanisms that promote coexistence of species with similar

ecological requirements is a central topic in community ecology, where competitive

interactions between species have the potential to affect diversity patterns by limiting or

promoting coexistence (Chesson, 2000). In order to mitigate the negative impact of

interspecific competition, species often partition resources along three main niche dimensions

(temporal, trophic and spatial), which eventually result in niche differentiation (Schoener,

1974). Most studies concerning species coexistence focus on the differential use of habitat

and food resources; nonetheless, the differential use of the diel cycle may enhance

coexistence of same-sized species, particularly among predator species (Di Bitetti et. al.,

2009; Wang & Fisher, 2012).

39   Diel activity patterns are cyclical behavioural patterns which can be influenced by

several factors (i.e., prey availability or thermoregulatory strategies) (Lode, 1995; Maloney

et. al., 2005). Within mammals, a large variability and plasticity in their activity patterns has

been documented (Bennie et al. 2014). This diversity of diel cycles could be due to the plastic

nature of behavioural responses to different pressures, which may in turn induce marked

variations in daily rhythms among different scenarios (Ensing et al., 2014; Gaynor et. al.,

2018). Broadly, activity patterns displayed by mammals are determined by circadian

endogenous boundaries (Kronfeld-Schor & Dayan, 2003), which, particularly in carnivores,

are shaped by external factors (Monterroso et.al., 2014; Zielinski, 1986), and thus their diel

cycle use is context-dependent. Hence, the co-occurrence of species that compete for a certain

resource has the potential to produce shifts in the daily activity patterns of competitively

surrogate species. For example, crab-eating and pampas foxes have been reported to change

their daily foraging activity in sympatric areas where both species are present (switching

from diurnal to nocturnal), compared to allopatric areas where only one fox species is present

(Di Bitetti et al., 2009).

54   In addition, predators' activity patterns could also adjust their daily activity rhythms to

those hours of the day in which the availability of prey increases (Foster et al., 2013), while,

at the same time, prey species could generate a temporal mismatch by centering their activity

at times when there is a lower risk of predation (Fenn & Macdonald, 1995). Consequently,

high temporal overlap between carnivores and their putative prey have been reported in some

predator-prey systems (Linkie & Ridout, 2011), whereas in others an asynchrony was found

in their respective activity peaks (Arias-Del Razo et al., 2011; Díaz⬜Ruiz et al., 2016). As a

consequence, the daily activity of a given predator could be a result of a cost-benefit trade-off

between maximizing their activity during periods with high prey availability (benefit) but

also higher mortality risk due to encounters with intraguild competitors (cost) (Santos et al.,

2019).

65   Interspecific interactions play an important role in structuring a mammalian

community (Holt et al., 1994), especially in carnivores where competition not only arises as a

consequence of exploiting the same resource simultaneously, but also due to the risk

associated with intraguild predation. Intraguild predation (IGP) is an antagonistic interaction

where a predator species kills another predator, or where both species prey upon each other

(Palomares & Caro, 1999; Polis et.al., 1989), being specially reported on top-predators

against surrogate competitors (Palomares & Caro, 1999). The IGP's effects in competitors

can be diverse, from abundance reductions and distributional shifts (in space and time)

(Jiménez et al., 2019; Newsome et al., 2017) to behavioural modifications through restriction

of their activity patterns (Wang & Fisher, 2012). Because apex predators play a key role in

ecosystem functioning due to their regulatory role on populations of prey and medium-sized

carnivores (top-down control), their extinction frequently triggers cascading effects through

direct and indirect interactions among trophic levels (Terborgh et. al., 2010). Hence, the

collapse of top predators can lead to an eventual increase of the distribution and abundance of

medium-sized carnivore populations, a phenomenon coined '*mesopredator release*' by Soulé

et al. (1988) that has been repeatedly documented (Prugh et al., 2009; Brashares et. al., 2010).

In the Iberian Peninsula, the Iberian lynx (*Lynx pardinus*) exerts its role as a top-

predator on the populations of surrogate competitors (Palomares et. al., 1998; Palomares

et.al., 1996). Sizeable reductions of mesocarnivore abundance have been reported after the

reintroduction of the Iberian lynx (Jiménez et al., 2019), reinforcing the idea that the

coexistence between these carnivores is supported by spatial structuring, where the

probability of occurrence of the lynx is negatively associated with the presence of

mesopredators (Monterroso et.al., 2020; Palomares et al., 1996). In Doñana National Park,

the carnivore guild is composed by a habitat and trophic specialist top predator (the Iberian

lynx, *Lynx pardinus*) and mesocarnivores such as the red fox (*Vulpes vulpes*), the European

badger (*Meles meles*) and the Egyptian mongoose (*Herpestes ichneumon*), which are

characterized by a wide niche breadth in terms of habitat and trophic preferences. This

assemblage is completed by the common genet (*Genetta genetta*), with a considerably

narrower diet. The negative effects that the Iberian lynx has on these mesocarnivores occur

mainly through intraguild predation, particularly on Egyptian mongooses (*Egyptian

mongoose), common genets (*Genetta genetta)* and red foxes (*Vulpes vulpes*) (Palomares et al.

1996, Palomares & Caro 1999, and references therein).

However, IGP interactions between the different mesocarnivores remain quite

unknown (Palomares et al., 1996; Valverde, 1957). In areas where the Iberian lynx is not

present, coexistence between mesopredators does not seem to be mediated by spatial

structuring (Soto & Palomares, 2015) and thus indirect competitive interactions may become

more relevant. In this sense, differences in activity shifts of mesocarnivores with similar

trophic requirements would alleviate resource overlap. Empirical data in this respect are

surprisingly very limited for the Mediterranean area, where situations of mesopredator release

may be common due to recent historical reductions in the Iberian lynx populations. Yet, most

previous studies have focused on spatial avoidance as the main mechanism allowing the

mesocarnivore coexistence, and only recently has the focus shifted towards temporal

segregation (Barrull et al., 2014; Monterroso et al., 2014; Vilella et al., 2020).

Hence, the main purpose of this study is to evaluate the ecological factors that allow

the coexistence of mesocarnivore species in the Doñana National Park without the influence

of a top-predator, focusing on the temporal segregation of four mesocarnivore species: the red

fox (*Vulpes vulpes*), the European badger (*Meles meles*), the common genet (*Genetta

genetta*), and the Egyptian mongoose (*Herpestes ichneumon)*. Thus, this study aims: 1) to

explore the relationship between mesocarnivores along the trophic niche axis; 2) to determine

the diel activity patterns of predator and prey species and their synchrony; and 3) to quantify

the temporal overlap or segregation between potential competitors. Following a niche

partitioning conceptual framework, we hypothesize that there would be greater temporal

segregation in those species that have a higher degree of overlap in their respective trophic

niches, while species that have similar use of the diel cycle would show differentiated trophic

preferences in order to promote coexistence by reducing competition along temporal or

trophic niche axes.

Material and methods

Study area

This study was carried out in Doñana National Park (37º1'N, 6º 33'W) in three 1-ha

sites characterized by juniper stands (*Juniperus phoenicea* subsp. *turbinata)* of different age

and ecological succession stages, ranging from mature stands with a higher degree of

vegetation cover to an advance front of juniper stands with lower vegetation cover. "*Sabinar*

del Marqués" (SABMAR) constitutes the most mature stage with 10520 junipers/ha,

"*Sabinar del Ojillo*" (SABOJI) is at an intermediate stage with 9010 junipers/ha, and

"*Sabinar de Colonización*" (SABCOL) represents the colonization front stage with 2030

junipers/ha (Supplementary Fig. S1).

Data collection

We deployed eight camera traps per site that were rotated periodically through 35

randomly selected focal juniper plants. Juniper plants were selected at random in each of five

subplots within each of the three main sites, with the subplots distributed regularly

throughout the main plot. The structure of this habitat consists of paths of variable width

between juniper individuals, where the cameras were installed at 1-m height focusing on the

ground and the bottom of the plant. Camera trap models employed in this study were

Browning Dark Ops Pro XD and Bushnell Aggressor. Camera traps were active for ca. 10

days in each of these rotations and then changed to other 8 locations, thus ensuring that the

whole site was sampled in 5 weeks. Following this procedure, we obtained data from a total

set of 24 camera traps sampling the three sites during two campaigns: October 2018 - May

2019 and October 2019 - June 2020. No attractants were used over the two years of sampling.

The diel activity cycle of each species was characterized by pooling the total number

of detections across all cameras during the whole study period. In order to avoid data

dependency, we discarded consecutive detections of a given species within a site (Davis,

Kelly, & Stauffer, 2011; Monterroso et al., 2014). When multiple photographs of the same

species were taken within a 15-min interval, we considered them as a single capture event to

ensure capture independence. Since the area covered by our study sites is limited (ca. 1

ha/site), we deem 15 min a reasonable time period for a species to abandon the site. In cases

where a given individual was detected repeatedly without leaving the camera detection zone,

only the time of its first detection was considered. In our analyses we used either the total,

absolute number of records per species (e.g., to compare the relative frequency of sunrise,

day, dusk, and nocturnal records within a given site) or the relative frequency of records

expressed as number of photographs per species per camera-day (e.g., when comparing

across sites that received different sampling intensity).

Our target species consisted of the four mesopredators which occur in Doñana

National Park namely the Egyptian mongoose, the red fox, the European badger and the

common genet (Valverde, 1967), and their respective prey species. Large top predators such

as the wolf (*Canis lupus*) and the Iberian lynx (*Lynx pardinus*) were not present in our study

sites. Potential prey species include lagomorphs (mainly European rabbits, *Oryctolagus*

cuniculus), small mammals and birds such as the red partridge *Alectoris rufa*. Detections

from ungulates (mainly *Cervus elaphus* and *Sus scrofa*) were not included in the analysis as

their presence in mesocarnivore diets is frequently associated with carrion consumption, with

direct predation being recorded sporadically on very young individuals (Valverde, 1967).

Conversely, we did not have sufficient detections from the other lagomorph present in RBD

(*Lepus granatensis*, N = 3) to include them in this study.

Trophic overlap data

We searched on Web of Science (WOS) and Google Scholar for studies that reported

trophic preferences of the four mesocarnivores in Doñana National Park using the following

search string: (trophic OR diet) AND (predator species) AND (Doñana), where "predator

species" was substituted by one of the mesopredator species considered in this study when

performing the search. We also gathered diet data for the common genet by consulting

experts (P. Ferreras). The selection of studies contains all the trophic information available

for the four species at different locations in the Doñana National Park and reflects the entire

trophic spectra of each of the different species (Supplementary Table S1). We collected all

food items per species and study without pooling seasons or periods in order to capture the

natural variability in dietary composition across the whole Doñana study area and its marked

seasonal component.

Some studies reported information collected in previous studies (e.g., Martín et al.

(1995) and Fedriani et al. (1998), or Palomares and Delibes (1991b) and Palomares (1993)).

To avoid pseudo-replication issues, we kept only those studies where the data strictly belong

to different samples and periods. From each study, period and species, we extracted the

frequency of occurrence of food items in mesopredator faeces. In order to estimate trophic

overlap between mesocarnivores, we carried out a correspondence analysis (CA) using the

frequency of occurrence of each prey item for each predator. The CA ordered the different

diet studies (cases) relative to the prey species composition reported by each study, resulting

in an ordination of the studies corresponding to each mesopredator species based on their

similarity of prey species composition. We used the coordinates on the first two CA axes

corresponding to each diet study to build convex polygons spanning each mesopredator group

of studies. Convex polygons illustrate the range of dietary variation shown by each

mesopredator species in relation to the diet composition of other species.

Camera trap data analysis

We extracted time and date from each detection of every camera trap in order to

estimate daily activity patterns and temporal overlap between mesocarnivores and their

potential prey, and between mesocarnivore potential competitors. As time and dates are

variables with a circular nature, diel activity cycles were estimated using non-parametric

Kernel density plots, which provide a density function of the daily activity pattern of a given

species (Rowcliffe et al., 2014). Raw data were subsampled in order to carry out analyses

only for those species with more than 10 detections (Fisher, 1995).

To determine the daily activity pattern of the species we resorted to the classification

used by van Schaik and Griffiths (1996) by which species are classified according to the

number of independent detections obtained as: diurnal (> 90% of the detections occurring

during the daylight), nocturnal (> 90% of the detections obtained at night), crepuscular (>

90% of their detections occur at dawn or dusk) or cathemeral if their activity pattern is

distributed uniformly throughout the daily cycle. We tested whether the target species

actually show a certain activity pattern and deviate from a random pattern using the Rao

Spacing Test (U) (Rao, 1976).

As the present study has been carried out over two years, daylight can vary from one

detection to another and induce bias to our results. Species could adjust their daily activity

levels depending on the day length, thus not taking this aspect into account when dealing with

strictly diurnal or nocturnal species could lead to the misinterpretation of the results and the

underestimation of peaks of daily activity (Vazquez et al., 2019). Therefore, we carried out a

double average anchoring time transformation to take into account the mean average sunrise

and sunset times for the study area (Vazquez et al., 2019).

We evaluated the extent of temporal segregation between mesopredators using the

overlap coefficient (Δ) described by Ridout and Linkie (2009). This coefficient ranges from 0

to 1, where 0 values represent completely different diel activity patterns while 1 represents

the maximum overlap between both species. Following Ridout and Linkie (2009), we used Δ_1

for cases with a sample size lower than 75 detections, and Δ_4 when sample size was equal or

greater than 75 detections.

Once the overlap coefficients between the different species were estimated, 99%

confidence intervals were calculated through smoothed bootstrap analysis with 10,000

replicates. Moreover, we evaluated differences in pairwise comparisons of daily activity

patterns using the non-parametric Mardia-Watson-Wheeler test (MWW, Batschelet, 1981).

Further, we used percentiles to establish threshold levels of overlap. Thus, $\Delta > $ 75th percentile

indicates a high temporal overlap between both species, $\Delta \leq$ 50th percentile a low degree of

overlap and, finally, intermediate values of these percentiles, 50th $< \Delta \leq$ 75th, denote

moderate levels of overlap (Monterroso et al. 2014).

All analyses were conducted in R Studio 1.2.5033 Statistical Software (R Core Team,

2020). The correspondence analysis (CA) was carried out with "*FactoMineR*" package (Lê,

Josse, & Husson, 2008); kernel density functions, overlap coefficients (Δ) and their

respective confidence intervals were calculated with the package "*overlap*" (Ridout & Linkie,

2009); MWW and Rao Spacing tests were conducted using the "*circular*" package

(Agostinelli & Lund, 2017). Finally, we calculated mean average sunrise and sunset times

and performed the average anchoring transformation with the "*activity*" package (Rowcliffe,

2019).

Results

Trophic segregation among mesocarnivores

The first two dimensions of the correspondence analysis explained 62% of the

variability of mesocarnivore diets (Fig. 1). Axis 1, which explained 34.5% of the variation,

clearly discriminates the trophic preferences of the European badger and the common genet

due to their eminently omnivorous and carnivorous diets, respectively, but not from the rest

of the mesocarnivores. The red fox and the European badger showed very close values due to

their frugivorous behaviour and the predominant consumption of lagomorphs and

invertebrates. Meanwhile, the Egyptian mongoose had the wider trophic preferences, with an

intermediate position between the omnivorous diets of the European badger and the red fox,

and the carnivorous diet of the common genet. Simultaneously, axis 2, which explained

27.3% of the trophic variation, is generated by variation between the diets of the red fox and

the common genet, with the latter preferring small mammals followed by birds as main food

items. However, the second dimension of the correspondence analysis does not discern

between the trophic niches of the red fox, the European badger and the Egyptian mongoose,

indicating that there is a greater trophic overlap between these three species with respect to

the common genet. Further, convex polygons indicate the ampler trophic diversification of

the generalist species (red fox, European badger and Egyptian mongoose) whose range of

prey types tends to be considerably wider compared with the common genet (Fig. 1).

Species detections and diel activity patterns

Results obtained in this study come from a total of 531 effective sampling days. All

sites were sampled with the same number of cameras per site and a similar sampling effort

(number of days): $\bar{x} \pm SD = 80.3 \pm 28.3$ d for SABCOL; 80.4 ± 27.3 d for SABOJI and

86.8 ± 26.0 d for SABMAR. Broadly, 2786 detections in total were obtained throughout the

study period, of which 96 (3.44%) could not be identified at the species level. Among all

detections, 16% corresponded to mesocarnivores, 28.7% to prey species, 15% to ungulates

and 38.7% to passeriformes, which were not included in this study (Supplementary Table

S2).

Regarding our target species, we recorded 301 detections of the red fox, with 54.8%

of them in SABCOL and the rest of the detections evenly distributed between the other two

sites (SABOJI: 23.9%; SABMAR: 21,3%). A large amount of the detections obtained from

this species were linked to its nocturnal habits while the rest were distributed throughout

daylight periods, demonstrating a cathemeral activity pattern with a tendency towards

bimodality (U = 142.26; p-value < 0.05) (Supplementary Table S3). Conversely, detections of

European badgers were very scarce in the three study sites (total: 22 detections), with 77% of

the detections occurring at night and the rest during the dawn or dusk periods (U = 178.95; p-

value < 0.01). As for the common genet, its activity pattern was strictly nocturnal (95% of the

records produced during nighttime; U = 165.81; p-value < 0.01), with most detections

occurring in SABCOL (62% of detections). Finally, detections of Egyptian mongooses were

mainly concentrated in the SABOJI (61%) while the other two sites maintained the same

proportion of records (ca. 20%), showing a marked diurnal activity pattern in each of them

(92% of the detections occurred during daylight; U = 199.78; p-value < 0.01) (Supplementary

Table S3).

For prey species, most of the detections belonged to the European rabbit (N = 650

detections). Rabbit detections concentrated similarly between SABOJI and SABMAR, while

SABCOL had only 6 detections during the entire sampling period (Supplementary Table S2).

Overall, the distribution of rabbit detections spanned throughout the day, with distinct activity

peaks around sunrise and sunset (U = 156.97; p-value < 0.01) (Supplementary Table S3).

Small mammals (*Mus spretus* and *Apodemus sylvaticus*) were detected 98 times,

mainly in SABMAR and only during nighttime (U = 198.27; p-value < 0.01). Finally, red-

legged partridges were mostly detected in SABCOL (89% of detections) and showed a

markedly diurnal activity pattern (U = 226.53; p-value < 0.01) (Supplementary Table S3).

Predator and prey diel activity patterns

Broadly, the activity patterns and their corresponding overlap differed between

predators and the potential prey items in the study area. The European rabbit showed a high

degree of overlap with the red fox ($\Delta_4 = 0.84$), followed by a moderate overlap with the

European badger ($\Delta_1 = 0.62$). Conversely, both the common genet and the Egyptian

mongoose showed low overlap coefficients ($\Delta_1 = 0.47$ and $\Delta_1 = 0.44$, respectively) with rabbit

activity patterns (Fi. 2).

We found a high temporal overlap between the activity patterns of small mammals

(with most detections at night) and the European badger ($\Delta_1 = 0.89$) and the common genet

($\Delta_1 = 0.76$), respectively. In turn, the red fox showed a moderate temporal overlap with small

mammals ($\Delta_4 = 0.68$), while the Egyptian mongoose showed a total temporal segregation

with respect to the detected micromammals ($\Delta_1 = 0.14$). Activity patterns of red-legged

partridges overlapped with those of the Egyptian mongoose ($\Delta_1 = 0.82$), but not with those of

European badgers ($\Delta_1 = 0.08$) and common genets ($\Delta_1 = 0.03$) (Fig. 2).

European rabbit activity patterns differed significantly between SABOJI and

SABMAR (W = 32.69; p-value < 0.01) (Fig. 3). Meanwhile, red fox, but not the Egyptian

mongoose, showed different responses in terms of diel cycle among sites with different rabbit

availability. The red fox showed a predominantly nocturnal activity in SABCOL, where

rabbits were absent, and a tendency towards bimodal activity patterns in SABOJI and

SABMAR. Red fox activity patterns differed significantly between SABCOL and SABMAR

(W = 11.14; p-value < 0.01), while no differences were found between SABCOL - SABOJI

(W = 3.97; p-value = 0.137) or SABOJI - SABMAR (W = 2.44, p-value = 0.295). In turn, no

statistically significant differences were found between activity patterns of the Egyptian

mongoose between SABCOL - SABOJI (W = 1.09; p-value = 0.58), SABCOL - SABMAR

(W = 2.06; p-value = 0.36) and SABOJI - SABMAR (W = 0.40; p-value = 0.82) (Fig. 3). The

temporal overlap between the activity patterns of the European rabbit and the red fox was Δ_1

= 0.79 for SABOJI and Δ_1 = 0.73 for SABMAR, while the temporal overlap between

European rabbit and Egyptian mongoose was sensibly lower (Δ_1 = 0.49 for SABOJI and Δ_1 =

0.37 for SABMAR) (Fig. 3).

Temporal overlap and segregation between mesocarnivores

The average overlap and standard deviation of the activity patterns of the different

species was Δ_1 = 0.48 and SD = 0.29, which reflect a low, although variable, temporal

overlap among these potential competitors. The average overlap values ($\Delta_1 \pm$ SD) of a species

with respect to the set of potential competitors were: 0.59 ± 0.15 for the fox; 0.57 ± 0.34 for

the badger; 0.50 ± 0.36 for the genet and 0.24 ± 0.18 for the Egyptian mongoose.

Strictly nocturnal species (European badger and common genet) showed a high

overlap coefficient and no significant differences between their activity patterns (Δ_1 = 0.79;

W = 1.89, p-value = 0.386). In contrast, the Egyptian mongoose showed a strong and

significant temporal segregation with the European badger (Δ_1 = 0.16; W = 41.75, p-value <

0.01) and the common genet (Δ_1 = 0.08; W = 65.79, p-value < 0.01) (Fig. 4). Simultaneously,

we found that the activity pattern of the red fox had intermediate levels of temporal

segregation with the common genet ($\Delta_1 = 0.57$; $W = 24.96$; p-value < 0.01) and the European

badger ($\Delta_1 = 0.71$; $W = 8.44$; p-value $= 0.014$) and a low overlap with the Egyptian

mongoose ($\Delta_1 = 0.44$; $W = 73.48$; p-value < 0.01) (Fig. 4). Among study sites, temporal

overlap between the red fox and the Egyptian mongoose was $\Delta_1 = 0.41$ for SABCOL, $\Delta_1 =$

0.45 for SABOJI and $\Delta_1 = 0.36$ for SABMAR (Fig. 3).

Discussion

In this study we explore the mechanisms underlying the structure of Mediterranean

mesocarnivore communities living in sympatry in a mesopredator release scenario. We show

that mesocarnivore coexistence could be mediated by temporal or trophic segregation when

there is a greater overlap in any of these two niche axes. Unlike previous studies which have

been traditionally focused on spatial responses (Fedriani et al. 1999, Soto and Palomares

2015, Monterroso et al. 2020, but see Barrull et al. 2014, Monterroso et al. 2014, Vilella et al.

2020), our study provides new insights on the importance of temporal segregation at

facilitating co-occurrence by reducing resource overlap. Overall, we found a moderate level

of temporal segregation within the mesocarnivore guild in our study area, which was

composed by two strictly nocturnal species - the European badger and the common genet

(Camps, 2008; Palomares & Delibes, 1994; Rodríguez et al., 1996), one diurnal - the

Egyptian mongoose (Palomares & Delibes, 1992) and one with plasticity in its daily activity

pattern - the red fox, which ranges from diurnal (Cavallini & Lovari, 1994) to nocturnal

habits (Baker et al., 2007).

The highest values of temporal overlap occurred between the European badger and the

common genet due to their predominantly nocturnal activity patterns. However, they showed

marked differences in their respective trophic niches, mostly because of the preferential

consumption of small mammals by the common genet. The common genet consumes almost

all species of rodents and insectivores, but it is certainly a specialized predator on *Apodemus*

sylvaticus when this prey is available (Palomares & Delibes, 1991a; Virgós et al., 1999).

According to the Optimal Foraging Theory (MacArthur & Pianka, 1966), predators maximize

its prey intake by showing an efficient foraging pattern. In the case of predators with

specialized diets such as the genet, they optimize their foraging behaviour by maximizing

synchrony of their daily activity patterns to their main and most profitable prey (Pyke et al.,

1977), as shown by our results. However, this pattern would not be so marked in predators

whose feeding behaviour is opportunistic and depends on the availability and accessibility of

several prey types (Pyke et al., 1977), which is the case for the rest of the mesocarnivores

studied.

The three other species that constitute the Doñana mesocarnivore assemblage are

known for their generalist feeding habits (Fedriani et al., 1999; Palomares, 1993; Rau, 1987).

Yet, we found trophic segregation of the Egyptian mongoose with respect to the red fox and

the European badger, with plant material (*e.g.*, fleshy fruits) virtually absent from its diet.

Meanwhile, the trophic overlap between European badger and red fox was the highest within

the mesocarnivore community due to their omnivorous behaviour based on the consumption

of fleshy fruits, invertebrates and lagomorphs. Despite their reduced differences in terms of

spatial and trophic niche, these species showed a marked temporal segregation, displaying

distinct activity patterns: nocturnal, cathemeral and diurnal, which could ease coexistence by

exploiting similar resources at different times and thus reducing intraguild antagonistic

encounters.

The ecological plasticity and the ample spectra of resource use shown by these species

in their trophic and habitat niche axes suggest a scenario with potentially reduced

interspecific competition (Fedriani et al., 1999; Monterroso et al., 2020; Palomares, 1993).

However, the temporal segregation of these species may become more relevant in situations

where those two niche axes exhibit ample overlap. For example, a reduction in the abundance

and availability of its main trophic resources at a given moment may lead to increasing

competition for the consumption of available secondary resources. A clear example is the

reported reduction in the consumption of European rabbits after a sudden population collapse

due to the emergence of the rabbit hemorrhagic disease (RHD) (Ferreras et. al., 2011), which

resulted in an increased trophic overlap among most generalist carnivore species. In such

cases, time partitioning becomes the main mechanism to facilitate coexistence by reducing

competitive stress in resource use among these species (Barrientos & Virgós, 2006).

The European rabbit in the study area accounts for a very high percentage of the diet

of Egyptian mongooses (Palomares & Delibes, 1991b) and also for European badgers. Some

authors even point out a possible specialization of badgers for this type of prey, a contentious

assertion which has been debated (Martín et al., 1995; Revilla & Palomares, 2002). The

activity pattern of European rabbits matches those previously reported, showing a bimodal

pattern with peaks of activity at sunrise and sunset (Díaz☐Ruiz et al., 2016; Monterroso et

al., 2013; Villafuerte et al., 1993). However, rabbit availability differed between study sites,

with almost no detections in the least mature, early successional juniper colonization front

site (SABCOL), and frequent detections in the other more mature sites (SABMAR and

SABOJI). Such differences may be due to its habitat requirements since, at the landscape

scale, the ideal habitat for rabbits is a juxtaposition of ecological units where the refuge areas

are intermingled with others where herbaceous food is more abundant (Dellafiore et al., 2014;

Monzón et al., 2004). The colonization site (SABCOL) represents a Mediterranean scrubland

in an early ecological stage composed mainly by a low-density population of *J. phoenicea*

(2030 junipers/ha) and therefore, a low vegetation cover which does not fulfil rabbit habitat

preferences. Meanwhile, SABMAR and SABOJI are sites characterized by late successional

mature vegetation and high vegetation cover, which may serve as refuge for rabbits. In these

two sites, we found differences between the diel cycle use of rabbits, showing two activity

peaks in SABOJI and only one crepuscular activity peak in SABMAR.

Interestingly enough, the activity pattern displayed by the red fox, showed a unimodal

pattern with an activity peak around midnight in SABCOL, but shifted towards bimodality in

sites where the rabbit was present, putatively to match the activity patterns of this prey and

thus increase the probability of encounter. This was particularly obvious in SABOJI, where

the temporal overlap between fox activity and that of the rabbits was the highest. Yet, this

strategy was less beneficial in SABMAR, where an increase in the red fox activity around

dawn may have triggered an anti-predatory behaviour in the European rabbit, which

displayed only one activity peak at twilight, thus leading to low temporal overlap. This

observation is in line with the Risk Allocation Hypothesis by which a temporal mismatch

between activity patterns of a prey species and its predator could be due to a higher

perception of mortality risk that drives anti-predatory behaviours along the time axis (Lima &

Bednekoff, 1999). Previous studies with Black-tailed jackrabbit (*Lepus californicus*), desert

cottontail (*Sylvilagus audubonii*) and the coyote (*Canis latrans*), have shown that lagomorphs

are able to shift their activity in order to reduce predation risk by the coyote (Arias-Del Razo

et al., 2011).

Dissimilarities in the plasticity of the activity patterns of the red fox and the Egyptian

mongoose may be explained by their different foraging strategies. Although, the European

rabbit is a profitable prey for the generalist mesopredators in Doñana (Ferreras et al., 2011;

Martín et al., 1995; Palomares & Delibes, 1991b; Fedriani, 1996), each predator displays

different hunting and consumption strategies of this prey item. Egyptian mongooses and

European badgers are species that preferentially consume younger rabbits by digging in

burrows (Martín et al., 1995; Palomares & Delibes, 1991b). Conversely, the red fox resorts to

a foraging strategy based mainly on the pursuit of prey items, although it also excavates in

rabbit burrows (Fedriani et al., 1999). Thus, these distinct foraging strategies could explain

why the Egyptian mongoose and European badgers display temporal asynchrony with the

prey's activity patterns while the red fox is expected to track and match the activity pattern of

its prey.

Hence, our study addresses for the first time the interspecific relationships among the

four mesocarnivores species present in Doñana National Park by integrating together the

different patterns of daily activity and their respective overlap, as well their trophic ecologies.

Yet, these results should be taken with caution as the observational nature of this study does

not allow us to establish a clear causal relationship in the observed patterns of the different

species (i.e., temporal avoidance between competitors). Future research should address

variations in the relative abundance of mesopredators across sites with different rabbit

availability. Unfortunately, our sample size was limited to address these questions and we

recommend complementing camera trap data with other census methods (i.e. pellet counts or

track surveys). Complementarily, trophic relationships between potential competitors could

be evaluated by studying the diet composition through the collection, identification and

analysis of faecal samples in the different study sites.

Conclusions

The activity patterns of the different mesopredators are ultimately restricted by the

circadian boundaries, but at the same time can be slightly modulated by interspecific

interactions with both prey and potential competitors. Our results show that the coexistence

of the different species that compose the mesocarnivore assemblage in the Doñana National

Park can be facilitated by the temporal segregation among generalist species due to a

divergence in the activity patterns (nocturnal, diurnal and cathemeral) but also by differences

in trophic preferences among specialist species, as suggested in previous studies (Fedriani et

al., 1999; Palomares & Delibes, 1991c; Soto & Palomares, 2015)

Our study further suggests that in order to clarify the coexistence mechanisms underlying

community structure, it is necessary to consider a multidimensional perspective, addressing

simultaneously multiple axes of the ecological niche. In markedly seasonal habitats like

Doñana, with patchy vegetation including ample gradients of vegetation cover and

successional status, even subtle differences in daily activity such as those documented here

represent multiple pathways to facilitate sympatry of coexisting mesopredator species. These

mechanisms might prove crucial for coexistence in settings where the apex predator is absent

and medium-sized carnivores live at high densities and show resource overlap. Recent

scenarios of rewilding with top predators may represent ideal conditions to experimentally

examine the effects of mesopredator release in diverse carnivore guilds like those of the

Mediterranean habitats studied here.

Acknowledgements

We thank J.P. González-Varó, T. Burgos, E. Quintero, C. Gutierrez, R. Tarifa and J.L.

Molina for technical support regarding species identification. We also thank P. Ferreras for

sharing his trophic data with us. I. Mendoza and B. Arroyo gave helpful comments on early

versions of the manuscript. Our work was funded by the project DISPMULTLAYER

(CGL2017-82847-P), Spanish Ministry of Science and Innovation (PJ). JI was supported by

the FPI grant PRE2018-085916 from the Spanish Ministry of Science and Innovation. ABL

was supported by a Juan de la Cierva-Incorporación grant (IJCI-2017-31419), Spanish

Ministry of Science, Innovation and Universities. We appreciate the support from ICTS-RBD

facilities during fieldwork.

Author contributions

ABL conceived the original idea; IF collected the diet data, performed the analyses, designed the figures and wrote the first draft of the manuscript; JI and PJ designed the methodology; JI and IF compiled the camera-trap data. All authors contributed to the writing of the manuscript.

Data accessibility

Data and R codes used in this study are deposited in GitHub (https://github.com/IagoFerreiroArias/TempPart)

References

Agostinelli, C., & Lund, U. (2017). *{R} package \texttt{circular}: Circular Statistics (version 0.4-93)*. CA: Department of Environmental Sciences, Informatics and Statistics, Ca' Foscari University, Venice, Italy. UL: Department of Statistics, California Polytechnic State University, San Luis Obispo, California, USA. Retrieved from https://r-forge.r-project.org/projects/circular/

Arias-Del Razo, I., Hernández, L., Laundré, J. W., & Myers, O. (2011). Do predator and prey foraging activity patterns match? A study of coyotes (*Canis latrans*), and lagomorphs (*Lepus californicus* and *Sylvilagus audobonii*). *Journal of Arid Environments*, *75*(2), 112–118.

Baker, P. J., Dowding, C. V., Molony, S. E., White, P. C. L., & Harris, S. (2007). Activity patterns of urban red foxes (*Vulpes vulpes*) reduce the risk of traffic-induced mortality. *Behavioral Ecology*, *18*(4), 716–724.

Barrientos, R., & Virgós, E. (2006). Reduction of potential food interference in two

sympatric carnivores by sequential use of shared resources. *Acta Oecologica*, *30*(1),

107–116.

Barrull, J., Mate, I., Ruiz-Olmo, J., Casanovas, J. G., Gosàlbez, J., & Salicrú, M. (2014).

Factors and mechanisms that explain coexistence in a Mediterranean carnivore

assemblage: an integrated study based on camera trapping and diet. *Mammalian*

Biology, *79*(2), 123–131.

Batschelet, E. (1981). *Circular statistics in biology*. New York: Academic Press.

Bennie, J. J., Duffy, J. P., Inger, R., & Gaston, K. J. (2014). Biogeography of time

partitioning in mammals. *Proceedings of the National Academy of Sciences of the*

United States of America, *111*(38), 13727–13732.

Brashares, J. S., Prugh, L. R., Stoner, C. J., & Epps, C. W. (2010). Ecological and

conservation implications of mesopredator release. In *Trophic cascades: predators,*

prey, and the changing dynamics of nature (pp. 221–240).

Camps, D. (2008). Activity patterns of adult common genets *Genetta genetta* (Linnaeus,

1758) in northeastern Spain. *Galemys*, *20*(1), 47–60.

Cavallini, P., & Lovari, S. (1994). Home range, habitat selection and activity of the red

fox in a Mediterranean coastal ecotone. *Acta Theriologica*, *39*(3), 279–287.

Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annual Review of*

Ecology and Systematics, *31*(1), 343–366.

Davis, M. L., Kelly, M. J., & Stauffer, D. F. (2011). Carnivore co☐existence and habitat

use in the Mountain Pine Ridge Forest Reserve, Belize. *Animal Conservation*,

14(1), 56–65.

Dellafiore, C. M., Rouco, C., Vallés, S. M., & Fernández, J. B. G. (2014). Seasonal

habitat use by the European rabbit (*Oryctolagus cuniculus*) in a coastal dune system

in SW Spain. *Animal Biodiversity and Conservation*, *2*, 233–242.

Di Bitetti, M. S., Di Blanco, Y. E., Pereira, J. A., Paviolo, A., & Pírez, I. J. (2009). Time

Partitioning Favors the Coexistence of Sympatric Crab-Eating Foxes (*Cerdocyon*

thous) and Pampas Foxes (*Lycalopex gymnocercus*). *Journal of Mammalogy*, *90*(2),

479–490.

Díaz☐Ruiz, F., Caro, J., Delibes☐Mateos, M., Arroyo, B., & Ferreras, P. (2016).

Drivers of red fox (*Vulpes vulpes*) daily activity: prey availability, human

disturbance or habitat structure? *Journal of Zoology*, *298*(2), 128–138.

Ensing, E. P., Ciuti, S., De Wijs, F. A. L. M., Lentferink, D. H., Hoedt, A., Boyce, M.

S., & Hut, R. A. (2014). GPS based daily activity patterns in european red deer and

North American elk (*Cervus elaphus*): Indication for a weak circadian clock in

ungulates. *PLoS ONE*, *9*(9), e106997.

Fedriani, J. M. (1996). Dieta anual del zorro, *Vulpes vulpes*, en dos hábitats del Parque

Nacional de Doñana. *Doñana Acta Vertebrata*, *23*(2), 143–152.

Fedriani, J. M., Ferreras, P., & Delibes, M. (1998). Dietary response of the Eurasian

badger, *Meles meles*, to a decline of its main prey in the Donana National Park.

Journal of Zoology, *245*(2), 214–128.

Fedriani, J. M., Palomares, F., & Delibes, M. (1999). Niche relations among three

sympatric Mediterranean carnivores. *Oecologia*, *121*(1), 138–148.

Fenn, M. G. P., & Macdonald, D. W. (1995). Use of middens by red foxes: risk reverses

rhythms of rats. *Journal of Mammalogy*, *76*(1), 130–136.

Ferreras, P., Travaini, A., Zapata, S. C., & Delibes, M. (2011). Short-term responses of

mammalian carnivores to a sudden collapse of rabbits in Mediterranean Spain.

Basic and Applied Ecology, *12*(2), 116–124.

Fisher, N. I. (1995). *Statistical analysis of circular data*. Cambridge University Press.

Foster, V. C., Sarmento, P., Sollmann, R., Tôrres, N., Jácomo, A. T. A., Negrões, N., …

Silveira, L. (2013). Jaguar and puma activity patterns and predator☐prey

interactions in four Brazilian biomes. *Biotropica*, *45*(3), 373–379.

Gaynor, K. M., Hojnowski, C. E., Carter, N. H., & Brashares, J. S. (2018). The influence

of human disturbance on wildlife nocturnality. *Science*, *360*(6394), 1232–1235.

Holt, R. D., Grover, J., & Tilman, D. (1994). Simple rules for interspecific dominance in

systems with exploitative and apparent competition. *The American Naturalist*,

144(5), 741–771.

Jiménez, J., Nuñez-Arjona, J. C., Mougeot, F., Ferreras, P., González, L. M., García-

Domínguez, F., … López-Bao, J. V. (2019). Restoring apex predators can reduce

mesopredator abundances. *Biological Conservation*, *238*, 108234.

Kronfeld-Schor, N., & Dayan, T. (2003). Partitioning of time as an ecological resource.

Annual Review of Ecology, Evolution, and Systematics, *34*(1), 153–181.

Lê, S., Josse, J., & Husson, F. (2008). {FactoMineR}: A Package for Multivariate
Analysis. *Journal of Statistical Software*, *25*(1), 1–18.

Lima, S. L., & Bednekoff, P. A. (1999). Temporal variation in danger drives
antipredator behavior: the predation risk allocation hypothesis. *The American*
Naturalist, *153*(6), 649–659.

Linkie, M., & Ridout, M. S. (2011). Assessing tiger–prey interactions in Sumatran
rainforests. *Journal of Zoology*, *284*(3), 224–229.

Lode, T. (1995). Activity pattern of polecats *Mustela putorius* L. in relation to food
habits and prey activity. *Ethology*, *100*(4), 295–308.

MacArthur, R. H., & Pianka, E. R. (1966). On optimal use of a patchy environment. *The*
American Naturalist, *100*(916), 603–609.

Maloney, S. K., Moss, G., Cartmell, T., & Mitchell, D. (2005). Alteration in diel activity
patterns as a thermoregulatory strategy in black wildebeest (*Connochaetes gnou*).
Journal of Comparative Physiology A, *191*(11), 1055–1064.

Martín, R., Rodríguez, A., & Delibes, M. (1995). Local feeding specialization by
badgers (*Meles meles*) in a Mediterranean environment. *Oecologia*, *101*(1), 45–50.

Monterroso, P., Alves, P. C., & Ferreras, P. (2013). Catch me if you can: Diel activity
patterns of mammalian prey and predators. *Ethology*, *119*(12), 1044–1056.

Monterroso, P., Alves, P. C., & Ferreras, P. (2014). Plasticity in circadian activity
patterns of mesocarnivores in Southwestern Europe: Implications for species
coexistence. *Behavioral Ecology and Sociobiology*, *68*(9), 1403–1417.

Monterroso, P., Díaz￮Ruíz, F., Lukacs, P. M., Alves, P. C., & Ferreras, P. (2020).

Ecological traits and the spatial structure of competitive coexistence among

carnivores. *Ecology*, *101*(8), e03059.

Monzón, A., Fernandes, P., & Rodrigues, N. (2004). Vegetation structure descriptors

regulating the presence of wild rabbit in the National Park of Peneda-Gerês,

Portugal. *European Journal of Wildlife Research*, *50*(1), 1–6.

Newsome, T. M., Greenville, A. C., Ćirović, D., Dickman, C. R., Johnson, C. N., Krofel,

M., … Wirsing, A. J. (2017). Top predators constrain mesopredator distributions.

Nature Communications, *8*, 15469.

Palomares, F. (1993). Opportunistic feeding of the Egyptian mongoose, *Herpertes*

ichneumon (L.) in Southwestern Spain. *Revue d'écologie*, *48*, 295–304.

Palomares, F., & Caro, T. M. (1999). Interspecific killing among mammalian carnivores.

The American Naturalist, *153*(5), 492–508.

Palomares, F., & Delibes, M. (1991a). Alimentación del meloncillo *Herpestes*

ichneumon y de la gineta *Genetta genetta* en la Reserva Biológica de Doñana, SO

de la Península Ibérica. *Doñana Acta Vertebrata*, *18*(1), 5–20.

Palomares, F., & Delibes, M. (1991b). Dieta del meloncillo (*Herpestes ichneumon*), en

el Coto del Rey (Norte del Parque Nacional de Doñana, SO España). *Doñana: Acta*

Vertebrata, *18*(2), 197–194.

Palomares, F., & Delibes, M. (1991c). Ecología comparada de la gineta *Genetta genetta*

(L.) y el meloncillo *Herpestes ichneumon* (L.) (Mammalia, Viverridae) en Doñana

(SO de la Península Ibérica). *Boletín de La Real Sociedad Española de Historia*

Natural (Sección Biología), *87*(1–4), 257–266.

Palomares, F., & Delibes, M. (1992). Circadian Activity Patterns of Free-Ranging Large

Gray Mongooses, *Herpestes ichneumon*, in Southwestern Spain. *Journal of*

Mammalogy, *73*(1), 173–177.

Palomares, F., & Delibes, M. (1994). Spatio-temporal ecology and behavior of European

genets in southwestern Spain. *Journal of Mammalogy*, *75*(3), 714–724.

Palomares, F., Ferreras, P., Fedriani, J. M., & Delibes, M. (1996). Spatial relationships

between Iberian lynx and other carnivores in an area of south-western Spain.

Journal of Applied Ecology, *33*, 5–13.

Palomares, F., Ferreras, P., Travaini, A., & Delibes, M. (1998). Co-existence between

Iberian lynx and Egyptian mongooses: estimating interaction strength by structural

equation modelling and testing by an observational study. *Journal of Animal*

Ecology, *67*(6), 967–978.

Polis, G. A., Myers, C. A., & Holt, R. D. (1989). The ecology and evolution of

intraguild predation: potential competitors that eat each other. *Annual Review of*

Ecology and Systematics, *20*(1), 297–330.

Prugh, L. R., Stoner, C. J., Epps, C. W., Bean, W. T., Ripple, W. J., Laliberte, A. S., &

Brashares, J. S. (2009). The rise of the mesopredator. *Bioscience*, *59*(9), 779–791.

Pyke, G. H., Pulliam, H. R., & Charnov, E. L. (1977). Optimal foraging: a selective

review of theory and tests. *The Quarterly Review of Biology*, *52*(2), 137–154.

R Core Team. (2020). *R: A Language and Environment for Statistical Computing*.

Vienna, Austria: R Foundation for Statistical Computing. Retrieved from

https://www.r-project.org/

Rao, J. S. (1976). Some tests based on arc-lengths for the circle. *Sankhyā: The Indian*

Journal of Statistics, Series B, 329–338.

Rau, J. R. (1987). *Ecología del zorro," Vulpes vulpes (L.)", en la reserva biológica de*

Doñana, SO de España. Universidad de Sevilla.

Revilla, E., & Palomares, F. (2002). Does local feeding specialization exist in Eurasian

badgers? *Canadian Journal of Zoology*, *80*(1), 83–93.

Ridout, M. S., & Linkie, M. (2009). Estimating overlap of daily activity patterns from

camera trap data. *Journal of Agricultural, Biological, and Environmental Statistics*,

14(3), 322–337.

Rodríguez, A., Martin, R., & Delibes, M. (1996). Space use and activity in a

Mediterranean population of badgers *Meles meles*. *Acta Theriologica*, *41*, 59–72.

Rowcliffe, J. M., Kays, R., Kranstauber, B., Carbone, C., & Jansen, P. A. (2014).

Quantifying levels of animal activity using camera trap data. *Methods in Ecology*

and Evolution, *5*(11), 1170–1179.

Rowcliffe, M. (2019). *activity: Animal Activity Statistics*. Retrieved from https://cran.r-

project.org/package=activity

Santos, F., Carbone, C., Wearn, O. R., Rowcliffe, J. M., Espinosa, S., Lima, M. G. M.,

… Alvarez-Loayza, P. (2019). Prey availability and temporal partitioning modulate

felid coexistence in Neotropical forests. *PloS One*, *14*(3), e0213671.

Schoener, T. W. (1974). The compression hypothesis and temporal resource

partitioning. *Proceedings of the National Academy of Sciences of the United States*

of America, *71*(10), 4169–4172.

Soto, C., & Palomares, F. (2015). Coexistence of sympatric carnivores in relatively

homogeneous Mediterranean landscapes: functional importance of habitat

segregation at the fine-scale level. *Oecologia*, *179*(1), 223–235.

Soulé, M. E., Bolger, D. T., Alberts, A. C., Wrights, J., Sorice, M., & Hill, S. (1988).

Reconstructed Dynamics of Rapid Extinctions of Chaparral Requiring Birds in

Urban Habitat Islands. *Conservation Biology*, *2*(1), 75–92.

Terborgh, J., Holt, R. D., & Estes, J. (2010). Trophic cascades: what they are, how they

work, and why they matter. In *Trophic Cascades: Predators, Prey and the*

Changing Dynamics of Nature.

Valverde, J. A. (1957). Notes écologiques sur le lynx d'Espagne feus Lynx Pardina

temminck. *Terre et Vie (Revue d'Ecologie)*, *104*, 51–67.

Valverde, J. A. (1967). *Estructura de una comunidad mediterránea de vertebrados*

terrestres. Madrid: Consejo Superior de Investigaciones Científicas (España).

van Schaik, C. P., & Griffiths, M. (1996). Activity Periods of Indonesian Rain Forest

Mammals. *Biotropica*, *28*(1), 105–112.

Vazquez, C., Rowcliffe, J. M., Spoelstra, K., & Jansen, P. A. (2019). Comparing diel

activity patterns of wildlife across latitudes and seasons: Time transformations

using day length. *Methods in Ecology and Evolution*, *10*(12), 2057–2066.

Vilella, M., Ferrandiz☐Rovira, M., & Sayol, F. (2020). Coexistence of predators in

time: Effects of season and prey availability on species activity within a

Mediterranean carnivore guild. *Ecology and Evolution*, *00*, 1–15.

Villafuerte, R., Kufner, M. B., Delibes, M., & Moreno, S. (1993). Environmental factors

influencing the seasonal daily activity of the European rabbit (*Oryctolagus*

cuniculus) in a Mediterranean area. *Mammalia*, *57*(3), 341–347.

Virgós, E., Llorente, M., & Cortésá, Y. (1999). Geographical variation in genet (*Genetta*

genetta L.) diet: a literature review. *Mammal Review*, *29*(2), 117–126.

Wang, Y., & Fisher, D. O. (2012). Dingoes affect activity of feral cats, but do not

exclude them from the habitat of an endangered macropod. *Wildlife Research*,

39(7), 611–620.

Zielinski, W. J. (1986). Circadian rhythms of small carnivores and the effect of

restricted feeding on daily activity. *Physiology & Behavior*, *38*(5), 613–620.

Figures

Figure 1. Niche segregation along the trophic axis among mesocarnivores in Doñana National Park. The first two CA dimensions (Dim 1, Dim 2) explain 66% of the variability in diet composition between species and studies. The first dimension represents a gradient of diet variability between the common genet and the European badger, while the second dimension separates the diet of the red fox and the common genet. Convex polygons represent trophic niche breadth.

Figure 2. Pairwise comparisons of daily activity patterns between mesocarnivores (rows, top to bottom: red fox, badger, genet, and Egyptian mongoose) and their main prey items in the study area (columns, left to right: *Oryctolagus cunniculus*, small mammals, and *Alectoris rufa*). Coloured lines describe prey activity patterns (blue for European rabbit, green for small mammals and yellow for red-legged partridge) and black lines indicate predator's activity patterns. Overlap coefficients (Δ) and their respective 99% confidence intervals are shown in brackets.

Figure 3. Diel activity patterns of two mesocarnivores (red fox and Egyptian mongoose; mid and bottom rows, respectively) and its main prey item (European rabbit, top row) along the gradient of ecological succession, from younger juniper stands (SABCOL) to mature, late successional stands (SABMAR).

Figure 4. Kernel density plots and temporal overlap between mesocarnivores in the study area: common genet (green line), red fox (blue line), European badger (purple line) and Egyptian mongoose (yellow line). Overlap coefficients (Δ) and their respective 99% confidence intervals are shown in brackets.

Landscape maturity