1 **Temporal and trophic partitioning promote coexistence between**
2 **mesocarnivores in a Mediterranean landscape**
3

4 Iago Ferreiro-Arias[1], Jorge Isla[1], Pedro Jordano[1], Ana Benítez-López[1]
5
6 [1]Department of Integrative Ecology, Estación Biológica de Doñana, Consejo Superior de
7 Investigaciones Científicas, Avda. Américo Vespucio, 26, 41092, Sevilla, Spain.
8
9 *Corresponding author: iago.ferreiro.arias@gmail.com

Abstract

Abstract

The partition of the ecological niche can enhance the coexistence of predators due to

differences in how they exploit three main resources: food, space, and time, the latter being

an axis that remains often unexplored. We studied niche segregation in a Mediterranean

mesocarnivore community in sites where the top predator, the Iberian lynx, is absent,

addressing simultaneously two niche axes: the temporal and the trophic axes. Temporal

overlap between prey and predator and potential competitors was analysed by comparing

daily activity patterns of predators and prey species through data obtained by using 24

camera-traps deployed in three different sites of the Doñana National Park. We further

examined the trophic dimension by compiling and analysing diet information in the study

area and exploring trophic preferences between potential competitors. We found an overall

temporal segregation between trophic generalist species while those species which showed

higher values of temporal overlap differ largely in their trophic preferences. Furthermore, we

observed an overall high degree of overlap between the activity patterns of predators and

their putative prey (the common genet vs small mammals and the red fox vs European

rabbit). Our study suggests that coexistence of the different species that compose the

mesocarnivore assemblage in Mediterranean landscapes can be facilitated by the marked

temporal segregation of the generalist species due to a divergence in the activity patterns, but

also by differences in trophic preferences by specialist species when their activity patterns

overlap.

Keywords

Introduction

Understanding the mechanisms that promote coexistence of species with similar ecological requirements is a central topic in community ecology, where competitive interactions between species have the potential to affect diversity patterns by limiting or promoting coexistence (Chesson 2000). In order to mitigate the negative impact of interspecific competition, species often partition resources along three main niche dimensions (temporal, trophic and spatial), which eventually results in niche differentiation (Schoener 1974). Most studies concerning species coexistence focused on the differential use of habitat and food resources; nonetheless, the differential use of the diel cycle may enhance coexistence of same-sized species, particularly among predator species (Di Bitetti et al. 2009, Wang and Fisher 2012).

Diel activity patterns are cyclical behavioural patterns which can be influenced by several factors (i.e., prey availability or thermoregulatory strategies) (Lode 1995, Maloney et al. 2005). Within mammals, a large variability and plasticity in their activity patterns has been documented (Bennie et al. 2014). This diversity of diel cycles could be due to the plastic nature of behavioural responses to different pressures which could induce marked variations in daily rhythms among different scenarios (Ensing et al. 2014, Gaynor et al. 2018). Broadly, activity patterns displayed by mammals are determined by circadian endogenous boundaries (Kronfeld-Schor and Dayan 2003) which are particularly shaped in carnivores by external factors (Zielinski 1986, Monterroso et al. 2014) setting a context-dependency in their diel cycle use. Hence, the co-occurrence of species that compete for a certain resource has the potential to produce shifts in the daily activity patterns of competitively surrogate species. For example, crab-eating and pampas foxes have been reported to change their daily foraging activity in sympatric areas where both species are present (switching from diurnal to

nocturnal), compared to allopatric areas where only one fox species is present (Di Bitetti et

al. 2009).

In addition, predators' activity patterns also could be influenced by rhythms of prey

activity where an arms race occurs through the temporal axis. Predators could adjust their

rhythms of daily activity to those hours of the day in which the availability of prey increases,

producing a synchrony between the daily activity patterns of the predator and their prey

(Foster et al. 2013), while at the same time, prey species could generate a temporal mismatch,

centering their activity at times when there is a lower risk of predation (Fenn and Macdonald

1995). Consequently, high temporal overlaps between carnivores and their putative prey have

been reported (Linkie and Ridout 2011) while in other predator-prey systems an asynchrony

was found in their respective activity peaks (Arias-Del Razo et al. 2011, Díaz-Ruiz et al.

2016). As a consequence, the daily activity of a given predator could be a result of a cost-

benefit trade-off between maximizing their activity during periods with high prey availability

(benefit) but also higher mortality risk due to encounters with intraguild competitors (cost)

(Santos et al. 2019).

Interspecific interactions play an important role in structuring a mammalian

community (Holt et al. 1994), especially in carnivores where competition not only arises for

exploiting the same resource simultaneously, but also for the risk associated with intraguild

predation. Indeed, interspecific relationships between different predators can be modulated

through direct or indirect competition. An example of a direct interaction is intraguild

predation (IGP), a type of interference competition where a predator species kills another

predator, or where both species prey upon each other (Polis et al. 1989, Palomares and Caro

1999). IGP is mediated mainly by species' biological characteristics such as body size, prey

availability as well as their trophic ecologies (Linnell and Strand 2000), being specially reported on top-predators (Palomares and Caro 1999).

The effects that top predators exert on surrogate competitors can be diverse, from abundance reductions and distributional shifts (in space and time) (Newsome et al. 2017, Jiménez et al. 2019) to behavioural modifications through restriction of their activity patterns (Wang and Fisher 2012). Apex predators play a key role in ecosystem functionality due to their regulatory role on prey populations (top-down control) and their extinction frequently triggers cascading effects through direct and indirect interactions among trophic levels (Terborgh et al. 2010). An effect of apex predator's extinction is the eventual ecological release of medium-sized carnivore populations, leading to increases in the distribution and abundance of their populations (Prugh et al. 2009) ; a phenomenon coined '*mesopredator release*' by Soulé et al. (1988), which has been repeatedly documented (Brashares et al. 2010).

In the Iberian Peninsula, the Iberian lynx (*Lynx pardinus*) exerts its role as a top-predator on the populations of surrogate competitors (Palomares et al. 1996, 1998). Sizeable reductions of mesocarnivore abundance have been reported after the reintroduction of the Iberian lynx (Jiménez et al. 2019), reinforcing the idea that the coexistence between these carnivores is supported by spatial structuring where the probability of occurrence of the lynx is negatively associated with the presence of mesopredators (Palomares et al. 1996, Monterroso et al. 2020). In Doñana National Park, the carnivore guild is composed by a habitat and trophic specialist top predator (the Iberian lynx, *Lynx pardinus*) and mesocarnivores such as the red fox (*Vulpes vulpes*), the European badger (*Meles meles*) and the Egyptian mongoose (*Herpestes ichneumon*), which are characterized by a wide niche breadth due to their generalist behaviour concerning habitat selection and trophic preferences.

The negative effects that the Iberian lynx has on these mesocarnivores occur mainly through

intraguild predation, particularly on Egyptian mongooses (*Egyptian mongoose)*, common

genets (*Genetta genetta)* and red foxes (*Vulpes vulpes*) (Palomares et al. 1996, Palomares and

Caro 1999, and references therein). However, aggressive interactions between the different

mesocarnivores remain quite unknown since most of the documented IGP interactions in

these scenarios were associated to the Iberian lynxes against surrogate competitors, and not

between mesopredator species (Valverde 1957, Palomares et al. 1996, Fedriani et al. 1999).

Therefore, in areas where the Iberian lynx is not present, indirect relationships may become

more relevant between these mesopredators since coexistence seems not to be mediated by

spatial features of their respective ecological niches (Soto and Palomares 2015).

Indirect competitive interactions can arise through competition for exploitation, which

occurs when one of the species consumes trophic or spatial resources more efficiently and

reduces their availability to the rest of the species (e.g., Holt et al. 1994). Within

mesocarnivore communities, these interactions can modulate activity patterns in order to

reduce spatial or temporal overlap between species with similar trophic requirements (Bu et

al. 2016). Empirical data in this respect are surprisingly very limited for the Mediterranean

area, where situations of mesopredator release may be common due to recent historical

reductions in the Iberian lynx populations. Yet, most previous studies have focused on spatial

avoidance as the main mechanism allowing the mesocarnivore coexistence, and only recently

has the focus shifted towards temporal segregation (Barrull et al. 2014, Monterroso et al.

2014, Vilella et al. 2020).

The main purpose of this study is to evaluate the ecological factors that allow the

coexistence of mesocarnivore species in the Doñana National Park without the presence and

influence of a top-predator, focusing in particular on the temporal segregation of four species

of mesocarnivores: the red fox (*Vulpes vulpes*), the European badger (*Meles meles*), the

common genet (*Genetta genetta*), and the Egyptian mongoose (*Herpestes ichneumon*). Thus,

this study aims: 1) to explore the relationship between mesocarnivores along the trophic

niche axis; 2) to determine the diel activity patterns of predator and prey species and their

synchrony; and 3) to quantify the temporal overlap or segregation between potential

competitors. Following a niche partitioning conceptual framework, we hypothesize that there

would be greater temporal segregation in those species that have a higher degree of overlap in

their respective trophic niches, while species that have similar use of the diel cycle should

show differentiated trophic preferences in order to promote coexistence by reducing

competition along temporal or trophic niche axes.

Material and methods

Study area

This study was carried out in Doñana National Park (37º1'N, 6º 33'W) in three 1-ha

sites characterized by juniper stands (*Juniperus phoenicea* subsp. *turbinata)* of different age

and ecological succession stages. The selected sites represent a gradient of habitat maturity in

terms of ecological succession that ranges from an advance front of juniper stands to mature

stands with a higher degree of vegetation cover: "*Sabinar de Colonización*" (SABCOL)

represents an early stage with 2030 junipers/ha, "*Sabinar del Ojillo*" (SABOJI) is regarded as

an intermediate stage with 9010 junipers/ha and "*Sabinar del Marqués*" (SABMAR)

constitutes the final stage with 10520 junipers/ha (Figure S1, Supplementary information).

Data collection

We located eight camera traps per site that were rotated periodically through 35

randomly selected focal juniper plants. Juniper locations were selected at random in each of

five subplots within each of the three main sites, with the subplots distributed regularly

throughout the main plot. The structure of this habitat consists of paths between juniper

individuals, so the cameras were installed at 1-m height focusing on the ground and the

bottom of the plant. Camera traps were active for ca. 10 days in each of these rotations and

then changed to other 8 locations, thus ensuring that the whole site was sampled in 5 weeks.

Following this procedure, data were obtained from a total set of 24 camera traps sampling the

three sites during two campaigns: the first one from October 2018 to May 2019 and the

second one from October 2019 to June 2020.

The diel activity cycle of each species was characterized by pooling the total number

of detections across all cameras during the whole study period. In order to avoid data

dependency, in our analyses we discarded consecutive detections of a given species within a

site (Davis et al. 2011, Monterroso et al. 2014). Since the area covered by our study sites is

limited (ca. 1 ha/site), we set a period of 15 min between consecutive detections of a given

species within a study site, which is a reasonable time period for a species to abandon the

site. Moreover, in such cases where a given individual was detected repeatedly without

leaving the camera detection zone, only the time of its first detection was considered. In

analyses we used either the total, absolute number of records (e.g., to compare the relative

frequency of sunrise, day, dusk, and nocturnal records within a given site) or the relative

frequency of records expressed as number per camera-day (e.g., when comparing across sites

that received different sampling intensity).

Target species consisted mainly of mesopredators which occur in Doñana National

Park and their respective prey species. The mesocarnivore guild in the study area is

composed by the Egyptian mongoose (*Herpestes ichneumon*), the red fox (*Vulpes vulpes*), the

Eurasian badger (*Meles meles*) and the common genet (*Genetta genetta*) (Valverde 1967).

Large top predators either extinct or at low density in the area include the wolf (*Canis lupus*)

and the Iberian lynx (*Lynx pardinus*) (Valverde 1967). Simultaneously, other species

recorded during the course of this study were considered as possible prey items for predators

such as lagomorphs (mainly European rabbits, *Oryctolagus cuniculus*), small mammals and

birds such as *Alectoris rufa*. Detections from ungulates (mainly *Cervus elaphus* and *Sus*

scrofa) were not included in the analysis as their presence in mesocarnivore diets is

frequently associated with carrion consumption, with direct predation being recorded

sporadically on very young individuals (Valverde 1967). No attractants were used over the

two years of sampling.

185   Trophic overlap data

With the aim of exploring trophic overlap between the predators mentioned above, we

extracted diet information from a set of studies carried out in Doñana National Park (Table

S1, Supplementary information). We searched WOS and Google Scholar for studies that

reported trophic preferences of the four mesocarnivores in Doñana National Park using the

following search string: (trophic OR diet) AND (*Vulpes vulpes* OR *Genetta genetta* OR

Meles meles OR *Herpestes ichneumon*) AND (Doñana). The selection of studies was aimed

at compiling all the trophic information available for the four species at different locations in

the Doñana National Park to better estimate the entire trophic spectra of each of the different

species. Given that the four target species are commonly reported as opportunists in their

feeding behaviour, this approach was taken in order to capture the natural variability in

dietary composition across the whole Doñana study area and its marked seasonal component.

To avoid pseudo-replication issues, we selected those studies in which the data come

from strictly different samples and periods. From each study, period and species, we

extracted the percentage of occurrence of food items. In order to estimate trophic overlap

between mesocarnivores, we carried out a correspondence analysis (CA) using the percentage

of occurrence of each prey item for each predator. The CA ordered the different diet studies

(cases) relative to the prey species composition reported by each study, resulting in an

ordination of the studies corresponding to each mesopredator species, based on their

similarity of prey species composition. We used the coordinates on the first two CA axes

corresponding to each diet study to build convex polygons spanning each mesopredator group

of studies. In this way, these polygons illustrate the range of dietary variation shown by each

mesopredator species in relation to the diet composition of other species.

Camera trap data analysis

Time and date were extracted from each detection of every camera trap to estimate

daily activity patterns and temporal overlap between mesocarnivores and their potential prey

and between mesocarnivore potential competitors. As time and dates are variables with a

circular nature, diel activity cycles were estimated using non-parametric Kernel density plots

which provide a density function of the daily activity pattern of a given species (Rowcliffe et

al. 2014). Raw data were subsampled in order to carry out analyses only for those species

with more than 10 detections (Fisher 1995, Gerber et al. 2012).

To determine the daily activity pattern of the species we resorted to the classification

used by van Schaik and Griffiths (1996) by which species are classified according to the

number of independent detections obtained as: diurnal (> 90% of the detections occurring

during the daylight), nocturnal (> 90% of the detections obtained at night), crepuscular (>

90% of their detections occur at dawn or dusk) or cathemeral if their activity pattern is

distributed uniformly throughout the daily cycle. We tested whether the target species

actually show a certain activity pattern and deviate from a random pattern using the Rao

Spacing Test (U) (Rao 1976).

Moreover, as the present study has been carried out over two years, daylight can vary

from one detection to another and induce bias to our results. Species could adjust their daily

activity levels depending on the day length, thus not taking this aspect into account when

dealing with strictly diurnal or nocturnal species could lead to the misinterpretation of the

results and the underestimation of peaks of daily activity (Vazquez et al. 2019). Therefore,

we carried out a double average anchoring time transformation to take into account the mean

average sunrise and sunset times for the study area (Vazquez et al. 2019).

We evaluated the extent of temporal segregation between mesopredators using the

overlap coefficient (Δ) described by Ridout and Linkie (2009). This coefficient ranges from 0

to 1, where 0 values represent completely different diel activity patterns while 1 represents

the maximum overlap between both species. Ridout and Linkie (2009) have reported

different estimators of the degree of overlap between the daily activity curves and tested

which of them represent a better estimate of the temporal overlap as a function of the sample

size. Briefly, Δ_1 represents accurately the degree of overlap between two curves when the

smallest size of the two samples is less than 50 detections while Δ_4 best fits when the smallest

sample size exceeds 75 detections. Following Ridout and Linkie (2009), we used Δ_1 for those

cases in which sample size was lower than 75 detections, and Δ_4 when sample sizes were

equal or greater than 75 detections.

Once the overlap coefficients between the different species were estimated, 99%

confidence intervals were calculated through smoothed bootstrap analysis with 10,000

replicates. Moreover, the significance of differences in pairwise comparisons of daily activity

patterns were evaluated by using the non-parametric Mardia-Watson-Wheeler test (MWW)

which allows us to detect whether two circular samples differ significantly from each other

(Batschelet 1981). However, deciding whether a certain degree of temporal overlap is high or

low based solely on this coefficient is somewhat subjective. Therefore, we followed

Monterroso et al. (2014) and used percentiles to establish threshold levels of overlap for our

discussion. Thus, when $\Delta >$ 75th percentile, a high temporal overlap exists between both

species, while $\Delta \leq$ 50th percentile values show a low degree of overlap. Intermediate values

of these percentiles, 50th $< \Delta \leq$ 75th, were interpreted as moderate levels of overlap.

Statistical software

All analyses were conducted in R Statistical Software 3.6.2 (R Core Team 2020). Data

handling was done with the package "dplyr" (Wickham et al. 2020); the correspondence

analysis (CA) was carried out with "*FactoMineR*" package (Lê et al. 2008); kernel density

functions, overlap coefficients (Δ) and their respective confidence intervals were calculated

with the package "*overlap*" (Ridout and Linkie 2009), and MWW and Rao Spacing tests

were conducted using the "*circular*" package (Agostinelli and Lund 2017). Finally, in order

to calculate mean average sunrise and sunset times and perform average anchoring

transformation we used the "*activity*" package (Rowcliffe 2019).

Results

Trophic segregation among mesocarnivores

The correspondence analysis explains approximately 66% of the variability of

mesocarnivore diets through the first two dimensions (Figure 1). Axis 1, which explains

51.6% of the variation, discriminates the set of mesocarnivores into two separate groups:

species whose diet is mainly omnivorous (red fox and European badger) and those with

essentially carnivorous diets (common genet and Egyptian mongoose). The red fox and the

European badger showed very close values due to their frugivorous behaviour and the

predominant consumption of lagomorphs and invertebrates. Simultaneously, axis 2, which

explains 15% of the trophic variation, is generated by variation between the diets of the

Egyptian mongoose and the common genet, with the latter having a trophic preference for

small mammals followed by a secondary intake of birds. However, the second dimension of

the correspondence analysis does not discern between the trophic niches of the red fox and

the European badger, indicating that there is a greater trophic overlap between these two

omnivorous species with respect to the rest of the mesocarnivore guild. Finally, the convex

polygons represent the more ample trophic diversification of the generalist species (red fox,

European badger and Egyptian mongoose) whose range of prey types tends to be

considerably wider compared with the common genet and its specialist feeding behaviour

(Figure 1).

Camera trapping results and diel activity patterns

Results obtained in this study come from a total of 531 effective sampling days. All

sites were sampled with same number of cameras per site and a similar effort (number of

days): $\bar{x} \pm SD = 80.3 \pm 28.3$ d for SABCOL; 80.4 ± 27.3 d for SABOJI and 86.8 ± 26.0 d for

SABMAR. Thus, sampling effort was quite evenly distributed among sites. Broadly, a total

of 2786 detections were obtained throughout the study period, of which 96 (3.44%) could not

be identified at the species level. Among all detections, 16% corresponded to

mesocarnivores, 28.7% to prey species (e.g. rabbit, partridge, small mammals…), 15% to

ungulates and 38.7% to passeriformes (Table S2, Supplementary information).

Regarding target species, detections accounted for by the red fox represented a total of

301 of which 55% were in SABCOL, while the rest of the detections were evenly distributed

between the other two sites. A large amount of the detections obtained from this species were

linked to its nocturnal habits while the rest were distributed throughout the periods with

daylight, demonstrating a cathemeral activity pattern with a tendency towards bimodality

(U = 142.26; p-value < 0.05) (Table S3, Supplementary information). Conversely, detections

of European badgers were very scarce in the three study sites, showing a predominantly

nocturnal activity pattern, with 77% of the detections occurring at night and the rest during

the dawn or dusk periods (U = 178.95; p-value < 0.01). Likewise, the activity pattern shown

by the common genet was strictly nocturnal (95% of the records produced during nighttime;

U = 165.81; p-value < 0.01) and linked to SABCOL where 62% of its detections took place.

Finally, detections of Egyptian mongooses were mainly concentrated in the SABOJI (54%)

while the other two sites maintained the same proportion of records, showing a marked

diurnal activity pattern in each of them (92% of the detections occurred during daylight; U =

199.78; p-value < 0.01) (Table S3, Supplementary information).

For prey species, most of the detections corresponded to the European rabbit

(*Oryctolagus cuniculus*), the red-legged partridge (*Alectoris rufa*) and small mammals.

European rabbits were detected 650 times with an even distribution between SABOJI and

SABMAR, while SABCOL had only 6 detections during the entire sampling period (Table

S2, Supplementary information). Rabbit detections were distributed throughout the day,

setting it as a cathemeral species with distinct activity peaks around sunrise and sunset (U =

156.97; p-value < 0.01) (Table S3, Supplementary information). Conversely, we did not have

sufficient detections from the other lagomorph present in RBD (*Lepus granatensis*, N = 3) to

include them in this study.

Small mammals accounted for a total of 98 detections, mainly associated with

SABMAR and detected only during nighttime (U = 198.27; p-value < 0.01). Red-legged

partridges were mainly detected in SABCOL (89% of detections) with a markedly diurnal

activity pattern (U = 226.53; p-value < 0.01) (Table S3, Supplementary information). Finally,

detections of ungulates present in the area (mainly *Cervus elaphus* and *Sus scrofa*)

represented approximately 14% of the total number of detections.

Predator and prey diel activity patterns

Broadly, the activity patterns and their corresponding overlap differed between

predators and the potential prey items in the study area. The European rabbit showed a high

degree of overlap with the red fox ($\Delta_4 = 0.84$), followed by a moderate overlap with the

European badger ($\Delta_1 = 0.62$). Conversely, both the common genet and the Egyptian

mongoose showed low overlap coefficients ($\Delta_1 = 0.47$ and $\Delta_1 = 0.44$, respectively) with

rabbit activity patterns (Figure 2).

We found a high temporal overlap between the activity patterns of small mammals

(with most detections at night) and the European badger ($\Delta_1 = 0.89$) and the common genet

($\Delta_1 = 0.76$). In turn, the red fox showed a moderate temporal overlap with small mammals (Δ_4

$= 0.68$), while the Egyptian mongoose showed a total temporal segregation with respect to the

detected micromammals ($\Delta_1 = 0.14$). Activity patterns of red-legged partridges overlapped

with those of the Egyptian mongoose ($\Delta_1 = 0.82$) while showing a large temporal segregation

with European badgers ($\Delta_1 = 0.08$) and common genets ($\Delta_1 = 0.03$) (Figure 2).

European rabbit activity patterns differed significantly between SABOJI and

SABMAR (W = 32.69; p-value < 0.01) (Figure 3). Conversely, the red fox showed shifts in

activity patterns between different sites with varying rabbit availability, with a predominantly

nocturnal activity in SABCOL and a tendency towards bimodal activity patterns in SABOJI

and SABMAR. Significative differences were found between activity patterns of red foxes

from SABCOL and SABMAR (W = 11.14; p-value < 0.01) while no differences were found

between SABCOL - SABOJI (W = 3.97; p-value = 0.137) or SABOJI - SABMAR (W =

2.44, p-value = 0.295). The temporal overlap for the European rabbit and the red fox was $\Delta_1 =$

0.79 for SABOJI and $\Delta_1 = 0.73$ for SABMAR while the temporal overlap between European

rabbit and Egyptian mongoose was $\Delta_1 = 0.49$ for SABOJI and $\Delta_1 = 0.37$ for SABMAR

(Figure 3). No statistically significant differences were found between activity patterns of the

Egyptian mongoose between SABCOL - SABOJI (W = 1.09; p-value = 0.58), SABCOL -

SABMAR (W = 2.06; p-value = 0.36) and SABOJI - SABMAR (W = 0.40; p-value = 0.82)

(Figure 3).

Temporal overlap and segregation between mesocarnivores

The mesocarnivore assemblage in the study area is composed mainly by two

predominantly nocturnal species, a strictly diurnal species and a species with a cathemeral

activity pattern. The average overlap and standard deviation of the activity patterns of the

different species was $\Delta_1 = 0.48$ and SD = 0.29, which reflect a low, although variable,

temporal overlap among these potential competitors. The average overlap values ($\Delta_1 \pm$ SD) of

a species with respect to the set of potential competitors were: 0.59 ± 0.15 for the fox; $0.57 \pm$

0.34 for the badger; 0.50 ± 0.36 for the genet and 0.24 ± 0.18 for the Egyptian mongoose.

Strictly nocturnal species (European badger and common genet) showed a high

overlap coefficient without significant differences between their activity patterns ($\Delta_1 = 0.79$;

W = 1.89, p-value = 0.386). On the contrary, the Egyptian mongoose, due to its diurnal

habits, showed a strong and significant temporal segregation with the European badger ($\Delta_1 =$

0.16; W = 41.75, p-value < 0.01) and the common genet ($\Delta_1 = 0.08$; W = 65.79, p-value <

0.01) (Figure 4). Simultaneously, the cathemeral activity pattern of the red fox sustained

intermediate levels of temporal segregation with the common genet ($\Delta_1 = 0.57$; W = 24.96; p-

value < 0.01) and the European badger ($\Delta_1 = 0.71$; W = 8.44; p-value = 0.014) and a low

overlap with the Egyptian mongoose ($\Delta_1 = 0.44$; W = 73.48; p-value < 0.01) (Figure 4).

Among study sites, temporal overlap between the red fox and the Egyptian mongoose was

$\Delta_1 = 0.41$ for SABCOL, $\Delta_1 = 0.45$ for SABOJI and $\Delta_1 = 0.36$ for SABMAR (Figure 3).

Discussion

In this study we explore how the sympatry of mesocarnivores in the Doñana National

Park could be mediated by temporal or trophic segregation when there is a greater overlap in

any of these two niche axes. Our study explores the mechanisms underlying the structure of

Mediterranean mesocarnivore communities, which have been traditionally focused on spatial

responses (Fedriani et al. 1999, Soto and Palomares 2015, Monterroso et al. 2020, but see

Barrull et al. 2014, Monterroso et al. 2014, Vilella et al. 2020), by providing new insights on

the importance of temporal segregation at facilitating co-occurrence by reducing resource

overlap. Overall, we found a moderate level of temporal segregation within the

mesocarnivore guild in our study area, which was composed by two strictly nocturnal species

- the European badger and the common genet (Palomares and Delibes 1994, Rodríguez et al.

1996, Camps 2008), one diurnal - the Egyptian mongoose (Palomares and Delibes 1992) and

one with plasticity in its daily activity pattern - the red fox, which ranges from diurnal

(Cavallini and Lovari 1994) to nocturnal habits (Baker et al. 2007).

The highest values of temporal overlap occurred between the European badger and the

common genet due to their predominantly nocturnal activity patterns. However, they

markedly differ in their respective trophic niches, with a segregation between them due to

preferential consumption of small mammals by the common genet. Indeed, the common

genet had a high temporal overlap with small mammals, its main prey item, which were

detected mostly at night. The common genet consumes almost all species of rodents and

insectivores, but it is certainly a specialized predator on *Apodemus sylvaticus*, the main prey

of its diet when this item is available (Palomares and Delibes 1991a, Virgós et al. 1999). This

synchrony between predator and prey is an expected outcome of Optimal Foraging Theory

constraints (MacArthur and Pianka 1966), by which an animal will try to maximize its prey

intake by showing an efficient foraging pattern. Predators whose diet is eminently specific are

expected to maximize synchrony of their daily activity patterns to their main and most

profitable prey (Pyke et al. 1977). However, this pattern would not be so marked in predators

whose feeding behaviour is opportunistic and depends on the availability and accessibility of

several prey types (Pyke et al. 1977), which is the case of the rest of mesocarnivores studied.

The rest of the species that constitute the Doñana mesocarnivore assemblage are

known for their generalist feeding habits (Rau 1987, Palomares 1993, Fedriani et al. 1999).

However, we found a distinct trophic segregation of the Egyptian mongoose with respect to

the red fox and the European badger, due to its eminently carnivorous behaviour, with plant

material (e.g., fleshy fruits) virtually absent from its diet. Meanwhile, the trophic overlap

between European badger and red fox was the highest within the mesocarnivore community

due to their omnivorous behaviour based on the consumption of fleshy fruits, invertebrates

and lagomorphs. Despite their reduced differences in terms of spatial and trophic niche, these

species showed a marked temporal segregation, displaying distinct activity patterns:

nocturnal, cathemeral and diurnal, which could ease coexistence by exploiting similar

resources at different times and thus reducing intraguild encounters.

The ecological plasticity and the ample spectra of resource use shown by these species

in trophic and habitat niche axes suggest a scenario with potentially reduced interspecific

competition (Palomares 1993, Fedriani et al. 1999, Monterroso et al. 2020). However, the

temporal segregation of these species may become more relevant, given that the other two

axes of their respective ecological niches can exhibit ample overlap. These situations could

be produced due to a reduction in the abundance and availability of its main trophic resources

at a given moment, increasing competition for the consumption of secondary resources

available. A clear example is the reduction in the consumption of European rabbits after a

sudden population collapse due to the emergence of the rabbit hemorrhagic disease (RHD)

(Ferreras et al. 2011), which resulted in an increased trophic overlap among most generalist

carnivore species. In such cases, time partitioning becomes the main mechanism to facilitate

coexistence by reducing competitive stress in resource use among these species (Barrientos

and Virgós 2006).

The European rabbit in the study area accounts for a very high percentage of the diet

of Egyptian mongooses (Palomares and Delibes 1991b) and also for European badgers. Some

authors even point out a possible specialization of badgers in this type of prey, a contentious

assertion which has been debated (Martín et al. 1995, Revilla and Palomares 2002). The

activity pattern of European rabbits matches those previously reported, showing a bimodal

pattern with peaks of activity at sunrise and sunset (Villafuerte et al. 1993, Monterroso et al.

2013, Díaz-Ruiz et al. 2016). However, rabbit availability differed between study sites since

during both sampling periods we did not obtain almost any detections in the least mature,

early successional juniper colonization front, the "*Sabinar de Colonization*" site. Conversely,

detections in "*Sabinar del Marqués*" and "*Sabinar del Ojillo*" were considerably more

frequent, which would reflect the differences in terms of prey availability among different

study sites. Such differences may be due to habitat requirements since, at the landscape scale,

the ideal habitat for rabbits is a juxtaposition of ecological units where the refuge areas are

intermingled with others where herbaceous food is more abundant (Monzón et al. 2004,

Dellafiore et al. 2014). The "*Sabinar de Colonización*" (SABCOL) represents a

Mediterranean scrubland in an early ecological stage composed mainly by a low-density

population of *J. phoenicea* (2030 junipers/ha) and therefore, a low vegetation cover which

does not fulfil rabbit habitat preferences. Meanwhile, "*Sabinar del Marqués*" (SABMAR)

and "*Sabinar del Ojillo*" (SABOJI) are sites characterized by late successional mature

vegetation and high vegetation cover, which may serve as refuge for rabbits. In these two

sites, we found differences between the diel cycle use of rabbits, showing two activity peaks

in SABOJI and only one crepuscular activity peak in SABMAR. Interestingly enough, the

activity pattern displayed by the red fox, showed a unimodal pattern with an activity peak

around midnight in SABCOL, but shifted towards bimodality in sites where the rabbit was

present, putatively to match the activity patterns of this prey and thus increase the probability

of encounter.

In SABMAR, the European rabbit had the lowest value of temporal overlap with its

main predators (red fox and Egyptian mongoose) since it concentrated its activity around

dusk periods while, in SABOJI, its bimodal pattern increased the overlap. This situation may

be a reflection of an anti-predatory behaviour of the European rabbit since, despite showing

two possible activity peaks in its daily cycle, its activity in SABMAR is centered at time

periods in which it does not overlap with its main predators. Meanwhile, in SABOJI, the

bimodal activity of the fox is less marked than in the SABMAR, matching with the activity

peaks of the rabbits and increasing the temporal overlap. In this way, the intensity of the red

fox activity could trigger a shift in the activity patterns of the European rabbit given that an

increase in the dawn activity of the predator in SABMAR is counteracted by rabbits which

display only one activity peak at twilight. This observation could support the Risk Allocation

Hypothesis by which a temporal mismatch between activity patterns of a prey species and its

predator could be due to a higher perception of mortality risk that drives anti-predatory

behaviours along the time axis (Lima and Bednekoff 1999). A similar situation has been

previously observed among other species of lagomorphs and canids, where the bimodal

patterns of the Black-tailed jackrabbit (*Lepus californicus*) and the desert cottontail

(*Sylvilagus audubonii*) showed a single peak of activity overlapping with the activity of their

predator, the coyote (*Canis latrans*), supporting the idea that these prey species are able to

shift their activity with a temporary mismatch that allows them to reduce predation risk

(Arias-Del Razo et al. 2011).

Regarding the differences between the plasticity of the activity patterns of the red fox

and the Egyptian mongoose, they may be due to differences in foraging strategies. The

European rabbit is a profitable prey for the generalist mesopredators in Doñana (Rau 1987,

Palomares and Delibes 1991b, Martín et al. 1995, Ferreras et al. 2011) but each predator

displays different hunting and consumption strategies of this prey item. Egyptian mongooses

and European badgers are species that preferentially consume younger rabbit age classes

since their foraging strategy consists of digging in burrows even though they can actively

hunt them (Palomares and Delibes 1991b, Martín et al. 1995). Conversely, the red fox resorts

to a foraging strategy based mainly on the pursuit of prey items, although it also excavates in

rabbit burrows (Fedriani et al. 1999). Thus, these distinct foraging strategies could allow the

Egyptian mongoose and European badgers to maintain a temporal asynchrony with the prey's

activity patterns while the red fox would be forced to track and match the activity pattern of

its prey.

The observational nature of this study does not allow us to establish a clear causal

relationship in the observed patterns of the different species (i.e., temporal avoidance

between competitors). Future research should address differences of rabbit availability for

predators through the study of their respective abundances among study sites, as well as the

relative abundances of each mesopredator by accounting for the probability of detection in

occupancy models. Unfortunately, our sample size did not allow us to fit these models. In

these situations, camera trap data could be complemented with other census methods (i.e.

pellet counts or track censuses). Complementarily, the study of the trophic relationships

between potential competitors could be evaluated by studying the diet composition through

the collection, identification and analysis of faecal samples in the different study sites

simultaneously to the camera-trapping surveys. Nevertheless, our study addresses for the first

time the interspecific relationships among the 4 species of mesocarnivores present in Doñana

National Park by integrating together the different patterns of daily activity and their

respective overlap, as well their trophic ecologies, since previous efforts have focused on

niche differentiation along other axes (spatial; (Palomares and Delibes 1991c, Palomares et

al. 1996, Soto and Palomares 2015); and trophic, (Palomares and Delibes 1991a, Fedriani et

al. 1999, Ferreras et al. 2011)).

Conclusions

The activity patterns of mesopredators are ultimately restricted by the circadian

boundaries, but at the same time can be slightly modulated by interspecific interactions with

both prey and potential competitors. Our results show that the coexistence of the different

species that compose the mesocarnivore assemblage in the Doñana National Park can be

facilitated by the temporal segregation among generalist species due to a divergence in the

activity patterns (nocturnal, diurnal and cathemeral) but also by differences in trophic

preferences among specialist species, as suggested in previous studies (Palomares and

Delibes 1991c, Fedriani et al. 1999, Soto and Palomares 2015).

Our study further suggests that in order to clarify the coexistence mechanisms

underlying community structure, it is necessary to consider a multidimensional perspective,

addressing simultaneously the differences in multiple axes of the ecological niches. In

markedly seasonal habitats like Doñana, with patchy vegetation including ample gradients of

vegetation cover and successional status, even subtle differences in daily activity such as

those documented here represent multiple ways to facilitate sympatry of coexisting

mesopredator species. These effects might prove crucial for such coexistence in settings

where the loss of apex predators may lead to mesopredator release, triggering increased

densities and resource overlap. Recent scenarios of rewilding with top predators may

represent ideal conditions to experimentally examine these effects of mesopredator release in

biodiverse carnivore guilds like those of the Mediterranean habitats studied here.

Acknowledgements

We thank Juan Pedro González Varó, Tamara Burgos, Elena Quintero, Carlos Gutierrez,

Rubén Tarifa and José Luís Molina for technical support regarding species identification. We

are also grateful to Irene Mendoza and Blanca Arroyo for helpful comments and suggestions

on early versions of the manuscript. Our work was funded by the project DISPMULTLAYER

(CGL2017-82847-P), Spanish Ministry of Science and Innovation (PJ). JI was supported by

the FPI grant PRE2018-085916 from the Spanish Ministry of Science and Innovation. ABL

was supported by a Juan de la Cierva-Incorporación grant (IJCI-2017-31419), Spanish

Ministry of Science, Innovation and Universities. We appreciate the support from ICTS-RBD

facilities during fieldwork.

Author contributions

ABL conceived the original idea; IF performed the analyses, designed the figures and led the

writing of the manuscript; JI and PJ designed the methodology; JI and IF compiled the camera-

trap data and IF collected the diet data. All authors contributed critically to subsequent drafts

and the final version of the manuscript.

Data and code accessibility

All data and R scripts used in this study are available in the GitHub repository:

https://github.com/IagoFerreiroArias/TempPart

References

References

Agostinelli, C., and U. Lund. 2017. {R} package \texttt{circular}: Circular Statistics

(version 0.4-93). CA: Department of Environmental Sciences, Informatics and

Statistics, Ca' Foscari University, Venice, Italy. UL: Department of Statistics,

California Polytechnic State University, San Luis Obispo, California, USA.

Arias-Del Razo, I., L. Hernández, J. W. Laundré, and O. Myers. 2011. Do predator and

prey foraging activity patterns match? A study of coyotes (*Canis latrans*), and

lagomorphs (*Lepus californicus* and *Sylvilagus audobonii*). Journal of Arid

Environments 75:112–118.

Baker, P. J., C. V Dowding, S. E. Molony, P. C. L. White, and S. Harris. 2007. Activity

patterns of urban red foxes (*Vulpes vulpes*) reduce the risk of traffic-induced

mortality. Behavioral ecology 18:716–724.

Barrientos, R., and E. Virgós. 2006. Reduction of potential food interference in two

sympatric carnivores by sequential use of shared resources. Acta Oecologica

30:107–116.

Barrull, J., I. Mate, J. Ruiz-Olmo, J. G. Casanovas, J. Gosàlbez, and M. Salicrú. 2014.

Factors and mechanisms that explain coexistence in a Mediterranean carnivore

assemblage: an integrated study based on camera trapping and diet. Mammalian

Biology 79:123–131.

Batschelet, E. 1981. Circular statistics in biology. Academic Press, New York.

Bennie, J. J., J. P. Duffy, R. Inger, and K. J. Gaston. 2014. Biogeography of time

partitioning in mammals. Proceedings of the National Academy of Sciences of the

United States of America 111:13727–13732.

Brashares, J. S., L. R. Prugh, C. J. Stoner, and C. W. Epps. 2010. Ecological and

conservation implications of mesopredator release. Pages 221–240 Trophic

cascades: predators, prey, and the changing dynamics of nature.

Bu, H., F. Wang, W. J. McShea, Z. Lu, D. Wang, and S. Li. 2016. Spatial Co-occurrence

and activity patterns of mesocarnivores in the temperate forests of Southwest

China. PLoS ONE 11:e0164271.

Camps, D. 2008. Activity patterns of adult common genets *Genetta genetta* (Linnaeus,

1758) in northeastern Spain. Galemys 20:47–60.

Cavallini, P., and S. Lovari. 1994. Home range, habitat selection and activity of the red

fox in a Mediterranean coastal ecotone. Acta Theriologica 39:279–287.

Chesson, P. 2000. Mechanisms of maintenance of species diversity. Annual Review of

Ecology and Systematics 31:343–366.

Davis, M. L., M. J. Kelly, and D. F. Stauffer. 2011. Carnivore co-existence and habitat

use in the Mountain Pine Ridge Forest Reserve, Belize. Animal Conservation

14:56–65.

Dellafiore, C. M., C. Rouco, S. M. Vallés, and J. B. G. Fernández. 2014. Seasonal

habitat use by the European rabbit (*Oryctolagus cuniculus*) in a coastal dune system

in SW Spain. Animal Biodiversity and Conservation 2:233–242.

Di Bitetti, M. S., Y. E. Di Blanco, J. A. Pereira, A. Paviolo, and I. J. Pírez. 2009. Time

Partitioning Favors the Coexistence of Sympatric Crab-Eating Foxes (*Cerdocyon*

thous) and Pampas Foxes (*Lycalopex gymnocercus*). Journal of Mammalogy

90:479–490.

Díaz-Ruiz, F., J. Caro, M. Delibes-Mateos, B. Arroyo, and P. Ferreras. 2016. Drivers of

580   red fox (*Vulpes vulpes*) daily activity: prey availability, human disturbance or

581   habitat structure? Journal of Zoology 298:128–138.

582   Ensing, E. P., S. Ciuti, F. A. L. M. De Wijs, D. H. Lentferink, A. Hoedt, M. S. Boyce,

583   and R. A. Hut. 2014. GPS based daily activity patterns in european red deer and

584   North American elk (*Cervus elaphus*): Indication for a weak circadian clock in

585   ungulates. PLoS ONE 9:e106997.

586   Fedriani, J. M. 1996. Dieta anual del zorro, *Vulpes vulpes*, en dos hábitats del Parque

587   Nacional de Doñana. Doñana Acta Vertebrata 23:143–152.

588   Fedriani, J. M., P. Ferreras, and M. Delibes. 1998. Dietary response of the Eurasian

589   badger, *Meles meles*, to a decline of its main prey in the Donana National Park.

590   Journal of Zoology 245:214–128.

591   Fedriani, J. M., F. Palomares, and M. Delibes. 1999. Niche relations among three

592   sympatric Mediterranean carnivores. Oecologia 121:138–148.

593   Fenn, M. G. P., and D. W. Macdonald. 1995. Use of middens by red foxes: risk reverses

594   rhythms of rats. Journal of Mammalogy 76:130–136.

595   Ferreras, P., A. Travaini, S. C. Zapata, and M. Delibes. 2011. Short-term responses of

596   mammalian carnivores to a sudden collapse of rabbits in Mediterranean Spain.

597   Basic and Applied Ecology 12:116–124.

Fisher, N. I. 1995. Statistical analysis of circular data. Cambridge University Press.

Foster, V. C., P. Sarmento, R. Sollmann, N. Tôrres, A. T. A. Jácomo, N. Negrões, C.

Fonseca, and L. Silveira. 2013. Jaguar and puma activity patterns and predator-prey

interactions in four Brazilian biomes. Biotropica 45:373–379.

Gaynor, K. M., C. E. Hojnowski, N. H. Carter, and J. S. Brashares. 2018. The influence

of human disturbance on wildlife nocturnality. Science 360:1232–1235.

Gerber, B. D., S. M. Karpanty, and J. Randrianantenaina. 2012. Activity patterns of

carnivores in the rain forests of Madagascar: implications for species coexistence.

Journal of Mammalogy 93:667–676.

Holt, R. D., J. Grover, and D. Tilman. 1994. Simple rules for interspecific dominance in

systems with exploitative and apparent competition. The American Naturalist

144:741–771.

Jiménez, J., J. C. Nuñez-Arjona, F. Mougeot, P. Ferreras, L. M. González, F. García-

Domínguez, J. Muñoz-Igualada, M. J. Palacios, S. Pla, C. Rueda, F. Villaespesa, F.

Nájera, F. Palomares, and J. V. López-Bao. 2019. Restoring apex predators can

reduce mesopredator abundances. Biological Conservation 238:108234.

Kronfeld-Schor, N., and T. Dayan. 2003. Partitioning of time as an ecological resource.

Annual review of ecology, evolution, and systematics 34:153–181.

Lê, S., J. Josse, and F. Husson. 2008. {FactoMineR}: A Package for Multivariate

Analysis. Journal of Statistical Software 25:1–18.

Lima, S. L., and P. A. Bednekoff. 1999. Temporal variation in danger drives

antipredator behavior: the predation risk allocation hypothesis. The American

Naturalist 153:649–659.

Linkie, M., and M. S. Ridout. 2011. Assessing tiger–prey interactions in Sumatran

rainforests. Journal of Zoology 284:224–229.

Linnell, J. D. C., and O. Strand. 2000. Interference interactions, co-existence and

conservation of mammalian carnivores. Diversity and Distributions 6:169–176.

Lode, T. 1995. Activity pattern of polecats *Mustela putorius* L. in relation to food habits

and prey activity. Ethology 100:295–308.

MacArthur, R. H., and E. R. Pianka. 1966. On optimal use of a patchy environment. The

American Naturalist 100:603–609.

Maloney, S. K., G. Moss, T. Cartmell, and D. Mitchell. 2005. Alteration in diel activity

patterns as a thermoregulatory strategy in black wildebeest (*Connochaetes gnou*).

Journal of Comparative Physiology A 191:1055–1064.

Martín, R., A. Rodríguez, and M. Delibes. 1995. Local feeding specialization by badgers

(*Meles meles*) in a Mediterranean environment. Oecologia 101:45–50.

Monterroso, P., P. C. Alves, and P. Ferreras. 2013. Catch me if you can: Diel activity

patterns of mammalian prey and predators. Ethology 119:1044–1056.

Monterroso, P., P. C. Alves, and P. Ferreras. 2014. Plasticity in circadian activity

patterns of mesocarnivores in Southwestern Europe: Implications for species

coexistence. Behavioral Ecology and Sociobiology 68:1403–1417.

Monterroso, P., F. Díaz-Ruíz, P. M. Lukacs, P. C. Alves, and P. Ferreras. 2020.

Ecological traits and the spatial structure of competitive coexistence among

carnivores. Ecology 101:e03059.

Monzón, A., P. Fernandes, and N. Rodrigues. 2004. Vegetation structure descriptors

regulating the presence of wild rabbit in the National Park of Peneda-Gerês,

Portugal. European Journal of Wildlife Research 50:1–6.

Newsome, T. M., A. C. Greenville, D. Ćirović, C. R. Dickman, C. N. Johnson, M.

Krofel, M. Letnic, W. J. Ripple, E. G. Ritchie, S. Stoyanov, and A. J. Wirsing.

2017. Top predators constrain mesopredator distributions. Nature Communications

8:15469.

Palomares, F. 1993. Opportunistic feeding of the Egyptian mongoose, *Herpertes*

ichneumon,(L.) in Southwestern Spain. Revue d'écologie 48:295–304.

Palomares, F., and T. M. Caro. 1999. Interspecific killing among mammalian carnivores.

The American Naturalist 153:492–508.

Palomares, F., and M. Delibes. 1991a. Alimentación del meloncillo *Herpestes*

ichneumon y de la gineta *Genetta genetta* en la Reserva Biológica de Doñana, SO

de la Península Ibérica. Doñana Acta Vertebrata 18:5–20.

Palomares, F., and M. Delibes. 1991b. Dieta del meloncillo (*Herpestes ichneumon*), en

el Coto del Rey (Norte del Parque Nacional de Doñana, SO España). Doñana: Acta

Vertebrata 18:197–194.

Palomares, F., and M. Delibes. 1991c. Ecología comparada de la gineta *Genetta genetta*

(L.) y el meloncillo *Herpestes ichneumon* (L.)(Mammalia, Viverridae) en Doñana

(SO de la Península Ibérica). Boletín de la Real Sociedad Española de Historia

Natural (Sección Biología) 87:257–266.

Palomares, F., and M. Delibes. 1992. Circadian Activity Patterns of Free-Ranging Large

664   Gray Mongooses, *Herpestes ichneumon*, in Southwestern Spain. Journal of

665   Mammalogy 73:173–177.

666   Palomares, F., and M. Delibes. 1994. Spatio-temporal ecology and behavior of European

667   genets in southwestern Spain. Journal of Mammalogy 75:714–724.

668   Palomares, F., P. Ferreras, J. M. Fedriani, and M. Delibes. 1996. Spatial relationships

669   between Iberian lynx and other carnivores in an area of south-western Spain.

670   Journal of Applied Ecology 33:5–13.

671   Palomares, F., P. Ferreras, A. Travaini, and M. Delibes. 1998. Co-existence between

672   Iberian lynx and Egyptian mongooses: estimating interaction strength by structural

673   equation modelling and testing by an observational study. Journal of Animal

674   Ecology 67:967–978.

675   Polis, G. A., C. A. Myers, and R. D. Holt. 1989. The ecology and evolution of intraguild

676   predation: potential competitors that eat each other. Annual review of ecology and

677   systematics 20:297–330.

678   Prugh, L. R., C. J. Stoner, C. W. Epps, W. T. Bean, W. J. Ripple, A. S. Laliberte, and J.

679   S. Brashares. 2009. The rise of the mesopredator. Bioscience 59:779–791.

680   Pyke, G. H., H. R. Pulliam, and E. L. Charnov. 1977. Optimal foraging: a selective

681   review of theory and tests. The quarterly review of biology 52:137–154.

682   R Core Team. 2020. R: A Language and Environment for Statistical Computing. R

683   Foundation for Statistical Computing, Vienna, Austria.

684 Rao, J. S. 1976. Some tests based on arc-lengths for the circle. Sankhyā: The Indian

685   Journal of Statistics, Series B:329–338.

686 Rau, J. R. 1987. Ecología del zorro," *Vulpes vulpes* (L.)", en la reserva biológica de

687   Doñana, SO de España. Universidad de Sevilla.

688 Revilla, E., and F. Palomares. 2002. Does local feeding specialization exist in Eurasian

689   badgers? Canadian Journal of Zoology 80:83–93.

690 Ridout, M. S., and M. Linkie. 2009. Estimating overlap of daily activity patterns from

691   camera trap data. Journal of Agricultural, Biological, and Environmental Statistics

692   14:322–337.

693 Rodríguez, A., R. Martin, and M. Delibes. 1996. Space use and activity in a

694   Mediterranean population of badgers *Meles meles*. Acta Theriologica 41:59–72.

695 Rowcliffe, J. M., R. Kays, B. Kranstauber, C. Carbone, and P. A. Jansen. 2014.

696   Quantifying levels of animal activity using camera trap data. Methods in Ecology

697   and Evolution 5:1170–1179.

698 Rowcliffe, M. 2019. activity: Animal Activity Statistics.

699 Santos, F., C. Carbone, O. R. Wearn, J. M. Rowcliffe, S. Espinosa, M. G. M. Lima, J. A.

700   Ahumada, A. L. S. Gonçalves, L. C. Trevelin, and P. Alvarez-Loayza. 2019. Prey

701   availability and temporal partitioning modulate felid coexistence in Neotropical

702   forests. PloS one 14:e0213671.

703 van Schaik, C. P., and M. Griffiths. 1996. Activity Periods of Indonesian Rain Forest

704   Mammals. Biotropica 28:105–112.

Schoener, T. W. 1974. The compression hypothesis and temporal resource partitioning.

Proceedings of the National Academy of Sciences of the United States of America

71:4169–4172.

Soto, C., and F. Palomares. 2015. Coexistence of sympatric carnivores in relatively

homogeneous Mediterranean landscapes: functional importance of habitat

segregation at the fine-scale level. Oecologia 179:223–235.

Soulé, M. E., D. T. Bolger, A. C. Alberts, J. Wrights, M. Sorice, and S. Hill. 1988.

Reconstructed Dynamics of Rapid Extinctions of Chaparral-Requiring Birds in

Urban Habitat Islands. Conservation Biology 2:75–92.

Terborgh, J., R. D. Holt, and J. Estes. 2010. Trophic cascades: what they are, how they

work, and why they matter. Page Trophic Cascades: Predators, Prey and the

Changing Dynamics of Nature.

Valverde, J. A. 1957. Notes écologiques sur le lynx d'Espagne feus Lynx Pardina

temminck. Terre et Vie (Revue d'Ecologie) 104:51–67.

Valverde, J. A. 1967. Estructura de una comunidad mediterránea de vertebrados

terrestres. Consejo Superior de Investigaciones Científicas (España), Madrid.

Vazquez, C., J. M. Rowcliffe, K. Spoelstra, and P. A. Jansen. 2019. Comparing diel

activity patterns of wildlife across latitudes and seasons: Time transformations

using day length. Methods in Ecology and Evolution 10:2057–2066.

Vilella, M., M. Ferrandiz-Rovira, and F. Sayol. 2020. Coexistence of predators in time:

Effects of season and prey availability on species activity within a Mediterranean

carnivore guild. Ecology and Evolution 00:1–15.

Villafuerte, R., M. B. Kufner, M. Delibes, and S. Moreno. 1993. Environmental factors

influencing the seasonal daily activity of the European rabbit (*Oryctolagus*

cuniculus) in a Mediterranean area. Mammalia 57:341–347.

Virgós, E., M. Llorente, and Y. Cortésá. 1999. Geographical variation in genet (*Genetta*

genetta L.) diet: a literature review. Mammal Review 29:117–126.

Wang, Y., and D. O. Fisher. 2012. Dingoes affect activity of feral cats, but do not

exclude them from the habitat of an endangered macropod. Wildlife Research

39:611–620.

Wickham, H., R. François, L. Henry, and K. Müller. 2020. dplyr: A Grammar of Data
Manipulation.

Zielinski, W. J. 1986. Circadian rhythms of small carnivores and the effect of restricted

feeding on daily activity. Physiology & behavior 38:613–620.

Figures

Figure 1. Niche segregation along the trophic axis among mesocarnivores in Doñana National

Park. Convex polygons represent trophic niche breadth for generalist species. Since only one

diet study could be used regarding common genet diet in the study area, convex polygons are

omitted for this species. The CA first two dimensions (Dim 1, Dim 2) explain 66% of the

variability in diet composition among species and studies. The first dimension separates

mesocarnivores into two groups: species whose diet is predominantly carnivorous and species

whose diet is omnivorous. The second dimension separates the Egyptian mongoose from the

genet while indicating a greater trophic overlap between the diets of the red fox and the

European badger.

758
759
760

Figure 2. Pairwise comparisons of daily activity patterns between mesocarnivores (rows, top to bottom: red fox, badger, genet, and Egyptian mongoose) and their main prey items in the study area (columns, left to right: *Oryctolagus cunniculus*, small mammals, and *Alectoris rufa*). Coloured lines describe prey activity patterns (blue for European rabbit, green for small mammals and yellow for red-legged partridge) and black lines indicate predator's activity patterns. Overlap coefficients (Δ) and their respective 99% confidence intervals are shown in brackets.

775   Figure 3. Diel activity patterns of two mesocarnivores (red fox and Egyptian mongoose; mid

776   and bottom rows, respectively) and its main prey item (European rabbit, top row) along the

777   gradient of ecological succession, from younger juniper stands (SABCOL) to mature, late

778   successional stands (SABMAR).

795 Figure 4. Kernel density plots and temporal overlap between mesocarnivores in the study

796 area: common genet (green line), red fox (blue line), European badger (purple line) and

797 Egyptian mongoose (yellow line). Overlap coefficients (Δ) and their respective 99%

798 confidence intervals are shown in brackets.

Supporting Information

Figure S1. Study area showing: A, location of Doñana National Park and Doñana Biological

Reserve (red rectangle); B, location of 3 sites selected (*Sabinar de Colonización - SABCOL,*

Sabinar del Ojillo - SABOJI and *Sabinar del Marqués - SABMAR*) and C, location of camera

trapping stations (points) and set of cameras active for ca. 10 days during a given sampling

period.

Table S1. Summary of the studies used to carry out the correspondence analysis and explore

the trophic relationships between mesocarnivores. Repetition of the studies is due to the

different data coming from different areas and/or periods sampled within the same study.

Table S2. Number of detections (N) of target species and trap success measured as number of

detections of each species per 100 camera-days for each study site.

Table S3. Number of detections of target species across three sites selected. Dawn: 1 hour

before and after sunrise; Day: 1 hour after sunrise until 1 hour before sunset; Dusk: 1 hour

before and after sunset; Night: 1 hour after sunset and 1 hour before sunrise. We set sunrise

as 8:00:00 and sunset as 20:00:00. Rao Spacing Test significance: p-value <0,01 **; p value

<0,05 *. Sampling effort per site reported as mean and standard deviation ($\bar{x} \pm$ SD) of

camera-days.

Landscape maturity

SABCOL   SABOJI   SABMAR

Density

Time