Title: Dorsomedial prefrontal neural ensembles reflect changes in task utility that culminate in task quitting

Abbreviated title: dmPFC activity corresponds to task utility

Authors: Blake S. Porter and Kristin L. Hillman

Department of Psychology and Brain Health Research Centre

University of Otago

275 Leith Walk, Dunedin 9016, New Zealand

Corresponding Author: Blake S. Porter, blakeporterneuro@gmail.com

Pages: 41

Figures: 8

Tables: 0

Word count Abstract: 198

Word count Introduction: 668

Word count Discussion: 1,401

Conflict of Interest: The authors declare no competing financial interests.

Funding: This study was supported by Marsden Fund grant U001617 (K.L.H.) from the Royal Society of New Zealand Te Apārangi.

Abstract

When performing a physically demanding behavior, sometimes the optimal choice is

to quit the behavior rather than persist in order to minimize energy expenditure for the

benefits gained. The dorsomedial prefrontal cortex (dmPFC), consisting of the anterior

cingulate cortex and secondary motor area, likely contributes towards such utility

assessments. Here, we examined how male rat dmPFC single unit and ensemble level activity

corresponded to changes in task utility and quitting in an effortful weight lifting task. Rats

carried out two task paradigms: one that became progressively more physically demanding

over time and a second fixed effort version. Rats could quit the task at any time. Dorsomedial

PFC neurons were highly responsive to each behavioral stage of the task, consisting of rope

pulling, reward retrieval, and reward area leaving. Activity was highest early in sessions,

commensurate with the highest relative task utility, then decreased until the point of quitting.

Neural ensembles consistently represented the sequential behavioral phases of the task.

However, these representations were modified over time and became more distinct over the

course of the session. These results suggest that dmPFC neurons represent behavioral states

that are dynamically modified as behaviors lose their utility, culminating in task quitting.

Key Words: prefrontal cortex, effort, quitting, motivation, decision making

New & Noteworthy

When carrying out a physically demanding task animals must continually assess whether to

persist or quit. In this study we recorded neurons in the dorsomedial prefrontal cortex

(dmPFC) of rats as they carried out a challenging weightlifting task, up to the point of

quitting. We demonstrate that dmPFC neurons form a representation of the task that is

modified, via a decrease in firing rate, by the decreasing utility of the task that may signal

quitting.

Introduction

When engaged in a certain behavior, animals must continuously assess whether to

persist in that behavior or change course. During foraging, for example, animals face

recurring decisions as to whether to stay in their current patch of diminishing food or to incur

the costs of exploration to search for a more abundant patch (1, 2). A third option also exists:

rather than adjusting the current course of action, an animal could quit the behavior all

together. The decision to quit may be driven by a multitude of factors including physiological

states (e.g., satiation, fatigue), the cognitive assessment of diminishing utility, and inter-

individual trait variability. However, the neurophysiological processes that underlie quitting

behaviors are not well understood.

The prefrontal cortex is likely a critical region in quitting processes due to its role in

encoding action-outcome contingencies and using these associations to drive goal-directed

behaviors (3–6). In particular, the anterior cingulate cortex (ACC or Anterior Cg1 in rodents)

plays a specialized role in evaluating the physical effort costs associated with certain

behaviors; this enables net utility determinations and behavioral motivation towards difficult

but worthwhile goals (7–10). The ACC is well connected to sensorimotor areas which may

translate decisions into actions (11–14).

The secondary motor area (M2) is laterally adjacent to Cg1 in rodents and is

considered a higher order motor area that participates in action plan selection based on utility

(9, 15–18). Together, these two regions in the rat dorsomedial prefrontal cortex (dmPFC) –

the anterior Cg1 and M2 – demonstrate important roles in determining the utility and

selection of behaviors that maximize benefit to the animal. However, it is not well-

understood how neurons in these areas respond to a prolonged task where quitting eventually

becomes the most beneficial action. In such a scenario, the continued time and energy costs

to pursue the challenging behavior are no longer worth the payout (e.g., calories gained).

In order to investigate quitting behaviors we recently developed an effortful weight

lifting task (WLT) for use with laboratory rats (19). To date, previous electrophysiological

studies on cost-benefit decision making have usually not been structured to allow a subject to

quit outright or, if subjects do quit, the quitting behavior may not be studied in-depth.

Generally, subjects are provided with different cost-benefit actions (e.g., operant actions

involving variable effort in rodent studies, (20–22) or cost-benefit contingencies (e.g.,

stimulus-outcome pictorials in primate studies, (23)) to choose between, and subjects make

recursive decisions over a fixed number of trials or a fixed duration. If subjects do not carry

out the requisite number of trials or duration, these data may not be analyzed. In our WLT,

rats are challenged to pull a weight-adjustable rope a fixed distance in order to earn rewards.

Weight-adjustment allows for dynamic control over the effort associated with the rope-

pulling behavior. Furthermore, the animals were free to quit the task at any point.

Here we recorded single units from the dmPFC while rats carried out the WLT in

order to characterize how neural activity patterns relate to quitting behaviors. Two different

paradigms of the WLT were used: a progressive weight (PW) version where the weight

increased over time, and a fixed weight (FW) version that used a constant, moderate weight.

In the PW version, task utility declines due to the increasing costs (weight) while task utility

in the FW version declines due to the internal states of the animal (e.g., fatigue, satiation).

We predicted that dmPFC neurons would be responsive to our effort-based task in that

different neurons should represent the actions, outcomes, and salient behavior transitions

associated with performing the task. Furthermore, these representations should be modulated

by changes in utility, brought about by heavier weight-adjustments and/or fatigue. We

predicted that Cg1 neurons would be more responsive during reward phases to monitor

action-outcome contingencies, while M2 neurons would be more responsive to rope pulling

and associated effort. Lastly, we predicted that neural populations in the dmPFC would

undergo state transitions that reflect the rats' behavioral transitions towards the point of

quitting.

Materials and Methods

Subjects

Male Sprague-Dawley rats (n=6, weighing between 400 - 480 g at the time of

experiment; Hercus Taieri Resource Unit, New Zealand) were single housed in individually

ventilated cages measuring 38 x 30 x 35 cm (Tecniplast, Italy). Animals were maintained on

a 12 h reverse light cycle (07:00 to 19:00), with experimentation occurring during the

animals' dark, active phase. Water was available *ad libitum* throughout training and

experimentation. Daily chow (Teklad diet; Envigo, USA) was rationed to promote food

motivation for training and during experimentation. Rats were weighed twice per week and

daily food rations were adjusted to maintain each animal's body weight $\geq 85\%$ of their free-

feeding body weight. All procedures were approved by the University of Otago's Animal

Ethics Committee, protocol 91/17.

Training

Prior to surgery, animals were trained in the WLT as previously described (19).

Briefly, the WLT is performed in a 120 x 90 x 60 cm arena (Figure 1a). An animal is

progressively shaped to pull a rope 30 cm to trigger the dispensing of 0.25 mL of 20%

sucrose solution. The rope can be weighted from 0 to 225 g to increase the effort demands of

the task, however only 0 g and 45 g are used in the training phase. Once an animal can

perform 20 successful pulls (10 pulls on 0 g followed immediately by 10 pulls on 45 g) in

under 10 minutes, they are considered trained in the WLT and ready for surgery. Rats took an

average of 26.3 days (Standard Deviation (S.D.) = 0.7 days) to reach the 20 successful trial

criterion.

Microdrives

All rats were implanted with Halo-10 Microdrives (Neuralynx, USA) consisting of eight

independently drivable tetrodes. Tetrodes were made from 18 micron Nichrome Formvar

insulated wire (A-M Systems, Inc., USA). Tetrodes were gold plated over three to five days

using a nanoZ (White Matter LLC, USA) such that electrode impedances were 200-300

kOhms prior to surgery. Tetrodes were secured to Neuralynx EIB-36's via gold pins to

interface with our headstage (see *Data collection*, below). Neuralynx Standard A exit tips

were used and tetrodes were positioned in the exit tip to minimize medial-lateral spread when

implanted in the brain.

Surgery

Anesthesia was induced with 5% Isoflurane in oxygen and then maintained at 2-3%

throughout the surgery. Rats were mounted in a Kopf (David Kopf Instruments, USA)

stereotaxic frame with non-puncture ear bars. One mL of 2% Lidocaine diluted in 4 mL of

0.9% saline was injected subcutaneously above the skull. A midline incision was made and

the skin over the skull retracted. A 3 mm diameter craniotomy was made centered at + 2.5

mm anterior-posterior and ± 0.7 mm medial-lateral from Bregma (Paxinos and Watson,

2007). Five rats were implanted on the right hemisphere and one on the left hemisphere. The

dura was removed and the microdrive was lowered onto the surface of the brain via a

stereotaxic arm such that the most medial tetrode was next to the central venous sinus. Two 4

mm long, 1 mm wide stainless steel screws, bilaterally placed over the cerebellum, served as

grounds. Six to eight additional 3 mm long, 1mm wide stainless steel screws were then put

partially into the skull and the implant was secured to the skull and skull screws with

ultraviolet cure self-etching dental cement (Maxcem Elite™, Kerr Corp., USA). Sutures were

used to close the skin around the implants.

Post-operatively (post-op), rats were subcutaneously administered prophylactic

antibiotic (30mg/kg of Amphoprim) and an anti-inflammatory (5 mg/kg of Carprofen) and

recovered in a heated cage until they regained consciousness and were active. Rats were then

returned to a clean home cage to recover for 10 days. The anti-inflammatory was

administered for a minimum of two additional days post-op and as needed if rats showed any

signs of pain. After the 10 days of post-op recovery, rats were re-introduced to the WLT and

tested to criterion to ensure that they could still carry out the task and habituate to their

implants. Rats took an average of 9.2 (S.D. = 4.9) days to return to criterion. Once rats

reestablished criterion performance, experimentation started.

Data collection

Single unit data was collected using a Neuralynx Digital Lynx SX system with

Cheetah software (version 6.3.2) via a Neuralynx HS-36-LED pre-amplifier headstage

connected to a Neuralynx Saturn commutator. Data from the 32 electrodes were sampled at

32 kHz and bandpass filtered between 600 and 6,000 Hz. Every tetrode was referenced to an

electrode on a different tetrode with no spiking activity. Single unit action potentials were

detected online with a threshold of 65 µV and stored for offline spike sorting. Spikes were

sorted offline using Plexon (USA) Offline Sorter (version 4) by primarily relying on peak-

valley amplitude and principal component analysis. Only well-isolated clusters of waveforms,

likely to be from the same neurons, were included for analyses. Tetrodes were not moved

day-to-day if they had single units on them. If a tetrode did not have any single units it was

advanced approximately 32 microns at the end of a recording session.

Each headstage contained two LEDs which enabled the animal's position to be

tracked by Neuralynx's Cheetah software via an overhead camera (JAI Inc., USA). Video

was sampled at 25 frames per second. An Arduino Uno (www.arduino.cc) was used to detect

rope pulls and to dispense sucrose solution reward; TTL signals of all events were sent to the

Digital Lynx SX system for timestamping and storage.

Progressive weight paradigm

Rats completed two WLT paradigms: a progressive weight (PW) paradigm and a

fixed weight (FW) paradigm (19). In the PW paradigm, a session started with a two minute

pre-baseline period where the rat was placed in the WLT arena but the rope was not extended

into the environment. After the pre-baseline, the rope was extended with 0 g of weight

attached and the rat could begin earning rewards; each successful 30 cm pull dispensed 0.2

mL of 20% sucrose solution. After every 10 successful (rewarded) pulls of the rope, 45 g of

weight was added to the rope until the rat quit or 225 g was reached. If a rat completed 10

trials of 225 g, a clip was placed on the rope in order to prevent the rat from pulling the

weight high enough to trigger reward dispensing. However, no rats in this study completed 10

trials of 225 g.

Quitting was defined as the rat making no attempts to pull the rope for two full

minutes. When a rat quit the rope was retracted from the apparatus at the end of the two

minute 'quit period', and a two minute post-baseline period started. Lastly, after the post-

baseline period, two sucrose rewards were manually dispensed to check if the rats were

satiated or not. Rats had one minute to consume the sucrose reward once dispensed. Every rat

on every session consumed these manually dispensed rewards, suggesting that satiation was

not occurring. After the sucrose satiation check the recording session was stopped and the

animal was returned to its home cage. Rats ran one session per day.

Fixed weight paradigm

In the FW paradigm, a session again started with a two minute pre-baseline period,

followed by the rope being extended into the arena with 0 g of weight attached. After 10

successful (rewarded) pulls on 0 g, the rope weight was immediately switched to a fixed

weight, calibrated to each specific rat. The initial fixed weight was the maximum weight that

the rat had completed in the PW paradigm (i.e., the weight on which they had successfully

performed 10 rewarded pulls). Rats were allowed to complete as many trials as they wanted

on the FW until they quit or until 60 minutes elapsed, whichever came first. After 2-3 days

(average 2.7 days, S.D. = 0.4 days), the weight was adjusted to be 45 g heavier or lighter

depending on the rats' performance: if rats were working for fewer than 15 minutes their

weight was reduced whereas if rats were working for more than 15 minutes their weight was

increased. Rats then carried out the adjusted FW paradigm for 2-5 further days (average 3.2

days, S.D. = 1.0 days) depending on their performance and the number of single units being

recorded.

Overall analyses

All analyses were carried out using Matlab (version R2018a; The MathWorks Inc.,

USA). Colorbrewer (https://github.com/axismaps/colorbrewer/) and mpl colormaps

(https://bids.github.io/colormap/) were used for graphical color maps.

Behavioral analyses

Behavioral metrics were calculated as previously reported (19). Briefly, for the PW

paradigm, the quit weight was calculated as the highest weight in a given session on which

the rat made at least one rope pulling attempt. For the FW paradigm, the quit duration was

calculated as the time from when the FW was attached (exclusive of 0 g) until the last pull

attempt.

For both the PW and FW paradigms, quartiles were determined based on the duration

of each individual session. For the PW paradigm, the duration was calculated starting from

when the 0 g weight was available until the rat quit. For the FW paradigm, quartiles were

calculated starting from the time the FW was available (thus excluding the initial period of 10

trials on 0 g). For all FW paradigm analyses this initial 0 g period was excluded so as to avoid

comparisons between 0 g-containing quartiles and FW-containing quartiles Thus, for

behavioral and neural analyses of the FW paradigm, every quartile has the same weight

attached to the rope. Every quartile had to contain at least five completed trials; recording

sessions that did not meet this minimum trial criterion were excluded from this manuscript.

The four quartiles are abbreviated throughout the manuscript as Q1, Q2, Q3, and Q4 to

denote the first, second, third, and fourth quartile.

Trials were defined as the sequence of events beginning with a rope pull attempt

followed by reward area entry and concluding with reward area leaving. This provided three

task phases: the attempt phase, the reward entry phase, and the reward leaving phase (Figure

1b). The attempt phase consisted of 1.5 seconds before and 2.0 seconds after a pull attempt.

Pull attempts were timestamped by a TTL code from the Arduino that logged when the

weight was lifted. Reward entry and reward leaving were based on the tracking data obtained

from LEDs on the headstage. A 20 x 20 cm area was manually placed around the reward cap

for each recording session based on the rat's tracking data and known cap location. Reward

entry was centered on when the rat entered the reward zone and their instantaneous speed

dropped below 15 cm/s. The reward entry phase consisted of 1.5 seconds before and 0.5

seconds after reward zone entry. Reward leaving consisted of 0.5 seconds before and 1.5

seconds after the point where the rat left the reward zone. Task-phase durations were chosen

based on the distribution of durations across all recording sessions in order to maximize

duration but minimize overlap between phases.

Trials were deemed "successful" when the rat pulled the weight sufficiently high

enough (30 cm) to trigger a reward, or deemed "failures" when the rat did not pull the weight

high enough but still went and sampled the reward cap. Events when the rat pulled the rope

but did not check the reward cap were excluded. Trials were also parsed according to whether

or not the rat "engaged" or "disengaged" from the task on the subsequent trial. Engagement

was determined based on how long it took from the rat leaving the reward cap until the next

pull attempt. If the rat made another attempt within 12 seconds the trial was considered an

engage trial; anything longer than 12 seconds was considered a disengage trial. Twelve

seconds was chosen as a cut off as 75% of all next-attempts occurred within 12 seconds of

leaving the reward cap.

The on-task ratio was calculated as the total time a rat spent working at the task

divided by the total time of the session when the rope was available. Rats were considered

working if they were both within the "work zone" and making attempts to pull the rope. The

work zone consisted of a 60 x 64 cm area encompassing the rope pulling area and the reward

cap. The reward interval was calculated as the time between earned rewards based on the

reward TTL signals. The success-to-attempt ratio was calculated as the number of successful

pull attempts divided by the total number of attempts made. For the on-task ratio and success-

to-attempt ratio we used a repeated-measures one-way ANOVA to test for significant

changes over the quartiles and Tukey-Kramer's test for post-hoc pairwise comparisons. Due

to some rats having no rewards in Q4, and thus no reward intervals, we instead used a

Kruskal-Wallis test to measure if reward interval times were affected by quartile and Tukey-

Kramer's test for post-hoc comparisons.

For every trial we calculated the animal's average speed as they traversed from the

attempt zone to the reward zone after an attempt was made. We carried out a one-way

ANOVA to determine if there were significant changes in speed over quartiles.

Neural analyses – single neurons

For every recording session, sorted spikes from Plexon Offline sorter were imported

into Matlab with the accompanying tracking and TTL data from that recording session.

Neurons were assigned a brain area based on what tetrode they were recorded from. We

tested whether or not neurons were task-responsive by randomly sampling every neuron's

firing rate across 10,000 two second epochs drawn from the entire recording session

(including the pre- and post-baselines). Neurons were considered to be task-responsive if,

during at least one task-phase, their firing rate was significantly different (Wilcoxon rank sum

test) to the random sample and they had a firing rate of at least 0.5 Hz. Furthermore, to avoid

including neurons that may have been lost or gained over the course of a recording session, a

neuron also had to have a non-zero firing rate in at least one task-phase for all four quartiles.

Neurons that did not meet this criteria were not included for analysis. Spike raster plots and

peri-event time histograms were created for task-responsive neurons for each of the three

task-phases across all trials. Spike times were aligned to the center of the task-phase event.

Histograms were created with 100 ms bins across each task-phase. For each task-phase, a

Chi-squared test against an equal distribution was used to test if there was an unequal

distribution of task-responsive neurons across brain areas.

Neurons were determined to be phasic or tonic based on how many bins of a trial they

fired an average of 2 standard deviations above their overall mean firing rate. Neurons that

fired for more than 2 bins but less than 10 bins were deemed phasic while all others were

deemed tonic. Neurons were considered to have an inhibitory response to the task phases if

they had an average response of less than -0.5 standard deviations for 10 or more bins.

Neurons that did not fit this criteria were deemed excitatory.

Neural analyses – population analyses

All task-responsive neurons from all sessions were pooled together for further

analysis. A sequence plot was created with all neurons where each neuron's binned trial

firing rates were z-scored then averaged across all trials in each 100 ms bin. All neurons were

then sorted based on the trial bin that contained their maximum firing rate. For graphical

purposes only, every neuron's activity was normalized to be between 0 and 1 where 1

corresponded to their maximum z-scored firing rate and 0 corresponded to their minimum z-

scored firing rate. All analysis was done on the z-scored firing rates. A cross-correlation

matrix was created using the sequence plot where the Pearson's correlation was calculated

pairwise for every 100 ms bin of a trial.

Principal component analysis (PCA) was carried out on the z-score averaged firing

rates of all task-responsive neuron in 100 ms bins across the trial using Matlab's pca function.

Trials were the same as described above, consisting of the 3.5 second attempt phase, 2.5

second reward entry, and 2.5 second reward leave phase for consistent behaviors and

durations. Thus PCA was carried out once on a matrix where every column was a neuron and

every row was the neuron's average z-scored firing rate for every 100 ms bin of a trial. PCA

was also carried out on the successful versus failed trials, engage versus disengage trials, and

quartiles. In these cases, each task-responsive neuron's firing rates were binned into 100 ms

bins, z-scored across all trial types, and then all neurons were pooled together. A matrix was

then constructed where every column was a neuron and every row was a 100 ms bin of a trial

for each trial type (successful and failed, engage and disengage, and each of the quartiles).

PCA was ran once on the matrix resulting in the same principal component space for all trial

types. We computed the distance between the PCA score trajectories by calculating the

distance between each 100 ms bin using the first three principal components. As a control, we

also randomly split trials of a recording session into halves (for engage versus disengage

comparisons, and success versus fail comparisons) or quarters (for quartile analyses) and

averaged a neuron's activity over the random trials. This was carried out 1,000 times per

session. For every iteration, all neurons were pooled together, PCA carried out, and the

distance between each 100 ms bin calculated. For the comparisons of engage versus

disengage and success versus fail, a paired t-test was used to determine if the observed PCA

distances were significantly different from the mean of the 1,000 random PCA score

distances. A one-way ANOVA, with Tukey-Kramer's post-hoc, was used to test whether or

not the observed PCA score distances across the three task-phases were significantly

different.

For graphical purposes only, the PCA score plots were smoothed with a 500 ms

moving average within each task-phase when comparing multiple trajectories. The PCA

trajectories of all trials from the PW and FW paradigms (Figures 4c and 7c) are plotted with

the raw scores. All distance calculations and statistics were done on the raw PCA scores.

Kruskal-Wallis tests were used on each task-responsive neuron to test firing rate

variability across quartiles for each task-phase. To keep trial numbers consistent across

sessions, we limited analysis to five trials in each quartile. For Q1, the first five trials were

used. For Q2 and Q3, the middle five trials were used. For Q4, the last five trials were used.

We then pooled the quartile averaged z-scored firing rates of all neurons with a significant (p

< 0.05) effect for quartile for each task-phase. For each task-phase we determined what

quartile each neuron was most active in based on their averaged fire rates in each quartile.

We carried out a Chi-squared test on the observed distribution of peak activity across the four

quartiles versus and equal distribution across all quartiles. Repeated-measures one-way

ANOVA with a post-hoc Tukey-Kramer's test was used on the populations of neurons for

each task-phase, with quartile as a factor.

As a control, we iteratively subsampled proportions of the total population of neurons

and repeated the PCA analyses. We carried this out 1,000 times for each subsampled

proportion and averaged the iterations together. The average PCA trajectories were visually

inspected and their pairwise distances were tested against distances from the above described

randomly sampled PCA distances.

Neuronal data was also broken up into two minute epochs across the entire recording

session, including the pre- and post-baseline epochs where the rope was not available to the

rat. Four more epochs consisted of the four quartiles. For quartile one, the first two minutes

of the 0 g (PW paradigm) or the fixed weight (FW paradigm) was used. For quartiles two and

three, the middle two minutes of the quartile were used. For quartile four, the two minutes

preceding the rats' final attempt were used. The quit epoch consisted of the two minutes

proceeding the rats' final attempt where the rope was still available but the rat was not

making attempts. For each task-responsive neuron a z-scored firing rate was calculated for

each two minute epoch and all neurons were pooled. Pairwise cross-correlations were

calculated using Pearson's correlation. Matlab's pdist function was used to calculate the

pairwise correlation distances between epochs. Matlab's linkage function was then used to

hierarchically cluster the correlation distances based on their average correlations. Matlab's

dendrogram function was used to plot the clustering.

As an additional control analyses, we tested whether session duration had an impact

on the changes in neural activity that we observed across quartiles. First, we tested if the

magnitude of change seen in neuron firing rates was influenced by time. If dmPFC neural

activity was changing due to time (24), we would expect a greater change in activity for

longer recording sessions. We calculated the difference in firing rate for each neuron from Q1

to Q4 for each task-phase and ran a linear regression with session duration as the explanatory

variable. We did this for all task-responsive neurons and then again for only task-responsive

neurons that had a significant effect of quartile.

We carried out a second control analysis for time where we again found the

difference between the mean firing rate of Q1 to Q4 in the 2 minute epoch periods. We again

fit a simple linear regression with time as the predictor.

Histology

After completion of all behavioral paradigms, rats were deeply anesthetized with

Isoflurane and transcardially perfused with 4% paraformaldehyde (PFA) in phosphate-

buffered saline (PBS). All tetrodes were wound up before the brain was removed to avoid

damage. Brains were then cryoprotected in 30% sucrose in 4% PFA PBS until they sunk.

Brains were sectioned at 40 μm using a Cryostat (CM1950, Lecia Biosystems, LLC.,

Germany). Brain sections were then mounted, stained with Thionine, and imaged with a light

microscope (Lecia Biosystems, LLC.). Tetrodes were mapped based on their tracks in

reference to (25) brain atlas, microdrive turn logs, and their relative locations in the

microdrive (Figure 1c).

Results

Progressive weight behavior

Rats first carried out the PW paradigm; here the rope weight started out at 0 g and was

increased by 45 g every 10 successful pulls, to a maximum weight of 225 g. Rats were free to

quit the task at any point. Six rats performed a total of 30 sessions of the PW task (3-7

sessions per rat; average 5.0 ± 1.7 S.D.). Consistent with our previous studies (19), rats were

more likely to quit on heavier weights (Figure 2a). No rat was able to complete 10 successful

pulls of 225 g. Due to the inter- and intra-individual variance of quit weights across sessions,

each session varied in time duration and number of trials. Sessions varied in length from 874

to 2,056 seconds (median = $1,589 \pm 373$). Across all sessions rats earned from 25 to 55

rewards (median = 47 ± 8). Because of this variability, we broke each session into quartiles

based on time and used those quartiles to analyze behavioral performance.

We computed a time-on-task ratio that reflected the proportion of time rats spent

engaged in the task. A ratio of 1 indicated that the rats were regularly making attempts to pull

the rope while a ratio of 0 reflected rats making no attempts. Across quartiles, rats spent

significantly less time on-task as the session progressed into the third and fourth quartiles (F

$(2.6, 76.4) = 177.0$, $p < 0.0001$; Figure 2b). When rats were on-task and working, it took

longer to earn rewards as the session progressed (H $(3, 107) = 56.0$, $p < 0.0001$; Figure 2c).

Increases in reward interval, notably in Q4, were likely in part due to an increasing number of

failed rope pulls as weights got heavier. Across quartiles, there was a significant decrease in

the success-to-attempt ratio ($F_{(2.4, 68.4)} = 88.1$, $p < 0.0001$; Figure 2d). Rats also slowed

down in their reward approach on the fourth quartile relative to the other three quartiles ($F_{(3)}$

$= 19$, $p < 0.0001$; Supplemental Figure S1a https://doi.org/10.6084/m9.figshare.14515644).

Fixed weight behavior

On the FW paradigm, the rope weight started out at 0 g and after 10 successful pulls

the rope weight was set to a moderate FW, calibrated to each rat (see Methods). Rats could

perform as many weighted rope pulls as desired, up to a maximum duration of 60 minutes.

Rats were free to quit the task at any point. Because we set a minimum trial number criterion

for data analysis inclusion (see Methods), only nine sessions of the FW task were available

for analysis (three rats performed these nine sessions, 2-4 sessions per rat; average 3.0 ± 1).

Quit durations ranged from 1,803 to 3,548 seconds (median $= 2,711 \pm 571$ seconds Figure 2e)

and rats earned from 24 to 106 rewards (median $= 81 \pm 27$ rewards) on the FW.

Similar to the PW paradigm, rats spent significantly less time on-task as the FW

sessions progressed ($F_{(1.7, 13.3)} = 47.5$, $p < 0.0001$; all pairwise comparisons to Q4, p's $<$

0.0001; Figure 2f). There was a main effect for quartile on reward interval ($H_{(3, 31)} = 9.8$, p

$= 0.02$; Figure 2g), with Q4 having a higher reward interval than Q3 ($p = 0.030$) and Q2 ($p =$

0.035). There was a significant main effect of quartile for the success-to-attempt ratio ($F_{(2.1,}$

$_{17.0)} = 5.9$, $p = 0.01$; Figure 2h). However, post-hoc pairwise analyses only showed modest

differences between quartiles: success-to-attempt ratio was significantly higher on Q2

compared to Q3 ($p = 0.041$) and Q4 ($p = 0.043$). Compared to behavior in the PW task, where

weights incrementally increased by 45 g every 10 trials, rats may have struggled with the FW

initially in Q1 where the weight suddenly increased from 0 g to the FW value greater than

135 g. This likely accounted for the lower success rate and greater reward interval observed

in Q1 in the FW task. There was no change in the rats' speed to leave the attempt zone and

approach the reward zone across quartiles (F (3) = 0.15, p = 0.93; Supplemental Figure S1b

https://doi.org/10.6084/m9.figshare.14515644).

Neuronal task selectivity – Progressive weight paradigm

For neuronal analysis we focused on three task-phases of interest: the attempt phase,

the reward entry phase, and the reward leaving phase (see Methods and Figure 1b). In the PW

paradigm, we recorded a total of 360 neurons across the 30 recording sessions. Based on

histological reconstruction (see Figure 1c), 123 of these neurons were from the Cg1, 93 were

between Cg1-M2, and 144 were from Motor cortices 2/1 (M2/1).

A variety of activity patterns were observed within and across task-phases as well as

over time (Figure 3a-h). Some neurons were especially active during the attempt phase (3a,

b). The Cg1 neuron shown in Figure 3a appeared to have increased its firing rate prior to the

attempt and then fired rhythmically at approximately 2 Hz. The motor cortex neuron in

Figure 3b was highly selective to the attempt phase with high firing right before the attempt

followed by a rhythmic firing pattern during rope pulling. Other neurons had phasic firing to

very specific time points of a trial. For example, the Cg1-M2 neuron shown in Figure 3c fired

just prior to reward entry while the M2/1 neuron in Figure 3d fired as the rat left the reward

zone. Furthermore, some neurons showed multiple phasic peaks such as the motor cortex

neuron in Figure 3e that was active both right before reward entry and at reward leaving.

More complex activity profiles were also observed such as the Cg1 neuron shown in Figure

3f that fired tonically during all three phases but appeared to be briefly inhibited prior to

reward entry and after reward leaving. Some neurons showed a clear change in firing rate

over the course of the recoding. For example, the neuron in Figure 3g was highly active

before reward entry on early trials (top lines of firing rate raster plot) but decreased its firing

rate at reward entry as the weight became heavier over the session (moving vertically down

the raster plot). The motor cortex neuron shown in Figure 3h doesn't have a clear preference

for a trial-phase but does increase in firing over time.

There were a total of 268 (74%) unique task-responsive neurons; 161 of these neurons

were responsive to the attempt phase, 174 to reward entry, and 153 to reward leave. Ninety-

seven neurons were responsive in only one epoch while 122 were responsive in two epochs

and 49 were responsive in all three. There was no significant difference in the proportion of

responsive neurons across the three brain locations (X^2 all p's > 0.51). Thus, unless otherwise

stated, all neurons responsive to at least one task-phase were pooled together across brain

locations for further analysis. Of the 92 neurons that were excluded, 56 were removed

because they did not have a mean firing rate over 0.5 Hz for any task phase while the

remaining 36 neurons were not responsive to any of the task phases compared to their

baselines (see Methods). However, we did repeat all analyses to include the 36 non-task

responsive neurons and found no qualitative differences in the results. Of the 268 neurons, 41

were classified as interneurons by their overall mean firing rate being greater than 10 Hz. The

response profiles of the putative pyramidal neurons and interneurons can be seen in

Supplemental Table 1 (https://doi.org/10.6084/m9.figshare.14755929.v1).

The normalized, trial averaged activity from these task-responsive neurons is shown

in Figure 4a. Trials were defined as the sequence of events beginning at a rope pull attempt

followed by reward zone entry and concluding with reward zone leaving (see Methods).

Neural activity during the attempt phase was highly correlated across task-responsive neurons

which may reflect tonic activity patterns during the rope pulling action (Figure 4b). In

contrast, neural activity during reward entry and reward leaving was more phasic with lower

correlation values from early compared to later time points within these phases. Activity in

the attempt phase was highly dissimilar compared to the two reward phases. To further

visualize neural activity across trials, we carried out PCA on the trial averaged activity of the

task-responsive neurons (Figure 4c). Supporting the cross-correlation analysis, principal

component 1 (PC1; 32% of variance) differentiated trial-bins by attempt phase versus the two

reward phases (reward entry and reward leave). PC2 and PC3 explained a further 16% and

11% of the observed variance, respectively. Trial-bins in the attempt phase clustered together

before transitioning to the reward phases. From the start of reward entry, the trajectory traces

out a path from reward entry to the end of reward leaving; there is some clustering around the

transition point from reward entry to reward leaving that may reflect reward consumption

(see Figure 4c). Overall, dmPFC neurons track task states; ensemble activity is sustained

during attempts and then smoothly transitions through the reward entry and leaving phases.

To test whether neural activity reflected differences in future behavior, we parsed

trials based on whether or not the rat went on to engage in another rope pull attempt after

reward consumption. A trial was deemed an "engage" trial if – after the reward leave phase –

the rat went on to perform another rope pull within 12 seconds. Twelve seconds was chosen

as a cut-off since, on approximately 75% of all trials, rats made another attempt within 12

seconds (Figure 5a). The PCA trial-phase trajectories for engage versus disengage trials were

highly similar (PC1, 24%; PC2, 11%; PC3, 8%; Figure 5b) and distances between bins

tended to be within 2 S.D. of the mean distance between randomly sampled trajectories

(Figure 5c). However, towards the end of the Reward leave phase, the engage and disengage

trajectories started to diverge. There was a main effect for task-phase ($F_{(2, 72)} = 34.3$, $p <$

0.0001). Post-hoc pairwise testing showed no significant difference between the distances of

engage-disengage trajectories for attempt versus reward entry ($p = 0.77$) but reward leaving

distances were significantly greater compared to both the attempt phase ($p < 0.0001$) and

reward entry (p < 0.0001). Thus, in dmPFC the choice to engage or disengage from the task

emerges relatively late as rats leave the reward zone.

To test whether neural activity reflected differences in rope pulling outcome, we also

parsed trials based on whether or not the trial was a "success" (i.e., the weight was lifted 30

cm to trigger reward dispensing) or a "failure" (i.e., rope pulling was initiated but the weight

was not lifted to the full required height of 30 cm). For 75% of recording sessions the average

success rate was above 50% (Figure 5d). The PCA trial phase trajectories of successful

versus failed trials were distinct but had similar overall shapes (PC1, 18%; PC2, 8%; PC3,

7%; Figure 5e). The distances between successful and failed time bins in PC space were

greater than 2 S.D. of randomly sampled trajectories (Figure 5f). Task phase had a significant

effect on successful versus failed trajectory distances ($F_{(2, 72)} = 8.68$, p = 0.0004) where

both the attempt phase (p < 0.0008) and reward leaving phase (p = 0.002) had greater

distances than reward entry. Thus, dmPFC ensemble states may reflect levels of task

engagement and the amount of effort put into a trial.

To test whether neural activity in each task-phase was changing over time as rats

carried out the PW paradigm, we parsed trials based on their temporal quartile of occurrence.

Thirty-five percent (57/161) of attempt phase neurons, 32% (56/174) of reward entry

neurons, and 22% (34/153) of reward leave neurons had a significant effect for quartile. Of

the neurons with a quartile effect, we determined in which quartile they had the highest firing

rate. For all three task-phases, the observed frequencies of peak activity across the quartiles

were significantly different than an equal distribution across the quartiles (Attempts, $X^2(3) =$

28.7, p < 0.0001; Entry, $X^2(3) = 14.7$, p = 0.0021; Leaving, $X^2(3) = 20.8$, p < 0.0002; Figure

6a). The majority of neurons, across task-phases had their highest firing rates in Q1. We also

analyzed each rat's neurons individually and the majority (3/5) of rats had the highest

proportion of neurons most active in Q1. To test if neural activity was changing over the

quartiles, we carried out one-way repeated measures ANOVAs on the population of task-

active quartile-effect neurons in each task-phase (Figure 6b). Across the four quartiles, there

was a significant effect for quartile for the attempt phase (F (1.8, 103.3) = 12.4, p < 0.0001),

reward entry (F (2.3, 128.3) = 3.2, p = 0.037), and reward leave phase (F (2.0, 66.6) = 10.2, p

< 0.0002). Specifically, pairwise post-hoc testing showed Q1 had significantly higher neural

activity compared to Q3 and Q4 during attempts and reward leaving (all p's < 0.05), but not

during reward entry (no pairwise comparisons were significant). For all three task-phases, we

did not see a difference in the observed numbers of quartile selective neurons across brain

regions (Chi-square test, all p's > 0.28; Figure 6c). We tested whether the magnitude of firing

rate change was impacted by session duration and found no effect of session duration for all

three task phases (all p's > 0.23, R^2's < 0.008; Supplemental Figure S2a,

https://doi.org/10.6084/m9.figshare.14515659). These data may indicate that a subpopulation

of dmPFC neurons have their peak activity early on in the session and then decrease firing

over the session.

To visualize any changes in neural activity over the quartiles, we carried out PCA on

the trial averaged activity of all 268 task-responsive neurons across the quartiles (PC1, 17%;

PC2, 8%; PC3, 7%). The PCA trial-phase trajectories were highly conserved across quartiles

(Figure 6d). Similar to the PCA trial phase trajectory that was observed when all trials across

a session were averaged together (see Figure 4c), each quartile's trajectory had activity

clustered tightly during attempts, then abrupt shifts to reward entry and then a smooth

transition through reward entry to reward leaving (Figure 6d). PC3, however, appears to

differentiate the trajectories by quartile in that distinct trajectories for each of the four

quartiles can be seen (Figure 6e, f). Pairwise distances for every trial-bin in PC space across

the four quartiles showed a significant effect for quartile pairs (F (5, 444) = 370, p < 0.0001;

Figure 6f). The distances between Q1-Q2 and Q2-Q3 were equivalent (p = 0.99) whereas all

other pairwise comparisons were significantly different from one another (all p's < 0.0001).

Q3-Q4 had the closest trajectories, whereas Q1-Q4 were the most distant. All distances from

real trajectories were significantly greater than those from randomly generated quartiles

(paired t-tests, all p's < 0.0001). The relative distances between pairs of quartiles remain even

when the population of neurons is subsampled down to 10% of the original number of

neurons (Supplemental Figure S3, https://doi.org/10.6084/m9.figshare.14515665.v1). We

repeated the PCA for the five individual rats that had more than 30 neurons. The general

trend of more-distant quartiles having more-distant trajectories was seen in 3/5 rats (data not

shown). For all five rats, the Q1-Q4 comparison had the farthest distances. Thus, across

sessions the dmPFC may have a consistent representation of the task-relevant behavioral

states that is modified over time by the current task utility.

To determine if neural activity changes could be seen more broadly across a recording

session – as rats went from entering the apparatus, to starting the task, to quitting the task –

we averaged the activity of the 268 task-active neurons in 2 minute epochs. These epochs

consisted of the pre-baseline period, the four quartiles of task performance, the quit period

where the rat could continue to do the task (rope available) but chose not to, and the final

post-baseline period (see Methods). We predicted that neural activity would be the most

similar when comparing the pre-baseline, quit period, and post-baseline epochs, as

behaviorally the rat is not engaged in the task during any of these periods. Cross-correlation

matrices revealed two distinct patterns which were contrary to our predictions: neural activity

was highly similar across the pre-baseline, Q1, and Q2 epochs, and highly similar across the

Q3, Q4, quit period, and post-baseline epochs (Figure 6g). Specifically, neural activity was

highly correlated between pre-baseline and Q1 (r = 0.48), Q4 and the quit period (r = 0.54),

as well as Q4 and the post-baseline period (r = 0.51). Clustering analyses confirmed these

patterns showing two distinct clusters relating to the first half of the session and the latter half

of the session (Figure 6h). The differences in firing rate activity between Q1 and Q4 in these

analyses was significantly influenced by recording duration (linear regression; F = 5.52, p =

0.0195; Supplemental Figure S2c, https://doi.org/10.6084/m9.figshare.14515659). However,

the model fit was poor (R^2 = 0.02). Thus, neural activity in the dmPFC appears to shift as the

session progresses, irrespective of the specific behaviors being performed.

552   We repeated the above analyses on the 227 putative pyramidal neurons and the 41

putative interneurons. The task-responsiveness of pyramidal neurons and interneurons can be

seen in Supplemental Table 1 (https://doi.org/10.6084/m9.figshare.14755929.v1). The

pyramidal neuron restricted dataset yielded very similar results to that of the whole

population and all statistical testing had the same outcomes for peak quartile across task

phases, decrease in firing rate over quartiles, and task phase responsiveness by brain region

(Supplemental Figure 4; https://doi.org/10.6084/m9.figshare.14757342).

559   In contrast, the interneuron restricted dataset did differ from the total and pyramidal

populations (Supplemental Figure 5; https://doi.org/10.6084/m9.figshare.14757876.v1).

Mainly, while interneurons were highly task responsive as shown by the proportion of

neurons selective to the three task phases (Supplemental Table 1), the neurons did not show a

change in firing rate across quartiles (all P's > 0.2; Supplemental Figure 5b). Our other

statistical tests were no longer appropriate (e.g., Chi-square tests) due to the low sample sizes

(Supplemental Figure 5a, c). The lack of robust decrease in firing rate across the quartiles

was also evident in the principal component analysis where PC1 (33% of variance), PC2

(11%), and PC3 (8%) reflected the three task phases (Supplemental Figure 5d, e) while PC4

(6%) showed modest quartile differences (Supplemental Figure 5f, g). Across the entire

session, the interneuron population correlations were similar to the overall and pyramidal

neuron data sets (Supplemental Figure 5h, i). Overall, interneurons may have a specialized

role in encoding the task structure (task phases) and show less modulation by task utility than

pyramidal neurons.

While there were no differences in the proportion of neurons active for a given task

phase across brain regions (Figure 6c), we repeated our analyses on each brain area

individually. The normalized trial activity of all the neurons from each brain regions is shown

in Supplemental Figure 6a-c (https://doi.org/10.6084/m9.figshare.14758143.v2). PCA was

carried out on the population of neurons from each brain region across quartiles during the

three task phases (Supplemental Figure 6d-f). The general trend of the top principal

components corresponding to differentiating the task phases was true for all three brain

regions. However, when comparing the pairwise distances between PC trajectories across

quartiles, Cg1 (Supplemental Figure 6g) and M2/1 (Supplemental Figure 6i) had more

distinct trajectories for each quartile than the intermediary Cg1-M2 (Supplemental Figure

6h). However, when looking at the neural responses over large, 2-minute epochs across

recording sessions, all three brain regions show the same general pattern of closer in time

epochs being more similar (Supplemental Figure 6j-l). These data may suggest that there are

subtle differences in the responsiveness of neurons in these different brain areas, though

targeted manipulation would be necessary to determine their functional roles.

Neuronal task selectivity – Fixed weight paradigm

The PW paradigm affords an animal continuously decreasing utility, due primarily to

the increasing effort costs of the increasing weight. Rats must decide when to quit the task

when the weight becomes too physically demanding. However, quitting in the PW task may

be spurred by other primary factors such as fatigue or satiation. In order to try and understand

the neural activity behind quitting behaviors more broadly, we analyzed dmPFC neural

activity as rats carried out the FW paradigm, which lacks progressive increases in weight.

Instead, the utility of the FW paradigm decreases due to changes in the internal state of the

animal, e.g. fatigue, motivation, satiation.

We recorded a total of 62 dmPFC neurons from the three rats that contributed the nine

FW sessions shown in Figure 3 (2-4 sessions per rat). Of these 62 neurons, a total of 53

(86%) met our task-responsive criteria for further analysis; trial averaged activity is shown in

Figure 7a. Thirty-eight of the task-responsive neurons were active in the attempt phase, 41 in

the reward entry phase, and 31 in the reward leave phase. Twelve of the task-responsive

neurons were active in one phase, 25 in two phases, and 16 in all three. A cross-correlation

matrix across all bins (Figure 7b) shows a similar pattern to that observed for the PW

paradigm (see Figure 4b). Despite the relatively small number of neurons, PCA of the trial

trajectory resulted in a similar low-dimensional trajectory (PC1, 40%; PC2, 16%; PC3, 10%;

Figure 7c) to that seen for the PW trajectory (see Figure 4c). Activity clustered tightly during

the attempt phase before transitioning through the reward entry phase with a relatively wide

trajectory. Activity then clustered again from the reward entry to reward leave transition.

To test whether neural activity was changing over the course of the FW paradigm, we

analyzed the activity of neurons across session quartiles. Kruskal-Wallis tests were used on

each task-responsive neuron to test firing rate variability across quartiles for each task-phase.

Of the 38 neurons active in the attempt phase, 14 (37%) showed a significant change in firing

rate across the quartiles. Twelve of the 41 (29%) reward entry and 9 of the 31 (29%) reward

leave active neurons showed a significant effect for quartile. As was seen in the PW

paradigm, the majority of task-active neurons with a significant effect of quartile had their

highest firing rate in Q1 of the FW task (Attempts, $X^2(3) = 16.3$, p < 0.0009; Entry, $X^2(3) =$

10.0, p < 0.019; Leaving, $X^2(3) = 9.2$, p < 0.027; Figure 8a). Across quartiles, the normalized

firing rates from the population of task-active neurons showed a significant effect for quartile

for all three task-phases (Attempts, F (2.6, 35.8) = 16.7 p = 0.001; Entry, F (2.0, 24.4) = 12.0,

p = 0.005; Leaving, F (2.2, 19.6) = 12.9, p = 0.007; Figure 8b). Post-hoc pairwise

comparisons revealed that the averaged population activity was significantly higher during

Q1 compared to Q4 (all p's < 0.0003) and during Q2 compared to Q4 (all p's < 0.004) in all

three task-phases. Session duration had no effect on the observed changes of firing rate from

Q1 to Q4 during reward entry or leaving (p's > 0.25). For the attempt phase, session duration

was negatively correlated to the differences in firing from Q1 to Q4 (F = -2.2; p = 0.04, R^2 =

0.11). Thus, similar to the PW paradigm, a subset of dmPFC neurons exhibit a decrease in

activity across quartiles in the FW paradigm. Quartile selective neurons by brain region are

shown in Figure 8c, however statistics are not provided as there were too few neurons to

conduct Chi-Square tests.

630   PCA was applied to the averaged normalized activity of the 53 task-active neurons

across the four quartiles during the FW task (PC1, 23%; PC2, 10%; PC3, 8%; Figure 8d, e).

Generally, the same trajectory shape across the whole trial can be seen across all four

quartiles (Figure 8d). However, as was seen in the PW task, the trial trajectories for each

quartile can be distinguished (Figure 8e). The attempt phase remains relatively similar across

quartiles, however, the wide neural trajectories during the reward entry phase show differing

paths. During reward leaving, trajectories become smaller over the quartiles reflective of

more stable, stationary neural states. Pairwise distances were calculated between every

quartile across trial bins (Figure 8f). As was seen in the PW, temporally distant trajectories

tended to be farther apart than temporally proximal trajectories (F (5, 444) = 120, p <

0.0001). Differing from the PW paradigm, distances between Q2-Q3 and Q3-Q4 (p = 0.99)

and Q1-Q2 and Q2-Q4 (p = 0.15) were not different in the FW paradigm, otherwise, all other

pairwise comparisons were significantly different (all p's < 0.0001). In order of mean

distance, Q1-Q2 was more distant than Q2-Q3 and Q3-Q4. Unique to the FW, distances

between trajectories tended to be farthest apart prior to the reward entry point and after

reward leaving point. The average distances of every real trajectory were significantly greater

than those from randomly sampled trajectories (paired t-tests, all p's < 0.0001).

Neural activity during the FW paradigm was also analyzed in broader two minute

epochs across the session, which included the pre- and post-baseline periods (Figure 8g, h).

Neural activity was highly correlated between pre-baseline and Q1 (r = 0.57), before

becoming less correlated quartile to quartile. Neural activity across the Q4, Quit, and post-

baseline periods was also highly correlated, with significant Pearson's r values observed

between Q4 and Quit (r = 0.64), Q4 and post-baseline (r = 0.72), and Quit versus post-

baseline (r = 0.80). Session duration had no impact on the firing rate differences between Q1

and Q4 (F = 0.96, p = 0.34, R^2 = 0.02). Overall, the FW paradigm results are similar to the

PW results in that a consistent representation of the behavioral states of the task is modified

across quartiles; in the FW paradigm, however, this cannot be directly attributed to increases

in absolute physical effort across time (i.e., increments in rope weight).

Discussion

We utilized two complimentary effortful behavioral paradigms to investigate how

dmPFC neural activity responded to changes in task utility, and to test whether dmPFC

ensemble activity reflected a rat's progression towards task quitting. In both the PW and FW

paradigms, behavioral metrics indicate that rats were highly engaged in the task until the final

quartile (Q4) where engagement waned. In the PW paradigm, rats took longer to earn rewards

and failed rope-pulling more often quartile-by-quartile as they approached their quit weight in

Q4. In the FW paradigm, reward intervals and failed attempts only increased in Q4. Overall,

these data show that the PW paradigm ramps up in difficulty with commensurate changes in

behavior across quartiles, leading towards the quit point. In contrast, the FW paradigm

presents a repetitive, consistent task where behavior is relatively stable until a change in Q4,

immediately preceding the quit point.

Despite the behavioral differences between the two paradigms, we observed similar

neural activity patterns. Neural activity in dmPFC neurons on average was highest in Q1 and

decreased across quartiles, leading to the quit point. Since these patterns were observed in

both paradigms they cannot be solely attributed to progressive increases in rope weight. PCA

showed conserved neural trajectories for trial phases (attempt, reward entry, reward leave),

which remained relatively consistent across the session. However, each quartile had a distinct

trajectory. The more temporally close quartiles had closer PCA trajectories, with Q1 and Q4

being the most distinct in both task paradigms. Broader temporal analysis of neural activity

encompassing the entire recording session – which included the baseline periods where the

rope was unavailable – revealed distinct shifts in activity from the start of the session to the

end. Overall, our results suggest that the dmPFC maintains a consistent representation of the

sequential behavioral states of the task that is responsive to repetitive task engagement. We

propose that the observed decrease in activity across a session reflects the decrease in task

utility across time, and that this corresponds to waning motivation and eventual quitting of

the task.

We observed similar activity patterns in neurons from Cg1, Cg1-M2, and M2/1 during

WLT performance. The proportion of neurons from each brain area that were active in each

task-phase were not significantly different. These results were somewhat surprising as we

expected M2 neurons to be more responsive to attempt phase rope pulling and Cg1 neurons

to be more responsive to reward phases. However, our results do align with the overlapping

roles ascribed to the two brain areas. Generally, M2 has been proposed to select and adapt

action plans based on sensory feedback (13, 15, 26) while the Cg1 monitors behaviors to

determine their utility and bias behavior to maximize value (27, 28). Both brain regions may

play complimentary roles in physical effort-based cost-benefit decision making: M2 selects

actions and evaluates sensorimotor feedback of the costs and outcomes of those actions,

while anterior Cg1 evaluates the overall utility of the selected action plan in comparison to

available alternatives (18, 29). We did find some subtle differences in how strongly the

regions differentiated their activity across the quartiles (Supplemental Figure 6). Further

manipulation experiments, specifically spanning the medial to lateral extent of the dmPFC,

would be needed to determine precise functional differences. Our results add further support

that anterior Cg1 and M2 play complimentary roles in cost-benefit decision making,

specifically at the intersection of action selection, action-outcome monitoring, and adjusting

future behavior.

Numerous studies have shown that the Cg1/ACC and M2 (or supplementary motor

area in primate) respond to the relative value of behaviors, with firing rates or BOLD

responses corresponding to high value options (18, 20, 21, 23, 30–34). However, in most

previous studies subjects had to choose between, or are at least were presented with, multiple

stimuli or options in a given task; quitting not being one of the options (though see 7)).

Furthermore, there has generally been an emphasis on the evaluation of the *relative* value of

behaviors in the context of all possible behaviors (21, 31, 32, 35) even if there is no overt

choice between multiple options (20, 34).

In our WLT rats could only choose between two options – carrying out the task or

not; there were no other extrinsically rewarded actions available in the environment. In

accordance with dmPFC encoding the relative value of a behavior, we observed a value

signal that decreased over the course of the session, culminating in the quit point (quitting

being the only alternative course of action). Overall our results further support the idea that

dmPFC encodes the relative value of behaviors in order to choose actions that maximize

utility (36) possibly through exerting cortex-wide control in task engagment (3). At a certain

point, in response to changing environmental demands and internal state, quitting becomes

the behavior with optimal utility when trying to minimize energy expenditure while trying to

satisfy internal drives (37–40).

When we compared trials where the animal went on to engage in another rope pull,

versus trials where the animal disengaged from the task, dmPFC activity in PC space showed

general overlap in neural trajectories. However, significant differences were observed

between engage and disengage trials after the rat left the reward cap. In separate analyses,

when we compared neural activity during successful trials versus failed, non-rewarded trials,

dmPFC activity in PC space showed highly distinct neural trajectories between trial types.

Taken together, one interpretation of these data is that dmPFC ensemble activity drives the

level of WLT engagement: one state correlates to effortful exertion and persistence in the task

(successful rope-pulling and/or re-engagement in a subsequent trial), whereas a state shift

correlates to faltering task performance (failed rope-pulling and/or disengagement in a

subsequent trial).

Previous studies that have examined quitting tended to use progressive ratio lever

pressing paradigms (41–44). Lesions to the Cg1 (43) or more ventral prelimbic cortex (44) do

not alter the quit-point/breakpoint of rats in these paradigms. However, lesions to the Cg1 of

rats has consistently resulted in avoidance of high-effort options (climbable-barriers, high-

ratio lever presses) when relatively lower effort options are available (45–48). We have

shown previously that electrical stimulation to Cg1 biases rats to quit earlier in our WLT

(49).These results seem to be at odds with one another: lesioning Cg1 results in rats shifting

their choice preference towards lower effort options, but does not affect high effort

performance on progressive ratio tasks. Wang et al. (50) provided some clarity on this

discrepancy by showing that inactivation of rat Cg1 impairs performance in a novel "do more

get more" task, suggesting that Cg1 is critical for *self-paced* effort expenditure. Thus, in the

context of quitting behaviors, Cg1 may provide a relative value signal of the current effortful

behavior and this signal influences the decision to persist or quit in a self-paced task. Future

research manipulating both Cg1 and M2 will be needed to elucidate the functional roles of

the two regions in effortful decision-making tasks. Furthermore, previous evidence has

shown neuromodulators such as dopamine (22, 50) and serotonin (51) influence decisions to

persist or quit behaviors.

Two proposed, complimentary functions of the dmPFC are reinforcement-learning

and action selection (27, 52, 53). These two functions work synergistically where the dmPFC

learns about and updates a mental model of the world forming an abstract task space in order

to select actions that maximize the value of an environment (54). Using PCA we

demonstrated that the dmPFC contains a consistent task-space representation of behaviors

during our task: effortful rope pulling (attempt phase), reward approach and consumption

(reward entry phase), and prospective task engagement at the end of a trial (reward leave

phase). Similar results have been reported showing the Cg1/ACC representing task-space (12,

20) including the Cg1 recalling the locations of unchosen rewards when mistakes were made,

providing evidence of an addressable task-space representations (55). In addition, Wang et al.

(56) reported Cg1 representations of the sequential steps of a delay-based cost-benefit task

where rats were free to nose-poke for longer durations to earn proportionally larger rewards.

We found similar results to Wang et al. (56) in the proportion of recorded neurons

representing task states (approximately 70%) and subpopulation responding to cost-benefit

contingencies or quartiles (approximately 35%). Thus, the dmPFC may have flexible,

sequential representations of all task-relevant behaviors that can be modified based on cost-

benefit contingencies. Together this enables optimal action selection, which sometimes

means quitting the task at hand and moving on.

References:

1.   **Hayden BY**. Economic choice: the foraging perspective. *Curr Opin Behav Sci* 24: 1–6, 2018. doi:
10.1016/j.cobeha.2017.12.002.

2.   **Stephens DW**, **Krebs JR**. *Foraging Theory*. Princeton (USA): Princeton University Press, 1986.

3.   **Allen WE**, **Kauvar IV**, **Chen MZ**, **Richman EB**, **Yang SJ**, **Chan K**, **Gradinaru V**, **Deverman BE**, **Luo
L**, **Deisseroth K**. Global Representations of Goal-Directed Behavior in Distinct Cell Types of
Mouse Neocortex. *Neuron* 94: 891-907.e6, 2017. doi: 10.1016/j.neuron.2017.04.017.

4.   **Euston DR**, **Gruber AJ**, **McNaughton BL**. The Role of Medial Prefrontal Cortex in Memory and
Decision Making. *Neuron* 76: 1057–1070, 2012. doi: 10.1016/j.neuron.2012.12.002.

5.   **Kennerley SW**, **Walton ME**. Decision making and reward in frontal cortex: Complementary
evidence from neurophysiological and neuropsychological studies. *Behav Neurosci* 125: 297–
317, 2011. doi: 10.1037/a0023575.

6.   **Zhao W**, **Kendrick KM**, **Chen F**, **Li H**, **Feng T**. Neural circuitry involved in quitting after repeated
failures: role of the cingulate and temporal parietal junction. *Sci Rep* 6: 24713, 2016. doi:
10.1038/srep24713.

7.   **Hosokawa T**, **Kennerley SW**, **Sloan J**, **Wallis JD**. Single-Neuron Mechanisms Underlying Cost-
Benefit Analysis in Frontal Cortex. *J Neurosci* 33: 17385–17397, 2013. doi:
10.1523/JNEUROSCI.2221-13.2013.

8.   **Kennerley SW**, **Walton ME**, **Behrens TEJ**, **Buckley MJ**, **Rushworth MFS**. Optimal decision
making and the anterior cingulate cortex. *Nat Neurosci* 9: 940–947, 2006. doi:
10.1038/nn1724.

9.   **Klein-Flügge MC**, **Kennerley SW**, **Friston K**, **Bestmann S**. Neural signatures of value comparison
in human cingulate cortex during decisions requiring an effort-reward trade-off. Neuroscience.

10. **Walton ME**, **Bannerman DM**, **Alterescu K**, **Rushworth MFS**. Functional Specialization within
Medial Frontal Cortex of the Anterior Cingulate for Evaluating Effort-Related Decisions. *J
Neurosci* 23: 6475–6479, 2003. doi: 10.1523/JNEUROSCI.23-16-06475.2003.

11. **Gu X**, **Staines WA**, **Fortier PA**. Quantitative analyses of neurons projecting to primary motor
cortex zones controlling limb movements in the rat. *Brain Res* 835: 175–187, 1999. doi:
10.1016/S0006-8993(99)01576-0.

12. **Lapish CC**, **Durstewitz D**, **Chandler LJ**, **Seamans JK**. Successful choice behavior is associated
with distinct and coherent network states in anterior cingulate cortex. *Proc Natl Acad Sci* 105:
11963–11968, 2008. doi: 10.1073/pnas.0804045105.

13. **Reep RL**, **Corwin JV**, **Hashimotos A**, **Watsonp RT**. Efferent Connections of the Rostral Portion of
Medial Agranular Cortex in Rats. *Brain Res Bull* 19: 203–221, 1987.

14.   **Rolls ET**. The cingulate cortex and limbic systems for emotion, action, and memory. *Brain*
805   *Struct Funct* 224: 3001–3018, 2019. doi: 10.1007/s00429-019-01945-2.

806   15.   **Barthas F**, **Kwan AC**. Secondary Motor Cortex: Where 'Sensory' Meets 'Motor' in the Rodent
807   Frontal Cortex. *Trends Neurosci* 40: 181–193, 2017. doi: 10.1016/j.tins.2016.11.006.

808   16.   **Ebbesen CL**, **Insanally MN**, **Kopec CD**, **Murakami M**, **Saiki A**, **Erlich JC**. More than Just a
809   "Motor": Recent Surprises from the Frontal Cortex. *J Neurosci* 38: 9402–9413, 2018. doi:
810   10.1523/JNEUROSCI.1671-18.2018.

811   17.   **Murakami M**, **Vicente MI**, **Costa GM**, **Mainen ZF**. Neural antecedents of self-initiated actions in
812   secondary motor cortex. *Nat Neurosci* 17: 1574–1582, 2014. doi: 10.1038/nn.3826.

813   18.   **Sul JH**, **Jo S**, **Lee D**, **Jung MW**. Role of rodent secondary motor cortex in value-based action
814   selection. *Nat Neurosci* 14: 1202–1208, 2011. doi: 10.1038/nn.2881.

815   19.   **Porter B**, **Hillman KL**. A Novel Weight Lifting Task for Investigating Effort and Persistence in
816   Rats. *Front Behav Neurosci* 13: 275, 2019. doi: 10.3389/fnbeh.2019.00275.

817   20.   **Cowen SL**, **Davis GA**, **Nitz DA**. Anterior cingulate neurons in the rat map anticipated effort and
818   reward to their associated action sequences. *J Neurophysiol* 107: 2393–2407, 2012. doi:
819   10.1152/jn.01012.2011.

820   21.   **Hillman KL**, **Bilkey DK**. Neurons in the Rat Anterior Cingulate Cortex Dynamically Encode Cost-
821   Benefit in a Spatial Decision-Making Task. *J Neurosci* 30: 7705–7713, 2010. doi:
822   10.1523/JNEUROSCI.1273-10.2010.

823   22.   **Salamone JD**, **Cousins MS**, **Bucher S**. Anhedonia or anergia? Effects of haloperidol and nucleus
824   accumbens dopamine depletion on instrumental response selection in a T-maze cost/benefit
825   procedure. *Behav Brain Res* 65: 221–229, 1994. doi: 10.1016/0166-4328(94)90108-2.

826   23.   **Kennerley SW**, **Wallis JD**. Evaluating choices by single neurons in the frontal lobe: outcome
827   value encoded across multiple decision variables. *Eur J Neurosci* 29: 2061–2073, 2009. doi:
828   10.1111/j.1460-9568.2009.06743.x.

829   24.   **Hyman JM**, **Ma L**, **Balaguer-Ballester E**, **Durstewitz D**, **Seamans JK**. Contextual encoding by
830   ensembles of medial prefrontal cortex neurons. *Proc Natl Acad Sci* 109: 5086–5091, 2012. doi:
831   10.1073/pnas.1114415109.

832   25.   **Paxinos G**, **Watson C**. *The rat brain in stereotaxic coordinates*. Amsterdam; Boston: Academic
833   Press/Elsevier, 2007.

834   26.   **Olson JM**, **Li JK**, **Montgomery SE**, **Nitz DA**. Secondary Motor Cortex Transforms Spatial
835   Information into Planned Action during Navigation. *Curr Biol* 30: 1845-1854.e4, 2020. doi:
836   10.1016/j.cub.2020.03.016.

837   27.   **Holroyd CB**, **McClure SM**. Hierarchical control over effortful behavior by rodent medial frontal
838   cortex: A computational model. *Psychol Rev* 122: 54–83, 2015. doi: 10.1037/a0038339.

839   28.   **Shenhav A**, **Botvinick MM**, **Cohen JD**. The Expected Value of Control: An Integrative Theory of
840   Anterior Cingulate Cortex Function. *Neuron* 79: 217–240, 2013. doi:
841   10.1016/j.neuron.2013.07.007.

29.   **Sul JH**, **Kim H**, **Huh N**, **Lee D**, **Jung MW**. Distinct Roles of Rodent Orbitofrontal and Medial
Prefrontal Cortex in Decision Making. *Neuron* 66: 449–460, 2010. doi:
10.1016/j.neuron.2010.03.033.

30.   **Croxson PL**, **Walton ME**, **O'Reilly JX**, **Behrens TEJ**, **Rushworth MFS**. Effort-Based Cost-Benefit
Valuation and the Human Brain. *J Neurosci* 29: 4531–4541, 2009. doi:
10.1523/JNEUROSCI.4515-08.2009.

31.   **Hart EE**, **Blair GJ**, **O'Dell TJ**, **Blair HT**, **Izquierdo A**. Chemogenetic Modulation and Single-Photon
Calcium Imaging in Anterior Cingulate Cortex Reveal a Mechanism for Effort-Based Decisions. *J*
Neurosci 40: 5628–5643, 2020. doi: 10.1523/JNEUROSCI.2548-19.2020.

32.   **Hillman KL**, **Bilkey DK**. Neural encoding of competitive effort in the anterior cingulate cortex.
Nat Neurosci 15: 1290–1297, 2012. doi: 10.1038/nn.3187.

33.   **Kennerley SW**, **Dahmubed AF**, **Lara AH**, **Wallis JD**. Neurons in the Frontal Lobe Encode the
Value of Multiple Decision Variables. *J Cogn Neurosci* 21: 1162–1178, 2009. doi:
10.1162/jocn.2009.21100.

34.   **Porter BS**, **Hillman KL**, **Bilkey DK**. Anterior cingulate cortex encoding of effortful behavior. *J*
Neurophysiol 121: 701–714, 2019. doi: 10.1152/jn.00654.2018.

35.   **Wallis JD**, **Kennerley SW**. Contrasting reward signals in the orbitofrontal cortex and anterior
cingulate cortex: Contrasting reward signals in the orbitofrontal cortex. *Ann N Y Acad Sci* 1239:
33–42, 2011. doi: 10.1111/j.1749-6632.2011.06277.x.

36.   **Rushworth M**, **Walton M**, **Kennerley S**, **Bannerman D**. Action sets and decisions in the medial
frontal cortex. *Trends Cogn Sci* 8: 410–417, 2004. doi: 10.1016/j.tics.2004.07.009.

37.   **Allen WE**, **Chen MZ**, **Pichamoorthy N**, **Tien RH**, **Pachitariu M**, **Luo L**, **Deisseroth K**. Thirst
regulates motivated behavior through modulation of brainwide neural population dynamics.
Science 364: 11, 2019.

38.   **Hockey R**. *The psychology of fatigue: work, effort, and control*. Cambridge: Cambridge
University Press, 2013.

39.   **Noakes TD**. Fatigue is a Brain-Derived Emotion that Regulates the Exercise Behavior to Ensure
the Protection of Whole Body Homeostasis. *Front Physiol* 3, 2012. doi:
10.3389/fphys.2012.00082.

40.   **van der Linden D**. The urge to stop: The cognitive and biological nature of acute mental
fatigue. In: *Cognitive fatigue: Multidisciplinary perspectives on current research and future*
applications., edited by Ackerman PL. American Psychological Association, p. 149–164.

41.   **Hodos W**. Progressive Ratio as a Measure of Reward Strength. *Sci New Ser* 134: 943–944, 1961.

42.   **Rickard JF**, **Body S**, **Zhang Z**, **Bradshaw CM**, **Szabadi E**. Effect of reinforecer magnitude on
performance maintained by progressive-ratio schedules. *J Exp Anal Behav* 91: 75–87, 2009. doi:
10.1901/jeab.2009.91-75.

43.   **Schweimer J**. Involvement of the rat anterior cingulate cortex in control of instrumental
responses guided by reward expectancy. *Learn Mem* 12: 334–342, 2005. doi:
10.1101/lm.90605.

44.　**Swanson K**, **Goldbach HC**, **Laubach M**. The rat medial frontal cortex controls pace, but not
breakpoint, in a progressive ratio licking task. *Behav Neurosci* 133: 385–397, 2019. doi:
10.1037/bne0000322.

45.　**Hart EE**, **Gerson JO**, **Zoken Y**, **Garcia M**, **Izquierdo A**. Anterior cingulate cortex supports effort
allocation towards a qualitatively preferred option. *Eur J Neurosci* 46: 1682–1688, 2017. doi:
10.1111/ejn.13608.

46.　**Hauber W**, **Sommer S**. Prefrontostriatal Circuitry Regulates Effort-Related Decision Making.
Cereb Cortex 19: 2240–2247, 2009. doi: 10.1093/cercor/bhn241.

47.　**Holec V**, **Pirot HL**, **Euston DR**. Not all effort is equal: the role of the anterior cingulate cortex in
different forms of effort-reward decisions. *Front Behav Neurosci* 8, 2014. doi:
10.3389/fnbeh.2014.00012.

48.　**Rudebeck PH**, **Walton ME**, **Smyth AN**, **Bannerman DM**, **Rushworth MFS**. Separate neural
pathways process different decision costs. *Nat Neurosci* 9: 1161–1168, 2006. doi:
10.1038/nn1756.

49.　**Silva C**, **Porter BS**, **Hillman KL**. Stimulation in the Rat Anterior Insula and Anterior Cingulate
During an Effortful Weightlifting Task. *Front Neurosci* 15: 643384, 2021. doi:
10.3389/fnins.2021.643384.

50.　**Wang S**, **Hu S-H**, **Shi Y**, **Li B-M**. The roles of the anterior cingulate cortex and its dopamine
receptors in self-paced cost–benefit decision making in rats. *Learn Behav* 45: 89–99, 2017. doi:
10.3758/s13420-016-0243-0.

51.　**Lottem E**, **Banerjee D**, **Vertechi P**, **Sarra D**, **Lohuis M oude**, **Mainen ZF**. Activation of serotonin
neurons promotes active persistence in a probabilistic foraging task. *Nat Commun* 9: 1000,
2018. doi: 10.1038/s41467-018-03438-y.

52.　**Hadland KA**, **Rushworth MFS**, **Gaffan D**, **Passingham RE**. The Anterior Cingulate and Reward-
Guided Selection of Actions. *J Neurophysiol* 89: 1161–1164, 2003. doi: 10.1152/jn.00634.2002.

53.　**Paus T**. Primate anterior cingulate cortex: Where motor control, drive and cognition interface.
Nat Rev Neurosci 2: 417–424, 2001. doi: 10.1038/35077500.

54.　**Heilbronner SR**, **Hayden BY**. Dorsal Anterior Cingulate Cortex: A Bottom-Up View. *Annu Rev
Neurosci* 39: 149–170, 2016. doi: 10.1146/annurev-neuro-070815-013952.

55.　**Mashhoori A**, **Hashemnia S**, **McNaughton BL**, **Euston DR**, **Gruber AJ**. Rat anterior cingulate
cortex recalls features of remote reward locations after disfavoured reinforcements. *eLife* 7:
e29793, 2018. doi: 10.7554/eLife.29793.

56.　**Wang S**, **Shi Y**, **Li B-M**. Neural representation of cost–benefit selections in rat anterior cingulate
cortex in self-paced decision making. *Neurobiol Learn Mem* 139: 1–10, 2017. doi:
10.1016/j.nlm.2016.12.003.

Figure Legends

Figure 1: Task overview and histology. a) Weight lifting task schematic as reported in Porter and Hillman (2020). b) Trial breakdown illustrating the three phases of interest. In the Attempt phase, rats are physically pulling the rope. In Reward entry, rats enter the reward cap area and consume sucrose reward if present. In the Reward leave, rats leave the reward cap area and choose to re-engage in a rope pull or disengage. c) Histological reconstruction of tetrode positions. Each color represents the tetrodes from one rat. Tetrodes spanned +3.72 mm to +2.28 mm relative to Bregma. Schematics modified from Paxinos & Watson (2007).

Figure 1: Behavioral measures on the PW and FW paradigms. a-d) Behavioral metrics for the PW task. e-h) Behavioral metrics for the FW task. For both tasks, each recording session was broken up into four equal duration quartiles, Q1-Q4. Histograms illustrate the weight that the rats quit on in the PW (a) or the duration until quitting in the FW (e). Each rat is assigned a different color. The time on-task ratio was calculated as the time rats spent working (making attempts) divided by the total time in each quartile. The success-to-attempt ratio was calculated as the number of successful attempts divided by the total number of attempts for each quartile. Box plots represent the first and third quartile with the middle line representing the median. Error bars represents $\pm$ 1.5 times the interquartile range. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Figure 2: Examples of single unit activity across task-phases during PW sessions. a-h) Each panel shows the activity of one neuron across the three task-phases with a raster plot and peri-event time histogram. Neuron location (Cg1, Cg1-M2, M2/1) is indicated in the title line for each panel. For the raster plots, each row is a trial where the top-most row corresponds to the first trial and descends chronologically. Vertical lines indicate the time of event: attempt, reward entry, reward leaving. Histograms were calculated with 100 ms bins. The y-tick number indicates the highest number of spikes across all three phases. The y-axis is consistent across all three phases.

Figure 3: Population activity of task-responsive neurons during the PW paradigm. a) Trial averaged activity of all 268 task-responsive neurons, sorted by trial-bin in which neurons had their highest activity. Each trial was segmented into 100ms time bins. For graphical purposes, each neuron's trial averaged activity was scaled between 0 and 1. b) Cross-correlation matrix of the average, normalized activity of all active neurons across trial phases. c) PCA scores from the average neural activity across trials in the first three principal components. Every point represents the top three PC scores from one 100ms bin. Red represents the start of the trial and progresses to blue at the end of the trial. The first time bin is highlighted with a star and last bin with a square. Black circles represent trial-phase transitions from Attempt to Reward Entry, and Reward Entry to Reward Leave.

Figure 4: dmPFC neural activity changes during task engagement and failure in the PW
paradigm. a) Cumulative distribution function of the time it took rats to engage in another
trial following reward zone leaving. Quartile boundaries indicated by text. b) Neural
trajectories in PC space during trials where rats subsequently engaged in a rope pulling trial
(blue) or disengaged (red). c) Distances between engage vs disengage trials in PC space (blue
line) for each 100ms time bin. Solid black line is the mean distance between random
trajectories and grey area represents ± 2 S.D. d) Cumulative distribution function for the
proportion of successful trials across sessions. Quartile boundaries indicated by text. e)
Neural trajectories in PC space during trials where rats succeeded in lifting the weight 30 cm
(blue) or failed (red). f) Distances between successful vs failed trials in PC space (blue line)
for each 100ms time bin. Solid black line is the mean distance between random trajectories
and grey area represents ± 2 S.D. b & e) Lighter color indicates earlier within the trial. The
first time bin is highlighted with a star and last bin with a square. Black circles represent trial-
phase transitions from Attempt to Entry and Entry to Leave.

Figure 5: dmPFC neural activity across quartiles during the PW paradigm. a) Distributions of

peak neural activity across the four quartiles for each task-phase. Dotted line indicates equal

distribution across quartiles. b) Average population activity of task-active neurons with a

quartile effect. Activity averaged over all neurons shown as solid black line with grey

shading representing S.E.M. Individual neuron activity plotted as light grey lines. c)

Proportion of task-responsive neurons with a significant quartile effect by brain region. d &

e) PCA of trial activity across the four quartiles, viewed from two different angles. Q1, green;

Q2, blue; Q3, purple; Q4, red. f) Pairwise distances between PC trajectories (colored) and

average distance between shuffled quartiles (black) with ± 2 S.D. in grey. Cooler colors

represent temporally closer pairs while warmer colors indicate more temporally distant pairs.

g) Cross-correlation matrix of 2 min epoch activity across sessions. Pre, pre-baseline; post,

post-baseline. h) Dendrogram of hierarchically clustered correlation distances.

1000

Figure 6: Population activity of task-responsive neurons during the FW paradigm. a) Trial

averaged activity of all 53 task-responsive neurons, sorted by trial-bin in which neurons had

their highest activity. Each trial was segmented into 100ms time bins. For graphical purposes,

each neuron's trial averaged activity was scaled between 0 and 1. b) Cross-correlation matrix

of the average, normalized activity of all active neurons across trial phases. c) PCA scores

from the average neural activity across trials in the first three principal components. Every

point represents the top three PC scores from one 100ms bin. Red represents the start of the

trial and progresses to blue at the end of the trial. The first time bin is highlighted with a star

and last bin with a square. Black circles represent trial-phase transitions from Attempt to

Reward Entry, and Reward Entry to Reward Leave.

1011

Figure 7: dmPFC neural activity across quartiles during the FW paradigm. a) Distributions of

peak neural activity across the four quartiles for each task-phase. Dotted line indicates equal

distribution across quartiles. b) Average population activity of task-active neurons with a

quartile effect. Activity averaged over all neurons shown as solid black line with grey

shading representing S.E.M. Individual neuron activity plotted as light grey lines. c)

Proportion of task-responsive neurons with a significant quartile effect by brain region. d &

e) PCA of trial activity across the four quartiles, viewed from two different angles. Q1, green;

Q2, blue; Q3, purple; Q4, red. f) Pairwise distances between PC trajectories (colored) and

average distance between shuffled quartiles (black) with $\pm$ 2 S.D. in grey. Cooler colors

represent temporally closer pairs while warmer colors indicate more temporally distant pairs.

g) Cross-correlation matrix of 2 minute epoch activity across sessions. Pre, pre-baseline; post,

post-baseline. h) Dendrogram of hierarchically clustered correlation distances.

1024

Figure 14

Bregma 2.28 mm

Progressive weight paradigm

Fixed weight paradigm