Direct regulation of cell cycle regulatory gene expression by NtrX to promote

Sinorhizobium meliloti **cell division**

Shenghui Xing, Fang An, Leqi Huang, Xinwei Yang, Shuang Zeng, Ningning Li,

Lanya Zhang,Wenjia Zheng, Khadidja Ouenzar, Junhui Yan, Liangliang Yu, Li Luo*

Shanghai Key Laboratory of Bio-energy Crops, School of Life Sciences, Shanghai

University, Shanghai 200444, China

Running title: Positive control of *Sinorhizobium* cell division by NtrX

*Correspondence: liluo@shu.edu.cn.

ABSTRACT

Cell division of the alfalfa symbiont, *Sinorhizobium meliloti,* is regulated by a signaling network centered on CtrA. The gene expression of regulatory proteins in the network can be regulated by nutrient signaling systems. In this study, we found that NtrX, one of the regulators of nitrogen metabolic response, can directly regulate the expression of several regulatory genes in the CtrA signaling network. Three groups of *S. meliloti ntrX* mutants showed similar cell division defects, such as slow growth, abnormal morphology of some cells and delayed DNA synthesis. Quantitative RT-PCR assays indicated that in these mutants, the transcription of genes such as *ctrA* and *gcrA* was up-regulated, while the transcription of genes such as *dnaA* and *ftsZ1* was down-regulated. Western blotting showed that the CtrA and GcrA proteins were apparently increased in the mutants. The 53rd aspartate conserved in NtrX homologs can be phosphorylated *in vitro* and *in vivo*. The phosphorylated NtrX protein can bind directly to the promoter regions of *ctrA*, *gcrA*, *dnaA* and *ftsZ1* by recognizing the characteristic sequence $CAAN_{1-5}TTG$. Therefore, phosphorylation of NtrX is essential for cell cycle regulation of *S. meliloti*. We expressed the NtrX protein carrying a phosphorylation site substitution in *Agrobacterium tumefaciens* and found that the expressed strains had different growth phenotypes, suggesting that NtrX also regulates cell division in this bacterial species. Our findings reveal that NtrX acts as a transcriptional regulator that positively affects bacterial cell division, associated with nitrogen metabolism.

IMPORTANCE

Sinorhizobium meliloti infects the host legume alfalfa and induces the formation of nitrogen-fixing nodules. The proliferation of rhizobia in plant tissues and cells is strictly controlled in the early stage of the interaction between symbiotic partners. However, the control mechanism is not very clear. Cell division of *S. meliloti* in the free-living state is regulated by the CtrA signaling network, but the molecular

mechanisms by which the CtrA system is associated with environmental nutrient

signals (e.g., ammonia nitrogen) need to be further explored. This study demonstrates

that NtrX, a regulator of nitrogen metabolism, required for symbiotic nodulation and

nitrogen fixation by *S. meliloti* 1021, can act as a transcriptional regulator of the CtrA

signaling system. It may link nitrogen signaling to cell cycle regulation in *Rhizobium*

species.

Key words:

NtrX; Rhizobium; transcriptional regulation; cell division; CtrA

Caulobacter crescentus is a model strain of α-proteobacteria in molecular cell biology

(1). It takes advantage of one cell division to produce two cells with different shapes

and sizes (2). In recent years, a complex cell cycle regulatory network has been

revealed in this species. This network consists of multiple histidine kinases such as

CckA, DivL, DivJ, and PleC, a histidine phosphotransfer protein ChpT, response

regulators DivK and CpdR, and transcription regulators CtrA, GcrA, DnaA, SciP, and

MucR (1-6). Although this network has been reported to possibly mediate nutritional

signals for regulating bacterial growth and proliferation, the exact molecular

mechanism is currently unclear.

Sinorhizobium meliloti is a model strain of rhizobia that can infect the host plant and

form nitrogen-fixing nodules. During symbiosis, the cell division of *S. meliloti* on the

surface of host plant alfalfa roots, at the front ends of extended infection threads and

in the infection zones of root nodules, is stringently controlled (7), but how cell

division is regulated is not very clear. Although NCR (Nodule Cysteine Rich) peptides

secreted by host plants are known to induce terminal differentiation of bacteroids in

host plant cells (8-11), several legumes do not produce NCR peptides. Therefore,

there may be other mechanisms that control cell division of symbiotic rhizobia in host

plants. Since *S. meliloti* and *C. crescentus* belong to α-proteobacteria, based on the

research results of *C. crescentus*, with the aid of DNA sequence homology analysis,

many cell cycle regulatory genes such as *ctrA*, *ccrM*, *cpdR1*, *divJ*, *divK*, *gcrA* and

pleC, have been identified in *S. meliloti* (12-14). In addition, the hybrid histidine

kinase CbrA is linked to the CtrA regulatory system, which is an important regulator

of cell division in *S. meliloti* (15, 16). However, it is still unclear whether these

regulatory proteins conduct environmental nutrition signals (e.g., ammonia nitrogen)

and whether they play a regulatory role in the symbiotic process.

The NtrY/NtrX two-component system, first discovered in *Azorhizobium caulinodans*,

regulates nitrogen metabolism under free-living conditions and affects nodulation and

nitrogen fixation in the host plant *Sesbania rostrata* (17). Subsequently, *ntrY/ntrX*

homologous genes were found in *Rhizobium tropici* to regulate nitrogen metabolism

and symbiotic nodulation (18). NtrY/NtrX homologs regulate nitrate uptake in

Azospirillum brasilense and *Herbaspirillum seropedicae* (19, 20), and this regulatory

system has been found to simultaneously control nitrogen metabolism and cellular

redox homeostasis in *Rhodobacter capsulatus* (21). Moreover, NtrX is involved in the

regulation of cell envelope formation in *R. sphaeroides*(22). In *Brucella abortus*, the

histidine kinase NtrY participates in micro-oxygen signaling and nitrogen respiration

(23), while the response regulator NtrX controls the expression of respiratory

enzymes in *Neisseria gonorrhoeae* (24). Interestingly, the NtrY/NtrX system regulates

cell proliferation, amino acid metabolism and CtrA degradation in *Ehrlichia*

chaffeensis (25). Finally, NtrX is required for the survival of *C. crescentus* cells and

its expression is induced by low pH (26). These findings show that NtrY/NtrX

appears to be a regulatory system for nitrogen metabolism, which may be involved in

the regulation of cell division.

NtrX is a NtrC family response regulator protein, which consists of a receiver (REC)

domain and a DNA-binding domain (27, 28). X-ray crystal diffraction results indicate

that the NtrX protein of *B. abortus* can form a dimer; the REC domain is composed of

α-helices and 5 β-sheets; the DNA-binding domain contains an HTH motif, which includes 4 α-helices. The three-dimensional structure of the C-terminus has not been resolved (28). *In vitro* experiments demonstrated that the NtrX protein of *B. abortus* can recognize and bind to the palindromic DNA sequence (**CAA**N_{3-5}**TTG**) in the *ntrY* promoter region via the HTH motif to regulate gene transcription (27, 28). In *S. meliloti* 1021, our previous work found that NtrX protein can regulate bacterial growth and proliferation, flagellar synthesis and motility, succinoglycan production, and symbiotic nodulation and nitrogen fixation with the host plant alfalfa (29, 30). In the present study, we investigated the control mechanism by which NtrX regulates *S. meliloti* cell division at the transcriptional level.

RESULTS

Defects of cell division resulting from *ntrX* mutation in *S. meliloti*. We previously constructed a plasmid insertion mutant of the *ntrX* gene in the background of *S. meliloti* 1021, SmLL1 (29). This mutant grew slowly even in the rich LB/MC medium compared to wild-type Sm1021 (29). According to the determined growth curve, the doubling time of bacterial cell proliferation was calculated to be 180 min for SmLL1 compared to 160 min for the wild-type strain as reported (34). Microscopic observation revealed that 5% to 10% of SmLL1 cells grown in the LB/MC broth up to the logarithmic phase exhibited morphological abnormalities (such as cell elongation, Y-shaped or V-shaped), whereas Sm1021 had almost no abnormally shaped cells (Fig. 1A). To determine whether the appearance of abnormally shaped cells of the SmLL strain is associated with the synthesis and segregation of genomic DNA, we synchronized the *S. meliloti* cells, inoculated them in LB/MC broth, grew them for 90 min and 180 min, respectively, and then harvested the cells for flow cytometric analysis. The results showed that Sm1021 synchronized cells were haploid containing 1C genomic DNA, while cells grown for 90 min were diploid containing 2C genomic DNA; cells grown for 180 min were both haploid and diploid (Fig. 1B). Cells of the SmLL mutant after synchronization were mainly haploid, and cells grown for 90 min

were mainly diploid, but the cells grown for 180 min were still mainly diploid (Fig.

1B), indicating a decelerated of replication and segregation of their genomic DNA as

compared to the wild type. These observations indicate that the SmLL mutant has cell

division defects.

It may be fatal due to the deletion of *ntrX,* and we have not yet successfully screened

the deletion mutant of this gene in *S. meliloti* 1021*,* so a depletion strain (where the

ntrX gene on the genome has been deleted and the NtrX protein in the cell is

expressed from an IPTG-inducible plasmid) was constructed to verify the above

results. Optical microscopic observation showed that more than 30% of the cells of

the depleted strain in LB/MC broth without IPTG showed abnormal shapes (such as

elongation and T-shaped), while in LB/MC broth with IPTG, almost no abnormal cells

were observed (Fig. 1C). The depletion strain barely proliferated in LB/MC broth

without IPTG (Figure 1D), indicating that *ntrX* gene expression is required for the cell

division of *S. meliloti.* Flow cytometric analysis showed that in addition to haploid

and diploid cells, polyploid cells were detected in the depletion strain in LB/MC broth

without IPTG (Fig.1E). After the addition of IPTG, polyploid cells were almost

undetectable (Fig. 1E), indicating that genomic DNA replication and segregation of *S.*

meliloti requires the expression of the *ntrX* gene.

NtrX, as a regulator of nitrogen metabolism, is composed of a REC domain and a

DNA-binding domain (28). The research results from *C. crescentus* and *B. abortus*

showed that the D53 residue on the REC domain is the phosphorylation site of the

protein (26, 28). We found that the REC domain of the NtrX protein from *S. meliloti*

also contains a conserved D53 residue. If the NtrX protein is indeed involved in the

regulation of *S. meliloti* cell division, as described above, then the mutation of the

D53 residue would affect its regulatory function. To test this hypothesis, we began

constructing amino acid substitution mutations (the codon of D was replaced by A, N

or E, respectively) in the *ntrX* gene on the genome of *S. meliloti* 1021, but were

unable to screen the mutants. As a result, we cloned the mutation gene into the

expression vector pSRK-Gm (31) and then introduced the recombinant plasmid into

Sm1021. On the LB/MC/IPTG plate, we found that the strain expressing NtrX^{D53A} or

NtrX^{D53N} did not form visible colonies; however, the strain expressing NtrX^{D53E} formed some colonies (Fig. S1). We observed GFP-labeled *S. meliloti* cells (32) cultured in LB/MC/IPTG broth up to the logarithmic phase under a fluorescence microscope and found that more than 20% of Sm1021/p*ntrX*^{D53E} cells had abnormal morphology, while Sm1021/p*ntrX* cells were normal (Fig. 1F). The growth curve determination also showed that the growth of Sm1021/p*ntrX*^{D53E} in LB/MC/IPTG broth was apparently slower than that of Sm1021/p*ntrX* (Fig. 1G). Flow cytometric analysis showed that synchronized Sm1021/p*ntrX* cells were haploid and grew for 90 min in LB/MC/IPTG broth, and were mainly diploid; cells grown for 180 min reverted to haploids (Fig. 1G). Synchronized Sm1021/p*ntrX*^{D53E} cells are haploids, while cells grown in LB/MC/IPTG broth for 90 minutes were mainly diploids (Fig. 1G). However, most of the cells cultured for 180 min were haploid and diploid, but some cells were polyploid (Fig. 1G). These results suggest that the phosphorylated NtrX protein is required for the regulation of cell division of *S. meliloti*.

Transcription of cell cycle regulated genes under the regulation of NtrX in *S. meliloti*. Since NtrX is involved in controlling cell division of *S. meliloti*, does it regulate the transcription of cell cycle regulatory genes such as *ctrA*, *gcrA* and *dnaA*? To answer this question, we applied quantitative RT-PCR technology to analyze the transcript levels of cell cycle regulatory genes in *S. meliloti* cells. First, the synchronized Sm1021 and SmLL cells were subcultured into LB/MC broth for shaking incubation, and then total RNA was extracted from cells that grew every half hour. The qRT-PCR results showed that the transcript level of the *ntrX* gene increased first, then decreased, and reached the maximum value in the cells cultured for 90 min, showing a trend of cyclical changes, while the *ntrY* gene cyclical transcription trend was not obvious (Fig. 2A). Known cell cycle regulatory genes, such as *ctrA*, *gcrA* and *dnaA*, also showed a cyclical transcription trend (Fig. 2A and S2A). Compared to the wild type, in the SmLL1 mutant, transcript levels of the *ntrX* gene were lower in cells grown at different times, but the cyclical trend was unchanged, and cell cycle regulatory genes with similar down-regulation trend of *ntrX* transcription included *dnaA*, *ftsZ1*, *pleC*, *chpT* and *cpdR1* (Fig. 2A and S2A). Contrary to these results, *ctrA*

was mainly up-regulated in SmLL1 mutants with similar trends to *gcrA*, *ccrM* and

ntrY (Fig. 2A and S2A). These findings suggest that the NtrX protein may repress the

transcription of genes such as *ctrA* and *gcrA* and activate the transcription of genes

such as *dnaA* and *ftsZ1*.

We analyzed the transcripts of cell cycle regulatory genes in cells of the depletion

strain to verify the above results. The qRT-PCR results showed that depleted cells

cultured in LB/MC broth without IPTG had extremely low levels of *ntrX* transcripts,

while transcripts of most cell cycle regulatory genes were available for detection (Fig.

2B and S2B). After culturing depleted cells in LB/MC broth containing IPTG for 1 h,

a large number of *ntrX* gene transcripts were detected (Fig. 2B and S2B). The

transcript levels of cell cycle regulatory genes were significantly altered in depleted

cells cultured in IPTG broth for 2 or 3 h compared to cells cultured without IPTG

broth: the transcription of *ctrA, gcrA* and *ccrM* was down-regulated; the transcription

of *dnaA*, *ftsZ1*, *pleC*, *chpT* and *cpdR1* was up-regulated (Fig. 2B and S2B). These

results further confirm that the NtrX protein represses the transcription of genes such

as *ctrA* and *gcrA* and simultaneously activates the transcription of genes such as *dnaA*

and *ftsZ1*.

We analyzed the transcript levels of major cell cycle regulatory genes in

Sm1021/p*ntrX*^{D53E} and Sm1021/p*ntrX* cells to determine whether the conserved D53

residue on NtrX is essential for transcriptional regulation. The qRT-PCR results

showed that transcripts of the *ntrX* gene were significantly increased in cells cultured

in LB/MC broth containing IPTG for 2 h compared to cells without IPTG; meanwhile,

the transcripts of *ctrA*, *gcrA* and *ntrY* were significantly reduced or tended to decrease,

while the transcripts of *dnaA* and *ftsZ1* genes were significantly increased (Fig. 2C).

Contrary to the above results, as transcripts of the *ntrX*^{D53E} gene increased

significantly under IPTG induction, transcripts of *ctrA*, *gcrA*, and *ntrY* also increased

significantly, while transcripts of *dnaA* and *ftsZ1* genes decreased significantly (Fig.

2C). These results further confirm that the NtrX protein represses the transcription of

genes such as *ctrA* and *gcrA* and activates the transcription of genes such as *dnaA* and

ftsZ1, and that this regulation depends on the D53 residue on NtrX.

Regulation of protein levels of CtrA and GcrA by NtrX in *S. meliloti*. To determine whether protein levels of key cell cycle regulators are affected by the *ntrX* mutation, we first expressed His-tagged NtrX, CtrA and GcrA proteins in *E. coli*. After purification by nickel columns, anti-rabbit polyclonal antibodies were prepared for immunoblotting assays (35). Sm1021-synchronized cells were subcultured in LB/MC broth for shaking incubation. Cells were collected every half hour and lysed for immunoblotting assays. The results showed a varying trend of increasing first and then decreasing NtrX protein levels and a maximum occurring in cells subcultured for 1.5 h (Fig. 3A). Unlike Sm1021, the total NtrX protein level in SmLL1 cells was apparently reduced and tended to increase gradually at different growth times (Fig. 3A). Contrary to the NtrX protein, the change trend of CtrA and GcrA protein levels in Sm1021 first decreased and then increased. The levels of these two proteins were apparently increased in SmLL1 cells compared to Sm1021 cells, (Fig. 3A). These results indicate that SmLL1 is a down-regulated mutant of *ntrX* and that NtrX protein levels are negatively correlated with CtrA and GcrA proteins.

We evaluated the NtrX protein level in cells of depleted strains grown in LB/MC broth by immunoblotting and found that cells cultured in the broth containing IPTG for 1 to 3 h expressed high levels of NtrX protein (Fig. 3B). CtrA and GcrA protein levels were highly expressed in cells cultured in LB/MC broth without IPTG, whereas the levels of these two proteins were reduced in cells cultured in broth containing IPTG for 1 to 3 h (Fig. 3B). These results also prove that NtrX protein levels are negatively correlated with CtrA and GcrA proteins.

To determine whether the D53 residue on the NtrX protein affects the protein levels of CtrA and GcrA, we performed immunoblot assays on Sm1021/p*ntrX*^{D53E} and Sm1021/p*ntrX* cells. The results showed that the protein levels of NtrX and NtrX^{D53E} increased significantly when cultured in LB/MC broth containing IPTG for 2-3 h (Fig. 3C). Under the same culture conditions, the protein levels of CtrA and GcrA were apparently reduced in Sm1021/p*ntrX* cells from the same cultured condition, while they were elevated in Sm1021/p*ntrX*^{D53E} cells (Fig. 3C). These results reaffirm that the level of NtrX protein is negatively correlated with CtrA and GcrA proteins,

depending on D53 residue on the NtrX protein.

The 53rd aspartate residue as a phosphorylation site of *S. meliloti* NtrX. The

homologous NtrX proteins in α-proteobacteria are composed of REC and DNA

binding domains. The three-dimensional structure of the NtrX protein from *B. abortus*

has been partially resolved (27-28). Using this as a template, we reconstructed the 3D

structure of the NtrX protein from *S. meliloti* and found that there were 5 α-helices

and 5 β-sheets connected by loops in the REC domain (Figure 4A-B). The conserved

D53 is located at the end of the third β-sheet and can form a negative charge center

with D9, D10 and E11 (Figure 4A-B). Mutation of D53 into A and N, but not E, may

disrupt the formation of this negative charge center.

The NtrY histidine kinase can phosphorylate NtrX from *B. abortus in vitro* (27-28).

We also expressed and purified the NtrY kinase domain and NtrX protein of *S.*

meliloti in *E. coli* for *in vitro* phosphorylation assays. Through Phos-Tag gel detection,

we found that the NtrY kinase domain can be autophosphorylated and can also

phosphorylate the NtrX protein *in vitro* (Fig. 4C). After mutating D53 to E,

phosphorylated NtrX protein could not be detected (data not shown). To further

determine whether NtrX was phosphorylated in cells, both Phos-Tag gel and western

blotting analysis were used. The results showed that the ratio of phosphorylated NtrX

protein compared to non-phosphorylated protein in both Sm1021 and SmLL1

remained unchanged, though fewer NtrX proteins were detected in SmLL1 compared

to SmLL1 (Fig. 4D). To further verify that the D53 residue is the phosphorylation site

of the NtrX protein, we applied the same method to analyze the phosphorylated NtrX

protein level of Sm1021/p$ntrX^{D53E}$ cells cultured in LB/MC/IPTG broth. The results

showed that the ratio of phosphorylated NtrX protein compared to unphosphorylated

protein in Sm1021/p$ntrX$ cells tended to increase, while the ratio in Sm1021/p$ntrX^{D53E}$

cells tended to decrease (Fig. 4D). Our results suggest that NtrX can be

phosphorylated in *S. meliloti* with the phosphorylation site being the D53 residue.

Direct binding of phosphorylated NtrX protein to the promoter regions of key

cell cycle regulatory genes. To determine whether the NtrX protein of *S. meliloti*

directly regulates the expression of cell cycle regulatory genes, we used anti-NtrX

polyclonal antibodies to perform chromatin immunoprecipitation experiments.

Sequencing results showed that a total of 82 DNA fragments were specifically

precipitated from Sm1021 cells, 60 of which were derived from the chromosome,

while the other 22 fragments originated from the plasmids SymA (10) and SymB (12,

Table S1). After careful sequence analysis of the precipitated DNA fragments, we

found that the promoter regions of some cell cycle regulatory genes (such as *ctrA*,

dnaA and *ftsZ1*) were enriched (Fig. 5A). Using the MEME program to scan the

precipitated DNA fragments, we found that some gene promoter regions contained

CAN_5TG motifs, which are similar to the characteristic sequence of the *ntrY* gene

promoter ($CAAN_{3-5}TTG$) reported in *B. abortus* (28). Therefore, we used this

sequence as a primer to scan the whole genomic DNA of *S. meliloti* 1021 by running

the   genome-scan   dna-pattern   program   on   the   RSAT   website

(http://embnet.ccg.unam.mx/rsat), and a total of 2155 $CAAN_{1-5}TTG$ motifs were

identified (Table S2). Fourteen of these motifs are located in the promoter regions of

some cell cycle regulatory genes, such as *ctrA*, *danA*, *gcrA* and *ftsZ1*(Fig. 5B). These

results suggest that the NtrX protein may recognize $CAAN_{1-5}TTG$ motifs in the

promoter regions of some cell cycle regulatory genes to directly regulate their

transcription. To verify the ChIP-Seq results, we applied quantitative PCR to evaluate

the level of genomic DNA fragments from Sm1021 cells enriched with anti-NtrX

polyclonal antibodies. The results showed that the promoter regions of *ctrA*, *dnaA*,

gcrA and *ftsZ1* genes were precipitated to different degrees (Fig. 5C), indicating that

the NtrX protein in Sm1021 cells can interact directly with the promoter regions of

the aforementioned cell cycle regulatory genes.

In Sm1021, the promoter region of the *ntrY* gene can directly interact with the NtrX

protein (Fig. 5A-C), which is consistent with the report in *B. abortus* (28). To further

confirm the above findings, we synthesized a biotin-labeled *ntrY* gene promoter DNA

probe (containing two **CAAN$_{3-5}$TTG** motifs: **CAA**CACCG**TTG** and

CAATGCG**TTG**) for gel retardation assays. The results showed that phosphorylated

NtrX specifically bound to it, forming two protein-DNA complexes (Fig. 5D). To

determine whether the D53 phosphorylation site of the NtrX protein is involved in the

protein-DNA binding reaction, we replaced the phosphorylated NtrX protein with the

NtrX^{D53E} protein. The gel retardation results showed almost no protein-DNA complex

formation (Fig. 5D), indicating that phosphorylation at position D53 is essential for

the binding of NtrX to the *ntrY* promoter region. The same method was used to

analyze the binding ability between the phosphorylated NtrX protein and the biotin-

labeled *dnaA* promoter DNA probe (containing the **CAA**AACCC**TTG** motif) and

found that they bound specifically to form a protein-DNA complex (Fig. 5F). We

mutated the **CAA**AACCC**TTG** motif of the DNA probe to **CGG**AACCC**CCG**, and

found that the mutant probe virtually did not bind to the phosphorylated NtrX protein

(Fig. 5F), indicating that the AA or TT bases conserved in the **CAA**AACCC**TTG**

motif of the *dnaA* promoter region are required for the probe binding to the

phosphorylated NtrX protein. In addition, we used gel retardation assays to confirm

that the phosphorylated NtrX protein can specifically bind to biotin-labeled *ctrA*, *gcrA*

and *ftsZ1* promoter probes (each containing a **CAA**N$_{1-5}$**TTG** motif: **CAACCTTG**,

CAAACC**TTG** and **CAACTTG**), and at least one protein-DNA complex is formed,

respectively (Fig. 5E, G and S3). These results indicate that the phosphorylated NtrX

protein can bind specifically to the promoter regions of *ctrA*, *gcrA*, *dnaA* and *ftsZ1 in*

vitro.

The role of *S. meliloti* NtrX proteins in *Agrobacterium tumefaciens*. The NtrX

proteins from *A. tumefaciens* C58 and *S. meliloti* 1021 are highly homologous (Fig.

4A-B). To determine whether the function of NtrX protein in regulating cell division

is conserved in bacteria, we introduced plasmids expressing *S. meliloti* NtrX or its

substitution mutants into *A. tumefaciens* C58. By observing the growth phenotype of

the transformed strain on the LB/IPTG agar plate, we found that the strain carrying

p*ntrX* or p*ntrX*^{D53E} grew slower than the strain with the control vector, p*ntrX*^{D53A} or

p*ntrX*^{D53N} (Fig. S4), indicating that the expression of NtrX or NtrX^{D53E} (partially

retained functions) proteins can suppress cell division of *A. tumefaciens* C58, while

suggesting that the NtrX protein of *A. tumefaciens* C58 is also involved in the cell

cycle regulation.

Since NtrX recognizes the conserved $CAAN_{1-5}TTG$ motif, do the promoter regions of

cell cycle regulatory genes from *A. tumefaciens* C58 contain these motifs? To answer

this question, we used the sequence as a primer to scan the whole genomic DNA of *A.*

tumefaciens C58 by running the genome-scan dna-pattern program on the RSAT

website (http://embnet.ccg.unam.mx/rsat), and a total of 2562 $CAAN_{1-5}TTG$ motifs were

identified. Among them, 9 motifs were located in the promoter regions of the cell

cycle regulatory genes *ctrA*, *dnaA*, *gcrA*, *ftsZ1* and *ftsZ2*, respectively (Fig. S5). This

result suggests that the NtrX protein in *A. tumefaciens* C58 may be involved in cell

cycle regulation by recognizing these motifs in the gene promoter regions.

DISCUSSION

In symbiotic nitrogen-fixing bacteria, rhizobia, the level of combined nitrogen as a

signal not only regulates the expression of nitrogen-fixing genes, but also can affect

cell growth and division. However, the molecular mechanism by which combined

nitrogen levels regulate bacterial cell division is unclear. This work first revealed in *S.*

meliloti that the phosphorylated nitrogen metabolism regulator NtrX directly regulates

the transcription of cell cycle regulatory genes such as *ctrA*, *gcrA*, *dnaA* and *ftsZ1* by

specifically recognizing the characteristic sequence of the promoter region ($CAAN_{1-}$

$_5TTG$) and promotes cell division (Fig. 6), which provides a preliminary answer to

the above question.

NtrX is a bacterial cell cycle regulator. Previous studies have suggested that NtrX is a

regulator of nitrogen metabolism in bacterial cells because its mutants affect the

utilization of nitrogen sources (17-20, 23, 25). Decreased utilization of nitrogen

sources would inevitably lead to weakened nucleic acid and protein synthesis, which

would subsequently affect the growth and proliferation of bacterial cells. This is one

explanation for the effect of NtrX on bacterial cell division. In *E. chaffeensis* cells,

NtrX affected the stability of CtrA protein through a post-translational mechanism

(25), indicating that NtrX may act directly on the cell cycle regulatory system for

regulating cell division. This work was carried out using *S. meliloti* as the study

material to reveal the transcriptional control mechanism of the cell cycle regulatory genes through which NtrX protein affects cell division for the first time. This conclusion is supported by multiple experimental evidences: 1) all three groups of *ntrX* gene mutant materials are defective in bacterial growth, cell division and genomic DNA synthesis (Fig. 1); 2) the transcript levels of cell cycle regulatory genes are differentially altered in *ntrX* mutants (Fig. 2; 29); 3) the protein levels of CtrA and GcrA are altered in the mutant and they were negatively correlated with the level of NtrX protein (Fig. 3); 4) the NtrX protein is phosphorylated *in vivo* and bind directly to the promoter regions of *ctrA*, *gcrA*, *dnaA* and *ftsZ1* (Fig.4-5). The degree of cell division defects and differential gene expression may be related to the level of *ntrX* gene expression in different genetic materials and the functional status of the NtrX protein. The ability of phosphorylated NtrX in binding to DNA may be associated with the difference in the composition of the target sequence (including the recognition site).

NtrX affects cell division through transcriptional regulation of the cell cycle control system, which may also exist in other α-proteobacteria. *A. tumefaciens* and *S. meliloti* belong to the *Rhizobium* family, and their NtrX proteins are highly homologous (Fig.4A-B). We initially speculated that *S. meliloti*'s NtrX might play the same role in *A. tumefaciens* cells, but heterologous expression results showed that *S. meliloti*'s NtrX or NtrX^{D53E} proteins suppressed the growth and proliferation of *A. tumefaciens* cells, while NtrX^{D53A} or NtrX^{D53N} protein did not (Fig. S4). This may reflect that the different modes of action of NtrX proteins in these two species of bacteria, which may include requirements for NtrX phosphorylation, target genes for transcriptional regulation, and even the number and location of recognition sites for the same cell cycle regulatory genes (Fig. S4). NtrX in *C. crescentus* has been reported not to affect cell growth and division during the logarithmic phase[26]. By comparing the amino acid sequences of NtrX proteins, we found that the NtrX protein of *C. crescentus* is not highly homologous to that of *S. meliloti* (Fig. 4A-B), so the functions of NtrX proteins may differ between the two species of bacteria. We attempted to express the

NtrX protein of *S. meliloti* in *C. crescentus*, but failed to screen transformed cells,

whereas cells expressing the NtrX[D53] substitution protein were successfully screened

(data not shown), suggesting that the endogenous NtrX protein of *C. crescentus* may

be an inactivated mutant protein whose function may be replaced by other proteins.

Whether the NtrX protein of *B. abortus* affects cell division requires further study.

The NtrY kinase derived from *S. meliloti* and *C. crescentus* can phosphorylate NtrX

protein *in vitro* (23; Fig.4-1), but there is no experimental evidence to demonstrate the

phosphorylation of NtrX by NtrY in the cell. Our previous study showed no

correlation between the genetic phenotypes of *ntrY* and *ntrX* mutants in *S. meliloti*

1021: the *ntrY* deletion mutant did not affect the growth and proliferation of bacterial

cells and inoculated hosts of alfalfa were not defective in symbiotic nitrogen fixation,

whereas the SmLL1 mutant grew slowly in LB/MC and inoculated host plants formed

inefficient nitrogen-fixing nodules (29). Therefore, the upstream kinase of NtrX in the

cell may not be NtrY. This inference is consistent with the repression of *ntrY* gene

expression by NtrX in *S. meliloti* (29; Fig. 2A). Furthermore, the study results from *B.*

abortus indicate that the expression of *ntrY* gene is induced by micro-oxygen and is

involved in micro-oxygen signal transduction (23). The primary function of NtrX in

bacteria is considered to regulate nitrogen metabolism. The nitrogen limitation signal

transduction in bacteria is mainly mediated by the NtrB/NtrC system (33), so it cannot

be ruled out that the NtrB/NtrC system can regulate the expression of *ntrX* under

nitrogen lacking conditions. Under nitrogen rich conditions, the activity of NtrX may

be regulated by an unknown kinase, which may be able to sense the level of ammonia

nitrogen.

Finally, *S. meliloti* can infect alfalfa and induce the formation of nitrogen-fixing

nodules. On the root surface of the plant, rhizobia cells grow and divide at the top of

the extended infection thread and in the nodule infection zone. Whether these division

events are regulated by NtrX requires further study.

MATERIALS AND METHODS

Strains and culture medium. *Escherichia coli* DH5α and BL21 were cultured in LB medium at 37 °C. *S. meliloti* (including Sm1021, SmLL1 and derivatives) (29) were cultured in LB/MC medium at 28 °C. MOPS-GS broth was utilized for the cell synchronization of *S. meliloti* (34). *A. tumefaciens* C58 was cultured in LB medium at 28 °C. The following antibiotics were added to the medium: kanamycin, 50 μg/ml; gentamicin, 10 μg/ml; chloramphenicol, 30 μg/ml; neomycin, 200 μg/ml; streptomycin, 200 μg/ml; tetracycline, 10 μg/ml.

Recombinant plasmid construction. Primers P*ntrX*1 and P*ntrX*2 bearing *Hind*III and *Xba*I digestion sites were used to amplify the *S. meliloti ntrX* gene (Table S3). The *ntrX* gene fragment was amplified using overlapping PCR primers NMF and NMR with substitution of aspartate for glutamate, asparagine or alanine (Table S3). Overlapping PCR was performed as described by Wang(29) in 2013. The PCR products were digested with *Hind*III and *Xba*I (Thermo) and ligated with digested pSRK-Gm (31) to obtain the recombinant plasmids p*ntrX*, p*ntrX*^{D53E}, p*ntrX*^{D53A} and p*ntrX*^{D53N}. Upon introducing p*ntrX* into SmLL1, the depletion strain was screened on LB/MC agar plates containing 1 mM IPTG. Primers for P*ntrY*k1, P*ntrY*k2, P*ntrX*1^{D53E}, P*ntrX*2^{D53E}, P*ctrA*1, P*ctrA*2, P*gcrA*1 and P*gcrA*2 were used to amplify the NtrY kinase domain, *ntrX*^{D53E}, *ctrA* and *gcrA*, respectively (Table S3). The PCR fragments were digested with the appropriate restriction enzymes and ligated into pET28b to harvest p*ntrY*k, p*ntrX*^{D53E}, p*ctrA* and p*gcrA* for recombinant protein purification.

Bacterial cell synchronization. De Nisco's method was used for bacterial cell synchronization (34). A sample of *S. meliloti* cells was selected from an agar plate, placed in 5 ml LB/MC broth and grown overnight. A 100-μl aliquot of the bacterial culture was transferred to 100 ml LB/MC broth and incubated overnight until $OD_{600} =$ 0.1-0.15; the culture was centrifuged (6,500 rpm, 5 min, 4°C) and the cell pellet was washed twice with sterilized 0.85% NaCl solution. Cells were resuspended in MOPS-GS synchronization medium and grown at 28°C for 270 min. After centrifugation, cells were washed twice with sterilized 0.85% NaCl solution, resuspended in LB/MC broth, and cultured at 28°C.

RNA extraction and purification. 20 ml of bacterial cultures were collected by centrifugation (6,000 rpm, 5 min, 4°C) and the cells were washed twice with DEPC-treated water. RNA extraction was performed using 1 ml of Trizol (Life Technology) for RNA extraction, and its purification was performed as described by Wang (29).

qRT-PCR and qPCR. RNA reverse transcription was performed using the PrimeScript RT reagent Kit with gDNA Eraser (TaKaRa). The qPCR reaction system included the following: SYBR® Green Real-time PCR Master Mix, 4.75 μl; cDNA or DNA, 0.25 μl; 10 pmol/μl primers, 0.5 μl; ddH$_2$O, 4.5 μl. The reaction procedure is as follows: 95°C, 5 min; 95°C, 30 s; 55°C, 30 s; 72°C, 1 kb/min. The selected reference gene was Smc00128. The 2$^{-\Delta\Delta CT}$ method was applied to the analysis of gene expression levels. All primers are listed in Table S3.

Chromatin immunoprecipitation (ChIP). ChIP was performed as described by Pini (4). Anti-NtrX antibodies prepared by Wenyuange (Shanghai) were used for ChIP assays (35).

Flow cytometry. Four milliliters of fresh bacterial cultures were centrifuged (6,000 rpm, 5 min, 4°C) and the cell pellets were washed twice with a 0.85% NaCl solution (stored at 4°C). De Nisco's flow cytometry protocol of was used (34). Each sample was assessed using a MoFlo XDF (Beckman Coulter) flow cytometer, and the results were analyzed using Summit 5.1 software (Beckman Coulter).

EMSA (electrophoretic mobility shift assay). EMSA was performed as described by Zeng (35). The LightShift^{TM} Chemiluminescent EMSA Kit (Thermo Fisher) was applied in the assays. Probes for *ntrY*, *ctrA*, *dnaA*, *gcrA* and *ftsZ1* promoter DNA labeled with biotin were synthesized (Invitrogen); the probes are listed in Table S3. Phosphorylated NtrXr proteins (from Ni^{2+} column purification) used for these assays were prepared. 6 μg of purified protein was mixed with 2 mM of acetyl phosphate (Sigma) in 100 μl of phosphorylation buffer (50 mM Tris-HCl pH 7.6, 50 mM KCl, 20 mM MgCl$_2$), and incubated for 1 h at 28°C. Acetyl phosphate was removed using an ultra-filtration tube (Millipore) resolved in 100 μl of phosphorylation buffer.

NtrX phosphorylation assay. 1 mg of His-NtrX fusion protein (NtrXr) and 1 mg of His-NtrY kinase domain fusion protein (NtrY-Kr) purified through a Ni^{2+} column

were used for *in vitro* NtrX phosphorylation assays. 2 mM of acetyl phosphate (Sigma) was mixed with 300 μg of NtrY-Kr in 1 ml of phosphorylation buffer and then incubated for 1 h at room temperature. Acetyl phosphate was removed using an ultra-filtration tube (Millipore) and resolved in 1 ml of phosphorylation buffer. 1, 3 and 10 μg of phosphorylated NtrY-Kr protein were added to 200 μl of phosphorylation buffer having 10 μg of NtrXr, and incubated overnight at 28℃. Samples were separated by a Phos-Tag gel (Mu Biotechnology, Guangzhou). Phosphorylated NtrX levels from *S. meliloti* cells were separated in a Phos-Tag gel. *S. meliloti* 1021 carrying p*ntrX* or p*ntrX*^{D53E} were cultured in LB/MC broth induced by 1 mM IPTG for 1 to 3 h. ~1 μg of total protein was input for each sample. The total protein was transferred to PVDF, and then detected by anti-NtrX antibodies from Western blotting assays (13).

Western blotting. Western blotting was performed as described by Tang (36). Proteins were detected using an ECL fluorescence colorimetric kit (Tiangen) and visualized using a Bio-Rad Gel Doc XR. Anti-NtrX antibodies were prepared at Wenyuange, Shanghai, China (35); Anti-CtrA and anti-GcrA antibodies were prepared by Hua'an Biotech, Hangzhou, China.

Microscopy. A 5-μl aliquot of fresh *S. meliloti* culture (OD$_{600}$ = 0.15) was placed on a glass slide and covered with a cover glass. The slide was slightly baked near the edge of the flame of an alcohol lamp for a few seconds. Cells expressing the pHC60 plasmid were observed in GFP mode (32), and images were acquired using a CCD camera Axiocam 506 color (Zeiss). Exposure times were set to 10 ms and 1,000 ms in order to capture bacterial morphology and motility, respectively. Images were analyzed with ZEN 2012 software (Zeiss).

ChIP sequencing. ChIP-Seq was performed by Bohao Biotech, Shanghai. The original data were evaluated by Bohao Biotech based on the standard protocol. Genome mapping was then performed using bowtie2 (version: 2-2.0.5) for local alignment of clean reads (37). The original genomic data of *S. meliloti* 1021 was obtained from NCBI. IGV software was employed to assess specific enrichment data based on ChIP-Seq results, and screenshots of peak maps were generated (38).

Bioinformatic analysis. Genome-wide CAAN$_{1-6}$TTG *cis*-acting elements of bacteria

were evaluated using the genome-scan dna pattern program on the RAST online

server (http://embnet.ccg.unam.mx/rsat) (39). The BDGP program was applied for

promoter prediction (http://www.fruitfly.org/seq_tools/promoter.html)

The NtrX protein 3D structure was reconstructed in Swiss-Model using the 4d6y

template (the 3D structure of the NtrX receiver domain from *Brucella*) in PDB (40).

The 3D structure of the NtrX receiver domains was analyzed by Pymol (Delano

Scientific).

ACKNOWLEDGMENTS

This research was supported by the Natural Science Foundation of China (31570241

to L.L).

AUTHOR CONTRIBUTIONS

L. L. designed research; S. X., F. A., X. Y., L. H., S. Z., L. Z., W. Z., and N. L.

performed research; S.X., F.A., J. Y., L. Y. and L. L. analyzed data; L. L. and K. O.

wrote the paper.

REFERENCES

1. Laub MT, Shapiro L, McAdams HH. 2007. Systems biology of Caulobacter. Annu Rev

Genet 41:429-41.

2. Laub MT, McAdams HH, Feldblyum T, Fraser CM, Shapiro L. 2000. Global analysis of

549   the genetic network controlling a bacterial cell cycle. Science 290:2144-8.

550   3.   Skerker JM, Laub MT. 2004. Cell-cycle progression and the generation of asymmetry in

551   Caulobacter crescentus. Nat Rev Microbiol 2:325-37.

552   4.   Panis G, Murray SR, Viollier PH. 2015. Versatility of global transcriptional regulators in

553   alpha-Proteobacteria: from essential cell cycle control to ancillary functions. FEMS

554   Microbiol Rev 39:120-33.

555   5.   Jacobs C, Ausmees N, Cordwell SJ, Shapiro L, Laub MT. 2003. Functions of the CckA

556   histidine kinase in Caulobacter cell cycle control. Mol Microbiol 47:1279-90.

557   6.   Biondi EG, Reisinger SJ, Skerker JM, Arif M, Perchuk BS, Ryan KR, Laub MT. 2006.

558   Regulation of the bacterial cell cycle by an integrated genetic circuit. Nature 444:899-

559   904.

560   7.   Jones KM, Kobayashi H, Davies BW, Taga ME, Walker GC. 2007. How rhizobial

561   symbionts invade plants: the Sinorhizobium-Medicago model. Nat Rev Microbiol

562   5:619-33.

563   8.   Van de Velde W, Zehirov G, Szatmari A, Debreczeny M, Ishihara H, Kevei Z, Farkas A,

564   Mikulass K, Nagy A, Tiricz H, Satiat-Jeunemaître B, Alunni B, Bourge M, Kucho K,

565   Abe M, Kereszt A, Maroti G, Uchiumi T, Kondorosi E, Mergaert P. 2010. Plant

566   peptides govern terminal differentiation of bacteria in symbiosis. Science 327:1122-6.

567   9.   Farkas A, Maróti G, Durgő H, Györgypál Z, Lima RM, Medzihradszky KF, Kereszt A,

568   Mergaert P, Kondorosi É. 2014. Medicago truncatula symbiotic peptide NCR247

569   contributes to bacteroid differentiation through multiple mechanisms. Proc Natl Acad

570   Sci U S A 111:5183-8.

10. Penterman J, Abo RP, De Nisco NJ, Arnold MF, Longhi R, Zanda M, Walker GC. 2014.

Host plant peptides elicit a transcriptional response to control the Sinorhizobium

meliloti cell cycle during symbiosis. Proc Natl Acad Sci U S A 111:3561-6.

11. Montiel J, Downie JA, Farkas A, Bihari P, Herczeg R, Bálint B, Mergaert P, Kereszt A,

Kondorosi É. 2017. Morphotype of bacteroids in different legumes correlates with the

number and type of symbiotic NCR peptides. Proc Natl Acad Sci U S A 114:5041-

5046.

12. Kobayashi H, De Nisco NJ, Chien P, Simmons LA, Walker GC. 2009. Sinorhizobium

meliloti CpdR1 is critical for co-ordinating cell cycle progression and the symbiotic

chronic infection. Mol Microbiol 73:586-600.

13. Pini F, Frage B, Ferri L, De Nisco NJ, Mohapatra SS, Taddei L, Fioravanti A, Dewitte F,

Galardini M, Brilli M, Villeret V, Bazzicalupo M, Mengoni A, Walker GC, Becker A,

Biondi EG. 2013. The DivJ, CbrA and PleC system controls DivK phosphorylation and

symbiosis in Sinorhizobium meliloti. Mol Microbiol 90:54-71.

14. Xue S, Biondi EG. 2019. Coordination of symbiosis and cell cycle functions in

Sinorhizobium meliloti. Biochim Biophys Acta Gene Regul Mech 1862:691-696.

15. Robinson MD, Oshlack A. 2010. A scaling normalization method for differential

expression analysis of RNA-seq data. Genome Biol 11:R25.

16. Schallies KB, Sadowski C, Meng J, Chien P, Gibson KE. 2015. Sinorhizobium meliloti

CtrA Stability Is Regulated in a CbrA-Dependent Manner That Is Influenced by

CpdR1. J Bacteriol 197:2139-2149.

17. Pawlowski K, Klosse U, de Bruijn FJ. 1991. Characterization of a novel Azorhizobium

caulinodans ORS571 two-component regulatory system, NtrY/NtrX, involved in

nitrogen fixation and metabolism. Mol Gen Genet 231:124-38.

18. Nogales J, Campos R, BenAbdelkhalek H, Olivares J, Lluch C, Sanjuan J. 2002.

Rhizobium tropici genes involved in free-living salt tolerance are required for the

establishment of efficient nitrogen-fixing symbiosis with Phaseolus vulgaris. Mol Plant

Microbe Interact 15:225-32.

19. Ishida ML, Assumpção MC, Machado HB, Benelli EM, Souza EM, Pedrosa FO. 2002.

Identification and characterization of the two-component NtrY/NtrX regulatory system

in Azospirillum brasilense. Braz J Med Biol Res 35:651-61.

20. Bonato P, Alves LR, Osaki JH, Rigo LU, Pedrosa FO, Souza EM, Zhang N, Schumacher

J, Buck M, Wassem R, Chubatsu LS. 2016. The NtrY-NtrX two-component system is

involved in controlling nitrate assimilation in Herbaspirillum seropedicae strain SmR1.

Febs j 283:3919-3930.

21. Gregor J, Zeller T, Balzer A, Haberzettl K, Klug G. 2007. Bacterial regulatory networks

include direct contact of response regulator proteins: interaction of RegA and NtrX in

Rhodobacter capsulatus. J Mol Microbiol Biotechnol 13:126-39.

22. Lemmer KC, Alberge F, Myers KS, Dohnalkova AC, Schaub RE, Lenz JD, Imam S,

Dillard JP, Noguera DR, Donohue TJ. 2020. The NtrYX Two-Component System

Regulates the Bacterial Cell Envelope. mBio 11.

23. Carrica Mdel C, Fernandez I, Martí MA, Paris G, Goldbaum FA. 2012. The NtrY/X two-

component system of Brucella spp. acts as a redox sensor and regulates the

expression of nitrogen respiration enzymes. Mol Microbiol 85:39-50.

24. Atack JM, Srikhanta YN, Djoko KY, Welch JP, Hasri NH, Steichen CT, Vanden Hoven

RN, Grimmond SM, Othman DS, Kappler U, Apicella MA, Jennings MP, Edwards JL,

McEwan AG. 2013. Characterization of an ntrX mutant of Neisseria gonorrhoeae

reveals a response regulator that controls expression of respiratory enzymes in

oxidase-positive proteobacteria. J Bacteriol 195:2632-41.

25. Cheng Z, Lin M, Rikihisa Y. 2014. Ehrlichia chaffeensis proliferation begins with

NtrY/NtrX and PutA/GlnA upregulation and CtrA degradation induced by proline and

glutamine uptake. mBio 5:e02141.

26. Fernandez I, Sycz G, Goldbaum FA, Carrica MD. 2018. Acidic pH triggers the

phosphorylation of the response regulator NtrX in alphaproteobacteria. Plos One 13.

27. Fernández I, Otero LH, Klinke S, Carrica MDC, Goldbaum FA. 2015. Snapshots of

Conformational Changes Shed Light into the NtrX Receiver Domain Signal

Transduction Mechanism. J Mol Biol 427:3258-3272.

28. Fernández I, Cornaciu I, Carrica MD, Uchikawa E, Hoffmann G, Sieira R, Márquez JA,

Goldbaum FA. 2017. Three-Dimensional Structure of Full-Length NtrX, an Unusual

Member of the NtrC Family of Response Regulators. J Mol Biol 429:1192-1212.

29. Wang D, Xue H, Wang Y, Yin R, Xie F, Luo L. 2013. The Sinorhizobium meliloti ntrX

gene is involved in succinoglycan production, motility, and symbiotic nodulation on

alfalfa. Appl Environ Microbiol 79:7150-9.

30. Calatrava-Morales N, Nogales J, Ameztoy K, van Steenbergen B, Soto MJ. 2017. The

NtrY/NtrX System of Sinorhizobium meliloti GR4 Regulates Motility, EPS I Production,

and Nitrogen Metabolism but Is Dispensable for Symbiotic Nitrogen Fixation. Mol

637   Plant Microbe Interact 30:566-577.

638   31. Khan SR, Gaines J, Roop RM, 2nd, Farrand SK. 2008. Broad-host-range expression

639   vectors with tightly regulated promoters and their use to examine the influence of

640   TraR and TraM expression on Ti plasmid quorum sensing. Appl Environ Microbiol

641   74:5053-62.

642   32. Cheng HP, Walker GC. 1998. Succinoglycan is required for initiation and elongation of

643   infection threads during nodulation of alfalfa by Rhizobium meliloti. Journal of

644   Bacteriology 180:5183-5191.

645   33. Szeto WW, Nixon BT, Ronson CW, Ausubel FM. 1987. Identification and

646   characterization of the Rhizobium meliloti ntrC gene: R. meliloti has separate

647   regulatory pathways for activation of nitrogen fixation genes in free-living and

648   symbiotic cells. J Bacteriol 169:1423-32.

649   34. De Nisco NJ, Abo RP, Wu CM, Penterman J, Walker GC. 2014. Global analysis of cell

650   cycle gene expression of the legume symbiont Sinorhizobium meliloti. Proc Natl Acad

651   Sci U S A 111:3217-24.

652   35. Zeng S, Xing S, An F, Yang X, Yan J, Yu L, Luo L. 2020. Sinorhizobium meliloti NtrX

653   interacts with different regions of the visN promoter. Acta Biochim Biophys Sin

654   (Shanghai) 52:910-913.

655   36. Tang G, Xing S, Wang S, Yu L, Li X, Staehelin C, Yang M, Luo L. 2017. Regulation of

656   cysteine residues in LsrB proteins from Sinorhizobium meliloti under free-living and

657   symbiotic oxidative stress. Environ Microbiol 19:5130-5145.

37. Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. Genome Biol 10:R25.

38. Thorvaldsdottir H, Robinson JT, Mesirov JP. 2013. Integrative Genomics Viewer (IGV):

high-performance genomics data visualization and exploration. Briefings in

Bioinformatics 14:178-192.

39. van Helden J. 2003. Regulatory sequence analysis tools. Nucleic Acids Res 31:3593-6.

40. Arnold K, Bordoli L, Kopp J, Schwede T. 2006. The SWISS-MODEL workspace: a web-

based environment for protein structure homology modelling. Bioinformatics 22:195-

201.

FIGURE LEGENDS

Fig. 1. Cell division defects of *S. meliloti ntrX* mutants in LB/MC medium.

(**A, C, F**) Cell shapes and sizes of *ntrX* mutants under a light or fluorescence

microscope. Red arrows, abnormal cells; bars, 2μm in **A** and **C**; Bar, 100 μm in **F**. (**B,**

E, H) Genomic DNA content of *ntrX* mutants was determined by flow cytometry. 1C,

haploid; 2C, diploid; >2C, polyploid. (**D, G**) Growth curves of the *ntrX* depletion

strain and Sm1021/p*ntrX*^{D53E}. Error bars, ±SD. 1 mM IPTG was added to the LB/MC

broth. SmLL1, *ntrX*::pK18*mobsacB*; wt, Sm1021.

Fig. 2. Differential expression of cell cycle regulatory genes in *ntrX* mutants.

Transcript levels of cell cycle regulatory genes in *S. meliloti* cells were evaluated by

qRT-PCR. Error bars, ±SD. The depletion strain was used in **B**. The genetic

background was Sm1021 in **C**. 1 mM IPTG was added to the LB/MC broth in **B** and

C.

Fig. 3. Protein levels of NtrX, CtrA and GcrA in the *ntrX* mutant as evaluated by

Western blotting.

N/I, C/I or G/I, the blot intensity of NtrX, CtrA or GcrA (the larger one)/the blotting

intensity of the input proteins, as collected in the original data by Image J. The

depletion strain was used in **B**. The genetic background was Sm1021 in **C**. 1 mM

IPTG was added to the LB/MC broth in **B** and **C**.

Fig. 4. Phosphorylation of the D53 residue in the NtrX protein.

(**A**) Alignment of f NtrX receiver domain from four bacterial species. The amino acid

sequence of each protein was obtained from NCBI. Secondary structures of the

receiver domain are shown as green lines (loops), yellow arrows (β- sheets) and red

bars (α-helixes). Ba2308, *B. abortus* bv. 1 str. 2308; Sm1021, *S. meliloti* 1021; AtC58,

A. tumefaciens C58; CcN1000, *C. crescentus* N1000. (**B**) 3D structures of the NtrX

receiver domain from three bacterial species were reconstructed using *B. abortus*

homolog protein (PDB:4d6y) as a template in Swiss-Model. Electrostatic interactions

of the carboxyl group from the conserved 53rd aspartate residues are shown as Arabic

numbers via Pymol. (**C**) *In vitro* NtrX phosphorylation catalyzed by the NtrY kinase

domain. NtrY-Kr, His-NtrY kinase domain fusion protein; NtrXr-P, phosphorylated

NtrXr; Ac-Pi, acetyl phosphate. (**D-E**) Phosphorylated NtrX proteins from *S. meliloti*

cells detected in Phos-Tag gel and by Western blotting. *S. meliloti* 1021 carrying

p*ntrX* or p*ntrX*^{D53E} cultured in LB/MC broth was induced by 1mM IPTG for 1 to 3

h. **~** 1 μg of total protein was input for each sample. P/NP, intensity of

phosphorylated protein/intensity of non-phosphorylated protein, as collected in the

original data by Image J.

Fig. 5. Phosphorylated NtrX binds to the promoter regions of key cell cycle

regulatory genes.

(**A**) Peak maps showing the promoter regions of some cell cycle genes precipitated by

anti-NtrX antibodies in two independent ChIP-Seq experiments. (**B**) The conserved

motif CAAN $_{(1-6)}$ TTG in the promoters of cell cycle regulatory genes. *, Predicted

translation initiation as indicated by +1. (**C**) Abundance of DNA fragments

specifically precipitated by anti-NtrX antibodies, as determined by qPCR. (**D-G**) The

promoter regions of *ntrY* (**D**), *ctrA* (**E**), *dnaA* (**F**), and *gcrA* (**G**) containing at least one

CAAN$_{(1-6)}$TTG motif are directly bound to the phosphorylated NtrX protein *in vitro*,

as based on EMSA. His-NtrX^{D53E}, the His-NtrX fusion protein containing a

substitution of D53 (phosphorylation site) for E in (**D**). Probe P*dnaA*s is the DNA

probe P*dnaA*, where **CAA**AACCC**TT**G is replaced by **CGG**AACCC**CC**G in (**F**). D/P

complex, DNA-protein complex; competitor, DNA is probed without biotin labeling.

The amounts of His-NtrX proteins per probe (2 nM) are 3, 6 and 15 ng, respectively. P,

putative transcriptional start sites as predicted by the BDGP program. Blue bars,

probes for EMSA; red bars, putative binding sites.

Fig. 6. An NtrX-mediated transcriptional control system for cell cycle progression of

S. meliloti

SUPPORTING INFORMATION

Table S1 DNA fragments of Sm1021 cells precipitated by anti-NtrX antibodies

Table S2 NtrX binding sites predicted by RSAT

Table S3 DNA oligonucleotides used in this study

Fig. S1. Colony formation of Sm1021 expressing NtrX substituted with D53 on

LB/MC/IPTG agar plates.

Fig. S2. Differential expression of cell cycle regulatory genes in *ntrX* mutants.

Transcript levels of cell cycle regulatory genes in *S. meliloti* cells were evaluated by

qRT-PCR. Error bars, ±SD. The depletion strain was used in **B**. 1 mM IPTG was

added to the LB/MC broth in **B**.

Fig. S3. The promoter regions of *ftsZ1* containing one CAAN$_{(1-6)}$TTG motif bind

directly to the phosphorylated NtrX protein *in vitro*.

D/P complex, DNA-protein complex; competitor, DNA is probed without biotin

labeling. The amounts of His-NtrX proteins per probe (2 nM) are 3, 6 and 15 ng,

respectively. P, the putative transcriptional start site was predicted by the BDGP

program. Blue bars, probes for EMSA; red bars, putative binding sites.

Fig. S4. Growth phenotypes of *A. tumefaciens* expressing *S. meliloti* NtrX

substitutions.

Fig. S5. Distribution of CAAN$_{(1-5)}$TTG motifs in three α-proteobacterial species by

scanning genomic DNA in RSAT.

These sites can be recognized by NtrX to affect gene transcription. *, the predicted

translation start is indicated by +1.

Fig.1

Fig.2

Fig.3

Fig.4

Fig.5

Fig.6