Ex vivo **osteochondral test sy**stem with **control over cartilage defect depth – a pilot**

study to investigate the effect of oxygen tension and chondrocyte-based

treatments in chondral and full thickness defects in an organ model

Andrea Schwab [1], Alexa Buss [1], Oliver Pullig [1, 2], Franziska Ehlicke [1,$]

[1] University Hospital Wuerzburg, Department Tissue Engineering and Regenerative Medicine,

Roentgenring 11, 97070 Wuerzburg, Germany

[2] Fraunhofer Institute for Silicate Research ISC, Translational Center Regenerative Therapies,

Roentgenring 11, 97070 Wuerzburg, Germany

Andrea Schwab

• University Hospital Wuerzburg, Department Tissue Engineering and Regenerative Medicine

• Present address: AO Research Institute Davos, Clavadelerstrasse 8, 7270 Davos, Switzerland

E-Mail: andrea.schwab@aofoundation.org

Alexa Buss

• University Hospital Wuerzburg, Department Tissue Engineering and Regenerative Medicine,

Roentgenring 11, 97070 Wuerzburg, Germany

• Present address: Klinikum Würzburg Mitte gGmbH, Juliusspital, Juliuspromenade 19, 97070

Wuerzburg, Germany

E-Mail: alexa.buss@kwm-klinikum.de

Oliver Pullig

• University Hospital Wuerzburg, Department Tissue Engineering and Regenerative Medicine,

Roentgenring 11, 97070 Wuerzburg, Germany

• Fraunhofer Institute for Silicate Research ISC, Translational Center Regenerative Therapies,

Roentgenring 11, 97070 Wuerzburg, Germany

E-Mail: oliver.pullig@isc.fraunhofer.de

Franziska Ehlicke ($corresponding author)

• University Hospital Wuerzburg, Department Tissue Engineering and Regenerative Medicine,

Roentgenring 11, 97070 Wuerzburg, Germany

E-Mail: franziska.ehlicke@uni-wuerzburg.de

Phone: +49 931 3181900

<u>Running title:</u>

Cell-based chondral regeneration *ex vivo*

Abstract

<u>Objective</u>

Cartilage defect treatment strategies are dependent on the lesion size and severity. Osteochondral

explants models are a platform to test cartilage repair strategies *ex vivo*. Current models lack in mimicking

the variety of clinically relevant defect scenarios. In this controlled laboratory study, an automated device

(artificial tissue cutter, ARTcut®) was implemented to reproducible create cartilage defects with controlled

depth. In a pilot study, the effect of cartilage defect depth and oxygen tension on cartilage repair was

investigated.

<u>Design</u>

Osteochondral explants were isolated from porcine condyles. 4 mm chondral and full thickness defects

were treated with either porcine chondrocytes (CHON) or co-culture of 20 % CHON and 80 % MSC (MIX)

embedded in collagen hydrogel. Explants were cultured with tissue specific media (without TGF-β) under

normoxia (20 % O_2) and physiological hypoxia (2 % O_2). After 28 days, immune-histological stainings

(Collagen II and X, Aggrecan) were scored (modified Bern-score, 3 independent scorer) to quantitatively

compare treatments outcome.

<u>Results</u>

ARTcut® represents a software-controlled device for creation of uniform cartilage defects. Comparing the

scoring results of the MIX and the CHON treatment, a positive relation between oxygen tension and defect

depth was observed. Low oxygen tension stimulated cartilaginous matrix deposition in MIX group in

chondral defects and CHON treatment in full thickness defects.

<u>Conclusion</u>

ARTcut® has proved a powerful tool to create cartilage defects and thus opens a wide range of novel

applications of the osteochondral model, including the relation between oxygen tension and defect depth

on cartilage repair.

Keywords

cartilage defect, *ex vivo* model, cartilage test system, chondrocytes, MSC, collagen type I

hydrogel

1 Introduction

Defects of the articulating surface are frequently occuring diseases in the field of orthopedics,

traumatology and sports medicine [1]. Around 14 % of patients who suffered from a trauma induced knee

injury during their middle ages, including fractures of the tibia, fibula, femur, or patella, develop

osteoarthritis in the same knee joint at later age (>65 years) [2]. High risk of failure of more than 20 % in

cartilage defect repair require to rethink and improve current cartilage treatment concepts with the aim

for long-term repair of cartilage defects [3, 4].

Due to the avascular and aneural character of adult articular cartilage, large defects do not heal

spontaneously [5]. Defects that are classified according to International Cartilage Repair Society (ICRS)

grade III (defect depth >50 % cartilage depth) and IV (defect reaches subchondral bone) need surgical

intervention to repair [6]. Gold standard for defects larger than 2-3 cm² is the autologous chondrocyte

implantation (ACI), resulting in similar outcome as microfracture [7, 8]. In a recent data analysis for

cartilage defects, matrix-associated chondrocyte implantation (M-ACI) showed a significant lower

reoperation rate than microfracture 2 years post-op [9]. M-ACI as well as ACI require two surgical

interventions; the first to take a cartilage tissue sample for subsequent chondrocyte isolation and the

second one for injection of isolated *in vitro* expanded autologous chondrocytes into defect site.

However, (M-)ACI has two main limitations: Due to the limited number of healthy and non-degenerated

chondrocytes, cells have to be expanded *in vitro* to achieve sufficient number of cells needed for the

implantation (ACI: 1 million cells per cm², M-ACI: 20 million cells per cm³) [10-12]. Further, chondrocyte-

based cartilage treatments, as well as microfracture techniques, result in formation of fibrocartilage,

characterized by its inferior mechanical properties and thus lacking functional restoration compared to

healthy hyaline cartilage [4, 13-17]. One approach to overcome the limitation of cell number for (M-)ACI

treatment are MSC-chondrocyte co-cultures, *e.g.*, 80 % MSC and 20 % chondrocytes, that reduces

autologous chondrocyte cell number at same total cell density. Co-cultures of MSCs and chondrocytes

have shown to increase cartilaginous matrix production and reduce hypertrophy, associated with MSCs

during chondrogenic differentiation [18, 19].

To bring cartilage treatments to the next level, cartilage repair needs to be studied and understood in

more detail, starting with basic research questions in a physiologically relevant environment.

An *ex vivo* cartilage defect test system based on osteochondral explants was established by Schwab *et al.*

[20]. This test system represents a valuable *ex vivo* platform for biomaterial evaluation in critical size

trauma-induced cartilage defects in terms of biocompatibility, biomaterial tissue integration and cartilage

repair with higher throughput compared to *in vivo* models. Separated media compartments of the culture

device allow for controlled, tissue- and cell-specific nutrient supply during *ex vivo* culture of osteochondral

explants. This model also allows direct comparison of different treatment approaches under controlled

conditions and has been shown to stimulate cartilage-like tissue formation of chondrocytes or MSCs in

osteochondral lesions [21, 22].

However, this defect model was limited to full thickness defects created with a biopsy punch. To date, the

creation of defects that do not fully penetrate the cartilage with the help of a biopsy punch or scalpel

suffer from low reproducibility and are dependent on the operator. Therefore, an automated device with

control over drilling depth to create defects with high reproducibility can overcome this limitation to study

chondral wound healing. Of note, the drilling should not harm surrounding cartilage tissue and cell

viability, neither by mechanical tissue disruption nor by friction induced heating. For the creation of

defects in hard tissues like cartilage or even bone, there is no device available for automated defect

creation that meets the above-mentioned requirements.

Following, in the present study the defect creation of the *ex vivo* defect model introduced by Schwab *et*

al. was modified by implementation of a semi-automated drilling device, originally developed to

mechanically induce standardized and reproducible wounds in full thickness skin equivalents [23]. Key

features of this artificial tissue cutter (ARTcut®) are the sensor controlled optical barrier in combination

with a moveable milling machine along x-, y- and z-axis. The light-barrier controlled drilling allows for

creation of defects with defined depths in a reproducible set-up. Creation of tissue defects with ARTcut®

can be performed under sterile conditions for subsequent *in vitro* or *ex vivo* culture with the possibility to

adjust defect geometry and control of defect creation process.

Another fact that has been neglected in the studies by Schwab *et al.* is the low oxygen tension present in

the articular capsule [24]. It has been shown in literature that low oxygen tension is an important stimulus

in chondrogenesis and increases the chondrogenic potential of articular cartilage progenitor cells and

MSCs [24-26]. Taken together, the present study aimed to implement the ARTcut® as tool for automated

wounding to the *ex vivo* model allowing to study the influence of defect depth (full thickness defects and

chondral defects) on tissue repair. In a pilot study, the relation between oxygen tension (normoxia 20 %

vs. physiological hypoxia 2 %), defect depth and treatment strategy was studied. For the treatment

strategies either chondrocytes only (CHON) or a mixture of 80 % MSC and 20 % CHON (MIX) were

embedded in a collagen type I hydrogel for implantation into cartilage defects in the *ex vivo* osteochondral

model without further supplementation of growth factors to elusively study the effect of the culture

conditions on cartilage repair.

126 2 Methods

127 2.1 Osteochondral cylinders: Isolation, defect creation and *ex vivo* culture

Osteochondral cylinders (diameter: 8 mm, height: 5 mm) were isolated from medial femoral condyles of

6-8-month-old domestic pigs as previously described [20]. Creation of defects in osteochondral cylinders

was carried out either manually or automatically, depending on the intended defect depth: Full thickness

defects were manually created with a biopsy punch (diameter 4 mm; Kai Medical, BPP-40F). Chondral

defects of 1 mm depth were created using custom-built ARTcut® (Figure 1A), a software-controlled device

for automatic wound placement in tissues developed by Fraunhofer ISC and IGB, Wuerzburg [23].

Osteochondral cylinders were fixed in support plate (Figure 1C) at specific x- and y-positions, and onset of

drilling 1 mm chondral defects (z-axis) was determined with light beam (Figure 1B). For more details on

the standardized defect creation process using ARTcut®, see chapter 1.2 in the supplementary

information.

After creation of full thickness or 1 mm chondral defects, the cartilage defects were filled with cell

embedded collagen type I hydrogels (Figure 2). For subsequent *ex vivo* culture, osteochondral explants

were transferred into custom-made culture platform [20] and cultured for 28 days with tissue specific

media (Table 1) changed every 3-4 days. The *ex vivo* culture was carried out in a humidified atmosphere

at 37°C and 5 % CO_2 (BBD 6220 CO_2 Incubator, Thermo Scientific™) either under normoxic (20 %) or

physiological hypoxic (2 %) conditions.

2.2   Chondrocyte and mesenchymal stromal cells (MSC): isolation and *in vitro* expansion

Porcine chondrocytes were isolated from lateral condyles of 6–8-month-old domestic pigs by enzymatic

digestion as previously described [20]. Chondrocytes were used at passage 0 – directly after isolation - for

embedding in collagen type I hydrogel.

Porcine MSC were isolated from bone marrow aspirate (iliac crest) under the approval (reference number:

55.2 2532-2-256) of the District Government of Lower Franconia and the local animal welfare committee

and performed according to the German Animal Welfare Act and the EU Directive 2010/63/EU according

to the procedure described in chapter 1.2 in the supplementary I.

2.3 Cell embedding in collagen type I hydrogel

In a pilot study, cartilage defects were treated with collagen type I hydrogel (see chapter 1.3 in the supplementary for details). For implantation, two different cell laden constructs, as illustrated in Figure 2, were prepared and each cultured under normoxic and physiological hypoxic conditions. 1) CHON were prepared by embedding of chondrocytes in collagen type I hydrogel without expansion at a final density of 20 million cells/mL hydrogel volume; 2) MIX were prepared by mixing 80 % MSC with 20 % chondrocytes at same cell density. CHON and MIX implants were filled into the cartilage defect for gelation at 37°C.

2.4 Live-dead viability staining

To visualize a possible effect of defect creation process on cell viability of tissue samples, live-dead viability staining was performed (live-dead staining kit for mammalian cells, L3224, Invitrogen). Osteochondral explants were incubated with 4 µM Calcein AM and 2 µM ethidium homodimer-1 in DMEM HG and visualized with fluorescence microscope (494 nm/517 nm and 517 nm/617 nm wavelength) Keyence BZ-9000 (Biorevo). Living cells are stained by Calcein AM in green, dead cells are stained by ethidium homodimer-1 in red.

2.5 Histological evaluation

Osteochondral explants, harvested on day 0 and day 28, were washed with phosphate buffered saline, fixed for 24 h with 4 % formalin (Roti Histofix, Carl Roth P087.3) and processed with plastic embedding (Technovit T9100, Heraeus Kulzer, 66006735) as detailed described in the supplementary.

For immune-histological stainings, antigens were enzymatically retrieved depending on antibody (Table 2). Before incubation with primary antibody over night at 4°C, slides were blocked with 3 % (v/v) H_2O_2 (Carl Roth, 8070) and incubated with 5 % (w/v) bovine serum albumin. All stainings were visualized with Horseradish-peroxidase kit (DCS Innovative, Dako, K3468) following manufacturer`s instructions. Cell nuclei were counterstained with Mayer's hematoxylin (Morphisto, 11895), rehydrated and mounted with

Entellan (Merck, 1079610500). All washing steps between blocking or chemical incubation were

performed with PBS supplemented with 0.5 % Tween-20 (VWR, 8.22184).

2.6   Histological scoring

Histological scoring for the evaluation of the immune histological stainings (collagen II, X and aggrecan)

was performed. For in vitro tissue engineered samples the Bern score has been introduced, while cartilage

tissue repair in clinical studies can be evaluated based on the international cartilage repair society (ICRS)

score II [27]. Further, Chang et al. introduced a modified International cartilage repair society score by

adding two new criteria, namely the visual assessment of collagen type I and II staining [28]. The inclusion

of extracellular markers is an important step to better evaluate the composition of the repair matrix.

Therefore, in this pilot study, the scoring category A (Bern-score), originally evaluating the uniformity and

darkness of Safranin O-fast green stain, was translated to immune histological stainings of collagen type I,

II, X and aggrecan) [29]. Stained sections (n=2 biological replicates) were evaluated blinded by three

independent operators according to the criteria summarized in Table 3. Maximum possible score for the

here reported study was 12 points.

2.7   Statistics

Statistical analysis was performed with GraphPad Prism 6.07. Scoring values obtained from two

independent experiments, each evaluated by three examinators were used for statistical analysis,

resulting in $n = 6$ for each experimental group (Figure 2). Two groups differing in only one culture

parameter (oxygen tension, defect depth and applied cell type) were directly compared. Following tests

for outlier and Gaussian distribution, the data was either compared using an unpaired t-test (normally

distributed values) or a Mann-Whitney test (not normally distributed values). A p-value <0.05 was

considered statistically significant.

3 Results

3.1 Chondral defect creation with ARTcut**®**

ARTcut® (Figure 1) represents an automated device for reproducible creation of chondral defects (1 mm in depth; Figure 3A). Main advantage of using the ARTcut® is the ability to create defined chondral defects with a flat bottom to allow a tide and uniform contact with any solid pre-formed cylindrical implant, for example bioprinted constructs with flat bottom.

The laser beam to detect the surface of every single osteochondral explant is the unique feature of the ARTcut® to define the onset of the drilling, thus all defects result with the same depth. Chondral cartilage defect induced with ARTcut® appeared as a well-defined rectangular defect boundary in cross section (diameter: 4 mm, height: 1 mm) with flat bottom (Figure 3B). In contrast, the manually created defect (using biopsy punch) resulted in a half-round shape and required removing of residual cartilage tissue with a second tool (*e.g.,* a sharp spoon) shown in Figure 3C. Following, the accuracy of chondral defect geometry was much lower in manual procedure - using biopsy punch - compared to software – using automated ARTcut®.

The automated defect creation took less than 5 sec and thus heating of tissue due to friction and rotation during drilling was very unlikely. Microscopic images of live-dead stainings of osteochondral explants with 1 mm cartilage defects (diameter: 4 mm) confirmed no indications of cell death after tissue defect creation, neither with ARTcut® nor with biopsy punch (Figure 3D-E).

For the creation of full thickness cartilage defects, the ARTcut® was not required, since the bottom of the defect was given by the cartilage-bone interface. This two-tissue interface was easily separated using biopsy punch by pulling out the upper cartilage part.

3.2 Evaluation of cartilage treatments in the *ex vivo* model

Macroscopic images of all treatment and culture groups of the osteochondral *ex vivo* model after 28 days of culture are illustrated in Figure 4 A and B. The results of the pilot study demonstrated the feasibility and

reproducibility of the *ex vivo* cartilage test system as a platform to compare cartilage repair strategies.

Results of the adapted Bern scoring are shown in Figure 4 and Table 4 with scoring results of the co-culture

(MIX) being close to the CHON treatment in chondral and full thickness defects. Representative sections

of the scored samples are illustrated in Figure 5 and 6.

Under physiological hypoxic conditions, scoring results of defect repair after co-culture treatment (MIX)

was higher (p = 0.0218) in chondral defect than in full thickness defects. Scoring values of CHON treatment

in chondral and full thickness experimental groups did not differ much (n.s. p = 0.1828). At physiological

hypoxic conditions the scoring value was higher for CHON treatment compared to MIX treatment in full

thickness defects (p = 0.0384).

Under normoxic conditions, full thickness defects reached higher scoring values compared to the chondral

defects.

Comparing the outcome of same defect depth and treatment group (cell type), but cultured under

physiological hypoxic or normoxic conditions, the oxygen tension plays only a significant role in chondral

defects using MIX (p = 0.0451, higher scoring values under physiological hypoxic conditions) and in full

thickness defects treated with CHON (p = 0.0316, lower scoring values under physiological hypoxic

conditions).

Overall, treatment of the two defect depths with MIX showed slightly reduced scoring values compared

to the CHON treatment group. Only exception is MIX chondral (physiological hypoxia) with a slightly higher

value as the corresponding one for CHON. The results of the MIX treatment groups also tend to spread

more than the CHON treatment group independently of defect depth and oxygen tension.

4   Discussion

The choice of optimal treatment approach for cartilage repair is dependent on the size and depth of the

cartilage defect and thus the severity of pathological condition.

To mimic these different pathological defect scenarios, the implementation of the ARTcut® allowed to

overcome the low reproducibility of manual induced chondral defects. So far, there is no other automated

defect creation device published that has comparable features, namely possibility to select different

drilling parameters within one run and the application of one device for a wide range of soft and hard

tissues or tissue equivalents, as it is demonstrated with the ARTcut®. The induction of defects in

osteochondral explants allows to compare different cell-based treatment approaches by implantation of

cell loaded materials into the defects or to study the effect of defect depth on repair in an *ex vivo* model.

The software-controlled drilling parameters allow for standardized and reproducible defect creation in

other soft and hard tissues including bone tissue as well. *In vitro* as well as *ex vivo* studies require sample

procession at high throughput rates under a negative microbial environment. ARTcut® device addresses

these demands, resulting in higher methodological accuracy and reproducibility, compared to manual

procedure using biopsy punch.

Current treatments of small or large cartilage defects mainly aim on pain and symptom relief rather than

on functional tissue repair [30]. From a scientific point of view, there is way for improvement of cartilage

defects, especially on long term results. Material-assisted as well as cell-based therapies, including (M-)

ACI, AMIC and microfracture, tend to result in formation of mechanical inferior fibrocartilage in long term

follow up [30-32]. Moreover, the treatment of large defects with autologous cells are limited due to the

high number of chondrocytes needed for the current techniques. Therefore, in this laboratory-controlled

study, an *ex vivo* osteochondral defect model was modified to screen cell-based approaches to treat

chondral and full thickness cartilage defects. Different cell types, namely chondrocytes and MSCs, were

embedded in collagen type I hydrogel (CHON or MIX) and implanted in trauma induced chondral and full

thickness defects. A tendency of a stimulative and beneficial effect of MIX treatment (20 % CHON and 80 %

MSCs) was observed regarding cartilaginous matrix formation with the main advantage to reduce the

overall number of CHON used for traditional ACI treatment. It has been demonstrated in previous studies

that the cells in cartilage and subchondral bone remained metabolic active up to 56 days of *ex vivo* culture

[20].

Promising results regarding cartilage matrix production were obtained in this pilot study comparing CHON

and MIX in chondral and full thickness defects. The most pronounced increase in proteoglycan content

normalized to DNA amount resulted for the MIX treatment in both conditions, chondral and full thickness

defect, cultured under physiological hypoxia (Supplementary Figure S1). The MIX treatment reduces the

amount of autologous cartilage tissue needed for chondrocyte isolation, since chondrocytes represented

only 20 % of the total cell number in our model. Several *in vitro* studies and *in silico* models have shown

that chondrocytes in co-culture with MSCs increase the chondrogenic differentiation potential [18, 19, 33,

34]. Scalzone *et al.* reported of an enhanced chondrocyte activity measured by proteoglycan and collagen

type II production after co-culture with MSC compared to chondrocyte monoculture both embedded in

chitosan based hydrogels [19].

One advantage of the co-culture approach – when clinically applied – would be the one-step procedure,

addressing the surgical intervention of the knee joint, for the treatment of the cartilage defect: The

reduced number of chondrocytes in the MIX treatment can be isolated intra-operatively without

expansion and thus reduced the risk of chondrocyte dedifferentiation [35]. Further, the required MSCs,

e.g., can be harvested independently form the knee surgery from bone marrow aspirate in a minimal

invasive procedure with subsequent expansion prior to the cartilage repair procedure to ensure a

sufficient number of MSCs.

Considering the absence of TGF-β in the culture media, which is known to drive MSC chondrogenesis and

matrix production [36], the here reported cartilage repair solely derives from the stimulating effect of the

surrounding osteochondral tissue, cell-cell signaling of the cell laden implants and the culture conditions.

Therefore, the overall matrix deposition is rather low due to missing stimuli of exogenous growth factors

in the here described model. Results of a similar *ex vivo* model, based on horse and bovine osteochondral

explants, suggested that the presence of the osteochondral tissue in the explants increase cartilage-like

matrix deposition compared to free swelling conditions of cell seeded material only [21, 22].

Beside the cell source, the influence of oxygen tension (normoxia 20 % O_2, physiological hypoxia 2 % O_2)

was studied during the *ex vivo* culture on chondrogenic matrix deposition, resulting in different outcomes

between the treatment groups in chondral and full thickness defects. Based on the histological scoring,

the results of the pilot study showed that the oxygen tension seems to only plays a role in chondral defects

- marked by elevated matrix production. In contrast, full thickness defects showed different response in

cartilage-like matrix deposition related to the treatment approach: While higher scoring values were

reached with CHON treatment under normoxic conditions, the pMIX treatment did not show differences

comparing physiological hypoxic and normoxic conditions. Physiological hypoxic conditions resulted in an

increase in cartilaginous matrix deposition in chondral defects compared to full thickness defects

independently of the used cell type (CHON, MIX). Once the defect reaches the cartilage-bone interface,

the cells in the implant receive stimuli from the subchondral bone that seem to counteract with the

hypoxic induced stimuli in the here presented model. One explanation of the differential cell response to

oxygen tension in chondral and full thickness defects may originate from the variation in oxygen tension

present in the human body. Chondrocytes are exposed to lower oxygen tension in avascular cartilage (2-

5 %) [37], synovial fluid and the synovial capsule (6.5-9.0 %) [25, 38], with an increasing oxygen level in the

bone marrow of subchondral bone (>7 %) [26, 39] and a maximum of 12 % in arterial blood [40]. Once the

cartilage defect progresses to the exposure of the subchondral bone in patients, chondrocytes within the

defect are exposed to higher levels of oxygen present in the bone supplied by vascular invasion [41]. It has

been shown that low oxygen inhibits the degradation of hypoxia induced factors (HIF) [42]. HIFs are

described to be essential for maintaining CHON homeostasis and extracellular matrix synthesis and

activate the transcription of genes [43].

While the here presented results of the pilot study are consistent, traceable and in line with literature,

there are still limitation in the current experimental setup: Due to limitations in availability of human

healthy osteochondral tissue – absence of any osteoarthritic phenotype – the authors decided to use

porcine tissue instead, as introduced by Schwab *et al.* [20]. The performance of an extensive comparative

ex vivo study comprising three parameters – namely 1) defect depth (chondral vs. full thickness), 2)

different cell types for defect filling and treatment (100 % chondrocytes vs. 20 % chondrocytes and 80 %

MSCs), and 3) oxygen tension (physiological hypoxia vs. normoxia) requires tissue material in the required

quantity and of similar quality. Due to the well-known donor issues associated with donor variability of

human tissue and the already mentioned limited availability of healthy human osteochondral tissue, this

could not be achieved for human tissue [44, 45]. In contrast, the use of porcine tissue isolated from one

pig population (similar age; grown up under equal conditions) minimizes the donor variation and

represents a tissue source for healthy and non-degenerated tissues with higher reproducibility. To proof

the findings and the trend of the here presented study and to show clear differences between treatment

groups, this experimental set-up needs to be repeated with more biological replicates.

The results obtained in this pilot study indicate that the surrounding tissue of the osteochondral explants

plays a crucial role in the defect repair, addressing chondral or full thickness defects, and that the oxygen

tension additionally stimulates the outcome in a defect depth dependent way. However, the exact

mechanism controlling success or failure of a treatment approach are not fully understood and require

further research. Despite the advantage of the here reported model, the mimicry of the complex 3D

environment present in the knee joint with the additional ability to reproducibly create (trauma induced)

defects, this model has its potential in pre-screening several biomaterials or treatment approaches, but

best candidates still require further pre-clinical and clinical testing to confirm obtained results.

To conclude, in this *ex vivo* study two cell-based approaches (CHON vs. MIX embedded in collagen type I

hydrogel) for the treatment of chondral and full thickness defects, cultured under normoxic and

physiological hypoxic conditions, were compared on their cartilage repair potential. Independently of

defect depth, the scoring results of the co-culture of chondrocytes and MSC (MIX) are close to those of

CHON treatment. Hence, usage of MIX could be a promising approach to reduce the number of

chondrocytes to one fifth and thus the amount of tissue for harvest and accompanied side morbidities for

the patient. Further, in this experimental set up the oxygen tension showed an influence in a defect depth

dependent way that has not been reported in literature before. The here presented advanced

osteochondral cartilage defect model provides a promising platform to deeper understand the underlying

mechanism.

347 Acknowledgements

The authors thank Andreas Diegeler (Fraunhofer ISC, Bronnbach) for technical support of the ARTcut®.

Thanks also to Sebastian Naczenski for establishing the protocol for wounding of osteochondral explants

using ARTcut® device. Finally, we would like to thank Heike Walles for her support. We acknowledge the

authors of the thesis (Andrea Schwab and Alexa Buss) for the permission to use the data in this manuscript.

Parts of the illustrations were produced in the light of a doctoral thesis (figure 1 and 3, with approval from

Schwab 2017, German copyright law) [46] and a doctoral thesis (figure 5 and 6, with approval from Buss

2021) [47].

356 Author contribution

357 Conception and design of the study: Franziska Ehlicke and Andrea Schwab; Acquisition of data: Alexa Buss

358 and Andrea Schwab; Analysis and interpretation of data: Andrea Schwab, Alexa Buss, Franziska Ehlicke and

359 Oliver Pullig; Manuscript drafting: Andrea Schwab and Alexa Buss; Revision of the manuscript: Franziska

360 Ehlicke and Oliver Pullig; Final approval: Franziska Ehlicke, Andrea Schwab, Alexa Buss and Oliver Pullig;

361 Funding: Franziska Ehlicke and Alexa Buss.

363 Role of funding Sources

364 This work was financially supported by the European Union Seventh Framework Programme (FP7/2007-

365 2013) under grant agreement no 309962 (HydroZONES).A.Buss was supported by a medical scholarship of

366 the German Excellence Initiative to the Graduate School of Life Sciences, University of Würzburg. This

367 publication was supported by the Open Access Publication Fund of the University of Wuerzburg.

Conflict of interest statement

369 The authors declare no conflict of interest.

Ethics approval and consent to participate

371 Animal experiment was approved (reference number: 55.2 2532-2-256) by the District Government of

372 Lower Franconia and the local animal welfare committee and performed according to the German Animal

373 Welfare Act and the EU Directive 2010/63/EU. Following heparinization of the pig, porcine MSCs were

374 obtained by bone marrow aspiration.

References

[1] W. Widuchowski, J. Widuchowski, T. Trzaska, Articular cartilage defects: study of 25,124 knee arthroscopies, Knee 14(3) (2007) 177-82. https://doi.org/10.1016/j.knee.2007.02.001.

[2] A.C. Gelber, M.C. Hochberg, L.A. Mead, N.Y. Wang, F.M. Wigley, M.J. Klag, Joint injury in young adults and risk for subsequent knee and hip osteoarthritis, Ann Intern Med 133(5) (2000) 321-8. https://doi.org/10.7326/0003-4819-133-5-200009050-00007.

[3] A.R. Martín, J.M. Patel, H.M. Zlotnick, J.L. Carey, R.L. Mauck, Emerging therapies for cartilage regeneration in currently excluded 'red knee' populations, npj Regenerative Medicine 4(1) (2019) 12. https://doi.org/10.1038/s41536-019-0074-7.

[4] J.D. Wylie, M.K. Hartley, A.L. Kapron, S.K. Aoki, T.G. Maak, Failures and Reoperations After Matrix-Assisted Cartilage Repair of the Knee: A Systematic Review, Arthroscopy 32(2) (2016) 386-92. https://doi.org/10.1016/j.arthro.2015.07.025.

[5] A.J. Sophia Fox, A. Bedi, S.A. Rodeo, The basic science of articular cartilage: structure, composition, and function, Sports Health 1(6) (2009) 461-8. https://doi.org/10.1177/1941738109350438.

[6] E.A. Makris, A.H. Gomoll, K.N. Malizos, J.C. Hu, K.A. Athanasiou, Repair and tissue engineering techniques for articular cartilage, Nat Rev Rheumatol 11(1) (2015) 21-34. https://doi.org/10.1038/nrrheum.2014.157.

[7] B.J. Cole, C. Pascual-Garrido, R.C. Grumet, Surgical management of articular cartilage defects in the knee, J Bone Joint Surg Am 91(7) (2009) 1778-90.

[8] A.K. Dewan, M.A. Gibson, J.H. Elisseeff, M.E. Trice, Evolution of autologous chondrocyte repair and comparison to other cartilage repair techniques, Biomed Res Int 2014 (2014) 272481. https://doi.org/10.1155/2014/272481.

[9] P. Niemeyer, T. Schubert, M. Grebe, A. Hoburg, Matrix-Associated Chondrocyte Implantation Is Associated With Fewer Reoperations Than Microfracture: Results of a Population-Representative, Matched-Pair Claims Data Analysis for Cartilage Defects of the Knee, Orthop J Sports Med 7(10) (2019) 2325967119877847. https://doi.org/10.1177/2325967119877847.

[10] M. Brittberg, A. Lindahl, A. Nilsson, C. Ohlsson, O. Isaksson, L. Peterson, Treatment of deep cartilage defects in the knee with autologous chondrocyte transplantation, N Engl J Med 331(14) (1994) 889-95. https://doi.org/10.1056/NEJM199410063311401.

[11] M. Mumme, A. Barbero, S. Miot, A. Wixmerten, S. Feliciano, F. Wolf, A.M. Asnaghi, D. Baumhoer, O. Bieri, M. Kretzschmar, G. Pagenstert, M. Haug, D.J. Schaefer, I. Martin, M. Jakob, Nasal chondrocyte-based engineered autologous cartilage tissue for repair of articular cartilage defects: an observational first-in-human trial, Lancet (London, England) 388(10055) (2016) 1985-1994. https://doi.org/10.1016/S0140-6736(16)31658-0.

[12] R.L. Davies, N.J. Kuiper, Regenerative Medicine: A Review of the Evolution of Autologous Chondrocyte Implantation (ACI) Therapy, Bioengineering (Basel) 6(1) (2019) https://doi.org/10.3390/bioengineering6010022.

[13] J.D. Lamplot, K.A. Schafer, M.J. Matava, Treatment of Failed Articular Cartilage Reconstructive Procedures of the Knee: A Systematic Review, Orthop J Sports Med 6(3) (2018) 2325967118761871. https://doi.org/10.1177/2325967118761871.

[14] G. Merkely, T. Ogura, J. Ackermann, A. Barbieri Mestriner, A.H. Gomoll, Clinical Outcomes
after Revision of Autologous Chondrocyte Implantation to Osteochondral Allograft
Transplantation for Large Chondral Defects: A Comparative Matched-Group Analysis, Cartilage
(2019) 1947603519833136. https://doi.org/10.1177/1947603519833136.
[15] K. Mithoefer, T. McAdams, R.J. Williams, P.C. Kreuz, B.R. Mandelbaum, Clinical efficacy of
the microfracture technique for articular cartilage repair in the knee: an evidence-based
systematic analysis, Am J Sports Med 37(10) (2009) 2053-63.
https://doi.org/10.1177/0363546508328414.
[16] P.C. Kreuz, M.R. Steinwachs, C. Erggelet, S.J. Krause, G. Konrad, M. Uhl, N. Sudkamp, Results
after microfracture of full-thickness chondral defects in different compartments in the knee,
Osteoarthritis Cartilage 14(11) (2006) 1119-25. https://doi.org/10.1016/j.joca.2006.05.003.
[17] J.L. Carey, Fibrocartilage Following Microfracture Is Not as Robust as Native Articular
Cartilage: Commentary on an article by Aaron J. Krych, MD, et al.: "Activity Levels Are Higher
After Osteochondral Autograft Transfer Mosaicplasty Than After Microfracture for Articular
Cartilage Defects of the Knee. A Retrospective Comparative Study", JBJS 94(11) (2012).
[18] L. Bian, D.Y. Zhai, R.L. Mauck, J.A. Burdick, Coculture of human mesenchymal stem cells and
articular chondrocytes reduces hypertrophy and enhances functional properties of engineered
cartilage, Tissue engineering. Part A 17(7-8) (2011) 1137-1145.
https://doi.org/10.1089/ten.TEA.2010.0531.
[19] A. Scalzone, A.M. Ferreira, C. Tonda-Turo, G. Ciardelli, K. Dalgarno, P. Gentile, The interplay
between chondrocyte spheroids and mesenchymal stem cells boosts cartilage regeneration
within a 3D natural-based hydrogel, Sci Rep 9(1) (2019) 14630. https://doi.org/10.1038/s41598-
019-51070-7.
[20] A. Schwab, A. Meeuwsen, F. Ehlicke, J. Hansmann, L. Mulder, A. Smits, H. Walles, L. Kock, Ex
vivo culture platform for assessment of cartilage repair treatment strategies, ALTEX 34(2) (2017)
267-277. https://doi.org/10.14573/altex.1607111.
[21] V.H.M. Mouser, N.M.M. Dautzenberg, R. Levato, M.H.P. van Rijen, W.J.A. Dhert, J. Malda, D.
Gawlitta, Ex vivo model unravelling cell distribution effect in hydrogels for cartilage repair,
ALTEX 35(1) (2018) 65-76. https://doi.org/10.14573/altex.1704171.
[22] M.L. de Vries-van Melle, R. Narcisi, N. Kops, W.J. Koevoet, P.K. Bos, J.M. Murphy, J.A.
Verhaar, P.M. van der Kraan, G.J. van Osch, Chondrogenesis of mesenchymal stem cells in an
osteochondral environment is mediated by the subchondral bone, Tissue Eng Part A 20(1-2)
(2014) 23-33. https://doi.org/10.1089/ten.TEA.2013.0080.
[23] A. Rossi, A. Appelt-Menzel, S. Kurdyn, H. Walles, F. Groeber, Generation of a three-
dimensional full thickness skin equivalent and automated wounding, J Vis Exp (96) (2015)
https://doi.org/10.3791/52576.
[24] G. Pattappa, R. Schewior, I. Hofmeister, J. Seja, J. Zellner, B. Johnstone, D. Docheva, P.
Angele, Physioxia Has a Beneficial Effect on Cartilage Matrix Production in Interleukin-1 Beta-
Inhibited Mesenchymal Stem Cell Chondrogenesis, Cells 8(8) (2019)
https://doi.org/10.3390/cells8080936.
[25] G. Pattappa, B. Johnstone, J. Zellner, D. Docheva, P. Angele, The Importance of Physioxia in
Mesenchymal Stem Cell Chondrogenesis and the Mechanisms Controlling Its Response, Int J Mol
Sci 20(3) (2019) https://doi.org/10.3390/ijms20030484.

[26] D.E. Anderson, B.D. Markway, D. Bond, H.E. McCarthy, B. Johnstone, Responses to altered
oxygen tension are distinct between human stem cells of high and low chondrogenic capacity,
Stem Cell Res Ther 7(1) (2016) 154. https://doi.org/10.1186/s13287-016-0419-8.
[27] M. Rutgers, M.J.P. van Pelt, W.J.A. Dhert, L.B. Creemers, D.B.F. Saris, Evaluation of
histological scoring systems for tissue-engineered, repaired and osteoarthritic cartilage,
Osteoarthritis and Cartilage 18(1) (2010) 12-23.
https://doi.org/https://doi.org/10.1016/j.joca.2009.08.009.
[28] F. Chang, T. Ishii, T. Yanai, H. Mishima, H. Akaogi, T. Ogawa, N. Ochiai, Repair of large full-
thickness articular cartilage defects by transplantation of autologous uncultured bone-marrow-
derived mononuclear cells, Journal of Orthopaedic Research 26(1) (2008) 18-26.
https://doi.org/https://doi.org/10.1002/jor.20470.
[29] S.P. Grogan, A. Barbero, V. Winkelmann, F. Rieser, J.S. Fitzsimmons, S. O'Driscoll, I. Martin,
P. Mainil-Varlet, Visual Histological Grading System for the Evaluation of in Vitro–Generated
Neocartilage, Tissue Engineering 12(8) (2006) 2141-2149.
https://doi.org/10.1089/ten.2006.12.2141.
[30] U. Horas, D. Pelinkovic, G. Herr, T. Aigner, R. Schnettler, Autologous chondrocyte
implantation and osteochondral cylinder transplantation in cartilage repair of the knee joint. A
prospective, comparative trial, J Bone Joint Surg Am 85(2) (2003) 185-92.
https://doi.org/10.2106/00004623-200302000-00001.
[31] J.R. Steadman, K.K. Briggs, J.J. Rodrigo, M.S. Kocher, T.J. Gill, W.G. Rodkey, Outcomes of
microfracture for traumatic chondral defects of the knee: average 11-year follow-up,
Arthroscopy 19(5) (2003) 477-84. https://doi.org/10.1053/jars.2003.50112.
[32] J.L. Carey, Fibrocartilage following microfracture is not as robust as native articular
cartilage: commentary on an article by Aaron J. Krych, MD, et al.: "Activity levels are higher after
osteochondral autograft transfer mosaicplasty than after microfracture for articular cartilage
defects of the knee. A retrospective comparative study", J Bone Joint Surg Am 94(11) (2012)
e80. https://doi.org/10.2106/JBJS.L.00319.
[33] M.J. Chen, J.P. Whiteley, C.P. Please, A. Schwab, F. Ehlicke, S.L. Waters, H.M. Byrne,
Inducing chondrogenesis in MSC/chondrocyte co-cultures using exogenous TGF-β: a
mathematical model, Journal of Theoretical Biology 439 (2018) 1-13.
https://doi.org/https://doi.org/10.1016/j.jtbi.2017.11.024.
[34] T.W. Kim, M.C. Lee, H.C. Bae, H.-S. Han, Direct Coculture of Human Chondrocytes and
Synovium-Derived Stem Cells Enhances In Vitro Chondrogenesis, Cell J 20(1) (2018) 53-60.
https://doi.org/10.22074/cellj.2018.5025.
[35] K. von der Mark, V. Gauss, H. von der Mark, P. Muller, Relationship between cell shape and
type of collagen synthesised as chondrocytes lose their cartilage phenotype in culture, Nature
267(5611) (1977) 531-2. https://doi.org/10.1038/267531a0.
[36] B. Johnstone, T.M. Hering, A.I. Caplan, V.M. Goldberg, J.U. Yoo, In vitro chondrogenesis of
bone marrow-derived mesenchymal progenitor cells, Exp Cell Res 238(1) (1998) 265-72.
https://doi.org/10.1006/excr.1997.3858.
[37] S. Zhou, Z. Cui, J.P.G. Urban, Factors influencing the oxygen concentration gradient from
the synovial surface of articular cartilage to the cartilage–bone interface: A modeling study,
Arthritis & Rheumatism 50(12) (2004) 3915-3924. https://doi.org/10.1002/art.20675.

[38] K. Lund-Olesen, Oxygen tension in synovial fluids, Arthritis Rheum 13(6) (1970) 769-76.
https://doi.org/10.1002/art.1780130606.
[39] A. Carreau, B. El Hafny-Rahbi, A. Matejuk, C. Grillon, C. Kieda, Why is the partial oxygen
pressure of human tissues a crucial parameter? Small molecules and hypoxia, J Cell Mol Med
15(6) (2011) 1239-53. https://doi.org/10.1111/j.1582-4934.2011.01258.x.
[40] M. Marenzana, T.R. Arnett, The Key Role of the Blood Supply to Bone, Bone Research 1(1)
(2013) 203-215. https://doi.org/10.4248/BR201303001.
[41] J. Filipowska, K.A. Tomaszewski, L. Niedzwiedzki, J.A. Walocha, T. Niedzwiedzki, The role of
vasculature in bone development, regeneration and proper systemic functioning, Angiogenesis
20(3) (2017) 291-302. https://doi.org/10.1007/s10456-017-9541-1.
[42] A. Kremer, M. Wussmann, M. Herrmann, M. Raghunath, H. Walles, Ciclopirox olamine
promotes the angiogenic response of endothelial cells and mesenchymal stem cells, Clin
Hemorheol Microcirc 73(2) (2019) 317-328. https://doi.org/10.3233/CH-190559.
[43] J. Fernandez-Torres, Y. Zamudio-Cuevas, G.A. Martinez-Nava, A.G. Lopez-Reyes, Hypoxia-
Inducible Factors (HIFs) in the articular cartilage: a systematic review, Eur Rev Med Pharmacol
Sci 21(12) (2017) 2800-2810.
[44] H.X. Chang, L. Yang, Z. Li, G. Chen, G. Dai, Age-related biological characterization of
mesenchymal progenitor cells in human articular cartilage, Orthopedics 34(8) (2011) e382-8.
https://doi.org/10.3928/01477447-20110627-06.
[45] M. Kim, I.E. Erickson, A.H. Huang, S.T. Garrity, R.L. Mauck, D.R. Steinberg, Donor Variation
and Optimization of Human Mesenchymal Stem Cell Chondrogenesis in Hyaluronic Acid, Tissue
Eng Part A 24(21-22) (2018) 1693-1703. https://doi.org/10.1089/ten.TEA.2017.0520.
[46] A. Schwab, Development of an osteochondral cartilage defect model, Julius-Maximilians
University, Wuerzburg 2017.
[47] A. Buss, Evaluation of cartilage regeneration strategies in an osteochondral ex vivo cartilage
defect model, Julius-Maximilians University, Wuerzburg 2021.

Figure Legends

Figure 1: Artificial tissue cutter (ARTcut®): A) View of the computerized numerical controlled device

equipped with B) optical barrier (white arrow indicates the laser light beam) and a C) supporting plate to

fix the samples and place on the bottom of the machine (marked with black star).

Figure 2: Schematic overview of all experimental groups: Full thickness and chondral defects of

osteochondral explants were treated with collagen type I hydrogel (grey) either containing chondrocytes

(CHON, blue) or MIX (80 % MSC [orange] and 20 % chondrocytes), implanted in chondral or full thickness

defects and cultured *ex vivo* under normoxic (20 % O_2, blue frame) or physiological hypoxic conditions (2 %

O_2, red frame).

547 Figure 3: Osteochondral explant cross section with chondral defect. A) Schematic illustration of A) 4 mm

548 chondral defect. B-E) Comparison of chondral defect created automated with ARTcut® (B) and manually

549 with biopsy punch (C) (scale bar 500 µm). D-E) Live-dead staining of explant cross section highlighting

550 accurate borders of defect induced with ARTcut® (D) and more irregular shaped geometry of manually

551 induced defect (E) (scale bar 200 µm).

Figure 4: Evaluation of defect repair after 28 days *ex vivo* culture. Macroscopic images of full thickness and chondral defects (diameter 4 mm) treated with A) porcine chondrocytes (CHON) or B) co-culture of porcine MSC and CHON (MIX). Red frame indicates culture under physiological hypoxic, blue frame under normoxic oxygen tension. Macroscopically, there are no obvious differences comparing full thickness and chondral defects. C) Scoring evaluation of cartilage repair strategies in the *ex vivo* model (mean ± standard deviation with single scoring results as dots resp. listed values for 1st and 2nd biological replicate). Unpaired t-test or Mann-Whitney test * $p < 0.05$, ** $p < 0.01$.

564 Figure 5: Chondral defects of the osteochondral defect model treated with either chondrocytes (CHON) or

565 MSC-chondrocyte co-culture (MIX) embedded in a collagen hydrogel. Histological stainings of cross

566 sections showing the defect area at day 0 and after 28 days of ex vivo culture under normoxic (20 % O_2) or

567 physiological hypoxic (2 % O_2 conditions. Scale bar 500 µm.

Figure 6: Full thickness cartilage defects of the osteochondral defect model treated with either

chondrocytes (CHON) or MSC-chondrocyte co-culture (MIX) embedded in a collagen hydrogel. Histological

stainings of cross sections showing the defect area at day 0 and after 28 days of ex vivo culture under

normoxic (20 % O_2) or physiological hypoxic (2 % O_2) conditions. Scale bar 500 μm.

574   Table 1: Composition of bone and cartilage media for *ex vivo* culture of osteochondral explants.

	Cartilage media	Bone media	Supplier
DMEM GlutaMAX (high glucose)	98 %	89 %	Gibco 61965
Antibiotics	10 U/mL penicillin, 10 µg/mL Streptomycin, 0.25 µg/mL Amphotericin (1 % (v/v))	10 U/mL penicillin, 10 µg/mL Streptomycin, 0.25 µg/mL Amphotericin (1 % (v/v))	Gibco, 15240
Sodium-pyruvate	1 mM	-	Gibco, 11360
L-ascorbic acid-2-phosphate	50 µg/mL	50 µg/mL	Wako chemicals, 013-19641
L-proline	40 µg/mL	-	Sigma-Aldrich, P5607
Dexamethasone	100 nM	100 nM	Sigma-Aldrich, D4902
fetal bovine serum (FBS)		10 % (v/v)	Gibco, 10270106
Insulin-transferrin-selenin	1 % (v/v)	-	ITS+-premix, Gibco, Life Technologies, 32430
β-glycerophosphate	-	10 mM	Sigma-Aldrich, G9422

576   Table 2: List of antibodies for immune-histological stainings.

Antibody	clone	Dilution and final antibody concentration	Antigen retrieval	Antibody supplier
Collagen type I	EPR7785	1:1000; 0.875 µg/mL	pepsin & hyaluronidase	Abcam, ab138492
Collagen type II	II-4C11	1:1000; 1 µg/mL	pepsin & hyaluronidase	MP Bio, 63171
Collagen type X	COL-10	1:4000; 0.25 µg/mL	pronase	Sigma-Aldrich, C7974
Aggrecan	EPR7785	1:2000; 0.5 µg/mL	pronase	Invitrogen, AHP0012

578 Table 3: Criteria for evaluating immune histological stainings with a maximum of 12 points (0-3 scores for each criteria assessing

579 the uniformity and darkness of aggrecan, collagen type I, II and X staining). Table adapted from Grogan et al. and Chang et al.

580 [28, 29].

Uniformity and darkness of aggrecan stain
0: No stain, 1: Weak staining of poorly formed matrix, 2: Moderately even staining, 3: Even dark stain
Uniformity and darkness of collagen type II stain
0: No stain, 1: Weak staining of poorly formed matrix, 2: Moderately even staining, 3: Even dark stain
Uniformity and darkness of collagen type I stain
3: No stain, 2: Weak staining of poorly formed matrix, 1: Moderately even staining, 0: Even dark stain
Uniformity and darkness of collagen type X stain
3: No stain, 2: Weak staining of poorly formed matrix, 1: Moderately even staining, 0: Even dark stain

582   Table 4: Scoring results to evaluate cartilage repair strategies in the ex vivo model. The table summarized the cumulative score

583   of the 3-independent scorer (mean ± standard deviation with single scoring results of both biological replicate) for the CHON

584   (100 % chondrocytes) and MIX (20 % chondrocytes, 80 % MSCs) treatment of the full thickness and chondral defects after 28

585   days culture. Values for samples cultured under physiological hypoxia (2 % O2) and normoxia (20 % O2) are displayed in

586   separate columns.

		CHON		MIX	
	Cumulative score (max 12)	Physiological hypoxia (2 % O_2)	Normoxia (20 % O_2)	Physiological hypoxia (2 % O_2)	Normoxia (20 % O_2)
full thickness defect	1st biological replicate	9; 8; 8	9; 9; 7	6; 7; 4	8; 8; 7
	2nd biological replicate	7; 6; 6	9; 11; 10	5; 7; 5	5; 5; 6
	Mean ± standard deviation	**7.33 ± 1.21**	**9.17 ± 1.33**	**5.67 ± 1.21**	**6.5 ± 1.38**
chondral defect	1st biological replicate	7; 8; 9	7; 7; 7	11; 9; 12	4; 2; 2
	2nd biological replicate	8; 9; 8	9; 8; 7	7; 7; 6	8; 8; 7
	Mean ± standard deviation	**8.17 ± 0.75**	**7.50 ± 0.84**	**8.67 ± 2.42**	**5.17 ± 2.86**