Pro chondrogenic effect of mesenchymal stromal cell-based treatment of chondral

defects under physioxia in a novel *ex vivo* organ model

Andrea Schwab [1], Alexa Buß [1], Oliver Pullig [1, 2], Franziska Ehlicke [1,$]

[1] University Hospital Wuerzburg, Department Tissue Engineering and Regenerative Medicine,

Roentgenring 11, 97070 Wuerzburg, Germany

[2] Fraunhofer Institute for Silicate Research ISC, Translational Center Regenerative Therapies,

Roentgenring 11, 97070 Wuerzburg, Germany

Andrea Schwab

• University Hospital Wuerzburg, Department Tissue Engineering and Regenerative Medicine

• Present address: AO Research Institute Davos, Clavadelerstrasse 8, 7270 Davos, Switzerland

E-Mail: andrea.schwab@aofoundation.org

Alexa Buß

• University Hospital Wuerzburg, Department Tissue Engineering and Regenerative Medicine,

Roentgenring 11, 97070 Wuerzburg, Germany

• Present address: Klinikum Würzburg Mitte gGmbH, Juliusspital, Juliuspromenade 19, 97070

Wuerzburg, Germany

E-Mail: alexa.buss@kwm-klinikum.de

Oliver Pullig

• University Hospital Wuerzburg, Department Tissue Engineering and Regenerative Medicine,

Roentgenring 11, 97070 Wuerzburg, Germany

• Fraunhofer Institute for Silicate Research ISC, Translational Center Regenerative Therapies,

Roentgenring 11, 97070 Wuerzburg, Germany

E-Mail: oliver.pullig@isc.fraunhofer.de

Franziska Ehlicke ([$]corresponding author)

• University Hospital Wuerzburg, Department Tissue Engineering and Regenerative Medicine,

Roentgenring 11, 97070 Wuerzburg, Germany

E-Mail: franziska.ehlicke@uni-wuerzburg.de

Phone: +49 931 3181900

Abstract

Objective

High failure rates of (trauma induced) knee injuries highlight the need to improve current treatment strategies aiming to decrease the number of secondary osteoarthritis developed by patients in later stage [1-3]. In this controlled laboratory study the stimulative effect of mesenchymal stromal cells (MSC) on chondrocyte (CHON) extracellular matrix production was investigated in an *ex vivo* cartilage defect model (chondral vs. full thickness defects) cultured under normoxic (20 % O_2) and physioxic (2 % O_2) conditions.

Design

Porcine CHON or co-culture of 20 % CHON and 80 % MSC (MIX) were embedded in collagen type I hydrogel, implanted into 4 mm diameter cartilage defects of osteochondral explants and cultured with tissue specific media without addition of TGF-β under normoxia and physioxia. Chondral defects were induced automatically, while full thickness defects were created with biopsy punch. After 28 days of culture, samples were histologically processed, and treatments outcome was evaluated using international cartilage repair society (ICRS)-II scoring.

Results

Under physioxic conditions, cartilage repair scoring results of the MIX treatment (chondral 8.67 ± 2.42, full thickness 5.67 ± 1.21) were close to those of CHON treatment (chondral 8.17 ± 0.75, full thickness

7.33 ± 1.21). Overall, scoring results were higher in physioxia compared to normoxia conditions in chondral

defects, but less or no prevalent for full thickness defects.

<u>Conclusion</u>

Co-culture of CHON with MSC represents a promising approach to stimulate chondrogenic repair and

tissue formation in our *ex vivo* model and reduces total amount of CHON needed for cell-based treatment.

<u>Keywords</u>

cartilage defect, *ex vivo* model, cartilage test system, chondrocytes, MSC, collagen type I

hydrogel

1   Introduction

Defects of the articulating surface are frequently occuring diseases in the field of orthopedics,

traumatology and sports medicine [4]. Around 14 % of patients who suffered from a trauma induced knee

injury during their middle ages, including fractures of the tibia, fibula, femur, or patella, develop

osteoarthritis in the same knee joint at later age (>65 years) [3]. High risk of failure of more than 20 % in

cartilage defect repair require to rethink and improve current cartilage treatment concepts with the aim

for long-term repair of cartilage defects [1, 2].

Due to the avascular and aneural character of adult articular cartilage, large defects do not heal

spontaneously [5]. Defects that are classified according to International Cartilage Repair Society (ICRS)

grade III (defect depth >50 % cartilage depth) and IV (defect reaches subchondral bone) need surgical

intervention to repair [6]. Gold standard for defects larger than 2-3 cm² is the autologous chondrocyte

implantation (ACI), resulting in similar outcome as microfracture [7, 8]. In a recent data analysis for

cartilage defects, matrix-associated chondrocyte implantation (M-ACI) showed a significant lower

reoperation rate than microfracture 2 years post-op [9]. M-ACI as well as ACI require two surgical

interventions; the first to take a cartilage tissue sample for subsequent chondrocyte isolation and the

second one for injection of isolated *in vitro* expanded autologous chondrocytes into defect site.

However, (M-)ACI has two main limitations: Due to the limited number of healthy and non-degenerated

chondrocytes, cells have to be expanded *in vitro* to achieve sufficient number of cells needed for the

implantation (ACI: 1 Mio/cm², M-ACI: 20 Mio/cm³) [10-12]. Further, chondrocyte-based cartilage

treatments, as well as microfracture techniques, result in formation of fibrocartilage, characterized by its

inferior mechanical properties and thus lacking functional restoration compared to healthy hyaline

cartilage [2, 13-17]. One approach to overcome the limitation of cell number for (M-)ACI treatment are

MSC-chondrocyte co-cultures, *e.g.*, 80 % MSC and 20 % chondrocytes, that reduces autologous

chondrocyte cell number at same total cell density. Co-cultures of MSCs and chondrocytes have shown to

increase cartilaginous matrix production and reduce hypertrophy, associated with MSCs during

chondrogenic differentiation [18, 19].

To bring cartilage treatments to the next level, cartilage repair needs to be studied and understood in

more detail, starting with basic research questions in a physiologically relevant environment.

An *ex vivo* cartilage defect test system based on osteochondral explants was established by Schwab *et al.*.

This test system represents a valuable *ex vivo* platform for biomaterial evaluation in critical size trauma-

induced cartilage defects in terms of biocompatibility, biomaterial tissue integration and cartilage repair

with higher throughput compared to *in vivo* models. Separated media compartments of the culture device

allow for controlled, tissue- and cell-specific nutrient supply during *ex vivo* culture of osteochondral

explants [20]. This model also allows direct comparison of different treatment approaches under

controlled conditions and has been shown to stimulate cartilage-like tissue formation of chondrocytes or

MSCs in osteochondral lesions [21, 22].

However, this defect model was limited to full thickness defects created with a biopsy punch. To date, the

creation of defects that do not fully penetrate the cartilage with the help of a biopsy punch or scalpel

suffer from low reproducibility and are dependent on the operator. Therefore, an automated device with

control over drilling depth to create defects with high reproducibility can overcome this limitation to study

chondral wound healing. Of note, the drilling should not harm surrounding cartilage tissue and cell

viability, neither by mechanical tissue disruption nor by friction induced heating. For the creation of

defects in hard tissues like cartilage or even bone, there is no device available for automated defect

creation that meets the above-mentioned requirements.

Following, in the present study the defect creation of the *ex vivo* defect model introduced by Schwab *et*

al. was modified by implementation of a semi-automated drilling device, originally developed to

mechanically induce standardized and reproducible wounds in full thickness skin equivalents [23]. Key

features of this artificial tissue cutter (ARTcut®) are the sensor controlled optical barrier in combination

with a moveable milling machine along x-, y- and z-axis. The light-barrier controlled drilling allows for

creation of defects with defined depths in a reproducible set-up. Creation of tissue defects with ARTcut®

can be performed under sterile conditions for subsequent *in vitro* or *ex vivo* culture with the possibility to

adjust defect geometry and control of defect creation process.

Another fact that has been neglected in the studies by Schwab *et al.* is the low oxygen tension present in

the articular capsule [24]. It has been shown in literature that low oxygen tension is an important stimulus

in chondrogenesis and increases the chondrogenic potential of articular cartilage progenitor cells and

MSCs [24-26].

Taken together, the present study aimed to investigate two cartilage treatment strategies and the

influence of defect depth (full thickness defects and chondral defects) on tissue repair. Additionally, the

effect of oxygen tension (normoxia 20 % vs. physioxia 2 %) was studied in both defect and treatment

approaches. For the treatment strategies either chondrocytes only (CHON) or a mixture of 80 % MSC and

20 % CHON (MIX) were embedded in a collagen type I hydrogel for implantation into cartilage defects in

the *ex vivo* osteochondral model without further supplementation of growth factors to elusively study the

effect of the culture conditions on cartilage repair.

2   Methods

2.1   Osteochondral cylinders: Isolation, defect creation and *ex vivo* culture

Osteochondral cylinders (diameter: 8 mm, height: 5 mm) were isolated from medial femoral condyles of 6-8-month-old domestic pigs as previously described [20]. Creation of defects in osteochondral cylinders was carried out either manually or automatically, depending on the intended defect depth: Full thickness defects were manually created with a biopsy punch (diameter 4 mm; Kai Medical, BPP-40F). Chondral defects of 1 mm depth were created using custom-built ARTcut® (Figure 1A), a software-controlled device for automatic wound placement in tissues developed by Fraunhofer ISC and IGB, Wuerzburg [23]. Osteochondral cylinders were fixed in support plate (Figure 1C) at specific x- and y-positions, and onset of drilling 1 mm chondral defects (z-axis) was determined with light beam (Figure 1B). For more details on the standardized defect creation process using ARTcut®, see supplementary information.

After creation of full thickness or 1 mm chondral defects, the cartilage defects were filled with cell embedded collagen type I hydrogels (**Error! Reference source not found.** 2). For subsequent *ex vivo* culture, osteochondral explants were transferred into custom-made culture platform [20] and cultured for 28 days with tissue specific media (Table 1) changed every 3-4 days. The *ex vivo* culture was carried out in a humidified atmosphere at 37°C and 5 % CO_2 (BBD 6220 CO_2 Incubator, Thermo Scientific™) either under normoxic (20 %) or physioxic (2 %) conditions.

2.2   Chondrocyte and mesenchymal stromal cells (MSC): isolation and *in vitro* expansion

Porcine chondrocytes were isolated from lateral condyles of 6-8 month-old domestic pigs by enzymatic digestion as previously described [20]. Chondrocytes were used at passage 0 – directly after isolation - for embedding in collagen type I hydrogel.

Porcine MSC were isolated from bone marrow aspirate (iliac crest) under the approval (reference number: 55.2 2532-2-256) of the District Government of Lower Franconia and the local animal welfare committee and performed according to the German Animal Welfare Act and the EU Directive 2010/63/EU. Following

gradient centrifugation using Ficoll-Paque (GE Healthcare Bio-Sciences, 17-5442-03), mononuclear cell

fraction was expanded in DMEM HG medium (GlutaMax, Gibco 61965) supplemented with 1 ng/mL FGF-

2 (R&D Systems, 233-FB-025), 10 % FCS (Gibco, 10270106) and 1 % Antibiotic-Antimycotic (10 U/mL

penicillin, 10 µg/mL Streptomycin, 0.25 µg/mL Amphotericin, Gibco, 15240) in humidified atmosphere

(37°C, 5 % CO_2). Media change was performed every 3-4 days. MSC were used at passage 2-3 for

embedding in collagen type I hydrogel.

2.3   Collagen type I hydrogel: hydrogel preparation and cell embedding

Cartilage defects were treated with collagen type I hydrogel. Collagen was isolated from rat tails (Wistar

IGS Rat; Charles River Laboratories), dissolved in 0.1 % (v/v) acetic acid to a concentration of 4.5 mg/mL,

and mixed with half volume of gel neutralizing solution (GNS) to achieve a final concentration of 3 mg/mL.

GNS consists of 2x DMEM high glucose (DMEM, PAA Laboratories, G0001,3010), 0.09 M HEPES (Sigma,

H3375), 3 % FCS (Bio&Sell, FCS.ADD.0500), 0.05 mg/mL chondroitin sulphate (Sigma, C4384) with pH value

adjusted to pH 8.5. Cell embedding was realized by mixing the porcine chondrocytes and/or MSCs in GNS,

resulting in a final total cell concentration of 20 Mio/mL hydrogel. Gelation was performed via incubation

of GNS collagen solution mixture for 20 min at 37°C under humidified atmosphere.

For implantation, two different cell laden constructs, as illustrated in Figure 2, were prepared and each

cultured under normoxic and physioxic conditions. 1) CHON were prepared by embedding of chondrocytes

in collagen type I hydrogel without expansion at a final density of 20 Mio/mL hydrogel volume; 2) MIX

were prepared by mixing 80 % MSC with 20 % chondrocytes at same cell density. CHON and MIX implants

were filled into the cartilage defect for gelation at 37°C.

2.4 Live-dead viability staining

To visualize a possible effect of defect creation process on cell viability of tissue samples, live-dead viability staining was performed (live-dead staining kit for mammalian cells, L3224, Invitrogen). Osteochondral explants were incubated with 4 µM Calcein AM and 2 µM ethidium homodimer-1 in DMEM HG and visualized with fluorescence microscope (494 nm/517 nm and 517 nm/617 nm wavelength) Keyence BZ-9000 (Biorevo). Living cells are stained by Calcein AM in green, dead cells are stained by ethidium homodimer-1 in red.

2.5 Histological evaluation

Osteochondral explants, harvested on day 0 and day 28, were washed with phosphate buffered saline, fixed for 24 h with 4 % formalin (Roti Histofix, Carl Roth P087.3) and processed with plastic embedding (Technovit T9100, Heraeus Kulzer, 66006735) as detailed described in the supplementary.

For immune-histological stainings, antigens were enzymatically retrieved depending on antibody (Table 2). Before incubation with primary antibody over night at 4°C, slides were blocked with 3 % (v/v) H_2O_2 (Carl Roth, 8070) and incubated with 5 % (w/v) bovine serum albumin. All stainings were visualized with Horseradish-peroxidase kit (DCS Innovative, Dako, K3468) following manufacturer`s instructions. Cell nuclei were counterstained with Mayer's hematoxylin (Morphisto, 11895), rehydrated and mounted with Entellan (Merck, 1079610500). All washing steps between blocking or chemical incubation were performed with PBS supplemented with 0.5 % Tween-20 (VWR, 8.22184).

2.6 Histological scoring

Histological scoring for the evaluation of the immunohistological stainings was performed according to the standardized guidelines of the international cartilage repair society (ICRS-II-Score). All stainings for collagens type I, II, X and aggrecan of this study ($n = 2$ biological replicates) were evaluated blinded by three independent operators according to the criteria summarized in Table 3. Maximum possible score for the here reported study was 12 points.

  ## 2.7   Statistics

Statistical analysis was performed with GraphPad Prism 6.07. Scoring values obtained from two independent experiments, each evaluated by three examinators were used for statistical analysis, resulting in $n = 6$ for each experimental group (Figure 2). Following tests for outlier and Gaussian distribution, the data was either compared using an unpaired t-test (normally distributed values) or a Mann-Whitney test (not normally distributed values). A p-value <0.05 was considered statistically significant.

3   Results

3.1   Chondral defect creation with ARTcut®

ARTcut® (Figure 1) represents an automated device for reproducible creation of chondral defects (1 mm in depth; Figure 3A). Main advantage of using the ARTcut® is the ability to create defined chondral defects with a flat bottom to allow a tide and uniform contact with any solid pre-formed cylindrical implant, for example bioprinted constructs with flat bottom.

The laser beam to detect the surface of every single osteochondral explant is the unique feature of the ARTcut® to define the onset of the drilling, thus all defects result with the same depth. Chondral cartilage defect induced with ARTcut® appeared as a well-defined rectangular defect boundary in cross section (diameter: 4 mm, height: 1 mm) with flat bottom (Figure 3B). In contrast, the manually created defect (using biopsy punch) resulted in a half-round shape and required removing of residual cartilage tissue with a second tool (*e.g.,* a sharp spoon) shown in Figure 3C. Following, the accuracy of chondral defect geometry was much lower in manual procedure - using biopsy punch - compared to software – using automated ARTcut®.

The automated defect creation took less than 5 sec and thus heating of tissue due to friction and rotation during drilling was very unlikely. Microscopic images of live-dead stainings of osteochondral explants with

1 mm cartilage defects (diameter: 4 mm) confirmed no indications of cell death after tissue defect

creation, neither with ARTcut® nor with biopsy punch (Figure 3D-E).

For the creation of full thickness cartilage defects, the ARTcut® was not required, since the bottom of the

defect was given by the cartilage-bone interface. This two-tissue interface was easily separated using

biopsy punch by pulling out the upper cartilage part.

3.2   Evaluation of cartilage treatments in the *ex vivo* model

The results of our study demonstrate the feasibility and reproducibility of the *ex vivo* cartilage test system

as a platform to compare cartilage repair strategies: Scoring results of the co-culture (MIX) are close to the

CHON treatment in chondral and full thickness defects. Results of the adapted ICRS-II scoring are shown

in Figure 4.

Under physioxic conditions, scoring results of defect repair after co-culture treatment (MIX) was

significantly higher (p = 0.0218) in chondral defect than in full thickness defects. Scoring values of CHON

treatment in chondral and full thickness experimental groups did not differ much (n.s. p = 0.1828). At

physioxic conditions the scoring value was significant higher for CHON treatment compared to MIX

treatment in full thickness defects (p = 0.0384).

Under normoxic conditions, full thickness defects reached higher scoring values compared to the chondral

defects.

Comparing the outcome of same defect depth and treatment group (cell type), but cultured under

physioxic or normoxic conditions, the oxygen tension plays only a significant role in chondral defects using

MIX (p = 0.0451, higher scoring values under physioxic conditions) and in full thickness defects treated

with CHON (p = 0.0316, lower scoring values under physioxic conditions).

Overall, treatment of the two defect depths with MIX showed slightly reduced scoring values compared

to the CHON treatment group. Only exception is MIX chondral (physioxia) with a slightly higher value as

the corresponding one for CHON. The results of the MIX treatment groups also tend to spread more than

the CHON treatment group independently of defect depth and oxygen tension.

4   Discussion

The choice of optimal treatment approach for cartilage repair is dependent on the size and depth of the

cartilage defect and thus the severity of pathological condition.

To mimic these different pathological defect scenarios, the implementation of the ARTcut® allowed to

overcome the low reproducibility of manual induced chondral defects. So far, there is no other automated

defect creation device published that has comparable features, namely possibility to select different

drilling parameters within one run and the application of one device for a wide range of soft and hard

tissues or tissue equivalents, as it is demonstrated with the ARTcut®. The induction of defects in

osteochondral explants allows to compare different cell-based treatment approaches by implantation of

cell loaded materials into the defects or to study the effect of defect depth on repair in an *ex vivo* model.

The software-controlled drilling parameters allow for standardized and reproducible defect creation in

other soft and hard tissues including bone tissue as well. *In vitro* as well as *ex vivo* studies require sample

procession at high throughput rates under a negative microbial environment. ARTcut® device addresses

these demands, resulting in higher methodological accuracy and reproducibility, compared to manual

procedure using biopsy punch.

Current treatments of small or large cartilage defects mainly aim on pain and symptom relief rather than

on functional tissue repair [27]. From a scientific point of view, there is way for improvement of cartilage

defects, especially on long term results. Material-assisted as well as cell-based therapies, including (M-)

ACI, AMIC and microfracture, tend to result in formation of mechanical inferior fibrocartilage in long term

follow up [27-29]. Moreover, the treatment of large defects with autologous cells are limited due to the

high number of chondrocytes needed for the current techniques. Therefore, in this laboratory-controlled

study, an *ex vivo* osteochondral defect model was modified to screen cell-based approaches to treat

chondral and full thickness cartilage defects. Different cell types, namely chondrocytes and MSCs, were

embedded in collagen type I hydrogel (CHON or MIX) and implanted in trauma induced chondral and full

thickness defects. A stimulative and beneficial effect of MIX treatment (20 % CHON and 80 % MSCs) was

observed regarding cartilaginous matrix formation with the main advantage to reduce the overall number

of CHON used for traditional ACI treatment.

Promising results regarding cartilage matrix production were obtained comparing CHON and MIX in

chondral and full thickness defects in the *ex vivo* model. The MIX treatment reduces the amount of

autologous cartilage tissue needed for chondrocyte isolation, since chondrocytes represented only 20 %

of the total cell number in our model. Several *in vitro* studies and *in silico* models have shown that

chondrocytes in co-culture with MSCs increase the chondrogenic differentiation potential [18, 19, 30, 31].

Scalzone *et al.* reported of an enhanced chondrocyte activity measured by proteoglycan and collagen type

II production after co-culture with MSC compared to chondrocyte monoculture both embedded in

chitosan based hydrogels [19].

One advantage of the co-culture approach – when clinically applied – would be the one-step procedure,

addressing the surgical intervention of the knee joint, for the treatment of the cartilage defect: The

reduced number of chondrocytes in the MIX treatment can be isolated intra-operatively without

expansion and thus reduced the risk of chondrocyte dedifferentiation [32]. Further, the required MSCs,

e.g., can be harvested independently form the knee surgery from bone marrow aspirate in a minimal

invasive procedure with subsequent expansion prior to the cartilage repair procedure to ensure a

sufficient number of MSCs.

Considering the absence of TGF-β in the culture media, which is known to drive MSC chondrogenesis [33],

the here reported cartilage repair solely derives from the stimulating effect of the surrounding

osteochondral tissue, cell-cell signaling of the cell laden implants and the culture conditions.

Results of a similar *ex vivo* model, based on horse and bovine osteochondral explants, suggested that the

presence of the osteochondral tissue in the explants increase cartilage-like matrix deposition compared to

free swelling conditions of cell seeded material only [21, 22].

Beside the cell source, the influence of oxygen tension (normoxia 20 % O_2, physioxia 2 % O_2) was studied

during the *ex vivo* culture on chondrogenic matrix deposition, resulting in different outcomes between the

treatment groups in chondral and full thickness defects. Based on the histological scoring, our results

showed that the oxygen tension only plays a role in chondral defects - marked by elevated matrix

production. In contrast, full thickness defects showed different response in cartilage-like matrix deposition

related to the treatment approach: While higher scoring values were reached with CHON treatment under

normoxic conditions, the pMIX treatment did not show differences comparing physioxic and normoxic

conditions. Physioxic conditions resulted in an increase in cartilaginous matrix deposition in chondral

defects compared to full thickness defects independently of the used cell type (CHON, MIX). Once the

defect reaches the cartilage-bone interface, the cells in the implant receive stimuli from the subchondral

bone that seem to counteract with the physioxic induced stimuli in the here presented model. One

explanation of the differential cell response to oxygen tension in chondral and full thickness defects may

originate from the variation in oxygen tension present in the human body. Chondrocytes are exposed to

lower oxygen tension in avascular cartilage (2-5 %) [34], synovial fluid and the synovial capsule (6.5-9.0 %)

[25, 35], with an increasing oxygen level in the bone marrow of subchondral bone (>7 %) [26, 36] and a

maximum of 12 % in arterial blood [37]. Once the cartilage defect progresses to the exposure of the

subchondral bone in patients, chondrocytes within the defect are exposed to higher levels of oxygen

present in the bone supplied by vascular invasion [38]. It has been shown that low oxygen inhibits the

degradation of hypoxia induced factors (HIF) [39]. HIFs are described to be essential for maintaining CHON

homeostasis and extracellular matrix synthesis and activate the transcription of genes [40].

While the here presented results are consistent, traceable and in line with literature, there is one limitation

in the current experimental setup: Due to limitations in availability of human healthy osteochondral tissue

– absence of any osteoarthritic phenotype – the authors decided to use porcine tissue instead, as

introduced by Schwab *et al.* [20]. The performance of an extensive comparative *ex vivo* study comprising

three parameters – namely 1) defect depth (chondral vs. full thickness), 2) different cell types for defect

filling and treatment (100 % chondrocytes vs. 20 % chondrocytes and 80 % MSCs), and 3) oxygen tension

(physioxia vs. normoxia) requires tissue material in the required quantity and of similar quality. Due to the

well-known donor issues associated with donor variability of human tissue and the already mentioned

limited availability of healthy human osteochondral tissue, this could not be achieved for human tissue

[41, 42]. In contrast, the use of porcine tissue isolated from one pig population (similar age; grown up

under equal conditions) minimizes the donor variation and represents a tissue source for healthy and non-

degenerated tissues with higher reproducibility.

Overall, the results obtained in this study strongly indicate that the surrounding tissue of the

osteochondral explants plays a crucial role in the defect repair, addressing chondral or full thickness

defects, and that the oxygen tension additionally stimulates the outcome in a defect depth dependent

way. However, the exact mechanism controlling success or failure of a treatment approach are not fully

understood and require further research. Despite the advantage of the here reported model, the mimicry

of the complex 3D environment present in the knee joint with the additional ability to reproducibly create

(trauma induced) defects, this model has its potential in pre-screening several biomaterials or treatment

approaches, but best candidates still require further pre-clinical and clinical testing to confirm obtained

results.

To conclude, in this *ex vivo* study two cell-based approaches (CHON vs. MIX embedded in collagen type I

hydrogel) for the treatment of chondral and full thickness defects, cultured under normoxic and physioxic

conditions, were compared on their cartilage repair potential. Independently of defect depth, the co-

culture of chondrocytes and MSC (MIX) has proven scoring results close to those of CHON treatment.

Hence, usage of MIX could be a promising approach to reduce the number of chondrocytes to one fifth

and thus the amount of tissue for harvest and accompanied side morbidities for the patient. Further, in

this experimental set up the oxygen tension showed an influence in a defect depth dependent way: an

increased extracellular matrix production resulted in chondral defect under physioxic conditions but was

not present in full thickness defects. This finding of defect dependent stimulus of oxygen tension on

cartilage repair has not been reported in literature before.

Acknowledgements

337 The authors thank Andreas Diegeler (Fraunhofer ISC, Bronnbach) for technical support of the ARTcut®.

338 Thanks also to Sebastian Naczenski for establishing the protocol for wounding of osteochondral explants

339 using ARTcut® device. Finally, we would like to thank Heike Walles for her support.

Author contribution

342 Conception and design of the study: Franziska Ehlicke and Andrea Schwab; Acquisition of data: Alexa

343 Buss and Andrea Schwab; Analysis and interpretation of data: Andrea Schwab, Alexa Buss, Franziska

344 Ehlicke and Oliver Pullig; Manuscript drafting: Andrea Schwab and Alexa Buss; Revision of the

345 manuscript: Franziska Ehlicke and Oliver Pullig; Final approval: Franziska Ehlicke, Andrea Schwab, Alexa

346 Buss and Oliver Pullig; Funding: Franziska Ehlicke and Alexa Buss.

Role of funding Sources

348 This work was financially supported by the European Union Seventh Framework Programme (FP7/2007-

349 2013) under grant agreement no 309962 (HydroZONES). Graduate School of Life Sciences (University of

350 Wuerzburg) is acknowledged for providing medical scholarship.

Competing interest statement

352 The authors declare no conflict of interest.

Ethics approval and consent to participate

354   Animal experiment was approved (reference number: 55.2 2532-2-256) by the District Government of

355   Lower Franconia and the local animal welfare committee and performed according to the German Animal

356   Welfare Act and the EU Directive 2010/63/EU. Following heparinization of the pig, porcine MSCs were

357   obtained by bone marrow aspiration.

References

[1] A.R. Martín, J.M. Patel, H.M. Zlotnick, J.L. Carey, R.L. Mauck, Emerging therapies for cartilage regeneration in currently excluded 'red knee' populations, npj Regenerative Medicine 4(1) (2019) 12.

[2] J.D. Wylie, M.K. Hartley, A.L. Kapron, S.K. Aoki, T.G. Maak, Failures and Reoperations After Matrix-Assisted Cartilage Repair of the Knee: A Systematic Review, Arthroscopy 32(2) (2016) 386-92.

[3] A.C. Gelber, M.C. Hochberg, L.A. Mead, N.Y. Wang, F.M. Wigley, M.J. Klag, Joint injury in young adults and risk for subsequent knee and hip osteoarthritis, Ann Intern Med 133(5) (2000) 321-8.

[4] W. Widuchowski, J. Widuchowski, T. Trzaska, Articular cartilage defects: study of 25,124 knee arthroscopies, Knee 14(3) (2007) 177-82.

[5] A.J. Sophia Fox, A. Bedi, S.A. Rodeo, The basic science of articular cartilage: structure, composition, and function, Sports Health 1(6) (2009) 461-8.

[6] E.A. Makris, A.H. Gomoll, K.N. Malizos, J.C. Hu, K.A. Athanasiou, Repair and tissue engineering techniques for articular cartilage, Nat Rev Rheumatol 11(1) (2015) 21-34.

[7] B.J. Cole, C. Pascual-Garrido, R.C. Grumet, Surgical management of articular cartilage defects in the knee, J Bone Joint Surg Am 91(7) (2009) 1778-90.

[8] A.K. Dewan, M.A. Gibson, J.H. Elisseeff, M.E. Trice, Evolution of autologous chondrocyte repair and comparison to other cartilage repair techniques, Biomed Res Int 2014 (2014) 272481.

[9] P. Niemeyer, T. Schubert, M. Grebe, A. Hoburg, Matrix-Associated Chondrocyte Implantation Is Associated With Fewer Reoperations Than Microfracture: Results of a Population-Representative, Matched-Pair Claims Data Analysis for Cartilage Defects of the Knee, Orthop J Sports Med 7(10) (2019) 2325967119877847.

[10] M. Brittberg, A. Lindahl, A. Nilsson, C. Ohlsson, O. Isaksson, L. Peterson, Treatment of deep cartilage defects in the knee with autologous chondrocyte transplantation, N Engl J Med 331(14) (1994) 889-95.

[11] M. Mumme, A. Barbero, S. Miot, A. Wixmerten, S. Feliciano, F. Wolf, A.M. Asnaghi, D. Baumhoer, O. Bieri, M. Kretzschmar, G. Pagenstert, M. Haug, D.J. Schaefer, I. Martin, M. Jakob, Nasal chondrocyte-based engineered autologous cartilage tissue for repair of articular cartilage defects: an observational first-in-human trial, Lancet (London, England) 388(10055) (2016) 1985-1994.

[12] R.L. Davies, N.J. Kuiper, Regenerative Medicine: A Review of the Evolution of Autologous Chondrocyte Implantation (ACI) Therapy, Bioengineering (Basel) 6(1) (2019).

[13] J.D. Lamplot, K.A. Schafer, M.J. Matava, Treatment of Failed Articular Cartilage Reconstructive Procedures of the Knee: A Systematic Review, Orthop J Sports Med 6(3) (2018) 2325967118761871.

[14] G. Merkely, T. Ogura, J. Ackermann, A. Barbieri Mestriner, A.H. Gomoll, Clinical Outcomes after Revision of Autologous Chondrocyte Implantation to Osteochondral Allograft Transplantation for Large Chondral Defects: A Comparative Matched-Group Analysis, Cartilage (2019) 1947603519833136.

[15] K. Mithoefer, T. McAdams, R.J. Williams, P.C. Kreuz, B.R. Mandelbaum, Clinical efficacy of
the microfracture technique for articular cartilage repair in the knee: an evidence-based
systematic analysis, Am J Sports Med 37(10) (2009) 2053-63.
[16] P.C. Kreuz, M.R. Steinwachs, C. Erggelet, S.J. Krause, G. Konrad, M. Uhl, N. Sudkamp, Results
after microfracture of full-thickness chondral defects in different compartments in the knee,
Osteoarthritis Cartilage 14(11) (2006) 1119-25.
[17] J.L. Carey, Fibrocartilage Following Microfracture Is Not as Robust as Native Articular
Cartilage: Commentary on an article by Aaron J. Krych, MD, et al.: "Activity Levels Are Higher
After Osteochondral Autograft Transfer Mosaicplasty Than After Microfracture for Articular
Cartilage Defects of the Knee. A Retrospective Comparative Study", JBJS 94(11) (2012).
[18] L. Bian, D.Y. Zhai, R.L. Mauck, J.A. Burdick, Coculture of human mesenchymal stem cells and
articular chondrocytes reduces hypertrophy and enhances functional properties of engineered
cartilage, Tissue engineering. Part A 17(7-8) (2011) 1137-1145.
[19] A. Scalzone, A.M. Ferreira, C. Tonda-Turo, G. Ciardelli, K. Dalgarno, P. Gentile, The interplay
between chondrocyte spheroids and mesenchymal stem cells boosts cartilage regeneration
within a 3D natural-based hydrogel, Sci Rep 9(1) (2019) 14630.
[20] A. Schwab, A. Meeuwsen, F. Ehlicke, J. Hansmann, L. Mulder, A. Smits, H. Walles, L. Kock, Ex
vivo culture platform for assessment of cartilage repair treatment strategies, ALTEX 34(2) (2017)
267-277.
[21] V.H.M. Mouser, N.M.M. Dautzenberg, R. Levato, M.H.P. van Rijen, W.J.A. Dhert, J. Malda, D.
Gawlitta, Ex vivo model unravelling cell distribution effect in hydrogels for cartilage repair,
ALTEX 35(1) (2018) 65-76.
[22] M.L. de Vries-van Melle, R. Narcisi, N. Kops, W.J. Koevoet, P.K. Bos, J.M. Murphy, J.A.
Verhaar, P.M. van der Kraan, G.J. van Osch, Chondrogenesis of mesenchymal stem cells in an
osteochondral environment is mediated by the subchondral bone, Tissue Eng Part A 20(1-2)
(2014) 23-33.
[23] A. Rossi, A. Appelt-Menzel, S. Kurdyn, H. Walles, F. Groeber, Generation of a three-
dimensional full thickness skin equivalent and automated wounding, J Vis Exp (96) (2015).
[24] G. Pattappa, R. Schewior, I. Hofmeister, J. Seja, J. Zellner, B. Johnstone, D. Docheva, P.
Angele, Physioxia Has a Beneficial Effect on Cartilage Matrix Production in Interleukin-1 Beta-
Inhibited Mesenchymal Stem Cell Chondrogenesis, Cells 8(8) (2019).
[25] G. Pattappa, B. Johnstone, J. Zellner, D. Docheva, P. Angele, The Importance of Physioxia in
Mesenchymal Stem Cell Chondrogenesis and the Mechanisms Controlling Its Response, Int J Mol
Sci 20(3) (2019).
[26] D.E. Anderson, B.D. Markway, D. Bond, H.E. McCarthy, B. Johnstone, Responses to altered
oxygen tension are distinct between human stem cells of high and low chondrogenic capacity,
Stem Cell Res Ther 7(1) (2016) 154.
[27] U. Horas, D. Pelinkovic, G. Herr, T. Aigner, R. Schnettler, Autologous chondrocyte
implantation and osteochondral cylinder transplantation in cartilage repair of the knee joint. A
prospective, comparative trial, J Bone Joint Surg Am 85(2) (2003) 185-92.
[28] J.R. Steadman, K.K. Briggs, J.J. Rodrigo, M.S. Kocher, T.J. Gill, W.G. Rodkey, Outcomes of
microfracture for traumatic chondral defects of the knee: average 11-year follow-up,
Arthroscopy 19(5) (2003) 477-84.

[29] J.L. Carey, Fibrocartilage following microfracture is not as robust as native articular
cartilage: commentary on an article by Aaron J. Krych, MD, et al.: "Activity levels are higher after
osteochondral autograft transfer mosaicplasty than after microfracture for articular cartilage
defects of the knee. A retrospective comparative study", J Bone Joint Surg Am 94(11) (2012)
e80.
[30] M.J. Chen, J.P. Whiteley, C.P. Please, A. Schwab, F. Ehlicke, S.L. Waters, H.M. Byrne,
Inducing chondrogenesis in MSC/chondrocyte co-cultures using exogenous TGF-β: a
mathematical model, Journal of Theoretical Biology 439 (2018) 1-13.
[31] T.W. Kim, M.C. Lee, H.C. Bae, H.-S. Han, Direct Coculture of Human Chondrocytes and
Synovium-Derived Stem Cells Enhances In Vitro Chondrogenesis, Cell J 20(1) (2018) 53-60.
[32] K. von der Mark, V. Gauss, H. von der Mark, P. Muller, Relationship between cell shape and
type of collagen synthesised as chondrocytes lose their cartilage phenotype in culture, Nature
267(5611) (1977) 531-2.
[33] B. Johnstone, T.M. Hering, A.I. Caplan, V.M. Goldberg, J.U. Yoo, In vitro chondrogenesis of
bone marrow-derived mesenchymal progenitor cells, Exp Cell Res 238(1) (1998) 265-72.
[34] S. Zhou, Z. Cui, J.P.G. Urban, Factors influencing the oxygen concentration gradient from
the synovial surface of articular cartilage to the cartilage–bone interface: A modeling study,
Arthritis & Rheumatism 50(12) (2004) 3915-3924.
[35] K. Lund-Olesen, Oxygen tension in synovial fluids, Arthritis Rheum 13(6) (1970) 769-76.
[36] A. Carreau, B. El Hafny-Rahbi, A. Matejuk, C. Grillon, C. Kieda, Why is the partial oxygen
pressure of human tissues a crucial parameter? Small molecules and hypoxia, J Cell Mol Med
15(6) (2011) 1239-53.
[37] M. Marenzana, T.R. Arnett, The Key Role of the Blood Supply to Bone, Bone Research 1(1)
(2013) 203-215.
[38] J. Filipowska, K.A. Tomaszewski, L. Niedzwiedzki, J.A. Walocha, T. Niedzwiedzki, The role of
vasculature in bone development, regeneration and proper systemic functioning, Angiogenesis
20(3) (2017) 291-302.
[39] A. Kremer, M. Wussmann, M. Herrmann, M. Raghunath, H. Walles, Ciclopirox olamine
promotes the angiogenic response of endothelial cells and mesenchymal stem cells, Clin
Hemorheol Microcirc 73(2) (2019) 317-328.
[40] J. Fernandez-Torres, Y. Zamudio-Cuevas, G.A. Martinez-Nava, A.G. Lopez-Reyes, Hypoxia-
Inducible Factors (HIFs) in the articular cartilage: a systematic review, Eur Rev Med Pharmacol
Sci 21(12) (2017) 2800-2810.
[41] H.X. Chang, L. Yang, Z. Li, G. Chen, G. Dai, Age-related biological characterization of
mesenchymal progenitor cells in human articular cartilage, Orthopedics 34(8) (2011) e382-8.
[42] M. Kim, I.E. Erickson, A.H. Huang, S.T. Garrity, R.L. Mauck, D.R. Steinberg, Donor Variation
and Optimization of Human Mesenchymal Stem Cell Chondrogenesis in Hyaluronic Acid, Tissue
Eng Part A 24(21-22) (2018) 1693-1703.
[43] A. Schwab, Development of an osteochondral cartilage defect model, Julius-Maximilians
University, Wuerzburg 2017.

Figure Legends

Figure 1: Artificial tissue cutter (ARTcut®): A) View of the computerized numerical controlled device equipped with B) optical barrier (white arrow indicates the laser light beam) and a C) supporting plate to fix the samples and place on the bottom of the machine (marked with black star). Reprinted from Schwab 2017 [43]

Figure 2: Schematic overview of all experimental groups: Full thickness and chondral defects of osteochondral explants were treated with collagen type I hydrogel (grey) either containing chondrocytes (CHON, blue) or MIX (80 % MSC and 20 % chondrocytes), implanted in chondral or full thickness defects and cultured *ex vivo* under normoxic (20 % O_2, blue frame) or physioxic conditions (2 % O_2, red frame).

Figure 3: Osteochondral explant cross section with chondral defect. A) Schematic illustration of A) 4 mm chondral defect. B-E) Comparison of chondral defect created automated with ARTcut® (B) and manually with biopsy punch (C) (scale bar 500 µm). D-E) Live-dead staining of explant cross section highlighting accurate borders of defect induced with ARTcut® (D) and more irregular shaped geometry of manually induced defect (E) (scale bar 200 µm). Reprinted from Schwab 2017 [43].

Figure 4: Evaluation of defect repair after 28 days *ex vivo* culture. Macroscopic images of full thickness and chondral defects (diameter 4 mm) treated with A) porcine Chondrocytes (CHON) or B) co-culture of porcine MSC and CHON (MIX). Red frame indicates culture under physioxic, blue frame under normoxic oxygen tension. Macroscopically, there are no obvious differences comparing full thickness and chondral defects. C-D) Scoring evaluation of cartilage repair strategies in the *ex vivo* model (mean ± standard deviation with single scoring results as dots resp. listed values for 1st and 2nd biological replicate). Unpaired t-test * $p < 0.05$, ** $p < 0.01$.

504   Figure 1

supporting plate

506   Figure 2

508   Figure 3

510   Figure 4

D

		CHON		MIX	
	Cumulative score (max 12)	Physioxia (2 % O_2)	Normoxia (20 % O_2)	Physioxia (2 % O_2)	Normoxia (20 % O_2)
full thickness defect	1st biological replicate	9; 8; 8	9; 9; 7	6; 7; 4	8; 8; 7
full thickness defect	2nd biological replicate	7; 6; 6	9; 11; 10	5; 7; 5	5; 5; 6
full thickness defect	Mean ± standard deviation	7.33 ± 1.21	9.17 ± 1.33	5.67 ± 1.21	6.5 ± 1.38
chondral defect	1st biological replicate	7; 8; 9	7; 7; 7	11; 9; 12	4; 2; 2
chondral defect	2nd biological replicate	8; 9; 8	9; 8; 7	7; 7; 6	8; 8; 7
chondral defect	Mean ± standard deviation	8.17 ± 0.75	7.50 ± 0.84	8.67 ± 2.42	5.17 ± 2.86

512   Table 1: Composition of bone and cartilage media for *ex vivo* culture of osteochondral explants.

	Cartilage media	Bone media	Supplier
DMEM GlutaMAX (high glucose)	98 %	89 %	Gibco 61965
Antibiotics	10 U/mL penicillin, 10 µg/mL Streptomycin, 0.25 µg/mL Amphotericin (1 % (v/v))	10 U/mL penicillin, 10 µg/mL Streptomycin, 0.25 µg/mL Amphotericin (1 % (v/v))	Gibco, 15240
Sodium-pyruvate	1 mM	-	Gibco, 11360
L-ascorbic acid-2-phosphate	50 µg/mL	50 µg/mL	Wako chemicals, 013-19641
L-proline	40 µg/mL	-	Sigma-Aldrich, P5607
Dexamethasone	100 nM	100 nM	Sigma-Aldrich, D4902
fetal bovine serum (FBS)		10 % (v/v)	Gibco, 10270106
Insulin-transferrin-selenin	1 % (v/v)	-	ITS$^+$-premix, Gibco, Life Technologies, 32430
β-glycerophosphate	-	10 mM	Sigma-Aldrich, G9422

514 Table 2: List of antibodies for immune-histological stainings.

Antibody	clone	Dilution and final antibody concentration	Antigen retrieval	Antibody supplier
Collagen type I	EPR7785	1:1000; 0.875 µg/mL	pepsin & hyaluronidase	Abcam, ab138492
Collagen type II	II-4C11	1:1000; 1 µg/mL	pepsin & hyaluronidase	MP Bio, 63171
Collagen type X	COL-10	1:4000; 0.25 µg/mL	pronase	Sigma-Aldrich, C7974
Aggrecan	EPR7785	1:2000; 0.5 µg/mL	pronase	Invitrogen, AHP0012

517   Table 3: Criteria for evaluating immune histological stainings with modified ICRS-II-Score with maximum of 12 points (0-3 scores

518   for each criteria judging the uniformity and darkness of aggrecan, collagen type I, II and X staining).

ICRS-II-Scoring	Uniformity and darkness* of Aggrecan stain
	0: No stain, 1: Weak staining of poorly formed matrix, 2: Moderately even staining ,3: Even dark stain
	Uniformity and darkness* of Collagen type II stain
	0: No stain, 1: Weak staining of poorly formed matrix, 2: Moderately even staining ,3: Even dark stain
	Uniformity and darkness* of Collagen type I stain
	3: No stain, 2: Weak staining of poorly formed matrix, 1: Moderately even staining 0: Even dark stain
	Uniformity and darkness* of Collagen type X stain
	3: No stain, 2: Weak staining of poorly formed matrix, 1: Moderately even staining 0: Even dark stain