Title: Genome-wide analysis of long noncoding RNAs, 24-nt siRNAs, DNA

methylation and H3K27me3 marks in *Brassica rapa* and related species

Hasan Mehraj[1], Daniel J. Shea[2], Satoshi Takahashi[3], Naomi Miyaji[1], Ayasha Akter[1],

Motoaki Seki[3,4,5], Elizabeth S. Dennis[6,7], Fujimoto Ryo[1], Kenji Osabe[8*]

Affiliation:

[1]Graduate School of Agricultural Science, Kobe University, Rokkodai, Nada-ku, Kobe

657-8501, Japan

[2] Bloomberg LP, Chiyoda-ku, Tokyo 100-0005, Japan

[3]Riken center for Sustainable Resource Science, Yokohama, Kanagawa, 230-0045,

Japan

[4]RIKEN Cluster for Pioneering Research, 2-1 Hirosawa, Wako, Saitama, 351-0198,

Japan

[5]Kihara Institute for Biological Research, Yokohama City University, 641-12

Maioka-cho, Totsuka-ku, Yokohama, Kanagawa 244-0813, Japan

[6]CSIRO Agriculture and Food, Canberra, ACT 2601, Australia

[7] University of Technology, Sydney, PO Box 123, Broadway, NSW 2007, Australia

[8]Institute of Scientific and Industrial Research, Osaka University, Mihogaoka,

Ibaraki-city, Osaka, Japan

*Corresponding author: o3b@sanken.osaka-u.ac.jp

21   1

Abstract

Long noncoding RNAs (lncRNAs) are RNA fragments that generally do not code for a protein but are involved in epigenetic gene regulation. In this study, lncRNAs of *Brassica rapa* were classified into long intergenic noncoding RNAs, natural antisense RNAs, and intronic noncoding RNAs and their expression were analyzed in relation to genome-wide 24-nt small interfering RNAs (siRNAs), DNA methylation, and histone H3 lysine 27 trimethylation marks (H3K27me3). More than 65% of the lncRNAs analyzed consisted of one exon, and more than 55% overlapped with inverted repeat regions (IRRs). Overlap of lncRNAs with IRRs or genomic regions encoding for 24-nt siRNAs resulted in increased DNA methylation levels and when both were present, there were further increase in DNA methylation levels. LncRNA did not overlap greatly with H3K27me3 marks, but the expression level of intronic noncoding RNAs that did coincide with H3K27me3 marks was higher than without H3K27me3 marks. The Brassica genus comprises important vegetables and oil seed crops grown across the world. *Brassica rapa* is a diploid (AA genome) thought to be one of the ancestral species of both *B. juncea* (AABB genome) and *B. napus* (AACC) through genome merging (allotetrapolyploidization). Complex genome restructuring and epigenetic alterations are thought to be involved in these allotetrapolyploidization events, but the detailed mechanism is not known. Comparison of lncRNAs between *B. rapa* and *B. nigra*, *B. oleracea*, *B. juncea*, and *B. napus* showed the highest conservation with *B.*

oleracea. This study presents a comprehensive analysis of the epigenome structure of

the *B. rapa* at multi-epigenetic levels (siRNAs, DNA methylation, H3K27me3, and

lncRNAs) and offers insights into the function of lncRNA in the Brassica genus.

Key words: Brassica rapa, noncoding RNA, siRNA, DNA methylation, H3K27

methylation, epigenome, Brassica nigra, Brassica oleracea, Brassica juncea, Brassica

napus

50   Introduction

The Brassica genus comprises vegetable and oil seed crops. The "Triangle of

U" proposed the genomic relationship among six major species of the Brassica genus.

Three allotetraploid species, each of which contains two complete diploid genomes

derived from two different parental species, *Brassica juncea* L. (AABB genome,

$2n = 4x = 36$), *B. napus* L. (AACC, $2n = 4x = 38$), and *B. carinata* L. (BBCC,

$2n = 4x = 34$) are derived from the natural hybridization of the diploid species, *B. rapa* L.

(AA, $2n = 2x = 20$)*, B. nigra* L. (BB, $2n = 2x = 16$), and *B. oleracea* L. (CC,

$2n = 2x = 18$) [1]. Some species in the Brassica genus show morphological divergence

(termed morphotype). *B. rapa* includes leafy vegetables including Chinese cabbage (var.

pekinensis), pak choi (var. *chinensis*), and komatsuna (var. *perviridis*), root vegetables

including turnip (var. *rapa*), and oilseed crops (var. *oleifera*) [2]. The first whole

genome sequence determined in the genus Brassica was that of *B. rapa* [3]. Later the

whole genome sequences of *B. oleracea, B. nigra*, *B. napus,* and *B. juncea* were

determined [4–7].

Plant transcriptome analyses have revealed transcribed RNAs devoid of

protein-coding potential, which are called noncoding RNAs (ncRNAs). Some ncRNAs

contain exons, which potentially code for short proteins or peptides, however,

experimental investigation is required to validate their functions. Two families of

ncRNAs are known; long noncoding RNAs (lncRNAs) that are longer than 200

nucleotides (nt), and small RNAs (sRNAs) that are ∼18-30 nt in length [8–12].

LncRNAs are classified by their position and orientation of transcription; long

intergenic noncoding RNAs (lincRNAs), intronic noncoding RNAs (incRNAs) derived

from introns, and natural antisense transcripts (NATs) transcribed from the

complementary DNA strand of their associated genes [8-12]. Recent studies have

shown that lncRNAs play crucial roles in various physiological processes such as

vernalization [13-15], photoperiod-sensitive male sterility [16], red-light-mediated

seedling photomorphogenesis [17], seed dormancy [18], and the transcriptional

regulation of plant innate immunity [19].

Epigenetics has been defined as ''the study of changes in genome expression

that are mitotically and/or meiotically heritable and that do not entail a change in DNA

sequence'' [20]. DNA methylation and histone modification are well-known epigenetic

modifications, and lncRNAs are considered to be involved in epigenetic regulation [21].

In plants, DNA methylation is established through RNA-directed DNA methylation

(RdDM) [22,23]. Plant-specific RNA POLYMERASE IV (Pol IV) and Pol V are

involved in RdDM. Pol IV-transcribed ncRNAs are cleaved into 24-nucleotide small

interfering RNAs (24-nt siRNAs) by DICER-LIKE 3 (DCL3), and 24-nt siRNAs are

loaded into ARGONAUTE 4 (AGO4). PolV-transcribed lncRNAs act as scaffold

molecules of AGO-siRNA complexes, which recruit DOMAINS REARRANGED

METHYLTRANSFERASE 2 (DRM2) for catalyzing *de novo* DNA methylation. In

Arabidopsis thaliana two cold-induced lncRNAs (COLDAIR and COLDWRAP)

regulate histone methylation by recruiting polycomb repressive complex 2 (PRC2),

which catalyzes the tri-methylation of histone H3 lysine 27 (H3K27me3), to the

chromatin region of *FLOWERING LOCUS C* (*FLC*) [14,24-26].

Previous studies have identified lncRNAs associated with different lines,

tissues or environmental changes in Brassica [27–33]. However, there are few reports of

the association between lncRNAs and epigenetic modifications in *B. rapa*. In this study,

we examined the DNA methylation levels and H3K27me3 levels in the region covering

lncRNAs, and found an association between lncRNAs and inverted repeat regions

(IRRs) or 24-nt siRNAs, but association with H3K27me3 to a specific type of lncRNA.

Materials and methods

Plant materials and growth conditions

Six lines of *B. rapa*, inbred lines of RJKB-T24 (var. *pekinensis*) [34] and Yellow sarson (var. *trilocularis*), doubled haploid lines of BRA2209 (var. *rapa*), Homei (var. *pekinensis*), and Osome (var. *perviridis*), and a commercial F_1 hybrid cultivar, 'Harunosaiten' (var. *pekinensis*) (Watanabe Seed Co., Ltd.) were used. Three *B. oleracea* cabbage (var. *capitata*) F_1 hybrid cultivars, 'Reiho', 'Matsunami' (Ishii Seed Growers CO., LTD), and 'Kinkei 201' (Sakata Seed Co., Ltd.) were also used.

Seeds were surface sterilized and grown on agar solidified Murashige and Skoog (MS) plates with 1 % (w/v) sucrose under long day (LD) condition (16h light / 8h dark) at 21 °C. Fourteen-day first and second leaves of *B. rapa* and 19-day first and second leaves of *B. oleracea* were harvested for isolation of genomic DNA or total RNA.

RJKB-T24 was used for RNA-sequencing (RNA-seq) for detection of lncRNAs. Briefly, after performing QC of the sequenced reads, putative mRNAs were identified by aligning the sequence reads to the *B. rapa* reference genome v1.5 using HISAT2 and then assembling transcripts with Stringtie. Assembled transcripts with a mapping code of 'u', indicating they are intergenic but not part of the annotated reference genome, were then compared to the SwissProt database using blastx (e-value 1e-10). Transcripts with hits were classified as putative mRNAs, while transcripts with

121 no hits were classified as putative lincRNAs. A more detailed description of this method

122 can be found in Shea et. al, 2018 [27].

124 DNA extraction and PCR

125   Genomic DNA was isolated by the Cetyl trimethyl ammonium bromide

126 (CTAB) method [35]. The PCR reaction was performed using the following conditions;

127 1 cycle of 94 °C for 3 min, 35 cycles of 94 °C for 30 s, 55 °C for 30 s, and 72 °C for 1

128 min, and final extension at 72 °C for 3 min. The PCR products were electrophoresed on

129 1.0% agarose gel. Primer sequences used in this study are shown in S1 Table.

131 RNA extraction and RT-PCR

Total RNA from the first and second leaves were isolated by SV Total RNA

Isolation System (Promega Co., WI, USA). To analyze lncRNA expression, cDNA was

synthesized from 500 ng total RNA using PrimeScript RT reagent Kit (Takara Bio.,

Shiga, JAPAN). The absence of genomic DNA contamination was confirmed by PCR

using a control without reverse transcriptase. The PCR conditions were 94 °C for 2 min

followed by 35 cycles of 94 °C for 30 s, 55 °C for 30 s, and 68 °C for 30 s. The primers

used in this study are listed in S1 Table.

140 Detection of epigenetic states in lncRNA regions

To examine the epigenetic states (DNA methylation levels, H3K27me3, and

24-nt siRNA levels) of lncRNA encoding regions in *B. rapa*, we used previous

sequence data of whole genome bisulfite sequencing (WGBS) [36], chromatin

immunoprecipitation sequencing (ChIP-seq) [37], and small RNA-sequencing

(sRNA-seq) data [36], which were generated using samples from the same line, tissue,

and developmental stages but harvested independently.

The reads of WGBS were mapped to the *B. rapa* reference genome v.1.5

using Bowtie2 version 2.2.3 and Bismark v0.14.3, and data covering genomic regions

encoding lncRNA in chromosomes A01 to A10 were extracted. In order to estimate the

methylation levels of CG, CHG, and CHH contexts, the numbers of methylated and

unmethylated reads were extracted for each cytosine position using bismark methylation

extractor script with the paired-end parameter. The methylation level at each cytosine

position was calculated by dividing the number of methylated cytosines (mC) reads by

the total number of reads.

The reads of sRNA-seq were mapped to the *B. rapa* reference genome v.1.5

using Bowtie2 version 2.2.3. We classified the alignment reads by length, and the 24-nt

aligned reads covering the lncRNA regions in chromosomes A01 to A10 were

extracted.

The reads of ChIP-seq using anti-H3K27me3 (Millipore, 07-449) antibodies

were mapped to the *B. rapa* reference genome v.1.5 using Bowtie2 version 2.2.3, and

data covering genomic regions encoding for lncRNA in chromosomes A01 to A10 were

extracted.

Comparison of putative mRNA and lncRNAs in *B. rapa* to other

related species of *Brassica*

The putative mRNAs and lncRNAs were first compared to the reference

genomes of *B. nigra*, *B. oleracea*, *B. juncea*, and *B. napus* by best-hit blastn (e-value

1e-10) to identify homologous regions in closely related Brassica species [6,7,38]. In

order to parse the local High-scoring segment pair (HSP) alignments produced by blastn,

genBlastA was used to produce a representative putative gene that is homologous to the

query [39]. The parsed local alignments were then analyzed by a custom python script

(available at http://www.github.com/danshea/lncRNA) to examine the overall alignment

length and computed coverage of the aligned homologous region to the putative mRNA

or lncRNA query transcript sequence. These results were then imported into R and

plotted to assess overall relative coverage of the sequences among the Brassica species.

Results

Characterization of lncRNA in *B. rapa*

We identified 1,444 lincRNAs, 551 NATs, and 93 incRNAs using RNA-seq

data of 14-day first and second leaves with and without four weeks of cold treatment

[27]. In order to investigate the relationship between lncRNAs and epigenetic

modifications or the species specificity of lncRNAs, we analyzed in more detail the

RNA-seq data of the 14-day first and second leaves without cold treatment. There was

no strong bias in the chromosomal distribution of expressed lncRNAs (S1 Fig). More

than 65% of lncRNAs contained one exon (lincRNAs, 65.7%; NATs, 72.4%; incRNAs,

71.0%), whereas the proportion of mRNAs containing only one exon is 15.1% (Fig 1A).

The mean transcript lengths of lincRNAs (725 nt) and incRNAs (779 nt) were shorter

than that of NATs (1,271 nt) and mRNAs (1,305 nt, Fig 1B). About 40% of lincRNAs

were located within 2 kb of the genic regions, and about 10% of lincRNAs were located

more than 20kb from the genic region (Fig 1C). In this study, we focused on lncRNAs

mapped to the chromosomes A01 to A10 as previous WGBS, ChIP-seq, and sRNA-seq

have omitted the placed scaffolds for their analyses [36,37]. 763 of 1,173 (65.0%)

lincRNAs, 291 of 529 (55.0%) NATs, and 66 of 92 (71.7%) incRNAs overlapped with

IRRs such as transposable elements (TEs) detected by RepeatMasker, suggesting that

IRRs are the source of more than half of lncRNAs in *B. rapa* (Fig 2).

Fig 1. Analysis of lncRNAs of 14-day first and second leaves of *B. rapa*. (A) Number

of exons in lncRNA. (B) Nucleotide length of lincRNA, NAT, incRNA, putative

mRNA and mRNA. "a" and "b" represent significant differences by one-way ANOVA

test ("*", $p<0.05$; "***", $p<0.001$). (C) The proportion lncRNA distances to the nearest

gene.

Fig 2. Proportion of each type of lncRNA and putative mRNAs that overlap with

IRRs and siRNA. "+" indicates with and "-" indicated without overlapping IRRs or

genomic regions encoding siRNAs.

Relationship between lncRNAs and siRNA or DNA methylation

We have performed sRNA-seq using 14-day first and second leaves, which

are identical developmental stages and tissues to previously analyzed samples [36], but

independently harvested for this study. We identified the lncRNAs having perfect

sequence identity to genomic regions encoding for 24-nt siRNAs. 219 of 1,173 (18.7 %)

lincRNAs, 74 of 529 (14.0%) NATs, and 16 of 92 (17.4%) incRNAs in A01-A10

overlapped with unique-mapped genomic regions encoding for 24-nt siRNAs, and more

than 80% of the lncRNAs overlapping with genomic regions encoding for 24-nt siRNAs

were from regions that harbored IRRs (Fig 2). 24-nt siRNAs were mapped in a similar

way to lncRNA and its 5' and 3' flanking regions; this mapping pattern is different from

those of genic regions or IRRs (Fig 3).

Fig 3. 24nt-siRNAs mapped to genic region, IRRs, putative mRNAs and each type

of lncRNAs. The x-axis represents the target region (genic region, IRRs, putative

mRNAs, or lncRNAs) and the flanking 5' and 3' regions. The y-axis represents the

reads per million (RPM) value of 24-nt siRNA that overlaps with the corresponding

target region.

We examined the whole genome DNA methylation state by WGBS using the

same 14-day first and second leaves [36]. The average DNA methylation levels in

regions covering lncRNAs were similar to those of the whole genome (S2 Fig). DNA

methylation level in regions producing NATs was lower than those producing any of

the three types of lncRNAs. Overlap of lncRNAs with IRRs or genomic regions

encoding for 24-nt siRNAs was associated with increased DNA methylation levels and

overlapping with both caused further increases in DNA methylation levels (Fig 4).

Fig 4. DNA methylation level of genomic regions encoding each RNA type with

siRNAs or IRRs. "+" indicates with and "-" indicates without overlapping genomic

regions encoding for siRNAs or IRRs.

The DNA methylation were similar in lincRNA and NAT encoding regions,

and DNA methylation levels in the 5' and 3' flanking regions were higher than in the

body regions encoding lincRNAs and NATs (S3 Fig). The DNA methylation levels

over the incRNA encoding regions and the flanking regions did not change (S3 Fig).

Overall, mapping of DNA methylation levels to lncRNA encoding regions were

different from those of genic regions or IRRs (S3 Fig).

We examined the influence on the lncRNA expression level when IRRs or

siRNA clusters on the genome spanned the lncRNA regions. IRRs did not affect the

expression levels of lncRNAs (Fig 5). The expression levels of lncRNAs covered with

siRNA clusters were higher than those not covered by siRNA clusters (Fig 5)

Fig 5. Expression level of each type of RNAs with (+) or without (-) overlapping

IRRs, 24nt-siRNAs, or H3K27me3. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$ (Student

t-test)

Relationship between lncRNAs and H3K27me3

We have examined the H3K27me3 distribution using the 14-day first and

second leaf samples [37] and identified the genomic regions encoding for lncRNAs that

have H3K27me3 marks. 127 of 1,173 (10.8 %) lincRNAs, 83 of 529 (15.7%) NATs,

and 15 of 92 (16.3%) incRNAs encoding genomic regions had H3K27me3 marks (S4

Fig). H3K27me3 was enriched in the transcribed region of lncRNAs, similar to the

genic regions, but H3K27me3 levels in lncRNA encoding genomic regions were lower

than in regions coding for mRNAs (S5 Fig). The expression level of incRNAs was

higher when the encoding regions had H3K27me3 than without H3K27me3 (Fig 5). In

NATs and lincRNAs, there was no difference of expression level with and without

H3K27me3 on their encoding regions (Fig 5).

Characterization of un-annotated genes

In 2,052 lncRNAs mapped to intergenic regions of the genome, 608

transcripts had hits (e-value < 1.0e-10) against the Swissport database using BLASTX,

indicating they could be un-annotated genes. The expression levels of the putative

mRNAs from these regions were similar to that of mRNA [27]. Putative mRNAs tended

to have fewer exons compared with annotated mRNAs (Fig 1A). The mean transcript

length of putative mRNA genes was similar to that of mRNAs (Fig 1B).

314 of 490 (64.1%) putative mRNAs in A01-A10 overlapped with IRRs (Fig

2) but the expression level of putative mRNAs overlapping with IRRs was similar to

those not-overlapping with IRRs (Fig 5). Mapping of 24-nt siRNAs and DNA

methylation levels to genomic regions encoding putative mRNAs were similar to that of

genic regions (Fig 3, S3 Fig). The average expression level of putative mRNAs with

overlapping 24-nt siRNAs was higher than that without any overlap (Fig 5). The pattern

of the average of DNA methylation level in the genomic region encoding putative

mRNAs and their flanking regions was similar to that to the average of the genic

regions (S3 Fig). Overlapping IRRs or 24-nt siRNAs resulted in an increase in DNA

methylation levels and overlapping with both causes further increase in DNA

methylation levels (Fig 4).

53 of 490 (10.8%) genomic regions corresponding to putative mRNAs had

H3K27me3. H3K27me3 was enriched in the genomic regions encoding for lncRNAs,

especially around the transcription start site, and this was similar to the pattern in the

genic region (S5 Fig). The expression levels of putative mRNAs with corresponding

genomic H3K27me3 were lower than those without H3K27me3, but there was no

significant difference (Fig 5).

Examination of the conservation of lncRNAs among the Brassica

genus

Using the sequences of the 2,088 lncRNAs of *B. rapa* , a best-blast hit search

against the *B. nigra*, *B. oleracea*, *B. juncea*, and *B. napus* reference genomes was

conducted using GenBlastA (e-value = 1e-10) to examine the conservation of lncRNAs.

B. rapa lncRNAs were most conserved in *B. oleracea* and moderate conservation was

observed in *B. nigra* and *B. napus*. LncRNAs were less conserved in *B. juncea*,

especially in the incRNAs (Fig 6).

Fig 6. BLAST search of each type of lncRNA from *B. rapa* against *B. nigra*, *B.*

***oleracea, B. juncea,* and *B. napus*.**

We selected twelve lncRNAs that showed high sequence similarity with the *B.*

oleracea reference genome. We tested whether these lncRNA coding genomic

sequences were conserved in three commercial cultivars of cabbage (*B. oleracea*) by

PCR using genomic DNA as template. PCR amplification of genomic regions

corresponding to all twelve lncRNAs was confirmed in all three *B. oleracea* lines. Next,

we tested by RT-PCR whether these putative lncRNAs in *B. oleracea* are expressed in

the first and second leaves. Six of twelve lncRNAs were expressed in all three cultivars.

One lncRNA was expressed in two of three lines, and one lncRNA was expressed in

one of three lines. The remaining four lncRNAs were not expressed or were slightly

expressed in all three lines (Table 1). We also examined the variation within *B. rapa*

species using six lines. Seven of the twelve lncRNAs were expressed in all six lines,

and the remaining five lncRNA showed line-specificity (Table 1).

Table 1. Conservation and diversity of the expression of lncRNAs within *B. rapa*

species or between species. RT-PCR results of each lncRNAs "+++" indicates strong

PCR band, "+" indicates weak PCR band, and "-" indicates no PCR band.

lncRNAs	B. rapa						B. oleracea		
	RJKB-T23	Homei	Harunosaiten	BRA2209	Osome	Yellow Sarson	Reiho	Matsunami	Kinkei
M15784	+++	-	+++	-	+++	+++	-	-	-
M26919	+++	+++	+++	+++	+++	+++	+++	+++	+++
M3316	+++	+++	+++	+++	+++	+++	+++	+++	-
M491	+++	-	+++	+++	+++	-	+	+	-
M17356	+++	+	-	-	-	-	-	-	-
M17153	+++	-	-	-	-	+	+++	+++	+++
M25534	+++	+++	+++	+++	+++	+++	+++	+++	+++
M259	+++	+++	+++	+++	+++	+++	+++	+++	+++
M4317	+++	+++	+++	+++	+++	+++	+++	+++	+++
M26796	+++	+++	+++	+++	+++	+++	+++	+++	+++
M4921	+++	+++	+	+	+++	+++	-	+++	-
M24531	+++	+++	+++	+++	+++	+++	-	+	-

Discussion

We analyzed 2,088 lncRNAs in leaves of *B. rapa*; their characteristics such as

number of exons, length of transcripts (except for NATs), and lower expression levels

were similar to those reported in other plant species [40–45]. The total number of

lncRNAs seems to be similar among species [46,47], but there tends to be low sequence

conservation between different species [48]. Lower conservation of lncRNAs than

mRNAs was observed between *B. napus* and its two ancestral species, *B. rapa* and *B.*

oleracea [29,42], even though the hybridization was relatively recent event,

~1,910–7,180 years ago [49]. We also found a lower sequence conservation of *B. rapa*

lncRNAs among other species of the genus Brassica (*B. nigra*, *B. oleracea*, *B. juncea*,

and *B. napus*). In 12 selected highly conserved lncRNAs, we found conservation of the

lncRNAs at both the sequence and transcriptional level between *B. rapa* and *B. oleracea*,

but there was transcriptional variation within species agreeing with the IRR

diversification of each species after the allotetraploidization of *B. napus* [50]. The

analysis of sequence homology of lncRNAs in *B. rapa* with other members of Brassica

genus may deepen the understanding of the evolutionary dynamics of lncRNAs in the

genus Brassica.

We found over 55% of each lncRNAs (65.0% lincRNAs, 55.0% NATs, and

71.7% incRNAs) overlapped with IRRs including TEs throughout the *B. rapa* genome.

TEs are considered a source of siRNAs [51], and lincRNAs identified in *A. thaliana*,

rice, and maize, were associated with 22.2%, 49.7%, and 51.5% of TEs, respectively

[52]. In this study, we found 18.7% lincRNAs, 14.0% NATs, and 17.4% of incRNAs

covered unique-mapped 24-nt siRNAs, and over 80% of lncRNAs covering the

genomic regions encoding for 24-nt siRNAs overlapped with IRRs with perfect

sequence match, suggesting that lncRNAs covering IRRs are also the likely source of

siRNAs in *B. rapa*. For mRNA, the average expression levels of lncRNAs having 24-nt

siRNAs was higher than that of lncRNAs not having 24-nt siRNAs [36]. LncRNAs and

24-nt siRNAs are known to be involved in the increase of DNA methylation, which

agrees with reports of increased gene expression when DNA methylation overlaps with

an exonic region [53]. The detailed mechanism is not clear, but our results also agrees

that lncRNAs may be involved in this gene regulation mechanism.

The epigenetic functions of different lncRNAs have been identified in diverse

organisms [21,22], but there is limited information in *B. rapa*. LncRNAs show a close

association with DNA methylation of the genomic region encoding the lncRNA [54],

but studies have focused more on the model plant species, *A. thaliana* [11,55]. In this

study, we found a similar level of DNA methylation in regions encoding for lncRNA

compared to previously described DNA methylation in the *B. rapa* genome [36]. Levels

of DNA methylation can be positively regulated by siRNAs through the RdDM

pathway [56], and Pol IV is important for biogenesis of 24-nt siRNAs [57].

Identification of long Pol IV-dependent transcripts is difficult because these transcripts

can be rapidly processed to produce 24-nt siRNAs. There are several reports that

detected Pol IV-dependent siRNA-precursor transcripts in *A. thaliana* of different

lengths [57–59]. Most Pol IV-dependent transcript regions overlapped with Pol

IV-dependent siRNA loci, and CHH methylation depends on the production of Pol

IV-dependent transcripts [59]. However, if Pol IV-dependent transcripts are short as

30-40 nucleotides [57,58], the lncRNAs that we identified are of much longer length

and not likely to be Pol-IV dependent transcripts that act as siRNA-precursors.

Pol V transcripts accumulate at very low levels and have been difficult to

identify. However, using RNA immunoprecipitation, Pol V-dependent lncRNAs were

identified, and CHH methylation and 24-nt siRNA accumulation were shown to be

restricted to Pol V transcribed regions [60]. In this study, about 13 % of lncRNAs

overlapped with genomic regions encoding for 24-nt siRNAs, and the DNA methylation

level of those regions was higher than the average. It has been reported that the median

length of Pol V transcripts was 689 nt [60]. The length of the lncRNAs overlapping

genomic regions encoding for 24-nt siRNAs identified in this study resembles the

length of the Pol V transcript precursors.

The level of H3K27me3 plays a role in tissue specific gene expression [61].

LncRNAs are also considered to be involved in regulating histone modifications.

Binding of lncRNAs to PRC2 has been observed in human/animal [62,63], and

COLDAIR and COLDWRAP have been shown to recruit the PRC2 complex to *FLC*

during vernalization in *A. thaliana* [13,14,64,65]. EMF2B is a component of the PRC2

complex and in rice, a mutant of *emf2b* was reported to lose H3K27me3 and derepress

some of the lincRNAs, suggesting that expression of these lincRNAs is regulated by

PRC2-mediated histone methylation of the region encoding the lncRNA [66]. We found

that about 10-16% of lncRNAs overlapped with H3K27me3 regions, which is lower

than the overlap with mRNAs. The pattern of H3K27me3 in lncRNA overlapping

regions was similar to mRNAs; H3K27me3 accumulated in the body region of

transcripts, especially around the transcription start site. However, there was no

negative relationship between the presence of H3K27me3 and lncRNA expression

levels, and incRNA coding regions having H3K27me3 marks showed higher expression

levels than incRNA coding regions without H3K27me3 marks. H3K27me3 over

lncRNA coding regions was also found in different tissues of maize, which was

responsible for the regulation of tissue-specific lncRNAs expression [67].

Coincidentally, DNA methylation in intronic regions is known to influence splicing

patterns [68], which may reflect epigenetic tissue specific gene regulation involving

RdDM and H3K27me3. However, as our study focused on the analyses in a single

tissue/developmental stage, we could not identify the lncRNAs that are transcriptionally

silenced by H3K27me3, leading to the underestimation of lncRNAs with H3K27me3

marks. Further study will be required to examine the transcriptional regulation of

lncRNAs by H3K27me3.

This study revealed that a small proportion of lncRNAs in *B. rapa* are conserved

with other Brassica species. The majority of lncRNAs in *B. rapa* overlap with IRRs and

there is some overlap with DNA methylation and 24-nt siRNAs, hinting at regulation of

lncRNA expression through the RdDM pathway. Interestingly, some of the lncRNAs

that overlapped with the genomic regions having H3K27me3 marks were more highly

expressed, which was unexpected because of the known gene suppression activity of

H3K27me3. This may indicate an unknown regulatory mechanism of lncRNAs by

H3K27me3. However, the current study focuses on the genome-wide quantification of

lncRNAs, 24-nt siRNAs, DNA methylation, and H3K27me3, and further quantitative

analysis at the locus specific level may reveal a clearer relationship between these

marks. Epigenetic regulation of the genome including lncRNA, 24-nt siRNAs, DNA

methylation and H3K27me3 depend on the stage of development, and abiotic stress

conditions. Further exploration of lncRNAs under different tissues and conditions may

reveal lncRNAs with specific functions.

Acknowledgement

We are grateful to Ms. Tomoko Kusumi for her technical assistance throughout this project.

References

1. U N. Genome analysis in Brassica with special reference to the experimental formation of *B. napus* and peculiar mode of fertilization. Japanese J Bot. 1935;7: 389–452.

2. Lv H, Miyaji N, Osabe K, Akter A, Mehraj H, Shea DJ, et al. The Importance of genetic and epigenetic research in the *Brassica* vegetables in the face of climate change. In: Kole C, editor. Genomic designing of climate-smart vegetable crops. Springer; 2020. pp. 161-255.

3. Wang X, Wang H, Wang J, Sun R, Wu J, Liu S, et al. The genome of the mesopolyploid crop species *Brassica rapa*. Nat Genet. 2011;43: 1035–1040. doi:10.1038/ng.919

4. Liu S, Liu Y, Yang X, Tong C, Edwards D, Parkin IAP, et al. The *Brassica oleracea* genome reveals the asymmetrical evolution of polyploid genomes. Nat Commun. 2014;5: 3930. doi:10.1038/ncomms4930

5. Parkin IAP, Koh C, Tang H, Robinson SJ, Kagale S, Clarke WE, et al. Transcriptome and methylome profiling reveals relics of genome dominance in the

mesopolyploid *Brassica oleracea*. Genome Biol. 2014;15: R77.

doi:10.1186/gb-2014-15-6-r77

6. Yang J, Liu D, Wang X, Ji C, Cheng F, Liu B, et al. The genome sequence of

allopolyploid *Brassica juncea* and analysis of differential homoeolog gene

expression influencing selection. Nat Genet. 2016;48: 1225–1232.

doi:10.1038/ng.3657

7. Chalhoub B, Denoeud F, Liu S, Parkin IAP, Tang H, Wang X, et al. Early

allopolyploid evolution in the post-Neolithic *Brassica napus* oilseed genome.

Science. 2014;345: 950–953. doi:10.1126/science.1253435

8. Karlik E, Ari S, Gozukirmizi N. LncRNAs: genetic and epigenetic effects in plants.

Biotechnol Biotechnol Equip. 2019;33: 429–439.

doi:10.1080/13102818.2019.1581085

9. Rai MI, Alam M, Lightfoot DA, Gurha P, Afzal AJ. Classification and experimental

identification of plant long non-coding RNAs. Genomics. 2019;111: 997–1005.

doi:10.1016/j.ygeno.2018.04.014

10. Mattick JS, Rinn JL. Discovery and annotation of long noncoding RNAs. Nat Struct

Mol Biol. 2015;22: 5–7. doi:10.1038/nsmb.2942

11. Chekanova JA. Long non-coding RNAs and their functions in plants. Curr Opin

Plant Biol. 2015;27: 207–216. doi:10.1016/j.pbi.2015.08.003

12. Liu J, Wang H, Chua NH. Long noncoding RNA transcriptome of plants. Plant

Biotechnol J. 2015;13: 319–328. doi:10.1111/pbi.12336

13. Heo JB, Sung S. Vernalization-mediated epigenetic silencing by a long intronic

noncoding RNA. Science 2011;331: 76–79. doi:10.1126/science.1197349

14. Kim DH, Sung S. Vernalization-triggered intragenic chromatin loop formation by

long noncoding RNAs. Dev Cell. 2017;40: 302-312.e4.

doi:10.1016/j.devcel.2016.12.021

15. Swiezewski S, Liu F, Magusin A, Dean C. Cold-induced silencing by long antisense

transcripts of an *Arabidopsis* Polycomb target. Nature. 2009;462: 799–802.

doi:10.1038/nature08618

16. Ding J, Lu Q, Ouyang Y, Mao H, Zhang P, Yao J, et al. A long noncoding RNA

regulates photoperiod-sensitive male sterility, an essential component of hybrid rice.

Proc Natl Acad Sci U S A. 2012;109: 2654–2659. doi:10.1073/pnas.1121374109

17. Wang Y, Fan X, Lin F, He G, Terzaghi W, Zhu D, et al. *Arabidopsis* noncoding

RNA mediates control of photomorphogenesis by red light. Proc Natl Acad Sci U S

A. 2014;111: 10359–10364. doi:10.1073/pnas.1409457111

18. Fedak H, Palusinska M, Krzyczmonik K, Brzezniak L, Yatusevich R, Pietras Z, et al.

Control of seed dormancy in *Arabidopsis* by a *cis*-acting noncoding antisense

transcript. Proc Natl Acad Sci U S A. 2016;113: E7846–E7855.

doi:10.1073/pnas.1608827113

19. Seo JS, Sun HX, Park BS, Huang CH, Yeh SD, Jung C, et al. ELF18-INDUCED

LONG-NONCODING RNA associates with mediator to enhance expression of

innate immune response genes in Arabidopsis. Plant Cell. 2017;29: 1024–1038.

doi:10.1105/tpc.16.00886

20. Wu CT, Morris JR. Genes, genetics, and epigenetics: A correspondence. Science.

2001;293: 1103–1105. doi:10.1126/science.293.5532.1103

21. Wang C, Wang L, Ding Y, Lu X, Zhang G, Yang J, et al. LncRNA structural

characteristics in epigenetic regulation. Int J Mol Sci. 2017;18: 2659.

doi:10.3390/ijms18122659

22. Matzke MA, Kanno T, Matzke AJM. RNA-directed DNA methylation: The

evolution of a complex epigenetic pathway in flowering plants. Annu Rev Plant

Biol. 2015;66: 243–267. doi:10.1146/annurev-arplant-043014-114633

23. Fujimoto R, Sasaki T, Ishikawa R, Osabe K, Kawanabe T, Dennis ES. Molecular

mechanisms of epigenetic variation in plants. Int J Mol Sci. 2012;13: 9900–9922.

doi:10.3390/ijms13089900

24. Liu ZW, Zhao N, Su YN, Chen SS, He XJ. Exogenously overexpressed intronic

long noncoding RNAs activate host gene expression by affecting histone

modification in Arabidopsis. Sci Rep. 2020;10: 3094.

doi:10.1038/s41598-020-59697-7

25. Kim DH, Xi Y, Sung S. Modular function of long noncoding RNA, COLDAIR, in

the vernalization response. PLoS Genet. 2017;13: e1006939.

doi:10.1371/journal.pgen.1006939

26. Itabashi E, Osabe K, Fujimoto R, Kakizaki T. Epigenetic regulation of agronomical

traits in Brassicaceae. Plant Cell Rep. 2018;27: 87-101.

doi.org/10.1007/s00299-017-2223-z

27. Shea DJ, Nishida N, Takada S, Itabashi E, Takahashi S, Akter A, et al. Long

noncoding RNAs in *Brassica rapa* L. following vernalization. Sci Rep. 2019;9:

9302. doi:10.1038/s41598-019-45650-w

28. Wang A, Hu J, Gao C, Chen G, Wang B, Lin C, et al. Genome-wide analysis of

long non-coding RNAs unveils the regulatory roles in the heat tolerance of Chinese

cabbage (*Brassica rapa* ssp.chinensis). Sci Rep. 2019;9: 5002.

doi:10.1038/s41598-019-41428-2

29. Zhang J, Wei L, Jiang J, Mason AS, Li H, Cui C, et al. Genome-wide identification,

putative functionality and interactions between lncRNAs and miRNAs in Brassica

species. Sci Rep. 2018;8: 4960. doi:10.1038/s41598-018-23334-1

30. Huang L, Dong H, Zhou D, Li M, Liu Y, Zhang F, et al. Systematic identification of

long non-coding RNAs during pollen development and fertilization in *Brassica rapa*.

Plant J. 2018;96: 203–222. doi:10.1111/tpj.14016

31. Song X, Liu G, Huang Z, Duan W, Tan H, Li Y, et al. Temperature expression

patterns of genes and their coexpression with LncRNAs revealed by RNA-Seq in

non-heading Chinese cabbage. BMC Genomics. 2016;17: 297.

doi:10.1186/s12864-016-2625-2

32. Liu X, Hao L, Li D, Zhu L, Hu S. Long non-coding RNAs and their biological roles

in plants. Genomics, Proteomics Bioinformatics. 2015;13: 137–147.

doi:10.1016/j.gpb.2015.02.003

33. Yu X, Yang J, Li X, Liu X, Sun C, Wu F, et al. Global analysis of cis-natural

antisense transcripts and their heat-responsive nat-siRNAs in *Brassica rapa*. BMC

Plant Biol. 2013;13: 208. doi:10.1186/1471-2229-13-208

34. Kawamura K, Kawanabe T, Shimizu M, Nagano AJ, Saeki N, Okazaki K, et al.

Genetic distance of inbred lines of Chinese cabbage and its relationship to heterosis.

Plant Gene. 2016,5: 1-7. doi:10.1016/j.plgene.2015.10.003

35. Murray MG, Thompson WF. Rapid isolation of high molecular weight plant DNA.

Nucleic Acids Res. 1980;8: 4321–4326. doi:10.1093/nar/8.19.4321

36. Takahashi S, Osabe K, Fukushima N, Takuno S, Miyaji N, Shimizu M, et al.

Genome-wide characterization of DNA methylation, small RNA expression, and

histone H3 lysine nine di-methylation in *Brassica rapa* L. DNA Res. 2018;25:

511–520. doi:10.1093/dnares/dsy021

37. Akter A, Takahashi S, Deng W, Shea DJ, Itabashi E, Shimizu M, et al. The histone

modification H3 lysine 27 tri-methylation has conserved gene regulatory roles in the

triplicated genome of *Brassica rapa* L. DNA Res. 2019;26: 433–443.

doi:10.1093/dnares/dsz021

38. Yu J, Zhao M, Wang X, Tong C, Huang S, Tehrim S, et al. Bolbase: A

comprehensive genomics database for *Brassica oleracea*. BMC Genomics. 2013;14:

664. doi:10.1186/1471-2164-14-664

39. She R, Chu JSC, Wang K, Pei J, Chen N. genBlastA: Enabling BLAST to identify

homologous gene sequences. Genome Res. 2009;19: 143–149.

doi:10.1101/gr.082081.108

40. Wang R, Zou J, Meng J, Wang J. Integrative analysis of genome-wide lncRNA and

mRNA expression in newly synthesized Brassica hexaploids. Ecol Evol. 2018;8:

6034–6052. doi:10.1002/ece3.4152

41. Wang H, Chung PJ, Liu J, Jang IC, Kean MJ, Xu J, et al. Genome-wide

identification of long noncoding natural antisense transcripts and their responses to

light in *Arabidopsis*. Genome Res. 2014;24: 444–453. doi:10.1101/gr.165555.113

42. Shen E, Zhu X, Hua S, Chen H, Ye C, Zhou L, et al. Genome-wide identification of

oil biosynthesis-related long non-coding RNAs in allopolyploid *Brassica napus*.

BMC Genomics. 2018;19: 745. doi:10.1186/s12864-018-5117-8

43. Liu J, Jung C, Xu J, Wang H, Deng S, Bernad L, et al. Genome-wide analysis

uncovers regulation of long intergenic noncoding RNAs in *Arabidopsis*. Plant Cell.

2012;24: 4333–4345. doi:10.1105/tpc.112.102855

44. Joshi RK, Megha S, Basu U, Rahman MH, Kav NNV. Genome wide identification

and functional prediction of long non-coding RNAs responsive to *Sclerotinia*

sclerotiorum infection in *Brassica napus*. PLoS One. 2016;11: e0158784.

doi:10.1371/journal.pone.0158784

45. Chen R, Li M, Zhang H, Duan L, Sun X, Jiang Q, et al. Continuous salt

stress-induced long non-coding RNAs and DNA methylation patterns in soybean

roots. BMC Genomics. 2019;20: 730. doi:10.1186/s12864-019-6101-7

46. Kopp F, Mendell JT. Functional classification and experimental dissection of long

noncoding RNAs. Cell. 2018;172: 393–407. doi:10.1016/j.cell.2018.01.011

47. Kung JTY, Colognori D, Lee JT. Long noncoding RNAs: Past, present, and future.

Genetics. 2013;193: 651–669. doi:10.1534/genetics.112.146704

48. Vandivier LE, Anderson SJ, Foley SW, Gregory BD. The conservation and function

of RNA secondary structure in plants. Annu Rev Plant Biol. 2016;67: 463–488.

doi:10.1146/annurev-arplant-043015-111754

49. Lu K, Wei L, Li X, Wang Y, Wu J, Liu M, et al. Whole-genome resequencing

reveals *Brassica napus* origin and genetic loci involved in its improvement. Nat

Commun. 2019;10: 1154. doi:10.1038/s41467-019-09134-9

50. Sampath P, Murukarthick J, Izzah NK, Lee J, Choi H-I, Shirasawa K, et al.

Genome-wide comparative analysis of 20 miniature inverted-repeat transposable

element families in *Brassica rapa* and *B. oleracea*. PLoS One. 2014;9: e94499.

doi.org/10.1371/journal.pone.0094499

51. Cho J. Transposon-derived non-coding RNAs and their function in plants. Front

Plant Sci. 2018;9: 600. doi:10.3389/fpls.2018.00600

52. Wang D, Qu Z, Yang L, Zhang Q, Liu ZH, Do T, et al. Transposable elements (TEs)

contribute to stress-related long intergenic noncoding RNAs in plants. Plant J.

2017;90: 133–146. doi:10.1111/tpj.13481

53. Bewick AJ, Schmitz RJ. Gene body DNA methylation in plants. Curr Opin Plant

Biol. 2017;36: 103–110. doi:10.1016/j.pbi.2016.12.007

54. Di Ruscio A, Ebralidze AK, Benoukraf T, Amabile G, Goff LA, Terragni J, et al.

DNMT1-interacting RNAs block gene-specific DNA methylation. Nature.

2013;503: 371–376. doi:10.1038/nature12598

55. Au PCK, Dennis ES, Wang MB. Analysis of argonaute 4-associated long

non-coding RNA in *Arabidopsis thaliana* sheds novel insights into gene regulation

through RNA-directed DNA methylation. Genes (Basel). 2017;8: 198.

doi:10.3390/genes8080198

56. Yan H, Bombarely A, Xu B, Frazier TP, Wang C, Chen P, et al. siRNAs regulate

DNA methylation and interfere with gene and lncRNA expression in the

heterozygous polyploid switchgrass. Biotechnol Biofuels. 2018;11: 208.

doi:10.1186/s13068-018-1202-0

57. Zhai J, Bischof S, Wang H, Feng S, Lee TF, Teng C, et al. A one precursor one

siRNA model for Pol IV-dependent siRNA biogenesis. Cell. 2015;163: 445–455.

doi:10.1016/j.cell.2015.09.032

58. Blevins T, Podicheti R, Mishra V, Marasco M, Wang J, Rusch D, et al.

Identification of Pol IV and RDR2-dependent precursors of 24 nt siRNAs guiding

de novo DNA methylation in Arabidopsis. Elife. 2015;4: e09591.

doi:10.7554/eLife.09591

59. Li S, Vandivier LE, Tu B, Gao L, Won SY, Li S, et al. Detection of Pol

IV/RDR2-dependent transcripts at the genomic scale in *Arabidopsis* reveals features

and regulation of siRNA biogenesis. Genome Res. 2015;25: 235–245.

doi:10.1101/gr.182238.114

60. Böhmdorfer G, Sethuraman S, Jordan Rowley M, Krzyszton M, Hafiz Rothi M,

Bouzit L, et al. Long non-coding RNA produced by RNA polymerase V determines

boundaries of heterochromatin. Elife. 2016;5: e19092. doi:10.7554/eLife.19092

61. Makarevitch I, Eichten SR, Briskine R, Waters AJ, Danilevskaya ON, Meeley RB,

et al. Genomic distribution of maize facultative heterochromatin marked by

trimethylation of H3K27. Plant Cell. 2013;25: 780–793.

doi:10.1105/tpc.112.106427

62. Tu S, Yuan GC, Shao Z. The PRC2-binding long non-coding RNAs in human and

mouse genomes are associated with predictive sequence features. Sci Rep. 2017;7:

41669. doi:10.1038/srep41669

63. Gaiti F, Degnan BM, Tanurdžić M. Long non-coding regulatory RNAs in sponges

and insights into the origin of animal multicellularity. RNA Biol. 2018;15: 696–702.

doi:10.1080/15476286.2018.1460166

64. Tian Y, Zheng H, Zhang F, Wang S, Ji X, Xu C, et al. PRC2 recruitment and

H3K27me3 deposition at *FLC* require FCA binding of *COOLAIR*. Sci Adv. 2019;5:

eaau7246. doi:10.1126/sciadv.aau7246

65. Yu T, Zhu H. Long non-coding RNAs: Rising regulators of plant reproductive

development. Agronomy. 2019;9: 53. doi:10.3390/agronomy9020053

66. Johnson C, Conrad LJ, Patel R, Anderson S, Li C, Pereira A, et al. Reproductive

long intergenic noncoding RNAs exhibit male gamete specificity and polycomb

repressive complex 2-mediated repression. Plant Physiol. 2018;177: 1198–1217.

doi:10.1104/pp.17.01269

67. Li L, Eichten SR, Shimizu R, Petsch K, Yeh CT, Wu W, et al. Genome-wide

discovery and characterization of maize long non-coding RNAs. Genome Biol.

2014;15: R40. doi:10.1186/gb-2014-15-2-r40

68. Regulski M, Lu Z, Kendall J, Donoghue MTA, Reinders J, Llaca V, et al. The maize

methylome influences mRNA splice sites and reveals widespread paramutation-like

switches guided by small RNA. Genome Res. 2013;23: 1651–1662.

doi:10.1101/gr.153510.112

Supporting information

S1 Fig. Chromosomal (A01-A10) distribution of lncRNA and mRNAs. $\log_2$ FPKM

of mRNAS represented in blue (positive values) and orange (negative values), and

lncRNAs in green (positive values) and pink (negative values).

S2 Fig. DNA methylation level of each type of lncRNAs, putative mRNA, total, and

IRRs. "Total" represents the methylation level of all regions.

S3 Fig. DNA methylation of the genic region, IRRs, or regions coding for putative

mRNA and lncRNAs.

S4 Fig. Proportion of lncRNAs, putative mRNA, mRNA with (K27+) or without

(K27-) H3K27me3 on encoding regions.

S5 Fig. H3K27me3 mapped to the genic region, IRRs, or encoding regions of

putative mRNA or lncRNAs.