Interspecies variation of larval locomotion kinematics in the

genus *Drosophila* and its relation to habitat temperature

Yuji Matsuo[a], Akinao Nose[a,b] and Hiroshi Kohsaka[a*]

[a] Department of Complexity Science and Engineering, Graduate School of

Frontier Science, the University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa, 277-

8561 Chiba, Japan

[b] Department of Physics, Graduate School of Science, the University of Tokyo, 7-

3-1 Hongo, Bunkyo-ku, 133-0033 Tokyo, Japan

* For correspondence: (kohsaka@edu.k.u-tokyo.ac.jp, +81-4-7136-3922)

Abstract

Speed and trajectory of locomotion are characteristic traits of individual species.

During evolution, locomotion kinematics is likely to have been tuned for survival in the

habitats of each species. Although kinematics of locomotion is thought to be influenced by

habitats, the quantitative relation between the kinematics and environmental factors has not

been fully revealed. Here, we performed comparative analyses of larval locomotion in 11

Drosophila species. We found that larval locomotion kinematics are divergent among the

species. The diversity is not correlated to the body length but is correlated instead to the

minimum habitat temperature of the species. Phylogenetic analyses using Bayesian

inference suggest that the evolutionary rate of the kinematics is diverse among phylogenetic

trees. The results of this study imply that the kinematics of larval locomotion has diverged in

the evolutionary history of the genus *Drosophila* and evolved under the effects of the

minimum ambient temperature of habitats.

Keywords: animal locomotion, larval crawling, the genus Drosophila, kinematics, evolution

Introduction

Kinematics of animal locomotion is a critical trait enabling each species to survive

in their habitats (Alexander, 2006). Movement patterns have been sculpted during their

evolution by adaptation to their environments, and could have diverged among species

(Kappeler, 2010). Comparative analyses have identified several examples of differences in

the kinematics of locomotion within a group of related species, including insects (Theunissen

et al., 2015), reptiles (Fuller et al., 2011), birds (Stoessel & Fischer, 2012), and primates

(Thompson et al., 2018). In general, differences in locomotion among species have been

attributed to the morphological differences and/or the ecological diversity among species. An

example of morphological differences is found in three stick insect species with distinct body

proportions but living in similar feeding environments, which show different kinematics in the

legged locomotion (Theunissen et al., 2015). In contrast, as an example of ecological

diversity, two gecko species inhabiting either sandy or rocky environments have been shown

to exhibit distinct postures (Fuller et al., 2011). Whereas the interspecies divergence in

locomotion patterns can be observed in various phylogenetic branches of the animal kingdom,

quantitative comparative analyses of locomotion kinematics remain limited.

46   Flies of the genus *Drosophila* have long been used as a model to study interspecific

diversification and evolution (Markow & O'Grady, 2005). One remarkable trait diversified in

the genus *Drosophila* is the food they eat: larvae in the subgenus *Sophophora* in the genus

Drosophila eat fruit, whereas larvae in the subgenus *Drosophila* in the genus *Drosophila* feed

on vegetables (Markow, 2015; Watada et al., 2020; Watanabe et al., 2019). Some eat multiple

kinds ("generalists" including *Drosophila melanogaster*) while others have strong preferences

in food ("specialists" including *Drosophila sechellia*) (Markow, 2015; Watada et al., 2020;

Watanabe et al., 2019). These diverged species in the genus *Drosophila* offer an opportunity

to perform interspecific comparisons of locomotion. Fly larvae, or maggots, have been widely

used in the study of the kinematics of locomotion (Berrigan & Lighton, 1993; Berrigan & Pepin,

1995; Fox et al., 2006; Green et al., 1983; Ruiz-Dubreuil et al., 1996; Sokolowski, 1985;

Sokolowski et al., 1997). Among the fly species, *Drosophila melanogaster* (*Dmel*) is one of

the most examined species, especially by virtue of the availability of resources in genetics

and connectomics (Ohyama et al., 2015; Schneider-Mizell et al., 2016; Simpson & Looger,

2018; Venken et al., 2011). *Dmel* larvae locomote by a sequence of forward crawling, and

changes of crawling direction achieved by bending their bodies (Green et al., 1983). The

kinematics of the larval crawling can be adaptively changed. Natural populations of *Dmel*

larvae possess a polymorphism in the crawling behaviour: larvae of "rover" strains crawl a

large area in feeding, whereas larvae of "sitter" strains stay in a patch of food during feeding

(Sokolowski, 1980). The gene responsible for this polymorphism is mapped to the foraging

gene, which encodes a cyclic guanosine monophosphate-dependent protein kinase

(Osborne et al., 1997). Ambient temperature is another factor affecting the kinematics of

larval locomotion (Kwon et al., 2010; Liu et al., 2003; Ohyama et al., 2013; Rosenzweig et

al., 2005a). In thermotaxis behaviour in temperature gradient environments, larvae regulate

the length of crawling runs between turns and the size and direction of turns (Luo et al., 2010),

and the probability of turns is also affected by ambient temperature gradients (Klein et al.,

2015). Cellular mechanisms in larval temperature sensation have been clarified. Different

temperatures are detected by specialized sensory organs and receptor proteins (Dillon et al.,

2009). Cold temperatures are sensed by two ganglia (the dorsal organ ganglion and the

terminal organ ganglion) (Klein et al., 2015; Liu et al., 2003; Ni et al., 2016a), and warm

temperatures are detected by dTRPA1-positive neurons in the brain (Rosenzweig et al.,

2005a). Some multi-dendritic (md) neurons in the body wall detect cold temperatures, while

other md neurons are sensitive to warm temperatures (Liu et al., 2003). Compared with this

remarkable level of understanding of the regulation of crawling behaviour in *Dmel* larvae,

locomotion kinematics in the larvae of sister species in the genus *Drosophila* remains unclear.

Here we conducted an interspecies comparison of the kinematics of larval

locomotion in the genus *Drosophila*. We address two questions in this study: are locomotion

kinematics of larvae similar among *Drosophila* species?; and, if the kinematics are diverged,

are morphological characteristics and/or ecological differences related to the diversity? To

this aim, we recorded locomotion of larvae of 11 *Drosophila* species and extracted kinematic

parameters using the tracking software FIMTrack (Risse et al., 2017). Clustering analysis

with Jensen-Shannon divergence and statistical analyses show that two kinematics

parameters (bend probability and crawling speed) are diverged among the *Drosophila*

species. We found that the minimum habitat temperature of each species is related to the

kinematics rather than the size of larvae. The relationship between the kinematics and

minimum habitat temperature held at two distinct ambient temperatures: 24°C and 32°C.

Phylogenetic analyses of these kinematics, based on Bayesian inference (Hohna et al.,

2016), suggests that the rate of evolution of the kinematics is diverged among phylogenetic

branches. Among eight kinematics parameters we tested, the crawling speed at 24°C and

32°C were suggested to have correlatively evolved. Consequently, our results suggest that

the kinematics of larval locomotion in the genus *Drosophila* evolved under the effects of

minimum ambient temperature in their habitats, and diverged in the evolutionary history.

Results

Kinematic analysis of crawling and bending behaviour in *Drosophila* larvae

In this study, we analyzed the kinematics of larval locomotion. To this aim, we

recorded the locomotion of fly larvae that were crawling freely on an agarose substrate stage

on a temperature-controlled plate (Figure 1A; see Method section for details). We placed

eight to ten larvae at the centre of the stage, which was kept at 24°C, illuminated them with

infrared light, which is invisible to fly larvae and does not affect their behaviour, and recorded

larval locomotion for three minutes at five frames per second (Figure 1B). Maximum

projection of the time-series images shows traces of larval locomotion as multiple curves of

multiple larvae (Figure 1C). In an example of *Drosophila melanogaster* (Figure 1D), the

traces show smooth curved lines interconnected with angles where larvae exhibit turning

behaviour and change crawling direction, which is consistent with previous studies (Berni et

al., 2012; Gershow et al., 2012; Gomez-Marin et al., 2011; Gomez-Marin & Louis, 2014;

Green et al., 1983; Sims et al., 2019; Tastekin et al., 2015). Larvae under these conditions

predominantly exhibit forward locomotion in which muscular contraction is propagating from

the posterior to anterior segments. Characteristics of larval locomotion can be described by

two measures: the bend angle and centroid speed (Gershow et al., 2012; Gomez-Marin et

al., 2011). The bend angle measures the angle of body axis bending (Figure 1E) and the

centroid speed is the speed of the position of the larval centroid (Figure 1F). To obtain these

values for each larva at each time frame, we used the object tracking software for small

animals, FIMTrack (Risse et al., 2017). The turning behaviour can be detected by the change

in the angle of the body axis (Figure 1E) and the reduction in the centroid speed (Figure 1F),

as reported previously (Gershow et al., 2012; Gomez-Marin et al., 2011). Accordingly, we

used the bend angle and centroid speed for quantitative analysis of larval locomotion in this

study.

Drosophila melanogaster

Figure 1. Measurement of larval crawling. (A) Setup of the recording of larval crawling with

infrared light and temperature control plate. (B) An example image of multiple larvae recorded

by the setup (A). (C) The trajectories of larval locomotion of *Drosophila melanogaster*

recorded for two minutes. (D) A trajectory of single *Drosophila melanogaster* locomotion.

Locations where the larva changes its direction are labelled (i) to (iv). (E and F). Bend angle

(E) and centroid speed (F) of the larva in (D). Labels (i) to (iv) correspond to those in (D).

Classification of locomotion properties in the genus *Drosophila*

134   For interspecies comparison of *Drosophila* larval locomotion, we collected 11 fly

species in the genus *Drosophila*, whose genome sequences have been read (A. G. Clark et

al., 2007; Miller et al., 2018), and for which living individuals are available from fly stock

centres (KYORIN-Fly, Fly Stocks of Kyorin University; KYOTO Stock Center (DGRC) at the

Kyoto Institute of Technology). The 11 *Drosophila* (*D.*, hereafter) species consist of *D.*

ananassae (*Dana*)*, D. erecta* (*Dere*)*, D. mauritiana* (*Dmau*)*, D. melanogaster* (*Dmel*)*, D.*

mojavensis (*Dmoj*)*, D. persimilis* (*Dper*)*, D. pseudoobscula* (*Dpse*)*, D. sechellia* (*Dsec*)*, D.*

willistoni (*Dwil*) *and D. yakuba* (*Dyak*). We plotted the centroid speed and bend angle of freely

crawling larvae of each species (Figure 2A). In all the cases, data points accumulated around

a region where the bend angle is zero (Figure 2A), which reflects an observation that larvae

do not bend during the majority of the time (Figure 1D). We noticed that the deviations from

zero in bend angle axis are different among species. For example, data points in *Dvir* are

scattered along the bend angle axis more than those in *Dwil* (Figure 2A). To quantify

similarities in the distribution of the two-dimensional plots among the species, we calculated

Jensen-Shannon divergence, which measures the similarity of two probability distributions

(Lin, 1991). We classified the 11 species based on the similarity in the probability distribution

of the crawling speed and bend angle plots by hierarchical clustering (Figure 2B and 2C).

The classification analysis shows that four species (*Dmel, Dvir, Dmoj and Dper*) form a cluster.

These four species show scattered data points along the bend angle axis compared with the

other seven species (Figure 2A). This observation suggests that the kinematics of larvae in

the genus *Drosophila* is diverse among species.

Figure 2. Interspecific comparison of kinematics of larval crawling in the genus

Drosophila. (A) Plots of the speed of the centroid of larvae and the bend angle in the 11

Drosophila species. Each point corresponds to a datum of a single larva of a species in a

single time frame. (B) Jensen-Shannon divergences of the probability density of (A). (C)

Hierarchical clustering of the kinematics of larval locomotion of the 11 *Drosophila* species.

Definition of the bend probability and crawling speed

The clustering analysis suggests that the kinematics is divergent among species.

To interpret the diversity in terms of locomotion behaviour, we defined two indices: the bend

probability and crawling speed. The bend probability measures how often larvae bend their

body laterally. We set the minimum angle of bend as 20 degrees, which was used previously

(Risse et al., 2013), and labelled larvae that bent more than this threshold angle to the right

or left side as "Bending" (Figure 3A and 3B). This threshold allowed us to extract the

difference in the frequency of bend among the species we examined, and we found that while

Dmel larvae exhibit larger bend angles than this threshold (Figure 3A), the bend angles in

Dwil larvae are mostly less than the threshold (Figure 3B). We defined the bend probability

of each single larva as a ratio of the number of time frames labelled "Bending" to the total

frame number. To define the second index crawling speed, we labelled larvae that bent less

than the threshold as "Crawling" (Figure 3A and 3B). We defined the median of centroid

speeds of larvae that were labelled "Crawling" as the crawling speed for each single larva.

By these definitions, we calculated the bend probability and crawling speed of each larva of

the species.

Diversity in the kinematics of larval locomotion among the *Drosophila* species

We plotted the bend probability of each species (Figure 3C). A statistical analysis

shows that the bend probability of the species is diverse ($p = 6.8 \times 10^{-25}$, Kruskal-Walls test).

While *Dvir* larvae exhibited frequent bending (bend probability: 0.355 ± 0.045, n = 24), *Dwil*

larvae rarely bend (bend probability: 0.003 ± 0.004, n = 18) (Figure 3C). We also plotted the

crawling speed of the *Drosophila* species (Figure 3D). The statistical analysis shows that the

speed is also diverse among the species ($p = 9.3 \times 10^{-24}$, Kruskal-Walls test). For example,

Dwil larvae crawl faster than *Dvir* larvae do (crawling speed: 1.46 ± 0.07 mm/sec, n = 24 in

Dwil; crawling speed: 0.95 ± 0.05 mm/sec, n = 18 in *Dvir*). These analyses indicate that the

bend probability and crawling speed are differentiated in the genus *Drosophila*.

Figure 3. Bend probability and crawling speed of larval locomotion in the 11 *Drosophila* species at 24°C and their relationship to habitat temperature of each species. (A and B) Plots of the speed of the centroid of larvae and the bend angle of *D. melanogaster* (A) and *D. willistoni* (B). Each point corresponds to a datum of a single larva of a species at a single time frame. Thick horizontal lines denote the threshold between crawling and bending. (C) Bend probability of individual larvae of each species. (D) Crawling

speed of individual larvae of each species. Sample numbers in (C) and (D) are the following:

Dvir: n=24; *Dmel*: n=24; *Dmoj*: n=30; *Dper*: n=27; *Dpse*: n=19; *Dsec*: n=21; *Dwil*: n=18; *Dere*:

n=22; *Dmau*: n=26; *Dana*: n=22; *Dyak*: n=16. (E) A world map of minimum habitat

temperature. (F) Scatter plot of bend probability at 24°C against the minimum habitat

temperature Tmin. (G) Scatter plot of crawling speed at 32°C against the minimum habitat

temperature Tmin. Median ± sem is shown in F and G.

Relationship between the kinematics of larval locomotion and minimum habitat

temperature of the *Drosophila* species

Next, we tried to find factors that relate to the diversification of larval kinematics

among species. Among the factors in the morphological differences and the ecological

diversity that might be involved in the kinematics diversity (Theunissen et al., 2015) we

examined the body length of larvae as a morphological factor and habitat temperature as an

ecological factor. A previous study reported an allometric relationship between the body size

and the crawling speed in *Diptera* larvae (Berrigan & Pepin, 1995). The authors analyzed

larvae in the order *Diptera*, the average size of which spans from 3.7 mm (*Dmel*) to 15.9 mm

(*Sarcophaga bullata*). We tested whether the relationship between the body length and the

crawling speed also holds within the genus *Drosophila*, a subgroup of the order *Diptera*. The

length of the *Drosophila* larvae we used spanned from 3.49 ± 0.05 mm (*Dyak*) to 5.68 ± 0.16

mm (*Dpse*) (Supplementary Figure 1A). We found no significant relationship between the

body length and crawling speed (Supplementary Figure 1B; Pearson correlation = -0.099).

The bend probability shows a positive correlation to body length (Supplementary Figure 1C;

Pearson correlation = 0.66). However, when omitting the data of *Dvir*, which has the highest

values in both the larva length and bend probability, the correlation became weak (Pearson

correlation = 0.28) suggesting that larval length is not a significant factor for larval kinematics

variation within the genus *Drosophila*.

Habitat temperature is one of the critical factors influencing species traits (Schmidt-

Nielsen, 1997). A previous study showed that variability of habitat temperatures of the genus

Drosophila is related to the proportions of duplicated genes in the genome (Makino & Kawata,

2012). To test the possible roles of habitat temperature in the evolution of *Drosophila* larval

locomotion, we examined the relationship between habitat temperatures and the locomotion

kinematics in the genus *Drosophila*. The habitat regions of the 11 species were obtained from

the literature (Brake & Baechli, 2008; Makino & Kawata, 2012; Markow & O'Grady, 2005),

and climate temperature data were obtained from a global climate dataset, WORLDCLIM

(Hijmans et al., 2005) (Figure 3E, Supplementary Figure 2). We examined the relation

between larval kinematics (bend probability or crawling speed) and indices of habitat

temperatures (Figure 3, Supplementary Figure 2 and 3). We found that the bend probability

and crawling speed are both correlated with minimum habitat temperature, Tmin (Figure 3F

and 3G. Pearson correlation: bend probability vs Tmin = -0.81, crawling speed vs Tmin =

0.74; for the definition of Tmin, see Materials and Methods). Even after omitting extreme

values (*Dvir* in the bend probability data and *Dpse* in the crawling speed data), the correlation

remained high (Pearson correlation: the bend probability vs Tmin (without *Dvir*) = -0.69, the

crawling speed vs Tmin (without *Dpse*) = 0.78). In species inhabiting moderate environments

(where Tmin is 15 to 25°C), larvae showed low bend probability and fast crawling (Figure 3F

and 3G). On the other hand, larvae of species inhabiting cold environments (where Tmin is

0 to 10°C) exhibited frequent bending and slow crawling (Figure 3F and 3G). This observation

implies that minimum temperature in the habitat may be one of the leading factors in sculpting

the kinematics of larval locomotion in the genus *Drosophila*.

Correlation between bend probability and minimum habitat temperature among

***Drosophila* species at high ambient temperature**

So far we had analyzed larval locomotion at 24°C. Larvae of *Dmel* are known to

crawl faster at 32°C than at 25°C (Ohyama et al., 2013). So we next examined whether the

relation between the kinematics indices and minimum habitat temperature hold or not, and

how the kinematics indices are changed, at a higher ambient temperature. We performed the

same set of measurements and analysis of larval locomotion at 32°C (Figure 4 and 5). The

interspecies comparison showed that the bend probabilities at 32°C were correlated with

minimum habitat temperature as they did at 24°C (Figure 4B; Pearson correlation = -0.85).

To examine the effects of ambient temperature on the bend probability in detail, we compared

the differences of bend probability at 24°C and 32°C among the species (Figure 4C). We

found no significant relationship between the change in bend probability and the minimum

habitat temperature (Figure 4C; Pearson correlation = 0.40). Seven species showed no

significant changes in the bend probability at 32°C (*Dmoj*, *Dper*, *Dpse*, *Dsec*, *Dmau*, *Dana*

and *Dyak* in Figure 4A - 4D), while *Dvir* and *Dmel* showed a decrease in the bend probability

at 32°C, and *Dwil* and *Dere* exhibited an increase (Figure 4A - 4C, 4E and 4F). Accordingly,

we concluded that while the shift of the bend probability between the distinct temperatures is

diverse among the *Drosophila* species, the overall trend between the bend probability in

relation to minimum habitat temperature holds at the higher ambient temperature of 32°C.

Figure 4. Bend probability in larval locomotion in the 11 *Drosophila* species at 32°C.

(A) Bend probability of the 11 species at 24°C and 32°C. The data at 24°C are the same as in Figure 3C. Sample numbers at 32°C are the following: *Dvir*: n=25; *Dmel*: n=20; *Dmoj*: n=29; *Dper*: n=27; *Dpse*: n=25; *Dsec*: n=20; *Dwil*: n=21; *Dere*: n=24; *Dmau*: n=17; *Dana*: n=23; *Dyak*: n=19. (B) Scatter plot of bend probability at 32°C against the minimum habitat temperature, Tmin. (C) Scatter plot of the difference in bend probability between 32°C and at 24°C (bend probability at 32°C – bend probability at 24°C) against the minimum habitat temperature Tmin. (D-F) Examples of trajectories (left) and the bend angle (right) at 24°C

(top) and 32°C (bottom) of larval locomotion of each species (D: *Dsec*, E: *Dmel*, and F: *Dwil*).

Median ± sem is shown in B and C.

Correlation between crawling speed and minimum habitat temperature among

***Drosophila* species at high temperature**

Next, we analyzed the crawling speed at 32°C (Figure 5A). Similarly to our analysis

of the bend probability, we found that the crawling speed at 32°C was correlated with the

minimum habitat temperature among the species, as it was at 24°C (Figure 5B; Pearson

correlation = 0.67). To examine the effects of ambient temperature on the crawling speed in

detail, we compared the difference in the crawling speed at 24°C to 32°C among the species

(Figure 5C). In contrast to the case in the bend probability, shifts in the crawling speed among

the species were common: ten species (all the species but *Dper*) showed a significant

increase in crawling speed at 32°C (Figure 5A). *Dper* also exhibited an increase in the

crawling speed but it was not statistically significant (Figure 5A). Accordingly, at 32°C, larvae

of all the species we tested increased their speed of crawling, which might reflect a general

demand to avoid malfunctions in metabolic reactions and dehydration of the body. We also

tested the relationship between the change in the speed at 32°C and the minimum habitat

temperature for each species (Figure 5C). We found no significant correlation between the

speed change and the habitat temperature (Figure 5C; Pearson correlation = 0.07), which

also implies that the speeding-up at high temperature is a general requirement among the

species for the larval survival. Accordingly, whereas the details of shifts are different (diverse

shift in the bend probability and common shift in the crawling speed among the species), the

overall relationship between the kinematic indices and minimum habitat temperature also

holds at the higher ambient temperature of 32°C.

Figure 5. Crawling speed of larval locomotion in the 11 *Drosophila* species at 32°C. (A) Crawling speed of the 11 species at 24°C and 32°C. The data at 24°C are the same as in Figure 3D. Sample numbers at 32°C are the following: *Dvir*: n=25; *Dmel*: n=20; *Dmoj*: n=29; *Dper*: n=27; *Dpse*: n=25; *Dsec*: n=20; *Dwil*: n=21; *Dere*: n=24; *Dmau*: n=17; *Dana*: n=23; *Dyak*: n=19. (B) Scatter plot of the crawling speed at 32°C against the minimum habitat temperature Tmin. (C) Scatter plot of the difference of the crawling speed between at 32°C and at 24°C (crawling speed at 32°C – crawling speed at 24°C) against the minimum habitat temperature Tmin. Median ± sem is shown in B and C.

A kinematic trend of larval locomotion in the genus Drosophila

To see any gross trend in the shift of larval kinematics in the different species at distinct ambient temperatures, we plotted the kinematics of individual larvae of all the species (Figure 6). At the ambient temperature of 24°C, a trend of kinematics was observed in which the data points were distributed from low crawling speed / high bend probability (the top-left

corner in the plot) to high crawling speed / low bend probability (the bottom-right corner in

the plot) as the minimum habitat temperature increased (Figure 6A). Intriguingly, the

kinematics indices between distinct species were not segregated but rather overlapped. This

continuum property can also be observed in the kinematics data at the ambient temperature

of 32°C (Figure 6B). According to the modern phylogenetic concept (Markow & O'Grady,

2005), the subgenera *Sophophora* and *Drosophila* (both belonging to the genus *Drosophila*)

are ancestral and descendant, respectively. The 11 genus *Drosophila* species in our study

consisted of nine ancestral subgenus *Sophophora* species (all but *Dvir* and *Dmoj*) and two

descendant subgenus *Drosophila* species (*Dvir* and *Dmoj*). In our plots of kinematics,

species in the descendant subgenera *Drosophila* (*Dvir* and *Dmoj*) were located at the top-

left corner in the continuum (Figure 6). This observation implies that the kinematics indices

of larvae in ancestral *Drosophila* species took values in this continuum and kinematics

indices in descendant species have gradually diverged along this continuum by adaptation

to minimum habitat temperature during evolution.

Figure 6. Scatter plot of kinematics in the 11 species at 24°C and 32°C. (A and B) Two

dimensional plots of the crawling speed and bend probability. Each point corresponds to data

of a single larva of the species. The colour of markers denotes minimum habitat temperature

(Tmin) as shown in the legend. A: 24°C, B: 32°C.

No correlation between crawling speed and habitat temperature among *Drosophila*

species at an extreme ambient temperature of 40°C

Temperatures of 24°C and 32°C are within the range of natural habitat temperatures

of most of the species we tested (Supplementary Figure 2). We next examined larval crawling

at an extreme temperature and investigated the relation between the kinematics indices and

minimum habitat temperature. Too high a temperature can be noxious for many animals,

including *Drosophila* larvae. When *D. melanogaster* larvae are stimulated with a probe

heated to 42°C, they exhibit a stereotyped rolling motion as an escape behaviour (Dason et

al., 2019; Petersen et al., 2018; Tracey et al., 2003). At below 40°C, on the other hand, larvae

do not exhibit rolling behaviour (Petersen et al., 2018), which allows us to study crawling

kinematics in this semi-noxious extreme environment. We recorded larval locomotion of the

Drosophila species at the ambient temperature of 40°C, which corresponds to the highest

edge of maximum habitat temperature histograms for many of the species (Supplementary

Figure 2). We measured the bend probability and crawling speed of the 11 *Drosophila*

species (Supplementary Figure 4 and 5). At the ambient temperature of 40°C, neither the

bend probability nor crawling speed was correlated with minimum habitat temperature

(Figure 7A and 7B; Pearson correlation: bend probability at 40°C vs Tmin = 0.17, crawling

speed at 40°C vs Tmin = -0.17). Since 40°C is close to the maximum habitat temperature

rather than the minimum habitat temperature, we analyzed the relation between the

kinematics indices and maximum habitat temperature instead of minimum habitat

temperature. However, neither the bend probability nor crawling speed was correlated with

maximum habitat temperature (Pearson correlation: bend probability at 40°C vs maximum

habitat temperature = -0.12, crawling speed vs maximum habitat temperature = -0.10).

Accordingly, the relation between the kinematic indices and habitat temperature does not

hold at the extreme ambient temperature of 40°C. This observation suggests that the

influence of minimum habitat temperature on the evolution of the locomotion kinematics is restricted within a specific range of ambient temperatures. To examine the effects of the extreme ambient temperature on the kinematics indices in detail, we examined the shift in the bend probability at the distinct temperatures of 40°C and 32°C among the species (Supplementary Figure 4). In seven species, bend probability increased at 40°C. Three species that inhabit moderate temperature areas (*Dvir*, *Dper* and *Dmoj*) and one species that lives on an isolated island (*Dsec*) showed little change in the bend probability, which may reflect a distinct strategy in evolution to cope with the semi-noxious environment. We also examined the shift in the crawling speed at the distinct temperatures of 40°C and 32°C among the species (Supplementary Figure 5). We found the crawling speed was reduced at 40°C in all the species (Supplementary Figure 5), which might be a common adaptation of the *Drosophila* larval locomotion at the extreme ambient temperature and/or due to an abnormal physiological reaction at the semi-noxious temperature. Intriguingly, in some species, backward crawling, which seldom occurs at 32°C, can be observed at 40°C (Figure 7C). Consequently, these observations show that at the extreme ambient temperature of 40°C, the relation between the kinematics indices and habitat temperature does not hold and larvae exhibit common (increase in the crawling speed) and diverse (changes in the bend probability and generation of backward crawling) shifts in the locomotion kinematics among the *Drosophila* species.

Figure 7. Larval behaviour in the 11 species at 40°C. (A) Scatter plot of the bend

probability at 40°C against the minimum habitat temperature Tmin. (B) Scatter plot of the

crawling speed at 40°C against the minimum habitat temperature Tmin. Median ± sem is

shown in (A) and (B). Sample numbers in (A) and (B) are the following: *Dvir*: n=20; *Dmel*:

n=28; *Dmoj*: n=23; *Dper*: n=21; *Dpse*: n=16; *Dsec*: n=27; *Dwil*: n=27; *Dere*: n=22; *Dmau*:

n=26; *Dana*: n=23; *Dyak*: n=20. (C) Backward crawling probability at 40°C of each species.

Sample numbers in (C) are the following: *Dvir*: n=20; *Dmel*: n=28; *Dmoj*: n=23; *Dper*: n=22;

Dpse: n=14; *Dsec*: n=27; *Dwil*: n=27; *Dere*: n=23; *Dmau*: n=26; *Dana*: n=23; *Dyak*: n=21.

Colouring of the species follows that in Figure 6A.

Phylogenetic analyses of interspecies differences in kinematics

The kinematics measurements of 11 *Drosophila* species at three temperatures have

demonstrated interspecies differentiation in the crawling kinematics. To obtain a hint on how

these kinematics parameters have developed during the evolutionary history of the genus

Drosophila, we conducted Bayesian phylogenetic analyses using RevBayes v. 1.0.7 (Hohna

et al., 2016; Höhna et al., 2017). At first, we inferred phylogenetic trees of the 11 *Drosophila*

species based on eight nuclear genes used previously (Turelli et al., 2018) (See Methods for

detail). Then by using our kinematics dataset, we estimated the following three factors in the

phylogenetic tree by Bayesian inference: the rate of evolution at each branch of the

phylogenetic trees, relative rates of evolution among the kinematic parameters, and the

correlation between the kinematic parameter evolutions (Kalay et al., 2020). To perform the

inference, we constructed a data matrix of eight parameters (the bend probability at 24, 32

and 40°C, the crawling speed at 24, 32 and 40°C, backward crawling probability at 40°C and

the body length), for each of the 11 *Drosophila* species. We assumed that these parameters

evolve under a multivariate Brownian-motion model (Huelsenbeck & Bruce, 2003; Kalay et

al., 2020; Lartillot & Poujol, 2011) (See Methods for detail). We estimated the three factors

(the evolution rates at phylogenetic trees, relative evolution rates among the kinematic

parameters, and the correlation between the parameter evolutions) by running a Markov

Chain Monte Carlo simulation.

The phylogenetic analyses indicated that the rates of evolution in the kinematics are
highly diverse over branches in the phylogenetic tree (Figure 8A). In addition, the
evolutionary rates are distinct among the eight kinematics parameters (Figure 8B). The bend
probability at 24°C and the probability of backward crawling at 40°C have a relatively high
evolution rate, which might reflect large diversification of these two parameters among the
species (Figure 6A and 7C).

Some kinematics traits might have evolved cooperatively. To test this possibility, we
calculated the correlation of the evolutionary changes between the kinematic parameters. In
the correlation analyses of all pairs of the parameters, the crawling speed at 24°C and the
crawling speed at 32°C were the most correlated (Figure 8C). On the other hand, the
evolutions of the bend probability between at 24°C and 32°C were less correlated. This
observation is consistent with the notion that while changes in the bend probability from 24°C
to 32°C among the species are diverse (Figure 4C), all the species show an increase in the
crawling speed by the temperature shift (Figure 5C). Consequently, the phylogenetic
analyses imply that the kinematics indices of larval locomotion have evolved differently in
distinct branches of a phylogenetic tree with keeping a correlation between specific
locomotion traits such as crawling speed at distinct temperatures.

Figure 8. Phylogenetic analyses of larval locomotion in the genus *Drosophila*. (A)

Phylogenetic tree estimated from Bayesian inference. The topology is inferred by coding

sequences of eight genes of the 11 *Drosophila* species. As a prior distribution for the

Bayesian inference, an uncorrelated exponential (UCED) relaxed-clock model was used.

Colour and a value of each branch denote the relative rate of the evolution of the eight

kinematic traits. (B) Relative evolution rates of the eight kinematics parameters estimated by

the Bayesian inference. (C) Correlation between the evolution of eight kinematics parameters.

Abbreviations in (B) and (C) are the following: Spd24: crawling speed at 24°C, Spd32:

crawling speed at 32°C, Spd40: crawling speed at 40°C, Bend24: bend probability at 24°C,

Bend32: bend probability at 32°C, Bend40: bend probability at 40°C, Bwd: probability of

backward crawling, Length: axial body length of larvae.

Discussion

In this work, we investigated interspecies differences in larval locomotion in the

genus *Drosophila*. We used the bend probability and crawling speed as measures to examine

larval locomotion. Despite the similar appearance of larval bodies in the different species, the

kinematics of larval locomotion is diverged (Figure 2). The body length is not a leading factor

for the diversity of kinematics (Supplementary Figure 1). In contrast, minimum habitat

temperature correlates with both the bend probability and crawling speed at both 24°C and

32°C ambient temperature (Figure 9), which implies the kinematics indices are adapted to

ambient temperature in evolution. Phylogenetic analysis by Bayesian inference suggests that

the rates of evolution are divergent among the branches of the phylogenetic tree (Figure 8A).

Among the kinematic parameters, the bend probability at the ambient temperature of 24°C

and the backward crawling probability at 40°C have relatively higher evolution rates than the

others and the evolution of the crawling speed at 24°C and 32°C are correlated (Figure 8B

and 8C). Regarding the questions raised in the Introduction section, our results suggest the

following: (1) Locomotion kinematics of larvae is divergent among *Drosophila* species. (2)

The minimum habitat temperature is more related to the kinematics indices (the bend

probability and crawling speed) than the body length (Figure 9).

Figure 9. Summary of this study. Disks and triangles denote kinematics of each species at

24°C and 32°C, respectively. Colour of markers indicates distinct minimum habitat

temperatures as shown in Figure 6A.

The correlation between habitat temperature and the kinematics raises a question

about the underlying mechanisms of adaptive change in kinematics to the minimum habitat

temperature. We presume that the adaptive change was likely to be driven by two factors:

the pupation positioning and competition over food. Regarding the pupation behaviour, a

previous study showed a correlation between path lengths of crawling and heights of

pupation location in *D. melanogaster* (Sokolowski, 1985). Since pupae are immobile,

adequate selection of pupation position is vital for their survival. Moisture is one of the key

factors that affects the selection of pupation site, since dry environments can cause pupal

desiccation, while soaked media might lead to the drowning of pupae (Sokolowski, 1985;

Sokolowski et al., 1984). It is reasonable to assume that habitat temperature is a leading

factor in determining the moisture levels of microenvironments of *Drosophila* larvae.

Accordingly, divergent temperatures in the habitats of flies leads to variability in moisture

levels, the ambient moisture affects the pupation position (Sokolowski et al., 1984), and the

pupation position is related to the path length of crawling (Sokolowski, 1985). These three

links might underlie the relationship between the larval locomotion kinematics and minimum

habitat temperature of each species.

The second possible mechanism underlying the relation between minimum habitat

temperature and locomotion kinematics is related to competition over food. It is suggested

that larval feeding is competitive between individuals (Rohlfs & Hoffmeister, 2003). There is

a positive relation between fitness and density (called Allee effects) in *D. melanogaster* larvae

(Wertheim et al., 2002) and so female flies aggregate eggs when laying them (Rohlfs &

Hoffmeister, 2003). Consequently, the larval population tends to be overcrowded (Burnet et

al., 1977) and larvae compete over the limited resources to meet their need to consume

sufficient amount of nutrient within limited larval periods (Nicholson, 1954). If the available

food is limited and larval density is high, larvae need to crawl further (Sokolowski et al., 1997),

which can be an evolutionary driving force to increase the crawling speed. In contrast, if there

is plenty of food in the habitat, the crawling speed remains unchanged or decreases during

evolution, because crawling behaviour is energetically costly (Berrigan & Lighton, 1993).

Accordingly, feeding conditions, especially the choice of which foods to eat, can affect

kinematics of larval locomotion in evolutionary processes so that the divergence of food to

eat can lead to the divergence in locomotion kinematics. In regard to this point, food for larvae

of the genus *Drosophila* is divergent and related to their living environments (Markow, 2015;

Watada et al., 2020; Watanabe et al., 2019). So, the nutritional conditions (nutrient balances

and fermentation etc.) and physical properties (hardness and wetness etc.) of food to eat

vary among the *Drosophila* species, which can drive evolutionary divergence in the

kinematics of larval locomotion. Then, the growth of plants, including fruits and vegetables,

for larvae is strongly affected by habitat temperature. Consequently, divergence in minimum

habitat temperature may affect the locomotion kinematics during evolution through

divergence in the foods that larvae feed on, and divergence in requisite feeding behaviour

for larval growth. Comprehensive kinematics studies of other *Drosophila* species and

quantitative analyses of microenvironments of wild larvae in nature will give us insights on

the relationships between ambient temperature and diverse larval locomotion kinematics.

How can we approach the neural circuit mechanisms underlying the interspecies

divergence in larval locomotion? Circuit mechanisms in larval locomotion have been

examined intensively in *Drosophila melanogaster*. Recent connectomics studies have

identified several key neurons for larval locomotion in *Drosophila melanogaster* (Carreira-

Rosario et al., 2018; M. Q. Clark et al., 2018; Fushiki et al., 2016; Heckscher et al., 2015;

Hughes & Thomas, 2007; Kohsaka et al., 2014, 2017, 2019; Schneider-Mizell et al., 2016;

Tastekin et al., 2015; Zwart et al., 2016). Regarding bending, the thoracic neuromere was

shown to be important in bending in chemotaxis (Tastekin et al., 2015). Especially, the

commissural connection is crucial for bending (Berni, 2015). In addition, a signal from the

chordotonal sensory organ is also required to generate bending (Ainsley et al., 2003).

Regarding the crawling speed, a group of inhibitory premotor neurons were identified to be

critical (Kohsaka et al., 2014). Proprioceptive feedback and neuromodulation are both

important for the normal crawling speed (Fox et al., 2006; Hughes & Thomas, 2007).

Furthermore, kinematics of the larval locomotion have been measured and investigated

quantitatively in detail (Gepner et al., 2015; Gershow et al., 2012; Gomez-Marin et al., 2011;

Gomez-Marin & Louis, 2014; Heckscher et al., 2012; Hernandez-Nunez et al., 2015; Lahiri

et al., 2011; Luo et al., 2010) and analysed by mathematical modelling (Gjorgjieva et al.,

2013; Loveless et al., 2019; Loveless & Webb, 2018; Pehlevan et al., 2016; Sun et al., 2020).

Regarding temperature sensing, neuronal and molecular mechanisms on temperature-

guided behaviour have been clarified (Dillon et al., 2009; Klein et al., 2015; Kwon et al., 2008,

2010; Liu et al., 2003; Ni et al., 2016b; Rosenzweig et al., 2005b, 2008; Shen et al., 2011).

These extensive findings on the cellular and molecular mechanisms on larval crawling in

Drosophila melanogaster will be an ideal starting point to investigate the evolution of larval

behaviour in the genus *Drosophila*. For example, differences in commissure fibre tracts in

the central nervous system (Berni, 2015) among the species might underlie the divergence

in the bend probability. Interspecies comparison of a group of interneurons PMSIs (*period-*

positive median segmental interneurons), which are involved in the crawling speed (Kohsaka

et al., 2014), would reveal neural mechanisms on the evolutionary diversification in the

crawling speed. Comparative analyses of neural network architectures and gene expression

among the *Drosophila* species will relate the evolution of the nervous system to the adaptive

diversification in larval behaviour.

Materials and Methods

***Drosophila* strains**

All animals were raised on standard cornmeal-based food. We used the following

fly stocks: *Drosophila ananassae* (k-s01), *Drosophila erecta* (k-s02), *Drosophila yakuba* (k-

s03), *Drosophila melanogaster* (k-s04), *Drosophila sechellia* (k-s10), *Drosophila persimilis*

(k-s11), *Drosophila pseudoobscura* (k-s12), *Drosophila willistoni* (k-s13), *Drosophila virilis* (k-

s14) and *Drosophila mojavensis* (k-s15) from KYORIN-Fly, Fly Stocks of Kyorin University

and *Drosophila mauritiana* (#900020) from KYOTO Stock Center (DGRC) at the Kyoto

Institute of Technology.

Recording larval locomotion

Third-instar wandering larvae of each strain were picked up and gently washed in

deionised water. Residual food on the larvae was brushed off with a paintbrush. An agarose

stage (size: 9cm x 9cm x 5mm; 1.5% agarose; RIKAKEN STAR agarose powder #RSV-

AGRP-100G) was placed on a temperature controlled plate (Cool Plate, AS ONE Corporation,

Japan) and the temperature of the surface was kept at 24±1°C, 32±1°C or 40±1°C. Eight to

ten larvae were placed on the agarose arena with a paintbrush. The larvae were illuminated

with infrared light, which larvae cannot see (LDQ-150IR2-850, wavelength = 850 nm, CCS

Inc., Japan) and recorded at five frames/sec with a CCD camera (CGE-B013-U, MIGHTEX,

Canada; resolution: 1280 pixels x 960 pixels) and an infrared filter (775LP filter, Omega

Optical, USA), and saved as a series of bitmap files. In our setting, the scale of image is 0.13

mm/pixel. For each of 11 species and at each of the three ambient temperatures, we repeated

the measurement three times.

Larva tracking

We obtained a time series of bitmap image files by the procedure described above.

The bitmap files were converted to tiff format by Fiji (https://imagej.net/fiji). In cases where

larvae accumulated at the first frame, we removed the first several images until the larvae

dispersed enough to be identified individually. The x and y coordinates of the centroid and

the bend angle (Figure 1E) of individual larvae were obtained by using the FIMTrack software

(Risse et al., 2017). We used tracking data of single larvae that were continuously tracked

for more than 300 frames (which corresponds to one minute). In case a larva collided with

another larva in the middle of the recording and the larva was traced differently before and

after the collision with distinct labels, we treated the two traces as two larvae. The coordinates

and bend angles were smoothed and plotted by Python 3.7.

Clustering analysis of kinematics

For clustering analysis of the plots in Figure 2A, we made a probability distribution.

We discretised the centroid speed axis by an interval of 0.1 mm/sec between 0 and 4

mm/sec and the bend angle axis by an interval of two degrees between 0 and 140 degrees,

then created 2800 bins in total (40 bins in the speed axis and 70 bins in the bend angle

axis). For each scatter plot, we counted the number of points in each bin and obtained

probability density by normalisation with the total number of the points. We calculated the

Kullback-Leibler divergence (KL) of probability distributions of two species p(bin) and q(bin)

as:

$$KL(p, q) = \sum_{bin=1}^{2800} p(bin) * \log \frac{p(bin)}{q(bin)}.$$

Since KL(p,q) is an asymmetric measure $(KL(p, q) \neq KL(q, p))$, we calculated the Jensen-

Shannon divergence JS, which is a symmetric alternative to the Kullback-Leibler

divergence:

$$JS(p, q) = \frac{1}{2} \{KL(p, q) + KL(q, p)\} = JS(q, p).$$

Based on the Jensen-Shannon divergence, we conducted the hierarchical clustering using

Python 3.7 and the SciPy library.

Habitat temperature of the *Drosophila* species

Global climate data were obtained from the WorldClim data website

(https://www.worldclim.org/data/worldclim21.html). We used minimum, mean and maximum

temperature in the world at a spatial resolution of five minutes (9.25 km x 9.25 km). Habitat

regions of each *Drosophila* species were obtained from the literature (Brake & Baechli, 2008;

Makino & Kawata, 2012; Markow & O'Grady, 2005) and the DrosWLD-Species database

(https://bioinfo.museum.hokudai.ac.jp/db/modules/stdb/index.php?action=tbl&tbl_id=21)

and traced them on the world map (2160 pixel x 4320 pixel) obtained from the WorldClim by

using Adobe Illustrator software (Adobe Inc., USA). The maps in Supplementary Figure 2

were generated by Adobe Illustrator and the statistics of the temperature within each species'

habitat were calculated by Python 3.7.

We examined the relation between the larval kinematics (the bend probability or

crawling speed) and indices of habitat temperatures (Figure 3, Supplementary Figure 2 and

3). We first used three indices of habitat temperatures: the mode (or the most frequent value)

of maximum, minimum and average temperature indices within the habitats. We found no

significant relationship between the crawling speed and these three temperature indices

(Pearson correlation: crawling speed vs average temperature = 0.43, vs minimum

temperature = 0.35, vs maximum temperature = 0.19). Although there are significant

relationships between the bend probability and the mode of minimum and maximum

temperature (Pearson correlation: bend probability vs average temperature = -0.82, vs

minimum temperature = -0.80, vs maximum temperature = -0.41), when omitting the data of

Dvir, which had the highest values in bend probability and the lowest values in the

temperature, the correlation became weak (Pearson correlation: bend probability vs average

temperature = -0.40, vs minimum temperature = -0.43). These results collectively suggest

that the mode values of maximum, minimum and average habitat temperatures are not

significant factors for larval kinematics variation within the genus *Drosophila*. In the minimum

habitat temperature data of *Dvir* and *Dpse*, we noticed that the mode temperatures were

below 0°C, which cannot be a representative habitat temperature for them (Supplementary

Figure 2). To test a milder and more representative index, we used the warmest peaks in the

minimum habitat temperature histogram ("Tmin" in Supplementary Figure 2). Considering

uneven distribution of flies within the region demarcated in Supplementary Figure 2, the

temperature of the warmest peak in the minimum habitat temperature may represent the

coldest temperature the majority of the population of each species experiences over many

years.

Statistical analysis

Statistical calculation was conducted by Python 3.7 in the following analyses:

Kruskal-Walls test in Figure 3C and 3D, Pearson correlation in Figure 3F, 3G, 4B, 4C, 5B and

5C, Supplementary Figure 1B, 1C, 3A, 3B, 4B and 5B, and Mann-Whitney U-test in Figure

4A and 5A and Supplementary Figure 4A and 5A.

Phylogenetic analyses

We adopted Bayesian statistics for phylogenetic analyses. In all the phylogenetic

analyses, we used the software package, RevBayes v.1.0.7 (Hohna et al., 2016).

We estimated the *Drosophila* phylogenetic trees (Figure 8A) by using eight nuclear

loci from Turelli et al. (2018): *aldolase*, *bicoid*, *enolase*, *esc*, *transaldolase*, *white*, *wingless*,

and *yellow*. We used *Scaptodrosophila lebanonensis* (*S. lebanonensis*) as an outgroup.

Coding sequences of these genes for all the species except for *D. mauritiana* and *S.*

lebanonensis were obtained from Kalay et al., (2020). Coding sequences of the genes of *D.*

mauritiana and *S. lebanonensis* were obtained from the NCBI web site

(https://www.ncbi.nlm.nih.gov/). The genes were aligned with MAFFT version 7 (Katoh &

Standley, 2013). We selected the eight genes out of twenty genes described in Turelli et al.,

(2018) because homologous genes of the other twelve genes were not identified in the

genome assembly of *D. mauritiana*. Based on the eight coding sequences of the twelve

species, we performed chronogram analyses described in Turelli et al. (2018) and obtained

the phylogenetic tree in Figure 8A, where the root is the branch of *S. lebanonensis* and the

length of each branch indicates relative time in the evolution. We repeated the phylogeny

inference four times to obtain four trees to check the robustness of this inference in the

following analysis.

Based on the phylogenetic tree obtained above, we analyzed the evolution of

kinematics parameters. We used eight parameters (bend probability at 24, 32 and 40°C,

crawling speed at 24, 32 and 40°C, backward crawling probability at 40°C, and body length),

for each of the 11 *Drosophila* species. In this analysis, while the topology of the phylogenetic

tree is unchanged, the rates of evolution of the kinematic parameters and the rates of

evolution of the branches of the phylogenetic tree were estimated by Bayesian inference. We

assumed the kinematics parameters evolve under a multivariate Brownian-motion model

(Huelsenbeck & Bruce, 2003; Lartillot & Poujol, 2011). This model consists of two

components: the relative rates of evolution among the kinematics parameters and the

correlation between each pair of the kinematics parameters (Kalay et al., 2020). The

Brownian model assumes that the changes in the kinematics parameters during evolution

are additive, which has two consequences: 1) parameters can become negative; 2) evolution

rates can depend on the scale of the kinematics parameters. To cope with these properties,

we adopted log-transformation to the kinematics parameters as described previously (Kalay

et al., 2020). This transformation guarantees that the original kinematics parameters (which

are exponentials of the log-transformation) remain positive and controls the size issue by

transforming additive changes to multiplicative ones (Kalay et al., 2020). For prior models for

the evolution rates of the branches, we specified an uncorrelated exponential (UCED)

relaxed-clock model, where the rate of each branch is drawn independently from an

exponential distribution. We tested three other models (an uncorrelated gamma (UCG)

relaxed-clock model, where the rate of each branch is drawn independently from a gamma

distribution, an uncorrelated lognormal (UCLN) relaxed-clock model, where the rate of each

branch is drawn independently from a lognormal distribution, and an autocorrelated

lognormal (ACLN) relaxed-clock model), but since the estimated sample sizes, which are

measures to check the quality of Markov-Chain Monte-Carlo (MCMC) simulation, were less

than 500 in our calculation, we did not use these models in our analyses. We ran the MCMC

simulation of UCED with each of the four phylogenetic trees described in the previous

paragraph.

Acknowledgement

We thank KYORIN-Fly, Fly Stocks of Kyorin University and KYOTO Stock Center (DGRC) at

the Kyoto Institute of Technology for providing fly lines, Dr. Takashi Makino for advice about

climate data in the world. We thank Ms. Shibahara for tracing habitats of the 11 species. We

thank Dr. Masayoshi Watada for critical comments on the paper. This work was supported

by MEXT/JSPS KAKENHI grants (22115002, 15H04255, 221S0003 to A.N. and 26430004,

17K07042, 217D0344 to H.K.).

Competing interests

686   We have no conflicts of interest with respect to the work.

References

692   Ainsley, J., Pettus, J., Bosenko, D., Gerstein, C., Zinkevich, N., Anderson, M., Adams, C.,
693   Welsh, M., & Johnson, W. (2003). Enhanced locomotion caused by loss of the Drosophila
694   DEG/ENaC protein Pickpocket1. *Current Biology*, *13*, 1557–1563.
695   https://doi.org/10.1016/S

Alexander, R. M. (2006). *Principles of Animal Locomotion*. Princeton University Press.

Berni, J. (2015). Genetic dissection of a regionally differentiated network for exploratory
698   behavior in drosophila larvae. *Current Biology*, *25*(10), 1319–1326.
699   https://doi.org/10.1016/j.cub.2015.03.023

Berni, J., Pulver, S. R., Griffith, L. C., & Bate, M. (2012). Autonomous circuitry for substrate
701   exploration in freely moving drosophila larvae. *Current Biology*, *22*(20), 1861–1870.
702   https://doi.org/10.1016/j.cub.2012.07.048

Berrigan, D., & Lighton, J. R. (1993). Bioenergetic and kinematic consequences of limblessness
704   in larval Diptera. *The Journal of Experimental Biology*, *179*, 245–259.

Berrigan, D., & Pepin, D. J. (1995). How maggots move: Allometry and kinematics of crawling in
706   larval Diptera. *Journal of Insect Physiology*, *41*(4), 329–337. https://doi.org/10.1016/0022-
707   1910(94)00113-U

Brake, I., & Baechli, G. (2008). *Drosophilidae (Diptera)*. Apollo Books.

Burnet, B., Sewell, D., & Bos, M. (1977). Genetic analysis of larval feeding behaviour in
710   Drosophila melanogaster: II. Growth relations and competition between selected lines.
711   *Genetical Research*, *30*(2), 149–161. https://doi.org/10.1017/S0016672300017559

Carreira-Rosario, A., Zarin, A. A., Clark, M. Q., Manning, L., Fetter, R. D., Cardona, A., & Doe,
713   C. Q. (2018). MDN brain descending neurons coordinately activate backward and inhibit
714   forward locomotion. *ELife*, *7*, 1–28. https://doi.org/10.7554/eLife.38554

Clark, A. G., Eisen, M. B., Smith, D. R., Bergman, C. M., Oliver, B., Markow, T. A., Kaufman, T.
716   C., Kellis, M., Gelbart, W., Iyer, V. N., Pollard, D. A., Sackton, T. B., Larracuente, A. M.,
717   Singh, N. D., Abad, J. P., Abt, D. N., Adryan, B., Aguade, M., Akashi, H., … MacCallum, I.
718   (2007). Evolution of genes and genomes on the Drosophila phylogeny. *Nature*, *450*(7167),
719   203–218. https://doi.org/10.1038/nature06341

Clark, M. Q., Zarin, A. A., Carreira-Rosario, A., & Doe, C. Q. (2018). Neural circuits driving larval
721   locomotion in Drosophila. *Neural Development*, *13*(1), 1–10.
722   https://doi.org/10.1186/s13064-018-0103-z

Dason, J. S., Cheung, A., Anreiter, I., Montemurri, V. A., Allen, A. M., & Sokolowski, M. B.
724   (2019).   Drosophila melanogaster foraging regulates a nociceptive-like escape behavior
725   through a developmentally plastic sensory circuit . *Proceedings of the National Academy
726   of Sciences*, 201820840. https://doi.org/10.1073/pnas.1820840116

Dillon, M. E., Wang, G., Garrity, P. A., & Huey, R. B. (2009). Thermal preference in Drosophila.
728   *Journal of Thermal Biology*, *34*(3), 109–119. https://doi.org/10.1016/j.jtherbio.2008.11.007

Fox, L. E., Soll, D. R., & Wu, C. F. (2006). Coordination and modulation of locomotion pattern

generators in Drosophila larvae: Effects of altered biogenic amine levels by the tyramine β hydroxlyase mutation. *Journal of Neuroscience*, *26*(5), 1486–1498. https://doi.org/10.1523/JNEUROSCI.4749-05.2006

Fuller, P. O., Higham, T. E., & Clark, A. J. (2011). Posture, speed, and habitat structure: Three-dimensional hindlimb kinematics of two species of padless geckos. *Zoology*, *114*(2), 104–112. https://doi.org/10.1016/j.zool.2010.11.003

Fushiki, A., Zwart, M. F., Kohsaka, H., Fetter, R. D., Cardona, A., & Nose, A. (2016). A circuit mechanism for the propagation of waves of muscle contraction in Drosophila. *ELife*, *5*, 1–23. https://doi.org/10.7554/elife.13253

Gepner, R., Skanata, M. M., Bernat, N. M., Kaplow, M., & Gershow, M. (2015). Computations underlying Drosophila photo- taxis, odor-taxis, and multi-sensory integration. *ELife*, *4*(MAY), 1–21. https://doi.org/10.7554/eLife.06229

Gershow, M., Berck, M., Mathew, D., Luo, L., Kane, E. A., Carlson, J. R., & Samuel, A. D. T. (2012). Controlling airborne cues to study small animal navigation. *Nature Methods*, *9*(3), 290–296. https://doi.org/10.1038/nmeth.1853

Gjorgjieva, J., Berni, J., Evers, J. F., & Eglen, S. J. (2013). Neural circuits for peristaltic wave propagation in crawling Drosophila larvae: analysis and modeling. *Frontiers in Computational Neuroscience*, *7*(April), 1–19. https://doi.org/10.3389/fncom.2013.00024

Gomez-Marin, A., & Louis, M. (2014). Multilevel control of run orientation in Drosophila larval chemotaxis. *Frontiers in Behavioral Neuroscience*, *8*(FEB), 1–14. https://doi.org/10.3389/fnbeh.2014.00038

Gomez-Marin, A., Stephens, G. J., & Louis, M. (2011). Active sampling and decision making in Drosophila chemotaxis. *Nature Communications*, *2*(1). https://doi.org/10.1038/ncomms1455

Green, C. H., Burnet, B., & Connolly, K. J. (1983). Organization and patterns of inter- and intraspecific variation in the behaviour of Drosophila larvae. *Animal Behaviour*, *31*(1), 282–291. https://doi.org/10.1016/S0003-3472(83)80198-5

Heckscher, E. S., Lockery, S. R., & Doe, C. Q. (2012). Characterization of Drosophila larval crawling at the level of organism, segment, and somatic body wall musculature. *Journal of Neuroscience*, *32*(36), 12460–12471. https://doi.org/10.1523/JNEUROSCI.0222-12.2012

Heckscher, E. S., Zarin, A. A., Faumont, S., Clark, M. Q., Manning, L., Fushiki, A., Schneider-Mizell, C. M., Fetter, R. D., Truman, J. W., Zwart, M. F., Landgraf, M., Cardona, A., Lockery, S. R., & Doe, C. Q. (2015). Even-Skipped+ Interneurons Are Core Components of a Sensorimotor Circuit that Maintains Left-Right Symmetric Muscle Contraction Amplitude. *Neuron*, *88*(2), 314–329. https://doi.org/10.1016/j.neuron.2015.09.009

Hernandez-Nunez, L., Belina, J., Klein, M., Si, G., Claus, L., Carlson, J. R., & Samuel, A. D. T. (2015). Reverse-correlation analysis of navigation dynamics in Drosophila larva using optogenetics. *ELife*, *4*(MAY), 1–16. https://doi.org/10.7554/eLife.06225

Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, *25*(15), 1965–1978. https://doi.org/10.1002/joc.1276

Höhna, S., Landis, M. J., & Heath, T. A. (2017). Phylogenetic inference using RevBayes. *Current Protocols in Bioinformatics*, *2017*(March), 6.16.1-6.16.34. https://doi.org/10.1002/cpbi.22

Hohna, S., Landis, M. J., Heath, T. A., Boussau, B., Lartillot, N., Moore, B. R., Huelsenbeck, J. P., & Ronquist, F. (2016). RevBayes: Bayesian phylogenetic inference using graphical models and an interactive model-specification language. *Systematic Biology*, *65*(4), 726–736. https://doi.org/10.1093/sysbio/syw021

Huelsenbeck, J., & Bruce, R. (2003). Detecting Correlation between Characters in a Comparative Analysis with Uncertain Phylogeny. *Evolution*, *57*(6), 1237–1247.

Hughes, C. L., & Thomas, J. B. (2007). A sensory feedback circuit coordinates muscle activity in Drosophila. *Molecular and Cellular Neuroscience*, *35*(2), 383–396. https://doi.org/10.1016/j.mcn.2007.04.001

Kalay, G., Atallah, J., Sierra, N. C., Tang, A. M., Crofton, A. E., Murugesan, M. K., Wykoff-Clary,

S., & Lott, S. E. (2020). Evolution of larval segment position across 12 Drosophila species*. *Evolution*, *74*(7), 1409–1422. https://doi.org/10.1111/evo.13911

Kappeler, P. (2010). *Animal Behaviour: Evolution and Mechanisms*. Springer.

Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution*, *30*(4), 772–780. https://doi.org/10.1093/molbev/mst010

Klein, M., Afonso, B., Vonner, A. J., Hernandez-Nunez, L., Berck, M., Tabone, C. J., Kane, E. A., Pieribone, V. A., Nitabach, M. N., Cardona, A., Zlatic, M., Sprecher, S. G., Gershow, M., Garrity, P. A., Samuel, A. D. T., & Sternberg, P. W. (2015). Sensory determinants of behavioral dynamics in Drosophila thermotaxis. *Proceedings of the National Academy of Sciences of the United States of America*, *112*(2), E220–E229. https://doi.org/10.1073/pnas.1416212112

Kohsaka, H., Guertin, P. A., & Nose, A. (2017). Neural Circuits Underlying Fly Larval Locomotion. *Current Pharmaceutical Design*, *23*(12), 1722–1733. https://doi.org/10.2174/1381612822666161208120835

Kohsaka, H., Takasu, E., Morimoto, T., & Nose, A. (2014). A group of segmental premotor interneurons regulates the speed of axial locomotion in drosophila larvae. *Current Biology*, *24*(22), 2632–2642. https://doi.org/10.1016/j.cub.2014.09.026

Kohsaka, H., Zwart, M. F., Fushiki, A., Fetter, R. D., Truman, J. W., Cardona, A., & Nose, A. (2019). Regulation of forward and backward locomotion through intersegmental feedback circuits in Drosophila larvae. *Nature Communications*, *10*(1), 2654. https://doi.org/10.1038/s41467-019-10695-y

Kwon, Y., Shen, W. L., Shim, H. S., & Montell, C. (2010). Fine thermotactic discrimination between the optimal and slightly cooler temperatures via a TRPV channel in chordotonal neurons. *Journal of Neuroscience*, *30*(31), 10465–10471. https://doi.org/10.1523/JNEUROSCI.1631-10.2010

Kwon, Y., Shim, H. S., Wang, X., & Montell, C. (2008). Control of thermotactic behavior via coupling of a TRP channel to a phospholipase C signaling cascade. *Nature Neuroscience*, *11*(8), 871–873. https://doi.org/10.1038/nn.2170

Lahiri, S., Shen, K., Klein, M., Tang, A., Kane, E., Gershow, M., Garrity, P., & Samuel, A. D. T. (2011). Two alternating motor programs drive navigation in Drosophila larva. *PLoS ONE*, *6*(8). https://doi.org/10.1371/journal.pone.0023180

Lartillot, N., & Poujol, R. (2011). A phylogenetic model for investigating correlated evolution of substitution rates and continuous phenotypic characters. *Molecular Biology and Evolution*, *28*(1), 729–744. https://doi.org/10.1093/molbev/msq244

Lin, J. (1991). Divergence Measures Based on the Shannon Entropy. *IEEE Transactions on Information Theory*, *37*(1), 145–151. https://doi.org/10.1109/18.61115

Liu, L., Yermolaieva, O., Johnson, W. A., Abboud, F. M., & Welsh, M. J. (2003). Identification and function of thermosensory neurons in Drosophila larvae. *Nature Neuroscience*, *6*(3), 267–273. https://doi.org/10.1038/nn1009

Loveless, J., Lagogiannis, K., & Webb, B. (2019). Modelling the neuromechanics of exploration and taxis in larval Drosophila. *PLoS Computational Biology*, 1–33. https://doi.org/10.5281/zenodo.1432637

Loveless, J., & Webb, B. (2018). A Neuromechanical Model of Larval Chemotaxis. *Integrative and Comparative Biology*, *58*(5), 906–914. https://doi.org/10.1093/icb/icy094

Luo, L., Gershow, M., Rosenzweig, M., Kang, K., Fang-Yen, C., Garrity, P. A., & Samuel, A. D. T. (2010). Navigational decision making in Drosophila thermotaxis. *Journal of Neuroscience*, *30*(12), 4261–4272. https://doi.org/10.1523/JNEUROSCI.4090-09.2010

Makino, T., & Kawata, M. (2012). Habitat variability correlates with duplicate content of drosophila genomes. *Molecular Biology and Evolution*, *29*(10), 3169–3179. https://doi.org/10.1093/molbev/mss133

Markow. (2015). The secret lives of Drosophila flies. *ELife*, *4*(JUNE), 1–9. https://doi.org/10.7554/eLife.06793

Markow, T. A., & O'Grady, P. M. (2005). *Drosophila: A guide to species identification and use*.

Academic Press.

Miller, D. E., Staber, C., Zeitlinger, J., & Hawley, R. S. (2018). Highly contiguous genome
assemblies of 15 drosophila species generated using nanopore sequencing. *G3: Genes,*
Genomes, Genetics, *8*(10), 3131–3141. https://doi.org/10.1534/g3.118.200160

Ni, L., Klein, M., Svec, K. V., Budelli, G., Chang, E. C., Ferrer, A. J., Benton, R., Samuel, A. D.
T., & Garrity, P. A. (2016a). The ionotropic receptors IR21a and IR25a mediate cool
sensing in Drosophila. *ELife*, *5*(APRIL2016), 1–12. https://doi.org/10.7554/eLife.13254

Ni, L., Klein, M., Svec, K. V., Budelli, G., Chang, E. C., Ferrer, A. J., Benton, R., Samuel, A. D.
T., & Garrity, P. A. (2016b). The ionotropic receptors IR21a and IR25a mediate cool
sensing in Drosophila. *ELife*, *5*(APRIL2016). https://doi.org/10.7554/eLife.13254

Nicholson, A. J. (1954). An outline of the dynamics of animal. *Australian Journal of Zoology*,
2(1), 9–65.

Ohyama, T., Jovanic, T., Denisov, G., Dang, T. C., Hoffmann, D., Kerr, R. A., & Zlatic, M.
(2013). High-Throughput Analysis of Stimulus-Evoked Behaviors in Drosophila Larva
Reveals Multiple Modality-Specific Escape Strategies. *PLoS ONE*, *8*(8).
https://doi.org/10.1371/journal.pone.0071706

Ohyama, T., Schneider-Mizell, C. M., Fetter, R. D., Aleman, J. V., Franconville, R., Rivera-Alba,
M., Mensh, B. D., Branson, K. M., Simpson, J. H., Truman, J. W., Cardona, A., & Zlatic, M.
(2015). A multilevel multimodal circuit enhances action selection in Drosophila. *Nature*,
520(7549), 633–639. https://doi.org/10.1038/nature14297

Osborne, K. A., Robichon, A., Burgess, E., Butland, S., Shaw, R. A., Coulthard, A., Pereira, H.
S., Greenspan, R. J., & Sokolowski, M. B. (1997). Natural behavior polymorphism due to a
cGMP-dependent protein kinase of Drosophila. *Science*, *277*(5327), 834–836.
https://doi.org/10.1126/science.277.5327.834

Pehlevan, C., Paoletti, P., & Mahadevan, L. (2016). Integrative neuromechanics of crawling in
D. Melanogaster Larvae. *ELife*, *5*(2016JULY), 1–23. https://doi.org/10.7554/eLife.11031

Petersen, M., Tenedini, F., Hoyer, N., Kutschera, F., & Soba, P. (2018). Assaying Thermo-
nociceptive Behavior in Drosophila Larvae. *Bio-Protocol*, *8*(4), 1–12.
https://doi.org/10.21769/bioprotoc.2737

Risse, B., Berh, D., Otto, N., Klämbt, C., & Jiang, X. (2017). FIMTrack: An open source tracking
and locomotion analysis software for small animals. *PLoS Computational Biology*, *13*(5),
1–15. https://doi.org/10.1371/journal.pcbi.1005530

Risse, B., Thomas, S., Otto, N., Löpmeier, T., Valkov, D., Jiang, X., & Klämbt, C. (2013). FIM, a
Novel FTIR-Based Imaging Method for High Throughput Locomotion Analysis. *PLoS ONE*,
8(1), 1–11. https://doi.org/10.1371/journal.pone.0053963

Rohlfs, M., & Hoffmeister, T. S. (2003). An evolutionary explanation of the aggregation model of
species coexistence. *Proceedings of the Royal Society B: Biological Sciences*,
270(SUPPL. 1), 33–35. https://doi.org/10.1098/rsbl.2003.0002

Rosenzweig, M., Brennan, K. M., Tayler, T. D., Phelps, P. O., Patapoutian, A., & Garrity, P. A.
(2005a). The Drosophila ortholog of vertebrate TRPA1 regulates thermotaxis. *Genes and*
Development, *19*(4), 419–424. https://doi.org/10.1101/gad.1278205

Rosenzweig, M., Brennan, K. M., Tayler, T. D., Phelps, P. O., Patapoutian, A., & Garrity, P. A.
(2005b). The Drosophila ortholog of vertebrate TRPA1 regulates thermotaxis. *Genes and*
Development, *19*(4), 419–424. https://doi.org/10.1101/gad.1278205

Rosenzweig, M., Kang, K. J., & Garrity, P. A. (2008). Distinct TRP channels are required for
warm and cool avoidance in Drosophila melanogaster. *Proceedings of the National*
Academy of Sciences of the United States of America, *105*(38), 14668–14673.
https://doi.org/10.1073/pnas.0805041105

Ruiz-Dubreuil, G., Burnet, B., Connolly, K., & Furness, P. (1996). Larval foraging behaviour and
competition in Drosophila melanogaster. *Heredity*, *76*(1), 55–64.
https://doi.org/10.1038/hdy.1996.7

Schmidt-Nielsen, K. (1997). *Animal Physiology: Adaptation and Environment*. Cambridge
University Press.

Schneider-Mizell, C. M., Gerhard, S., Longair, M., Kazimiers, T., Li, F., Zwart, M. F., Champion,

A., Midgley, F. M., Fetter, R. D., Saalfeld, S., & Cardona, A. (2016). Quantitative
neuroanatomy for connectomics in Drosophila. *ELife*, *5*(MARCH2016), 1–36.
https://doi.org/10.7554/eLife.12059
Shen, W. L., Kwon, Y., Adegbola, A. A., Luo, J., Chess, A., & Montell, C. (2011). Function of
rhodopsin in temperature discrimination in Drosophila. *Science*, *331*(6022), 1333–1336.
https://doi.org/10.1126/science.1198904
Simpson, J. H., & Looger, L. L. (2018). *Functional Imaging and Optogenetics in Drosophila*.
208(April), 1291–1309.
Sims, D. W., Humphries, N. E., Hu, N., Medan, V., & Berni, J. (2019). Optimal searching
behaviour generated intrinsically by the central pattern generator for locomotion. *ELife*, *8*,
1–31. https://doi.org/10.7554/eLife.50316
Sokolowski, M. B. (1980). Foraging strategies of Drosophila melanogaster: a chromosomal
analysis. *Behavior Genetics*, *10*(3), 291–302.
Sokolowski, M. B. (1985). Genetics and ecology of Drosophila melanogaster larval foraging and
pupation behaviour. *Journal of Insect Physiology*, *31*(11), 857–864.
https://doi.org/10.1016/0022-1910(85)90103-9
Sokolowski, M. B., Kent, C., & Wong, J. (1984). Drosophila larval foraging behaviour:
Developmental stages. *Animal Behaviour*, *32*(3), 645–651. https://doi.org/10.1016/S0003-
3472(84)80139-6
Sokolowski, M. B., Pereira, H. S., & Hughes, K. (1997). Evolution of foraging behavior in
drosophila by density-dependent selection. *Proceedings of the National Academy of
Sciences of the United States of America*, *94*(14), 7373–7377.
https://doi.org/10.1073/pnas.94.14.7373
Stoessel, A., & Fischer, M. S. (2012). Comparative intralimb coordination in avian bipedal
locomotion. *Journal of Experimental Biology*, *215*(23), 4055–4069.
https://doi.org/10.1242/jeb.070458
Sun, X., Liu, Y., Liu, C., Mayumi, K., Ito, K., Nose, A., & Kohsaka, H. (2020). A neuromechanical
model and kinematic analyses for Drosophila larval crawling based on physical
measurements. *BioRxiv*.
Tastekin, I., Riedl, J., Schilling-Kurz, V., Gomez-Marin, A., Truman, J. W., & Louis, M. (2015).
Role of the subesophageal zone in sensorimotor control of orientation in drosophila larva.
Current Biology, *25*(11), 1448–1460. https://doi.org/10.1016/j.cub.2015.04.016
Theunissen, L. M., Bekemeier, H. H., & Dürr, V. (2015). Comparative whole-body kinematics of
closely related insect species with different body morphology. *Journal of Experimental
Biology*, *218*(3), 340–352. https://doi.org/10.1242/jeb.114173
Thompson, N. E., O'Neill, M. C., Holowka, N. B., & Demes, B. (2018). Step width and frontal
plane trunk motion in bipedal chimpanzee and human walking. *Journal of Human
Evolution*, *125*, 27–37. https://doi.org/10.1016/j.jhevol.2018.09.006
Tracey, W. D., Tracey, W. D., Wilson, R. I., Wilson, R. I., Laurent, G., Laurent, G., Benzer, S., &
Benzer, S. (2003). Gene Essential for Nociception. *Cell*, *113*, 261–273.
Turelli, M., Cooper, B. S., Richardson, K. M., Ginsberg, P. S., Peckenpaugh, B., Antelope, C. X.,
Kim, K. J., May, M. R., Abrieux, A., Wilson, D. A., Bronski, M. J., Moore, B. R., Gao, J. J.,
Eisen, M. B., Chiu, J. C., Conner, W. R., & Hoffmann, A. A. (2018). Rapid Global Spread
of wRi-like Wolbachia across Multiple Drosophila. *Current Biology*, *28*(6), 963-971.e8.
https://doi.org/10.1016/j.cub.2018.02.015
Venken, K. J. T., Simpson, J. H., & Bellen, H. J. (2011). Genetic manipulation of genes and
cells in the nervous system of the fruit fly. *Neuron*, *72*(2), 202–230.
https://doi.org/10.1016/j.neuron.2011.09.021
Watada, M., Hayashi, Y., Watanabe, K., Mizutani, S., Mure, A., Hattori, Y., & Uemura, T. (2020).
Divergence of Drosophila species: longevity and reproduction under different nutrient
balances . *Genes to Cells*, *June*. https://doi.org/10.1111/gtc.12798
Watanabe, K., Kanaoka, Y., Mizutani, S., Uchiyama, H., Yajima, S., Watada, M., Uemura, T., &
Hattori, Y. (2019). Interspecies Comparative Analyses Reveal Distinct Carbohydrate-
Responsive Systems among Drosophila Species. *Cell Reports*, *28*(10), 2594-2607.e7.

https://doi.org/10.1016/j.celrep.2019.08.030

Wertheim, B., Marchais, J., Vet, L. E. M., & Dicke, M. (2002). Allee effect in larval resource
exploitation in Drosophila: An interaction among density of adults, larvae, and micro-
organisms. *Ecological Entomology*, *27*(5), 608–617. https://doi.org/10.1046/j.1365-
2311.2002.00449.x

Zwart, M. F., Pulver, S. R., Truman, J. W., Fushiki, A., Fetter, R. D., Cardona, A., & Landgraf,
M. (2016). Selective Inhibition Mediates the Sequential Recruitment of Motor Pools.
Neuron, *91*(3), 615–628. https://doi.org/10.1016/j.neuron.2016.06.031

Supplementary Figure 1. Interspecies comparison between larva length and kinematics parameters. (A) Axial body length of larvae of the 11 species. Sample numbers are the following: *Dvir*: n=69; *Dmel*: n=72; *Dmoj*: n=82; *Dper*: n=76; *Dpse*: n=60; *Dsec*: n=68; *Dwil*: n=66; *Dere*: n=69; *Dmau*: n=69; *Dana*: n=68; *Dyak*: n=56. (B) Scatter plot of the crawling speed at 24°C against the larva length of the 11 species. (C) Scatter plot of the bend probability at 24°C against the larva length of the 11 species.   Median ± sem are shown in B and C.

Supplementary Figure 2. World maps of histograms of habitat temperature of the ***Drosophila*** **species.** The top of each panel shows a map of the habitat of the labelled species. The bottom on each panel shows a histogram of minimum (green) and maximum (magenta) temperatures within the habitat. Asterisks denote the warmest peak in the minimum habitat temperature. The temperature at the peak is called Tmin in this study.

Supplementary Figure 3. Scatter plots of kinematics parameters and the modes of average, minimum, and maximum habitat temperatures. (A) Scatter plot of bend probability of the 11 species at 24°C against the modes of the average (left), minimum (center), and maximum (right) habitat temperatures. (B) Scatter plot of the crawling speed of the 11 species at 24°C against the modes of the average (left), minimum (center), and maximum (right) habitat temperatures. Median ± sem are shown.

Supplementary Figure 4. Bend probability in the 11 *Drosophila* species at 40°C. (A)

Bend probability of the 11 species at 32°C and 40°C. The data at 32°C are the same as in

Figure 4A. (B) Scatter plot of the bend probability at 40°C against at 32°C. Sample numbers

at 40°C are the following: *Dvir*: n=20; *Dmel*: n=28; *Dmoj*: n=23; *Dper*: n=21; *Dpse*: n=16;

Dsec: n=27; *Dwil*: n=27; *Dere*: n=22; *Dmau*: n=26; *Dana*: n=23; *Dyak*: n=20. Sample

numbers at 32°C are described in the legend of Figure 4. Median $\pm$ sem are shown in B.

Supplementary Figure 5. Crawling speed in the 11 *Drosophila* species at 40°C. (A)
Crawling speed of the 11 species at 32°C and 40°C. The data at 32°C are the same as in
Figure 5A. (B) Scatter plot of the crawling speed at 40°C against at 32°C. Sample numbers
at 40°C are the following: *Dvir*: n=20; *Dmel*: n=28; *Dmoj*: n=23; *Dper*: n=21; *Dpse*: n=16;
Dsec: n=27; *Dwil*: n=27; *Dere*: n=22; *Dmau*: n=26; *Dana*: n=23; *Dyak*: n=20. Sample
numbers at 32°C are described in the legend of Figure 5. Median $\pm$ sem are shown in B.

47