Interspecies variation of larval locomotion kinematics in the

genus *Drosophila* and its relation to habitat temperature

Yuji Matsuo[a], Akinao Nose[a,b] and Hiroshi Kohsaka[a*]

[a] Department of Complexity Science and Engineering, Graduate School of

Frontier Science, the University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa, 277-

8561 Chiba, Japan

[b] Department of Physics, Graduate School of Science, the University of Tokyo, 7-

3-1 Hongo, Bunkyo-ku, 133-0033 Tokyo, Japan

* For correspondence: (kohsaka@edu.k.u-tokyo.ac.jp, +81-4-7136-3922)

Abstract

Speed and trajectory of locomotion are characteristic traits of individual species.

During evolution, locomotion kinematics is likely to have been tuned for survival in the

habitats of each species. Although kinematics of locomotion is thought to be influenced by

habitats, the quantitative relation between the kinematics and environmental factors has not

been fully revealed. Here, we performed comparative analyses of larval locomotion in 11

Drosophila species. We found that larval locomotion kinematics are divergent among the

species. The diversity is not correlated to the body length but is correlated instead to the

habitat temperature of the species. Phylogenetic analyses using Bayesian inference suggest

that the evolutionary rate of the kinematics is diverse among phylogenetic trees. The results

of this study imply that the kinematics of larval locomotion has diverged in the evolutionary

history of the genus *Drosophila* and evolved under the effects of the ambient temperature of

habitats.

Keywords: animal locomotion, larval crawling, the genus Drosophila, kinematics, evolution

Introduction

Kinematics of animal locomotion is a critical trait enabling each species to survive

in their habitats (Alexander, 2006). Movement patterns have been sculpted during their

evolution by adaptation to their environments, and could have diverged among species

(Kappeler, 2010). Comparative analyses have identified several examples of differences in

the kinematics of locomotion within a group of related species, including insects (Theunissen

et al., 2015), reptiles (Fuller et al., 2011), birds (Stoessel & Fischer, 2012), and primates

(Thompson et al., 2018). To take an example, two gecko species inhabiting either sandy or

rocky environments have been shown to exhibit distinct postures (Fuller et al., 2011).

Whereas the interspecies divergence in locomotion patterns can be observed in various

phylogenetic branches of the animal kingdom, quantitative comparative analyses of

locomotion kinematics remain limited.

Flies of the genus *Drosophila* have long been used as a model to study interspecific

diversification and evolution (Markow & O'Grady, 2005). One salient example of the

interspecific variations in the genus *Drosophila* is food for larvae. Some species eat multiple

kinds of foods ("generalists" including *Drosophila melanogaster*) while others have strong

preferences in food ("specialists" including *Drosophila sechellia*, which has specialized to

feed on Morinda fruits) (Markow, 2015; Watada et al., 2020; Watanabe et al., 2019). These

diverged species in the genus *Drosophila* offer an opportunity to perform interspecific

comparisons in various animal traits including larval locomotion.

Fly larvae, or maggots, have been widely used in the study of the kinematics of

locomotion (Berrigan & Lighton, 1993; Berrigan & Pepin, 1995; Fox et al., 2006; Green et al.,

1983; Ruiz-Dubreuil et al., 1996; Sokolowski, 1985; Sokolowski et al., 1997). Among the fly

species, *Drosophila melanogaster* (*Dmel*) is one of the most examined species, especially

by virtue of the availability of resources in genetics and connectomics (Ohyama et al., 2015;

Schneider-Mizell et al., 2016; Simpson & Looger, 2018; Venken et al., 2011). *Dmel* larvae

locomote by a sequence of forward crawling, and changes of crawling direction achieved by

bending their bodies (Green et al., 1983).   The kinematics of larval locomotion is affected

by ambient temperature (Kwon et al., 2010; Liu et al., 2003; Ohyama et al., 2013;

Rosenzweig et al., 2005a). In thermotaxis behaviour in temperature gradient environments,

larvae regulate the length of crawling runs between turns and the size and direction of turns

(Luo et al., 2010), and the probability of turns is also affected by ambient temperature

gradients (Klein et al., 2015). In contrast to the intensive studies on larval behavior in *Dmel*,

locomotion kinematics in the larvae of its sister species in the genus *Drosophila* remains

unclear.

Here we conducted an interspecies comparison of the kinematics of larval

locomotion in the genus *Drosophila*. We address two questions in this study: are locomotion

kinematics of larvae similar among *Drosophila* species?; and, if the kinematics are diverged,

what factors are related to the diversity? To this aim, we recorded locomotion of larvae of 11

Drosophila species and extracted kinematic parameters using the tracking software

FIMTrack (Risse et al., 2017). Clustering analysis with Jensen-Shannon divergence and

statistical analyses show that two kinematics parameters (bend probability and crawling

speed) differ among the *Drosophila* species. We found that kinematics varies with habitat

temperature but not with body size. The relationship between the kinematics and minimum

habitat temperature held at two distinct ambient temperatures: 24°C and 32°C. Phylogenetic

analyses of these kinematics, based on Bayesian inference (Hohna et al., 2016), suggests

that the rate of evolution of the kinematics is diverged among phylogenetic branches. Among

eight traits we tested, the evolution of the crawling speed at 24°C and 32°C were correlated.

Consequently, our results suggest that the kinematics of larval locomotion in the genus

Drosophila diverged in response to environmental variation in ambient temperature.

Results

Kinematic analysis of crawling and bending behaviour in *Drosophila* larvae

In this study, we analyzed the kinematics of larval locomotion. To this aim, we

recorded the locomotion of fly larvae that were crawling freely on an agarose substrate stage

on a temperature-controlled plate (Figure 1A; see Method section for details). We placed

eight to ten larvae at the centre of the stage, which was kept at 24°C, illuminated them with

infrared light, which is invisible to fly larvae and does not affect their behaviour, and recorded

larval locomotion for three minutes at five frames per second (Figure 1B). Maximum

projection of the time-series images shows traces of larval locomotion as multiple curves of

multiple larvae (Figure 1C). In an example of *Drosophila melanogaster* (Figure 1D), the

traces show smooth curved lines interconnected with angles where larvae exhibit turning

behaviour and change crawling direction, which is consistent with previous studies (Berni et

al., 2012; Gershow et al., 2012; Gomez-Marin et al., 2011; Gomez-Marin & Louis, 2014;

Green et al., 1983; Sims et al., 2019; Tastekin et al., 2015). Larvae under these conditions

predominantly exhibit forward locomotion in which muscular contraction is propagating from

the posterior to anterior segments. Characteristics of larval locomotion can be described by

two measures: the bend angle and centroid speed (Gershow et al., 2012; Gomez-Marin et

al., 2011). The bend angle measures the angle of body axis bending (Figure 1E) and the

centroid speed is the speed of the position of the larval centroid (Figure 1F). To obtain these

values for each larva at each time frame, we used the object tracking software for small

animals, FIMTrack (Risse et al., 2017). The turning behaviour can be detected by the change

in the angle of the body axis (Figure 1E) and the reduction in the centroid speed (Figure 1F),

as reported previously (Gershow et al., 2012; Gomez-Marin et al., 2011). Accordingly, we

used the bend angle and centroid speed for quantitative analysis of larval locomotion in this

study.

Figure 1. Measurement of larval crawling. (A) Setup of the recording of larval crawling with

infrared light and temperature control plate. (B) An example image of multiple larvae recorded

by the setup (A). (C) The trajectories of larval locomotion of *Drosophila melanogaster*

recorded for two minutes. (D) A trajectory of single *Drosophila melanogaster* locomotion.

Locations where the larva changes its direction are labelled (i) to (iv). (E and F). Bend angle

(E) and centroid speed (F) of the larva in (D). Labels (i) to (iv) correspond to those in (D).

Classification of locomotion properties in the genus *Drosophila*

116   For interspecies comparison of *Drosophila* larval locomotion, we collected 11 fly

species in the genus *Drosophila*, whose genome sequences have been read (A. G. Clark et

al., 2007; Miller et al., 2018), and for which living individuals are available from fly stock

centres (KYORIN-Fly, Fly Stocks of Kyorin University; KYOTO Stock Center (DGRC) at the

Kyoto Institute of Technology). The 11 *Drosophila* (*D.*, hereafter) species consist of *D.*

ananassae (*Dana*)*, D. erecta* (*Dere*)*, D. mauritiana* (*Dmau*)*, D. melanogaster* (*Dmel*)*, D.*

mojavensis (*Dmoj*)*, D. persimilis* (*Dper*)*, D. pseudoobscula* (*Dpse*)*, D. sechellia* (*Dsec*)*, D.*

virilis (*Dvir*)*, D. willistoni* (*Dwil*) *and D. yakuba* (*Dyak*). Nine species (all but *Dvir* and *Dmoj*)

are classified as subgenus *Sophophora* and the remaining two (*Dvir* and *Dmoj*) are *non-*

Sophophora species. We plotted the centroid speed and bend angle of freely crawling larvae

of each species (Figure 2A). In all the cases, data points accumulated around a region where

the bend angle is zero (Figure 2A), which reflects an observation that larvae do not bend

during the majority of the time (Figure 1D). We noticed that the deviations from zero in bend

angle axis are different among species. For example, data points in *Dvir* are scattered along

the bend angle axis more than those in *Dwil* (Figure 2A). To quantify similarities in the

distribution of the two-dimensional plots among the species, we calculated Jensen-Shannon

divergence, which measures the similarity of two probability distributions (Lin, 1991). We

classified the 11 species based on the similarity in the probability distribution of the crawling

speed and bend angle plots by hierarchical clustering (Figure 2B and 2C). The classification

analysis shows that four species (*Dmel, Dvir, Dmoj and Dper*) form a cluster. These four

species show scattered data points along the bend angle axis compared with the other seven

species (Figure 2A). This observation suggests that the kinematics of larvae in the genus

Drosophila is diverse among species.

Figure 2. Interspecific comparison of kinematics of larval crawling in the genus

Drosophila. (A) Plots of the speed of the centroid of larvae and the bend angle in the 11

Drosophila species. Each point corresponds to a datum of a single larva of a species in a

single time frame. The number of larvae in each species is the following: *Dvir*: n=24; *Dmel*:

n=24; *Dmoj*: n=30; *Dper*: n=27; *Dpse*: n=19; *Dsec*: n=21; *Dwil*: n=18; *Dere*: n=22; *Dmau*:

n=26; *Dana*: n=22; *Dyak*: n=16. (B) Jensen-Shannon divergences of the probability density

of (A). (C) Hierarchical clustering of the kinematics of larval locomotion of the 11 *Drosophila*

species.

Definition of the bend probability and crawling speed

The clustering analysis suggests that the kinematics is divergent among species.

To interpret the diversity in terms of locomotion behaviour, we defined two indices: the bend

probability and crawling speed. The bend probability measures how often larvae bend their

body laterally. We set the minimum angle of bend as 20 degrees, which was used previously

(Risse et al., 2013), and labelled larvae that bent more than this threshold angle to the right

or left side as "Bending" (Figure 3A and 3B). This threshold allowed us to extract the

difference in the bending rate among the species we examined, and we found that while

Dmel larvae exhibit larger bend angles than this threshold (Figure 3A), the bend angles in

Dwil larvae are mostly less than the threshold (Figure 3B). We defined the bend probability

of each single larva as a ratio of the number of time frames labelled "Bending" to the total

frame number. To define the second index crawling speed, we labelled larvae that bent less

than the threshold as "Crawling" (Figure 3A and 3B). We defined the median of centroid

speeds of larvae that were labelled "Crawling" as the crawling speed for each single larva.

By these definitions, we calculated the bend probability and crawling speed of each larva of

the species.

Kinematics of larval locomotion is diverse among the *Drosophila* species

We plotted the bend probability of each species (Figure 3C). A statistical analysis

shows that the bend probability of the species is diverse (p = 6.8 x 10^{-25}, Kruskal-Walls test).

While *Dvir* larvae exhibited frequent bending (bend probability: 0.36 ± 0.05, n = 24), *Dwil*

larvae rarely bend (bend probability: 0.003 ± 0.004, n = 18) (Figure 3C). We also plotted the

crawling speed of the *Drosophila* species (Figure 3D). The statistical analysis shows that the

speed is also diverse among the species (p = 9.3 x 10^{-24}, Kruskal-Walls test). For example,

Dwil larvae crawl faster than *Dvir* larvae do (crawling speed: 1.46 ± 0.07 mm/sec, n = 24 in

Dwil; crawling speed: 0.95 ± 0.05 mm/sec, n = 18 in *Dvir*). These analyses indicate that the

bend probability and crawling speed are differentiated in the genus *Drosophila*.

To capture the trends in the diversity of larval kinematics, we plotted the data in the

space of the crawling speed and bend probability (Figure 3E). The graph shows a negative

correlation between them (Pearson correlation = -0.76, p = 0.0064). While we used 20

degrees as the threshold for defining the bend probability (Figure 3A and 3B), the negative

correlation between the crawling speed and bend probability is robust to the choice of the

threshold (Pearson correlation = -0.80, p = 0.0030, when the threshold is 10 degrees;

Pearson correlation = -0.71, p = 0.014, when the threshold is 30 degrees; Supplementary

Figure 1A - 1C). Furthermore, the median of absolute values of bend angle instead of the

bend probability exhibits a negative correlation to the crawling speed (Pearson correlation =

-0.66, p = 0.026; Supplementary Figure 1D). To sum, the crawling speed and bend probability

are diverse among the *Drosophila* species and negatively correlated.

Figure 3. Bend probability and crawling speed of larval locomotion in the 11

Drosophila **species at 24°C and their relationship to habitat temperature of each**

species. (A and B) Plots of the speed of the centroid of larvae and the bend angle of *D.*

melanogaster (A) and *D. willistoni* (B). Each point corresponds to a datum of a single larva

of a species at a single time frame. Thick horizontal lines denote the threshold between

crawling and bending. (C) Bend probability of individual larvae of each species. (D) Crawling

speed of individual larvae of each species. Sample numbers in (C) and (D) are the following:

Dvir: n=24; *Dmel*: n=24; *Dmoj*: n=30; *Dper*: n=27; *Dpse*: n=19; *Dsec*: n=21; *Dwil*: n=18; *Dere*:

n=22; *Dmau*: n=26; *Dana*: n=22; *Dyak*: n=16. (E) Scatter plot of bend probability at 24°C

against crawling speed at 24°C. Median ± sem is shown. The red line shows the linear

regression function and the shaded area represents the 95% confidence band. The point

estimate of the Pearson correlation and its 95% confidence interval is -0.76 and [-0.94, -0.30].

Comparison between intraspecies and interspecies variability

The comparison of the 11 *Drosophila* species represents the natural variability in

larval locomotion across them. Here, it should be noted that we used a single strain for each

species in the analysis. For that, the variability appeared above could result not only from

interspecies but also intraspecies diversity. Even if there was no interspecies variability,

sampling data from population with high intraspecies variation could lead to an apparent

diversity across the species. To address this issue, we compared intraspecific deviation with

interspecific variability. If intraspecific diversity is a dominant factor for the variability, the

deviation within species should be comparable to that among species. In contrast, if

interspecific diversity is a major cause, the deviation within species would be smaller than

that across species. We examined intraspecies variation in two species in subgenus

Sophophora (*Dmel* and *Dana*) and one *non-Sophophora* species (*Dvir*) (Figure 4). Two

isofemale strains of each species were obtained from the Kyorin stock centre and the larval

kinematics of them were measured. To evaluate the contribution of the interspecific

deviations to the total deviations, we conducted the analysis of variance (ANOVA). We found

a significant interspecific difference in the bend probability (p=0.008, the one-way ANOVA)

although the difference in the crawling speed was marginal (p = 0.08, the one-way ANOVA).

This observation implies the existence of interspecies diversity in larval locomotion.

Figure 4. Intraspecific comparison in larval locomotion. Scatter plot of bend probability

at 24°C against crawling speed at 24°C of nine strains from three species. Sample numbers

are the following: *Dvir*: n=44; *Dvir* (Hokkaido): n=32; *Dvir* (Toyama): n=35; *Dmel*: n=33; *Dmel*

(Kyoto): n=47; *Dmel (Iriomote)*: n=33; *Dana*: n=33; *Dana (Ogasawara)*: n=40; *Dana*

(Iriomote): n=20.

Crawling distance is related to crawling speed and bend probability

In the interspecies variability, there is a negative correlation between the crawling

speed and bend probability (Figure 3E). We noticed that this correlation could reflect the

control of the crawling distance by a coordinated change of the crawling speed along with

the bend probability. The bend probability is negatively correlated with the crawling distance

since frequent bending shortens the distance larvae progress in one direction (Figure 5A)

whereas it is obvious that the crawling speed is positively correlated with the crawling

distance (Figure 5B). Accordingly, high crawling speed and low bend probability, which are a

combination appeared in the negative correlation of them (Figure 3E) both contribute to the

increase in the crawling distance. To check whether these geometrical speculations hold in

the locomotion of larvae, we examined the relationship between the kinematic parameters

and the crawling distance (Figure 5C and 5D). Consistent with the conjecture, the crawling

distance is larger when either the bend probability is lower (Figure 5C; Pearson correlation =

-0.78) or the crawling speed is higher (Figure 5D; Pearson correlation = 0.67). These

observations imply that the coordinated changes in the crawling speed and bend probability

would be related to the change in the crawling distance of larvae of each species.

Figure 5. Relationship between crawling distance and the two kinematic parameters in the 11 Drosophila species. (A) Schematics of the relationship between the bend probability and crawling distance. (B) Schematics of the relationship between the crawling speed and crawling distance. (C) Scatter plot of the bend probability and crawling distance in one minute in the 11 species. (D) Scatter plot of the crawling speed and crawling distance in one minute in the 11 species. The source locomotion data are the same as in Figure 3C and 3D. In C and D, the red lines show the linear regression functions and the shaded areas represent the 95% confidence bands. The point estimates of the Pearson correlation and their 95% confidence intervals are -0.78 and [-0.94, -0.34] in C and 0.67 and [0.11, 0.91] in D.

No relationship between the kinematics of larval locomotion and the body length nor phylogenetic relationship

Next, we tried to find factors that relate to the diversification of larval kinematics among species. Among the factors in the morphological differences and the ecological diversity that might be involved in the kinematics diversity (Theunissen et al., 2015) we examined the body length of larvae as a morphological factor and habitat temperature as an ecological factor. A previous study reported an allometric relationship between the body size and the crawling speed in *Diptera* larvae (Berrigan & Pepin, 1995). The authors analyzed larvae in the order *Diptera*, the average size of which spans from 3.7 mm (*Dmel*) to 15.9 mm (*Sarcophaga bullata*). We tested whether the relationship between the body length and the crawling speed also holds within the genus *Drosophila*, a subgroup of the order *Diptera*. The length of the *Drosophila* larvae we used spanned from 3.49 ± 0.05 mm (*Dyak*) to 5.68 ± 0.16 mm (*Dpse*) (Supplementary Figure 2A). We found no significant relationship between the body length and crawling speed (Supplementary Figure 2B; Pearson correlation = 0.04, p = 0.91) and between the body length and the bend probability (Supplementary Figure 2C;

Pearson correlation = 0.55, p = 0.077). These data suggest that larval length is not a

significant factor for larval kinematics variation within the genus *Drosophila*.

Phylogenetic relationship can also be a factor that affects the kinematics. To test

this issue, we focused on two species groups: the obscura species group and replete species

group. *Dper* and *Dpse* belongs to the obscura species group (See Figure 11A). The

kinematics of these two sister species are separated in the distribution of the crawling speed

and bend probability (Figure 3E). Similarly, two species of the replete species group, *Dvir*

and *Dmor* exhibit distinct kinematics (Figure 3E). These observations suggest that

phylogenetic relationship is not a major factor in the divergence of the kinematics.

Relationship between the kinematics of larval locomotion and habitat temperature of

the *Drosophila* species

Habitat temperature is one of the critical factors influencing species traits (Schmidt-

Nielsen, 1997). To test the possible roles of habitat temperature in the evolution of *Drosophila*

larval locomotion, we examined the relationship between habitat temperatures and the

locomotion kinematics in the genus *Drosophila*. The habitat regions of the 11 species were

obtained from the literature (Brake & Baechli, 2008; Makino & Kawata, 2012; Markow &

O'Grady, 2005), and climate temperature data were obtained from a global climate dataset,

WORLDCLIM (Hijmans et al., 2005) (Figure 6A, Supplementary Figure 3). We examined the

relation between larval kinematics (bend probability or crawling speed) and indices of habitat

temperatures (Figure 6B and 6C). We used the mode (the most frequent value) of three

indices of habitat temperatures: average, maximum, and minimum temperature. In the

minimum habitat temperature data of *Dvir* and *Dpse*, we noticed that the mode temperatures

were below 0°C, which cannot be a representative habitat temperature for them

(Supplementary Figure 3). Accordingly, for a milder and more representative index, we used

the warmest peaks in the minimum habitat temperature histogram ("Tmin" in Supplementary

Figure 3). Considering uneven distribution of flies within the region demarcated in

Supplementary Figure 3, the temperature of the warmest peak in the minimum habitat

temperature may represent the coldest temperature the majority of the population of each

species experiences over many years.

We found that the bend probability and crawling speed are both correlated with

average habitat temperature Tave (Figure 6B and 6E. Pearson correlation: bend probability

vs Tave = -0.73, crawling speed vs Tave = 0.66). We further analyzed whether the maximum

and minimum temperature contribute to the correlation. Maximum habitat temperature, Tmax

shows no obvious correlation to the kinematics (Figure 6C and 6F. Pearson correlation: bend

probability vs Tmax = 0.37, crawling speed vs Tmax = -0.31). In contrast, minimum habitat

temperature, Tmin, exhibit stronger correlation than Tmax (Figure 6D and 6G. Pearson

correlation: bend probability vs Tmin = -0.81, crawling speed vs Tmin = 0.74). Even after

omitting extreme values (*Dvir* in the bend probability data and *Dpse* in the crawling speed

data), the correlation remained high (Pearson correlation: the bend probability vs Tmin

(without *Dvir*) = -0.69, the crawling speed vs Tmin (without *Dpse*) = 0.78). Furthermore, even

considering multiple comparisons (four factors: larval length, Tave, Tmax and Tmin), the

correlations between the kinematic parameters and Tmin were statistically significant (p =

0.039 in crawling speed vs Tmin, p = 0.0010 in bend probability vs Tmin; Bonferroni

correction).

The range (or variability) of habitat temperature could also be a critical factor to

determine larval kinematics because species that inhabits in highly variable temperature area

would be insensitive to the change in the ambient temperature whereas those breeding in a

narrow range of temperature would be sensitive to the small shift in the ambient temperature.

To test this point, we examined the relationship between the larval kinematics and the range

of habitat temperature, the difference between maximum and minimum temperature in their

habitat area (Supplementary Figure 4). The range of habitat temperature is correlated with

both the bend probability (Pearson correlation = 0.65, p = 0.030) and the crawling speed

(Pearson correlation = -0.69, p = 0.0019). Consequently, the range of habitat temperature

could affect the larval kinematics. In the following analysis, we will focus on the minimum

temperature Tmin since it showed the most evident correlation to the larval kinematics.

It should be noted that the strains we used have been kept at the housed stocks at

23 or 20°C (See Materials and Methods), which might affect the innate behaviour in wild type

strains. However, comparison between time period after speciation (at least million years ~

10^7 generations (Campbell & Ganetzky, 2012; Russo et al., 1995) and that in laboratories

(100 years ~ $10^{3.5}$ generations) indicates that the duration in laboratories occupies as small

as 0.03% of the time for the evolution of these *Drosophila* strains. In addition, the rearing

temperatures in the stock centres were within the range of habitat temperature of each

species (Supplementary Figure 3). These rearing conditions imply that the strains we

collected from the stock center should possess innate behaviour that was evolved in their

habitat.

To sum, larvae of species inhabiting moderate environments (where Tmin is 15 to

25°C) showed low bend probability and fast crawling, or long crawling distance, whereas

those inhabiting cold environments (where Tmin is 0 to 10°C) exhibited frequent bending and

slow crawling, or short crawling distance. Especially, species that has lower Tmin than the

ambient temperature in this assay (24°C) showed shorter crawling distance, which implies

that excess heat stimuli to these species should reduce their crawling distance.

In the analysis above, we estimated the indices of habitat temperature from the

temperature data in the entire potential habitat of each species because the original habitat

of each strain was unclear. To further check the correlation between the larval kinematics and

habitat temperature, we examined the relationship between the kinematics and habitat

temperature among the strains of which habitat were recorded (Figure 4 and 6H). We used

two *Dmel* strains (collected at Kyoto and Iriomote in Japan), two *Dvir* strains (collected at

Hokkaido and Toyama in Japan), and two *Dana* strains (collected at Ogasawara and Iriomote

in Japan). The habitat temperature at these location for collecting flies is diverse (Figure 6H

right). Intriguingly, the strains originated from the north part of Japan, where the minimum

habitat temperature is low, exhibit slower crawling and high bend probability whereas those

from the south part of Japan, where the minimum habitat temperature is high, show the

opposite trend (Figure 6H left), which is consistent with the observation based on the

temperature of the worldwide statistics (Figure 6D and 6G). Consequently, these

observations imply that habitat temperature should be one of the leading factors in sculpting

the kinematics of larval locomotion in the genus *Drosophila*.

Figure 6. Relationship between the kinematics of larval locomotion and habitat temperature of the Drosophila species. (A) A world map of minimum habitat temperature. (B - D) Scatter plot of the bend probability of the 11 species at 24°C against the average (B), maximum (C), and minimum (D) habitat temperatures. (E - G) Scatter plot of the crawling speed of the 11 species at 24°C against the average (E), maximum (F), and minimum (G)

habitat temperatures. In B to G, median $\pm$ sem are shown and the source locomotion data

are the same as in Figure 3C and 3D. The red lines show the linear regression functions and

the shaded areas represent the 95% confidence bands. The point estimates of the Pearson

correlation and their 95% confidence intervals are -0.73 and [-0.93, -0.23] in B, 0.37 and [-

0.30, 0.79] in C, and -0.81 and [-0.95, -0.41] in D, 0.66 and [0.09, 0.90] in E, -0.31 and [-0.77,

0.36] in F, and 0.74 and [0.25, 0.93] in G. (H) Left: A scatter plot of intraspecific comparison

(the same as Figure 4). Right: A map of Japan representing the minimum habitat

temperatures of the six strains shown in the left panel

Correlation between bend probability and habitat temperature among *Drosophila*

species at high ambient temperature

So far, we have analyzed larval locomotion at 24°C, which is close to the rearing

temperature (See Materials and Methods). Larvae of *Dmel* are known to crawl faster at 32°C

than at 25°C (Ohyama et al., 2013). So, we next examined whether the relation between the

kinematics indices and habitat temperature holds or not, and how the kinematics indices are

changed, at a higher ambient temperature. We performed the same set of measurements

and analysis of larval locomotion at 32°C (Figure 7 and 8). The interspecies comparison

showed that the bend probabilities at 32°C were correlated with minimum habitat temperature

(Figure 7B; Pearson correlation = -0.85, p = 0.0009). To examine the effects of ambient

temperature on the bend probability in detail, we compared the differences of bend probability

at 24°C and 32°C among the species (Figure 7C). We found no significant relationship

between the change in bend probability and the minimum habitat temperature (Figure 7C;

Pearson correlation = 0.40, p = 0.22). Seven species showed no significant changes in the

bend probability at 32°C (Mann-Whitney U-test of *Dmoj*, *Dper*, *Dpse*, *Dsec*, *Dmau*, *Dana* and

Dyak in Figure 7A - 7D), while *Dvir* and *Dmel* showed a decrease in the bend probability at

32°C, and *Dwil* and *Dere* exhibited an increase (Mann-Whitney U-test in Figure 7A - 7C, 7E

and 7F). Accordingly, we concluded that while the shift of the bend probability between the

distinct temperatures is diverse among the *Drosophila* species, the overall trend between the

bend probability in relation to habitat temperature holds at the higher ambient temperature of

32°C.

Figure 7. Bend probability in larval locomotion in the 11 *Drosophila* species at 32°C.

(A) Bend probability of the 11 species at 24°C and 32°C. The data at 24°C are the same as

in Figure 3C. Sample numbers at 32°C are the following: *Dvir*: n=25; *Dmel*: n=20; *Dmoj*:

n=29; *Dper*: n=27; *Dpse*: n=25; *Dsec*: n=20; *Dwil*: n=21; *Dere*: n=24; *Dmau*: n=17; *Dana*:

n=23; *Dyak*: n=19. P values present the results of Mann-Whitney U-test. (B) Scatter plot of

bend probability at 32°C against the minimum habitat temperature, Tmin. (C) Scatter plot of

the difference in bend probability between 32°C and at 24°C (bend probability at 32°C – bend

probability at 24°C) against the minimum habitat temperature Tmin. In B and C, median ±

sem is shown. The red lines show the linear regression functions and the shaded areas

represent the 95% confidence bands. The point estimates of the Pearson correlation and

their 95% confidence intervals are -0.850 and [-0.96, -0.51] in B and 0.40 and [-0.26, 0.81] in

C. (D-F) Examples of trajectories (left) and the bend angle (right) at 24°C (top) and 32°C

(bottom) of larval locomotion of each species (D: *Dsec*, E: *Dmel*, and F: *Dwil*).

Correlation between crawling speed and habitat temperature among *Drosophila*

species at high temperature

Next, we analyzed the crawling speed at 32°C (Figure 8A). Similar to the bend

probability, we found that the crawling speed at 32°C was correlated with the minimum habitat

temperature among the species, as it was at 24°C (Figure 8B; Pearson correlation = 0.67, p

= 0.024). To examine the effects of ambient temperature on the crawling speed in detail, we

compared the difference in the crawling speed at 24°C to 32°C among the species (Figure

8C). In contrast to the case in the bend probability, shifts in the crawling speed among the

species were common: ten species (all the species but *Dper*) showed a significant increase

in crawling speed at 32°C (Mann-Whitney U-test in Figure 8A). *Dper* also exhibited an

increase in the crawling speed but it was not statistically significant (Mann-Whitney U-test in

Figure 8A). Accordingly, at 32°C, larvae of all the species we tested increased their speed of

crawling, which might reflect a general demand to avoid malfunctions in metabolic reactions

and dehydration of the body. We also tested the relationship between the change in the

speed at 32°C and the minimum habitat temperature for each species (Figure 8C). We found

no significant correlation between the speed change and the habitat temperature (Figure 8C;

Pearson correlation = 0.07, p = 0.84), which also implies that the speeding-up at high

temperature is a general requirement among the species for the larval survival. Accordingly,

whereas the details of shifts are different (diverse shift in the bend probability and common

shift in the crawling speed among the species), the overall relationship between the kinematic

indices and minimum habitat temperature also holds at the higher ambient temperature of

32°C.

Figure 8. Crawling speed of larval locomotion in the 11 *Drosophila* species at 32°C. (A)

Crawling speed of the 11 species at 24°C and 32°C. The data at 24°C are the same as in

Figure 3D. Sample numbers at 32°C are the following: *Dvir*: n=25; *Dmel*: n=20; *Dmoj*: n=29;

Dper: n=27; *Dpse*: n=25; *Dsec*: n=20; *Dwil*: n=21; *Dere*: n=24; *Dmau*: n=17; *Dana*: n=23;

Dyak: n=19. P values present the results of Mann-Whitney U-test. (B) Scatter plot of the

crawling speed at 32°C against the minimum habitat temperature Tmin. (C) Scatter plot of

the difference of the crawling speed between at 32°C and at 24°C (crawling speed at 32°C –

crawling speed at 24°C) against the minimum habitat temperature Tmin. In B and C, median

± sem is shown. The red lines show the linear regression functions and the shaded areas

represent the 95% confidence bands. The point estimates of the Pearson correlation and

their 95% confidence intervals are 0.67 and [0.12, 0.91] in B and 0.07 and [-0.56, 0.64] in C.

A kinematic trend of larval locomotion in the genus Drosophila

To see any gross trend in the shift of larval kinematics in the different species at

distinct ambient temperatures, we plotted the kinematics of individual larvae of all the species

(Figure 9). At the ambient temperature of 24°C, a trend of kinematics was observed in which

the data points were distributed from low crawling speed / high bend probability (the top-left

corner in the plot) to high crawling speed / low bend probability (the bottom-right corner in

the plot) as the minimum habitat temperature increased (Figure 9A). Intriguingly, the

kinematics indices between distinct species were not segregated but rather overlapped. This

continuum property can also be observed in the kinematics data at the ambient temperature

of 32°C (Figure 9B). According to the modern phylogenetic concept (Markow & O'Grady,

2005), the 11 genus *Drosophila* species in our study consisted of nine subgenus *Sophophora*

species (all but *Dvir* and *Dmoj*) and two *non-Sophophora* species (*Dvir* and *Dmoj*). In our

plots of kinematics, species in the *non-Sophophora* (*Dvir* and *Dmoj*) were located at the left

side in the continuum (Figure 9). This observation implies that the kinematics indices of larvae

in *Sophophora* species in the genus *Drosophila* took values in this continuum and kinematics

indices in *non-Sophophora* species in the genus *Drosophila* have diverged along this

continuum by adaptation to habitat temperature during evolution.

Figure 9. Scatter plot of kinematics in the 11 species at 24°C and 32°C. (A and B) Two
dimensional plots of the crawling speed and bend probability. Each point corresponds to data
of a single larva of the species. The colour of markers denotes minimum habitat temperature
(Tmin) as shown in the color bar. Circles denote *Sophophora* species and squares denote
non-Sophophora species. The source locomotion data are the same as Figure 3, 7, and 8.
A: 24°C, B: 32°C.

No correlation between crawling speed and habitat temperature among *Drosophila*
species at an extreme ambient temperature of 40°C

Temperatures of 24°C and 32°C are within the range of natural habitat temperatures
of most of the species we tested (Supplementary Figure 3). We next examined larval crawling
at an extreme temperature and investigated the relation between the kinematics indices and
minimum habitat temperature. Too high a temperature can be noxious for many animals,
including *Drosophila* larvae. When *D. melanogaster* larvae are stimulated with a probe
heated to 42°C, they exhibit a stereotyped rolling motion as an escape behaviour (Dason et
al., 2019; Petersen et al., 2018; Tracey et al., 2003). At below 40°C, on the other hand, larvae

do not exhibit rolling behaviour (Petersen et al., 2018), which allows us to study crawling

kinematics in this semi-noxious extreme environment. We recorded larval locomotion of the

Drosophila species at the ambient temperature of 40°C, which corresponds to the highest

edge of maximum habitat temperature histograms for many of the species (Supplementary

Figure 3). We measured the bend probability and crawling speed of the 11 *Drosophila*

species (Supplementary Figure 5 and 6). At the ambient temperature of 40°C, neither the

bend probability nor crawling speed was correlated with minimum habitat temperature

(Figure 10A and 10B; Pearson correlation: bend probability at 40°C vs Tmin = 0.17, p = 0.61;

crawling speed at 40°C vs Tmin = -0.17, p = 0.61). Since 40°C is close to the maximum

habitat temperature rather than the minimum habitat temperature, we analyzed the relation

between the kinematics indices and maximum habitat temperature instead of minimum

habitat temperature. However, neither the bend probability nor crawling speed was correlated

with maximum habitat temperature (Pearson correlation: bend probability at 40°C vs

maximum habitat temperature = -0.13, p = 0.71; crawling speed vs maximum habitat

temperature = -0.10, p = 0.77). Accordingly, the relation between the kinematic indices and

habitat temperature does not hold at the extreme ambient temperature of 40°C. This

observation suggests that the influence of habitat temperature on the evolution of the

locomotion kinematics is restricted within a specific range of ambient temperatures. To

examine the effects of the extreme ambient temperature on the kinematics indices in detail,

we examined the shift in the bend probability at the distinct temperatures of 40°C and 32°C

among the species (Supplementary Figure 5). In seven species, bend probability increased

at 40°C. Three species that inhabit moderate temperature areas (*Dvir*, *Dper* and *Dmoj*) and

one species that lives on an isolated island (*Dsec*) showed little change in the bend probability,

which may reflect a distinct strategy in evolution to cope with the semi-noxious environment.

We also examined the shift in the crawling speed at the distinct temperatures of 40°C and

32°C among the species (Supplementary Figure 5). We found the crawling speed was

reduced at 40°C in all the species (Supplementary Figure 6), which might be a common

adaptation of the *Drosophila* larval locomotion at the extreme ambient temperature and/or

due to an abnormal physiological reaction at the semi-noxious temperature. The similar

tendency, the increase in the bend probability and the decrease in the crawling speed at

40°C, was also observed when compared with the kinematics at 24°C (Figure 10C and 10D).

Intriguingly, in some species, backward crawling, which seldom occurs at 32°C, can be

observed at 40°C (Figure 10E). Consequently, these observations show that at the extreme

ambient temperature of 40°C, the relation between the kinematics indices and habitat

temperature does not hold and larvae exhibit common (increase in the crawling speed) and

diverse (changes in the bend probability and generation of backward crawling) shifts in the

locomotion kinematics among the *Drosophila* species.

Figure 10. Larval behaviour in the 11 species at 40°C. (A) Scatter plot of the bend

probability at 40°C against the minimum habitat temperature Tmin. (B) Scatter plot of the

crawling speed at 40°C against the minimum habitat temperature Tmin. (C) Scatter plot of

the difference of the bend probability between at 40°C and at 24°C (bend probability at 40°C

– bend probability at 24°C) against the minimum habitat temperature Tmin. (D) Scatter plot

of the difference of the crawling speed between at 40°C and at 24°C (crawling speed at 40°C

– crawling speed at 24°C) against the minimum habitat temperature Tmin. Median ± sem is

shown in A - D. Sample numbers for data at 40°C are the following: *Dvir*: n=20; *Dmel*: n=28;

Dmoj: n=23; *Dper*: n=21; *Dpse*: n=16; *Dsec*: n=27; *Dwil*: n=27; *Dere*: n=22; *Dmau*: n=26;

Dana: n=23; *Dyak*: n=20. Sample numbers for data at 24°C are the same as Figure 3C and

3D. The red lines show the linear regression functions and the shaded areas represent the

95% confidence bands. The point estimates of the Pearson correlation and their 95%

confidence intervals are 0.17 and [-0.48, 0.70] in A and -0.17, [-0.70, 0.48] in B, 0.17, [-0.48,

0.70] in C, and -0.50, [-0.84, 0.15] in D. (E) Backward crawling probability at 40°C of each

species. Probability $\pm$ standard error based on the binomial distribution are shown. Sample

numbers in E are the following: *Dvir*: n=20; *Dmel*: n=28; *Dmoj*: n=23; *Dper*: n=22; *Dpse*:

n=14; *Dsec*: n=27; *Dwil*: n=27; *Dere*: n=23; *Dmau*: n=26; *Dana*: n=23; *Dyak*: n=21.

Phylogenetic analyses of interspecies differences in kinematics

542   Finally, to obtain a hint on how the kinematics parameters diversified across the

species have developed during the evolutionary history of the genus *Drosophila*, we

conducted Bayesian phylogenetic analyses using RevBayes v. 1.0.7 (Hohna et al., 2016;

Höhna et al., 2017). At first, we inferred phylogenetic trees of the 11 *Drosophila* species

based on eight nuclear genes used previously (Turelli et al., 2018) (See Methods for detail).

Then by using our kinematics dataset, we estimated the following three factors in the

phylogenetic tree by Bayesian inference: the rate of evolution at each branch of the

phylogenetic trees, relative rates of evolution among the kinematic parameters, and the

correlation between the kinematic parameter evolutions (Kalay et al., 2020). To perform the

inference, we constructed a data matrix of eight parameters (the bend probability at 24, 32

and 40°C, the crawling speed at 24, 32 and 40°C, backward crawling probability at 40°C and

the body length), for each of the 11 *Drosophila* species. We assumed that these parameters

evolve under a multivariate Brownian-motion model (Huelsenbeck & Bruce, 2003; Kalay et

al., 2020; Lartillot & Poujol, 2011) (See Methods for detail). We estimated the three factors

(the evolution rates at phylogenetic trees, relative evolution rates among the kinematic

parameters, and the correlation between the parameter evolutions) by running a Markov

Chain Monte Carlo simulation.

The phylogenetic analyses indicated that the rates of evolution in the kinematics are

highly diverse over branches in the phylogenetic tree (Figure 11A and Supplementary Figure

7). In addition, the evolutionary rates are distinct among the eight kinematics parameters

(Figure 11B). The bend probability at 24°C and the probability of backward crawling at 40°C

have a relatively high evolution rate, which might reflect large diversification of these two

parameters among the species (Figure 9A and 10E).

Some kinematics traits might have evolved cooperatively. To test this possibility, we

calculated the correlation of the evolutionary changes between the kinematic parameters. In

the correlation analyses of all pairs of the parameters, the crawling speed at 24°C and the

crawling speed at 32°C were the most correlated (Figure 11C). On the other hand, the

evolutions of the bend probability between at 24°C and 32°C were less correlated. This

observation is consistent with the notion that while changes in the bend probability from 24°C

to 32°C among the species are diverse (Figure 7C), all the species show an increase in the

crawling speed by the temperature shift (Figure 8C). Consequently, the phylogenetic

analyses imply that the kinematics indices of larval locomotion have evolved differently in

distinct branches of a phylogenetic tree with keeping a correlation between specific

locomotion traits such as crawling speed at distinct temperatures.

Figure 11. Phylogenetic analyses of larval locomotion in the genus *Drosophila*. (A)

Phylogenetic tree estimated from Bayesian inference. The topology is inferred by coding

sequences of eight genes of the 11 *Drosophila* species. As a prior distribution for the

Bayesian inference, an uncorrelated exponential (UCED) relaxed-clock model was used.

Colour and a value of each branch denote the relative rate of the evolution of the eight

kinematic traits. (B) Relative evolution rates of the eight kinematics traits estimated by the

Bayesian inference. Mean $\pm$ 95% highest posterior density is shown. (C) Correlation

between the evolution of eight traits. Numbers represent the range of 95% highest posterior

density. Abbreviations in (B) and (C) are the following: Spd24: crawling speed at 24°C,

Spd32: crawling speed at 32°C, Spd40: crawling speed at 40°C, Bend24: bend probability at

24°C, Bend32: bend probability at 32°C, Bend40: bend probability at 40°C, Bwd: probability

of backward crawling, Length: axial body length of larvae.

Discussion

In this work, we investigated interspecies differences in larval locomotion in the

genus *Drosophila*. We used the bend probability and crawling speed as measures to examine

larval locomotion. Despite the similar appearance of larval bodies in the different species, the

kinematics of larval locomotion is diverged (Figure 2). The body length is not a leading factor

for the diversity of kinematics (Supplementary Figure 2). Phylogenetic relationship is also not

a major determinant for the kinematics (Figure 3E and 11A). Considering a previous study

showing that phylogenetic relationship does not correlate with the divergence in the

morphology of larval neuromuscular junctions (Campbell & Ganetzky, 2012), genetic drift with

random accumulation of neutral mutations is unlikely to underlie the divergence in the larval

crawling patterns. In contrast, habitat temperature correlates with both the bend probability

and crawling speed at both 24°C and 32°C ambient temperature (Figure 12), which implies

the kinematics indices are adapted to ambient temperature in evolution. Phylogenetic

analysis by Bayesian inference suggests that the rates of evolution are divergent among the

branches of the phylogenetic tree (Figure 11A). Among the kinematic parameters, the bend

probability at the ambient temperature of 24°C and the backward crawling probability at 40°C

have relatively higher evolution rates than the others and the evolution of the crawling speed

at 24°C and 32°C are correlated (Figure 11B and 11C). Regarding the questions raised in

the Introduction section, our results suggest the following: (1) Locomotion kinematics of

larvae is divergent among *Drosophila* species. (2) The habitat temperature is more related

to the kinematics indices (the bend probability and crawling speed) than the body length

(Figure 12).

Figure 12. Summary of this study. Disks and triangles denote the kinematics of each

species at 24°C and 32°C, respectively.

To conduct the quantitative analyses, we measured the larval crawling behaviour in

simple experimental conditions: at the constant temperatures and humidity levels and on a

flat surface of agarose gel. However, the environmental conditions in the wild for *Drosophila*

larvae are far more complicated and diverse (Markow, 2015). For example, *Dmoj* breeds in

cacti in the desert while *Dvir* inhabits in the slime fluxes in temperate and subarctic climate

(Supplementary Figure 3; (Markow, 2015)). Our data in this paper imply that the variation of

the kinematics should be related to the diversity of environmental conditions including habitat

temperature. Comprehensive and quantitative analyses of larval locomotion in nature would

shed light on the causes and meanings of the changes in larval kinematics appeared in the

evolution.

The correlation between habitat temperature and the kinematics raises a question

about the underlying mechanisms of adaptive change in kinematics to the habitat

temperature. We presume that the adaptive change was likely to be driven by two factors:

the pupation positioning and competition over food. Regarding the pupation behaviour, a

previous study showed a correlation between path lengths of crawling and heights of

pupation location in *D. melanogaster* (Sokolowski, 1985). Since pupae are immobile,

adequate selection of pupation position is vital for their survival. Moisture is one of the key

factors that affects the selection of pupation site, since dry environments can cause pupal

desiccation, while soaked media might lead to the drowning of pupae (Sokolowski, 1985;

Sokolowski et al., 1984). It is reasonable to assume that habitat temperature is a leading

factor in determining the moisture levels of microenvironments of *Drosophila* larvae.

Accordingly, divergent temperatures in the habitats of flies leads to variability in moisture

levels, the ambient moisture affects the pupation position (Sokolowski et al., 1984), and the

pupation position is related to the path length of crawling (Sokolowski, 1985). These three

links might underlie the relationship between the larval locomotion kinematics and habitat

temperature of each species.

The second possible mechanism underlying the relation between habitat

temperature and locomotion kinematics is related to competition over food. It is suggested

that larval feeding is competitive between individuals (Rohlfs & Hoffmeister, 2003). There is

a positive relation between fitness and density (called Allee effects) in *D. melanogaster* larvae

(Wertheim et al., 2002) and so female flies aggregate eggs when laying them (Rohlfs &

Hoffmeister, 2003). Consequently, the larval population tends to be overcrowded (Burnet et

al., 1977) and larvae compete over the limited resources to meet their need to consume

sufficient amount of nutrient within limited larval periods (Nicholson, 1954). If the available

food is limited and larval density is high, larvae need to crawl further (Sokolowski et al., 1997),

which can be an evolutionary driving force to increase the crawling distance. In contrast, if

there is plenty of food in the habitat, the crawling distance remains unchanged or even

decreases during evolution, because crawling behaviour is energetically costly (Berrigan &

Lighton, 1993). Consistent with this implication, the diversification in the larval kinematics in

the 11 *Drosophila* species gives rise to the variation in the crawling distance (Figure 5).

Accordingly, feeding conditions, especially the choice of which foods to eat, can affect

kinematics of larval locomotion in evolutionary processes so that the divergence of food to

eat can lead to the divergence in locomotion kinematics. In regard to this point, food for larvae

of the genus *Drosophila* is divergent and related to their living environments (Markow, 2015;

Watada et al., 2020; Watanabe et al., 2019). So, the nutritional conditions (nutrient balances

and fermentation etc.) and physical properties (hardness and wetness etc.) of food to eat

vary among the *Drosophila* species, which can drive evolutionary divergence in the

kinematics of larval locomotion. Then, the growth of plants, including fruits and vegetables,

for larvae is strongly affected by habitat temperature. Therefore, divergence in habitat

temperature may affect the locomotion kinematics during evolution through divergence in the

foods that larvae feed on, and divergence in requisite feeding behaviour for larval growth.

Comprehensive kinematics studies of other *Drosophila* species and quantitative analyses of

microenvironments of wild larvae in nature will give us insights on the relationships between

ambient temperature and diverse larval locomotion kinematics.

How can we approach the neural circuit mechanisms underlying the interspecies

divergence in larval locomotion? Circuit mechanisms in larval locomotion have been

examined intensively in *Drosophila melanogaster*. Recent connectomics studies have

identified several key neurons for larval locomotion in *Drosophila melanogaster* (Carreira-

Rosario et al., 2018; M. Q. Clark et al., 2018; Fushiki et al., 2016; Heckscher et al., 2015;

Hughes & Thomas, 2007; Kohsaka et al., 2014, 2017, 2019; Schneider-Mizell et al., 2016;

Tastekin et al., 2015; Zwart et al., 2016). Regarding bending, the thoracic neuromere was

shown to be important in bending in chemotaxis (Tastekin et al., 2015). Especially, the

commissural connection is crucial for bending (Berni, 2015). In addition, a signal from the

chordotonal sensory organ is also required to generate bending (Ainsley et al., 2003).

Regarding the crawling speed, a group of inhibitory premotor neurons were identified to be

critical (Kohsaka et al., 2014). Proprioceptive feedback and neuromodulation are both

important for the normal crawling speed (Fox et al., 2006; Hughes & Thomas, 2007).

Furthermore, kinematics of the larval locomotion have been measured and investigated

quantitatively in detail (Gepner et al., 2015; Gershow et al., 2012; Gomez-Marin et al., 2011;

Gomez-Marin & Louis, 2014; Heckscher et al., 2012; Hernandez-Nunez et al., 2015; Lahiri

et al., 2011; Luo et al., 2010) and analysed by mathematical modelling (Gjorgjieva et al.,

2013; Loveless et al., 2019; Loveless & Webb, 2018; Pehlevan et al., 2016; Sun et al., 2020).

Regarding temperature sensing, neuronal and molecular mechanisms on temperature-

guided behaviour have been clarified (Dillon et al., 2009; Klein et al., 2015; Kwon et al., 2008,

2010; Liu et al., 2003; Ni et al., 2016; Rosenzweig et al., 2005b, 2008; Shen et al., 2011).

These extensive findings on the cellular and molecular mechanisms on larval crawling in

Drosophila melanogaster will be an ideal starting point to investigate the evolution of larval

behaviour in the genus *Drosophila*. For example, differences in commissure fibre tracts in

the central nervous system (Berni, 2015) among the species might underlie the divergence

in the bend probability. Interspecies comparison of a group of interneurons PMSIs (*period-*

positive median segmental interneurons), which are involved in the crawling speed (Kohsaka

et al., 2014), would reveal neural mechanisms on the evolutionary diversification in the

crawling speed. Comparative analyses of neural network architectures and gene expression

among the *Drosophila* species will relate the evolution of the nervous system to the adaptive

diversification in larval behaviour.

Materials and Methods

***Drosophila* strains**

708   We used the following fly stocks (17 strains from 11 species): *Drosophila ananassae* (k-s01), *Drosophila erecta* (k-s02), *Drosophila yakuba* (k-s03), *Drosophila melanogaster* (k-s04), *Drosophila sechellia* (k-s10), *Drosophila persimilis* (k-s11), *Drosophila pseudoobscura* (k-s12), *Drosophila willistoni* (k-s13), *Drosophila virilis* (k-s14) and *Drosophila mojavensis* (k-s15), *Drosophila melanogaster* collected at Kyoto (k-aba029) and Iriomote (k-aba032), *Drosophila ananassae* collected at Ogasawara (k-aaa027) and Iriomote (k-aaa309), and *Drosophila virilis* collected at Hokkaido (E-15601) and Toyama (E-15605) from KYORIN-Fly, Fly Stocks of Kyorin University and *Drosophila mauritiana* (#900020) from KYOTO Stock Center (DGRC) at the Kyoto Institute of Technology. These strains have been maintained at 23°C (except for *Drosophila persimilis* at 20°C) in the stock centres for more than 10 years. All animals were raised on standard cornmeal-based food at 25°C in the authors' laboratory before the experiments for less than three months after the acquisition from the stock centres.

Recording larval locomotion

722   Third-instar wandering larvae of each strain were picked up and gently washed in deionised water. Residual food on the larvae was brushed off with a paintbrush. An agarose stage (size: 9cm x 9cm x 5mm; 1.5% agarose; RIKAKEN STAR agarose powder #RSV-AGRP-100G) was placed on a temperature-controlled plate (Cool Plate, AS ONE Corporation, Japan) and the temperature of the surface was kept at 24±1°C, 32±1°C or 40±1°C. Eight to ten larvae were placed on the agarose arena with a paintbrush. The larvae were illuminated with infrared light, which larvae cannot see (LDQ-150IR2-850, wavelength = 850 nm, CCS Inc., Japan) and recorded at five frames/sec with a CCD camera (CGE-B013-U, MIGHTEX, Canada; resolution: 1280 pixels x 960 pixels) and an infrared filter (775LP filter, Omega

Optical, USA), and saved as a series of bitmap files. In our setting, the scale of image is 0.13

mm/pixel. For each of the 17 strains (from 11 species) and at each of the three ambient

temperatures, we repeated the measurement three times.

Larva tracking

We obtained a time series of bitmap image files by the procedure described above.

The bitmap files were converted to tiff format by Fiji (https://imagej.net/fiji). In cases where

larvae accumulated at the first frame, we removed the first several images until the larvae

dispersed enough to be identified individually. The x and y coordinates of the centroid and

the bend angle (Figure 1E) of individual larvae were obtained by using the FIMTrack software

(Risse et al., 2017). We used tracking data of single larvae that were continuously tracked

for more than 300 frames (which corresponds to one minute). For the intraspecific analysis

in Figure 4, single larvae with more than 150 continuous frames were tracked. In case a larva

collided with another larva in the middle of the recording and the larva was traced differently

before and after the collision with distinct labels, we treated the two traces as two larvae.

Crawling speed of each larva was obtained as the median of crawling speed in the whole

trace of the larva. Bend probability of each larva was calculated as the ratio of the number of

frames showing bending (Figure 3 for the definition of bending) to the total number of the

frames of its trace. The data of the three movies for each species/strain were merged. The

coordinates and bend angles were smoothed and plotted by Python 3.7. For the visualization

of the traces in Figure 1E and 1F, we smoothed the data by a uniform window function with

five frames. The number of pixels in single larva ranges from about 150 (*Dyak*) to 500 (*Dpse*)

that is sufficient to capture the body bend angles.

Clustering analysis of kinematics

For clustering analysis of the plots in Figure 2A, we made a probability distribution.

We discretised the centroid speed axis by an interval of 0.1 mm/sec between 0 and 4

mm/sec and the bend angle axis by an interval of two degrees between 0 and 140 degrees,

then created 2800 bins in total (40 bins in the speed axis and 70 bins in the bend angle

axis). For each scatter plot, we counted the number of points in each bin and obtained

probability density by normalisation with the total number of the points. We calculated the

Kullback-Leibler divergence (KL) of probability distributions of two species p(bin) and q(bin)

as:

764
$$KL(p, q) = \sum_{bin=1}^{2800} p(bin) * \log \frac{p(bin)}{q(bin)}.$$

765   Since KL(p,q) is an asymmetric measure $(KL(p, q) \neq KL(q, p))$, we calculated the Jensen-

766   Shannon divergence JS, which is a symmetric alternative to the Kullback-Leibler

767   divergence:

768
$$JS(p, q) = \frac{1}{2} \{KL(p, q) + KL(q, p)\} = JS(q, p).$$

Based on the Jensen-Shannon divergence, we conducted the hierarchical clustering using

Python 3.7 and the SciPy library.

Habitat temperature of the *Drosophila* species

Global climate data were obtained from the WorldClim data website

(https://www.worldclim.org/data/worldclim21.html). We used minimum, mean and maximum

temperature in the world at a spatial resolution of five minutes (9.25 km x 9.25 km). Habitat

regions of each *Drosophila* species were obtained from the literature (Brake & Baechli, 2008;

Makino & Kawata, 2012; Markow & O'Grady, 2005) and the DrosWLD-Species database

(https://bioinfo.museum.hokudai.ac.jp/db/modules/stdb/index.php?action=tbl&tbl_id=21)

and traced them on the world map (2160 pixel x 4320 pixel) obtained from the WorldClim by

using Adobe Illustrator software (Adobe Inc., USA). The maps in Supplementary Figure 2

were generated by Adobe Illustrator and the statistics of the temperature within each species'

habitat were calculated by Python 3.7.

Statistical analysis

Statistical calculation was conducted by Python 3.7 in the following analyses: Kruskal-Walls test in Figure 3C and 3D, Pearson correlation in Figure 6B, 6C, 7B, 7C, 8B and 8C, Supplementary Figure 2B, 2C, 4B and 5B, and Mann-Whitney U-test in Figure 7A and 8A and Supplementary Figure 5A and 6A.

Phylogenetic analyses

We adopted Bayesian statistics for phylogenetic analyses. In all the phylogenetic analyses, we used the software package, RevBayes v.1.0.7 (Hohna et al., 2016).

We estimated the *Drosophila* phylogenetic trees (Figure 8A) by using eight nuclear loci from Turelli et al. (2018): *aldolase*, *bicoid*, *enolase*, *esc*, *transaldolase*, *white*, *wingless*, and *yellow*. We used *Scaptodrosophila lebanonensis* (*S. lebanonensis*) as an outgroup. Coding sequences of these genes for all the species except for *D. mauritiana* and *S. lebanonensis* were obtained from Kalay et al., (2020). Coding sequences of the genes of *D. mauritiana* and *S. lebanonensis* were obtained from the NCBI web site (https://www.ncbi.nlm.nih.gov/). The genes were aligned with MAFFT version 7 (Katoh & Standley, 2013). We selected the eight genes out of twenty genes described in Turelli et al., (2018) because homologous genes of the other twelve genes were not identified in the genome assembly of *D. mauritiana*. Based on the eight coding sequences of the twelve species, we performed chronogram analyses described in Turelli et al. (2018) and obtained the phylogenetic tree in Figure 8A, where the root is the branch of *S. lebanonensis* and the length of each branch indicates relative time in the evolution. We repeated the phylogeny inference four times to obtain four trees to check the robustness of this inference in the following analysis.

Based on the phylogenetic tree obtained above, we analyzed the evolution of kinematics parameters. We used eight parameters (bend probability at 24, 32 and 40°C, crawling speed at 24, 32 and 40°C, backward crawling probability at 40°C, and body length),

for each of the 11 *Drosophila* species. In this analysis, while the topology of the phylogenetic

tree is unchanged, the rates of evolution of the kinematic parameters and the rates of

evolution of the branches of the phylogenetic tree were estimated by Bayesian inference. We

assumed the kinematics parameters evolve under a multivariate Brownian-motion model

(Huelsenbeck & Bruce, 2003; Lartillot & Poujol, 2011). This model consists of two

components: the relative rates of evolution among the kinematics parameters and the

correlation between each pair of the kinematics parameters (Kalay et al., 2020). The

Brownian model assumes that the changes in the kinematics parameters during evolution

are additive, which has two consequences: 1) parameters can become negative; 2) evolution

rates can depend on the scale of the kinematics parameters. To cope with these properties,

we adopted log-transformation to the kinematics parameters as described previously (Kalay

et al., 2020). This transformation guarantees that the original kinematics parameters (which

are exponentials of the log-transformation) remain positive and controls the size issue by

transforming additive changes to multiplicative ones (Kalay et al., 2020). For prior models for

the evolution rates of the branches, we specified an uncorrelated exponential (UCED)

relaxed-clock model, where the rate of each branch is drawn independently from an

exponential distribution. We ran the Markov Chain Monte Carlo (MCMC) simulation of UCED

with each of the four phylogenetic trees described in the previous paragraph (Figure 11).

Consistent results were obtained by another prior model, an uncorrelated gamma (UCG)

relaxed clock model (Supplementary Figure 7), which indicates the predictions are robust to

the choice of priors.

Acknowledgement

We thank KYORIN-Fly, Fly Stocks of Kyorin University and KYOTO Stock Center (DGRC) at

the Kyoto Institute of Technology for providing fly lines, Dr. Takashi Makino for advice about

climate data in the world. We thank Ms. Shibahara for tracing habitats of the 11 species. We

thank Dr. Masayoshi Watada for critical comments on the paper. This work was supported

by MEXT/JSPS KAKENHI grants (17K19439, 19H04742, 20H05048 to A.N. and 17K07042,

20K06908 to H.K.).

Competing interests

We have no conflicts of interest with respect to the work.

References

Ainsley, J., Pettus, J., Bosenko, D., Gerstein, C., Zinkevich, N., Anderson, M., Adams, C.,
Welsh, M., & Johnson, W. (2003). Enhanced locomotion caused by loss of the Drosophila
DEG/ENaC protein Pickpocket1. *Current Biology*, *13*, 1557–1563.
https://doi.org/10.1016/S
Alexander, R. M. (2006). *Principles of Animal Locomotion*. Princeton University Press.
Berni, J. (2015). Genetic dissection of a regionally differentiated network for exploratory
behavior in drosophila larvae. *Current Biology*, *25*(10), 1319–1326.
https://doi.org/10.1016/j.cub.2015.03.023
Berni, J., Pulver, S. R., Griffith, L. C., & Bate, M. (2012). Autonomous circuitry for substrate
exploration in freely moving drosophila larvae. *Current Biology*, *22*(20), 1861–1870.
https://doi.org/10.1016/j.cub.2012.07.048
Berrigan, D., & Lighton, J. R. (1993). Bioenergetic and kinematic consequences of limblessness
in larval Diptera. *The Journal of Experimental Biology*, *179*, 245–259.
Berrigan, D., & Pepin, D. J. (1995). How maggots move: Allometry and kinematics of crawling in
larval Diptera. *Journal of Insect Physiology*, *41*(4), 329–337. https://doi.org/10.1016/0022-
1910(94)00113-U
Brake, I., & Baechli, G. (2008). *Drosophilidae (Diptera)*. Apollo Books.
Burnet, B., Sewell, D., & Bos, M. (1977). Genetic analysis of larval feeding behaviour in
Drosophila melanogaster: II. Growth relations and competition between selected lines.
Genetical Research, *30*(2), 149–161. https://doi.org/10.1017/S0016672300017559
Campbell, M., & Ganetzky, B. (2012). Extensive morphological divergence and rapid evolution
of the larval neuromuscular junction in Drosophila. *Proceedings of the National Academy
of Sciences of the United States of America*, *109*(11).
https://doi.org/10.1073/pnas.1201176109
Carreira-Rosario, A., Zarin, A. A., Clark, M. Q., Manning, L., Fetter, R. D., Cardona, A., & Doe,
C. Q. (2018). MDN brain descending neurons coordinately activate backward and inhibit
forward locomotion. *ELife*, *7*, 1–28. https://doi.org/10.7554/eLife.38554
Clark, A. G., Eisen, M. B., Smith, D. R., Bergman, C. M., Oliver, B., Markow, T. A., Kaufman, T.
C., Kellis, M., Gelbart, W., Iyer, V. N., Pollard, D. A., Sackton, T. B., Larracuente, A. M.,
Singh, N. D., Abad, J. P., Abt, D. N., Adryan, B., Aguade, M., Akashi, H., … MacCallum, I.
(2007). Evolution of genes and genomes on the Drosophila phylogeny. *Nature*, *450*(7167),
203–218. https://doi.org/10.1038/nature06341
Clark, M. Q., Zarin, A. A., Carreira-Rosario, A., & Doe, C. Q. (2018). Neural circuits driving larval
locomotion in Drosophila. *Neural Development*, *13*(1), 1–10.

https://doi.org/10.1186/s13064-018-0103-z

Dason, J. S., Cheung, A., Anreiter, I., Montemurri, V. A., Allen, A. M., & Sokolowski, M. B.
(2019).   Drosophila melanogaster foraging regulates a nociceptive-like escape behavior
through a developmentally plastic sensory circuit . *Proceedings of the National Academy*
of Sciences, 201820840. https://doi.org/10.1073/pnas.1820840116

Dillon, M. E., Wang, G., Garrity, P. A., & Huey, R. B. (2009). Thermal preference in Drosophila.
Journal of Thermal Biology, *34*(3), 109–119. https://doi.org/10.1016/j.jtherbio.2008.11.007

Fox, L. E., Soll, D. R., & Wu, C. F. (2006). Coordination and modulation of locomotion pattern
generators in Drosophila larvae: Effects of altered biogenic amine levels by the tyramine β
hydroxlyase mutation. *Journal of Neuroscience*, *26*(5), 1486–1498.
https://doi.org/10.1523/JNEUROSCI.4749-05.2006

Fuller, P. O., Higham, T. E., & Clark, A. J. (2011). Posture, speed, and habitat structure: Three-
dimensional hindlimb kinematics of two species of padless geckos. *Zoology*, *114*(2), 104–
112. https://doi.org/10.1016/j.zool.2010.11.003

Fushiki, A., Zwart, M. F., Kohsaka, H., Fetter, R. D., Cardona, A., & Nose, A. (2016). A circuit
mechanism for the propagation of waves of muscle contraction in Drosophila. *ELife*, *5*, 1–
23. https://doi.org/10.7554/elife.13253

Gepner, R., Skanata, M. M., Bernat, N. M., Kaplow, M., & Gershow, M. (2015). Computations
underlying Drosophila photo- taxis, odor-taxis, and multi-sensory integration. *ELife*,
4(MAY), 1–21. https://doi.org/10.7554/eLife.06229

Gershow, M., Berck, M., Mathew, D., Luo, L., Kane, E. A., Carlson, J. R., & Samuel, A. D. T.
(2012). Controlling airborne cues to study small animal navigation. *Nature Methods*, *9*(3),
290–296. https://doi.org/10.1038/nmeth.1853

Gjorgjieva, J., Berni, J., Evers, J. F., & Eglen, S. J. (2013). Neural circuits for peristaltic wave
propagation in crawling Drosophila larvae: analysis and modeling. *Frontiers in*
Computational Neuroscience, *7*(April), 1–19. https://doi.org/10.3389/fncom.2013.00024

Gomez-Marin, A., & Louis, M. (2014). Multilevel control of run orientation in Drosophila larval
chemotaxis. *Frontiers in Behavioral Neuroscience*, *8*(FEB), 1–14.
https://doi.org/10.3389/fnbeh.2014.00038

Gomez-Marin, A., Stephens, G. J., & Louis, M. (2011). Active sampling and decision making in
Drosophila chemotaxis. *Nature Communications*, *2*(1).
https://doi.org/10.1038/ncomms1455

Green, C. H., Burnet, B., & Connolly, K. J. (1983). Organization and patterns of inter- and
intraspecific variation in the behaviour of Drosophila larvae. *Animal Behaviour*, *31*(1), 282–
291. https://doi.org/10.1016/S0003-3472(83)80198-5

Heckscher, E. S., Lockery, S. R., & Doe, C. Q. (2012). Characterization of Drosophila larval
crawling at the level of organism, segment, and somatic body wall musculature. *Journal of*
Neuroscience, *32*(36), 12460–12471. https://doi.org/10.1523/JNEUROSCI.0222-12.2012

Heckscher, E. S., Zarin, A. A., Faumont, S., Clark, M. Q., Manning, L., Fushiki, A., Schneider-
Mizell, C. M., Fetter, R. D., Truman, J. W., Zwart, M. F., Landgraf, M., Cardona, A.,
Lockery, S. R., & Doe, C. Q. (2015). Even-Skipped+ Interneurons Are Core Components
of a Sensorimotor Circuit that Maintains Left-Right Symmetric Muscle Contraction
Amplitude. *Neuron*, *88*(2), 314–329. https://doi.org/10.1016/j.neuron.2015.09.009

Hernandez-Nunez, L., Belina, J., Klein, M., Si, G., Claus, L., Carlson, J. R., & Samuel, A. D. T.
(2015). Reverse-correlation analysis of navigation dynamics in Drosophila larva using
optogenetics. *ELife*, *4*(MAY), 1–16. https://doi.org/10.7554/eLife.06225

Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high
resolution interpolated climate surfaces for global land areas. *International Journal of*
Climatology, *25*(15), 1965–1978. https://doi.org/10.1002/joc.1276

Höhna, S., Landis, M. J., & Heath, T. A. (2017). Phylogenetic inference using RevBayes.
Current Protocols in Bioinformatics, *2017*(March), 6.16.1-6.16.34.
https://doi.org/10.1002/cpbi.22

Hohna, S., Landis, M. J., Heath, T. A., Boussau, B., Lartillot, N., Moore, B. R., Huelsenbeck, J.
P., & Ronquist, F. (2016). RevBayes: Bayesian phylogenetic inference using graphical

936   models and an interactive model-specification language. *Systematic Biology*, *65*(4), 726–
937   736. https://doi.org/10.1093/sysbio/syw021

938 Huelsenbeck, J., & Bruce, R. (2003). Detecting Correlation between Characters in a
939   Comparative Analysis with Uncertain Phylogeny. *Evolution*, *57*(6), 1237–1247.

940 Hughes, C. L., & Thomas, J. B. (2007). A sensory feedback circuit coordinates muscle activity in
941   Drosophila. *Molecular and Cellular Neuroscience*, *35*(2), 383–396.
942   https://doi.org/10.1016/j.mcn.2007.04.001

943 Kalay, G., Atallah, J., Sierra, N. C., Tang, A. M., Crofton, A. E., Murugesan, M. K., Wykoff-Clary,
944   S., & Lott, S. E. (2020). Evolution of larval segment position across 12 Drosophila
945   species*. *Evolution*, *74*(7), 1409–1422. https://doi.org/10.1111/evo.13911

946 Kappeler, P. (2010). *Animal Behaviour: Evolution and Mechanisms*. Springer.

947 Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7:
948   Improvements in performance and usability. *Molecular Biology and Evolution*, *30*(4), 772–
949   780. https://doi.org/10.1093/molbev/mst010

950 Klein, M., Afonso, B., Vonner, A. J., Hernandez-Nunez, L., Berck, M., Tabone, C. J., Kane, E.
951   A., Pieribone, V. A., Nitabach, M. N., Cardona, A., Zlatic, M., Sprecher, S. G., Gershow,
952   M., Garrity, P. A., Samuel, A. D. T., & Sternberg, P. W. (2015). Sensory determinants of
953   behavioral dynamics in Drosophila thermotaxis. *Proceedings of the National Academy of*
954   *Sciences of the United States of America*, *112*(2), E220–E229.
955   https://doi.org/10.1073/pnas.1416212112

956 Kohsaka, H., Guertin, P. A., & Nose, A. (2017). Neural Circuits Underlying Fly Larval
957   Locomotion. *Current Pharmaceutical Design*, *23*(12), 1722–1733.
958   https://doi.org/10.2174/1381612822666161208120835

959 Kohsaka, H., Takasu, E., Morimoto, T., & Nose, A. (2014). A group of segmental premotor
960   interneurons regulates the speed of axial locomotion in drosophila larvae. *Current Biology*,
961   *24*(22), 2632–2642. https://doi.org/10.1016/j.cub.2014.09.026

962 Kohsaka, H., Zwart, M. F., Fushiki, A., Fetter, R. D., Truman, J. W., Cardona, A., & Nose, A.
963   (2019). Regulation of forward and backward locomotion through intersegmental feedback
964   circuits in Drosophila larvae. *Nature Communications*, *10*(1), 2654.
965   https://doi.org/10.1038/s41467-019-10695-y

966 Kwon, Y., Shen, W. L., Shim, H. S., & Montell, C. (2010). Fine thermotactic discrimination
967   between the optimal and slightly cooler temperatures via a TRPV channel in chordotonal
968   neurons. *Journal of Neuroscience*, *30*(31), 10465–10471.
969   https://doi.org/10.1523/JNEUROSCI.1631-10.2010

970 Kwon, Y., Shim, H. S., Wang, X., & Montell, C. (2008). Control of thermotactic behavior via
971   coupling of a TRP channel to a phospholipase C signaling cascade. *Nature Neuroscience*,
972   *11*(8), 871–873. https://doi.org/10.1038/nn.2170

973 Lahiri, S., Shen, K., Klein, M., Tang, A., Kane, E., Gershow, M., Garrity, P., & Samuel, A. D. T.
974   (2011). Two alternating motor programs drive navigation in Drosophila larva. *PLoS ONE*,
975   *6*(8). https://doi.org/10.1371/journal.pone.0023180

976 Lartillot, N., & Poujol, R. (2011). A phylogenetic model for investigating correlated evolution of
977   substitution rates and continuous phenotypic characters. *Molecular Biology and Evolution*,
978   *28*(1), 729–744. https://doi.org/10.1093/molbev/msq244

979 Lin, J. (1991). Divergence Measures Based on the Shannon Entropy. *IEEE Transactions on*
980   *Information Theory*, *37*(1), 145–151. https://doi.org/10.1109/18.61115

981 Liu, L., Yermolaieva, O., Johnson, W. A., Abboud, F. M., & Welsh, M. J. (2003). Identification
982   and function of thermosensory neurons in Drosophila larvae. *Nature Neuroscience*, *6*(3),
983   267–273. https://doi.org/10.1038/nn1009

984 Loveless, J., Lagogiannis, K., & Webb, B. (2019). Modelling the neuromechanics of exploration
985   and taxis in larval Drosophila. *PLoS Computational Biology*, 1–33.
986   https://doi.org/10.5281/zenodo.1432637

987 Loveless, J., & Webb, B. (2018). A Neuromechanical Model of Larval Chemotaxis. *Integrative*
988   *and Comparative Biology*, *58*(5), 906–914. https://doi.org/10.1093/icb/icy094

989 Luo, L., Gershow, M., Rosenzweig, M., Kang, K., Fang-Yen, C., Garrity, P. A., & Samuel, A. D.

T. (2010). Navigational decision making in Drosophila thermotaxis. *Journal of Neuroscience*, *30*(12), 4261–4272. https://doi.org/10.1523/JNEUROSCI.4090-09.2010

Makino, T., & Kawata, M. (2012). Habitat variability correlates with duplicate content of drosophila genomes. *Molecular Biology and Evolution*, *29*(10), 3169–3179. https://doi.org/10.1093/molbev/mss133

Markow. (2015). The secret lives of Drosophila flies. *ELife*, *4*(JUNE), 1–9. https://doi.org/10.7554/eLife.06793

Markow, T. A., & O'Grady, P. M. (2005). *Drosophila: A guide to species identification and use*. Academic Press.

Miller, D. E., Staber, C., Zeitlinger, J., & Hawley, R. S. (2018). Highly contiguous genome assemblies of 15 drosophila species generated using nanopore sequencing. *G3: Genes, Genomes, Genetics*, *8*(10), 3131–3141. https://doi.org/10.1534/g3.118.200160

Ni, L., Klein, M., Svec, K. V., Budelli, G., Chang, E. C., Ferrer, A. J., Benton, R., Samuel, A. D. T., & Garrity, P. A. (2016). The ionotropic receptors IR21a and IR25a mediate cool sensing in Drosophila. *ELife*, *5*(APRIL2016). https://doi.org/10.7554/eLife.13254

Nicholson, A. J. (1954). An outline of the dynamics of animal. *Australian Journal of Zoology*, *2*(1), 9–65.

Ohyama, T., Jovanic, T., Denisov, G., Dang, T. C., Hoffmann, D., Kerr, R. A., & Zlatic, M. (2013). High-Throughput Analysis of Stimulus-Evoked Behaviors in Drosophila Larva Reveals Multiple Modality-Specific Escape Strategies. *PLoS ONE*, *8*(8). https://doi.org/10.1371/journal.pone.0071706

Ohyama, T., Schneider-Mizell, C. M., Fetter, R. D., Aleman, J. V., Franconville, R., Rivera-Alba, M., Mensh, B. D., Branson, K. M., Simpson, J. H., Truman, J. W., Cardona, A., & Zlatic, M. (2015). A multilevel multimodal circuit enhances action selection in Drosophila. *Nature*, *520*(7549), 633–639. https://doi.org/10.1038/nature14297

Pehlevan, C., Paoletti, P., & Mahadevan, L. (2016). Integrative neuromechanics of crawling in D. Melanogaster Larvae. *ELife*, *5*(2016JULY), 1–23. https://doi.org/10.7554/eLife.11031

Petersen, M., Tenedini, F., Hoyer, N., Kutschera, F., & Soba, P. (2018). Assaying Thermo-nociceptive Behavior in Drosophila Larvae. *Bio-Protocol*, *8*(4), 1–12. https://doi.org/10.21769/bioprotoc.2737

Risse, B., Berh, D., Otto, N., Klämbt, C., & Jiang, X. (2017). FIMTrack: An open source tracking and locomotion analysis software for small animals. *PLoS Computational Biology*, *13*(5), 1–15. https://doi.org/10.1371/journal.pcbi.1005530

Risse, B., Thomas, S., Otto, N., Löpmeier, T., Valkov, D., Jiang, X., & Klämbt, C. (2013). FIM, a Novel FTIR-Based Imaging Method for High Throughput Locomotion Analysis. *PLoS ONE*, *8*(1), 1–11. https://doi.org/10.1371/journal.pone.0053963

Rohlfs, M., & Hoffmeister, T. S. (2003). An evolutionary explanation of the aggregation model of species coexistence. *Proceedings of the Royal Society B: Biological Sciences*, *270*(SUPPL. 1), 33–35. https://doi.org/10.1098/rsbl.2003.0002

Rosenzweig, M., Brennan, K. M., Tayler, T. D., Phelps, P. O., Patapoutian, A., & Garrity, P. A. (2005a). The Drosophila ortholog of vertebrate TRPA1 regulates thermotaxis. *Genes and Development*, *19*(4), 419–424. https://doi.org/10.1101/gad.1278205

Rosenzweig, M., Brennan, K. M., Tayler, T. D., Phelps, P. O., Patapoutian, A., & Garrity, P. A. (2005b). The Drosophila ortholog of vertebrate TRPA1 regulates thermotaxis. *Genes and Development*, *19*(4), 419–424. https://doi.org/10.1101/gad.1278205

Rosenzweig, M., Kang, K. J., & Garrity, P. A. (2008). Distinct TRP channels are required for warm and cool avoidance in Drosophila melanogaster. *Proceedings of the National Academy of Sciences of the United States of America*, *105*(38), 14668–14673. https://doi.org/10.1073/pnas.0805041105

Ruiz-Dubreuil, G., Burnet, B., Connolly, K., & Furness, P. (1996). Larval foraging behaviour and competition in Drosophila melanogaster. *Heredity*, *76*(1), 55–64. https://doi.org/10.1038/hdy.1996.7

Russo, C. A. M., Takezaki, N., & Nei, M. (1995). Molecular phylogeny and divergence times of drosophilid species. *Molecular Biology and Evolution*, *12*(3), 391–404.

1044  https://doi.org/10.1093/oxfordjournals.molbev.a040214

Schmidt-Nielsen, K. (1997). *Animal Physiology: Adaptation and Environment*. Cambridge
  University Press.

Schneider-Mizell, C. M., Gerhard, S., Longair, M., Kazimiers, T., Li, F., Zwart, M. F., Champion,
  A., Midgley, F. M., Fetter, R. D., Saalfeld, S., & Cardona, A. (2016). Quantitative
  neuroanatomy for connectomics in Drosophila. *ELife*, *5*(MARCH2016), 1–36.
  https://doi.org/10.7554/eLife.12059

Shen, W. L., Kwon, Y., Adegbola, A. A., Luo, J., Chess, A., & Montell, C. (2011). Function of
  rhodopsin in temperature discrimination in Drosophila. *Science*, *331*(6022), 1333–1336.
  https://doi.org/10.1126/science.1198904

Simpson, J. H., & Looger, L. L. (2018). *Functional Imaging and Optogenetics in Drosophila*.
  208(April), 1291–1309.

Sims, D. W., Humphries, N. E., Hu, N., Medan, V., & Berni, J. (2019). Optimal searching
  behaviour generated intrinsically by the central pattern generator for locomotion. *ELife*, *8*,
  1–31. https://doi.org/10.7554/eLife.50316

Sokolowski, M. B. (1985). Genetics and ecology of Drosophila melanogaster larval foraging and
  pupation behaviour. *Journal of Insect Physiology*, *31*(11), 857–864.
  https://doi.org/10.1016/0022-1910(85)90103-9

Sokolowski, M. B., Kent, C., & Wong, J. (1984). Drosophila larval foraging behaviour:
  Developmental stages. *Animal Behaviour*, *32*(3), 645–651. https://doi.org/10.1016/S0003-
  3472(84)80139-6

Sokolowski, M. B., Pereira, H. S., & Hughes, K. (1997). Evolution of foraging behavior in
  drosophila by density-dependent selection. *Proceedings of the National Academy of
  Sciences of the United States of America*, *94*(14), 7373–7377.
  https://doi.org/10.1073/pnas.94.14.7373

Stoessel, A., & Fischer, M. S. (2012). Comparative intralimb coordination in avian bipedal
  locomotion. *Journal of Experimental Biology*, *215*(23), 4055–4069.
  https://doi.org/10.1242/jeb.070458

Sun, X., Liu, Y., Liu, C., Mayumi, K., Ito, K., Nose, A., & Kohsaka, H. (2020). A neuromechanical
  model and kinematic analyses for Drosophila larval crawling based on physical
  measurements. *BioRxiv*.

Tastekin, I., Riedl, J., Schilling-Kurz, V., Gomez-Marin, A., Truman, J. W., & Louis, M. (2015).
  Role of the subesophageal zone in sensorimotor control of orientation in drosophila larva.
  Current Biology, *25*(11), 1448–1460. https://doi.org/10.1016/j.cub.2015.04.016

Theunissen, L. M., Bekemeier, H. H., & Dürr, V. (2015). Comparative whole-body kinematics of
  closely related insect species with different body morphology. *Journal of Experimental
  Biology*, *218*(3), 340–352. https://doi.org/10.1242/jeb.114173

Thompson, N. E., O'Neill, M. C., Holowka, N. B., & Demes, B. (2018). Step width and frontal
  plane trunk motion in bipedal chimpanzee and human walking. *Journal of Human
  Evolution*, *125*, 27–37. https://doi.org/10.1016/j.jhevol.2018.09.006

Tracey, W. D., Tracey, W. D., Wilson, R. I., Wilson, R. I., Laurent, G., Laurent, G., Benzer, S., &
  Benzer, S. (2003). Gene Essential for Nociception. *Cell*, *113*, 261–273.

Turelli, M., Cooper, B. S., Richardson, K. M., Ginsberg, P. S., Peckenpaugh, B., Antelope, C. X.,
  Kim, K. J., May, M. R., Abrieux, A., Wilson, D. A., Bronski, M. J., Moore, B. R., Gao, J. J.,
  Eisen, M. B., Chiu, J. C., Conner, W. R., & Hoffmann, A. A. (2018). Rapid Global Spread
  of wRi-like Wolbachia across Multiple Drosophila. *Current Biology*, *28*(6), 963-971.e8.
  https://doi.org/10.1016/j.cub.2018.02.015

Venken, K. J. T., Simpson, J. H., & Bellen, H. J. (2011). Genetic manipulation of genes and
  cells in the nervous system of the fruit fly. *Neuron*, *72*(2), 202–230.
  https://doi.org/10.1016/j.neuron.2011.09.021

Watada, M., Hayashi, Y., Watanabe, K., Mizutani, S., Mure, A., Hattori, Y., & Uemura, T. (2020).
  Divergence of Drosophila species: longevity and reproduction under different nutrient
  balances . *Genes to Cells*, *June*. https://doi.org/10.1111/gtc.12798

Watanabe, K., Kanaoka, Y., Mizutani, S., Uchiyama, H., Yajima, S., Watada, M., Uemura, T., &

Hattori, Y. (2019). Interspecies Comparative Analyses Reveal Distinct Carbohydrate-
Responsive Systems among Drosophila Species. *Cell Reports*, *28*(10), 2594-2607.e7.
https://doi.org/10.1016/j.celrep.2019.08.030
Wertheim, B., Marchais, J., Vet, L. E. M., & Dicke, M. (2002). Allee effect in larval resource
exploitation in Drosophila: An interaction among density of adults, larvae, and micro-
organisms. *Ecological Entomology*, *27*(5), 608–617. https://doi.org/10.1046/j.1365-
2311.2002.00449.x
Zwart, M. F., Pulver, S. R., Truman, J. W., Fushiki, A., Fetter, R. D., Cardona, A., & Landgraf,
M. (2016). Selective Inhibition Mediates the Sequential Recruitment of Motor Pools.
Neuron, *91*(3), 615–628. https://doi.org/10.1016/j.neuron.2016.06.031

1108