Exosome-Mediated mRNA Delivery For SARS-CoV-2 Vaccination

[1]Shang Jui Tsai, [1]Chenxu Guo, [2]Alanna Sedgwick, [2]Saravana Kanagavelu, [2]Justin Nice,

[2]Sanjana Shetty, [2]Connie Landaverde, [2]Nadia A. Atai, and [1]*Stephen J. Gould

[1]Department of Biological Chemistry, The Johns Hopkins University School of Medicine,

725 North Wolfe Street, Baltimore, MD, 21205

[2]Capricor Therapeutics, Inc. 8840 Wilshire Blvd. 2nd Floor, Beverly Hills, CA 90211

Correspondence:

Name: Stephen J. Gould

Address: 725 North Wolfe Street

13   Physiology Building, Room 409

14   Baltimore, MD 21205

15   USA

16 Phone number: 443 847 9918

17 Email: sgould@jhmi.edu

19 Running title: Exosomal SARS-CoV-2 vaccine

21 Key Words: COVID19, spike, nucleocapsid, exosomes, mRNA, lipid, antibody, T-cell,

22 extracellular vesicles

Abstract

Expression-dependent, Spike-only vaccines have been developed, deployed, and shown to be effective in the fight against SARS-CoV-2. However, additional approaches to vaccine development may be needed to meet existing and future challenges posed by emerging Spike variant strains, as well as a likely need for different antigen-delivery systems that are safe and effective for regular, periodic re-administration. We report here the development of mRNA-loaded exosomes, demonstrate that they can mediate the functional expression of heterologous proteins *in vitro* and *in vivo*, and have fewer adverse effects than comparable doses of lipid nanoparticles. Furthermore, we applied this approach to the development of an exosome-based, multiplexed mRNA vaccine that drives expression of immunogenic SARS-CoV-2 Nucleocapsid and Spike proteins. This vaccine elicited long-lasting cellular and humoral responses to Nucleocapsid and to Spike, demonstrating that exosome-based mRNA formulations represent a previously unexplored platform in the fight against COVID-19 and other infectious diseases.

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is the causative agent of COVID-19 [1,2]. COVID-19 typically presents with symptoms common to many respiratory infections, including fever and cough, but in many cases progresses to more severe disease that may include acute respiratory distress, disseminated disease, and death [34-8]. SARS-CoV-2 entered the human population in late 2019 as the result of a zoonotic leap and is most closely related to coronaviruses endemic to bats (Chiroptera) [9]. SARS-CoV-2 is the third recent zoonotic betacoronavirus to enter the human population, the others being responsible for the outbreaks of severe acute respiratory syndrome (SARS-CoV) in 2002 [10] and middle east respiratory syndrome (MERS-CoV) in 2012 (Memish et al., 2013), indicative of a generally susceptibility of human populations to coronavirus zoonoses. These zoonoses are more distantly to human endemic betacoronaviruses (OC43, HKU1, etc.) that also cause respiratory infections of milder effect [11]). While SARS-CoV-2 infection is associated with lower mortality than SARS-CoV or MERS-CoV, SARS-CoV-2 initially displayed a higher rate of transmission, quickly became a major cause of morbidity and mortality worldwide (https://www.cdc.gov/coronavirus/2019-ncov/hcp/clinical-guidance-management-patients.html)(coronavirus.jhu.edu), and has continued to evolve into numerous variant strains that display even more-elevated rates of transmission and an emerging resistance to antibody-based neutralization [12-15].

SARS-CoV-2 enters host cells via a multistep pathway that begins with binding between the Spike protein on the virus surface and its cognate receptor proteins on the host cell

surface. These include angiotensin-converting enzyme II (ACE2) [1,16,17], neuropilin-1 [18,19],

and perhaps also CD147 [20]. Following virus-cell binding, host cell proteases (e.g.

TMPRSS2 [16], cathepsins [21], etc.) cleave Spike, potentiating Spike-catalyzed fusion

between the viral and cellular membranes and functional infection of the host cell. Not

surprisingly, SARS-CoV-2 receptors and proteases are expressed within the respiratory

tract, consistent with its respiratory mode of transmission [22]. However, they are also

expressed in many other cell types, allowing SARS-CoV-2 to spread within the body and

impact multiple organ systems (brain, heart, gastrointestinal tract, circulatory system,

immune system, etc. [18,19,23-26]).

Following virus-cell membrane fusion, the viral genomic RNA (gRNA) is translated to

generate two large polyproteins, open reading frame 1 (orf1a) and orf1ab, which are

processed to release 16 nonstructural proteins (nsp1-16) [27]. These early proteins prime

the host cell for virus replication and mediate the synthesis of subgenomic viral RNAs and

their unique protein products. These include a dozen or more additional proteins,

including the SARS-CoV-2 structural proteins Nucleocapsid (N), Spike (S), Membrane

(M), and Envelope (E). Spike, Membrane and Envelope are integral membrane proteins,

co-translationally translocated into the endoplasmic reticulum (ER), that subsequently

drive virion formation while also incorporating the Nucleocapsid and its bound gRNA as

well as some other ancillary proteins [28,29], with virus release via lysosomal exocytosis

(Ghosh et al., 2020) []. The released viral particles are ~100 nm diameter, display

prominent Spike protrusions from the cell surface and a lumen containing Nucleocapsid-

gRNA complexes[30]. SARS-CoV-2 biogenesis also involves extensive processing of its

Spike protein at a polybasic site, generating S1 and S2 forms of Spike, with the N-

terminal, receptor-binding S1 fragment bound non-covalently to the fusogenic,

membrane-anchored S2 fragment [16,31,32].

Vaccine design should mirror, and ideally improve on, the correlates of protective

immunity that arise from natural infections. It is now well-established that SARS-CoV-2

infection generates potent cellular and humoral immune responses to viral proteins that

in most cases reverse the course of disease, clear the viral infection, and confer

resistance to reinfection in both people and in animal models [33-3637-39]. Disease-preventing

vaccines have previously been developed for animal coronaviruses [40] and have been

successfully developed and deployed for SARS-CoV-2 [41-47], a development that is likely

to save millions of lives. However, these first-generation SARS-CoV-2 vaccines only elicit

immunity to a single viral protein, Spike, the rapid evolution of which may impair vaccine

efficacy [12-15]. Furthermore, the Spike-only vaccine approach ignores the fact that a

primary correlate of immunity in COVID-19 patients is the array of potent immune

reactions to the SARS-CoV-2 Nucleocapsid protein [48].

Here we describe an expression-dependent SARS-CoV-2 vaccine that combines

exosome-based delivery, multiplexed mRNA formulation, induction of immunity to both

Spike and Nucleocapsid, and antigen design that involves expressing Nucleocapsid in a

form designed for improved antigen presentation. Exosomes are small extracellular

vesicles (sEVs) of ~30-150 nm in diameter that are made by all cells, abundant in all

biofluids, and mediate intercellular transmission of signals and macromolecules, including

RNAs [49]. Allogenic exosome transplantations and transfusions have been practices in

one form or another for more than a century and have never been associated with any

adverse effects. Moreover, exosomes have already been shown effective for delivery of

RNA-based therapeutics [50,51]. The remainder of this report describes the production of

engineered exosome/mRNA formulations, their ability to drive protein expression in

cultured cells and animals, their improved safety relative to LNPs, and their use as a

multiplexed, exosome-based SARS-CoV-2 vaccine that elicited immunity to multiple viral

antigens, including Nucleocapsid as well as Spike.

Results

Exosomes display robust ability to deliver functional mRNAs in vitro and in vivo

Exosomes are capable of delivering functional RNAs to target cells [50,51], but so too are synthetic lipid vesicles, often referred to as lipid nanoparticles (LNPs) [52]. To better understand the dynamics of mRNA delivery by these two natural and synthetic forms of soluble vesicles, we generated matched formulations of mRNA-loaded exosomes and mRNA-loaded LNPs. Exosomes were purified from the culture of 293F cells (***Fig. 1***), LNPs [52] were obtained from a commercial provider, and equal amounts of each (by vesicle number) were loaded with a synthetic mRNA encoding the hybrid luciferase/fluorescent protein Antares2 (Antares2 is comprised of the luciferase teLuc fused to two copies of the fluorescent protein CyOFP1 (CyOFP1-teLuc-CyOFP1), emits far-red shift light via bioluminescent resonance energy transfer[53]). Equal amounts of these matched exo-mRNA and LNP-mRNA formulations were then incubated at low and high doses with human cells, followed by an overnight incubation to allow for Antares2 protein expression. The next day, the cells were incubated with diphenylterazine (DTZ), a cell-permeable substrate (luciferin) for Antares2, and assayed for DTZ-dependent, Antares2-catalyzed light emission (***Fig. 2***). At low-dose administration, Antares2 expression was 25% higher in cells treated with the exo-mRNA formulation than with the LNP-mRNA formulation (n = 6, *p* = 0.0016). The difference in Antares2 expression was even more pronounced at high-dose administration, as the exo-mRNA-treated cells expressed far more Antares2 activity than the LNP-exo-treated cells (16-fold; n = 6; *p* = 0.00035).

This large difference in particle-mediated Antares2 expression was caused by a drop in

LNP-mRNA-mediated expression, raising the possibility that LNP administration is

inhibitory at high levels of administration. This in turn raised the possibility of general

toxicity of LNP administration, which we addressed by following the short-term

consequences of exosome and LNP injections in mice. Animals were injected (i.m.) with

equal numbers of either exosomes or LNPs (50 ml of ??? particles/ml), returned to their

cages for three days, and then sacrificed and processed for organ histology by an

independent testing laboratory (**Fig. 3A**). No abnormalities were detected in control

animals (5/5) or in animals injected with exosomes (5/5). In contrast, only one of the LNP-

injected animals (1/5) displayed normal spleen histology, as 4/5 animals showed an

increase in red pulp. Adverse LNP effects may also explain the ~5% reduction in body

mass (n = 5; p = 0.05) we observed at 3 days post-injection (**Fig. 3B**).

The robust expression of exosome-delivered mRNA *in vitro* and the absence of exosome-

associated adverse effects led us to next test whether RNA-loaded exosomes might also

be able to drive Antares2 expression *in vivo*. Towards this end, we injected adult mice

(0.05 ml volume, intramuscular (i.m.) administration) with Antares2 mRNA-loaded

exosomes, returned the animals to cages to allow for Antares2 expression. 24 hours later,

the control (uninjected) and treated mice were injected (i.p.) with a solution of the

Antares2 luciferin DTZ and imaged immediately using a real-time bioluminescent imaging

(BLI) system to visualize exosome-mediated, mRNA-directed Antares2 expression.

Control animals displayed no significant light emission upon DTZ injections whereas

animals that had been injected with the mRNA-loaded exosome formulation displayed

robust light emission (**Fig. 3**). These observations demonstrate that RNA-loaded

exosomes can deliver functional mRNAs into cells in live animals in a way that leads to

mRNA translation, protein expression, and directed enzyme activity.

Design and validation of S^{W1} and LSNME mRNAs

We next tested whether exosome-mRNA formulations can be used to elicit immune

responses to mRNA-encoded antigens. Towards this end, we synthesized a pair of

mRNAs, one of which expresses the form of SARS-CoV-2 Spike (S^{W1}) encoded by the

initial viral isolate [1]. The second mRNA expresses a fusion protein (LSNME) comprised

of the SARS-CoV-2 Nucleocapsid protein, as well as fragments of the Spike, Membrane,

and Envelope proteins, all inserted in the extracellular domain of human Lamp1 (this

Lamp1-based fusion protein aims to induce anti-SARS-CoV-2 immunity by targeting viral

protein fragments to the MHC Class I and II antigen presentation pathways [54,5556]).

Transfection of these mRNAs into HEK293 cells (**Fig. 4**) resulted in expression of Spike

at the cell surface but also at internal organelles (shown elsewhere to be lysosomes [57]),

whereas expression of LSNME led to its accumulation in what appears to the

endoplasmic reticulum, the site of MHC Class I peptide loading and maturation.

The LSNME/S^{W1} vaccine induces antibody responses to N and S

A single exosome-mRNA formulation containing both the LSNME and S^{W1} mRNAs

(hereafter referred to as the LSNME/S^{W1} vaccine) was injected (i.m.) into 13 weeks-old

male C57BL/6J mice (**Fig. 5**). The vaccine was dosed at 4 ug or 0.25 ug equivalents of

each mRNA and injections were performed on day 1 (primary immunization), day 21 (1st

boost), and day 42 (2nd boost). Blood (0.1 mL) was collected on days 14, 35, 56, 70 and

84. On day 84 the animals were sacrificed to obtain tissue samples for histological

analysis and splenocytes for blood cell studies. Using ELISA kits adapted for the detection

of mouse antibodies, we observed that vaccinated animals displayed a dose-dependent

antibody response to both the SARS-CoV-2 N protein and S protein. These antibody

reactions were not particularly robust but they were long-lasting, persisting to 7 weeks

after the final boost with little evidence of decline. It should be noted that the modest

antibody production was expected in the case of the N protein, as the LSNME mRNA is

designed to stimulate cellular immune responses rather than the production of anti-N

antibodies.

198   *LSNME/S^{W1} vaccination induces cellular immune responses to N and S*

Vaccinated and control animals were also interrogated for the presence of antigen-

reactive CD4+ and CD8+ T-cells. This was carried out by collecting splenocytes at the

completion of the trial (day 84) using a CFSE proliferation assay in the presence or

absence of recombinant N and S proteins. These experiments revealed that vaccination

had induced a significant increase in the percentages of CD4$^+$ T-cells and CD8$^+$ T-cells

that proliferated in response to addition of either recombinant N protein or recombinant S

protein to the culture media (***Fig. 6A-D***). These vaccine-specific, antigen-induced

proliferative responses demonstrate that the LSNME/S^{W1} vaccine achieved its primary

goal, which was to prime the cellular arm of the immune system to generate N-reactive

CD4$^+$ and CD8$^+$ T-cells, and also S-reactive CD4$^+$ and CD8$^+$ T-cells. In additional

experiments, we stained antigen-induced T-cells cells for the expression of interferon

gamma (IFNγ) and interleukin 4 (IL4). These experiments revealed that the S-reactive

CD4$^+$ T-cell population displayed elevated expression of the Th1-associated cytokine

IFNγ, and to a lesser extent, the Th2-associated cytokine IL4 (**Fig 7**). In contrast, N-

reactive T-cells failed to display an N-induced expression of either IFNγ or IL4.

Absence of vaccine-induced adverse reactions

Control and vaccinated animals were examined regularly for overall appearance, general

behavior, and injection site inflammation (redness, swelling). No vaccine-related

differences were observed in any of these variables, and animals from all groups

displayed similar age-related increases in body mass (***supplemental figure 1***).

Vaccination also had no discernable effect on blood cell counts (***supplemental figure 2***).

Histological analyses were performed on all animals at the conclusion of the study by an

independent histology service, which reported that vaccinated animals showed no

difference in overall appearance of any of the tissues that were examined. Representative

images are presented for brain, lung, heart, liver, spleen, kidney, and side of injection

skeletal muscle in an animal from each of the trial groups (***Fig. 8***).

Discussion

Exosomes are natural products of human cells that are more 'self' than 'non-self'. Immune

systems are tolerant of the high levels of exosomes that are continuously present in all

biofluids (e.g. blood, lymph, cerebrospinal fluid, vitreous, interstitial fluids, etc.) [49,58].

Furthermore, there is no evidence of adverse effects of allogeneic exosome transfer,

whether of purified exosomes (from amniotic fluid, blood, etc.) or of inadvertent exosome

transfer during tissue transplantation, blood transfusion, plasma injection, etc. In this

context, the fact that exosomes normally participate in pathways of vesicle-mediated,

intercellular RNA traffic [59-61] indicates that exosomes may be an ideal vehicle for clinical

RNA delivery. The data presented here support this hypothesis by showing that that

exosome-mRNA formulations can support the *in vivo*, functional expression of proteins

as diverse as soluble cytoplasmic enzymes, viral structural proteins, and synthetic fusion

proteins.

Our findings are also relevant to the ongoing battle against SARS-CoV-2. Current vaccine

strategies are all centered on inducing immunity to Spike, but Spike-only vaccines are

susceptible to escape effects whenever and antigenically shifted Spike variants starts to

spread in susceptible populations. While we are developing strategies designed to

address this challenge by improved design of expression-dependent Spike vaccines, we

are also working to address it by generating a multiplexed mRNA vaccine that delivers

two or more mRNAs, one encoding Spike and the others encoding Nucleocapsid and

perhaps fragments of other proteins as well. One limitation of this approach is that

Nucleocapsid is a cytoplasmic protein rather than a surface antigen, a topology that limits

its efficacy in vaccination studies. However, this limitation can be overcome by expressing

Nucleocapsid as part of a fusion with the lysosomal resident protein Lamp1, which places

Nucleocapsid protein in the correct compartments for Class I and Class II antigen

presentation (ER and lysosome/MHC Class II compartment, respectively). This approach

was realized in our LSNME/S^{W1} vaccine, which elicited strong cellular immune responses

to Nucleocapsid as well as to Spike. Vaccinated animals displayed antigen-induced CD4$^+$

and CD8$^+$ T-cell responses to both Nucleocapsid and to Spike that persisted for nearly

two months after immunization. Furthermore, when these cell populations were

interrogated for antigen-induced expression of the cytokines IFNγ and IL4, we detected

elevated expression of IFNγ in CD4+ T-cells exposed to exogenous Spike protein, as well

as a more modest Spike-induced expression of IL4. These results raise the possibility

that the exosome-based LSNME/S^{W1} vaccine induces the kind of Th1-skewed cellular

immune response desired for an anti-viral vaccine. Vaccinated animals also developed

durable antibody responses to the Nucleocapsid and the Spike proteins that were

sustained at relatively constant levels over the 7 weeks following immunization. This

multi-antigen immune response bodes well for this approach in the next generation of

SARS-CoV-2 vaccines that will be needed to protect against the emerging array of

antigenically distinct SARS-CoV-2 viral strains and their ever-increasing spectrum of

Spike protein mutations.

In conclusion, the results presented in this study validate the use of multiplexed exosome-

mRNA formulations for functional delivery of mRNAs both in cultured cells and in live

animals. The successful use of exosomes to deliver Antares2 mRNA opens the door to

follow-on studies aimed at optimizing exosome-RNA formulation conditions, as well as

for characterizing the time-dependence of Antares2 expression, biodistribution of

exosome-mediated RNA expression, injection site effects, and exosome-mediated tissue

tropism. As for the future development of exosome-based SARS-CoV-2 mRNA vaccines,

we anticipate that follow-on studies will demonstrate multiple advantages of exosome-

based delivery, improved antigen designed, and most importantly, improved protective

effects that arise from immunization with multiple viral antigens, and particularly

Nucleocapsid, which is a main target of anti-SARS-CoV-2 immunity in COVID-19 patients

[48] and has proven effective in vaccine studies of other coronaviruses [54]. Furthermore, the

fact that exosomes can be deployed at high concentrations without adverse effects on

cells or animals bodes well for their future use in dosing regimens that require higher-

level or ongoing repeated injections.

Methods

Cell culture

293F cells (Gibco, Cat.# 51-0029) were tested for pathogens and found to be free of viral (cytomegalovirus, human immunodeficiency virus I and II, Epstein Barr virus, hepatitis B virus, and parvovirus B19), and bacterial (*Mycoplasma*) contaminants. Cells were maintained in FreeStyle 293 Expression Medium (Gibco, #12338-018) and incubated at 37°C in 8% CO_2. For exosome production, 293F cells were seeded at a density of 1.5 x 10^6 cells/ml in shaker flasks in a volume of ~1/4 the flask volume and grown at a shaking speed of 110 rpm. HEK293 cells were grown in Dulbecco's modified Eagle's medium supplemented with 10% fetal calf serum.

Exosome purification

293F cells were grown in shaker cultures for a period of three days. Cells and large cell debris were removed by centrifugation at 300 x *g* for 5 minutes followed by 3000 x *g* for 15 minutes. The resulting supernatant was passed through a 0.22 µm sterile filtration filter unit (Thermo Fisher, #566-0020) to generate a clarified tissue culture supernatant (CTCS). The CTCS was concentrated by centrifugal filtration (Centricon Plus-70, Ultracel-PL Membrane, 100 kDa size exclusion, Millipore Sigma # UFC710008), with ~120 mLs CTCS concentrated to ~0.5 mLs. Concentrated CTCS was then purified by size exclusion chromatography (SEC) in 1x PBS (qEV original columns/35 nm: Izon Science, #SP5), with the exosomes present in each 0.5 mL starting sample eluting in three 0.5 mL fractions. Purified exosomes were reconcentrated using Amicon® Ultra-4 100 kDa cutoff

spin columns (#UFC810024). This process yielded a population of exosomes/small EVs

that have the expected ultrastructure and size distribution profile of human exosomes and

contain the exosomal marker proteins CD9 and CD63 (*Fig. 8*), at a concentrating effect

of ~500-fold, to a final concentration of ~2 x 10^{12} exosomes/ml, representing an average

recovery of 35%.

Nanoparticle Tracking Analysis (NTA)

Vesicle concentrations and size distribution profiles of exosome preparations were

measured by nanoparticle tracking analysis (NTA) using a NanoSight NS300 (Malvern

Panalytical, United Kingdom) in 1x PBS clarified by filtration through a 0.22 µm sterile

filtration unit. Measurements were carried out in triplicates at ambient temperature with

fixed camera settings (level of 14, screen gain of 10, detection threshold 3, and

temperature of 21.7-22.2 °C). Immunostaining nanoparticle tracking analysis (NTA) was

performed using fluorescently labeled antibody conjugate directed against human CD63

(AlexaFluor488-conjugated clone 460305; R&D Systems (Minneapolis, USA)). The

fluorescently labeled anti-CD63-antibody (1 µl) was incubated with exosomes (9 µl) for 2

hours at room temperature in the dark, then diluted by addition of 1 ml of sterile-filtered

PBS (Thermo Fisher, USA) and examined for exosome abundance, size, and CD63

immunoreactivity using a Particle Metrix ZetaView® TWIN device. Samples were

visualized in scatter mode using the 488 nm laser and standard instrument settings

(sensitivity: 80, shutter: 100, min. brightness: 30; min. area: 10; max. area: 1000) in

fluorescence mode with standard fluorescence settings (sensitivity: 88, shutter: 100, min.

brightness: 25; min. area: 10; max. area: 1000). The resulting videos were analysed with

the ZetaView® software 8.05.10 (Particle Metrix, Germany).

Immunoblots

Exosome and cell lysates were separated by SDS-PAGE using pre-cast, 4-15% gradient

gels (Bio-Rad 4561086) and transferred to PVDF membranes (ThermoFisher, #88518).

Membranes were blocked, probed with antibodies directed against CD9 (clone HI9a;

BioLegend), CD63 (MX-49.129.5), CD81 (555675; BD Pharmingen), or HSP90 (sc-

13119; Santa Cruz Biotechnology), then washed, exposed to HRP-conjugates of goat

secondary antibodies (Jackson Immunoresearch), washed, and processed for

chemiluminescent imaging using HRP-activated chemiluminescence detection solution

(Amersham ECL Western Blotting Detection Reagents; cat# RPN2106), and imaged

using a GE Amersham Imager 600. Images were exported as JPEG files, analyzed using

ImageJ software, and processed using Photoshop (Adobe).

Electron Microscopy and light microscopy

Exosomes were fixed by addition of formaldehyde to a final concentration of 4%. Carbon-

coated grids were placed on top of a drop of the exosome suspension. Next, grids were

placed directly on top of a drop of 2% uranyl acetate. The resulting samples were

examined with a Tecnai-12 G2 Spirit Biotwin transmission electron microscope (John

Hopkins University, USA). Fluorescence micrographs of Antares2 expression in

transfected HEK293 cells were captured as PNG files using an EVOS M7000 microscope

equipped with an Olympus UPlanSAPo 40x/0.95 objective.

Production of mRNA-loaded exosomes and LNPs

mRNAs were obtained from a commercial provider (Trilink). mRNAs were purified using RNeasy columns (Qiagen) and resuspended in DNase-free, RNase-free water using nuclease-free tips and tubes. Purified mRNAs were pre-incubated with a coating of polycationic lipids and then mixed with equal amounts of either purified exosomes or LNPs (DOTAP/DOPE, #F50102, FormuMAx Scientific Inc) at 4°C for 10 minutes. Formulations were either used immediately or frozen at -80°C and thawed rapidly prior to use.

Luciferase measurements and bioluminescent imaging

HEK293 cells were incubated with exosome-mRNA formulations overnight under standard culture conditions. Antares2 luciferase activity was measured by Live cell bioluminescence was collected after incubating with substrate diphenylterazine (MCE, HY-111382) at final concentration of 50 µM for 3 minutes. Readings were collected using a SpectraMax i3x (Molecular Devices). For in vivo studies, thirteen months-old, female Balb/c mice (Jackson Laboratory) housed under pathogen-free conditions at the Cedars-Sinai Medical Center animal facility were used to study the expression of Exosome-Anteres2 mRNA expression 24 hours after injection. Intramuscular injections were at a volume of 50 µls per mouse containing 5 ug mRNA. After 24 hours the animals were imaged using an IVIS Spectrum imager (PerkinElmer, Waltham, MA) (All animal experimentation was performed following institutional guidelines for animal care and were approved by the Cedars-Sinai Medical Center IACUC (#8602).

Animal experimentation

All animal experimentation was performed following institutional guidelines for animal care and were approved by the Cedars-Sinai Medical Center IACUC (#8602). All injections were at a volume of 50 μls. Experiments involved injection of exosomes, LNPS, and Antares2 mRNA-loaded exosomes were performed with BALB/c mice (Jackson Laboratory). Immunization with mRNA-loaded exosomes were performed on thirteen weeks-old, male C57BL/6J mice (Jackson Laboratory) housed under pathogen-free conditions at the Cedars-Sinai Medical Center animal facility. Blood (~0.1 mL) was collected periodically from the orbital vein. At day 84, mice were deeply anesthetized using isoflurane, euthanized by cervical dislocation, and processed using standard surgical procedures to obtain spleen, lung, brain, heart, liver, kidney, muscle, and other tissues. Spleens were processed for splenocyte analysis, and all tissues were processed for histological analysis by fixation in 10% neutral buffered formalin. Histological analysis was performed by the service arm of the HIC/Comparative Pathology Program of the University of Washington.

ELISA for SARS-CoV-2 antigen-specific antibody responses

Mouse IgG antibody production against SARS-CoV-2 antigens was measured by enzyme-linked immunosorbent assays (ELISA). For antigens S1 (RBD) and N, pre-coated ELISA plates from RayBiotech were utilized (IEQ-CoV S RBD-IgG; IEQ-CoVN-IgG), and the experiments were performed according to the manufacturer's instructions, with modification. Briefly, mouse plasmas at dilutions of 1:50 were added to antigen pre-

coated wells in duplicates and incubated at room temperature (RT) for 2 hours on a

shaker (200 rpm). The plates were washed 4 times with wash buffer followed by blocking

for 2 hours at RT with 1% BSA in PBS. Mouse antibodies bound to the antigens coated

on the ELISA plates were detected using HRP-conjugated goat anti-mouse secondary

antibodies (Jackson Immuno Research Inc.) Plates were washed 4 times with washing

buffer, and developed using TMB substrate (RayBiotech). Microplate Reader was used

to measure the absorbance at 650 nm (SpectraMaxID3, Molecular Devices, with SoftMax

Pro7 software).

Single cell splenocyte preparation

After terminal blood collection, mice were euthanized, and part of fresh spleens were

harvested. Single cell splenocyte preparation was obtained by machinal passage through

a 40 μm nylon cell strainer (BD Falcon, #352340). Erythrocytes were depleted using

Ammonium-Chloride-Potassium (ACK) lysis buffer (Gibco, #A10492-01), and

splenocytes were washed using R10 media by centrifuging at 300x *g* for 5 minutes at RT.

R10 media (RPMI 1640 media (ATCC, Cat#302001) supplemented with 10% fetal bovine

serum (FBS) (Atlas, #E01C17A1), 50 μM 2-mercaptoethanol (Gibco, #21985-023),

penicillin/streptomycin (VWR life sciences, #K952), and 10 mM HEPES (Gibco, #15630-

080)) was used for all analyses of blood cells. The cells were resuspended in fresh media

and counted in hemocytometer counting chamber to be used in subsequent experiments.

Spleen lymphocyte population characterization

Splenocytes (2×10^5 cells/mouse) were resuspended in 100 µL of 10% FBS in 1x PBS

and incubated with fluorochrome-conjugated antibodies for surface staining of CD3

(Invitrogen, #17-0032-82) CD4 (Biolegend, #100433), CD8 (Biolegend, #100708), B220

(BD, #552771) CD11c (Invitrogen, #17-0114-81), F4/80 (Invitrogen, #MF48004) Ly6G

(Invitrogen, #11-9668-80) and Ly6C (BD, #560592)) for 30 minutes at 4 °C in the dark.

Following incubation, samples were washed twice with 200 µLs 10% FBS in 1x PBS and

centrifuged at 300 x g for 5 minutes at RT to remove unbound antibodies. Next the cells

were fixed with 100 µLs ICS fixation buffer (Invitrogen, #00-8222-49). Samples were

analyzed on a FACS Canto II (BD Biosciences) with 2,000 – 10,000 recorded

lymphocytes . The data analysis was performed using FlowJo 10 software (FlowJo, LLC)

and presented as a percentage change in the immune cell population compared to the

vehicle-treated group.

SARS-CoV-2 antigen-specific T cell proliferation assay using CFSE

Splenocytes were resuspended at 10^6 cells/mL in 10% FBS in 1xPBS and stained with

carboxyfluorescein succinimidyl ester (CFSE) (Invitrogen, #C34554) by rapidly mixing

equal volume of cell suspension with 10 µM CFSE in 10% FBS in 1x PBS for 5 minutes

at 37°C. The labeled cells were washed three times with R10 complete medium. The cells

were incubated for 96 hours in the presence of 10 µg/mL SARS-CoV-2 antigens N or S1

(Acro Biosystems, #NUN-C5227; SIN-C52H4) or medium alone as negative control. After

96 hours, cells were washed with 200 µLs 10% FBS in 1xPBS and centrifuged at 300 x g

for 5 minutes at RT. Cells were then stained with anti-CD3-APC (Invitrogen, #17-0032-

82), anti-CD4-PerCP-Cy5.5 (Biolegend, #100433), and anti-CD8-PE antibodies

(Biolegend, #MCD0801) for 30 minutes at 4°C. The stained cells were washed twice with

200 µLs 1x PBS and analyzed on a FACS Canto II (BD Biosciences). For analysis,

lymphocytes were first gated for CD3+ T-cells, then for CD4+/CD8− or CD8+/CD4−

populations. The data analysis was performed using FlowJo 10 software (FlowJo LLC).

Intracellular staining for cytokines

2.0×10^5 splenocytes/mouse were incubated for 72 hours in the presence of 10 µg/mL

SARs-CoV2 antigens N or S1 (Acro Biosystems) or R10 medium alone (negative control).

After 72 hours, the cells were washed with fresh R10 medium and incubated with phorbol

myristate acetate (PMA) at concentration of 50 ng/mL (Sigma, #P1585), ionomycin at

concentration of 350 ng/mL (Invitrogen, #124222), and GogiPlug at concentration of 0.8

µL/mL (Invitrogen, #51-2301KZ) for 4 hours to amplify cytokine expression in T cells. The

cells were then washed with 10% FBS in 1x PBS and stained with anti-CD3-APC, anti-

CD4-PerCP-Cy5.5, and anti-CD8-PE antibodies (Added above) for 30 minutes at 4°C in

dark. The cells were washed twice with 1xPBS followed by permeabilization step using

ready-to-use buffer (Invitrogen #00-8333-56). Next the cells were fixed with ICS fixation

bufferAdded above for 10 minutes at RT in dark and stained intracellular for IFN-γ

(eBioscience, #11-7311-82), IL-10 (eBioscience, #11-7101-82), IL-4 (Invitrogen, #12-

7041-41) and Foxp3 (Invitrogen, #12-5773-80) overnight at 4°C in permeabilization buffer.

The stained cells were analyzed on a BD FACS Canto II with 5,000 – 10,000 recorded

lymphocytes. The data analysis was performed using FlowJo 10 software.

Statistical Analysis

Statistical analysis was performed using GraphPad Prism 8 software for Windows/Mac

(GraphPad Software, La Jolla California USA) or Excel. Results are reported as mean ±

standard deviation or mean ± standard error, and the differences were analyzed using

Student's t-test or one-way analysis of variance.

Sources of Support: This study was conducted with support from Capricor and from

Johns Hopkins University.

Disclosures: S.J.G is a paid consultant for Capricor, holds equity in Capricor, and is co-

inventor of intellectual property licensed by Capricor. S.J.T. is co-inventor of intellectual

property licensed by Capricor. C.G. is co-inventor of intellectual property licensed by

Capricor. A.S., S.K., J.N., S.S., C.L., and N.A. are employees of Capricor.

Acknowledgments: We thank Omid Sheikh for outstanding technical assistance during

the course of these studies.

Figure Legends

Figure 1. Exosome purification and characterization. (A) Schematic of exosome

purification from cultures of 293F cells grown in chemically defined media. (B) NTA

analysis of purified exosomes showed a mean exosome diameter of ~115 nm. (C)

Negative stain electron micrograph of purified exosomes. Bar, 100 nm. (D)

Immunofluorescent NTA analysis of 293F-derived exosomes that had been labeled

previously using fluorescently labeled anti-CD63 antibody. (E) Immunoblot analysis of

equal proportions of 293F cell and exosome lysates using antibodies specific for the

exosomal markers CD81, CD9, & CD63, as well as the control cytoplasmic protein Hsp90.

Figure 2. Exosomes display superior mRNA delivery characteristics. Relative luciferase

activities (average +/- standard error of the mean) of cells treated with low or high

concentrations of mRNA-loaded exosomes or mRNA-loaded LNPs.

Figure 3. Effect of exosome and LNP injections on organ histology and body mass. (A)

H&E staining of tissue sections from BALB/c mice that had been injected three days

earlier with 50 ml of PBS, exosomes (10^{12}/ml), or LNPs (10^{12}/ml). (B) Body mass

measurements prior to and at 3 days after injection. All animals were subjected to analysis

by an independent pathology service, which noted spleen abnormalities in 4/5 LNP-

treated animals but no abnormalities in control or exosome-treated animals.

Figure 4. Real-time imaging of exosome-mediated, functional mRNA delivery. Combined

bioluminescent and light images of control mice and treated mice immediately following

i.p. administration of DTZ. Treated mice had been injected with Antares2 mRNA-loaded

exosomes 24 hours prior to imaging.   Radiance is in photons/second/area

(cm^2)/steradian.

Figure 5. Expression of S^{W1} and LSNME following mRNA transfection. (A, B)

Fluorescence micrographs of HEK293 cells stained with DAPI and a plasma from a

COVID-19 patient. (C-F) Fluorescence micrographs of HEK293 cells stained with DAPI

and plasmas from a COVID-19 patient following their transfection with the (C, D) S^{W1}-

encoding mRNA and (E, F) the LSNME-encoding mRNA. Bar, 50 µm.

Figure 6. LSNME/S^{W1} vaccination induces antibody responses to SARS-CoV-2 N and S

protein. (A) Schematic of immunization and blood/tissue collection timeline. (B) Anti-N

ELISA results of diluted plasma from (grey bars and black circles) individual six control

mice, (orange bars and black squares) six mice immunized with 0.25 µg equivalents of

each mRNA, and (rust bars and black triangles) six mice immunized with 4 µg equivalents

of each mRNA. (C) Anti-S1 ELISA results of diluted plasma from (grey bars and black

circles) individual six control mice, (orange bars and black squares) six mice immunized

with 0.25 µg equivalents of each mRNA, and (rust bars and black triangles) six mice

immunized with 4 µg equivalents of each mRNA. Height of bars represents the mean,

error bars represent +/- one standard error of the mean, and the statistical significance of

differences between different groups is reflected in Student's t-test values of * for <0.05,

** for <0.005, and *** for <0.0005.

Figure 7. LSNME/S^{W1} vaccination induces CD4^+ and CD8^+ T-cell responses. CFSE-

labeled splenocytes were interrogated by flow cytometry following incubation in the

absence or presence of (A, B) purified, recombinant N protein or (C, D) purified,

recombinant S protein, and for antibodies specific for CD4 and CD8. Differences in

proliferation of CD4^+ cells and CD8^+ cells were plotted for (grey bars and black circles)

individual six control mice, (orange bars and black squares) six mice immunized with 0.25

μg equivalents of each mRNA, and (rust bars and black triangles) six mice immunized

with 4 μg equivalents of each mRNA. Height of bars represents the mean, error bars

represent +/- one standard error of the mean, and the statistical significance of differences

between different groups is reflected in Student's t-test values of * for <0.05 and ** for

<0.005.

Figure 8. LSNME/S^{W1} vaccination leads to S-induced expression of IFNγ and IL4 by CD4^+

T-cells. Splenocytes were interrogated by flow cytometry following incubation in the

absence or presence of (A, B) purified, recombinant N protein or (C, D) purified,

recombinant S protein, and labeling with antibodies specific for CD4 or CD8, and for IFNγ

or IL4. Differences in labeling for IFNγ or IL4 in CD4^+ CD8^+ cell populations were plotted

for (grey bars and black circles) individual six control mice, (orange bars and black

squares) six mice immunized with 0.25 μg equivalents of each mRNA, and (rust bars and

black triangles) six mice immunized with 4 μg equivalents of each mRNA. Height of bars

represents the mean, error bars represent +/- one standard error of the mean, and the

statistical significance of differences between different groups is reflected in Student's t-

test values of * for <0.05.

Figure 9. Absence of tissue pathology upon LSNME/S^{W1} vaccination. Representative

micrographs from histological analysis (hematoxylin and eosin stain) of lung, brain, heart,

liver, kidney, spleen, and muscle (side of injection) of animals from (upper row) control

mice, (middle row) mice immunized with the lower dose of the LSNME/S^{W1} vaccine, and

(lower row) mice immunized with the higher dose of the LSNME/S^{W1} vaccine.

Supplemental Figure 1. Equivalent growth of vaccinated and control animals. Body mass

of all mice was measured over the course of the study and plotted as average +/- the

standard error of the mean, relative to the body mass at the initiation of the trial, with

groups reported as (grey lines and circles) control mice, (orange lines and squares) lower

dose-treated mice, and (rust lines and triangles) higher dose-treated mice.

Supplemental Figure 2. Vaccination does not induce changes in the proportional

representation of key blood cell populations. Splenocytes were interrogated by flow

cytometry using antibodies specific for (A) B220, (B) Ly6C, (C) CD11c, and (D) CD3.

CD3$^+$ cells were further differentiated by staining for (E) CD4 and (F) CD8. No statistically

significant differences were detected in these subpopulations of white blood cells.

References

Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270-273, doi:10.1038/s41586-020-2012-7 (2020).

Coronaviridae Study Group of the International Committee on Taxonomy of, V. The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nat Microbiol* **5**, 536-544, doi:10.1038/s41564-020-0695-z (2020).

Gandhi, R. T., Lynch, J. B. & Del Rio, C. Mild or Moderate Covid-19. *N Engl J Med* **383**, 1757-1766, doi:10.1056/NEJMcp2009249 (2020).

Wang, D. *et al.* Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA* **323**, 1061-1069, doi:10.1001/jama.2020.1585 (2020).

Zhou, F. *et al.* Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* **395**, 1054-1062, doi:10.1016/S0140-6736(20)30566-3 (2020).

Force, A. D. T. *et al.* Acute respiratory distress syndrome: the Berlin Definition. *JAMA* **307**, 2526-2533, doi:10.1001/jama.2012.5669 (2012).

Richardson, S. *et al.* Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA* **323**, 2052-2059, doi:10.1001/jama.2020.6775 (2020).

Guo, T. *et al.* Cardiovascular Implications of Fatal Outcomes of Patients With Coronavirus Disease 2019 (COVID-19). *JAMA Cardiol* **5**, 811-818, doi:10.1001/jamacardio.2020.1017 (2020).

Boni, M. F. *et al.* Evolutionary origins of the SARS-CoV-2 sarbecovirus lineage responsible for the COVID-19 pandemic. *Nat Microbiol* **5**, 1408-1417, doi:10.1038/s41564-020-0771-4 (2020).

Graham, R. L. & Baric, R. S. Recombination, reservoirs, and the modular spike: mechanisms of coronavirus cross-species transmission. *J Virol* **84**, 3134-3146, doi:10.1128/JVI.01394-09 (2010).

Corman, V. M., Muth, D., Niemeyer, D. & Drosten, C. Hosts and Sources of Endemic Human Coronaviruses. *Adv Virus Res* **100**, 163-188, doi:10.1016/bs.aivir.2018.01.001 (2018).

Korber, B. *et al.* Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the COVID-19 Virus. *Cell* **182**, 812-827 e819, doi:10.1016/j.cell.2020.06.043 (2020).

Alam, I. *et al.* CovMT: an interactive SARS-CoV-2 mutation tracker, with a focus on critical variants. *Lancet Infect Dis*, doi:10.1016/S1473-3099(21)00078-5 (2021).

Ho, D. *et al.* Increased Resistance of SARS-CoV-2 Variants B.1.351 and B.1.1.7 to Antibody Neutralization. *Res Sq*, doi:10.21203/rs.3.rs-155394/v1 (2021).

15   Diamond, M. *et al.* SARS-CoV-2 variants show resistance to neutralization by many
monoclonal and serum-derived polyclonal antibodies. *Res Sq*, doi:10.21203/rs.3.rs-
228079/v1 (2021).
16   Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked
by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280 e278,
doi:10.1016/j.cell.2020.02.052 (2020).
17   Matheson, N. J. & Lehner, P. J. How does SARS-CoV-2 cause COVID-19? *Science* **369**, 510-
511, doi:10.1126/science.abc6156 (2020).
18   Daly, J. L. *et al.* Neuropilin-1 is a host factor for SARS-CoV-2 infection. *Science*,
doi:10.1126/science.abd3072 (2020).
19   Cantuti-Castelvetri, L. *et al.* Neuropilin-1 facilitates SARS-CoV-2 cell entry and infectivity.
Science, doi:10.1126/science.abd2985 (2020).
20   Wang, K. *et al.* CD147-spike protein is a novel route for SARS-CoV-2 infection to host cells.
Signal Transduct Target Ther **5**, 283, doi:10.1038/s41392-020-00426-x (2020).
21   Wei, J. *et al.* Genome-wide CRISPR Screens Reveal Host Factors Critical for SARS-CoV-2
Infection. *Cell* **184**, 76-91 e13, doi:10.1016/j.cell.2020.10.028 (2021).
22   Mason, R. J. Thoughts on the alveolar phase of COVID-19. *Am J Physiol Lung Cell Mol*
Physiol **319**, L115-L120, doi:10.1152/ajplung.00126.2020 (2020).
23   Nicin, L. *et al.* Cell type-specific expression of the putative SARS-CoV-2 receptor ACE2 in
human hearts. *Eur Heart J* **41**, 1804-1806, doi:10.1093/eurheartj/ehaa311 (2020).
24   Li, M. Y., Li, L., Zhang, Y. & Wang, X. S. Expression of the SARS-CoV-2 cell receptor gene
ACE2 in a wide variety of human tissues. *Infect Dis Poverty* **9**, 45, doi:10.1186/s40249-
020-00662-x (2020).
25   Ziegler, C. G. K. *et al.* SARS-CoV-2 Receptor ACE2 Is an Interferon-Stimulated Gene in
Human Airway Epithelial Cells and Is Detected in Specific Cell Subsets across Tissues. *Cell*
181, 1016-1035 e1019, doi:10.1016/j.cell.2020.04.035 (2020).
26   Singh, M., Bansal, V. & Feschotte, C. A Single-Cell RNA Expression Map of Human
Coronavirus Entry Factors. *Cell Rep* **32**, 108175, doi:10.1016/j.celrep.2020.108175 (2020).
27   V'Kovski, P., Kratzel, A., Steiner, S., Stalder, H. & Thiel, V. Coronavirus biology and
replication: implications for SARS-CoV-2. *Nat Rev Microbiol*, doi:10.1038/s41579-020-
00468-6 (2020).
28   Ujike, M. & Taguchi, F. Incorporation of spike and membrane glycoproteins into
coronavirus virions. *Viruses* **7**, 1700-1725, doi:10.3390/v7041700 (2015).
29   Ruch, T. R. & Machamer, C. E. The coronavirus E protein: assembly and beyond. *Viruses*
4, 363-382, doi:10.3390/v4030363 (2012).
30   Yao, H. *et al.* Molecular Architecture of the SARS-CoV-2 Virus. *Cell* **183**, 730-738 e713,
doi:10.1016/j.cell.2020.09.018 (2020).
31   Hoffmann, M., Kleine-Weber, H. & Pohlmann, S. A Multibasic Cleavage Site in the Spike
Protein of SARS-CoV-2 Is Essential for Infection of Human Lung Cells. *Mol Cell* **78**, 779-784
e775, doi:10.1016/j.molcel.2020.04.022 (2020).
32   Walls, A. C. *et al.* Structure, Function, and Antigenicity of the SARS-CoV-2 Spike
Glycoprotein. *Cell* **181**, 281-292 e286, doi:10.1016/j.cell.2020.02.058 (2020).
33   Deng, W. *et al.* Primary exposure to SARS-CoV-2 protects against reinfection in rhesus
macaques. *Science* **369**, 818-823, doi:10.1126/science.abc5343 (2020).

34   Chandrashekar, A. *et al.* SARS-CoV-2 infection protects against rechallenge in rhesus
macaques. *Science* **369**, 812-817, doi:10.1126/science.abc4776 (2020).

35   Shan, C. *et al.* Infection with novel coronavirus (SARS-CoV-2) causes pneumonia in Rhesus
macaques. *Cell Res* **30**, 670-677, doi:10.1038/s41422-020-0364-z (2020).

36   Bosco-Lauth, A. M. *et al.* Experimental infection of domestic dogs and cats with SARS-CoV-
2: Pathogenesis, transmission, and response to reexposure in cats. *Proc Natl Acad Sci U S*
A **117**, 26382-26388, doi:10.1073/pnas.2013102117 (2020).

37   St John, A. L. & Rathore, A. P. S. Early Insights into Immune Responses during COVID-19. *J*
Immunol **205**, 555-564, doi:10.4049/jimmunol.2000526 (2020).

38   Poland, G. A., Ovsyannikova, I. G. & Kennedy, R. B. SARS-CoV-2 immunity: review and
applications to phase 3 vaccine candidates. *Lancet*, doi:10.1016/S0140-6736(20)32137-1
(2020).

39   Zost, S. J. *et al.* Potently neutralizing and protective human antibodies against SARS-CoV-
2. *Nature* **584**, 443-449, doi:10.1038/s41586-020-2548-6 (2020).

40   Tizard, I. R. Vaccination against coronaviruses in domestic animals. *Vaccine* **38**, 5123-
5130, doi:10.1016/j.vaccine.2020.06.026 (2020).

41   Walsh, E. E. *et al.* Safety and Immunogenicity of Two RNA-Based Covid-19 Vaccine
Candidates. *N Engl J Med*, doi:10.1056/NEJMoa2027906 (2020).

42   Keech, C. *et al.* Phase 1-2 Trial of a SARS-CoV-2 Recombinant Spike Protein Nanoparticle
Vaccine. *N Engl J Med*, doi:10.1056/NEJMoa2026920 (2020).

43   Jackson, L. A. *et al.* An mRNA Vaccine against SARS-CoV-2 - Preliminary Report. *N Engl J*
Med, doi:10.1056/NEJMoa2022483 (2020).

44   Hassan, A. O. *et al.* A Single-Dose Intranasal ChAd Vaccine Protects Upper and Lower
Respiratory Tracts against SARS-CoV-2. *Cell* **183**, 169-184 e113,
doi:10.1016/j.cell.2020.08.026 (2020).

45   Zhu, F. C. *et al.* Immunogenicity and safety of a recombinant adenovirus type-5-vectored
COVID-19 vaccine in healthy adults aged 18 years or older: a randomised, double-blind,
placebo-controlled, phase 2 trial. *Lancet* **396**, 479-488, doi:10.1016/S0140-
6736(20)31605-6 (2020).

46   Bos, R. *et al.* Ad26 vector-based COVID-19 vaccine encoding a prefusion-stabilized SARS-
CoV-2 Spike immunogen induces potent humoral and cellular immune responses. *NPJ*
Vaccines **5**, 91, doi:10.1038/s41541-020-00243-x (2020).

47   Graham, S. P. *et al.* Evaluation of the immunogenicity of prime-boost vaccination with the
replication-deficient viral vectored COVID-19 vaccine candidate ChAdOx1 nCoV-19. *NPJ*
Vaccines **5**, 69, doi:10.1038/s41541-020-00221-3 (2020).

48   Burbelo, P. D. *et al.* Sensitivity in Detection of Antibodies to Nucleocapsid and Spike
Proteins of Severe Acute Respiratory Syndrome Coronavirus 2 in Patients With
Coronavirus Disease 2019. *J Infect Dis* **222**, 206-213, doi:10.1093/infdis/jiaa273 (2020).

49   Pegtel, D. M. & Gould, S. J. Exosomes. *Annu Rev Biochem* **88**, 487-514,
doi:10.1146/annurev-biochem-013118-111902 (2019).

50   Kamerkar, S. *et al.* Exosomes facilitate therapeutic targeting of oncogenic KRAS in
pancreatic cancer. *Nature* **546**, 498-503, doi:10.1038/nature22341 (2017).

51   Li, L. *et al.* Extracellular vesicles carry microRNA-195 to intrahepatic cholangiocarcinoma
and improve survival in a rat model. *Hepatology* **65**, 501-514, doi:10.1002/hep.28735
(2017).
52   Hassett, K. J. *et al.* Optimization of Lipid Nanoparticles for Intramuscular Administration
of mRNA Vaccines. *Mol Ther Nucleic Acids* **15**, 1-11, doi:10.1016/j.omtn.2019.01.013
(2019).
53   Yeh, H. W. *et al.* Red-shifted luciferase-luciferin pairs for enhanced bioluminescence
imaging. *Nat Methods* **14**, 971-974, doi:10.1038/nmeth.4400 (2017).
54   Gupta, V. *et al.* SARS coronavirus nucleocapsid immunodominant T-cell epitope cluster is
common to both exogenous recombinant and endogenous DNA-encoded immunogens.
Virology **347**, 127-139, doi:10.1016/j.virol.2005.11.042 (2006).
55   Vyas, J. M., Van der Veen, A. G. & Ploegh, H. L. The known unknowns of antigen processing
and presentation. *Nat Rev Immunol* **8**, 607-618, doi:10.1038/nri2368 (2008).
56   Imai, J., Otani, M. & Sakai, T. Distinct Subcellular Compartments of Dendritic Cells Used
for Cross-Presentation. *Int J Mol Sci* **20**, doi:10.3390/ijms20225606 (2019).
57   Guo, C. *et al.* The D614G Mutation Enhances the Lysosomal Trafficking of SARS-CoV-2
Spike. *bioRxiv*, doi:10.1101/2020.12.08.417022 (2020).
58   Yu, W. *et al.* Exosome-based Liquid Biopsies in Cancer: Opportunities and Challenges. *Ann*
Oncol, doi:10.1016/j.annonc.2021.01.074 (2021).
59   Skog, J. *et al.* Glioblastoma microvesicles transport RNA and proteins that promote
tumour growth and provide diagnostic biomarkers. *Nat Cell Biol* **10**, 1470-1476,
doi:10.1038/ncb1800 (2008).
60   Ratajczak, J. *et al.* Embryonic stem cell-derived microvesicles reprogram hematopoietic
progenitors: evidence for horizontal transfer of mRNA and protein delivery. *Leukemia* **20**,
847-856, doi:10.1038/sj.leu.2404132 (2006).
61   O'Brien, K., Breyne, K., Ughetto, S., Laurent, L. C. & Breakefield, X. O. RNA delivery by
extracellular vesicles in mammalian cells and its applications. *Nat Rev Mol Cell Biol* **21**,
585-606, doi:10.1038/s41580-020-0251-y (2020).
740

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A N-induced cytokine expression

B S1-induced cytokine expression

Figure 8

Figure 9

Supplemental Figure 1

Supplemental Figure 2