Reciprocal social ties drive stability within a social network

Roslyn Dakin[1*] roslyn.dakin@gmail.com, Paisley Clunis[1*] PaisleyClunis@cmail.carleton.ca, Emily M. Cornthwaite[1] EmilyCornthwaite@cmail.carleton.ca, and T. Brandt Ryder[2,3] rydert@si.edu

*co-first authors

[1]Department of Biology
Carleton University
1125 Colonel By Drive
Ottawa, Ontario, Canada K1S 5B6

[2]Migratory Bird Center
Smithsonian Conservation Biology Institute
National Zoological Park
Washington, DC, USA 20013

[3]Current address:
Bird Conservancy of the Rockies
Fort Collins, Colorado, USA 80525

To whom correspondence should be addressed:
Roslyn Dakin
roslyn.dakin@gmail.com

KEYWORDS: dynamic network, directed network, manakin, social connectivity, reciprocity, bidirectional, sociality

LAY SUMMARY

When animals are free to choose their partners, how does reciprocity influence social dynamics? In the wire-tailed manakin, males form coalitions with other males to help attract mates. We tracked male social networks and found that some males form reciprocal partnerships where each male leaves its territory to display at the other's territory. These bidirectional partnerships are particularly strong and stable, indicating that reciprocity is associated with stability in complex social networks.

38 **ABSTRACT**

Social reciprocity is thought to be the most important driver of cooperation among non-kin, but

40 its effects within complex social networks where individuals choose their partners have not been

studied. Here, we test whether reciprocation can explain social dynamics in a bird species where

42 males form coalitions to court females, the wire-tailed manakin (*Pipra filicauda*). In our study

population, approximately a third of male-male social interactions involve one male leaving his

44 territory to interact and display with the owner of another territory; this directional behaviour

provides benefits to the recipient at a cost to the visiting male. We characterize social ties among

46 pairs of territory holders according to the directionality of their interactions, and show here that

territory holding males engaged in far more reciprocated (bidirectional) partnerships with other

48 territory holders than expected by chance. Reciprocated partnerships were also stronger (i.e., the

partners interacted more frequently) than non-reciprocated partnerships, controlling for spatial

50 proximity of the territories. An individual-level analysis revealed that a male's social

contribution to a given partner was predictive of the number of social interactions he received

52 from that same partner. Finally, we show that reciprocation predicted the long-term stability of

social partnerships one year later. Together, these results demonstrate that reciprocity is

54 associated with stability in complex social networks where partner choice is flexible.

Introduction

58 Helping behavior among non-kin is found in many taxa, including mammals, birds, fish, and

invertebrates (Hart and Hart 1992; McDonald and Potts 1994; Griffiths and Magurran 1997;

60 Backwell et al. 1998; DuVal 2007; Rutte and Taborsky 2007; Brucks and von Bayern 2020;

Moore et al. 2020). One of the primary mechanisms thought to drive the evolution of cooperative

62 behaviour in non-kin is social reciprocity (Trivers 1971; Roberts and Sherratt 1998; Taborsky et

al. 2016). Recently, several pioneering studies have used careful experiments to demonstrate that

64 animals can both detect and respond to help from unrelated social partners (Rutte and Taborsky

2007; Cheney et al. 2010; Brucks and von Bayern 2020; Carter et al. 2020; Engelhardt and

66 Taborsky 2022). In nature, social relationships often occur in the context of complex social

networks, where individuals are free to choose their partners, and where social structure changes

68 dynamically through time (Hart and Hart 1992; Croft et al. 2008; Fehl et al. 2011; Ilany et al.

2015). In natural social systems, an individual's social investment into one partner (or social tie)

70 often occurs at the cost to other partnerships and to self-maintenance behaviour (e.g., foraging,

reproduction). Consequently, social reciprocity is predicted to drive long-term patterns of group-

72 level social structure and stability (Dakin and Ryder 2020; Yamamoto and Ishibashi 2022).

Despite the well-known importance of reciprocity in evolutionary and ecological theory (Fehl et

74 al. 2011), previous studies have not yet examined how mutual social investment affects

dynamics within a complex social network.

76   Here, we test the hypothesis that mutual social investment can explain social dynamics in

a complex social system of wire-tailed manakins *Pipra filicauda* (Ryder et al. 2008). This

78 species is part of a large clade of manakins characterized by elaborate male displays that are

considered to be some of the most complex of all passerine birds (Schwartz and Snow 1978;

80   Prum 1990; Heindl 2002; Ryder et al. 2008). In the wire-tailed manakin, males form cooperative

partnerships with non-kin and perform coordinated displays at specific display perches within

82   their lek territories. A typical male will have several social partners that he displays with

repeatedly, within and across breeding seasons (Ryder et al. 2008; Ryder et al. 2011; Dakin and

84   Ryder 2020). Previous studies have shown that an individual male's social centrality within the

broader male coalition network is a strong predictor of his ability to acquire a territory and

86   achieve long-term siring success (Ryder et al. 2008; Ryder et al. 2009). Male-male partnerships

in this system are also temporally stable across multiple years (Ryder et al. 2011; Dakin and

88   Ryder 2020), as observed in other manakins (Trainer 2002).

In the wire-tailed manakin, there are two male status classes: the dominant males who

90   hold territories, and the subordinate floater males who do not yet hold a territory (Heindl 2002;

Ryder et al. 2008; Ryder et al. 2009; Ryder et al. 2011). Although most social interactions occur

92   between a floater and a territory-holder, approximately a third of social interactions involve one

male leaving his territory to interact and display with the owner of another territory in a directed

94   interaction (Dakin and Ryder 2020). Importantly, the visiting male in this scenario takes on a

crucial cost: by leaving his own territory, he foregoes potential mating opportunities and

96   beneficial social ties that could have occurred there (Ryder et al. 2008; Ryder et al. 2009). On the

other hand, the recipient male in this scenario is expected to gain a fitness benefit, because social

98   displays on a male's own territory are associated with increased siring success (Ryder et al.

2008; Ryder et al. 2009). These directed interactions provide an opportunity to investigate how

100   often reciprocity occurs, and to test the prediction that reciprocity is associated with stability in

social network dynamics (Dakin and Ryder 2020).

We categorized social partnerships among territory-holding wire-tailed manakins as either mutually reciprocal or one-directional (Fig. 1A). A mutually reciprocal partnership is a

bidirectional relationship in which the two territory-holders interacted with each other mutually on their two respective territories. One-directional partnerships are those in which only one of

the territorial males visited the other male's territory to interact socially with him, and the reverse did not occur. We determined the directionality of manakin partnerships over both weekly and

annual timescales. Our analysis here evaluates the following four questions: (1) Do reciprocated cooperative partnerships occur more often than expected by chance, based on individual

spatiotemporal behavior? (2) Are reciprocated partnerships preferred, as defined by having a higher social interaction frequency, as compared to one-directional partnerships? (3) At an

individual level, does the magnitude of one male's social contribution to a given partner predict the contribution he received from that partner? (4) Are reciprocated partnerships more

temporally stable in the long-term than one-directional partnerships? Together, these analyses allow us to test whether reciprocity is a mechanism of social preference and temporal stability

within animal social networks.

METHODS

(a) Field work

120   We studied a population of wire-tailed manakins at the Tiputini Biodiversity Station in Orellana Province, Ecuador (0º 38' S, 76º 08' W, ~20m elevation). Field work took place over three

122   breeding seasons (December to March) from 2015 to 2018. At the beginning of each field season, wire-tailed manakins were captured using mist nets and tagged with a unique

124   combination of color bands as well as a coded nanotag (NTQB-2, Lotek Wireless; 0.35 g).

Nanotags transmitted a unique VHF radio signal throughout the field season. We sampled males

126   from 11 leks at the field site, with lek sizes ranging from 4-14 territories (and typically, the

number of males using a lek is about 2x the number of territories). To record male social

128   interactions on lek territories, we placed proximity data-loggers (SRX-DL800, Lotek Wireless)

within each territory on a lek for a recording session of ~6 consecutive days (± SD 1 day; loggers

130   were active between 6:00 to 16:00 each day). Each lek was monitored for 3–4 recording sessions

per field season, distributed approximately evenly throughout the reproductive period from

132   December to March. Territory ownership was determined through direct observation and

confirmed in the proximity data. To determine the distance between territories, GPS latitude and

134   longitude coordinates were recorded at a location near the display perch within each territory. In

total, we monitored between 48–54 territory-holders per year (see Table S1).

(b) Social data

138   We used an algorithm described in (Dakin and Ryder 2018) to identify social interactions

between two male manakins within a display territory. The algorithm defined a social interaction

140   to have occurred when two males pinged a data-logger within a threshold of 45 s and 10 signal

strength units, which indicates spatial proximity of two males < 5 m of each other near the

142   display perch (Dakin and Ryder 2018; Dakin et al. 2021). This method has been shown to

reliably detect social interactions between male wire-tailed manakins in ground-truthing

144   experiments (Ryder et al. 2012; Dakin and Ryder 2018). All data processing and analyses were

performed in R 3.6.3 (R Core Team 2020). Here, we focus on interactions of interest between

146   pairs of territory holding males. We define a directed social interaction among territory holders

as occurring when two territory-holders engaged in a social interaction on one of their respective

148 territories (Pinter-Wollman et al. 2014). A partnership between two territory-holders was defined

as one-directional if all of their social interactions during a given time period occurred on only

150 one of their respective territories. A partnership was defined reciprocated (or bidirectional) if the

two males interacted on both of their territories. We computed both directed and undirected edge

152 weights based on the frequencies of these social interactions. The undirected edge weight was

the total frequency of social interactions for a dyad within a given time period (either weekly or

154 annual); the directed edge weights refer to the frequencies occurring on each recipient male's

territory. An important indicator of directional symmetry for reciprocated partnerships is the

156 minimum edge weight that occurred in both directions. Hence, we defined an additional metric

that we refer to as the "mutual edge weight", defined as the smaller of the two directed edge

158 weights for a given dyad. Recent experiments have established that spatial proximity is an

important driver of reciprocal relationships (Razik et al. 2022). To consider spatial distances as a

160 potential determinant of behavior, we estimated the Euclidean spatial distance between each pair

of territories from their latitude and longitude coordinates using the sp package 1.4-2 (Pebesma

162 and Roger 2020) in R.

164 *(c) Raw data stream permutation*

We compared the occurrence of reciprocated partnerships with the number expected to occur by

166 chance using a null model of territory visitation (Farine 2017; Dakin et al. 2021). The null model

was based on a permutation of the territory visits performed by each male, as described in (Dakin

168 et al. 2021). Note that we use "territory visit" here to refer to the detection of an individual male

on *any* territory, independent of social context. The null permutation preserved each male's

170 spatial and temporal distribution of territory visits, including the dates of each visit, and the visit

durations, but randomly sampled the visit start times from the distribution of actual start times

172   observed in the data (Dakin et al. 2021). Hence, the null model preserved each male's

spatiotemporal behavior (i.e., the same territories were visited, the same number of times, and for

174   the same duration), but ensured that the exact timing of visits was independent of the presence of

other males. This approach previously demonstrated that manakin social partnerships have much

176   greater undirected edge weights than expected (Dakin et al. 2021). For the present study, we used

100 permuted null datasets to determine the expected number and directionality of annual

178   partnerships among territory holders, and their edge weights (undirected and mutual; based on

frequency of social interactions, as defined above). We then compared these expected values

180   from the permutation analysis with the actual values observed in the manakin data.

182   *(d) Predictors of whether a partnership was reciprocated*

We used multilevel models at the dyad level to determine the factors that could predict whether

184   or not a partnership was reciprocated within a particular recording session. The response variable

for this analysis was the partnership type (reciprocated vs. one-directional), modelled in a

186   binomial analysis in the lme4 package 1.1-23 (Bates et al. 2020) in R. Because this analysis

focused on a dyad-level property, the random effect accounted for the dyad identity, based on the

188   combined identities of the two participating males. To examine the factors that predict

reciprocation, we examined two fixed effect predictors: (i) the undirected edge weight (log-

190   transformed) and (ii) the spatial distance between the two males' territories. We also accounted

for fixed effects of field season (three levels, one for each year) and sampling effort, as defined

192   by the total number of hours that the two territories were recorded. To ensure that this analysis

only considered partnerships that could have been reciprocated, we limited it to the 377 session-

194   partnerships that had an edge weight ≥2 (the minimum required for reciprocation), and where

both males' territories were monitored throughout the season.

(e) Magnitude of individual contributions in reciprocated partnerships

198   The goal of this individual-level analysis was to describe the factors that predict a given

territory-holder's social contribution to his territorial partners within a recording session. Hence,

200   we use 'strength' here to refer to the social interaction frequency of a particular individual (i.e.,

the sum of the edge weights for that individual's node in the social network). This analysis was

202   run using a multilevel model in the lme4 package in R, with the response variable as the relative

amount that a focal male contributed to a specific partner. To calculate this variable (relative

204   amount contributed), we took the focal male's social interactions contributed to partner x_i,

divided by the focal male's total social interactions away from his own territory. Note that

206   "contributed" here refers to times when the focal male went to the partner's territory and

interacted with him there. The main predictor of interest was the relative amount that the focal

208   male received on his territory from partner x_i (i.e., strength received from x_i expressed relative to

the focal male's total social interactions received). Thus, this analysis tested whether the focal

210   male's social contribution to a particular partner was correlated with the amount received

(reciprocal social investment) from that partner. Additional fixed effects accounted for field

212   season (three levels, one for each year), the spatial distance between territories, and sampling

effort. We also included the duration of visits by floater males (i.e., males that do not hold

214   territories) to the focal male, and the duration of visits by floater males to partner x_i as two

additional fixed effect predictors (both expressed as the natural log of the total number of floater

216   pings). This allowed us to account for the possibility that social contributions among territory

holders may depend on the activity of these other male social partners within the lek. The

218   identities of the focal and partner male were included as crossed random effects. We limited this

analysis to reciprocated session-partnerships, represented in a sample size of 326 contributions

220   within 163 partnerships among 72 males.

222   *(f) Predictors of long-term partnership stability*

To evaluate whether reciprocity leads to long-term stability of male-male partnerships, we

224   analyzed the interaction frequency of a partnership in the following year, after an initial

designation of partnership type. The response variable was the partnership's undirected edge

226   weight in the second year, $t + 1$, analyzed in a Gaussian model in lme4. To meet the assumptions

of the Gaussian model, the response variable was log-transformed. To test whether previous

228   reciprocation could predict long-term dynamics, the model included a fixed effect of partnership

directionality in the initial year t (either reciprocated or one-directional). The other fixed effects

230   in the model included field season, the spatial distance between territories, and the edge weight

of the partnership in the initial year t (log-transformed). Hence, this analysis tested whether edge

232   weight in year $t + 1$ was associated with previous reciprocation in year t, after accounting for the

previous edge weight. Because this analysis focused on a dyad-level property, the random effect

234   accounted for identity of the dyad as describe above. The criteria for inclusion in this analysis

was that the year t edge weight had to be ≥ 2 (i.e., the partnership had to have occurred often

236   enough that it could have been reciprocated in year t), and that both males had to be tagged and

monitored in the subsequent field season; the resulting sample size was 109 annual-partnerships.

RESULTS

240 *(a) Occurrence of reciprocated partnerships*

We recorded over 9,500 directed social interactions between two territorial male wire-tailed

242 manakins and identified 122 bidirectional annual-partnerships that were reciprocated at least

once. The number of reciprocated partnerships was 56% more than expected, based on a null

244 model (Fig. 1c). By contrast, we observed far fewer one-directional partnerships among territory

holders than expected by chance (Table S1). This indicates that reciprocity is not occurring at

246 random; rather, it is a common feature of the wire-tailed manakin social system. The average

territory holder had about 4 ($\pm$ 2.0 SD) territorial partners per year, 2 ($\pm$ 1.4 SD) of which were

248 reciprocated, and 2 ($\pm$ 1.6 SD) of which were one-directional.

The mutual edge weights of reciprocated partnerships were typically 6-to-14-fold greater

250 than expected in the null model, demonstrating that these social partnerships are far more

symmetrical than expected (Fig. 1d). For example, 58 partnerships had an annual mutual edge

252 weight $\geq$ 5, a level of symmetry that was never observed in the null model (Fig. 1d). Moreover,

we found that reciprocated social partners interacted 10-to-20-fold more than expected by chance

254 (based on undirected edge weights; see Table S1). One-directional social partners also interacted

more than expected, but to a much lesser extent, with only a 3-to-7-fold difference as compared

256 to the expected values (see Table S1).

258 *(b) Predictors of whether a partnership was reciprocated*

Reciprocation was more likely to occur within partnerships that interacted frequently, as

260 compared to partners that interacted rarely (Fig. 2). This strong association between

reciprocation and edge weight also holds while accounting for the spatial proximity of the two

262   territories, which was another significant predictor of reciprocation. Pairs with closer territories

were more often reciprocal (Table S2).

(c) Magnitude of individual contributions in reciprocated partnerships

266   There were two main factors that predicted a male's social contribution to a given partner. The

first predictor was the relative social contribution received from that partner: focal males directed

268   more of their social interactions toward partners they had received more from (Fig. 3; Table S3).

The second major predictor was the spatial proximity of their territories: males contributed more

270   to closer neighbors (Table S3). The magnitude of social contribution was not significantly

associated with the frequency of floater male occurrence at either territory (Table S3).

(d) Predictors of long-term partnership stability

274   Partnerships that were reciprocated in year *t* maintained far greater edge weight (interaction

frequency) the following year than did one-directional partnerships (Fig. 4). This result held even

276   when accounting for territory proximity and the previous year's edge weight, both of which were

also positive predictors of edge weight in the following year (Table S4).

DISCUSSION

280   Reciprocation has been proposed to drive the evolution of cooperation and determine the

structure and dynamics of partner choice within cooperative social networks (Fehl et al. 2011;

282   Dakin and Ryder 2020). We used an analysis of directed social ties to test how reciprocation

predicts social network dynamics in a wild population of cooperative birds, where individuals

284   were free to choose their social partners. We found that the social networks of territory holding

male manakins contained many more reciprocated (bidirectional) partnerships than expected, and

286   fewer one-directional partnerships than predicted by a null model of individual spatiotemporal

behaviour. Moreover, mutual edge weight, a key measure of social symmetry, was also found to

288   be much higher than expected. Our analysis demonstrates that there are two main predictors of

whether or not a social interaction was reciprocated: the overall frequency of social interactions

290   between the two partners, and the spatial proximity of their territories. Male-male pairs that were

located physically closer to each other, and those with a higher frequency of social interaction,

292   were more likely to have a reciprocated relationship. Together, these findings indicate that

reciprocation is either a cause of, or a marker of, social preference. They also indicate that spatial

294   structure can also influence the costs and benefits of an individual's social choices (Backwell et

al. 1998; Webber and Vander Wal 2018; Bonar et al. 2020).

296   Previous studies of wire-tailed manakins have shown that a male's social connectivity on

his own territory is correlated with reproductive success (Ryder et al. 2008; Ryder et al. 2009).

298   Here, we show that a male manakin's social contribution to another territory-holding partner was

positively correlated with the number of social interactions he received from that partner,

300   independent of spatial proximity. Collectively, these findings suggest that males who form strong

social bonds with other territory holders may receive delayed benefits through reciprocated

302   social interactions that "return the favour". This result adds to the growing literature that animals

can respond in-kind to others' social contributions (Rutte and Taborsky 2007; Cheney et al.

304   2010; Brucks and von Bayern 2020; Carter et al. 2020; Mishra et al. 2020; Lalot et al. 2021).

Importantly, the present findings demonstrate that reciprocation can drive social investment in

306   natural systems where individuals allocate their time towards multiple partnerships within a

complex network.

308   How does reciprocity influence the long-term dynamics of social ties? We found that

reciprocated cooperative partnerships retained greater social interaction frequencies one year

310   later, even after accounting for positive effects of the initial social interaction frequency and

territory spatial proximity. This finding indicates that cooperative relationships that are

312   reciprocal are more likely to be maintained at a high rate in the future. Although we lack the

ability to experimentally manipulate social bonds, this longitudinal analysis is consistent with

314   two non-exclusive hypotheses, as mentioned above: (1) reciprocation may directly enhance

social preference for particular partners, and (2) reciprocation may be a marker of (or

316   consequence of) more general social compatibility between two individuals. Under the first

hypothesis, reciprocation in the past is a direct cause of future social preference (Roberts and

318   Sherratt 1998). The fact that reciprocation predicted future dynamics after accounting for the

previous social interaction frequency is suggestive of this first possibility, and is consistent with

320   recent controlled experiments in other species (Carter et al. 2020). An additional possibility is

that reciprocation in year t may be a consequence of more general social compatibility (Busia et

322   al. 2017; Ellis et al. 2019), which may in turn drive the enhanced stability of particular social

bonds. Although we are unable to experimentally disentangle these two possibilities with the

324   present data, future studies may be able to address this by the comparison of data on the fine-

scale temporal dynamics of natural systems with carefully constructed simulation models.

326   Our results raise further questions about the behavioral mechanisms underlying

reciprocation in the wild. For example, the results of our individual-level analysis suggest that

328   wire-tailed manakins may recognize and distinguish the quality of cooperative social

partnerships with particular individuals (Cheney et al. 2010). This raises the question of whether

330   individuals within complex social networks can allocate their social contributions strategically

(Ellis et al. 2019). Although not a main focus of our study, our results also emphasize the

332   importance of physical proximity in determining the directionality and dynamics of social

network ties (Webber and Vander Wal 2018; Webber et al. 2023). Because a positive correlation

334   exists between proximity and the probability of an initial interaction occurring within a pair,

spatially constrained social systems may have a greater potential for reciprocated interactions.

336   Historically, research on the ecology of cooperation among non-kin has focused on

distinguishing reciprocal altruism from other mechanisms, such as by-product mutualism

338   (Nowak 2006). Recent studies highlight how reciprocation can often be graded and dynamic in

time (Roberts and Sherratt 1998; Carter et al. 2020). We show here that reciprocation occurs

340   often in a complex social network, that it is either a cause of or a marker of social preference,

and that it is associated with long-term social stability of cooperative partnerships (Fehl et al.

342   2011; Dakin and Ryder 2020). More broadly, social network structures are an important

framework for understanding the movement ecology and behaviour in the wild. We propose that

344   reciprocal network ties may often be stronger and more stable, and that the emergent properties

of social networks will often depend on the composition and symmetry of these bidirectional

ties.

FIGURE LEGENDS

Figure 1. Social relationships can be one-directional or reciprocated. (A) Male manakins form cooperative partnerships with other males to perform coordinated courtship displays. A
reciprocated partnership occurs when two territory holders interact with each other mutually on their two respective territories. In the network diagrams, the circles represent each male's
territory, and the arrows represent directional social interactions that originate from the visitor and point toward the recipient. (B) Example of a directed social network among 10 territory-
holding males. (C) In total, we observed 122 reciprocated annual partnerships (indicated in red), which is 56% more than expected based on a null permutation. (D) The mutual edge weights of
reciprocated partnerships were also far greater than expected by chance. The boxplots show mutual edge weights with lines connecting the observed value for each partnership with its null
expectation (n = 122 reciprocated annual partnerships). See also Table S1.

Figure 2. Reciprocated dyads interact more frequently. Reciprocation was more likely to occur among partners that interacted more frequently with each other (greater undirected edge
weight). The positive association between reciprocation and edge weight holds while accounting for the spatial proximity of the two males' territories, field season, recording duration, and the
identities of the two partner males. The number of session partnerships in each column is provided along the top of the graph (total = 377 session partnerships with an edge weight >= 2,
involving 80 males). Error bars show the 95% confidence intervals for the proportion estimated in each column. Note that while edge weight is binned here for visualization purposes, it was
treated as a continuous variable in the statistical analysis (see Table S2).

Figure 3. A male's contribution to a given social partner is correlated with the amount he received. Following the diagram on the top left, this analysis evaluated whether a focal male's
contribution to partner x was predicted by the relative number of social visits received from partner x. The y-axis shows partial residuals for a male's social contribution while controlling for
other factors in the analysis, including proximity, field season, recording duration, floater male visits, and the identities of the two partner males (see Table S3 for details). Note that this
analysis is limited to session partnerships that were reciprocated (n = 326 individual contributions within 163 session partnerships, among 72 males).
380
Figure 4. Reciprocated dyads maintain greater stability through time. Social dyads that were
382 reciprocated in year t had significantly stronger edge weights (interaction frequencies) in year $t + 1$ (n = 109 annual partnerships among 38 males). This positive association holds while
384 accounting for the undirected edge weight in year t, spatial proximity of the two males' territories, field season, and the identities of the two partner males (see Table S4). Note that this
386 analysis is limited to dyads that interacted at least twice in year t and that were both tagged and monitored over the following year.

388 ETHICS STATEMENT
All research was approved by the Smithsonian ACUC (protocols #12-23, 14-25, and 17-11) and
390 the Ecuadorean Ministry of Environment (MAE-DNB-CM-2015-0008).

392 DATA ACCESSIBILITY
All data and R scripts are available at: https://figshare.com/s/f225fefb1ec4906e0353
394 The repository will be made public when the final version of the study is published.

396 AUTHORS' CONTRIBUTION
All authors designed the study. TBR collected the data. RD and PC analyzed the data and wrote
398 the manuscript. All authors edited the manuscript.

400 COMPETING INTERESTS
We have no competing interests.
402

FUNDING
404 Supported by the National Science Foundation (NSF) IOS 1353085, the Smithsonian Migratory
Bird Center, the Natural Sciences and Engineering Research Council of Canada (NSERC), and
406 Carleton University.

408 ACKNOWLEDGEMENTS
We thank Ben Vernasco, Camilo Alfonso, Brent Horton, Ignacio Moore, Brian Evans, David and
410 Consuelo Romo, Kelly Swing, Diego Mosquera, Gabriela Vinueza, and the Tiputini Biodiversity
Station of the Universidad San Francisco de Quito.

412 **REFERENCES**

414 Backwell P, Jennions M, Passmore N, Christy J. 1998. Synchronized courtship in fiddler crabs. Nature. 391(6662):31–32. doi:10.1038/34076.

416 Bates D, Maechler M, Bolker B, Walker S, Haubo R, Christenson B. 2020. lme4: linear mixed-effects models using 'Eigen' and S4. https://cran.r-project.org/web/packages/lme4/index.html.

418 Bonar M, Lewis KP, Webber QMR, Dobbin M, Laforge MP, Vander Wal E. 2020. Geometry of the ideal free distribution: individual behavioural variation and annual reproductive success in
420 aggregations of a social ungulate. Ecol Lett. 23(9):1360–1369. doi:10.1111/ele.13563.

Brucks D, von Bayern AMP. 2020. Parrots voluntarily help each other to obtain food rewards.
422 Current Biology. 30(2):292-297.e5. doi:10.1016/j.cub.2019.11.030.

Busia L, Schaffner CM, Aureli F. 2017. Relationship quality affects fission decisions in wild
424 spider monkeys (*Ateles geoffroyi*). Ethology. 123(6–7):405–411. doi:10.1111/eth.12609.

Carter GG, Farine DR, Crisp RJ, Vrtilek JK, Ripperger SP, Page RA, Carter GG, Farine DR,
426 Crisp RJ, Vrtilek JK, et al. 2020. Development of new food-sharing relationships in vampire bats. Current Biology.:1–5. doi:10.1016/j.cub.2020.01.055.

428 Cheney DL, Moscovice LR, Heesen M, Mundry R, Seyfarth RM. 2010. Contingent cooperation between wild female baboons. Proc Natl Acad Sci U S A. 107(21):9562–9566.
430 doi:10.1073/pnas.1001862107.

Croft DP, James R, Krause J. 2008. Exploring Animal Social Networks. Princeton University
432 Press.

Dakin R, Moore IT, Horton BM, Vernasco BJ, Ryder TB. 2021. Testosterone□mediated
434 behaviour shapes the emergent properties of social networks. Journal of Animal Ecology. 90(1):131–142. doi:10.1111/1365-2656.13305.

436 Dakin R, Ryder TB. 2018. Dynamic network partnerships and social contagion drive cooperation. Proceedings of the Royal Society B: Biological Sciences. 285:20181973.
438 doi:10.1098/rspb.2018.1973.

Dakin R, Ryder TB. 2020. Reciprocity and behavioral heterogeneity govern the stability of social
440 networks. Proceedings of the National Academy of Sciences. 117(6):2993–2999. doi:10.1073/pnas.1913284117.

442 DuVal EH. 2007. Adaptive advantages of cooperative courtship for subordinate male lance-tailed manakins. American Naturalist. 169(4):423–432. doi:10.1086/512137.

444 Ellis S, Snyder-Mackler N, Ruiz-Lambides A, Platt ML, Brent LJN. 2019. Deconstructing sociality: The types of social connections that predict longevity in a group-living primate.

446   Proceedings of the Royal Society B: Biological Sciences. 286(1917):20191991.
       doi:10.1098/rspb.2019.1991.

448   Engelhardt SC, Taborsky M. 2022. Food-exchanging Norway rats apply the direct reciprocity
       decision rule rather than copying by imitation. Anim Behav. 194:265–274.
450   doi:10.1016/j.anbehav.2022.09.005.

       Farine DR. 2017. A guide to null models for animal social network analysis. Methods Ecol Evol.
452   8(10):1309–1320. doi:10.1111/2041-210X.12772.

       Fehl K, van der Post DJ, Semmann D. 2011. Co-evolution of behaviour and social network
454   structure promotes human cooperation. Ecol Lett. 14(6):546–551. doi:10.1111/j.1461-
       0248.2011.01615.x.

456   Griffiths SW, Magurran AE. 1997. Schooling preferences for familiar fish vary with group size
       in a wild guppy population. Proceedings of the Royal Society B: Biological Sciences.
458   264(1381):547–551. doi:10.1098/rspb.1997.0078.

       Hart BL, Hart LA. 1992. Reciprocal allogrooming in impala, *Aepyceros melampus*. Anim Behav.
460   44(6):1073–1083. doi:10.1016/S0003-3472(05)80319-7.

       Heindl M. 2002. Social organization on leks of the wire-tailed manakin in southern Venezuela.
462   Condor. 104(4):772–779. doi:10.1650/0010-5422(2002)104[0772:soolot]2.0.co;2.

       Ilany A, Booms AS, Holekamp KE. 2015. Topological effects of network structure on long-term
464   social network dynamics in a wild mammal. Ecol Lett. 18(7):687–695. doi:10.1111/ele.12447.

       Lalot M, Liévin-Bazin A, Bourgeois A, Saint Jalme M, Bovet D. 2021. Prosociality and
466   reciprocity in capybaras (Hydrochoerus hydrochaeris) in a non-reproductive context.
       Behavioural Processes. 188:104407. doi:10.1016/j.beproc.2021.104407.

468   McDonald DB, Potts WK. 1994. Cooperative display and relatedness in a lek-mating bird.
       Science (1979). 266(5187):1030–1032.

470   Mishra PS, Pal A, Velankar AD, Kumara HN, Singh M. 2020. Do males bond? A study of male-
       male relationships in Nicobar long-tailed macaques Macaca fascicularis umbrosus. J Biosci.
472   45(1):22. doi:10.1007/s12038-020-9995-y.

       Moore BL, Connor RC, Allen SJ, Krützen M, King SL. 2020. Acoustic coordination by allied
474   male dolphins in a cooperative context. Proceedings of the Royal Society B: Biological Sciences.
       287(1924):20192944. doi:10.1098/rspb.2019.2944.

476   Nowak MA. 2006. Five rules for the evolution of cooperation. Science (1979). 314(5805):1560–
       1563. doi:10.1126/science.1133755.

478   Pebesma E, Roger B. 2020. sp: Classes and Methods for Spatial Data.

Pinter-Wollman N, Hobson EA, Smith JE, Edelman AJ, Shizuka D, De Silva S, Waters JS, Prager SD, Sasaki T, Wittemyer G, et al. 2014. The dynamics of animal social networks: Analytical, conceptual, and theoretical advances. Behavioral Ecology. 25(2):242–255. doi:10.1093/beheco/art047.

Prum RO. 1990. Phylogenetic analysis of the evolution of display behavior in the neotropical manakins (Aves: Pipridae). Ethology. 84(3):202–231. doi:10.1111/j.1439-0310.1990.tb00798.x.

R Core Team. 2020. R: A Language and Environment for Statistical Computing.

Razik I, Brown BKG, Carter GG. 2022. Forced proximity promotes the formation of enduring cooperative relationships in vampire bats. Biol Lett. 18(4). doi:10.1098/rsbl.2022.0056.

Roberts G, Sherratt T. 1998. Development of cooperative relationships through increasing investment. Nature. 394(12):175–179.

Rutte C, Taborsky M. 2007. Generalized reciprocity in rats. PLoS Biol. 5(7):1421–1425. doi:10.1371/journal.pbio.0050196.

Ryder TB, Blake JG, Parker PG, Loiselle BA. 2011. The composition, stability, and kinship of reproductive coalitions in a lekking bird. Behavioral Ecology. 22(2):282–290. doi:10.1093/beheco/arq213.

Ryder TB, Horton BM, Van Den Tillaart M, De Dios Morales J, Moore IT. 2012. Proximity data-loggers increase the quantity and quality of social network data. Biol Lett. 8(6):917–920. doi:10.1098/rsbl.2012.0536.

Ryder TB, McDonald DB, Blake JG, Parker PG, Loiselle BA. 2008. Social networks in the lek-mating wire-tailed manakin (*Pipra filicauda*). Proceedings of the Royal Society B: Biological Sciences. 275(1641):1367–1374. doi:10.1098/rspb.2008.0205.

Ryder TB, Parker PG, Blake JG, Loiselle BA. 2009. It takes two to tango: reproductive skew and social correlates of male mating success in a lek-breeding bird. Proceedings of the Royal Society B: Biological Sciences. 276(1666):2377–2384. doi:10.1098/rspb.2009.0208.

Schwartz P, Snow D. 1978. Display and related behavior of the wire-tailed manakin. Living Bird. 17:51–78.

Taborsky M, Frommen JG, Riehl C. 2016. Correlated pay-offs are key to cooperation. Philosophical Transactions of the Royal Society B: Biological Sciences. 371(1687):1357–4316. doi:10.1098/rstb.2015.0084.

Trainer JM. 2002. The development of coordinated singing in cooperatively displaying long-tailed manakins. Behavioral Ecology. 13(1):65–69. doi:10.1093/beheco/13.1.65.

Trivers BYRL. 1971. The evolution of reciprocal altruism. Q Rev Biol. 46(1):35–57.

512   Webber QMR, Albery GF, Farine DR, Pinter☐Wollman N, Sharma N, Spiegel O, Vander Wal E, Manlove K. 2023. Behavioural ecology at the spatial–social interface. Biological Reviews.
514   98(3):868–886. doi:10.1111/brv.12934.

       Webber QMR, Vander Wal E. 2018. An evolutionary framework outlining the integration of
516   individual social and spatial ecology. Journal of Animal Ecology. 87(1):113–127. doi:10.1111/1365-2656.12773.

518   Yamamoto C, Ishibashi T. 2022. Flipper rubbing reciprocity and partner choice in common bottlenose dolphins. J Ethol. 40(1):49–59. doi:10.1007/s10164-021-00729-8.

Figure 1

(a)

(b)

(c) # Reciprocated

Null   Observed

40 60 80 100 120 140

(d)

Mutual edge weight
(# interactions)

200
100
50

10
5

Observed   Null

Figure 2

Figure 3

Figure 4