Resuscitation of soil microbiota after > 70-years of desiccation

Jun Zhao[1‡], Dongfeng Chen[1‡*], Wei Gao[1,2], Zhiying Guo[1,2], Zhongjun Jia[1], Marcela Hernández[3*]

[1]State Key Laboratory of Soil and Sustainable Agriculture. Institute of Soil Science, Chinese Academy of Sciences. Nanjing, 210008, Jiangsu Province, China

[2]University of Chinese Academy of Sciences, Beijing 100049, China

[3]School of Environmental Sciences, Norwich Research Park, University of East Anglia, Norwich, UK

‡These authors contribute equally to this work

* Correspondence:

Dongfeng Chen: dfchen@issas.ac.cn

Marcela Hernández: marcela.hernandez@uea.ac.uk

Running title: *Soil microbes after long-term desiccation*

Abstract

The abundance and diversity of bacteria in 24 historical soil samples under air-dried storage conditions for more than 70 years were assessed by quantification and high-throughput sequencing analysis of 16S rRNA genes. All soils contained a measurable abundance of bacteria varying from 10^3 to 10^8 per gram of soil and contrasting community compositions were observed in different background soils, suggesting that the bacteria detected were indigenous to the soil. Following a 4-week soil rewetting event, the bacterial abundance significantly increased in soils, indicating strong adaptation of soil bacteria to extreme osmotic change and high resuscitation potential of some bacteria over long periods of desiccation. *Paenibacillus, Cohnella* and two unclassified *Bacillales* genera within the phylum *Firmicutes* represented the most ubiquitously active taxa, which showed growth in the highest number of soils ($\geq$12 soils), while genera *Tumebacillus, Alicyclobacillus* and *Brevibacillus* in the phylum *Firmicutes* displayed the highest growth rates in soils (with >1000-fold average increase) following rewetting. Additionally, some *Actinobacteria* and *Proteobacteria* genera showed relatively high activity following rewetting, suggesting that the resilience to long-term desiccation and rewetting is a common trait across phylogenetically divergent microbes. The present study thus demonstrated that diversified groups of microbes are present and potentially active in historically desiccated soils, which might be of importance in the context of microbial ecology.

Key words: Long-term soil desiccation; Dry-rewet; 16S rRNA gene; Quantitative PCR; Amplicon sequencing; soil archives.

1 Introduction

Although frozen storage is widely accepted as the optimal method for preserving the chemical and biological components of soils, desiccation (i.e. air-drying) is more commonly used for large scale or cross-generational soil archives [1]. The desiccated soils, even after decades of storage, have been frequently used for analyzing the chemical composition [2-6], but the persistence of the biological components, especially the microbes in these soils remain largely untested. Microbes are famously known for tolerance to extreme environmental stresses, exemplified by growth of a few isolated strains after millions of years of dormancy under extremely low thermal conditions [7-10]. In respect to water stress, microbes have demonstrated strong resistance to long periods of drought based on both the cultivation of pure strains and the ecological investigation of samples from arid ecosystems [11, 12]. For instance, taxonomically diverse groups of bacteria are abundantly present and detectable in perennial drylands, including members from phyla of *Actinobacteria*, *Bacteroidetes* and *Proteobacteria* [13-15], suggesting that the ability of adaptation to long periods of drought is not restricted to a few particular taxa [16]. Compared to natural arid soils, the desiccated soils systematically archived in the laboratory have endured persistent

water depletion without intermittent rain precipitation as occurs in the field. The careful
curation of the soils in the laboratory prevents the occurrence of other major
environmental perturbations, such as vegetation and animal activities. Additionally, the
air-dried soil samples can also provide information on drought tolerance of microbial
communities that originated from non-arid ecosystems. Previous studies on historical
samples after decades of storage in air-dried conditions has indeed identified different
groups of bacterial and eukaryotic phylotypes, and suggested the possibility to use these
samples to detect systematic differences in soil microbial community composition [17-
22]. However, due to the low depth of sequencing, the information on microbial
abundance and diversity revealed from these studies is limited.

Water availability is a key factor controlling soil microbial activity and regulating
terrestrial ecosystem functioning. A decrease in soil moisture following desiccation can
cause high osmotic stress to the cells and prevent the diffusion of soluble nutrient
substrates [23], leading to dormancy of many soil microbes and subsequent decline in
many important biogeochemical processes including nitrification [24], denitrification
[25, 26] and methanotrophy [27, 28]. Some soil microbes can be reactivated soon after
the alleviation of a short period of drought stress, usually following a rewetting event
[24, 29-31]. Different phylotypes of microbes appear to respond differentially to
rewetting, with their abundance recovered at different rates from within one hour to
after a few days [32]. However, it is not known how different taxa respond to a rewet
event following a long-term, continuous desiccation.

In the present study, we acquired 24 historical soil samples collected from across
mainland China during the years of 1934-1939 from the Soil Archives at the Institute
of Soil Science, Chinese Academy of Sciences. By assessing the abundances (by
quantitative PCR of 16S rRNA genes) and community compositions (by
pyrosequencing) of bacteria in desiccated condition and after a rewet incubation, our
study aims to assess the resuscitation potentials of microbes through decades of
desiccation and identify the most responsive taxa in these soils.

2 Material and methods

2.1 Soil description and preparation

Samples were collected from seven provinces across the mainland of China during
the years of 1934-1939 (Fig. S1), and stored in air-dried conditions. The available
records by the collectors indicate widely spread field origins and/or properties for most
of the soils, including different locations, soil taxonomies and vegetation (Table 1).
Soon after each collection, a small part of the soil was kept in a reservation box in air-
dried conditions as part of the soil archives at the Institute of Soil Science, Chinese
Academy of Sciences. The weight of the archive soil was about 15-50 g for each sample
and completely dried by visual inspection. Due to the scarcity in quantity and historic
values of the samples, we obtained no more than 10 g of each soil for following
molecular analysis and did not perform tests on the soil chemical and physical
properties.

2.2 Rewetting microcosm incubation

For each microcosm, 1 g of soils were added into sterilized 5-ml centrifuge tubes and rewetted up to 30% of water holding capacity with sterile water. The tubes were sealed and incubated at 28°C for 28 days. Soil samples were destructively collected and frozen at -80°C for molecular analyses. All soils were subjected to rewetting microcosm incubations in triplicate, except for S01, S05, S23 and S24, which were incubated in duplicate due to a low soil quantity obtained.

2.3 DNA extraction

DNA was extracted from 1 g of soil, using FastDNA spin kit for DNA extraction (MP Biomedicals, USA), according to the manufacturer's instructions, and dissolved in a final volume of 50 μl sterile water. The quantity and quality of DNA extracts were measured using a NanoDrop ND-1000 UV-visible light spectrophotometer (NanoDrop Technologies, USA).

2.4 Real-time quantitative PCR

Real-time quantitative PCR (qPCR) was conducted on a CFX96 Optical Real-Time detection system (Bio-Rad Laboratories, Inc., Hercules, CA, USA) to determine the abundance of 16S rRNA genes in total DNA extracts, using universal PCR primers 515F (GTGCCAGCMGCCGCGG) and 907R (CCGTCAATTCMTTTRAGTTT) [33]. The thermal process started with 3 min of incubation at 95°C, followed by 40 cycles of 95°C for 30s, 55°C for 30s, and 72°C for 30s with plate read, and ended with a melt curve from 65 to 95°C with increments of 0.5°C per 5s. The real-time quantitative PCR standards were generated using plasmid DNA from one representative clone containing 16S rRNA genes, and a dilution series of standard template over eight orders of magnitude per assay was used. Amplification efficiencies ranged from 96% to 105%, with R^2 values of 0.992 to 0.996.

2.5 High-throughput sequencing and analysis

High-throughput sequencing was performed to investigate the bacterial community composition in soils. 16S rRNA gene fragments were amplified using the same universal 515F-907R primer pair but with an adaptor, key sequence, and unique tag sequence for each sample. The thermal conditions were as follows: 5 min of incubation at 94°C, followed by 30 cycles of 94°C for 30s, 54°C for 30s, and 72°C for 45s, with a 10 min extension at 72°C. The resulting PCR products were gel purified and combined in equimolar ratios into single tubes before pyrosequencing. The pyrosequencing was conducted on a Roche 454 GS FLX Titanium sequencer (Roche Diagnostics Corporation, Bran-ford, CT, USA).

All sequencing data were processed using multiple software packages. Reads from each sample were first sorted by specific tag sequences using Mothur software [34]. Sequences were further trimmed, and only reads of > 350 bp in length with an average quality score > 25 and no ambiguous base calls were included for subsequent analyses. The operational taxonomic units (OTUs) were clustered at 97% similarity cutoff by

138 UPARSE algorithm after the removal of chimeras using QIIME software [35]. A
139 representative sequence of each OTU was classified using the SILVA-132 16S rRNA
140 gene database (bootstrap confidence threshold of 80 %) in Mothur [34].

141 The raw sequencing data of all samples have been deposited in the European
142 Nucleotide Archive (ENA) with accession numbers ERR1520684 and ERR1520685.

2.6 Statistical analyses

The absolute abundance of total bacteria or a specific taxon in each soil microcosm
was calculated by multiplying the total 16S rRNA gene number (results from qPCR) by
the relative abundance of bacteria or the taxon (results of amplicon sequencing). The
nonparametric Wilcoxon test was used to determine the significance in difference in
bacterial abundance between all the dry and rewet soils. The fold change in abundance
of total bacteria or a specific taxon in each soil was further calculated as the ratio of
abundance after rewetting to that in the desiccated condition. The abundance of a taxon
was considered to increase in the soil when the fold change was significantly higher
than 1 following a t-test ($p < 0.05$), and the corresponding taxon was defined as active.

Heatmaps were constructed as previously described in Hernández et al., [36] using
the vegan package in R. The samples and OTUs were clustered according to Euclidean
distances between all Hellinger transformed data.

3 Results

3.1 Change in bacterial abundance

Total bacterial abundance was calculated based on the qPCR and high-throughput
sequencing of 16S rRNA genes. The 28-day rewet incubation led to significant increase
in bacterial abundance in most soils (n=24) (Fig. 1a). The total bacterial abundances in
soils were 1.9×10^3 to 1.7×10^8 per gram of dry weight soil (g^{-1} *d.w.s*), and reached
2.6×10^3 to 4.1×10^8 g^{-1} *d.w.s* following the rewetting incubation (Fig. S2). Specifically,
bacterial abundance showed significant increase in 18 soils ($P < 0.05$), remained
unchanged in 5 soils and only in 1 soil (S21) declined following the rewet. Notably,
bacterial abundance in two soils (S03 and S05) increased by > 100-fold and in another
seven soils (S04, S06, S08, S09, S12, S14 and S16) increased by > 10-fold following
the rewetting incubation (Fig. 1b). These results demonstrated that some bacterial
communities had high growth rates after regaining water in the soils. However, we
cannot rule out the possibility that a large portion of the detected genes in the dry soils
were residues from already dead cells, i.e. extracellular DNA [37, 38]. These cell
residues could potentially become accessible nutrients for the living cells in the soils
following rewetting [39], which might explain the decreased total bacterial abundance
in one of the soils (S21) after the incubation.

3.2 Change in bacterial composition

175 Taxonomic classification of 16S rRNA genes revealed a broad spectrum of
176 bacterial taxa in soils. A total of 32 bacterial phyla were identified in soils at the phylum

level. In the desiccated soils, the most abundant bacterial phyla identified were *Firmicutes* (averagely 55.4% of total bacteria), *Actinobacteria* (19.9%), *Proteobacteria* (9.7%), *Chloroflexi* (6.3%), *Acidobacteria* (1.7%) and *Planctomycetes* (1.1%) (Fig. 2 and S3a). Following the rewetting incubation, *Firmicutes* showed the highest growth rate and became the most abundant phylum in soils (80.4%), followed by *Proteobacteria* (9.4%), *Actinobacteria* (3.5%), *Acidobacteria* (1.6%) and *Chloroflexi* (1.0%) (Fig. 2 and S3b).

3.3 Identification active taxa

The activity of a specific bacterial taxon in each soil was assessed by the fold change in abundance after the rewetting incubation. At the genus level, a total of 759 taxa were identified in all soils, and 183 of them showed significant increase in abundance in at least one soil following the rewet (with fold change > 1 relative to the desiccated condition). Fig. 3 shows the top 21 genera with significant increases in abundance in more than three of the soils. *Paenibacillus*, *Cohnella* and two other unclassified *Bacillales* genera represented the most ubiquitously activated genera in soils, which showed increases in at least half of the soils following rewetting incubations (Fig. 3). On the other hand, *Bacillales* genera *Tumebacillus*, *Alicyclobacillus* and *Brevibacillus* showed the highest growth rate following rewetting, with on average >1,000-fold increase in abundance in these soils (Fig. 3). Although *Bacillales* comprised the most active microbial communities in our soils, other bacterial taxa, represented by one unclassified *Actinobacteria* genus, two *Gammaproteobacteria* genera (*Halomonas* and *Ralstonia*) and four genera belonging to *Firmicutes* order *Clostridia*, also showed marked growth in soils (Fig. 3).

Discussion

Our study described both the quantitative and compositional patterns of soil bacterial communities after continuous desiccation for decades. Phylogenetically diverse guilds of bacteria with measurable abundance were identified in these historical soils (Fig. 2 and S3a), demonstrating high resistance to the long-term desiccation. Driven by the stress from both water and nutrient deprivation for more than seven decades, the bacteria abundantly detected in the dry soils were most likely in a dormant state [23]. This was supported by the dominant presence of *Firmicutes* in most of the desiccated soils (Fig. 2), which consisted of *Paenibacillus*, *Cohnella, Bacillus* and *Clostridium* species (Fig. S3a) that are well known for producing resting bodies of endospores. In addition, in many soils, the frequent presence of taxa from *Actinobacteria*, *Proteobacteria* and *Chloroflexi* (Fig. S3a), which are less known for tolerance to extreme drought stress, implies the ubiquity of dormant life strategies against long-term desiccation across phylogenetically diverse microbial populations [16, 40]. This might be important in the maintenance of microbial biodiversity and the restoration of many ecosystem functions when conditions become favorable [41, 42].

Following the rewetting incubation, the growth of several genera, including *Paenibacillus* and *Cohnella* from the phylum *Firmicutes*, were stimulated in more than half of soils (Fig. 3). Some genera within the *Firmicutes* (*Tumebacillus*,

Alicyclobacillus and *Brevibacillus*) also showed the highest growth rate in many of the
soils after rewetting. It is noted that all these genera belonged to order *Bacillales* within
the *Firmicutes.* These endospore-forming bacteria thus represented the most active
microbes or rapidly resuscitated organisms in these soils. Culture-based studies on a
desert soil revealed growth of *Firmicutes* in cultivation medium, despite their low
relative abundance in the soil [13]. Similarly, *Firmicutes* in many of our soils (S01, S03,
S06, S09, S12 and S23) were not the most abundant group under desiccated conditions,
but became dominant after the incubation (Fig. 2). These results suggested that the
formation of endospores is not only beneficial for the survival of *Firmicutes* through
water deprivation, but also an efficient strategy for recovery of populations when the
environmental stress is relieved. Additionally, two genera within the
Gammaproteobacteria, *Halomonas* and *Ralstonia*, also showed marked growth after
the rewet (Fig. 3). Although members of *Halomonas* have not been reported to have
strong resilience to extreme drought-rewet occurrences, they are halotolerant and
capable of producing ectoine as osmolytes to maintain cell-osmotic balance [43]. The
ectoine can protect the cells against salinity, desiccation and high temperature [44],
which might facilitate cell survival in our soils following extreme moisture changes.
Similarly, disaccharide trehalose contribute to osmotic stress tolerance of *Ralstonia*
species [45], which might have implications for resilience to water stress.

It is noteworthy that *Actinobacteria* represent the second-most abundant group in
desiccated soils (Fig. 2), suggesting high resistance to long-term desiccation as
previously documented [13-15]. Some members of *Actinobacteria* contain genes for
trace-gas scavenging, including H_2 and CO, which can support a dormant lifestyle in
arid deserts [15]. However, the growth rate of *Actinobacteria* appears much lower than
Firmicutes and *Proteobacteria*, leading to sharp declines in average relative abundance
in many of the soils after rewetting (Fig. 2). Similar trends were also found in a short
period of a dry-rewet experiment, where the relative abundance of *Actinobacteria*
increased after desiccation but then decreased following a rewet [30, 46]. It is also
possible that the activity of *Actinobacteria* might be stimulated at the early stage of the
28-day incubation and outcompeted later by the stronger growth of *Firmicutes* and
Proteobacteria. A previous study defined *Actinobacteria* as rapid responders with fast
reactivation within one hour following rewetting, whereas *Firmicutes* and many
Proteobacteria classes (*Alpha-*, *Beta-* and *Gamma-proteobacteria*) were consistently
delayed responders [32]. This can be assessed by monitoring temporal changes in
relative abundance of different phyla or classes in soils following rewetting in future
studies.

Although all soil samples analyzed in the present study have been processed by the
same laboratory treatment (long-term air-dry and storage in the same room conditions
since 1939), it appeared that the microbial community compositions between different
background soils can still be clearly distinguished, suggesting that the key bacteria
identified in these soils should be indigenous to the soil field habitats rather than
originating from the laboratory environment. This gained further support from analysis
of some niche specialized taxa. For instance, members from class *Clostridia* (e.g. genus

Clostridium), consisting of well-documented obligate anaerobes, were frequently detected (16.3-62.2%, Fig. S3a) in the desiccated rice paddy soils (S03 and S13-16), which should have endured constant anaerobic conditions by irrigation. In addition, although archaea were not detected or were only detected at low relative abundance in most of the soils, the class *Halobacteria*, known for salt tolerance, was exclusively detected in the two coastal saline soils (S22 and S23) in our study, accounting for >99% of the archaea (Fig. S4). These results suggest that genetic fingerprints of some microbes in desiccated soils is preserved (either as intact cell component or dead cell residues) and can be a used to indicate important field history, such as land use and soil formation. These results collectively validate the value of desiccated soil samples for systematic study on differences in soil microbial community composition, as previously proposed [18].

5 Conclusion

The present study revealed a diversified guild of bacteria resistant to seven decades of desiccation in 24 historical soils and nearly 25% of the taxa (at genus level) still show capability of growth. A dissimilar response of different phylotypes to the rewetting was observed, confirming to previous study suggesting the resuscitation strategies of soil microbes may be a phylogenetically conserved ecological trait [32]. Considering long-term starvation of microbes by desiccation and the fact that only water was added for incubation in our study, we expect growth of more diverse microbes with the additional supply of nutrients, such as carbon and nitrogen. Moreover, the activity of microbes was assessed by only DNA-based molecular analyses, which cannot capture all the active microbes in the soils, especially for those that did not quickly propagate within our incubation time. In addition, in future studies microbial viability would be more accurately depicted using more sensitive molecular tools, e.g. rRNA molecules rather than genes as a better indicator for cell activity [47]. Other methods would also be of great potential in distinguishing growing, dormant and dead cells, e. g. isotope tracing [31, 48-50] and single-cell based imaging [51, 52]. The removal of extracellular nucleic acids should also be considered in the future study to validate our conclusions [37]. Finally, future studies targeting specific microbial groups could enhance our understanding of some important functional microbes and their response to extreme stress [53]. By adopting more sensitive techniques, the desiccated historical soils could serve as precious "fossils" for soil microbiologists to discover tenacious species and retrieve undocumented site-specific historical events.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We gratefully acknowledge the soil science predecessors of the Chinese Academy of

Sciences for the original soil collection, the Institute of Soil Science for devotion to historical sample archives and all the former faculties in charge of the soil archiving. This work was supported by the National Science Foundation of China [41701302 and 41530857].

References

[1] J. Dolfing, Y. Feng, The importance of soil archives for microbial ecology, Nature reviews. Microbiology, 13 (2015) 1.

[2] S. Meijer, C. Halsall, T. Harner, A. Peters, W. Ockenden, A. Johnston, K. Jones, Organochlorine pesticide residues in archived UK soil, ENVIRONMENTAL SCIENCE & TECHNOLOGY, 35 (2001) 1989-1995.

[3] I. Cousins, H. Kreibich, L. Hudson, W. Lead, K. Jones, PAHs in soils: contemporary UK data and evidence for potential contamination problems caused by exposure of samples to laboratory air, SCIENCE OF THE TOTAL ENVIRONMENT, 203 (1997) 141-156.

[4] W. Lead, E. Steinnes, J. Bacon, K. Jones, Polychlorinated biphenyls in UK and Norwegian soils: Spatial and temporal trends, SCIENCE OF THE TOTAL ENVIRONMENT, 193 (1997) 229-236.

[5] S. Meijer, T. Harner, P. Helm, C. Halsall, A. Johnston, K. Jones, Polychlorinated naphthalenes in UK soils: Time trends, markers of source, and equilibrium status, ENVIRONMENTAL SCIENCE & TECHNOLOGY, 35 (2001) 4205-4213.

[6] R. Alcock, C. Halsall, C. Harris, A. Johnston, W. Lead, G. Sanders, K. Jones, Contamination of environmental-samples prepared for PCB analysis, ENVIRONMENTAL SCIENCE & TECHNOLOGY, 28 (1994) 1838-1842.

[7] R. Cano, M. Borucki, Revival and identification of bacterial spores in 25- to 40-million-year-old Dominican amber, Science, 268 (1995) 1060-1064.

[8] R. Vreeland, W. Rosenzweig, D. Powers, Isolation of a 250 million-year-old halotolerant bacterium from a primary salt crystal Nature, 407 (2000) 897-900.

[9] K.D. Bidle, S. Lee, D.R. Marchant, P.G. Falkowski, Fossil genes and microbes in the oldest ice on earth, Proceedings of the National Academy of Sciences of the United States of America, 104 (2007) 13455-13460.

[10] S.T. Jaakkola, F. Pfeiffer, J.J. Ravantti, Q. Guo, Y. Liu, X. Chen, H. Ma, C. Yang, H.M. Oksanen, D.H. Bamford, The complete genome of a viable archaeum isolated from 123-million-year-old rock salt, Environmental microbiology, 18 (2016) 565-579.

[11] T.P. Makhalanyane, A. Valverde, E. Gunnigle, A. Frossard, J.B. Ramond, D.A. Cowan, Microbial ecology of hot desert edaphic systems, FEMS microbiology reviews, 39 (2015) 203-221.

[12] G. Zhu, S. Wang, C. Wang, L. Zhou, S. Zhao, Y. Li, F. Li, M.S.M. Jetten, Y. Lu, L. Schwark, Resuscitation of anammox bacteria after >10,000 years of dormancy, The ISME journal, 13 (2019) 1098-1109.

[13] A. Chanal, V. Chapon, K. Benzerara, M. Barakat, R. Christen, W. Achouak, F. Barras, T. Heulin, The desert of Tataouine: an extreme environment that hosts a wide diversity of microorganisms and radiotolerant bacteria, Environmental microbiology, 8 (2006) 514-525.

[14] S.A. Connon, E.D. Lester, H.S. Shafaat, D.C. Obenhuber, A. Ponce, Bacterial diversity in hyperarid Atacama Desert soils, Journal of Geophysical Research: Biogeosciences, 112 (2007) n/a-n/a.

[15] M. Ji, C. Greening, I. Vanwonterghem, C.R. Carere, S.K. Bay, J.A. Steen, K. Montgomery, T. Lines, J. Beardall, J. van Dorst, I. Snape, M.B. Stott, P. Hugenholtz, B.C. Ferrari, Atmospheric trace gases

support primary production in Antarctic desert surface soil, Nature, 552 (2017) 400-403.

[16] W. Borken, E. Matzner, Reappraisal of drying and wetting effects on C and N mineralization and fluxes in soils, Global Change Biology, 15 (2009) 808-824.

[17] S.Y. Moon-van der Staay, V.A. Tzeneva, G.W. van der Staay, W.M. de Vos, H. Smidt, J.H. Hackstein, Eukaryotic diversity in historical soil samples, FEMS microbiology ecology, 57 (2006) 420-428.

[18] J. Dolfing, A. Vos, J. Bloem, P.A. Ehlert, N.B. Naumova, P.J. Kuikman, Microbial diversity in archived soils, Science, 306 (2004) 813.

[19] I.M. Clark, P.R. Hirsch, Survival of bacterial DNA and culturable bacteria in archived soils from the Rothamsted Broadbalk experiment, Soil Biology and Biochemistry, 40 (2008) 1090-1102.

[20] M. Denobili, M. Contin, P. Brookes, Microbial biomass dynamics in recently air-dried and rewetted soils compared to others stored air-dry for up to 103 years, Soil Biology and Biochemistry, 38 (2006) 2871-2881.

[21] C. Knapp, J. Dolfing, P. Ehlert, D. Graham, Evidence of Increasing Antibiotic Resistance Gene Abundances in Archived Soils since 1940 ENVIRONMENTAL SCIENCE & TECHNOLOGY, 44 (2010) 580-587.

[22] L.A. Lewis, F.R. Trainor, Survival of Protosiphon botryoides (Chlorophyceae, Chlorophyta) from a Connecticut soil dried for 43 years, Phycologia, 51 (2012) 662-665.

[23] J. Schimel, T. Balser, M. Wallenstein, Microbial stress-response physiology and its implications for ecosystem function Ecology, 88 (2007) 1386-1394.

[24] C. Thion, J.I. Prosser, Differential response of nonadapted ammonia-oxidising archaea and bacteria to drying-rewetting stress, FEMS Microbiol Ecol, 90 (2014) 380-389.

[25] E.L. Aronson, M.L. Goulden, S.D. Allison, Greenhouse gas fluxes under drought and nitrogen addition in a Southern California grassland, Soil Biology and Biochemistry, 131 (2019) 19-27.

[26] A. Bollmann, R. Conrad, Influence of O2 availability on NO and N2O release by nitrification and denitrification in soils, Global Change Biology 4(1998) 387-396.

[27] A. Ho, C. Lüke, A. Reim, P. Frenzel, Resilience of (seed bank) aerobic methanotrophs and methanotrophic activity to desiccation and heat stress, Soil Biology and Biochemistry, 101 (2016) 130-138.

[28] S. Collet, A. Reim, A. Ho, P. Frenzel, Recovery of paddy soil methanotrophs from long term drought, Soil Biology and Biochemistry, 88 (2015) 69-72.

[29] M. Chen, M. Alexander, Survival of soil bacteria during prolonged desiccation, Soil Biology & Biochemistry, 5 (1973) 213-221.

[30] R.L. Barnard, C.A. Osborne, M.K. Firestone, Responses of soil bacterial and fungal communities to extreme desiccation and rewetting, The ISME journal, 7 (2013) 2229-2241.

[31] Z.T. Aanderud, S.E. Jones, N. Fierer, J.T. Lennon, Resuscitation of the rare biosphere contributes to pulses of ecosystem activity, Frontiers in microbiology, 6 (2015) 24.

[32] S.A. Placella, E.L. Brodie, M.K. Firestone, Rainfall-induced carbon dioxide pulses result from sequential resuscitation of phylogenetically clustered microbial groups, Proceedings of the National Academy of Sciences of the United States of America, 109 (2012) 10931-10936.

[33] S. Stubner, Enumeration of 16S rDNA of Desulfotomaculum lineage 1 in rice field soil by real-time PCR with SybrGreen™ detection, Journal of Microbiological Methods, 50 (2002) 155-164.

[34] P.D. Schloss, S.L. Westcott, T. Ryabin, J.R. Hall, M. Hartmann, E.B. Hollister, R.A. Lesniewski, B.B. Oakley, D.H. Parks, C.J. Robinson, J.W. Sahl, B. Stres, G.G. Thallinger, D.J. Van Horn, C.F. Weber, Introducing mothur: open-source, platform-independent, community-supported software for describing

388 and comparing microbial communities, Appl Environ Microbiol, 75 (2009) 7537-7541.

389 [35] J. Caporaso, J. Kuczynski, J. Stombaugh, K. Bittinger, F. Bushman, E. Costello, N. Fierer, A. Pena,

390 J. Goodrich, J. Gordon, G. Huttley, S. Kelley, D. Knights, J. Koenig, R. Ley, C. Lozupone, D. McDonald,

391 B. Muegge, M. Pirrung, J. Reeder, J. Sevinsky, P. Tumbaugh, W. Walters, J. Widmann, T. Yatsunenko, J.

392 Zaneveld, R. Knight, QIIME allows analysis of high-throughput community sequencing data, NATURE

393 METHODS, 7 (2010) 335-336.

394 [36] M. Hernández, R. Conrad, M. Klose, K. Ma, Y. Lu, Structure and function of methanogenic

395 microbial communities in soils from flooded rice and upland soybean fields from Sanjiang plain, NE

396 China, Soil Biology and Biochemistry, 105 (2017) 81-91.

397 [37] G. Pietramellara, J. Ascher, F. Borgogni, M.T. Ceccherini, G. Guerri, P. Nannipieri, Extracellular

398 DNA in soil and sediment: fate and ecological relevance, Biology and Fertility of Soils, 45 (2008) 219-

399 235.

400 [38] S.H. Sirois, D.H. Buckley, Factors governing extracellular DNA degradation dynamics in soil,

401 Environ Microbiol Rep, 11 (2019) 173-184.

402 [39] D.J. Levy-Booth, R.G. Campbell, R.H. Gulden, M.M. Hart, J.R. Powell, J.N. Klironomos, K.P. Pauls,

403 C.J. Swanton, J.T. Trevors, K.E. Dunfield, Cycling of extracellular DNA in the soil environment, Soil

404 Biology and Biochemistry, 39 (2007) 2977-2991.

405 [40] D. Roszak, R. Colwell, Survival strategies of bacteria in the natural environment, Microbiological

406 Reviews, 51 (1987) 365-379.

407 [41] J.T. Lennon, S.E. Jones, Microbial seed banks: the ecological and evolutionary implications of

408 dormancy, Nature reviews. Microbiology, 9 (2011) 119-130.

409 [42] S.E. Jones, J.T. Lennon, Dormancy contributes to the maintenance of microbial diversity,

410 Proceedings of the National Academy of Sciences of the United States of America, 107 (2010) 5881-

411 5886.

412 [43] K. Schwibbert, A. Marin-Sanguino, I. Bagyan, G. Heidrich, G. Lentzen, H. Seitz, M. Rampp, S.C.

413 Schuster, H.P. Klenk, F. Pfeiffer, D. Oesterhelt, H.J. Kunte, A blueprint of ectoine metabolism from the

414 genome of the industrial producer Halomonas elongata DSM 2581 T, Environmental microbiology, 13

415 (2011) 1973-1994.

416 [44] C. Vargas, M. Argandona, M. Reina-Bueno, J. Rodriguez-Moya, C. Fernandez-Aunion, J.J. Nieto,

417 Unravelling the adaptation responses to osmotic and temperature stress in Chromohalobacter salexigens,

418 a bacterium with broad salinity tolerance, Saline Systems, 4 (2008) 14.

419 [45] A.M. MacIntyre, J.X. Barth, M.C. Pellitteri Hahn, C.O. Scarlett, S. Genin, C. Allen, Trehalose

420 Synthesis Contributes to Osmotic Stress Tolerance and Virulence of the Bacterial Wilt Pathogen

421 Ralstonia solanacearum, Mol Plant Microbe Interact, (2020) MPMI08190218R.

422 [46] X. Zhou, D. Fornara, M. Ikenaga, I. Akagi, R. Zhang, Z. Jia, The Resilience of Microbial Community

423 under Drying and Rewetting Cycles of Three Forest Soils, Frontiers in microbiology, 7 (2016) 1101.

424 [47] S.J. Blazewicz, R.L. Barnard, R.A. Daly, M.K. Firestone, Evaluating rRNA as an indicator of

425 microbial activity in environmental communities: limitations and uses, The ISME journal, 7 (2013) 2061-

426 2068.

427 [48] E. Schwartz, Characterization of growing microorganisms in soil by stable isotope probing with

428 H218O, Applied and environmental microbiology, 73 (2007) 2541-2546.

429 [49] E. Schwartz, D.J. Van Horn, H.N. Buelow, J.G. Okie, M.N. Gooseff, J.E. Barrett, C.D. Takacs-

430 Vesbach, Characterization of growing bacterial populations in McMurdo Dry Valley soils through stable

431 isotope probing with (18) O-water, FEMS microbiology ecology, 89 (2014) 415-425.

[50] K. Papp, R.L. Mau, M. Hayer, B.J. Koch, B.A. Hungate, E. Schwartz, Quantitative stable isotope
probing with H2(18)O reveals that most bacterial taxa in soil synthesize new ribosomal RNA, ISME J,
12 (2018) 3043-3045.
[51] D. Berry, E. Mader, T. Lee, D. Woebken, Y. Wang, D. Zhu, M. Palatinszky, A. Schintimeister, M.
Schmid, B. Hanson, N. Shterzer, I. Mizrahi, I. Rauch, T. Decker, T. Bocklitz, J. Popp, C. Gibson, P.
Fowler, W. Huang, M. Wagner, Tracking heavy water (D2O) incorporation for identifying and sorting
active microbial cells, Proceedings of the National Academy of Sciences of the United States of America,
112 (2015) E194-E203.
[52] N. Musat, R. Foster, T. Vagner, B. Adam, M.M. Kuypers, Detecting metabolic activities in single
cells, with emphasis on nanoSIMS, FEMS microbiology reviews, 36 (2012) 486-511.
[53] S. Placella, M. Firestone, Transcriptional response of nitrifying communities to wetting of dry soil,
Applied and environmental microbiology, 79 (2013) 3294-3302.

Figure captions:

Figure 1. (a) Box-plot of bacterial abundance in soils in desiccated conditions (Dry) and after a rewetting incubation (Rewet). Centre lines show median value and boxes show upper and lower quartiles. Nonparametric Wilcoxon test was used to determine the significant difference of bacterial abundance in soils and $p < 0.05$ indicates significant difference between dry and rewetted soils (n=24 for both conditions). (b) The fold change of bacterial abundance following the rewetting incubation compared to the desiccated soils. The open black cycle indicates a fold change significantly greater than 1 ($p < 0.05$), the solid green circles represent no significant difference from 1 ($p > 0.05$), and the solid red circle indicates a change significantly lower than 1 ($P < 0.05$).

Figure 2. Composition of the bacteria in soils under desiccated conditions (Dry) and after the rewetting incubation (Rewet). The bacterial composition was plotted at phylum level and the most abundant phyla (with relative abundance >1% in average) are shown. Taxa not seen more than three times in at least 20 % of the samples were removed using phyloseq package in *R*. The relative abundances of minor phyla or classes were summed up and shown as "others" which do not exceed 7.5% of total bacterial reads in any given soil.

Figure 3. Heatmap showing significant fold change ($p < 0.05$) in abundance of active taxa in different soils following the 28-day rewetting incubation. A total of 21 taxa at the genus level are listed that displayed growth in the highest number of soils (> 3 soils). Each taxa designation contains the phylum (p), class (c) and genus (g) affiliation. Whenever a genus designation is not defined (unclassified genus), a defined family (f) or order (o) affiliation is specified instead. The values in the column "Average" indicates the average fold change in abundance of each taxa in soils following rewetting. Blank (white) square indicates no significant increase in abundance in a soil after rewetting ($p > 0.05$).

Table 1. The documented information of archive soils

Sample No.	Series No.	Collecting date	Collector(s)	Province	City/County	Land use	Vegetation	Soil genetic classification	Soil parent materials	Soil taxonomy
S01	5151	1934/10/9	Po Suo, Guangjiong Hou	Qinghai	Dulan	livestock farming	-	calcareous clay	-	uap-ustic isohumisols
S02	5213	1935/12/24	Lianqing Zhu et. al	Sichuan	Huayang	dry land	-	brown soil	granite	hapli-udic argosols
S03	5009	1939/11/15	Lianqing Zhu, Fanqi Zeng	Yunnan	Mengzi	farmland	rice	red soil	-	Fe-leachi-stagnic anthrosols
S04	5157	1934/11/13	Lianjie Li et. al	Guangxi	Yongning	wasteland	miscellaneous tree e.g. Chinese fir, pine	red soil	sandstone	hap-udic ferrisols
S05	HN	1935/1/1	Changyun Zhou	Hunan	Changsha	-	-	red soil	-	-
S06	5001	1934/11/13	Guangjiong Hou	Jiangxi	Nanchang	-	-	red soil	-	hap-udic ferrisols
S07	5289	1938/5/24	Daquan Song et. al	Fujian	Jianyang	vegetable garden	vegetables	fluvisols	river alluvium	acidic humid alluvial entisols
S08	5143	1938/10/1	Daquan Song et. al	Fujian	Longyan	wasteland	-	lime soil	limestone	black lithologic isohumisols
S09	5277	1938/5/24	Daquan Song et. al	Fujian	Chongan	tea orchard	tea	purple soil	calcareous purple shale	calcareous purpli-udic cambosols
S10	5061	1938/12/1	Daquan Song et. al	Fujian	Jianyang	wasteland	-	podzol	-	typic spodosol
S11	5062	1938/12/1	Daquan Song et. al	Fujian	Dehua	wasteland	grass	podzol	volcanic rock	typic spodosol
S12	5275	1938/5/3	Daquan Song et. al	Fujian	Chongan	wasteland	weeds	purple soil	purple sandshale	purpli-udic cambosols
S13	5030	1938/10/23	Daquan Song, Zhenyu Yu	Fujian	Longxi	farmland	rice	red soil	river alluvium	typic Fe-accumuli-stagnic anthrosols
S14	5031	1938/10/23	Daquan Song et. al	Fujian	Jianou	farmland	rice	red soil	river alluvium	typic Fe-accumuli-stagnic anthrosols
S15	5032	1938/12/23	Daquan Song et. al	Fujian	Longxi	farmland	rice	red soil	-	typic Fe-accumuli-stagnic anthrosols
S16	5033	1938/12/23	Daquan Song et. al	Fujian	Longxi	farmland	rice	red soil	-	Fe-accumuli-stagnic anthrosols
S17	5171	1938/8/5	Daquan Song et. al	Fujian	Yongchun	wasteland	shrubs and weeds	red soil	white effusive rock	hap-udic ferrisols
S18	5172	1938/5/5	Daquan Song et. al	Fujian	Yongchun	wasteland	evergreen broad-leaved trees	red soil	quartzite and sandstone	humic udic ferrisols
S19	5174	1938/5/5	Daquan Song et. al	Fujian	Yongchun	wasteland	evergreen broad-leaved	red soil	quartzite	hap-udic ferrisols
S20	5274	1938/10/3	Changyun Zhou et. al	Fujian	Yongchun	wasteland	grass	acidic residual soil	granite	acidic ochri-aquic cambosols
S21	5278	1938/5/24	Changyun Zhou et. al	Fujian	Yongan	wasteland	-	red soil	sandy lake sediment	semi-humic organic soil
S22	5119	1938/10/22	Chengfan Xi et. al	Fujian	Putian	wasteland	natural vegetation	coastal saline soil	sea mudalluvium	halogenic soil
S23	5120	1938/10/22	Chengfan Xi et. al	Fujian	Putian	wasteland	natural vegetation	coastal saline soil	-	halogenic soil
S24	5173	1938/5/5	Daquan Song et. al	Fujian	Zhangping	wasteland	weeds	yellow soil	granite	humic udic ferrisols

Figure 1

485 **Figure 2**

Relative abundance of bacterial phyla (%)

493   **Figure 3**