Environment-imposed constraints can make Brownian walkers efficient searchers

Harshana Rajakaruna[1], James O'Connor[2], Ian A. Cockburn[2] and Vitaly V. Ganusov[1,*]

[1]Department of Microbiology, University of Tennessee, Knoxville, TN 37996, USA
[2]Department of Immunology and Infectious Disease, John Curtin School of Medical Research, Australian National University, Canberra, 2601, Australia
[*]Corresponding author: vitaly.ganusov@gmail.com

May 28, 2021

Abstract

Pathogen-specific CD8 T cells face the problem of finding rare cells that present their cognate antigen either in the lymph node or infected tissue. To optimize the search for rare targets it has been proposed that T cells might perform a random walk with long displacements called Levy walks enabling superdiffusive behavior and shorter search times[1–4]. Many agents ranging from cells to large animals have been found to perform Levy walks[4–6] suggesting that Levy walk-based search strategies may be evolutionary selected[7–9]. However, whether random walk patterns are driven by agent-intrinsic programs or shaped by environmental factors remains largely unknown[6,10]. We examined the behavior of activated CD8 T cells in the liver where both the movement of the cells and the underlying structural constraints can be clearly defined. We show that *Plasmodium*-specific liver-localized CD8 T cells perform short displacement, Brownian-like walks and yet display transiently superdiffusive displacement, the cardinal feature of efficient Levy walks. Because liver-localized CD8 T cells are mainly associated with liver sinusoids, simulations of Brownian or Levy walkers in structures derived from the liver sinusoids illustrate that structure together with preferred forward movement results in similar superdiffusive displacement. Interestingly, Brownian walkers require less time to find a rare target when T cells search for the infection in physiologically-derived liver structures. Importantly, analysis of fibroblastic reticular cell networks on which CD8 T cells move in lymph nodes also allows for superdiffusion in simulations, though this is not observed experimentally, suggesting that structure is not the only factor determining movement patterns of T cells. Our results strongly suggest that observed patterns of movement of CD8 T cells are likely to result from a combination of a cell-intrinsic movement program, physical constraints imposed by the environmental structures, and other environmental cues. Future work needs to focus on quantifying the relative contributions of these factors to the overall observed movement patterns of biological agents.

Abbreviations: MSD - mean squared displacement, LW - Levy walk, BW - Brownian walk, RAS - radiation attenuated sporozoites, FRC - fibroblastic reticular cells, DC - dendritic cells.

Introduction

Search for resources is a fundamental property of many biological agents starting from individual cells to large animals. While different agents involve various search strategies, two types of search are of particular importance due to their fundamental nature: Brownian walks and Levy walks/flights[11]. Brownian walkers display short displacements, typically consistent with a thin-tailed distribution that has a finite mean and variance, with random turning angles; in Brownian walks mean squared displacement (MSD) increases linearly with time[11–14]. In contrast, Levy walkers display both short and long displacements consistent with a fat-tailed distribution that has either finite mean and infinite variance or both infinite mean and variance; in Levy walks/flights MSD increases faster than linear with time[11,13]. Understanding what types of walk various biological agents employ has been a subject of many investigations[1–10].

For adequate protection the adaptive immune system of vertebrates needs to efficiently survey tissues for infection. However, how such efficiency is achieved remains poorly understood. In lymph nodes (LNs) naive T cells seem to search randomly for antigen-presenting cells, acting as Brownian walkers undergoing short displacements, although some role of attracting chemokines has been proposed[13,15–18]. Even less information exists on infection search strategies in peripheral, nonlymphoid tissues. In the skin, activated or memory CD8 T cells tend to perform Brownian walks[19], and in explanted murine lungs T cells display intermittent migration resulting in normal diffusion and a large variability in average velocities between individual cells[20]. Surprisingly, *Toxoplasma gondii*-specific activated CD8 T cells in murine brains undergo Levy walks[4]. Levy walks may allow T cells in the brain to locate rare parasite-infected cells efficiently[4].

Another apicomplexan parasite *Plasmodium*, the causative agents of malaria infects the liver following inoculation of parasites by probing mosquitoes[21–23]. At this stage of the *Plasmodium* lifecycle the intracellular liver stage parasites are susceptible to clearance by vaccine-induced CD8 T cells. Remarkably, liver-localized CD8 T cells alone are capable of eliminating all liver stages and preventing clinical malaria[24,25]. Given that a murine liver has about 10^8 hepatocytes how exactly CD8 T cells locate all parasites within the 48 hour span of the parasite liver stages is unknown[26]. Furthermore, while we and others have documented movement of liver-localized CD8 T cells, the walk types employed by these cells and their potential underlying mechanisms remain unknown[27–29].

Typical intravital imaging experiments allow to collect 3D coordinates of T cells in tissues over time, and with such data one can construct several movement characteristics to determine the type of walk cells exhibit[4,5,18,30–32]. Specifically, the rate of MSD increase with time, distributions of movement lengths and turning angles often allow to determine the walk type[5,31,33]. Scaling the distribution of movement displacements generated for different imaging frequencies also allows to determine the type of cell walk because Levy walks-based movement distributions are thought to be relatively invariant to such rescaling[4,18,32]. Determining the walk type of T cells is important because Levy walks have been shown to be more efficient than Brownian walks in some cases suggesting that Levy walk-based strategies may be evolutionary selected[1–4,34,35]. However, defining search efficiency may be specific system-dependent, and many studies identified conditions when Levy walks are not optimal[6–10,32,36–40].

While several studies provided strong evidence that T cells undergo a specific movement strategy mechanisms explaining such movement strategy in general remain unknown. For example, activated CD8 T cells in murine brains undergo Levy walks[4], however, whether such a walk type is driven by cell-intrinsic program (as suggested by Harris *et al.*[4]) or arises due to other factors has not been determined. Indeed, previous ecological studies suggested that turbulence of fluids or air may

explain patterns of movement of individual cells or birds, respectively, but the actual features of turbulences that allow agents to undergo Levy walks were not directly measured[41,42]. Furthermore, in a theoretical study Volpe & Volpe[38] also showed that environmental constrains impact the movement patterns of agents and efficiency of the search suggesting that in addition to cell-intrinsic programs other factors are likely to impact movement patterns of T cells[17].

In this study with the use of intravital microscopy of live mice we provide the first accurate quantification of the movement pattern of liver-localized CD8 T cells (and openly share these data with the community). We show that in most circumstances CD8 T cells undergo short, Brownian-like movements characterized by thin-tailed distributions that yet result in highly superdiffusive displacements; these superdiffusive displacements occur for longer times than, for example, movements observed for CD8 T cells in murine brains (20 vs. 8 min, respectively). By accurately quantifying the structure of liver sinusoids we show that superdiffusive behavior of liver-localized T cells naturally emerges for Brownian walkers when simulations of movements are done in the liver. Interestingly, our analysis of the structure of fibroblastic reticular cells of LNs suggested that LN-localized T cells should be as superdiffusive as liver-localized T cells; however, LN-localized T cells are superdiffusive for only few (~ 5) minutes suggesting that other factors such as chemical cues may also play a role in determining movement patterns of T cells in tissues. We finally demonstrate that the often cited advantage of Levy walkers in search for rare targets in free, unconstrained 3D space disappears when walks are done in the liver sinusoids. Taken together, our results highlight that movement patterns of T cells observed in various tissues is likely to be impacted by the underlying physical constraints and that future studies should focus on discriminating between the relative contribution of cell-intrinsic programs, environmental constraints, and chemotactic cues in determining the overall patterns of T cell movements.

Results

Majority of liver-localized CD8 T cells crawl in liver sinusoids. To understand what strategies *Plasmodium*-specific CD8 T cells may be using to survey the liver for infection we have established a protocol to generate stable numbers of liver-localized CD8 T cells[28]. In these experiments, naive T cells are activated *in vitro*, and then transferred i.v. into naive mice that either left uninfected or can be later infected with *Plasmodium* sporozoites (Fig. 1A)[28]. By injecting fluorescent label (rhodamine-dextran) i.v. immediately prior to surgery and imaging we can efficiently label the liver sinusoids (Fig. 1B). Intravital imaging showed that CD8 T cells in the liver are highly motile cells with the average speed of 6.6 μm/min, a high meandering index of 0.52, and a low arrest coefficient of 0.25 (Fig. 1C and Movie S1). Importantly, these CD8 T cells were observed to be located in liver sinusoids most of the time (89-100%, Fig. 1D). Due to incomplete labeling of the sinusoids and reduced red signal from deeper liver areas it is likely that a lower bound is an underestimate and in this system nearly all T cell movements (and by inference, most T cells) were associated with the sinusoids. As compared with the blood flow rate in the liver, a relatively low average speed of these liver-localized T cells and T cells in our previous experiments with high frequency intravital imaging suggests that the cells are crawling on the endothelial wall of the sinusoids and not flowing with the blood[28]. Thus, our robust experimental set-up allows for the first time to accurately quantify movement patterns of tissue-associated CD8 T cells along with the well defined structures (i.e., hepatic sinusoids) constraining and defining walking patterns of T cells.

Liver-localized CD8 T cells undergo correlated random walks. In our experiments move-

Figure 1: The majority of *in vitro* activated CD8 T cells transferred to naive recipients move in liver sinusoids. We performed intravital imaging of the livers of mice that had previously received *in vitro* activated CD8 T cells and i.v. injection of rhodamine dextran (see Materials and Methods for more detail). (A) 4 hours after i.v. cell transfer of 5×10^6 OT-I GFP CD8 T-lymphocytes (green), mice were injected with 50 µl (20 µg/ml) of rhodamine dextran 70kDa (red) and immediately prepared for intravital imaging. The mice were imaged using a two-photon microscopy with a standard galvanometer scanner in order to acquire a 50 µm deep z-stack every 13 or 26 sec. (B) A sequence of images taken over time to demonstrate the movement of GFP cells (green) through the sinusoids (red) in the direction indicated by the white lines (Movie S1) with star and arrow indicating displacement of two different cells. We show the overall scanned area (Bi) and time lapse for 2 cells (indicated by an arrow and a star) moving in the sinusoids (Bii-Bv). Scale bar is 50 µm (Bi) or 20 µm (Bii-Bv). Time in min:sec:milliseconds is shown. (C) Tracks of randomly chosen 30 cells are shown. (D) Movement parameters of OT-I GFP cells in the liver sinusoids such as average track speed, meandering index, and arrest coefficient ($n = 269$). For every cell movement we calculated the percent of time each cell spent in the sinusoids by calculating signal overlap between cell fluorescence (GFP) and sinusoid fluorescence (rhodamine dextran, $n = 231$). The imaging data in B is representative of 3 experiments, and analysis shown in D is for datasets pooled together (means and SD are shown by bars).

ments of liver-localized *Plasmodium*-specific CD8 T cells were recorded with two photon microscopy. Interestingly, T cells in the liver displayed transiently superdiffusive overall displacement characterized by MSD increasing faster than linearly with time (with slope $\gamma = 1.64$) for up to 20 min (Fig. 2A). Saturation in MSD with time in these and other experiments typically arises in part due to rapidly moving T cells leaving the imaging area (Fig. S1). Superdiffusion is a hallmark feature of Levy walks[13,43], and liver-localized CD8 T cells displayed superdiffusive displacement for much longer times than brain-localized CD8 T cells undergoing Levy walks (see Figure 3A in Harris *et al.*[4]). Rare long movements are the cardinal feature of Levy walks[3,4,11] and yet liver-localized T cells only displayed short, Brownian-like movements (Fig. 2B). Tail analysis[33] of the distribution of actual or scaled movement lengths (movement lengths normalized by mean length) suggested the shape parameter of the powerlaw distribution $\mu = 5.6$ — an indication of Brownian ($\mu > 3$) and not Levy ($\mu \leq 3$) walk. Interestingly, scaling the movement length distribution for sub-sampled data (i.e., for data sampled at a lower frequency than was used at imaging) resulted in relatively similar distributions (Fig. S2) suggesting that even thin-tailed movement length distributions may be similar for sub-sampled, scaled data. Thus, large overall cell displacements with small movement lengths arose not due to Levy-like cell-intrinsic behavior but due to small turning angles and a strong correlation between initial and subsequent velocity vectors for at least 10-15 min, resulting in cells moving in one direction for substantial periods of time (Fig. S3).

To complement the movement length distribution tail analysis we fitted a series of alternative mathematical models characterizing the distribution of movement lengths of T cells. We found that generalized Pareto distribution fitted the data with best quality (Fig. 2C-D and Tab. S2). Best fit parameters of the generalized Pareto distribution predicted a finite mean and variance for the movement length distribution – which again are the chief features of the Brownian and not Levy walkers[11]. Thus, in our experiments liver-localized CD8 T cells performed correlated (persistent) random walk with superdiffusive behavior that was longer than that observed for CD8 T cells in the brain.

To verify that this result was not due to a potential impact of the injected rhodamine-dextran dye we analyzed four other datasets from our previous study on movement of liver-localized CD8 T cells[28]. These experiments involved transfer of *in vitro* generated wild-type and LFA-1-deficient activated CD8 T cells or *in vivo* generation of liver-localized T cells by transfer of naive OT-I cells to mice followed by immunization of these mice with radiation-attenuated malaria sporozoites (RAS) expressing the SIINFEKL epitope recognized by these cells[26,28]. Both *in vitro* wild-type activated cells and *in vivo* activated cells imaged at a memory time point (4 weeks) after immunization also displayed transiently superdiffusive behavior with $\gamma > 1$ for 10-20 min (Fig. 2E). Interestingly, LFA1-deficient T cells which cannot strongly attach to endothelial cells and roll under the blood pressure[28], displayed displacement closest to Brownian and yet were transiently superdiffusive ($\gamma = 1.18$ in Fig. 2E). We also observed high heterogeneity in MSD in mice imaged one week after RAS immunization (Fig. 2E). At this time point after *in vivo* priming, many CD8 T cells do not undertake the crawling behavior characteristic of liver-localized T cells, most likely due to intermediate expression of LFA-1 shortly after immunization[28].

Likewise all but one cell population had relatively small movement lengths. The exception was for CD8 T cells in mice one week post RAS immunization (Fig. 2F) where cells do not attach firmly to the liver endothelium and often flow in the bloodstream, thus, performing Levy flights[28]. Transiently superdiffusive behavior with small, Brownian-like movements was also observed for *Plasmodium*-specific activated CD8 T cells and activated CD8 T cells with irrelevant specificity in *Plasmodium berghei*-infected mice (Fig. S5). Taken together, our results establish that most crawling liver-localized CD8

Figure 2: Activated CD8 T cells in the liver are Brownian walkers displaying super-diffusive displacement. For experiments described in Fig. 1 we calculated basic movement characteristics such as the mean squared displacement (MSD, A) and distribution of scaled movement lengths (B) for all T cells ($n = 268$). Resulting slopes for linear regressions for the log(MSD) with log(t) and movement length distribution (γ and μ, respectively) are shown on individual panels. (C)-(D). We fit several alternative models of movement length distribution to the raw (C) or scaled (D) displacement data from experiments in Fig. 1 (eqns. (S.4)–(S.9)). Parameters for the fitted models along with AIC values are given in Tabs. S1 and S2. (E)-(F). We performed similar analyses with other published data for *in vitro* or *in vivo* activated CD8 T cells in the liver[28]. Cell displacement in the liver was characterized by MSD with slope γ (E) and movement lengths characterized by the tail slope μ (F). Additional details on analysis of movement length distribution are given in Fig. S4. Scaling in (B) and (F) was done for all tracks per dataset using mean displacement length $\bar{r}$ using relationship $\rho = r/\bar{r}$ (see Materials and Methods for detail).

169 T cells display superdiffusive behavior with small movements; these are the hallmarks of correlated
170 random walks[44].

Figure 3: Liver sinusoids have relatively short lengths and small branching angles. We performed experiments in which liver sinusoids were labeled by i.v. injection of Evans blue or rhodamine dextran and then imaged immediately by using intravital microscopy. (A) Z-stacks of 2D images of the liver used for analysis. (B) Computer-digitized 3D sinusoidal maps were generated by contrast thresholding and by stacking up the 2D images. Red shows the center paths of the sinusoids. (C)-(D) Computer-assisted rendering of liver sinusoids for a subset of the data (C, shown in red) used to determine the nodes (D, yellow circles) and the edges/branches (red) of the structure (turquoise circles are the end points). (E)-(H) Characterization of the liver sinusoids. (E) We fit several alternative models to the distribution of lengths of sinusoids using likelihood approach (see Materials and Methods and Tab. S1). (F) Cumulative distribution of the sinusoid lengths was also analyzed using tail analysis with the indicated linear regression slope μ [33]. (G) Distribution of branching angles between sinusoids calculated for nodes ($n = 4013$) in D. Parameter $\overline{\theta_a}$ is the average branching angle for acute angles. (H) Straightness of the liver sinusoids (computed as the ratio of point-to-point branch length to actual branch length for all near neighbors in D); $\bar{S}$ is the average straightness.

Liver sinusoids have short lengths and small branching angles. Given the network struc-
ture of liver sinusoids (Fig. 1B) and universal pattern of transient but relatively long superdiffusive
behavior of liver-localized CD8 T cells we hypothesized that a linear structure of the sinusoids may be
forcing T cells to exhibit persistent walks. Therefore, we analyzed 3D structure of liver sinusoids. For
these intravital imaging experiments we injected a contrast dye (Evans blue or rhodamine-dextran)
and imaged murine livers (Fig. 3A). Z-stacks of the images were processed in Matlab to generate
highlighted sinusoidal maps (Fig. 3B). A subset of the full image was further processed to generate
nodes and edges of the sample (Fig. 3C-D). Analysis of the processed digital structures suggested
that the branch length distribution was best described by the half-normal distribution with gener-
alized Pareto distribution being second best (Fig. 3E and Tabs. S1 and S2). The longest branch
detected did not exceed 100 μm, and the tail of the branch length distribution was characterized by
a relatively high shape parameter of the powerlaw distribution ($\mu = 7.65$, Fig. 3F)[33]. Importantly,
sinusoids tend to branch at a small angle which is likely to improve blood flow (Fig. 3G)[45]. Finally,
analysis of the neighboring nodes in the network (Fig. 3D) suggests that sinusoids connecting two
adjacent nodes tend to be straight (Fig. 3H). Taken together, it seems remarkable how properties of
the liver sinusoids such as small branching angles, straightness, and shortness match the observed

pattern of movement of liver-localized T cells (e.g., Fig. 2F and Fig. 3F).

Liver sinusoids alone allow for superdiffusion of Brownian walkers. To investigate how the structure of the liver sinusoids may impact the movement patterns of cells with an intrinsic movement program we simulated movement of cells assuming that they are either Brownian or Levy walkers (Fig. 4). We assumed that distribution of movement lengths follows a powerlaw distribution with the shape parameter μ being different between Brownian ($\mu = 4.5$) and Levy ($\mu = 2.5$) walkers while fixing the average movement length for two walk types[11]. As expected in free 3D space Brownian walkers displayed normal diffusion ($\gamma = 1$) while Levy walkers superdiffused ($\gamma = 1.5$, Fig. 4A-C). In contrast, when cells were run on the digitized network of liver sinusoids (Fig. 3A-D) with cells allowed to make a random choice of turning when reaching a branching point the movement pattern changed dramatically (compared Fig. 4A vs. D and B vs. E). Impressively, both Brownian and Levy walkers displayed superdiffusion in the liver structure for substantial period of time (~ 55 time steps, Fig. 4F) with the diffusion becoming normal at longer time scales. This pattern is similar to small particles displaying transient superdiffusion early and normal diffusion later due to motion of self-propelling bacteria in media[46]. Thus, the structure of the liver sinusoids seems to be sufficient to allow transient superdiffusion of Brownian walkers such as crawling liver-localized CD8 T cells.

It is surprising that structure of liver sinusoids was sufficient to allow Brownian walkers to superdiffuse without the need for chemical cues in the environment (e.g., chemokines). Because previous studies have suggested that T cells in LNs are close to Brownian walkers[13,18,37,47] we sought to determine if this is because of the structural constraints of the LN environment. It has been suggested that T cells use the network of fibroblastic reticular cells (FRCs) or dendritic cells (DCs) as a guide for movement[48,49]. Recently, the FRC network of the LNs has been imaged using confocal microscopy[50]. Using a similar computational pipeline we digitized the FRC network using previously published data[50] and calculated its basic characteristics (Fig. S7A-C). Interestingly, similarly to the liver sinusoids the FRC network had relatively short branch lengths with the tail of the distribution being well described by the powerlaw distribution with a shape parameter $\mu = 6.7$ (Fig. S7D). A half-normal distribution fitted the data on branch length distribution with best quality (Fig. S7E and Tabs. S1 and S2). Similarly to the liver sinusoids FRC network has relatively small branching angles at the nodes, and connections between adjacent nodes are mostly straight (Fig. S7F-G). By assuming that T cells can only walk in the middle of the digitized FRC networks with preferred movement forward simulations of T cell walks suggested that independently whether cells do Brownian ($\mu = 4.5$) or Levy ($\mu = 2.5$) walks they display transient superdiffusion similar to that of T cells simulated to walk in the liver (Fig. S7H). However, analysis of the experimental data on movement of naive CD8 T cells in murine LNs[50] suggested that such T cells are superdiffusive for very short time period (Fig. S8). The duration of a persistent walk of CD8 T cells in LNs is around 4-5 min while CD8 T cells in the liver superdiffuse for nearly 20 min (Fig. S9). Analysis of the actual moving tracks of T cells in LNs suggests that many T cells tend to turn after a short directed movement (Fig. S8A&E). Indeed, it is well known that naive T cells in LNs tend to be restricted to the T cell areas, most likely by the gradient of chemokines. In contrast, in the liver in the absence of specific infection liver-localized T cells are likely to be unconstrained in movement by chemokines but are constrained by the physical structure of the sinusoids. Our results, thus, suggest that while structural constraints of the liver sinusoids or FRC network in LNs may allow for long-term superdiffusion of T cells other factors such as attraction cues may impair cell's ability to superdiffuse.

Brownian and Levy walkers have similar search efficiency in the liver. One of the advantages of Levy walks as often cited is a high efficiency at locating rare targets[1,3,4,32]. However, evaluation of search efficiency is often done in free space ignoring physical constraints or pathways

Figure 4: Liver structure imposes strong constraints on the movement pattern and overall displacement for Brownian or Levy walkers. We simulated movement of 200 T cells assuming that their movement length distribution is described by Pareto (powerlaw) distribution (see eqn. (S.9)) with $\mu = 4.5$ (i.e., Brownian walkers, A&D) or $\mu = 2.5$ (i.e., Levy walkers, B&E). T cells moved in free environment with no constraints (A-C) or are constrained by liver sinusoids (D-F). A sample of 20 cell trajectories is shown in A-B & D-E. (C)&(F). We calculated the MSD for the walkers and estimated the displacement slope γ by linear regression of the initially linear part of the ln-transformed data. In free space simulations turning angle distribution for moving T cells was generated using uniform spherical distribution, while in liver sinusoids movement was set forward along the sinusoidal mean path. The average movement length per time step was $\bar{r} = 2.1$ μm as in our data with time step of 26 sec. An example of a simulated T cell movement in the liver is shown in Fig. S6 (see Materials and Methods for more details).

agents may encounter when searching for targets. It has been shown theoretically that presence of obstacles may influence optimality of Levy walks[38]; however, whether similar results hold for physiologically constrained environments was not investigated. Furthermore, previous studies did not attempt to normalize distribution of movement lengths, so that the average speeds are similar for Brownian and Levy walkers. In case of ballistic movements ($\mu \leq 2$ of the Pareto distribution) this is also impossible because mean of the displacement distribution does not exist, and therefore, comparing BW and LW with $\mu \leq 2$ is biologically unreasonable.

We simulated the search for a single *Plasmodium* parasite by a liver-localized CD8 T cell assuming that the parasite induces no attraction to the infection site and by restricting the average movement speed to be the same for Brownian or Levy walkers. We used Pareto (powerlaw) distribution to describe the movement lengths with scale parameter $\mu = 4.5$ for BW and $\mu = 2.5$ for LW (see Materials and Methods for more detail). When a CD8 T cell searches for the parasite in constraint-free 3D space, on average Levy walkers require less time ($\bar{T}$) to locate the target (Fig. 5A&C) which is consistent with previous results[1,4]. Levy walks also result in smaller variability (characterized by the standard deviation of the time to target σ) in the time to find the target. Interestingly, when the search is performed in the digitized liver structures, Brownian walkers are on average more efficient at finding targets than Levy walkers (Fig. 5B&D) with smaller variance in the time to target. Furthermore, in 55% of simulations Levy walkers would take longer than the slowest Brownian walker to find the target (Fig. 5D). This may be particularly important in the case of malaria where a failure to find all the targets in the liver within a fixed time period (e.g., 48 hours) will result in blood stage infection and fulminant disease. Alternatively, in the liver Brownian walkers may be unable to locate the infection within a given time period if they start their search too far away from the parasite, while Levy walkers have a non-zero chance (Fig. 5D). It is well established experimentally that efficient vaccination protocols need to induce a relatively large number of Plasmodium-specific CD8 T cells to provide sterilizing protection against infection with Plasmodium sporozoites[24,26]. Whether such a requirement of a large number of T cells to control the infection comes from T cells undergoing a correlated random walk with long-term superdiffusion remains to be determined.

The difference in search efficiency between Levy and Brownian walkers in free 3D space and in the liver is due to differences in the paths to the target. In the digitized liver there are essentially few short pathways between a T cell and the target, and Brownian walkers make consistently small movements moving closer to the target. In contrast, due to small average movement length Levy walkers spend long periods of time pausing before performing a long run sometimes resulting in them reaching the target rapidly. However, long pauses also result in much longer overall time to the target, which increases the average search time substantially. Thus, our results suggest that as in the tale of "The Tortoise and the Hare" in the liver a slow and steady Brownian walker reaches the target on average faster than an irregular but rapid Levy walker.

In our simulations so far we assumed that a single T cell searches for the single infection site. However, in reality multiple T cells may be searching for the same parasite. Indeed, our recent analysis suggests that for 10^5 liver-localized CD8 T cells at least 1 T cell should be within 100 μm of the parasite with 99% certainty[51]. In our experiments we rarely see parasites surrounded by more than 5-8 T cells suggesting that in many situations fewer than 10 T cells are close enough to the parasite to find it within several hours[52,53]. As expected the average time to find the target declines with increasing number of searchers, independently whether the search is done in free 3D space or constrained liver environment (Fig. S10) but the overall reduction in search time is larger when search is done in 3D. Intuitively, the reduction in time to infection arises due to variability in search time since we assume that search by individual T cells is done independently; therefore, with many T cells

Figure 5: Levy walkers are less efficient in search for a single target in constrained environment of the liver. We simulated search of a T cell for a single target assuming Brownian (BW, $\mu = 4.5$) or Levy (LW, $\mu = 2.5$) walks in free space (A&C) or in liver sinusoids (B&D) and calculated the time it took for the T cell to locate the target (C&D). We assume that search occurs randomly (no attraction of the T cell to the target). (A)&(C) Levy walkers are more efficient in free space. We show distribution of times to the target for simulations when T cell started the search 100 μm away from the target. (B)&(D) Brownian walkers are more efficient in the liver. We show distribution of times to the target for simulations when T cell started the search 500 μm away from the target (mean $\bar{T}$ and standard deviation $\pm\sigma$ of the time is shown in C&D). In all simulations T cell was assumed to find the target when it was within 40 μm from the parasite (an approximate size of the murine hepatocyte). We simulated 1000 cells (3D) or 3000 cells (liver) under each scenario with average movement length per time step being $\bar{r} = 2.1$ μm in Pareto (powerlaw) distribution with the shape parameter μ (eqn. (S.9)).

searching for infection, the one T cell having shortest search time will define success of the search and LW allows for a larger fraction of faster searches. Yet, for physiologically relevant numbers of T cells searching for the infection site in the liver, Levy walks do not provide any substantially increased efficiency, and large enough numbers of Brownian walkers may be fully sufficient to locate all the targets within a fixed period of time.

Discussion

T cells of the adaptive immune system are very efficient at locating and eliminating pathogens. However, the search strategies used by such efficient T cells remains poorly defined. It has been suggested that generalized Levy walks may be an evolutionary-selected cell-intrinsic strategy that allows brain-localized activated CD8 T cells to efficiently locate and eliminate *Toxoplasma gondii*-infected cells[4]. By analyzing novel data on movement of liver-localized CD8 T cells and linking these movement patterns to the digitized structure of liver sinusoids we showed that superdiffusion of T cells, the hallmark of Levy walks, may occur for longer durations as compared to superdiffusion

observed for T cells in the brain; yet, most liver-localized T cells make only short movements. Such transient superdiffusion naturally arises due to the physical constraints imposed by narrow branching liver sinusoids instructing forward movement-biased T cells to move in a persistent direction with small turning angles. Interestingly, activated CD8 T cells may perform Levy flights in the liver in situations when they poorly attach to endothelial wall of the sinusoids and float with the blood flow, thus, creating an impression of long range movements[11,28]. We have observed this phenomenon for liver-localized CD8 T cells one week after immunization with RAS; under these conditions CD8 T cells exhibit Levy walks with a high frequency of long movements (Fig. 2F). Overall, however, our results strongly suggest that observed pattern of superdiffusion or rare long range movements need not arise from cell-intrinsic movement programs but can be explained by the physical properties of the environment in which T cells move and preference of T cells to move forward. Importantly, we found that the widely cited advantages of Levy walks in searching for rare targets disappear if the search is done by a few T cell in biologically defined liver structures which effectively reduces target search problem from 3D to 1D further challenging the universality of the Levy walk-based search strategies[36,38,39].

Physical structures are unlikely to be the only constraint impacting the properties of T cell movements in tissues. We showed that FRC network of the LNs also allows for superdiffusion of agents in simulations; however, actual T cells in LNs superdiffuse for very short time periods (Figs. S8 and S9). Thus, other factors such as chemical cues (chemokines), finer details of the physical paths on which T cells walk such as FRCs/DCs, infection-induced inflammation, and preference of moving forward in the absence of external disturbances are likely to be also important in determining overall patterns of T movement in tissues. In particular, the role of environmental cues in determining agent movement patterns has been suggested in several systems[13,20,54].

Our results have broad implications for other systems in which superdiffusion and rare long range displacements have been observed or suspected[3,55]. It will be important to re-examine the evidence supporting intrinsic search programs by matching movement patterns of agents to the experimentally measured environmental details (e.g., airflows for albatrosses or currents for marine predators) and evaluating the role of cues (chemical or otherwise) in defining overall movement patterns of agents[6,10,38]. Ultimately, future studies will need to quantify the relative contribution of intrinsic programs, environmental physical constraints, and cues in determining movement pattern of agents[17].

Given our results it is tempting to speculate about potential mechanisms resulting in Levy walks of brain-localized CD8 T cells observed previously[4]. One possibility is that Levy-like displacements are due to special structures in the brain used by T cells for crawling. Such structures will then have to be poised at the threshold of percolation[56,57]. Another possibility is that some brain-localized CD8 T cells are in fact located in brain vasculature and thus undergo Levy flights with the blood flow — similarly as we have observed for liver-localized CD8 T cells under some circumstances. Finally, Levy walks may arise in the analysis due to data generation or processing artifacts. We have found that when many T cells are present in the tissue it is sometimes difficult to segment experimental data to identify individual cells and their movements. We have also found in our and other publicly available datasets that cell coordinate data often have missing time frames, and if such tracks are not cleaned, data analysis may generate an impression of a long displacement between two assumed-to-be-sequential time frames. Whether such artifacts contributed to the overall detection of Levy walks of T cells in the brain remains to be determined, and in general it can be beneficial when raw data from imaging data analysis are made publicly available following publication (i.e., the data of Harris *et al.*[4] are not publicly available).

Our study has several limitations. Only a small area of the liver (approximately $500 \times 500 \times 50~\mu$m) is generally imaged with intravital microscopy which restricts the number of cells recorded and creates a bias for retention of slowly moving cells in longer movies. While we show that the digitized structure of liver sinusoids can override the intrinsic movement program of T cells, we did not estimate how much of the experimentally observed movement pattern of liver-localized T cells is due to a cell-intrinsic program such as a rare ability of cells to turn back, physical constraints of the liver sinusoids, or other environmental cues. Methods for discriminating between these alternatives will have to be developed. Finally, we have focused on explaining our data with simplest possible movement models involving movement length distribution, MSD, and turning angles. Other, more complicated Levy walk models have been suggested such as generalized Levy walks (GLW) or Zigzag-GLW which allow for runs and pauses and assume or generate heterogeneity in cell velocities[4,32,43]. However, such models in general have more parameters, and fitting GLW-like models to our data is likely to result in overfitting. Our results, nevertheless, suggest that future studies focusing on benefits or evolutionary optimality of Levy walks would gain from more thorough investigation of the structural constraints and environmental cues that may be responsible for observed Levy-like movement pattern.

Our analyses raise many additional questions for future research. In particular, the impact of the *Plasmodium* infection on movement patterns of liver-localized CD8 T cells remains to be explored further. Our analysis of one unpublished dataset suggests similar long-term superdiffusion and short displacements for CD8 T cells following the infection (Fig. S5); however, this remains to be more fully investigated in the future. In particular, our recent work suggests that while initially, *Plasmodium*-specific CD8 T cells search for the infection site randomly, following formation of a T cell cluster near the parasite, other T cells become moderately attracted to the cluster[51,53]. Whether movement patterns of liver-localized T cells change following formation of T cell cluster near the parasite remains to be investigated. Furthermore, infection may also influence blood flow allowing T cells in circulation to be recruited to the infection site. Therefore, it may be advantageous for T cells that are crawling on endothelial cells of the liver sinusoids to de-attach from the endothelium and rapidly float with the blood, thus, undergoing Levy flights. Whether a combination of local search by crawling and then large displacements with the blood flow allows for a more efficient localization of all parasites in the liver also remains to be determined.

Ethics statement

Author contributions

VVG had the original idea for the study. VVG, HR, IAC, and JOC worked on various details of the experiments and analyses to be performed. JOC performed experiments (under supervision of IAC) and provided digitized data on movement of CD8 T cells in labeled liver sinusoids and separately

provided images of labeled sinusoids. HR performed data curations, developed methods to digitize liver sinusoids and FRC networks and to analyze track data for T cells in the liver and LNs, ran most of data analyses and simulations, and wrote summaries of the methods and results. VVG wrote the first draft of the paper. All authors participated in the discussion of interim analyses and results. All authors contributed to the writing of the final version of the paper via changes, comments, and suggestions.

Data sources

T cell movement/track data generated in this paper or in our previous publication are provided as an Excel spreadsheet with different tabs for different datasets. Imaging data of liver sinusoids (TIFF images) are also provided. The data from a previous publication on movement pattern of naive CD8 T cells in murine lymph nodes and imaging data of FRC networks in the LNs[50] are also provided. These files are stored together with the Matlab codes used to process and analyze the data on Github.

Code sources

All analyses have been primarily performed in Matlab (ver 2019b) and codes used to generate most of figures in the paper are provided on GitHub: `https://github.com/vganusov/Levy_walks_liver.git`. All code questions should be addressed to H.R. (hanrajakaruna@gmail.com).

Acknowledgments

We would like to thank Viktor Zenkov to contributing to the start of this work by manually digitizing liver sinusoidal maps and Marco Novkovic for providing imaging data of lymph nodes and data on movements of naive CD8 T cells in murine lymph nodes. This work was supported by the NIH grant (R01 GM118553) to VVG.

Methods summary

To generate liver-localized CD8 T cells we followed our previously established protocol[28]. In short, naive TCR-transgenic OT-I cells, specific to an epitope from *Plasmodium berghei* (Pb-CS^{5M}) were stimulated *in vitro* with specific peptide (SIINFEKL) in the presence of IL-2. Four days after the culture, 5×10^6 cells were transferred into individual mice via tail vein, and prepared for liver imaging 2-4h after the injection (Fig. 1A). It takes few hours hours for activated CD8 T cells to attach and retain in the liver[28]. To label liver sinusoids we injected rhodamine-dextran i.v. 5-15 min prior to imaging (to allow for mouse preparation for liver imaging). Z-stacked images were processed using Imaris (Bitplane) to provide 3D coordinates of digitized cells over time. Imaging area was approximately $500 \times 500 \times 50$ μm. Cell positions were curated to make sure that measurements were done in consistent time intervals of 13 or 26 sec. For liver structure analysis another set of experiments involved injection of rhodamine-dextran or Evans blue 5-15 min prior to imaging (2 mice). Imaging was done using two photon microscope. Analyses of the imaging data or of cell

coordinate data were done in Matlab or Mathematica (see extended description of the methods in Extended Methods online). T cell coordinate data and liver sinusoids imaging data are available on Github (see above).

References

1. Viswanathan, G. M., Buldyrev, S. V., Havlin, S., da Luz, M. G., Raposo, E. P. & Stanley, H. E. 1999 Optimizing the success of random searches. *Nature*, **401**(6756), 911–914. doi:10.1038/44831.

2. Bartumeus, F., Catalan, J., Fulco, U. L., Lyra, M. L. & Viswanathan, G. M. 2002 Optimizing the encounter rate in biological interactions: Lévy versus Brownian strategies. *Phys Rev Lett*, **88**(9), 097 901.

3. Viswanathan, G., Raposo, E. & Da Luz, M. 2008 Lévy flights and superdiffusion in the context of biological encounters and random searches. *Physics of Life Reviews*, **5**(3), 133–150.

4. Harris, T. H., Banigan, E. J., Christian, D. A., Konradt, C., Wojno, E. D. T., Norose, K., Wilson, E. H., John, B., Weninger, W. *et al.* 2012 Generalized Lévy walks and the role of chemokines in migration of effector CD8+ T cells. *Nature*, **486**(7404), 545–548. doi:10.1038/nature11098.

5. Edwards, A. M., Phillips, R. A., Watkins, N. W., Freeman, M. P., Murphy, E. J., Afanasyev, V., Buldyrev, S. V., da Luz, M. G. E., Raposo, E. P. *et al.* 2007 Revisiting Lévy flight search patterns of wandering albatrosses, bumblebees and deer. *Nature*, **449**(7165), 1044–1048. doi: 10.1038/nature06199.

6. Reynolds, A. M. 2018 Current status and future directions of Lévy walk research. *Biology open*, **7**. doi:10.1242/bio.030106.

7. Bartumeus, F. 2007 Levy processes in animal movement: an evolutionary hypothesis. *Fractals*, **15**(02), 151–162. doi:10.1142/S0218348X07003460.

8. Ribeiro-Neto, P. J., Raposo, E. P., Arajo, H. A., Faustino, C. L., da Luz, M. G. E. & Viswanathan, G. M. 2012 Dissipative Lévy random searches: universal behavior at low target density. *Phys Rev E Stat Nonlin Soft Matter Phys*, **86**(6 Pt 1), 061 102.

9. Wosniack, M. E., Santos, M. C., Raposo, E. P., Viswanathan, G. M. & da Luz, M. G. E. 2017 The evolutionary origins of Lévy walk foraging. *PLoS computational biology*, **13**, e1005 774. doi:10.1371/journal.pcbi.1005774.

10. Reynolds, A. 2015 Liberating Lévy walk research from the shackles of optimal foraging. *Physics of life reviews*, **14**, 59–83. doi:10.1016/j.plrev.2015.03.002.

11. Zaburdaev, V., Denisov, S. & Klafter, J. 2015 Lévy walks. *Rev. Mod. Phys.*, **87**, 483–530. doi:10.1103/RevModPhys.87.483.

12. Beltman, J. B., Marée, A. F. M. & de Boer, R. J. 2009 Analysing immune cell migration. *Nat Rev Immunol*, **9**(11), 789–798. doi:10.1038/nri2638.

13. Krummel, M. F., Bartumeus, F. & Gérard, A. 2016 T cell migration, search strategies and mechanisms. *Nature reviews. Immunology*, **16**, 193–201. doi:10.1038/nri.2015.16.

14. Genthon, A. 2020 The concept of velocity in the history of brownian motion. *The European Physical Journal H*, **45**(1), 49–105.

15. Miller, M., Wei, S., Parker, I. & Cahalan, M. 2002 Two-photon imaging of lymphocyte motility and antigen response in intact lymph node. *Science*, **296**(5574), 1869–73.

16. Castellino, F., Huang, A. Y., Altan-Bonnet, G., Stoll, S., Scheinecker, C. & Germain, R. N. 2006 Chemokines enhance immunity by guiding naive cd8+ t cells to sites of cd4+ t cell-dendritic cell interaction. *Nature*, **440**, 890–895. doi:10.1038/nature04651.

17. Mrass, P., Petravic, J., Davenport, M. P. & Weninger, W. 2010 Cell-autonomous and environmental contributions to the interstitial migration of T cells. *Seminars in immunopathology*, **32**, 257–274. doi:10.1007/s00281-010-0212-1.

18. Banigan, E. J., Harris, T. H., Christian, D. A., Hunter, C. A. & Liu, A. J. 2015 Heterogeneous CD8+ T cell migration in the lymph node in the absence of inflammation revealed by quantitative migration analysis. *PLoS Comput Biol*, **11**(2), e1004 058. doi:10.1371/journal.pcbi.1004058.

19. Ariotti, S., Beltman, J. B., Chodaczek, G., Hoekstra, M. E., van Beek, A. E., Gomez-Eerland, R., Ritsma, L., van Rheenen, J., Marée, A. F. M. *et al.* 2012 Tissue-resident memory CD8+ T cells continuously patrol skin epithelia to quickly recognize local antigen. *Proc Natl Acad Sci U S A*, **109**(48), 19 739–19 744.

20. Mrass, P., Oruganti, S. R., Fricke, G. M., Tafoya, J., Byrum, J. R., Yang, L., Hamilton, S. L., Miller, M. J., Moses, M. E. *et al.* 2017 ROCK regulates the intermittent mode of interstitial T cell migration in inflamed lungs. *Nature communications*, **8**, 1010. doi:10.1038/s41467-017-01032-2.

21. Ménard, R., Tavares, J., Cockburn, I., Markus, M., Zavala, F. & Amino, R. 2013 Looking under the skin: the first steps in malarial infection and immunity. *Nat Rev Microbiol*, **11**(10), 701–712. doi:10.1038/nrmicro3111.

22. Gething, P. W., Casey, D. C., Weiss, D. J., Bisanzio, D., Bhatt, S., Cameron, E., Battle, K. E., Dalrymple, U., Rozier, J. *et al.* 2016 Mapping plasmodium falciparum mortality in africa between 1990 and 2015. *The New England journal of medicine*, **375**, 2435–2445. doi: 10.1056/NEJMoa1606701.

23. Aleshnick, M., Ganusov, V. V., Nasir, G., Yenokyan, G. & Sinnis, P. 2020 Experimental determination of the force of malaria infection reveals a non-linear relationship to mosquito sporozoite loads. *PLoS pathogens*, **16**, e1008 181. doi:10.1371/journal.ppat.1008181.

24. Schmidt, N. W., Podyminogin, R. L., Butler, N. S., Badovinac, V. P., Tucker, B. J., Bahjat, K. S., Lauer, P., Reyes-Sandoval, A., Hutchings, C. L. *et al.* 2008 Memory CD8 T cell responses exceeding a large but definable threshold provide long-term immunity to malaria. *Proc Natl Acad Sci U S A*, **105**(37), 14 017–22.

25. Krzych, U., Dalai, S., Zarling, S. & Pichugin, A. 2012 Memory CD8 T cells specific for plasmodia liver-stage antigens maintain protracted protection against malaria. *Front Immunol*, **3**, 370. doi: 10.3389/fimmu.2012.00370.

26. Fernandez-Ruiz, D., Ng, W. Y., Holz, L. E., Ma, J. Z., Zaid, A., Wong, Y. C., Lau, L. S., Mollard, V., Cozijnsen, A. *et al.* 2016 Liver-resident memory CD8(+) T cells form a front-line defense against malaria liver-stage infection. *Immunity*, **45**, 889–902. doi:10.1016/j.immuni.2016.08.011.

27. Guidotti, L. G., Inverso, D., Sironi, L., Di Lucia, P., Fioravanti, J., Ganzer, L., Fiocchi, A., Vacca, M., Aiolfi, R. *et al.* 2015 Immunosurveillance of the liver by intravascular effector CD8(+) T cells. *Cell*, **161**, 486–500. doi:10.1016/j.cell.2015.03.005.

28. McNamara, H. A., Cai, Y., Wagle, M. V., Sontani, Y., Roots, C. M., Miosge, L. A., O'Connor, J. H., Sutton, H. J., Ganusov, V. V. *et al.* 2017 Up-regulation of LFA-1 allows liver-resident memory T cells to patrol and remain in the hepatic sinusoids. *Science Immunology*, pp. 1–10.

29. Holz, L. E., Prier, J. E., Freestone, D., Steiner, T. M., English, K., Johnson, D. N., Mollard, V., Cozijnsen, A., Davey, G. M. *et al.* 2018 CD8 T cell activation leads to constitutive formation of liver tissue-resident memory T cells that seed a large and flexible niche in the liver. *Cell reports*, **25**, 68–79.e4. doi:10.1016/j.celrep.2018.08.094.

30. Wu, P.-H., Giri, A., Sun, S. X. & Wirtz, D. 2014 Three-dimensional cell migration does not follow a random walk. *Proceedings of the National Academy of Sciences of the United States of America*, **111**, 3949–3954. doi:10.1073/pnas.1318967111.

31. Wu, P.-H., Giri, A. & Wirtz, D. 2015 Statistical analysis of cell migration in 3D using the anisotropic persistent random walk model. *Nature protocols*, **10**, 517–527. doi:10.1038/nprot.2015.030.

32. Li, H., Qi, S., Jin, H., Qi, Z., Zhang, Z., Fu, L. & Luo, Q. 2015 Zigzag generalized lévy walk: the in vivo search strategy of immunocytes. *Theranostics*, **5**, 1275–1290. doi:10.7150/thno.12989.

33. Clauset, A., Shalizi, C. R. & Newman, M. E. J. 2009 Power-law distributions in empirical data. *SIAM Rev.*, **51**(4), 661–703. doi:10.1137/070710111.

34. Buldyrev, S. V., Raposo, E. P., Bartumeus, F., Havlin, S., Rusch, F. R., da Luz, M. G. E. & Viswanathan, G. M. 2021 Comment on "Inverse square Levy walks are not optimal search strategies for D=2". *Physical review letters*, **126**, 048 901. doi:10.1103/PhysRevLett.126.048901.

35. Guinard, B. & Korman, A. 2021 Intermittent inverse-square Levy walks are optimal for finding targets of all sizes. *Science advances*, **7**. doi:10.1126/sciadv.abe8211.

36. Palyulin, V. V., Chechkin, A. V. & Metzler, R. 2014 Levy flights do not always optimize random blind search for sparse targets. *Proceedings of the National Academy of Sciences of the United States of America*, **111**, 2931–2936. doi:10.1073/pnas.1320424111.

37. Letendre, K., Donnadieu, E., Moses, M. E. & Cannon, J. L. 2015 Bringing statistics up to speed with data in analysis of lymphocyte motility. *PloS one*, **10**, e0126 333. doi:10.1371/journal.pone.0126333.

38. Volpe, G. & Volpe, G. 2017 The topography of the environment alters the optimal search strategy for active particles. *Proceedings of the National Academy of Sciences of the United States of America*, **114**, 11 350–11 355. doi:10.1073/pnas.1711371114.

39. Levernier, N., Textor, J., Benichou, O. & Voituriez, R. 2020 Inverse square Levy walks are not optimal search strategies for D=2. *Physical review letters*, **124**, 080 601. doi:10.1103/PhysRevLett.124.080601.

40. Levernier, N., Textor, J., Bénichou, O. & Voituriez, R. 2021 Reply to "Comment on 'Inverse square Levy walks are not optimal search strategies for D=2"'. *Physical review letters*, **126**, 048 902. doi:10.1103/PhysRevLett.126.048902.

41. Reynolds, A. M., Cecere, J. G., Paiva, V. H., Ramos, J. A. & Focardi, S. 2015 Pelagic seabird flight patterns are consistent with a reliance on olfactory maps for oceanic navigation. *Proceedings. Biological sciences*, **282**. doi:10.1098/rspb.2015.0468.

42. Reynolds, A. M. 2018 Passive particles lévy walk through turbulence mirroring the diving patterns of marine predators. *Journal of Physics Communications*, **2**(8), 085 003.

43. Zumofen & Klafter 1995 Laminar-localized-phase coexistence in dynamical systems. *Physical review. E, Statistical physics, plasmas, fluids, and related interdisciplinary topics*, **51**, 1818–1821. doi:10.1103/physreve.51.1818.

44. Kareiva, P. M. & Shigesada, N. 1983 Analyzing insect movement as a correlated random walk. *Oecologia*, **56**, 234–238. doi:10.1007/BF00379695.

45. Murray, C. D. 1926 The physiological principle of minimum work applied to the angle of branching of arteries. *The Journal of general physiology*, **9**, 835–841. doi:10.1085/jgp.9.6.835.

46. Kanazawa, K., Sano, T. G., Cairoli, A. & Baule, A. 2020 Loopy lévy flights enhance tracer diffusion in active suspensions. *Nature*, **579**, 364–367. doi:10.1038/s41586-020-2086-2.

47. Miller, M., Wei, S., Cahalan, M. & Parker, I. 2003 Autonomous T cell trafficking examined in vivo with intravital two-photon microscopy. *Proc Natl Acad Sci U S A*, **100**(5), 2604–9.

48. Bajénoff, M., Egen, J. G., Koo, L. Y., Laugier, J. P., Brau, F., Glaichenhaus, N. & Germain, R. N. 2006 Stromal cell networks regulate lymphocyte entry, migration, and territoriality in lymph nodes. *Immunity*, **25**, 989–1001. doi:10.1016/j.immuni.2006.10.011.

49. Krishnamurty, A. T. & Turley, S. J. 2020 Lymph node stromal cells: cartographers of the immune system. *Nature immunology*, **21**, 369–380. doi:10.1038/s41590-020-0635-3.

50. Novkovic, M., Onder, L., Cupovic, J., Abe, J., Bomze, D., Cremasco, V., Scandella, E., Stein, J. V., Bocharov, G. *et al.* 2016 Topological small-world organization of the fibroblastic reticular cell network determines lymph node functionality. *PLoS biology*, **14**, e1002 515. doi: 10.1371/journal.pbio.1002515.

51. Zenkov, V. S., O'Connor, J., Cockburn, I. A. & Ganusov, V. V. 2020 A minority of liver-localized CD8 T cells demonstrate difficult-to-detect attraction to Plasmodium-infected hepatocytes. *bioRxiv*. doi:https://doi.org/10.1101/2020.12.12.422451.

52. Cockburn, I. A., Amino, R., Kelemen, R. K., Kuo, S. C., Tse, S.-W., Radtke, A., MacDaniel, L., Ganusov, V. V., Zavala, F. *et al.* 2013 In vivo imaging of CD8+ T cell-mediated elimination of malaria liver stages. *Proc Natl Acad Sci U S A*, **110**(22), 9090–9095. doi: 10.1073/pnas.1303858110.

53. Kelemen, R. K., Rajakaruna, H., Cockburn, I. A. & Ganusov, V. V. 2019 Clustering of activated CD8 T cells around malaria-infected hepatocytes is rapid and is driven by antigen-specific cells. *Frontiers in immunology*, **10**, 2153. doi:10.3389/fimmu.2019.02153.

54. Moses, M. E., Cannon, J. L., Gordon, D. M. & Forrest, S. 2019 Distributed adaptive search in t cells: Lessons from ants. *Frontiers in immunology*, **10**, 1357. doi:10.3389/fimmu.2019.01357.

55. Wosniack, M. E., Santos, M. C., Raposo, E. P., Viswanathan, G. M. & da Luz, M. G. E. 2015 Robustness of optimal random searches in fragmented environments. *Physical review. E, Statistical, nonlinear, and soft matter physics*, **91**, 052 119. doi:10.1103/PhysRevE.91.052119.

56. Stanley, H., Birgeneau, R., Reynolds, P. & Nicoll, J. 1976 Thermally driven phase transitions near the percolation threshold in two dimensions. *Journal of Physics C: solid State Physics*, **9**(20), L553.

57. Shlesinger, M. F. 1982 Weierstrassian Levy flights and self-avoiding random walks. *J Chem Phys*, **78**, 416–419. doi:10.1063/1.444518.

58. Ta-Chih Lee, R. L. K. & Chu, C.-N. 1994 Building skeleton models via 3-D medial surface/axis thinning algorithms. *Computer Vision, Graphics, and Image Processing*, **56**, 462–478.

59. Kollmannsberger, P., andFelix Repp, M. K., Wagermaier, W., Weinkamer, R. & Fratzl, P. 2017 The small world of osteocytes: connectomics of the lacuno-canalicular network in bone. *New Journal of Physics*, **19**, 073 019.

60. Newton, I. 1846 *Newton's Principia: the mathematical principles of natural philosophy.* New York: Daniel Adee.

61. Jerison, E. R. & Quake, S. R. 2020 Heterogeneous T cell motility behaviors emerge from a coupling between speed and turning in vivo. *eLife*, **9**. doi:10.7554/eLife.53933.

62. Cockburn, I. A., Tse, S.-W. & Zavala, F. 2014 CD8+ T cells eliminate liver-stage Plasmodium berghei parasites without detectable bystander effect. *Infect Immun*, **82**(4), 1460–1464. doi: 10.1128/IAI.01500-13.

Supplemental Information
Environment-imposed constraints can make Brownian walkers efficient searchers

Harshana Rajakaruna, James O'Connor, Ian A. Cockburn, Vitaly V. Ganusov

Materials and Methods

Experimental Details

T cell movement in the liver. Experimental data on movement of wild type (WT), GFP-expressing, TCR transgenic CD8 T cells (OT-I) have been generated similarly as in our previous study[28]. OT-I cells were activated *in vitro* as follows. Splenocytes were incubated with 1 μg/ml SIINFEKL peptide (Biomatik) at 37^0C for 30 minutes, washed, and cultured in complete RPMI 1640 (10% FCS, 2mm L-glutamine, 1mm Na-Pyruvate, 100U/ml Penicillin/Streptomicin, 5mm HEPES, 20 μg/ml Gentimicin and 50 μM β-mercaptoethanol) at 37^0 C, 5% CO_2. Two days later cells were passaged and cultured in fresh media supplemented with (12.5U/ml) recombinant IL-2 (PeproTech). Cells were passaged again after 24 hours, and given fresh media with IL-2. Four days after activation cells were ready for collection and use. Cells were collected from tissue culture flasks, washed, layered over Histopaque® 1083 (Sigma) and centrifuged at 400rcf, for 30 minutes at 21^0C to remove dead cells. Live lymphocytes were collected from the interface between the Histopaque® and cell suspension media. Cells were washed a further 3 times prior to transfer to mice.

To label the sinusoids, Evans blue or Rhodamine-dextran were injected i.v. into mice (50 μg/mouse) immediately prior to imaging of the livers (Fig. 1A&B). In some cases, static images of the livers were acquired using two phone microscope. In total, we have generated images of the liver sinusoids from 5 mice (with or without T cells).

Mice were prepared for multi-photon microscopy essentially as previously described[28]. Briefly, mice were anesthetized with a mix of Ketamine (100 mg/kg) and Xylazine (10 mg/kg). Throughout the surgery and imaging procedure the mouse temperature was maintained at 37^oC using a heating mat attached to feedback probe inserted in the mouse rectum. A lateral incision was made over the left lobe of the liver and any vessels cauterized by applying light pressure to the vessel until clotting occurred naturally. The mouse was then placed in a custom made holder. The liver was then exposed and directly adhered to a coverslip that was secured in the holder. Once stable the preparation was transferred to a Fluoview FVMPE-RS multiphoton microscope system (Olympus) equipped with a XLPLN25XWMP2 objective (25x; NA1.05; water immersion; 2mm working distance). For analysis of the motility of liver-localized CD8 T cells a 50 μm z-stack (2 μm/ slice) was typically acquired using a standard galvano-scanner at a frame rate of 2-4 frames per minute.

Other datasets for movement of WT and LFA-1-deficient CD8 T cells, or WT CD8 T cells 1 or 4 weeks after immunization with radiation-attenuated *Plasmodium berghei* sporozoites Pb-CS[5M] were generated previously[28].

Imaging data processing and basic analyses. Raw imaging data was analyzed with Imaris x64 software (Bitplane) v9.12. Tracking of individual cells in a z-stack was performed using the 'Spots' function in the surpass mode, and in a single slice by the 'Surfaces' function in surpass mode. Detection of individual cells relied upon their relative fluorescence intensity, size (diameter ≥ 9 μm) and used the remaining default function settings for the calculation of the motion tracks using

an autoregressive motion algorithm. Tracks were also manually inspected and curated. Basic cell characteristics such as average track speed, meandering index and arrest coefficient were calculated using Imaris. To calculate the percentage of T-cell movements that were occurring in the sinusoids we generated a co-localization channel which calculated colocalized signal of both red and green fluorescence within each frame. To do this, we utilised the colocalization function on Imaris with the threshold defined in Imaris as 16 for uGFP OT-1 cells which subtracted background fluorescence of the liver, and 3000 for rhodamine dextran. We then calculated the total number of frames a T-cell was tracked using the unique cell ID, and compared this to the total frames each individual T-cell was detected in the designed colocalization channel. Rhodamine dextran fluorescence was dependent on the depth of imaging, and in the lower Z-stacks the frequency of T-cells with colocalized signal was poor (results not shown).

Data details and data cleaning steps. In our analysis we used 3D coordinates of Plasmodium-specific CD8 T cells in the liver over time from the following experiments:

1. Novel data on movement of GFP-expressing activated CD8 T cells called "New13" and "New26". In the "New13" dataset data were generated from three mice and frames were sampled at 14.63 (14.63-14.64) sec apart. In the "New26" dataset data were generated from two mice, and frames were sampled on average at 25 (24.29, 25.51) sec apart. To increase the power in the analysis we generated a combined dataset "New1326" by merging data from datasets "New13" and "New26". We ignored the mid time-points in time-frames in the "New13" dataset, thus, computing time-frames approximately at 26 sec frequency.

2. Previously published data on movement of WT and LFA1-deficient CD8 T cells in the liver[28] resulted in "Lfa1 WT" and "Lfa1 KO" datasets. Data were from 7 mice, and frames were sampled, on average, at 27 (24.65 - 29.56) sec apart.

3. Previously published data on movement of *Plasmodium*-specific CD8 T cells following vaccination of mice with radiation-attenuated sporozoites[28]. In "Wk1" dataset data were generated from two mice immunized with RAS 1 week before. Frames were sampled, on average, at 6.51 (6.51-6.51) sec apart. In "Wk4" dataset five mice were RAS-immunized 4 weeks prior to imaging, and frames were sampled, on average, at 3.25 (3.25-3.26) sec apart.

4. Novel data on movement of *Plasmodium*-specific (OT-1) and lymphocytic choriomeningitis virus-specific (P14) CD8 T cells moving in livers of *Plasmodium berghei*-infected mice[51].

In the Imaris-processed movies we found that there were random missing time-frames in tracks in nearly all data. This could happen when Imaris would associate a cell that went out of focus for one time step with other time steps. Because having a consistent time step to determine cell movement characteristics such as mean squared displacement and movement length distribution was critical we cleaned the data in the following ways. We split tracks with missing time frames and converted them to individual cells for the analyses. For example, if one time frame was missing for a cell, the track was split into two with the second part being assigned as a new "cell". This created many more cells with shorter tracks. Taken together, cleaning up the data yielded the following. For "New13" dataset we originally had 56 T cells out of which we generated 14 new split cells with 1-2 splits resulting in total of 70 T cells with 2565 time-frames (movements). For "New26" dataset we had 213 T cells with 51 new split cells with 1-2 splits from the original tracks totaling to 264 T cells, with 7736 time-frames. For the combined "New1326" dataset we had 334 T cells with 9018 time-frames. For "Lfa1 WT" dataset we originally had 198 T cells out of which we generated 19 new

split cells with 1-2 splits from the original tracks totaling to 217 T cells, with 6210 time-frames. For "Lfa1 KO" dataset we originally had 154 T cells out of which we generated 16 new split cells with 1-2 splits from the original tracks totaling to 170 T cells, with 5097 time-frames. For "Wk1" dataset we had 99 T cells and generated 3 new split cells with 1-2 splits from the original tracks, totaling to 102 T cells, with 2525 time-frames. For "Wk4" dataset we had 44 T cells with 4169 time-frames that required no cleaning. Most of the analyses have been repeated on "clean" data in which tracks with missing time-frames were removed with similar conclusions reached.

T cell movement in the LNs. Data have been provided by Mario Novkovic from their previous publication[50]. In short, 3×10^6 naive P14 TCR transgenic T cells were transferred i.v. into naive mice. Three to twenty-four hours after T cell injection, intravital imaging was performed on the right popliteal LNs. To detect FRCs, anti-YFP mAbs were used and samples were imaged using a confocal microscope (LSM-710; Carl Zeiss)[50]. In total we had 3 sets of images of LNs with FRCs labeled (about 30 images per set with 2 μm depth). In addition, we analyzed 3D movement data of T cells in the LNs (in total 6 movies) produced in the same publication[50]. The data were also cleaned and tracks for which there were not always consecutive timeframes available were split as described above. Analyses were repeated with "clean" data in which cells with missing time-frames were removed; similar results were found. These data were generated at frequency of 20 sec per frame.

Modeling and Statistical Analytical Methods

Mean squared displacement, turning angles, and velocity correlations. For each of the datasets we calculated several basic characteristics. To calculate mean squared displacement (MSD) we used all tracks in the dataset in which time frequency of measurements was approximately the same:

$$\text{MSD}(t) = \frac{1}{N(t)} \sum_{i=1}^{N(t)} r_{i,t}^2, \tag{S.1}$$

where $r_{i,t}$ is the displacement of an i^{th} cell from its initial position to the position at time t and $N(t)$ is the number of cells for which coordinate data were available at time t. Note that this is different from movement lengths — which are movements T cells make between sequential time frames. To characterize the MSD we used the following relation $\text{MSD}(t) = ct^{\gamma}$ where t are lag-times or delays, and γ was estimated by log-log transforming MSD and t. Estimate of $\gamma < 1$ suggests subdiffusion of cells, whose walks are constrained by boundaries, regardless of their step length distribution, $\gamma = 1$ suggests Brownian (or normal) diffusion, whose walks are affected by neither boundaries nor transportation, whereas, $\gamma > 1$ suggests superdiffusion, whose walks are indicative of transportation or directed movements. Root mean squared displacement, $\zeta = \text{RMSD}$ was calculated as

$$\zeta(t) = \text{RMSD}(t) = \sqrt{\text{MSD}}, \tag{S.2}$$

where MSD is defined in eqn. (S.1). To calculate turning angle for cell movement between time steps $(t, t+1)$ and $(t+1, t+2)$ we calculated the vectors of movements between two time points and then calculated the angle between these two movement vectors. To estimate decay in the correlation movements of a T cell over time we calculated the vector for the initial T cell velocity $\bar{v}_1$ (movement

vector for a cell from $t = 0$ to $t = 1$), vector for later movement $\bar{v}_t$ (movement vector for a cell between times $t - 1$ and t), and the angle $\phi_{1,t}$ between the two vectors $\bar{v}_1$ and $\bar{v}_t$:

$$c_V = \cos(\phi_{1,t}), \tag{S.3}$$

where by definition c_V varies between -1 and 1.

Basic statistical distributions to describe movement length data. To investigate which model is best consistent with the distribution of movement/step lengths of activated CD8 T cells in the liver or distribution of branch lengths of liver sinusoids we fit several alternative distributions to the data. Cauchy distribution is defined as

$$f(r|\gamma) = \frac{1}{\pi\gamma} \left[\frac{\gamma^2}{\gamma^2 + (r - r_{\min})^2} \right], \tag{S.4}$$

where $r \geq r_{\min}$ is the displacement, $r_{\min}$ is the location parameters, and the mean of the distribution is undefined. Half-gaussian distribution with $r \geq 0$ is defined as

$$f(r|\sigma) = \frac{\sqrt{2}}{\sqrt{\pi}\sigma} e^{-\frac{r^2}{2\sigma^2}}, \tag{S.5}$$

where mean of the half-normal distribution is $\bar{r} = \sigma\sqrt{2/\pi}$.

Stable Levy (SL) distribution is defined as

$$f(r|\gamma) = \frac{\sqrt{\gamma}}{\sqrt{2\pi}(r - r_{\min})^{3/2}} e^{-\frac{\gamma}{2(r - r_{\min})}}, \tag{S.6}$$

where γ and $r_{\min}$ are the shape parameter and location parameter, respectively. The SL distribution has infinite mean and variance.

Generalized Pareto (GP) distribution is defined as

$$f(r|k, \sigma, \theta) = \frac{1}{\sigma} \left(1 + \frac{k(r - \theta)}{\sigma} \right)^{-1 - \frac{1}{k}}, \tag{S.7}$$

where k is the shape parameter, σ is the scale parameter, and θ is the location parameter. In this distribution, the mean is $\bar{r} = \theta + \sigma/(1 - k)$.

When $k = 0$ and $\theta = 0$, GP becomes exponential distribution

$$f(r|\lambda) = \lambda e^{-\lambda r}, \tag{S.8}$$

where $\lambda = 1/\sigma$, and mean being equal to $\bar{r} = \sigma$. When $k > 0$, $\theta \to 0$ GP becomes the Pareto (or Powerlaw) distribution

$$f(r|r_{\min}, \alpha) = \frac{\alpha r_{\min}^{\alpha}}{r^{\alpha+1}} = \frac{(\mu - 1)r_{\min}^{\mu-1}}{r^{\mu}}, \tag{S.9}$$

where $r_{\min} = \sigma/k$ (scale parameter), $\alpha = 1/k$, $\mu = \alpha + 1$ (shape parameter), and $\bar{r} = (\mu-1)r_{\min}/\mu$.
Note that in Pareto distribution $r \geq r_{\min}$. The important property of the Pareto distribution,
describing either Brownian or Levy walks, is dependence of the mean and variance on the value of
μ (and thus α). For $\mu > 3$ ($\alpha > 2$), both mean and variance of the Pareto distribution are finite
corresponding to Brownian walk. In contrast, when $\mu \leq 3$ ($\alpha \leq 2$) mean is finite and variance is
infinite corresponding to Levy walks. Finally, when $\mu \leq 2$ ($\alpha \leq 1$), both mean and variance are
infinite corresponding to bullet motion [11].

Generalized extreme value (GeV) distribution is defined as

$$f(r|\mu, \sigma, \xi) = \frac{1}{\sigma}t(r)^{\xi+1}e^{-t(r)}, \tag{S.10}$$

where $t(r) = \begin{cases} (1 + \xi\frac{(r-\mu)}{\sigma})^{-1/\xi}, & \text{if } \xi \neq 0, \\ e^{-(r-\mu)/\sigma}, & \text{otherwise.} \end{cases}$

Statistical methodology of fitting models to data. We fit the models (eqns. (S.4)–(S.9))
to either actual displacements r of cells between two time points (i.e., movement lengths) or scaled
displacements ρ. Scaled displacements are defined as $\rho = r/\bar{r}$, where $\bar{r}$ is the average displacement in
the particular dataset ($\bar{r} = \frac{1}{n}\sum_{i=1}^{n} r_i$ for n movements). Using scaling of the displacements by square
room of mean of squared movement lengths ($\sqrt{r^2}$) gave similar results. It is important to emphasize
that scaled displacements ρ allows to convert cell tracks to be independent of the effect of different
mean speeds of cells and independent of the effect of different sampling frequency [4]. Likelihood of
the model parameters given the data is given as

$$L(m_1, m_2, ...|x_1, x_2, ..., x_n) = \prod_{i=1}^{n} f(x_i|m_1, m_2, ...), \tag{S.11}$$

where x_i are either the cell movement length data $x_i = r_i$ or the scaled displacement data $x_i = \rho_i(t) = r_i/\zeta_i(t)$, for $i = 1...n$, for n data points, for fixed time intervals t. The probability density
functions $f(x_i|m_1, m_2, ...)$ are given in eqns. (S.4)–(S.9). Parameters $m_1, m_2, ...$ of the models were
estimated by minimizing the log of negative likelihood, $\mathcal{L} = -\ln L$.

Tail analysis of the step length distribution. In addition to fitting different distributions to
the movement length/displacement data we performed an analysis to determine the shape parameter
of the powerlaw/Pareto distribution μ that is best consistent with tail of the displacement distribution
data [33]. Given that Pareto distribution has two parameters ($r_{\min}$ and μ, eqn. (S.9)) one needs to define
$r_{\min}$ that determine the cut-off in the data and that allows for the best description of the tail of the
data with Pareto distribution. The steps to define $r_{\min}$ are as follows [33]: (1) for each possible choice
of $r_{\min}$ from the smallest to the largest unique values of r, we computed μ by maximum likelihood
estimate given by the analytic expression $\hat{\mu} = 1 + n\left(\sum_{i=1}^{n} \ln\frac{r_i}{r_{\min}}\right)^{-1}$, where r_i, $i = 1...n$, are the
observed values of r and $r_i \geq r_{\min}$. In doing so, we also set a limit to the number of data points
in the analysis by setting a criteria by which the analysis is terminated if the standard error on $\hat{\mu}$,
approximately given by $\sigma = \frac{\hat{\mu}-1}{\sqrt{n}}$, was greater than 0.1, as higher order corrections of the standard
error are generally positive. This condition was set because the estimates of μ can be erroneously
large when $r_{\min}$ is so large that the number of observed $r \geq r_{\min}$ is very small. It was also assumed
that $\mu > 1$, since the distributions with $\mu \leq 1$ are not normalizable, and hence not realistic in
nature. We selected $r_{\min}$, at which the Kolmogorov-Smirnov (K-S) goodness-of-fit statistic D was
the minimum out of all D values given by the fits over all possible unique $r_{\min}$'s of the data. Here,

we also made a finite correction to the estimates of μ for small sample sizes ($n < 50$) in addition to
the condition set by the standard error of μ.

From the above analyses, we carried forward the estimated $r_{\min}$, μ, and D to the next stage of
testing the probability of the data, given that the estimated model is the plausible one, by the method
of Kolmogorov-Smirnov goodness-of-fit test. To do that we generated data from the hypothesized
(or estimated) Pareto model, sampling using a semi-parametric bootstrap method, computing D for
each sample, which is the synthetic D, and thus, computing a fraction of those were greater than
the empirical D of the given data, which yields the p value of the hypothesis. We then computed
the uncertainty in the estimates of $r_{\min}$, μ, and n, given by standard deviations, by resampling the
data with replacements by Markov chain Monte Carlo (MCMC) simulations.

Digitizing and skeletonizing the 3D sinusoidal structure. We analyzed 23 z-stacks of
Evans blue stained 512×512 μm images of the liver (mouse 1: 171207 and mouse 2: 190624), as
well as 3 sets of images/movies with T cells in which liver sinusoids were labeled with rhodamine-
dextran. Resulting structures were similar based on calculated characteristics such as branching angle
distribution and straightness index (results not shown). Light-adjusted 2D images were converted
from RGB into grayscale using rgb2gray, filtered through Matlab filters (1) imfilter with Gaussian
N(10,2) to remove noise, (2) medfilt2, with 'symmetric' setting to remove all the lines from the image,
(3) uint8 to subtract image without lines from image with lines, (4) setting threshold at > 130 of
the subtracted image, and (5) removing more noise by imopen, and saving all 2D images in a stack
converting to a digitized 3D image, with "1" giving the sinusoids and "0" giving the hepatocytes
and other unstained tissues (Fig. 3). The digitized 3D image was then skeletonized with 1's giving
the mean path of sinusoids and 0's elsewhere, converting to a voxel image, using parallel medial axis
thinning method as per the MATLAB vectorized implementation of the algorithm[58]. The 3D binary
voxel skeleton was converted to a network graph, computing nodes (junctions) and edges (branches),
using previously published methods[59], and mapping the respective skeleton to 3D graphics function.
The nodes and the branches were used to compute branch angles, branch lengths, node-to-node
direct distances, number of branches per node. Straightness of branches metric was computed as
the node-to-node direct distance over the branch length. The binary 3D liver image was used to run
simulations of cells walking on the 1's on the lattice, scale of which can be decided, for e.g., 1 unit
equals 0.01 μm (Fig. 4).

Simulation methods of cell walks on digitized sinusoidal structures and analyses
of re-sampled step-length distribution and MSD. We simulated cell walks on (i) digitized
3D real sinusoidal branch structures (5 imaging sets), (ii) simulated sinusoidal branch structures
generated from both Levy-like (5 simulations) or truncated Pareto and Gaussian (5 simulations)
branch length distributions, with branch angles generated from random (spherical uniform) and
empirical distributions from real structural samples, and (iii) no structures (5 simulations) imposing
no physical constraints giving freedom for cells to turn any direction after every step. For each
generated structure we selected a random point on a mean sinusoidal path, i.e. on the digital
skeleton given by 1's, on the matrices generated from digitized 3D sinusoidal structures of 1's and
0's in the real case, and on a branch position coordinate in the simulated case, as the starting
point of a T cell. Then we selected a cell step-length from the following specified displacement
distributions in unit time: (i) phenomenologically best-fitted truncated Generalized Extreme Value
distribution with parameter $1 > \xi > 0.5$, i.e, heavy-tailed Levy-like with infinite variance, yet
finite mean (eqn. (S.10)); (ii) half-Gaussian (eqn. (S.5)); (iii) Composite models with Pareto tails,
approximating Levy and Gaussian tails, with Gamma heads (eqn. (S.9)), and (iv) Convoluted model
(generalized Levy walks as described previously[4,32]), approximating Levy tails in (Pareto) runs and

pauses, or observed and unobserved step-lengths), and walked the cell digitally along the sinusoidal path to the next point on the matrix, sampled the step-end coordinates at fixed time intervals. Note that we considered all movements as forward movements, as reverse movements of cells along the sinusoids were the minority in the data (about 20%, Fig. S3). Ability of cells to move in one preferred direction in constant external conditions can be viewed similarly to the physics postulate of momentum conservation (or "biological inertia",[60]). We disregarded deviation of tracking cell centers from the mean sinusoidal path because the cell diameter, on average, 10 μm, was greater than the cylindrical diameter of the sinusoids, 7 μm, and thus, T cells likely occupied most of the cross-sectional area of the sinusoids. Note also that step-lengths and branch lengths were thus simulated disregarding their autocorrelations assuming they are independent and identically distributed (i.d.d.). Cell walk simulations on no-structure, the cell turning angles were random from a uniformly spherical distribution. Note that the walk simulations of cells and sampling coordinate were terminated once a cell hits the boundary of the structure before the simulation time was elapsed in the case of real sinusoidal structures.

We simulated cell walks on i) no structure (i.e., free 3D space), or ii) digitized 3D real sinusoidal structure. To simulate T cell walk in digitized liver sinusoids, we selected a random point on a mean sinusoidal path, i.e. on the digital skeleton given by 1's, on the matrices generated from digitized 3D sinusoidal structures of 1's and 0's. We selected step-lengths of the simulated walks from a Pareto distribution with $\mu = 2.5$ and $\mu = 4.5$ with the same mean (eqn. (S.9)), assuming Levy vs. Brownian walk scenarios, and motion of cells strictly forward along the sinusoid mean paths. We chose 200 cells with each walking 50 steps to match the average time observations of the real data within the limited 3 dimensions of the structure. Simulation with no structure we did the same and selected turning angles from a uniform spherical distribution. We computed MSD of all four scenarios. To simulate T cell walks in digitized 3D network of FDCs in LNs we proceeded similarly to the method for liver sinusoids assuming that T cells can only walk in the middle of the network path and turns can be taken only when a T cell reaches a branching point.

Simulation methods of computing efficiency of cells finding a target. We computed two metrics of efficiency of cells finding a target on the structure vs. no structure: (i) mean and variance of time-to-reach a random target (100 targets) on the walk path in a structure, given cells with two types of walk characteristics (at least 1,000 cells in each): powerlaw/Pareto with different values of the shape parameter μ; using the same path on (a) Gaussian and truncated-Levy branch structures, with random and empirical branch angles; (b) real structures; and (c) no-structure with uniformly spherically distributed turning angles; (ii) mean and variance of time-to-hit a given random target (300 targets), given that they did not miss the target, for cells (10,000 for each), walking Gaussian vs truncated-Levy step-lengths, starting from a random point on the same structures vs. no-structure, given cell-absorbing structural boundaries in the structures, assuming cells absorbed missed the target; and (iii) the probability of a cell hitting the target, simulating walks as same as above. We repeated the simulation with 5 different distances to targets, taken geometrically, from the initial points, and also with multiple targets with different densities in space (results not shown). We started simulations at different distances between a T cell and the target and assumed that T cell found the infected cell when it was within 40 μm from the target, an approximate size of the murine hepatocyte[52,53]. For 3D simulations if searching T cell reached the boundary we assumed that it missed the target (and thus was not included in the calculation of the average time). For simulating T cells searching for a target in the liver we simulated T cells moving forward on one path, effectively reducing search problem to 1D. This is reasonable given that difference between Brownian and Levy walks is only on the movement length distributions and not on turning angles that cells take[11]. To simulate n T cells searching for a single sample, assuming that search is done by different cells

independently, we resampled times to target with replacement n times from the dataset on a single T cell searching for a single target, and calculated minimal of the time in the sample. This procedure was them repeated 1,000 times to generate a distribution to target when n T cells search for the infection site.

Detailed characterization of the movement patterns of T cells. In the main text we focused on characterizing movement patterns of CD8 T cells in the liver or the LNs by calculating MSD for cells and the distribution of movement lengths and comparing the latter data to several well-characterized parametric distributions. This is a standard approach to detect whether moving agents undergo Brownian or Levy walks[4,5,61]. However, characterizing walk type exclusively using MSD is inappropriate since several alternative movement types may result in similar (or identical) MSD and movement length distributions. Other characteristics such as RMSD, distribution of turning angles or T cell velocities may provide additional information that may explain specific patterns of cell movement[4,32]. Our finding that liver-localized CD8 T cells undergo only small movement steps is fully consistent with the Brownian walk; however, this does not prove that walks are indeed Brownian. We found that T cell movement program in which cells move as generalized Levy walkers with long but rare movements/runs ($\mu_r < 3$ for the Pareto distribution of runs) and with pauses ($\mu_p < 3$ for the Pareto distribution of pauses) can be fully consistent with the overall short displacements observed in our data (results not shown). The only possibility to truly discriminate between these alternatives would be to record movements at extremely high frequencies to see if long distance movements do occur. Importantly, we previously found that liver-localized memory CD8 T cells after 4 weeks post RAS immunization never made long distance jumps when recording was done at 3 frames/sec[28]. However, it is possible that long displacements may occur when liver-localized T cells detach from the endothelial wall and float with the blood flow as we observed with T cells after 1 week post RAS immunization (Fig. 2)[28]. Such movements, however, are better described as Levy flights and not Levy walks[11] because the speed at which T cells float with the blood in the liver dramatically exceeds average movement velocity of cells crawling on endothelium.

 # Additional movies, figures, and tables

Movie S1: Example of movement of liver-localized CD8 T cells with labeled liver sinusoids. Experiments were performed as described in Fig. 1 in 3 individual mice. In the movie red denotes rhodamine dextran-labeled liver sinusoids and green are *Plasmodium*-specific liver-localized CD8 T cells (OT-I). Cells turn yellow when there is an overlap between green and red. Bar scale is 50 μm. The volume of $512 \times 512 \times 44$ μm was scanned every 13 sec.

Figure S1: Departure from linear (on log-log scale) change in MSD or root mean squared displacement (RMSD) is in part due to decline in the number of cells with long tracks. For the New1326 datasets of activated CD8 T cells in the liver we calculated the MSD (panel A), RMSD (RMSD = $\sqrt{\text{MSD}}$, panel B), or the number of T cell tracks used to calculate these characteristics (panel C). Note that panel A lists a different slope from that given in Fig. 2A because different number of data points are taken into account in regression (T_p is the persistence time defining which data points were taken for linear regression).

Non scaled data	Dataset	Levy γ	Cauchy γ	GP k	GP σ	Exp λ	H-N σ	Pareto rmin (μm)	Pareto μ-tail	Sampling frequency (sec)
Liver	New13	0.35	1.14	-0.07	1.64	1.53	2.01	3.03	4.63	13.00
Liver	New26	0.48	1.69	0.00	2.51	2.50	3.50	7.09	5.32	26.00
Liver	New1326	0.49	1.69	-0.01	2.51	2.49	3.46	7.09	5.47	26.00
Liver	Wk1	0.13	0.40	0.28	0.50	0.76	2.15	0.61	2.55	6.50
Liver	Wk4	0.10	0.33	0.01	0.46	0.46	0.66	0.94	4.42	3.25
Liver	Lfa1_WT	0.33	1.11	0.18	1.68	2.03	3.11	8.60	6.55	27.50
Liver	Lfa1_KO	0.18	0.57	0.11	0.80	0.91	1.46	2.10	3.30	27.50
LN	0209_Doc6	2.31	7.01	-0.19	10.14	8.72	10.58	22.02	6.67	20.00
LN	0409_Doc7	1.20	3.67	-0.48	5.77	4.16	4.71	6.14	6.74	20.00

Scaled data	Dataset	Levy γ	Cauchy γ	General Pareto k	General Pareto σ	Exp λ	H-N σ	Pareto ρmin	Pareto μ-tail	Sampling frequency (sec)
Liver	New13	0.23	0.74	-0.07	1.07	1.00	1.31	2.01	4.68	13.00
Liver	New26	0.19	0.68	0.00	1.01	1.00	1.41	2.90	5.39	26.00
Liver	New1326	0.20	0.68	-0.01	1.01	1.00	1.39	2.91	5.55	26.00
Liver	Wk1	0.17	0.52	0.28	0.66	1.00	2.82	0.78	2.52	6.50
Liver	Wk4	0.22	0.71	0.01	0.99	1.00	1.44	1.93	4.28	3.25
Liver	Lfa1_WT	0.16	0.55	0.18	0.83	1.00	1.53	4.42	6.47	27.50
Liver	Lfa1_KO	0.20	0.63	0.11	0.88	1.00	1.60	2.29	3.30	27.50
LN	0209_Doc6	0.29	0.87	-0.38	1.30	1.00	1.14	2.05	9.34	20.00
LN	0409_Doc7	0.29	0.88	-0.48	1.39	1.00	1.13	1.55	7.00	20.00

Structure	Dataset	Levy γ	Cauchy γ	GP k	GP σ	Exp λ	H-N σ	Pareto rmin (μm)	Pareto μ-tail	
Liver sinusoids	Evans Blue	3.48	10.54	-0.20	15.71	13.38	16.37	36.36	7.45	
LN FRCs	CCL19day5	2.31	7.01	-0.19	10.14	8.72	10.58	22.02	6.67	

Table S1: Estimates of parameters providing best fits of various datasets. We fitted Levy (eqn. (S.6)), Cauchy (eqn. (S.4)), General Pareto (eqn. (S.7)), exponential (eqn. (S.8)), or half-normal (eqn. (S.5)) to the data on movement of T cells in the liver, lymph nodes, or to branch length distribution data for liver sinusoids or fibroblastic retricular cell (FRC) network in lymph nodes. For distribution with a location parameter $r_{\min}$ we set $r_{\min} = 0$. The same models were also fitted to scaled data (step length distribution data scaled by the RSMD). For imaging data we also list frequency of the imaging in the movies. See Extended Materials and Methods for the models and how models were fit to data.

Non scaled data	Dataset	Levy	Cauchy	GP	Exp	H-N	Levy	Cauchy	GP	Exp	H-N
Liver	New13	11397	11433	7269	7302	7279	0.00	0.00	0.99	0.00	0.01
Liver	New26	40929	41951	29623	29621	30631	0.00	0.00	0.32	0.68	0.00
Liver	New1326	45728	46801	33002	33003	33968	0.00	0.00	0.58	0.42	0.00
Liver	Wk1	5868	5968	2746	3398	6955	0.00	0.00	1.00	0.00	0.00
Liver	Wk4	8184	8275	1860	1859	2575	0.00	0.00	0.39	0.61	0.00
Liver	Lfa1_WT	26909	28181	19649	19789	21556	0.00	0.00	1.00	0.00	0.00
Liver	Lfa1_KO	15307	15615	8554	8668	10549	0.00	0.00	1.00	0.00	0.00
LN	0209_Doc6	42313	41380	28946	30515	28685	0.00	0.00	0.00	0.00	1.00
LN	0409_Doc7	3352	3278	2281	2433	2281	0.00	0.00	0.53	0.00	0.47
	Dataset	AIC					Akaike Weights				
Scaled data		Levy	Cauchy	GP	Exp	H-N	Levy	Cauchy	GP	Exp	H-N
Liver	New13	9219	9254	5090	5123	5099	0.00	0.00	0.99	0.00	0.01
Liver	New26	26806	27820	15491	15489	16494	0.00	0.00	0.32	0.68	0.00
Liver	New1326	30021	31086	17287	17287	18248	0.00	0.00	0.59	0.41	0.00
Liver	Wk1	7141	7242	4020	4673	8231	0.00	0.00	1.00	0.00	0.00
Liver	Wk4	14498	14595	8181	8180	8901	0.00	0.00	0.41	0.59	0.00
Liver	Lfa1_WT	18723	19988	11456	11596	13359	0.00	0.00	1.00	0.00	0.00
Liver	Lfa1_KO	16191	16499	9439	9554	11435	0.00	0.00	1.00	0.00	0.00
LN	0209_Doc6	24765	23829	11394	12965	11133	0.00	0.00	0.00	0.00	1.00
LN	0409_Doc7	1924	1849	853	1005	853	0.00	0.00	0.54	0.00	0.46
Structure	Dataset	Levy	Cauchy	GP	Exp	H-N	Levy	Cauchy	GP	Exp	H-N
Liver sinusoids	Evans Blue	47544	47114	37936	38375	37599	0.00	0.00	0.00	0.00	1.00
LN FRCs	CCL19day5	29759	29402	23058	23349	22758	0.00	0.00	0.00	0.00	1.00

Table S2: AIC values and Akaike weights of alternative models fit to different datasets (as described in Tab. S1). See Extended Materials and Methods for the models and how models were fit to data.

Figure S2: Liver-localized display similar displacement distribution for sub-sampled data. For the cleaned New1326 dataset we calculated the cumulative distribution of displacements for all data (panel A) or when we sub-sample the data, i.e., by taking every time frame ($\Delta t = 26$ sec), every second time frame ($\Delta t = 52$ sec), etc (panel B). In panel B we rescaled all displacements by the average displacement $\bar{r}$ in a given sub-sampled dataset. Results were similar if we scaled the data by dividing displacements by the relative duration of the sub-sampling period ($k = 1, 2 \ldots$). Parameter μ in panel A is from the fit of the Pareto distribution (with $r_{\min} = 9.1$ μm) to the subset of the data (fail fit in which data with $r \geq r_{\min}$ were used).

Figure S3: Activated CD8 T cells in the liver perform correlated random walk with a high degree of persistence. A: For every track in our dataset that includes measurement of T cell movement in the liver at 26 sec time steps, we calculated the turning angles for each subsequent cell movement. The data are from 3 individual mice involving 332 cleaned tracks and 8305 movements ("New1326" dataset). Fraction f denotes the frequency of turning angles between specific cutoff values. B: We calculated the correlation between vectors determining the first movement and every another movement for all 329 cell tracks in our data for which velocity correlations could be calculated. Specifically, we calculated the angle ϕ between the vector determining the first cell movement and every another movement at different times and plotted $\cos(\phi)$ vs. the time difference between first movement and another movements. Insert in B shows the data for the first 20 min of imaging. The exponential decay model ($v = v_0 e^{-t/T_v}$ was fitted to the data and estimated parameter T_v is shown. C: Distribution of average speed per cell in these data. D: cumulative distribution of speeds of all cells int he population. Parameter μ denotes the shape parameter of the Pareto distribution fitted to the tail of these data.

Figure S4: Fits of the Pareto (powerlaw) distributions to the CD8 T cell in the liver displacement data. We used previously published method [33] to estimate the shape parameter μ of the Pareto distribution (eqn. (S.9)) for 6 different datasets (see Materials and Methods for dataset description) by fitting a linear function (on log-log scale) to the tails of the data distribution using likelihood method. We fit the model to actual displacements r (A-C) or to scaled displacements ρ (D-F). The cut-off $r_{\min}$ (for actual displacements) or $\rho_{\min}$ (for normalized displacements) for the linear regressions was chosen using a goodness-of-fit based method [33]. In short, for multiple choices of $r_{\min}$ slope μ was estimated via the method of maximum likelihood and the Kolmogorov-Smirnov goodness of fit statistic, D, was calculated. Value of $r_{\min}$ that resulted in minimal fit statistics D was chosen for the final fit. Indicated p values were estimated by resampling the data 1000 times and performing Kolmogorov-Smirnov Goodness of Fit Test [33].

Figure S5: Activated CD8 T cells are superdiffusive Brownian walkers in the liver during *Plasmodium* infection. We performed experiments in which *in vitro* activated OT-I (*Plasmodium berghei*-specific) and P14 (LCMV-specific) CD8 T cells were transferred i.v. into naive mice and these mice were 1.5-2 hours later infected with 10^5 *Plasmodium berghei*-CS[5M] sporozoites expressing SIINFEKL epitope from chicken ovalbumin[28,62]. Ten to twenty minutes after infection, livers of infected mice were exposed and movements of T cells were recorded using intravital microscopy[51]. We performed the same analyses as in the main text (Fig. 2) including (A) regression analysis of the MSD with time, (B) distribution of movement lengths, (C)-(D) fitting alternative distributions to the movement data for OT1 (C) or P14 (D) T cells. Generalized Pareto distribution was best describing data in C-D (results not shown).

Figure S6: Example of the simulation in which T cells were moving in sinusoids of the liver. Grey denotes sinusoids, red dots denote the mean path of the sinusoids, and green denotes moving T cell.

Figure S7: Characterizing structure of the fibroblastic reticular cell network in the murine lymph nodes. We used previously published data in which FRCs were imaged using Ccl19[eyfp] reporter mouse[50]. As with analysis of liver sinusoids z-stacks of LN images were contrasted and mean paths in the FRCs were calculated. To simulate T cell walk on digitized FRC network we assumed that movement length distribution is described by the powerlaw (Pareto) distribution with $r_{\min} = 0.2$ μm and $\mu = 4.5$ (for Brownian walks) or $\mu = 2.5$ (for Levy walks).

Figure S8: Naive CD8 T cells in lymph nodes are not superdiffusive. We analyzed experimental data in which naive CD8 T cells moving in resting lymph nodes were imaged using two photon microscope[50]. Example of the analysis of data from one of 6 such datasets is show. A: individuals tracks of T cells digitized using Imaris software; B: MSD vs. time for all T cels indicating short-term (2-3 min) persistence/superdiffusion; C: movement length distribution with tail analysis performed[33]; D: fits of several alternative distributions to the movement length distribution data; E: distribution of turning angles of T cells in the LNs; F: velocity autocorrelation function for T cells moving in the LNs.

Figure S9: T cells in the liver display persistent movement for a longer time as compared to T cells in the LNs. Data from all datasets on movement of activated CD8 T cells in the liver (blue) or naive CD8 T cells in the LNs (red) are plotted together. A: difference in MSD for T cells in the LNs vs. liver; B: difference in movement length distribution of T cells in the liver vs. LNs.

Figure S10: Benefits of Levy walks depend on the environment (open space/3D vs. liver/1D) and the number of T cells n searching for the infection site. We simulated T cells searching for an infection similar to that in Fig. 5 either in free 3D space (panel A, starting at $100\mu m$ from the target) or in the liver sinusoids (panel B, starting at $500\mu m$ from the target) assuming that there are different numbers of T cells all independently searching for the infection. For each simulation the time to target was calculated as the minimal time to find the targeted by any T cell in n cells. Brownian walks (BW) were described by the Pareto distribution with $\mu = 4.5$ and Levy walks (LW) were described with $\mu = 2.5$ (eqn. (S.9)). In both cases, the average step length distribution was set to $\bar{r} = 2.1$ μm per time step. Error bars denote 95% range for the estimated time to target. A given T cell reaches the target when it is within $40\mu m$ from the target[52,53].