Predatory selection of mucoid, antibiotic resistant *Pseudomonas putida* phenotype by

myxobacterium *Cystobacter ferrugineus*

Shukria Akbar[a] and D. Cole Stevens[a,#]

Department of BioMolecular Sciences, University of Mississippi, University, MS, USA[a]

Running Head: Predatory selection of antibiotic resistant *P. putida*

#Address correspondence to D. Cole Stevens, stevens@olemiss.edu.

Abstract: Predation contributes to the structure and diversity of microbial communities.

Predatory myxobacteria are ubiquitous to a variety of microbial habitats and capably

consume a broad diversity of microbial prey. Predator-prey experiments utilizing

myxobacteria have provided details into predatory mechanisms and features that facilitate

consumption of prey. However, prey resistance to myxobacterial predation remains

underexplored, and prey resistances have been observed exclusively from predator-prey

experiments that included the model myxobacterium *Myxococcus xanthus*. Utilizing a

predator-prey pairing that instead included the myxobacterium, *Cystobacter ferrugineus,*

with *Pseudomonas putida* as prey, we infrequently observed surviving phenotypes

capable of eluding predation. Comparative transcriptomics between *P. putida* unexposed

to *C. ferrugineus* and the survivor phenotype suggested that increased expression of

efflux pumps, genes associated with mucoid conversion, and various membrane features

contribute to predator avoidance. The *P. putida* survivor phenotype was confirmed to be

resistant to the antibiotics kanamycin, gentamicin, and tetracycline and to produce more

alginate, pyoveridine, and phenazine-1-carboxylic acid than predator-unexposed *P.*

putida. Unique features observed from the survivor phenotype including small colony

variation, efflux-mediated antibiotic resistance, phenazine-1-carboxylic acid production,

and increased mucoid conversion overlap with traits associated with *Pseudomonas*

aeruginosa predator avoidance and pathogenicity. The survivor phenotype also benefited

from increased predator resistance during subsequent predation assays. These results

demonstrate the utility of myxobacterial predator-prey models and provide insight into

prey resistances in response to predatory stress might contribute to the phenotypic

diversity and structure of bacterial communities.

Introduction

Abundant within soils and marine environments, predatory myxobacteria contribute to nutrient cycling within the microbial food web (1-4). Myxobacteria are prolific producers of antimicrobial specialized metabolites and display a cooperative, swarming predation strategy that can be readily reproduced and monitored within laboratory settings making them uniquely appropriate for assessment of predator-prey interactions (1, 3-5). Broadly considered generalist predators, myxobacteria capably predate a range of prey including both Gram-negative and Gram-positive bacteria as well as fungi (5, 6). Constitutive production of specialized metabolites and lytic proteins are often associated with this predatory range, and a variety of metabolites and enzymatic features have been reported to benefit predation (7-10).

However, relatively few examples of prey resistance to myxobacterial predation have been reported (11). Compared to features associated with bacterial strategies to avoid protozoan predators, prey avoidance of predatory myxobacteria remains underexplored (12-16). Examples of prey responses correlated with resistance to myxobacterial predation include *Escherichia coli* biofilm formation (17), *Bacillus subtilis* sporulation and production of bacillaene (18, 19), *Bacillus licheniformus* glycosylation of the predation-associated metabolite myxovirescin A (20), galactoglucan exopolysaccharide production and increased melanin production by *Sinorhizobium meliloti* (21, 22), and formaldehyde secretion by *Pseudomonas aeruginosa* (11). All of these features were discovered from predator-prey experiments utilizing the model myxobacterium *M. xanthus*.

We suspect the development of predator-prey pairings including myxobacteria other than *M. xanthus* might provide additional insight into prey resistance to myxobacterial

predation. For our predator-prey experiments, the soil dwelling myxobacterium

Cystobacter ferrugineus strain Cbfe23 was included due to a favorable growth profile and

capability to quickly consume *P. putida* during standard predation assays (23-25). Also

found within soils, root colonizing *P. putida* was chosen as prey due to an established

ability to resist protozoan grazers (14). Initial predator-prey experiments provided a *P.*

putida phenotype capable of avoiding *C. ferrugineus* predation. Herein we report the

generation and predator avoidance of a *P. putida* phenotype resistant to myxobacterial

predation using standard predator-prey experiments, differential gene expression data

comparing the survivor phenotype with predator-unexposed *P. putida*, and traits observed

to potentially contribute to predator avoidance.

Results

Selection of predation resistant *P. putida* phenotype

Utilizing predation assays where an inoculum of *C. ferrugineus* was introduced to the

edge of an established spot of *P. putida* (24), we considered swarming overtaking the *P.*

putida spot with no visible prey remaining to be an endpoint for assessment of remaining

viable prey cells using colony forming unit (CFU) assays. Typically, *C. ferrugineus*

swarmed predation sensitive *P. putida* with no visible prey after 3 days of co-cultivation

on nutrient restrictive WAT agar (Figure 1A and 1D). However, from 100 total assays 5

provided minimally consumed and readily observable *P. putida* remaining after 10-14

days (Figure 1B and 1D). Additionally, instead of frontal attack at the predator-prey

interface when exposed to predation sensitive *P. putida*, *C. ferrugineus* swarmed the

perimeter of the survivor phenotype (Supplemental Figure 1). Exponentially fewer viable

colonies of predation sensitive *P. putida* were observed from endpoint CFU assays (3-4

days for predation sensitive *P. putida*; 10-14 days for *P. putida* survivor phenotype)

supporting the assumption that swarming resistance correlates with prey survival (Figure

1D and 1E). Also apparent from CFU assays, colonies of the survivor phenotype were

consistently smaller when compared to predation sensitive *P. putida* (Figure 1C).

Predation resistance of the survivor phenotype was also observed from additional

predation assays with the optical density of prey inoculum doubled (OD_{600}=1.0 versus

OD_{600}=0.5). (Supplemental Figure 2). Stocks of each *P. putida* survivor observed from

predation assays were stored at -80°C as individual aliquots and utilized to determine

features associated with the survivor phenotype that might contribute to predator

avoidance.

Differential gene expression of predator-unexposed *P. putida* and survivor

phenotype

Utilizing RNA sequencing, we conducted comparative transcriptomic experiments

between the predator unexposed (control) *P. putida* and the survivor phenotype (n=3

biological replicates with biological replicates for the survivor phenotype defined as an

individually observed phenotype resistant to *C. ferrugineus* predation for ≥10 days). This

comparative analysis determined that 1,178 genes were down-regulated ≥4-fold and 122

genes were up-regulated ≥4-fold across the survivor phenotype replicates when

compared to control *P. putida* replicates (n=3; p≤0.05). Considering up-regulated genes

from the survivor phenotype with the highest fold change values, apparent differences in

overexpressed features specific to the survivor phenotype included genes involved in

mucoid conversion, siderophore production, and a variety of membrane-associated

proteins (Figure 2A). A 4 gene operon including a ferric uptake regulatory (*furA*)

associated gene (WP_016501343.1), a fumarate hydratase (WP_016501342.1), a

manganese superoxide dismutase (*sodA*) (WP_016501340.1), and a hypothetical protein

with a conserved DUF2753 domain (WP_06501341.1) was significantly up-regulated in

the survivor phenotype (Figure 2B and 2C). Expression of this operon from *P. aeruginosa*

in response to iron limitation leads to elevated alginate production, an exopolysaccharide

composed of mannuronic acid and guluronic acid monomers, and increased mucoidy (26,

27) and corroborates our observation of small colony variance from the survivor

phenotype. Interestingly, the alginate regulatory elements MucA (WP_016498171.1) and

MucB (WP_016498172.1) were both significantly down regulated in the survivor

phenotype (Figure 2B). The anti-sigma factor MucA and the associated binding partner

MucB inhibit mucoid conversion by sequestering the alternative sigma factor AlgU

required for alginate production in *P. aeruginosa* (28-31). Increased transcription from the

4 gene, *furA* operon combined with decreased transcription of MucA and MucB, indicates

that mucoid conversion might contribute to predator avoidance which parallels *P.*

aeruginosa predatory avoidance mechanisms when exposed to grazing amoeba (32).

Also associated with iron limitation, regulatory and biosynthetic genes contributing to

production of the siderophore pyoverdine were significantly up-regulated in the survivor

phenotype. These included PvdA an L-ornithine monooxygenase (WP_016499102.1)

and PvdH a diaminobutyrate-2-oxoketoglutarate transaminase (WP_016498669.1) both

involved in pyoverdine precursor biosynthesis, a regulatory transcription factor PvdS

(WP_016498655.1), a proximal MbtH-like protein (WP_016499090.1), and a core

nonribosomal peptide synthetase homologous to PvdL from *P. aeruginosa*

(NP_746359.1) (Supplemental Figure 3) (33-36). Also associated with iron limiting

conditions, a bacterioferritin-associated ferredoxin (WP_016488781.1) was upregulated

in the survivor phenotype.

A variety of membrane-associated proteins were also overexpressed in the survivor

phenotype including an osmotically inducible protein OsmY (WP_0016501632.1), an

outer membrane protein A (OmpA) family protein (WP_016501196.1), an FKBP-type

peptidyl-prolyl cis-trans isomerase (WP_0165010671), an aquaporin (WP_016498616.1),

and a DUF1282 domain-containing hypothetical protein from the Yip1 superfamily

(WP_016499858.1) (Figure 2A). While not a membrane associated feature, we also found

the overexpression of a formaldehyde dehydrogenase (WP_016497631.1) by the

survivor phenotype to be of note due to the recent observation that suggests

formaldehyde secretion to be a predation-resistance trait of *P. aeruginosa* (11). The

abundance of up-regulated genes that encode membrane-associated products in the

survivor phenotype including those responsible for increased mucoid production provides

insight into features involved in predation avoidance.

Numerous efflux and transport proteins were also among the most up-regulated genes

observed from the survivor phenotype (Figure 2D). Upregulated efflux genes included

inner (WP_016498130 and WP_016499477.1) and outer (WP_016498128.1) membrane

components highly homologous to the AcrAD-TolC-type multidrug resistance-nodulation-

division (RND) family of efflux pumps and the Mex RND efflux pumps known to facilitate

P. aeruginosa resistance to aminoglycosides and tetracyclines (37-39). Genes encoding

features homologous to multidrug efflux pumps contributing to virulence of human

pathogens were also up-regulated in the survivor phenotype including a P-type ATPase

(WP_016502100) and an additional RND/MmpL (Mycobacterial membrane protein

Large) protein (Figure 2D). Afforded these observed differences in gene expression, we

sought to determine if the survivor phenotype demonstrated antibiotic resistance,

increased mucoidy, and increased production of the siderophore pyoverdine which have

all been previously associated with predator avoidance strategies.

Antibiotic resistance of survivor phenotype

Provided the observation that numerous transport proteins were overexpressed in the

survivor phenotype (Figure 2D) including RND-type efflux pumps known to contribute to

P. aeruginosa antibiotic resistances, we were interested to determine if the survivor

phenotype exhibited antibiotic resistance. The survivor phenotype was capable of initially

forming colonies on LB agar supplemented with the antibiotics gentamicin (10 μg/ml),

kanamycin (50 μg/ml), and tetracycline (10 μg/ml), with no colonies from predation

sensitive *P. putida* observed on identical medias. Subsequent growth curve assays

comparing predation sensitive *P. putida* with the survivor phenotype when grown in LB

and LB supplemented with antibiotics (Figure 3A and 3B) confirmed that the survivor

phenotype was uniquely resistant to gentamicin, kanamycin, and tetracycline. These

results combined with the overexpression of efflux-associated proteins observed from the

survivor phenotype confirms that *C. ferrugineus* selection provides prey that benefit from

antibiotic resistance and suggests efflux-mediated antibiotic resistance to be involved in

P. putida response to predatory stress.

Increased production of alginate by survivor phenotype

Alginate-based formation of mucoid biofilms has previously been associated with *P.*

aeruginosa avoidance of the protozoan grazer *Rhynchomonas nasuta* (32, 40-42).

Utilizing an established carbazole assay as well as alginate antibody dot blots (43-45),

we sought to determine if the increased transcription of genes known to impact alginate

production observed from the survivor phenotype resulted in increased production of

alginate when compared to predator unexposed *P. putida*. Comparing supernatants from

predation sensitive *P. putida* with the survivor phenotype, we observed that the survivor

phenotype indeed produced more alginate when compared to *P. putida* not exposed to

C. ferrugineus (Figure 4).

Increased pyoverdine production by survivor phenotype

Significantly upregulated features associated with pyoverdine biosynthesis previously

noted from the survivor phenotype in our transcriptomic data suggested that increased

pyoverdine production might be an observable survivor trait. Fluorescence of extracellular

pyoverdine was quantified from clarified medias of predator unexposed *P. putida* and the

survivor phenotype using methodology established by Imperi *et al* (46, 47). Indicative of

increased pyoverdine secretion, extracellular fractions from the survivor phenotype

exhibited a 3-5 fold increase in fluorescence when compared to extracellular fractions

from predator unexposed *P. putida* (Figure 5). Although the genome of the *P. putida* type

strain includes a pyoverdine biosynthetic gene cluster, no structurally elucidated

pyoverdines have been characterized from the *P. putida* type strain, and these results

merely suggest the presence of fluorescent pyoverdine-like metabolites potentially

produced by the strain. Regardless, these results corroborate overexpression of

pyoverdine biosynthetic pathway components observed in our transcriptomic analysis

and suggest that pyoverdine production may contribute to the predator avoidance trait of

the survivor phenotype.

Increased production of phenazine-1-carboxylic acid by survivor phenotype

Untargeted mass spectrometry was used to assess additional differences in detectable

quantities of *P. putida* metabolites comparing organic phase extracts from predator

unexposed *P. putida* and the survivor phenotype. Initial analysis of resulting data using

Global Natural Product Social Molecular Networking (GNPS) (48) resulted in a library hit

for phenazine-1-carboxylic acid ([M+H]=225.066) that was only observed in extracts from

the survivor phenotype (Figure 6A). Subsequent comparison utilizing XCMS-MRM (49)

provided the statistical significance (n=3; p=0.0509) in detected phenazine-1-carboxylic

acid between extracts from predator unexposed *P. putida* and the survivor phenotype

(Figure 6B). Interestingly, phenazine production in *P. aeruginosa* contributes to redox

maintenance during biofilm formation and promotes efflux-based antibiotic resistance

(50-53). Despite no significant change in expression of putative phenazine biosynthesis

genes *phzD* (SUD74861.1) and *phzF* (SUD71919.1) from our comparative transcriptomic

data (54-57), we suspect that higher quantities of phenazine-1-carboxylic acid detected

in extracts from the survivor phenotype may contribute to the increased transcription of

efflux pump components and antibiotic resistance observed from the survivor phenotype.

However, associations between phenazine production and efflux-mediated resistance

have only been reported from *P. aeruginosa*, and any parallels potentially observed in our

data require further investigation. Also of note, phenazine production has been observed

to protect *Pseudomonas aureofaciens* from predatory myxobacteria (58). These results

suggest that the survivor phenotype produces significantly greater quantities of

phenazine-1-carboxylic acid when compared to predator-unexposed *P. putida*.

Predation resistance of survivor phenotype in subsequent predation assays

Additional subsequent predation assays were done to assess the predator avoidance trait

of the *P. putida* survivor phenotype with *P. putida* not exposed to *C. ferrugineus* included

as a control. These predation assays included an assay as previously described with

predator cells directly introduced to the edge of established prey, a similar slightly

modified assay with predator introduced at a distance of 5-7 mm from an established spot

of *P. putida*, and a lawn culture assay (23) monitoring predator swarming diameters when

placed at the center of prey embedded in agar. For direct contact assays, time to complete

swarming was recorded. As observed in our prior predator-prey experiments, compared

to control *P. putida* the survivor demonstrated significantly increased predator avoidance

(Figure 7A). The survivor phenotype also demonstrated resistance to swarming

compared to predation sensitive *P. putida* during identical direct-contact predation assays

conducted on nutrient-rich VY/2 agar (Supplemental Figure 4) For indirect contact

predation assays, time to predator-prey contact and time to complete swarming were

recorded. Time to predator-prey contact was between 1-2 days for all samples, and as

observed in the directed contact predation assays the survivor phenotype demonstrated

increased predator avoidance when compared to control *P. putida* (Figure 7B and 7C).

However, no differences between swarming diameters of *C. ferrugineus* grown on lawns

of control *P. putida* and the *P. putida* survivor phenotype were observed (Figure 7D).

These data suggest the *P. putida* survivor phenotype benefits from a predator avoidance

trait when subjected to 2 out of 3 subsequent predation assays.

Discussion

Building upon previously reported resistance strategies to avoid predatory myxobacteria

discovered from predator-prey experiments (11, 17-22), predation experiments including

C. ferrugineus and *P. putida* provided a prey phenotype significantly resistant to

predation. Traits associated with prey resiliency during bacterial predator-prey

interactions remain underexplored despite growing evidence that predatory stress

contributes to the development of phenotypic traits associated with persistent clinical

pathogens (12, 16, 59). As an example, clinical isolates of persistent *P. aeruginosa* from

cystic fibrosis patients exhibit increased mucoidy and small colony variation as well as

efflux-mediated antibiotic resistance (60-62). Other human pathogens that have

demonstrated overlapping factors involved in both virulence and anti-predation strategies

include *Legionella pneumophila*, *Escherichia coli*, *Vibrio cholerae*, *Campylobacter jejuni*,

Listeria monocytogenes, *Mycobacterium leprae*, and *Yersinia enterocolitica* (12, 16).

Overlapping factors thus far observed to be associated with both virulence and predator

avoidance include biofilm formation (15, 32, 42), quorum signal (63) and siderophore (64)

small molecule biosynthesis, toxin production (65-67), expression of transport proteins

such as type III and type VI secretion systems (68, 69) and antibiotic resistance-

associated efflux pumps (70, 71). While the impact of predatory selection on virulence

factors have primarily been reported from predator-prey experiments with grazing

amoeba, efforts to determine the influence of generalist predators on microbial

communities have suggested that bacterial predators also drive adaptation of prey traits

that overlap with virulence in bacterial pathogens (5). Recently, Nair *et al.* reported that

during predator-prey interactions with *E. coli* prey, the predatory myxobacterium

Myxococcus xanthus selects for 2 traits associated with virulence, mucoidy and increased

expression of the outer membrane protease OmpT (5). Compared to features associated

with bacterial strategies to avoid protozoan predators, prey avoidance of myxobacterial

predators remains underexplored. Although predator-prey experiments that utilize

myxobacteria with various prey such as *Bacillus subtilis* (9, 18, 19), *E. coli* (5, 7),

Sinorhizobium melilotti (21, 22), *P. aeruginosa* (11), and *Streptomyces coelicolor* (72)

have been reported, all myxobacterial resistances have been discovered from predator-

prey experiments that included the model myxobacterium *M. xanthus*. Utilizing an

alternative myxobacterium *C. ferrugineus* provides the opportunity to further compare

predatory selection and adaptive prey avoidance strategies.

Comparative transcriptomics provided key features up-regulated in the *P. putida* survivor

phenotype that overlap with traits associated with *P. aeruginosa* predator avoidance and

pathogenicity. Of these, 3 putative RND family transporter components with high

homology to the MexAB-OprM and MexCD-OprM efflux systems known to mediate *P.*

aeruginosa antibiotic resistance were up-regulated in the survivor phenotype (38, 39).

Growth curve analysis determined that the *P. putida* survivor phenotype was indeed

resistant to the aminoglycosides kanamycin and gentamicin as well as tetracycline;

susceptibility to all 3 of these antibiotics was observed from predator unexposed *P. putida*.

Also of note, the multidrug efflux pump CmeABC serves as a survival mechanism for *C.*

jejuni when engulfed by the predatory amoeba *Acanthamoeba polyphaga* (71).

Several features associated with mucoid conversion were also up-regulated in the

survivor phenotype including a 4 gene operon responsible for elevated alginate

production in *P. aeruginosa* and the regulatory transcription factors MucA and MucB (26-

28, 30). Adaptive mucoid phenotypes have been previously observed in response to

myxobacterial predation (5), amoebal grazing (40), macrophages (73), and lytic phages

(74). Increased alginate production by the survivor phenotype was confirmed using

established carbazole assays and anti-alginate dot blots. Alginate-based mucoid

conversion and biofilm formation have previously been associated with *P. aeruginosa*

avoidance of grazing amoeba and could also contribute to the small colony variation

observed from our survivor phenotype (32, 40, 41). These results and the observed

alternative predatory pattern of *C. ferrugineus* during predator-prey interactions with the

survivor phenotype parallel previous observation of mucoid *S. meliloti* phenotypes

avoiding *M. xanthus* predation (22). Additionally, upregulation of an OmpA family protein

observed in the survivor phenotype might also correlate with mucoid conversion as *P.*

aeruginosa phenotypes that upregulate expression of OmpA also overproduce alginate

(75). These results provide clear overlap between features contributing to avoidance of

myxobacterial and amoebal predation and also afford additional evidence that the

coincidental selection of prey virulence factors reported from amoebal predatory stress

(12) might also be observed from predator-prey interactions involving myxobacteria. Both

efflux-mediated antibiotic resistance and mucoid conversion have also been attributed to

virulence of clinical isolates of *P. aeruginosa* (29, 59, 60, 76).

Additionally, our data provided key areas for follow-up investigations to assess the

chemical ecology of the predator-prey interface and its contribution to predator avoidance

such as the increased production of pyoverdines and phenazine-1-carboxylic acid

detected from the survivor phenotype. For example, up-regulation of the pyoverdine

biosynthetic pathway and a bacterioferritin-associated ferredoxin as well as increased

pyoverdine in extracellular extracts observed from the survivor phenotype suggests an

iron-depleted predator-prey environment similar to the previously reported interaction

between *M. xanthus* and *Streptomyces coelicolor* (72). Enhanced production of the

primary myxobacterial siderophore myxochelin by *M. xanthus* contributed to an iron-

restrictive environment and triggered production of the pigment actinorhodin from *S.*

coelicolor. Further investigation focused on the chemical ecology of predator-prey

interfaces might also shed light on the various membrane features observed to be up-

regulated in the survivor phenotype.

The survivor phenotype was also assessed with subsequent predation assays to

determine the resiliency of the predator avoidance trait. While comparative transcriptomic

data provided insight into key up-regulated features potentially involved in prey survival,

down-regulation of 1,178 genes in the survivor phenotype when compared to predator

unexposed *P. putida* suggests potentially diminished survivor fitness. Despite this, the

survivor phenotype maintained the predator avoidance trait in 2 out of 3 subsequent

predation assays. While it is unclear why no significant predator avoidance was observed

from lawn culture predation assays, we suspect the introduction of predator to the center

of prey lawns as opposed to consistent introduction to the peripheral interface of prey

potentially reduce any predator avoidance mechanisms. Predator avoidance observed

from direct and indirect contact predation assays and the heterogeneity associated with

differences between the growth and metabolism of peripheral cells and central cells of

bacterial colonies supports this suspicion (77, 78). Perhaps belaboring, we also suggest

that these results again highlight the importance of chemical exchange at the predator-

prey interface. We anticipate that *P. putida* phenotypes resistant to myxobacterial

predation continually subjected to such predatory selection will afford adapted

phenotypes for further scrutiny of any resulting mutations and genotypes. Ultimately,

these results demonstrate the utility of predator-prey experiments using lesser studied

myxobacteria and provide insight into how predatory stress might contribute to adaptive

phenotypic diversity of bacterial communities.

Materials and Methods

Bacterial Strains and Cultivation

The myxobacterium *Cystobacter ferrugineus* strain Cbfe23, DSM 52764 was employed

as the predator. *Pseudomonas putida* type strain, ATCC 12633, and the discussed *P.*

putida survivor phenotype resulted from predation assays with this *P. putida* type strain

were included as prey. *C. ferrugineus* was grown on VY/2 solid (1.4% w/v agar, 0.1% w/v

$CaCl_2 * 2\ H_2O$, 0.5% w/v Baker's yeast, 500 µM vitamin B12) media for 5-7 days. Luria-

Bertani (LB) solid (1.5% agar) and liquid media were utilized for the cultivation of *P.*

putida. Predation experiments were performed on WAT agar (1.5% w/v agar, 0.1% w/v

$CaCl_2$, 20 mM HEPES) plates. All bacteria were grown at 30 °C.

Predation assays

Prey spot assay was performed as described by Seccareccia et al. (24). Briefly, *P. putida*

prey was grown in LB liquid media on a rotary shaker (150 rpm) at 30 °C for 16-18 h.

Next, the bacterial cultures were sedimented by centrifugation at 4,000 g for 15 min, and

the sedimented cells were washed and resuspended in TM buffer (50 mM Tris, pH 7.8,

10 mM $MgSO_4$) to an OD_{600} of 0.5. Then the WAT agar plates were spotted with 150 µl

of the freshly prepared bacterial suspensions, and the spots were dried in a laminar flow

hood. Finally, *C. ferrugineus* was inoculated at the edge of the prey spot. The assay plates

were incubated at 30 °C for 14 days or until the full visible swarming of *C. ferrugineus* on

the prey spot. The assay was performed with 100 replicates. Any prey spot with visible

biomass of *P. putida* after day 14 was considered as survivor phenotype of *P. putida*, and

five observed *P. putida* survivor stocks were acquired that remained visible after day 14

of the assay.

For subsequent predation assays, *P. putida* predation sensitive (control) and the survivor

phenotypes were employed as prey. The direct spot assay was performed as mentioned

above. Statistical significance of recorded time to predation reported in was calculated

using an unpaired t test with Welch's correction in Prism 7.0d. The subsequent directed

contact predation assay was performed on VY/2 (1.4% w/v agar, 0.1% w/v $CaCl_2$ * 2 H_2O,

0.5% w/v Baker's yeast, 500 μM vitamin B12) media as well, considered as nutrient-rich

media. For indirect contact spot assay, *C. ferrugineus* was inoculated at a distance of 5-

7 mm from the established *P. putida* control and *P. putida* survivor prey spots. Next, time

C. ferrugineus took to reach the prey spots and the time of its swarming over the spots

was recorded. Indirect predation assays were performed with 8 replicates per phenotype.

Statistical significance of recorded time to predation reported in days and recorded time

to initial contact between *C. ferrugineus* and prey reported in days was calculated using

an unpaired t test with Welch's correction in Prism 7.0d.

Swarming predation assays were based on established lawn culture methods (23). In

brief, *P. putida* prey phenotypes were grown in LB liquid media for 16-18 h at 30 °C with

shaking at 150 rpm. The cultures were then subjected to centrifugation at 4,000 g for 15

min, and the sedimented cells were washed and resuspended in TM buffer (50 mM Tris,

pH 7.8, 10 mM $MgSO_4$) to an OD_{600} 0.5. Next, 1 ml of the cell suspension was spread

onto a 14 cm diameter WAT agar plate and dried to form a uniform prey lawn. Finally, *C.*

ferrugineus previously cultured on VY/2 agar media was spotted onto the center of the

prey lawn. The assay plates were incubated at 30 °C, and the expansion of the *C.*

ferrugineus swarm was measured on day 4. Lawn culture predation assays were

performed with 8 replicates per phenotype with replicates from each survivor stock

included. All 3 predation assays were performed simultaneously for all replicates.

Statistical significance of *C. ferrugineus* swarm diameters reported in millimeters on day

4 was calculated using an unpaired t test with Welch's correction in Prism 7.0d

CFU assays.

The colony forming units (CFU) were calculated as described by DePas *et al.* (17). Briefly,

after the full swarming of *C. ferrugineus* over the prey spot, the agar slab under the prey

spot was excised and suspended in 2 mL phosphate buffer saline (PBS). Next, a dilution

series from this solution was inoculated on the LB plates, and the plates were incubated

at 30 °C for 18 hr. Then, CFU/mL was calculated from the viable colonies that appeared

on the LB plates. Finally, the CFU/mL of the survivor was compared with CFU/mL of the

unexposed *P. putida,* considered as a control. Statistical significance of all CFU assays

was calculated using an unpaired t test with Welch's correction in Prism 7.0d.

RNA sequencing

Triplicate samples of *P. putida* control and *P. putida* survivor phenotype grown from

stocks in LB at 37°C for 18 hr and pelleted via centrifugation were stored in RNA*later*

solution for sequencing. The DNA from *P. putida* type strain cells (200 μl) was extracted

using MagAttract HMW DNA Kit (Qiagen). The DNA was eluted in 100 uL AE buffer. The

concentration of DNA was evaluated (Supplementary Table 1) using the Qubit® dsDNA

HS Assay Kit (Life Technologies). The library was prepared using Nextera DNA Flex

library preparation kit (Illumina) following the manufacturer's user guide. 50 ng DNA was

used to prepare the library. The sample underwent the simultaneous fragmentation and

addition of adapter sequences. These adapters are utilized during a limited-cycle (6

cycles) PCR in which unique index was added to the sample. Following the library

preparation, the final concentration of the library (Supplementary Table 1) was measured

using the Qubit® dsDNA HS Assay Kit (Life Technologies), and the average library size

(Supplementary Table 1) was determined using the Agilent 2100 Bioanalyzer (Agilent

Technologies). The library was diluted (to 6.5 pM) and sequenced paired end for 500

cycles using the MiSeq system (Illumina).

Total RNA was isolated from triplicate samples of the *P. putida* control, the *P. putida*

survivor phenotype from stocks, and the kanamycin resistant phenotype using the

RNeasy PowerSoil Total RNA Kit (Qiagen) following the manufacturer's instructions. 400

μl cell sample was used for extractions. The concentration of total RNA was determined

(Supplementary Table 2) using the Qubit® RNA Assay Kit (Life Technologies). For rRNA

depletion, first, 1000 ng of total RNA was used to remove the DNA contamination using

Baseline-ZERO™ DNase (Epicentre) following the manufacturer's instructions followed

by purification using the RNA Clean & Concentrator-5 columns (Zymo Research). DNA

free RNA samples were used for rRNA removal by using RiboMinus™ rRNA Removal Kit

(Bacteria; Thermo Fisher Scientific) and final purification was performed using the RNA

Clean & Concentrator-5 columns (Zymo Research). rRNA depleted samples were used

for library preparation using the KAPA mRNA HyperPrep Kits (Roche) by following the

manufacturer's instructions. Following the library preparation, the final concentrations of

all libraries (Supplementary Table 2) were measured using the Qubit® dsDNA HS Assay

Kit (Life Technologies), and the average library size was determined using the Agilent

2100 Bioanalyzer (Agilent Technologies). The libraries were then pooled in equimolar

ratios of 0.6 nM, and sequenced paired end for 300 cycles using the NovaSeq 6000

system (Illumina). Differential expression between the resulting transcriptomes was

calculated from pair-wise analysis of trimmed mean of M-values (TMM) normalized read

counts (79) including fold change, counts-per-million (CPM), and associated p-values

using the R-package, edgeR (80). Differential expression data with p-values ≤ 0.05 were

considered statistically significant. Genome and RNA sequencing was conducted by MR

DNA (Molecular Research LP).   All fold change, CPM, and associated p-values resulting

from differential expression experiments as described included as supplementary data

files TABLE 1 (*P. putida* survivor compared with *P. putida* control). All raw RNAseq data

is publicly available at the NCBI Sequence Read Archive (PRJNA577468).

Antibiotic Susceptibility Assays.

The standard growth curve assay was employed to determine the sensitivity of the

unexposed *P. putida* and the survivor phenotype towards kanamycin (50 mg/mL),

gentamicin (10 mg/mL), and tetracycline (10 mg/mL) antibiotics. The Lauria Bertani (LB)

broth without any supplemented antibiotics was used as a control. The test was

performed in a 96-well microtiter plate. In a 96-well plate, each well having 200 μL of LB

or LB supplemented with an antibiotic was inoculated with a colony of *P. putida.* Next, the

absorbance at OD_{620} was recorded using a plate reader every 30 min for 10 hr to examine

the growth dynamics of each phenotype. The test was performed in 12 replicates per

condition for both phenotypes. Recorded OD_{620} values were plotted using Prism version

7.0d with error bars depicting standard deviation of replicate data.

Carbazole Assays.

The alginate was isolated according to Jones *et al*. (81). In brief, both *P. putida* phenotype

unexposed to the predatory stress and the survivor phenotype were grown overnight in 2

ml LB media and adjusted to an OD_{600} of 1.00. Following centrifugation, 1 ml of

supernatant was treated with 2% cetyl pyridinium chloride to precipitate the alginate. The

precipitated alginate was further collected by centrifugation. It was resuspended in 1 M

NaCl and then re-precipitated in cold isopropanol. Precipitated alginate was suspended

in 150 μl 0.9% (w/v) saline solution. Following established protocols (43, 44), a 50 µL

aliquot of the isolated alginate and a dilution series of standard alginic acid (Sigma) were

mixed with 200 µL of a solution of 25 mM sodium tetraborate in sulfuric acid and added

in a 96-well plate. Next, the plate was heated for 10 min at 100 °C in an oven. After cooling

at room temperature for 15 min, 50 μl of a 0.125% carbazole solution in absolute ethanol

was added. Then, the plate was re-heated at 100 °C for 10 min in an oven and cooled

down at room temperature for 15 min. Finally, the plate was read in a CLARIOstar

microplate reader (BMG Labtech Inc., Cary, NC, USA) at a wavelength of 550 nm.

Quantities of alginate per sample (μg/ml) were calculated from the resulting standard

curve of purchase alginic acid, and statistical significance was calculated using an

unpaired t test with Welch's correction in Prism 7.0d.

Anti-alginate Dot Plots.

Alginate dot blots were generated using methodology from Lorenz *et al.* with slight

modifications (45). Alginate was isolated as previously described, suspended in 150 μl

0.9% (w/v) saline solution, and 2 μl aliquots were pipetted onto a nitrocellulose

membrane. After air-drying for 30 min, Tris buffered saline-Tween 20 (TBST) with 5%

bovine serum albumin (BSA) was used for membrane blocking for 1 h. Membrane was

then washed 3x for 10 min with TBST and subsequently incubated with the alginate

antibody (Sigma, monoclonal, anti-mouse) overnight at 4°C. The following day, the

membrane was washed 3x for 10 min with TBST, incubated with IRDye 800CW Goat

anti-mouse (Sigma) for 30 min, rinsed 2x with TBST, and imaged on a LI-COR Odyssey.

Average relative intensities for imaged blots comparing isolated alginate from predator

unexposed *P. putida* with alginate isolated from the survivor phenotype (n=10) are

reported with statistical significance was calculated using an unpaired t test with Welch's

correction in Prism 7.0d.

Assessment of Extracellular Pyoverdine

Extracellular pyoverdine fluorescence was detected from supernatants following

methodologies Imperi *et al.* and Barrientos-Moreno *et al.* (46, 47). Clarified supernatants

from overnight cultivation of 200 μl LB cultures of predator unexposed *P. putida* and the

survivor phenotype adjusted to an OD_{600} of 0.07 were generated by centrifugation (10,000

rpm, 15m). Fluorescence was recorded at 455 nm upon excitation at 400 nm using a

CLARIOstar microplate reader (BMG Labtech Inc., Cary, NC, USA). Sterile LB was used

as a negative control to assess background fluorescence. Average relative fluorescence

for supernatants from predator unexposed *P. putida* and survivor phenotype cultures

(n=12) are reported with statistical significance was calculated using an unpaired t test

with Welch's correction in Prism 7.0d.

Metabolite Extraction and Analysis.

For metabolomic analysis, the survivor phenotype and predator unexposed *P. putida*

were cultivated on VY/2 agar plates as described in the predation assay. VY/2 agar plates

with TM buffer were used as a negative control. The plates were incubated at 30°C for 14

days. After the incubation period, agar was chopped and extracted with ethyl acetate

(EtOAC). The EtOAc extracts were dried in vacuo to produce crude extracts for LC-

MS/MS analysis. The crude extracts from each condition were generated in triplicate and

analyzed as previously described (82). LC-MS/MS generated data was converted to

.mzML files using MS-Convert, and the GNPS platform (48) was utilized as a dereplication

tool to look for the library hits within publicly available natural products libraries. For the

statistical analysis, using XCMSonline (49), multigroup analysis with HPLC orbitrap

default settings was employed. MZmine 2.53 was used to generate extracted ion

chromatograms (83-85).

Author Contributions.

SA performed the experiments. DCS supervised the project. SA and DCS conceived the

project, designed the experiments, and wrote the manuscript.

Acknowledgements.

The authors appreciate funding and support from the National Institute of Allergy and

Infectious Diseases (R15AI137996) and the National Institute of General Medical

Sciences of the National Institutes of Health (P20GM130460). The authors appreciate

Scot E. Dowd at MR DNA, Molecular Research LP for assistance throughout sequencing

services provided, Dr. Sandeep Misra Manager of the Analytical and Biophysical

Chemistry Core associated with the Glycoscience Center of Research Excellence

(GlyCORE; P20GM130460) for LC-MS/MS analysis, and Dr. Nicole Ashpole for her

assistance with anti-alginate dot blot generation and imaging.

References.

1.   Albataineh H, Stevens DC. 2018. Marine Myxobacteria: A Few Good Halophiles. Mar Drugs
16.

2.   Findlay BL. 2016. The Chemical Ecology of Predatory Soil Bacteria. ACS Chem Biol 11:1502-
10.
3.   Munoz-Dorado J, Marcos-Torres FJ, Garcia-Bravo E, Moraleda-Munoz A, Perez J. 2016.
Myxobacteria: Moving, Killing, Feeding, and Surviving Together. Front Microbiol 7:781.
4.   Mohr KI. 2018. Diversity of Myxobacteria-We Only See the Tip of the Iceberg.
Microorganisms 6.
5.   Nair RR, Vasse M, Wielgoss S, Sun L, Yu YN, Velicer GJ. 2019. Bacterial predator-prey
coevolution accelerates genome evolution and selects on virulence-associated prey
defences. Nat Commun 10:4301.
6.   Perez J, Moraleda-Munoz A, Marcos-Torres FJ, Munoz-Dorado J. 2016. Bacterial
predation: 75 years and counting! Environ Microbiol 18:766-79.
7.   Livingstone PG, Millard AD, Swain MT, Whitworth DE. 2018. Transcriptional changes when
Myxococcus xanthus preys on Escherichia coli suggest myxobacterial predators are
constitutively toxic but regulate their feeding. Microb Genom 4.
8.   Ellis BM, Fischer CN, Martin LB, Bachmann BO, McLean JA. 2019. Spatiochemically
Profiling Microbial Interactions with Membrane Scaffolded Desorption Electrospray
Ionization-Ion Mobility-Imaging Mass Spectrometry and Unsupervised Segmentation.
Anal Chem 91:13703-13711.
9.   Muller S, Strack SN, Ryan SE, Shawgo M, Walling A, Harris S, Chambers C, Boddicker J,
Kirby JR. 2016. Identification of Functions Affecting Predator-Prey Interactions between
Myxococcus xanthus and Bacillus subtilis. J Bacteriol 198:3335-3344.
10.   Xiao Y, Wei X, Ebright R, Wall D. 2011. Antibiotic production by myxobacteria plays a role
in predation. J Bacteriol 193:4626-33.
11.   Sutton D, Livingstone PG, Furness E, Swain MT, Whitworth DE. 2019. Genome-Wide
Identification of Myxobacterial Predation Genes and Demonstration of Formaldehyde
Secretion as a Potentially Predation-Resistant Trait of Pseudomonas aeruginosa. Front
Microbiol 10:2650.
12.   Erken M, Lutz C, McDougald D. 2013. The rise of pathogens: predation as a factor driving
the evolution of human pathogens in the environment. Microb Ecol 65:860-8.
13.   Justice SS, Hunstad DA, Cegelski L, Hultgren SJ. 2008. Morphological plasticity as a
bacterial survival strategy. Nat Rev Microbiol 6:162-8.
14.   Seiler C, van Velzen E, Neu TR, Gaedke U, Berendonk TU, Weitere M. 2017. Grazing
resistance of bacterial biofilms: a matter of predators' feeding trait. FEMS Microbiol Ecol
93.
15.   Weitere M, Bergfeld T, Rice SA, Matz C, Kjelleberg S. 2005. Grazing resistance of
Pseudomonas aeruginosa biofilms depends on type of protective mechanism,
developmental stage and protozoan feeding mode. Environ Microbiol 7:1593-601.
16.   Sun S, Noorian P, McDougald D. 2018. Dual Role of Mechanisms Involved in Resistance to
Predation by Protozoa and Virulence to Humans. Front Microbiol 9:1017.
17.   DePas WH, Syed AK, Sifuentes M, Lee JS, Warshaw D, Saggar V, Csankovszki G, Boles BR,
Chapman MR. 2014. Biofilm formation protects Escherichia coli against killing by
Caenorhabditis elegans and Myxococcus xanthus. Appl Environ Microbiol 80:7079-87.

18.   Muller S, Strack SN, Hoefler BC, Straight PD, Kearns DB, Kirby JR. 2014. Bacillaene and
560   sporulation protect Bacillus subtilis from predation by Myxococcus xanthus. Appl Environ
561   Microbiol 80:5603-10.
562   19.   Muller S, Strack SN, Ryan SE, Kearns DB, Kirby JR. 2015. Predation by Myxococcus xanthus
563   induces Bacillus subtilis to form spore-filled megastructures. Appl Environ Microbiol
564   81:203-10.
565   20.   Wang C, Liu X, Zhang P, Wang Y, Li Z, Li X, Wang R, Shang Z, Yan J, He H, Wang J, Hu W, Li
566   Y. 2019. Bacillus licheniformis escapes from Myxococcus xanthus predation by
567   deactivating myxovirescin A through enzymatic glucosylation. Environ Microbiol 21:4755-
568   4772.
569   21.   Contreras-Moreno FJ, Munoz-Dorado J, Garcia-Tomsig NI, Martinez-Navajas G, Perez J,
570   Moraleda-Munoz A. 2020. Copper and Melanin Play a Role in Myxococcus xanthus
571   Predation on Sinorhizobium meliloti. Front Microbiol 11:94.
572   22.   Perez J, Jimenez-Zurdo JI, Martinez-Abarca F, Millan V, Shimkets LJ, Munoz-Dorado J.
573   2014. Rhizobial galactoglucan determines the predatory pattern of Myxococcus xanthus
574   and protects Sinorhizobium meliloti from predation. Environ Microbiol 16:2341-50.
575   23.   Livingstone PG, Morphew RM, Whitworth DE. 2017. Myxobacteria Are Able to Prey
576   Broadly upon Clinically-Relevant Pathogens, Exhibiting a Prey Range Which Cannot Be
577   Explained by Phylogeny. Front Microbiol 8:1593.
578   24.   Seccareccia I, Kost C, Nett M. 2015. Quantitative Analysis of Lysobacter Predation. Appl
579   Environ Microbiol 81:7098-105.
580   25.   Akbar S, Dowd SE, Stevens DC. 2017. Draft Genome Sequence of Cystobacter ferrugineus
581   Strain Cbfe23. Genome Announc 5.
582   26.   Hassett DJ, Howell ML, Ochsner UA, Vasil ML, Johnson Z, Dean GE. 1997. An operon
583   containing fumC and sodA encoding fumarase C and manganese superoxide dismutase is
584   controlled by the ferric uptake regulator in Pseudomonas aeruginosa: fur mutants
585   produce elevated alginate levels. J Bacteriol 179:1452-9.
586   27.   Hassett DJ, Howell ML, Sokol PA, Vasil ML, Dean GE. 1997. Fumarase C activity is elevated
587   in response to iron deprivation and in mucoid, alginate-producing Pseudomonas
588   aeruginosa: cloning and characterization of fumC and purification of native fumC. J
589   Bacteriol 179:1442-51.
590   28.   Damron FH, Goldberg JB. 2012. Proteolytic regulation of alginate overproduction in
591   Pseudomonas aeruginosa. Mol Microbiol 84:595-607.
592   29.   Deretic V, Martin DW, Schurr MJ, Mudd MH, Hibler NS, Curcic R, Boucher JC. 1993.
593   Conversion to mucoidy in Pseudomonas aeruginosa. Biotechnology (N Y) 11:1133-6.
594   30.   Li S, Lou X, Xu Y, Teng X, Liu R, Zhang Q, Wu W, Wang Y, Bartlam M. 2019. Structural basis
595   for the recognition of MucA by MucB and AlgU in Pseudomonas aeruginosa. FEBS J
596   286:4982-4994.
597   31.   Martin DW, Schurr MJ, Mudd MH, Deretic V. 1993. Differentiation of Pseudomonas
598   aeruginosa into the alginate-producing form: inactivation of mucB causes conversion to
599   mucoidy. Mol Microbiol 9:497-506.
600   32.   Matz C, Bergfeld T, Rice SA, Kjelleberg S. 2004. Microcolonies, quorum sensing and
601   cytotoxicity determine the survival of Pseudomonas aeruginosa biofilms exposed to
602   protozoan grazing. Environ Microbiol 6:218-26.

33. Drake EJ, Cao J, Qu J, Shah MB, Straubinger RM, Gulick AM. 2007. The 1.8 A crystal
structure of PA2412, an MbtH-like protein from the pyoverdine cluster of Pseudomonas
aeruginosa. J Biol Chem 282:20425-34.
34. Felnagle EA, Barkei JJ, Park H, Podevels AM, McMahon MD, Drott DW, Thomas MG. 2010.
MbtH-like proteins as integral components of bacterial nonribosomal peptide
synthetases. Biochemistry 49:8815-7.
35. Parker DL, Lee SW, Geszvain K, Davis RE, Gruffaz C, Meyer JM, Torpey JW, Tebo BM. 2014.
Pyoverdine synthesis by the Mn(II)-oxidizing bacterium Pseudomonas putida GB-1. Front
Microbiol 5:202.
36. Ringel MT, Bruser T. 2018. The biosynthesis of pyoverdines. Microb Cell 5:424-437.
37. Garneau-Tsodikova S, Labby KJ. 2016. Mechanisms of Resistance to Aminoglycoside
Antibiotics: Overview and Perspectives. Medchemcomm 7:11-27.
38. Li XZ, Plesiat P, Nikaido H. 2015. The challenge of efflux-mediated antibiotic resistance in
Gram-negative bacteria. Clin Microbiol Rev 28:337-418.
39. Aeschlimann JR. 2003. The role of multidrug efflux pumps in the antibiotic resistance of
Pseudomonas aeruginosa and other gram-negative bacteria. Insights from the Society of
Infectious Diseases Pharmacists. Pharmacotherapy 23:916-24.
40. Matz C, Deines P, Jurgens K. 2002. Phenotypic variation in Pseudomonas sp. CM10
determines microcolony formation and survival under protozoan grazing. FEMS Microbiol
Ecol 39:57-65.
41. Matz C, Kjelleberg S. 2005. Off the hook--how bacteria survive protozoan grazing. Trends
Microbiol 13:302-7.
42. Matz C, McDougald D, Moreno AM, Yung PY, Yildiz FH, Kjelleberg S. 2005. Biofilm
formation and phenotypic variation enhance predation-driven persistence of Vibrio
cholerae. Proc Natl Acad Sci U S A 102:16819-24.
43. Al Ahmar R, Kirby BD, Yu HD. 2020. Culture of Small Colony Variant of Pseudomonas
aeruginosa and Quantitation of its Alginate. J Vis Exp doi:10.3791/60466.
44. Knutson CA, Jeanes A. 1968. A new modification of the carbazole analysis: application to
heteropolysaccharides. Anal Biochem 24:470-81.
45. Lorenz C, Dougherty TJ, Lory S. 2020. Transcriptional Responses of Pseudomonas
aeruginosa to Inhibition of Lipoprotein Transport by a Small Molecule Inhibitor. J Bacteriol
202.
46. Barrientos-Moreno L, Molina-Henares MA, Pastor-Garcia M, Ramos-Gonzalez MI,
Espinosa-Urgel M. 2019. Arginine Biosynthesis Modulates Pyoverdine Production and
Release in Pseudomonas putida as Part of the Mechanism of Adaptation to Oxidative
Stress. J Bacteriol 201.
47. Imperi F, Tiburzi F, Visca P. 2009. Molecular basis of pyoverdine siderophore recycling in
Pseudomonas aeruginosa. Proc Natl Acad Sci U S A 106:20440-5.
48. Wang M, Carver JJ, Phelan VV, Sanchez LM, Garg N, Peng Y, Nguyen DD, Watrous J,
Kapono CA, Luzzatto-Knaan T, Porto C, Bouslimani A, Melnik AV, Meehan MJ, Liu WT,
Crusemann M, Boudreau PD, Esquenazi E, Sandoval-Calderon M, Kersten RD, Pace LA,
Quinn RA, Duncan KR, Hsu CC, Floros DJ, Gavilan RG, Kleigrewe K, Northen T, Dutton RJ,
Parrot D, Carlson EE, Aigle B, Michelsen CF, Jelsbak L, Sohlenkamp C, Pevzner P, Edlund A,
McLean J, Piel J, Murphy BT, Gerwick L, Liaw CC, Yang YL, Humpf HU, Maansson M, Keyzers

RA, Sims AC, Johnson AR, Sidebottom AM, Sedio BE, et al. 2016. Sharing and community
curation of mass spectrometry data with Global Natural Products Social Molecular
Networking. Nat Biotechnol 34:828-837.

49. Domingo-Almenara X, Montenegro-Burke JR, Ivanisevic J, Thomas A, Sidibe J, Teav T,
Guijas C, Aisporna AE, Rinehart D, Hoang L, Nordstrom A, Gomez-Romero M, Whiley L,
Lewis MR, Nicholson JK, Benton HP, Siuzdak G. 2018. XCMS-MRM and METLIN-MRM: a
cloud library and public resource for targeted analysis of small molecules. Nat Methods
15:681-684.

50. Dietrich LE, Price-Whelan A, Petersen A, Whiteley M, Newman DK. 2006. The phenazine
pyocyanin is a terminal signalling factor in the quorum sensing network of Pseudomonas
aeruginosa. Mol Microbiol 61:1308-21.

51. Sakhtah H, Koyama L, Zhang Y, Morales DK, Fields BL, Price-Whelan A, Hogan DA, Shepard
K, Dietrich LE. 2016. The Pseudomonas aeruginosa efflux pump MexGHI-OpmD transports
a natural phenazine that controls gene expression and biofilm development. Proc Natl
Acad Sci U S A 113:E3538-47.

52. Schiessl KT, Hu F, Jo J, Nazia SZ, Wang B, Price-Whelan A, Min W, Dietrich LEP. 2019.
Phenazine production promotes antibiotic tolerance and metabolic heterogeneity in
Pseudomonas aeruginosa biofilms. Nat Commun 10:762.

53. Sporer AJ, Beierschmitt C, Bendebury A, Zink KE, Price-Whelan A, Buzzeo MC, Sanchez LM,
Dietrich LEP. 2018. Pseudomonas aeruginosa PumA acts on an endogenous phenazine to
promote self-resistance. Microbiology (Reading) 164:790-800.

54. Blankenfeldt W, Kuzin AP, Skarina T, Korniyenko Y, Tong L, Bayer P, Janning P, Thomashow
LS, Mavrodi DV. 2004. Structure and function of the phenazine biosynthetic protein PhzF
from Pseudomonas fluorescens. Proc Natl Acad Sci U S A 101:16431-6.

55. Blankenfeldt W, Parsons JF. 2014. The structural biology of phenazine biosynthesis. Curr
Opin Struct Biol 29:26-33.

56. Mavrodi DV, Peever TL, Mavrodi OV, Parejko JA, Raaijmakers JM, Lemanceau P, Mazurier
S, Heide L, Blankenfeldt W, Weller DM, Thomashow LS. 2010. Diversity and evolution of
the phenazine biosynthesis pathway. Appl Environ Microbiol 76:866-79.

57. Parsons JF, Song F, Parsons L, Calabrese K, Eisenstein E, Ladner JE. 2004. Structure and
function of the phenazine biosynthesis protein PhzF from Pseudomonas fluorescens 2-79.
Biochemistry 43:12427-35.

58. Bull CT, Shetty KG, Subbarao KV. 2002. Interactions Between Myxobacteria, Plant
Pathogenic Fungi, and Biocontrol Agents. Plant Dis 86:889-896.

59. Leong W, Lutz, C., Williams, J., Yan, H.P., Benny, Y.K.Y., Chua, C., Rice, S.A., Givskov, M.,
Sanderson-Smith, M., McDougald, D. 2020. *Pseudomonas aeruginosa* isolates co-
incubated with *Acanthamoeba castellanii* exhibit phenotypes similar to chronic cystic
fibrosis isolates. *bioRxiv* doi:https://doi.org/10.1101/2020.02.25.964320.

60. Winstanley C, O'Brien S, Brockhurst MA. 2016. Pseudomonas aeruginosa Evolutionary
Adaptation and Diversification in Cystic Fibrosis Chronic Lung Infections. Trends Microbiol
24:327-337.

61. Malone JG. 2015. Role of small colony variants in persistence of Pseudomonas aeruginosa
infections in cystic fibrosis lungs. Infect Drug Resist 8:237-47.

62. Maunders E, Welch M. 2017. Matrix exopolysaccharides; the sticky side of biofilm
formation. FEMS Microbiol Lett 364.
63. Sun S, Kjelleberg S, McDougald D. 2013. Relative contributions of Vibrio polysaccharide
and quorum sensing to the resistance of Vibrio cholerae to predation by heterotrophic
protists. PLoS One 8:e56338.
64. Butt AT, Thomas MS. 2017. Iron Acquisition Mechanisms and Their Role in the Virulence
of Burkholderia Species. Front Cell Infect Microbiol 7:460.
65. Chekabab SM, Daigle F, Charette SJ, Dozois CM, Harel J. 2013. Shiga toxins decrease
enterohaemorrhagic Escherichia coli survival within Acanthamoeba castellanii. FEMS
Microbiol Lett 344:86-93.
66. Lainhart W, Stolfa G, Koudelka GB. 2009. Shiga toxin as a bacterial defense against a
eukaryotic predator, Tetrahymena thermophila. J Bacteriol 191:5116-22.
67. Steinberg KM, Levin BR. 2007. Grazing protozoa and the evolution of the Escherichia coli
O157:H7 Shiga toxin-encoding prophage. Proc Biol Sci 274:1921-9.
68. Matz C, Moreno AM, Alhede M, Manefield M, Hauser AR, Givskov M, Kjelleberg S. 2008.
Pseudomonas aeruginosa uses type III secretion system to kill biofilm-associated
amoebae. ISME J 2:843-52.
69. Riquelme S, Varas M, Valenzuela C, Velozo P, Chahin N, Aguilera P, Sabag A, Labra B,
Alvarez SA, Chavez FP, Santiviago CA. 2016. Relevant Genes Linked to Virulence Are
Required for Salmonella Typhimurium to Survive Intracellularly in the Social Amoeba
Dictyostelium discoideum. Front Microbiol 7:1305.
70. Pombinho R, Camejo A, Vieira A, Reis O, Carvalho F, Almeida MT, Pinheiro JC, Sousa S,
Cabanes D. 2017. Listeria monocytogenes CadC Regulates Cadmium Efflux and Fine-tunes
Lipoprotein Localization to Escape the Host Immune Response and Promote Infection. J
Infect Dis 215:1468-1479.
71. Vieira A, Ramesh A, Seddon AM, Karlyshev AV. 2017. CmeABC Multidrug Efflux Pump
Contributes to Antibiotic Resistance and Promotes Campylobacter jejuni Survival and
Multiplication in Acanthamoeba polyphaga. Appl Environ Microbiol 83.
72. Perez J, Munoz-Dorado J, Brana AF, Shimkets LJ, Sevillano L, Santamaria RI. 2011.
Myxococcus xanthus induces actinorhodin overproduction and aerial mycelium
formation by Streptomyces coelicolor. Microb Biotechnol 4:175-83.
73. Miskinyte M, Sousa A, Ramiro RS, de Sousa JA, Kotlinowski J, Caramalho I, Magalhaes S,
Soares MP, Gordo I. 2013. The genetic basis of Escherichia coli pathoadaptation to
macrophages. PLoS Pathog 9:e1003802.
74. Scanlan PD, Buckling A. 2012. Co-evolution with lytic phage selects for the mucoid
phenotype of Pseudomonas fluorescens SBW25. ISME J 6:1148-58.
75. Yang F, Gu J, Zou J, Lei L, Jing H, Zhang J, Zeng H, Zou Q, Lv F, Zhang J. 2018. PA0833 Is an
OmpA C-Like Protein That Confers Protection Against Pseudomonas aeruginosa Infection.
Front Microbiol 9:1062.
76. Mulcahy LR, Burns JL, Lory S, Lewis K. 2010. Emergence of Pseudomonas aeruginosa
strains producing high levels of persister cells in patients with cystic fibrosis. J Bacteriol
192:6191-9.

77.   Jeanson S, Floury J, Gagnaire V, Lortal S, Thierry A. 2015. Bacterial Colonies in Solid Media
and Foods: A Review on Their Growth and Interactions with the Micro-Environment. Front
Microbiol 6:1284.
78.   Reyrolle J, Letellier F. 1979. Autoradiographic study of the localization and evolution of
growth zones in bacterial colonies. J Gen Microbiol 111:399-406.
79.   Dillies MA, Rau A, Aubert J, Hennequet-Antier C, Jeanmougin M, Servant N, Keime C,
Marot G, Castel D, Estelle J, Guernec G, Jagla B, Jouneau L, Laloe D, Le Gall C, Schaeffer B,
Le Crom S, Guedj M, Jaffrezic F, French StatOmique C. 2013. A comprehensive evaluation
of normalization methods for Illumina high-throughput RNA sequencing data analysis.
Brief Bioinform 14:671-83.
80.   Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics 26:139-40.
81.   Jones CJ, Ryder CR, Mann EE, Wozniak DJ. 2013. AmrZ modulates Pseudomonas
aeruginosa biofilm architecture by directly repressing transcription of the psl operon. J
Bacteriol 195:1637-44.
82.   Adaikpoh BI, Akbar S, Albataineh H, Misra SK, Sharp JS, Stevens DC. 2020. Myxobacterial
Response to Methyljasmonate Exposure Indicates Contribution to Plant Recruitment of
Micropredators. Front Microbiol 11:34.
83.   Katajamaa M, Miettinen J, Oresic M. 2006. MZmine: toolbox for processing and
visualization of mass spectrometry based molecular profile data. Bioinformatics 22:634-
6.
84.   Olivon F, Grelier G, Roussi F, Litaudon M, Touboul D. 2017. MZmine 2 Data-Preprocessing
To Enhance Molecular Networking Reliability. Anal Chem 89:7836-7840.
85.   Pluskal T, Castillo S, Villar-Briones A, Oresic M. 2010. MZmine 2: modular framework for
processing, visualizing, and analyzing mass spectrometry-based molecular profile data.
BMC Bioinformatics 11:395.
758

Figure 1: A) 4 day sequence of *C. ferrugineus* predation of *P. putida* with complete swarming of prey (t= d3). B) 4 day sequence depicting predator avoidance observed from *P. putida* survivor phenotype including an additional image from day 9 to demonstrate length of avoidance. C) Comparison of colony sizes of predator unexposed *P. putida* (left) and *P. putida* survivor phenotype (right) depicting small colony variance of the survivor phenotype. D) Time to predation data for *P. putida* control and *P. putida* survivor phenotype (n=3; p≤0.002) with E) end point colony forming unit data (n=3; p≤0.006) to determine differences in cell viabilities post-swarming. Unpaired t-test with Welch's correction used for statistical analyses included in D and E; all scale bars depict 1mm.

Figure 2: A) Most abundantly over-expressed genes from survivor phenotype when compared to predator unexposed *P. putida*. B) Up-regulated features associated with alginate production and the associated genomic context for each (C). D) Up-regulated transport proteins including those associated with antibiotic efflux. All data depicted as an averages from 3 biological replicates (p≤0.05).

Figure 3: Growth curves for predator unexposed *P. putida* (A) and *P. putida* survivor phenotype (B) depicting antibiotic resistance unique to the survivor phenotype (n=12). LB supplemented with gentamicin (10 μg/ml), kanamycin (50 μg/ml), and tetracycline (10 μg/ml) where indicated.

Figure 4: A) Isolated alginate concentration determined from carbazole assays depicting
increased alginate production observed from the survivor phenotype (n=16; p≤0.02). B)
Relative intensities of alginate dot blots generated from isolated alginate also depicting
increased alginate production observed from the survivor phenotype (n=10; p=0.039).
Relative units determined from average of control values with statistical significance
calculated from an unpaired t test with Welch's correction.

Figure 5: Relative fluorescence of extracellular pyoverdine depicting increased

pyoverdine production from the survivor phenotype (n=12; p≤0.0001). Relative units

determined from average of control values with statistical significance calculated from an

unpaired t test with Welch's correction.

Figure 6: A) Extracted ion chromatograph (224.9-225.1 m/z) depicting presence of

phenazine-1-carboxylic acid in extracts from survivor phenotype replicates (n=3;

magenta, yellow, cyan) and absence in extracts from predator unexposed *P. putida*

replicates (n=3; red, blue, green). Chromatograph rendered with MZmine v2.37. B)

Comparison of detected phenazine-1-carboxylic acid in predator unexposed *P. putida*

extracts (labelled control) and survivor phenotype extracts (blue). Box plot data and

statistical analysis (n=3; p=0.0509) generated by XCMS v3.7.1 and figure rendered with

Prism v7.0d.

Figure 7: A) Direct contact predation assay as depicted in Figure 1 comparing control

(predator unexposed) *P. putida* with freezer stocks of the survivor phenotype (biological

replicates) (n=8; p≤0.0001). B) Modified indirect contact predation assay with prey

introduced 2 cm from predator (n=8; p≤0.01) and time to predator-prey contact recorded

(C). Swarming assays with *C. ferrugineus* plated on prey lawns and swarming diameters

recorded after 7 days (n=8). Statistical significance calculated using an unpaired t test

with Welch's correction.

821