Predatory selection of mucoid, antibiotic resistant *Pseudomonas putida* phenotype by

myxobacterium *Cystobacter ferrugineus*

Shukria Akbar[a] and D. Cole Stevens[a,#]

Department of BioMolecular Sciences, University of Mississippi, University, MS, USA[a]

Running Head: Predatory selection of antibiotic resistant *P. putida*

#Address correspondence to D. Cole Stevens, stevens@olemiss.edu.

Abstract: Predation contributes to the structure and diversity of microbial communities. Predatory myxobacteria are ubiquitous to a variety of microbial habitats and capably consume a broad diversity of microbial prey. Predator-prey experiments utilizing myxobacteria have provided details into predatory mechanisms and features that facilitate consumption of prey. However, prey resistance to myxobacterial predation remains underexplored, and prey resistances have been observed exclusively from predator-prey experiments that included the model myxobacterium *Myxococcus xanthus*. Utilizing a predator-prey pairing that instead included the myxobacterium, *Cystobacter ferrugineus,* with *Pseudomonas putida* as prey, we infrequently observed surviving phenotypes capable of eluding predation. Comparative transcriptomics between *P. putida* unexposed to *C. ferrugineus* and the survivor phenotype suggested that increased expression of efflux pumps, genes associated with mucoid conversion, and various membrane features contribute to predator avoidance. The *P. putida* survivor phenotype was confirmed to be resistant to the antibiotics kanamycin, gentamicin, and tetracycline and to produce more alginate than predator-unexposed *P. putida*. Unique features observed from the survivor phenotype including small colony variation, efflux-mediated antibiotic resistance, and increased mucoid conversion overlap with traits associated with *Pseudomonas aeruginosa* predator avoidance and pathogenicity. The survivor phenotype also benefited from increased predator resistance during subsequent predation assays. These results demonstrate the utility of myxobacterial predator-prey models and provide insight into prey resistances in response to predatory stress might contribute to the phenotypic diversity and structure of bacterial communities.

Introduction

Abundant within soils and marine environments, predatory myxobacteria contribute to nutrient cycling within the microbial food web (Findlay, 2016; Munoz-Dorado et al., 2016; Albataineh and Stevens, 2018; Mohr, 2018). Myxobacteria are prolific producers of antimicrobial specialized metabolites and display a cooperative, swarming predation strategy that can be readily reproduced and monitored within laboratory settings making them uniquely appropriate for assessment of predator-prey interactions (Munoz-Dorado et al., 2016; Albataineh and Stevens, 2018; Mohr, 2018; Nair et al., 2019). Broadly considered generalist predators, myxobacteria capably predate a range of prey including both Gram-negative and Gram-positive bacteria as well as fungi (Perez et al., 2016; Nair et al., 2019). Constitutive production of specialized metabolites and lytic proteins are often associated with this predatory range, and a variety of metabolites and enzymatic features have been reported to benefit predation (Xiao et al., 2011; Muller et al., 2016; Livingstone et al., 2018; Ellis et al., 2019).

However, relatively few examples of prey resistance to myxobacterial predation have been reported (Sutton et al., 2019). Compared to features associated with bacterial strategies to avoid protozoan predators, prey avoidance of predatory myxobacteria remains underexplored (Weitere et al., 2005; Justice et al., 2008; Erken et al., 2013; Seiler et al., 2017; Sun et al., 2018). Examples of prey responses correlated with resistance to myxobacterial predation include *Escherichia coli* biofilm formation (DePas et al., 2014), *Bacillus subtilis* sporulation and production of bacillaene (Muller et al., 2014; Muller et al., 2015), *Bacillus licheniformus* glycosylation of the predation-associated metabolite myxovirescin A (Wang et al., 2019), galactoglucan exopolysaccharide production and

increased melanin production by *Sinorhizobium meliloti* (Perez et al., 2014; Contreras-

Moreno et al., 2020), and formaldehyde secretion by *Pseudomonas aeruginosa* (Sutton

et al., 2019). All of these features were discovered from predator-prey experiments

utilizing the model myxobacterium *M. xanthus*.

We suspect the development of predator-prey pairings including myxobacteria other than

M. xanthus might provide additional insight into prey resistance to myxobacterial

predation. For our predator-prey experiments, the soil dwelling myxobacterium

Cystobacter ferrugineus strain Cbfe23 was included due to a favorable growth profile and

capability to quickly consume *P. putida* during standard predation assays (Seccareccia

et al., 2015; Akbar et al., 2017; Livingstone et al., 2017). Also found within soils, root

colonizing *P. putida* was chosen as prey due to an established ability to resist protozoan

grazers (Seiler et al., 2017). Initial predator-prey experiments provided a *P. putida*

phenotype capable of avoiding *C. ferrugineus* predation. Herein we report the generation

and predator avoidance of a *P. putida* phenotype resistant to myxobacterial predation

using standard predator-prey experiments, differential gene expression data comparing

the survivor phenotype with predator-unexposed *P. putida*, and traits observed to

potentially contribute to predator avoidance.

Results

Selection of predation resistant *P. putida* phenotype

Utilizing predation assays where an inoculum of *C. ferrugineus* was introduced to the

edge of an established spot of *P. putida* (Seccareccia et al., 2015), we considered

swarming overtaking the *P. putida* spot with no visible prey remaining to be an endpoint

for assessment of viable prey cells using colony forming unit (CFU) assays. Typically, *C.*

ferrugineus swarmed predation sensitive *P. putida* with no visible prey after 3 days of co-

cultivation on nutrient restrictive WAT agar (Figure 1A and 1D). However, from 100 total

assays 5 provided minimally consumed and readily observable *P. putida* remaining after

10-14 days (Figure 1B and 1D). Additionally, instead of frontal attack at the predator-prey

interface when exposed to predation sensitive *P. putida*, *C. ferrugineus* swarmed the

perimeter of the survivor phenotype (Supplemental Figure 1). Exponentially fewer viable

colonies of predation sensitive *P. putida* were observed from endpoint CFU assays (3-4

days for predation sensitive *P. putida*; 10-14 days for *P. putida* survivor phenotype)

supporting the assumption that swarming resistance correlates with prey survival (Figure

1D and 1E). Also apparent from CFU assays, colonies of the survivor phenotype were

consistently smaller when compared to predation sensitive *P. putida* (Figure 1C).

Predation resistance of the survivor phenotype was also observed from additional

predation assays with the optical density of prey inoculum doubled (OD_{600}=1.0 versus

OD_{600}=0.5). (Supplemental Figure 2). Stocks of each *P. putida* survivor observed from

predation assays were stored at -80°C as individual aliquots and utilized to determine

features associated with the survivor phenotype that might contribute to predator

avoidance.

Differential gene expression of predator-unexposed *P. putida* and survivor

phenotype

Utilizing RNA sequencing, we conducted comparative transcriptomic experiments

between the predator unexposed (control) *P. putida* and the survivor phenotype (n=3

biological replicates with biological replicates for the survivor phenotype defined as an

individually observed phenotype resistant to *C. ferrugineus* predation for ≥10 days). This

comparative analysis determined that 1,178 genes were down-regulated ≥4-fold and 122

genes were up-regulated ≥4-fold across the survivor phenotype replicates when

compared to control *P. putida* replicates (n=3; p≤0.05). Considering up-regulated genes

from the survivor phenotype with the highest fold change values, apparent differences in

overexpressed features specific to the survivor phenotype included genes involved in

mucoid conversion, siderophore production, and a variety of membrane-associated

proteins (Figure 2A). A 4 gene operon including a ferric uptake regulatory (*furA*)

associated gene (WP_016501343.1), a fumarate hydratase (WP_016501342.1), a

manganese superoxide dismutase (*sodA*) (WP_016501340.1), and a hypothetical protein

with a conserved DUF2753 domain (WP_06501341.1) was significantly up-regulated in

the survivor phenotype (Figure 2B and 2C). Expression of this operon from *P. aeruginosa*

in response to iron limitation leads to elevated alginate production, an exopolysaccharide

composed of mannuronic acid and guluronic acid monomers, and increased mucoidy

(Hassett et al., 1997b; Hassett et al., 1997a) and corroborates our observation of small

colony variance from the survivor phenotype. Interestingly, the alginate regulatory

elements MucA (WP_016498171.1) and MucB (WP_016498172.1) were both

significantly down regulated in the survivor phenotype (Figure 2B). The anti-sigma factor

MucA and the associated binding partner MucB inhibit mucoid conversion by

sequestering the alternative sigma factor AlgU required for alginate production in *P.*

aeruginosa (Deretic et al., 1993; Martin et al., 1993; Damron and Goldberg, 2012; Li et

al., 2019). Increased transcription from this operon combined with decreased transcription

of MucA and MucB, indicates that mucoid conversion might contribute to predator

avoidance which parallels *P. aeruginosa* predatory avoidance mechanisms when

exposed to grazing amoeba (Matz et al., 2004). Also associated with iron limitation,

regulatory and biosynthetic genes contributing to production of the siderophore

pyoverdine were significantly up-regulated in the survivor phenotype. These included

PvdA an L-ornithine monooxygenase (WP_016499102.1) and PvdH a diaminobutyrate-

2-oxoketoglutarate transaminase (WP_016498669.1) both involved in pyoverdine

precursor biosynthesis, a regulatory transcription factor PvdS (WP_016498655.1), a

proximal MbtH-like protein (WP_016499090.1), and a core nonribosomal peptide

synthetase homologous to PvdL from *P. aeruginosa* (NP_746359.1) (Supplemental

Figure 3) (Drake et al., 2007; Felnagle et al., 2010; Parker et al., 2014; Ringel and Bruser,

2018). Also associated with iron limiting conditions, a bacterioferritin-associated

ferredoxin (WP_016488781.1) was upregulated in the survivor phenotype. A variety of

membrane-associated proteins were also overexpressed in the survivor phenotype

including an osmotically inducible protein OsmY (WP_0016501632.1), an outer

membrane protein A (OmpA) family protein (WP_016501196.1), an FKBP-type peptidyl-

prolyl cis-trans isomerase (WP_0165010671), an aquaporin (WP_016498616.1), and a

DUF1282 domain-containing hypothetical protein from the Yip1 superfamily

(WP_016499858.1) (Figure 2A). While not a membrane associated feature, we also found

the overexpression of a formaldehyde dehydrogenase (WP_016497631.1) by the

survivor phenotype to be of note due to the recent observation that suggests

formaldehyde secretion to be a predation-resistance trait of *P. aeruginosa* (Sutton et al.,

2019). The abundance of up-regulated genes that encode membrane-associated

products in the survivor phenotype including those responsible for increased mucoid

production provides insight into features involved in predation avoidance. Numerous

efflux and transport proteins were also among the most up-regulated genes observed

from the survivor phenotype (Figure 2D). Upregulated efflux genes included inner

(WP_016498130 and WP_016499477.1) and outer (WP_016498128.1) membrane

components highly homologous to the AcrAD-TolC-type multidrug resistance-nodulation-

division (RND) family of efflux pumps and the Mex RND efflux pumps known to facilitate

P. aeruginosa resistance to aminoglycosides and tetracyclines (Aeschlimann, 2003; Li et

al., 2015; Garneau-Tsodikova and Labby, 2016). Genes encoding features homologous

to multidrug efflux pumps contributing to virulence of human pathogens were also up-

regulated in the survivor phenotype including a P-type ATPase (WP_016502100) and an

additional RND/MmpL (Mycobacterial membrane protein Large) protein (Figure 2D).

Provided theses observed differences in gene expression, we sought to determine if the

survivor phenotype demonstrated antibiotic resistance and increased mucoidy, which

have both been previously associated with predator avoidance strategies.

Antibiotic resistance of survivor phenotype

Provided the observation that numerous transport proteins were overexpressed in the

survivor phenotype (Figure 2D) including RND-type efflux pumps known to contribute to

P. aeruginosa antibiotic resistances, we were interested to determine if the survivor

phenotype exhibited antibiotic resistance. The survivor phenotype was capable of initially

forming colonies on LB agar supplemented with the antibiotics gentamicin (10 μg/ml),

kanamycin (50 μg/ml), and tetracycline (10 μg/ml), with no colonies from predation

sensitive *P. putida* observed on identical medias. Subsequent growth curve assays

comparing predation sensitive *P. putida* with the survivor phenotype when grown in LB

and LB supplemented with antibiotics (Figure 3A and 3B) confirmed that the survivor

phenotype was uniquely resistant to gentamicin, kanamycin, and tetracycline. These

results combined with the overexpression of efflux-associated proteins observed from the

survivor phenotype confirms that *C. ferrugineus* selection provides prey that benefit from

antibiotic resistance and suggests that efflux-mediated antibiotic resistance is a *P. putida*

predation avoidance feature.

Increased production of alginate by survivor phenotype

Alginate-based formation of mucoid biofilms has previously been associated with *P.*

aeruginosa avoidance of the protozoan grazer *Rhynchomonas nasuta* (Matz et al., 2002;

Matz et al., 2004; Matz and Kjelleberg, 2005; Matz et al., 2005). Utilizing established

carbazole assays on isolated alginate (Knutson and Jeanes, 1968; Al Ahmar et al., 2020),

we sought to determine if the increased transcription of genes known to impact alginate

production observed from the survivor phenotype resulted in increased production of

alginate when compared to predator unexposed *P. putida*. Comparing isolated alginate

from predation sensitive *P. putida* with the survivor phenotype, we observed that the

survivor phenotype indeed produced more alginate when compared to *P. putida* not

exposed to *C. ferrugineus* (Figure 4).

Predation resistance of survivor phenotype in subsequent predation assays

Additional subsequent predation assays were done to assess the predator avoidance trait

of the *P. putida* survivor phenotype with *P. putida* not exposed to *C. ferrugineus* included

as a control. These predation assays included an assay as previously described with

predator cells directly introduced to the edge of established prey, a similar slightly

modified assay with predator introduced at a distance of 5-7 mm from an established spot

of *P. putida*, and a lawn culture assay (Livingstone et al., 2017) monitoring predator

swarming diameters when placed at the center of prey embedded in agar. For direct

contact assays, time to complete swarming was recorded. As observed in our prior

predator-prey experiments, compared to control *P. putida* the survivor demonstrated

significantly increased predator avoidance (Figure 5A). The survivor phenotype also

demonstrated resistance to swarming compared to predation sensitive *P. putida* during

identical direct-contact predation assays conducted on nutrient-rich VY/2 agar

(Supplemental Figure 4) For indirect contact predation assays, time to predator-prey

contact and time to complete swarming were recorded. Time to predator-prey contact

was between 1-2 days for all samples, and as observed in the directed contact predation

assays the survivor phenotype demonstrated increased predator avoidance when

compared to control *P. putida* (Figure 5B and 5C). However, no differences between

swarming diameters of *C. ferrugineus* grown on lawns of control *P. putida* and the *P.*

putida survivor phenotype were observed (Figure 5D). These data suggest the *P. putida*

survivor phenotype benefits from a predator avoidance trait when subjected to 2 out of 3

subsequent predation assays.

Discussion

Building upon previously reported resistance strategies to avoid predatory myxobacteria

discovered from predator-prey experiments (DePas et al., 2014; Muller et al., 2014; Perez

et al., 2014; Muller et al., 2015; Sutton et al., 2019; Wang et al., 2019; Contreras-Moreno

et al., 2020), predation experiments including *C. ferrugineus* and *P. putida* provided a

prey phenotype significantly resistant to predation. Traits associated with prey resiliency

during bacterial predator-prey interactions remain underexplored despite growing

evidence that predatory stress contributes to the development of phenotypic traits

associated with persistent clinical pathogens (Erken et al., 2013; Sun et al., 2018; Leong,

2020). As an example, clinical isolates of persistent *P. aeruginosa* from cystic fibrosis

patients exhibit increased mucoidy and small colony variation as well as efflux-mediated

antibiotic resistance (Malone, 2015; Winstanley et al., 2016; Maunders and Welch, 2017).

Other human pathogens that have demonstrated overlapping factors involved in both

virulence and anti-predation strategies include *Legionella pneumophila*, *Escherichia coli*,

Vibrio cholerae, *Campylobacter jejuni*, *Listeria monocytogenes*, *Mycobacterium leprae*,

and *Yersinia enterocolitica* (Erken et al., 2013; Sun et al., 2018). Overlapping factors thus

far observed to be associated with both virulence and predator avoidance include biofilm

formation (Matz et al., 2004; Matz et al., 2005; Weitere et al., 2005), quorum signal (Sun

et al., 2013) and siderophore (Butt and Thomas, 2017) small molecule biosynthesis, toxin

production (Steinberg and Levin, 2007; Lainhart et al., 2009; Chekabab et al., 2013), and

expression of transport proteins such as type III and type VI secretion systems (Matz et

al., 2008; Riquelme et al., 2016) and antibiotic resistance-associated efflux pumps

(Pombinho et al., 2017; Vieira et al., 2017). While the impact of predatory selection on

virulence factors have primarily been reported from predator-prey experiments with

grazing amoeba, efforts to determine the influence of generalist predators on microbial

communities have suggested that bacterial predators also drive adaptation of prey traits

that overlap with virulence in bacterial pathogens (Nair et al., 2019). Recently, Nair *et al.*

reported that during predator-prey interactions with *E. coli* prey, the predatory

myxobacterium *Myxococcus xanthus* selects for 2 traits associated with virulence,

mucoidy and increased expression of the outer membrane protease OmpT (Nair et al.,

2019). Compared to features associated with bacterial strategies to avoid protozoan

predators, prey avoidance of myxobacterial predators remains underexplored. Although

predator-prey experiments that utilize myxobacteria with various prey such as *Bacillus*

subtilis (Muller et al., 2014; Muller et al., 2015; Muller et al., 2016), *E. coli* (Livingstone et

al., 2018; Nair et al., 2019), *Sinorhizobium melilotti* (Perez et al., 2014; Contreras-Moreno

et al., 2020), *P. aeruginosa* (Sutton et al., 2019), and *Streptomyces coelicolor* (Perez et

al., 2011) have been reported, all myxobacterial resistances have been discovered from

predator-prey experiments that included the model myxobacterium *M. xanthus*. Utilizing

an alternative myxobacterium *C. ferrugineus* provides the opportunity to further compare

predatory selection and adaptive prey avoidance strategies.

Comparative transcriptomics provided key features up-regulated in the *P. putida* survivor

phenotype that overlap with traits associated with *P. aeruginosa* predator avoidance and

pathogenicity. Of these, 3 putative RND family transporter components with high

homology to the MexAB-OprM and MexCD-OprM efflux systems known to mediate *P.*

aeruginosa antibiotic resistance were up-regulated in the survivor phenotype

(Aeschlimann, 2003; Li et al., 2015). Growth curve analysis determined that the *P. putida*

survivor phenotype was indeed resistant to the aminoglycosides kanamycin and

gentamicin as well as tetracycline; susceptibility to all 3 of these antibiotics was observed

from predator unexposed *P. putida*. Also of note, the multidrug efflux pump CmeABC

serves as a survival mechanism for *C. jejuni* when engulfed by the predatory amoeba

Acanthamoeba polyphaga (Vieira et al., 2017). Several features associated with mucoid

conversion were also up-regulated in the survivor phenotype including a 4 gene operon

responsible for elevated alginate production in *P. aeruginosa* and the regulatory

transcription factors MucA and MucB (Hassett et al., 1997b; Hassett et al., 1997a;

Damron and Goldberg, 2012; Li et al., 2019). Adaptive mucoid phenotypes have been

previously been observed in response to myxobacterial predation (Nair et al., 2019),

amoebal grazing (Matz et al., 2002), macrophages (Miskinyte et al., 2013), and lytic

phages (Scanlan and Buckling, 2012). Increased alginate production by the survivor

phenotype was confirmed using established carbazole assays. Alginate-based mucoid

conversion and biofilm formation have previously been associated with *P. aeruginosa*

avoidance of grazing amoeba and could also contribute to the small colony variation

observed from our survivor phenotype (Matz et al., 2002; Matz et al., 2004; Matz and

Kjelleberg, 2005). These results and the observed alternative predatory pattern of *C.*

ferrugineus during predator-prey interactions with the survivor phenotype parallel

previous observation of mucoid *S. meliloti* phenotypes avoiding *M. xanthus* predation

(Perez et al., 2014). Also of note, upregulation of an OmpA family protein observed in

the survivor phenotype might also correlate with mucoid conversion as *P. aeruginosa*

phenotypes that upregulate expression of OmpA also overproduce alginate (Yang et al.,

2018). These results provide clear overlap between features contributing to avoidance of

myxobacterial and amoebal predation and also afford additional evidence that the

coincidental selection of prey virulence factors reported from amoebal predatory stress

(Erken et al., 2013) might also be observed from predator-prey interactions involving

myxobacteria. Both efflux-mediated antibiotic resistance and mucoid conversion have

also been attributed to virulence of clinical isolates of *P. aeruginosa* (Deretic et al., 1993;

Mulcahy et al., 2010; Winstanley et al., 2016; Leong, 2020). Additionally, transcriptomic

data provided key features for follow-up investigations to assess any direct impact on

predator avoidance including biosynthetic genes involved in pyoverdine biosynthesis and

numerous features associated with outer membranes. For example, up-regulation of the

pyoverdine biosynthetic pathway and a bacterioferritin-associated ferredoxin observed

from the survivor phenotype suggests an iron-depleted predator-prey environment similar

to the previously interaction between *M. xanthus* and *Streptomyces coelicolor* (Perez et

al., 2011). Enhanced production of the primary myxobacterial siderophore myxochelin by

M. xanthus contributed to an iron-restrictive environment and triggered production of the

pigment actinorhodin from *S. coelicolor.*

The survivor phenotype observed to be resistant to *C. ferrugineus* predation was

assessed with subsequent predation assays to determine the resiliency of the predator

avoidance trait. While comparative transcriptomic data provided insight into key up-

regulated features potentially involved in prey survival, down-regulation of 1,178 genes in

the survivor phenotype when compared to predator unexposed *P. putida* suggests

potentially diminished survivor fitness. Despite this, the survivor phenotype maintained

the predator avoidance trait in 2 out of 3 subsequent predation assays. While it is unclear

why no significant predator avoidance was observed from lawn culture predation assays,

we suspect the introduction of predator to the center of prey lawns as opposed to

consistent introduction to the peripheral interface of prey potentially reduce any predator

avoidance mechanisms. Predator avoidance observed from direct and indirect contact

predation assays and the heterogeneity associated with differences between the growth

and metabolism of peripheral cells and central cells of bacterial colonies supports this

suspicion (Reyrolle and Letellier, 1979; Jeanson et al., 2015). We anticipate that *P. putida*

phenotypes resistant to myxobacterial predation continually subjected to such predatory

selection will afford adapted phenotypes for further scrutiny of any resulting mutations

and genotypes. Ultimately, these results demonstrate the utility of predator-prey

experiments using lesser studied myxobacteria and provide insight into how predatory

stress might contribute to adaptive phenotypic diversity of bacterial communities.

Materials and Methods

Bacterial Strains and Cultivation

The myxobacterium *Cystobacter ferrugineus* strain Cbfe23, DSM 52764 was employed

as the predator. *Pseudomonas putida* type strain, ATCC 12633, and the discussed *P.*

putida survivor phenotype resulted from predation assays with this *P. putida* type strain

were included as prey. *C. ferrugineus* was grown on VY/2 solid (1.4% w/v agar, 0.1% w/v

$CaCl_2$ * 2 H_2O, 0.5% w/v Baker's yeast, 500 µM vitamin B12) media for 5-7 days. Luria-

Bertani (LB) solid (1.5% agar) and liquid media were utilized for the cultivation of *P.*

putida. Predation experiments were performed on WAT agar (1.5% w/v agar, 0.1% w/v

$CaCl_2$, 20 mM HEPES) plates. All bacteria were grown at 30 °C.

Predation assays

Prey spot assay was performed as described by Seccareccia et al. (Seccareccia et al.,

2015). Briefly, *P. putida* prey was grown in LB liquid media on a rotary shaker (150 rpm)

at 30 °C for 16-18 h. Next, the bacterial cultures were sedimented by centrifugation at

4,000 g for 15 min, and the sedimented cells were washed and resuspended in TM buffer

(50 mM Tris, pH 7.8, 10 mM $MgSO_4$) to an OD_{600} of 0.5. Then the WAT agar plates were

spotted with 150 µl of the freshly prepared bacterial suspensions, and the spots were

dried in a laminar flow hood. Finally, *C. ferrugineus* was inoculated at the edge of the prey

spot. The assay plates were incubated at 30 °C for 14 days or until the full visible

swarming of *C. ferrugineus* on the prey spot. The assay was performed with 100

replicates. Any prey spot with visible biomass of *P. putida* after day 14 was considered

as survivor phenotype of *P. putida*, and five observed *P. putida* survivor stocks were

acquired that remained visible after day 14 of the assay.

For subsequent predation assays, *P. putida* predation sensitive (control) and the survivor

phenotypes were employed as prey. The direct spot assay was performed as mentioned

above. Statistical significance of recorded time to predation reported in was calculated

using an unpaired t test with Welch's correction in Prism 7.0d. The subsequent directed

contact predation assay was also performed on VY/2 (1.4% w/v agar, 0.1% w/v $CaCl_2$ *

2 H_2O, 0.5% w/v Baker's yeast, 500 µM vitamin B12) media as well, considered as

nutrient-rich media. For indirect contact spot assay, *C. ferrugineus* was inoculated at a

distance of 5-7 mm from the established *P. putida* control and *P. putida* survivor prey

spots. Next, time *C. ferrugineus* took to reach the prey spots and the time of its swarming

over the spots was recorded. Indirect predation assays were performed with 8 replicates

per phenotype. Statistical significance of recorded time to predation reported in days and

recorded time to initial contact between *C. ferrugineus* and prey reported in days was

calculated using an unpaired t test with Welch's correction in Prism 7.0d.

Swarming predation assays were based on established lawn culture methods

(Livingstone et al., 2017). In brief, *P. putida* prey phenotypes were grown in LB liquid

media for 16-18 h at 30 °C with shaking at 150 rpm. The cultures were then subjected to

centrifugation at 4,000 g for 15 min, and the sedimented cells were washed and

resuspended in TM buffer (50 mM Tris, pH 7.8, 10 mM $MgSO_4$) to an OD_{600} 0.5. Next, 1

ml of the cell suspension was spread onto a 14 cm diameter WAT agar plate and dried to

form a uniform prey lawn. Finally, *C. ferrugineus* previously cultured on VY/2 agar media

was spotted onto the center of the prey lawn. The assay plates were incubated at 30 °C,

and the expansion of the *C. ferrugineus* swarm was measured on day 4. Lawn culture

predation assays were performed with 8 replicates per phenotype with replicates from

each survivor stock included. All 3 predation assays were performed simultaneously for

all replicates. Statistical significance of *C. ferrugineus* swarm diameters reported in

millimeters on day 4 was calculated using an unpaired t test with Welch's correction in

Prism 7.0d

CFU assays.

The colony forming units (CFU) were calculated as described by DePas *et al.* (DePas et

al., 2014). Briefly, after the full swarming of *C. ferrugineus* over the prey spot, the agar

slab under the prey spot was excised and suspended in 2 mL phosphate buffer saline

(PBS). Next, a dilution series from this solution was inoculated on the LB plates, and the

plates were incubated at 30 °C for 18 hr. Then, CFU/mL was calculated from the viable

colonies that appeared on the LB plates. Finally, the CFU/mL of the survivor was

compared with cfu/mL of the unexposed *P. putida,* considered as a control. Statistical

significance of all CFU assays was calculated using an unpaired t test with Welch's

correction in Prism 7.0d.

RNA sequencing

Triplicate samples of *P. putida* control and *P. putida* survivor phenotype grown from

stocks in LB at 37°C for 18 hr and pelleted via centrifugation were stored in RNA*later*

solution for sequencing. The DNA from *P. putida* type strain cells (200 µl) was extracted

using MagAttract HMW DNA Kit (Qiagen). The DNA was eluted in 100 uL AE buffer. The

concentration of DNA was evaluated (Supplementary Table 1) using the Qubit® dsDNA

HS Assay Kit (Life Technologies). The library was prepared using Nextera DNA Flex

library preparation kit (Illumina) following the manufacturer's user guide. 50 ng DNA was

used to prepare the library. The sample underwent the simultaneous fragmentation and

addition of adapter sequences. These adapters are utilized during a limited-cycle (6

cycles) PCR in which unique index was added to the sample. Following the library

preparation, the final concentration of the library (Supplementary Table 1) was measured

using the Qubit® dsDNA HS Assay Kit (Life Technologies), and the average library size

(Supplementary Table 1) was determined using the Agilent 2100 Bioanalyzer (Agilent

Technologies). The library was diluted (to 6.5 pM) and sequenced paired end for 500

cycles using the MiSeq system (Illumina).

Total RNA was isolated from triplicate samples of the *P. putida* control, the *P. putida*

survivor phenotype from stocks, and the kanamycin resistant phenotype using the

RNeasy PowerSoil Total RNA Kit (Qiagen) following the manufacturer's instructions. 400

μl cell sample was used for extractions. The concentration of total RNA was determined

(Supplementary Table 2) using the Qubit® RNA Assay Kit (Life Technologies). For rRNA

depletion, first, 1000 ng of total RNA was used to remove the DNA contamination using

Baseline-ZERO™ DNase (Epicentre) following the manufacturer's instructions followed

by purification using the RNA Clean & Concentrator-5 columns (Zymo Research). DNA

free RNA samples were used for rRNA removal by using RiboMinus™ rRNA Removal Kit

(Bacteria; Thermo Fisher Scientific) and final purification was performed using the RNA

Clean & Concentrator-5 columns (Zymo Research). rRNA depleted samples were used

for library preparation using the KAPA mRNA HyperPrep Kits (Roche) by following the

manufacturer's instructions. Following the library preparation, the final concentrations of

all libraries (Supplementary Table 2) were measured using the Qubit® dsDNA HS Assay

Kit (Life Technologies), and the average library size was determined using the Agilent

2100 Bioanalyzer (Agilent Technologies). The libraries were then pooled in equimolar

ratios of 0.6 nM, and sequenced paired end for 300 cycles using the NovaSeq 6000

system (Illumina). Differential expression between the resulting transcriptomes was

calculated from pair-wise analysis of trimmed mean of M-values (TMM) normalized read

counts (Dillies et al., 2013) including fold change, counts-per-million (CPM), and

associated p-values using the R-package, edgeR (Robinson et al., 2010). Differential

expression data with p-values ≤ 0.05 were considered statistically significant. Genome

and RNA sequencing was conducted by MR DNA (Molecular Research LP).   All fold

change, CPM, and associated p-values resulting from differential expression experiments

as described included as supplementary data files TABLE 1 (*P. putida* survivor compared

with *P. putida* control). All raw RNAseq data is publicly available at the NCBI Sequence

Read Archive (PRJNA577468).

Antibiotic Susceptibility Assays.

The standard growth curve assay was employed to determine the sensitivity of the

unexposed *P. putida* and the survivor phenotype towards kanamycin (50 mg/mL),

gentamicin (10 mg/mL), and tetracycline (10 mg/mL) antibiotics. The Lauria Bertani (LB)

broth without any supplemented antibiotics was used as a control. The test was

performed in a 96-well microtiter plate. In a 96-well plate, each well having 200 µL of LB

or LB supplemented with an antibiotic was inoculated with a colony of *P. putida.* Next, the

absorbance at OD_{620} was recorded using a plate reader every 30 min for 10 hr to examine

the growth dynamics of each phenotype. The test was performed in 12 replicates per

condition for both phenotypes. Recorded OD_{620} values were plotted using Prism version

7.0d with error bars depicting standard deviation of replicate data.

Carbazole Assays.

The alginate was isolated according to Jones *et al*. (Jones et al., 2013). In brief, both *P.*

putida phenotype unexposed to the predatory stress and the survivor phenotype were

grown overnight in 2 ml LB media and adjusted to an OD_{600} of 1.00. Following

centrifugation, 1 ml of supernatant was treated with 2% cetyl pyridinium chloride to

precipitate the alginate. The precipitated alginate was further collected by centrifugation.

It was resuspended in 1 M NaCl and then re-precipitated in cold isopropanol. Precipitated

alginate was suspended in 150 μl 0.9% (w/v) saline solution. Following established

protocols (Knutson and Jeanes, 1968; Al Ahmar et al., 2020), a 50 µL aliquot of the

isolated alginate and a dilution series of standard alginic acid (Sigma) were mixed with

200 µL of a solution of 25 mM sodium tetraborate in sulfuric acid and added in a 96-well

plate. Next, the plate was heated for 10 min at 100 °C in an oven. After cooling at room

temperature for 15 min, 50 μl of a 0.125% carbazole solution in absolute ethanol was

added. Then, the plate was re-heated at 100 °C for 10 min in an oven and cooled down

at room temperature for 15 min. Finally, the plate was read in a microplate reader at a

wavelength of 550 nm. Quantities of alginate per sample (μg/ml) were calculated from the

resulting standard curve of purchase alginic acid, and statistical significance was

calculated using an unpaired t test with Welch's correction in Prism 7.0d.

Author Contributions.

SA performed the experiments. DCS supervised the project. SA and DCS conceived the

project, designed the experiments, and wrote the manuscript.

Acknowledgements.

The authors appreciate funding and support from the National Institute of Allergy and

Infectious Diseases (R15AI137996) and the National Institute of General Medical

Sciences of the National Institutes of Health (P20GM130460). The authors appreciate

Scot E. Dowd at MR DNA, Molecular Research LP for assistance throughout sequencing

services provided.

References.

Aeschlimann, J.R. (2003) The role of multidrug efflux pumps in the antibiotic resistance of
Pseudomonas aeruginosa and other gram-negative bacteria. Insights from the Society of
Infectious Diseases Pharmacists. *Pharmacotherapy* **23**: 916-924.

Akbar, S., Dowd, S.E., and Stevens, D.C. (2017) Draft Genome Sequence of Cystobacter
ferrugineus Strain Cbfe23. *Genome Announc* **5**.

Al Ahmar, R., Kirby, B.D., and Yu, H.D. (2020) Culture of Small Colony Variant of Pseudomonas
aeruginosa and Quantitation of its Alginate. *J Vis Exp*.

Albataineh, H., and Stevens, D.C. (2018) Marine Myxobacteria: A Few Good Halophiles. *Mar*
Drugs **16**.

Butt, A.T., and Thomas, M.S. (2017) Iron Acquisition Mechanisms and Their Role in the Virulence
of Burkholderia Species. *Front Cell Infect Microbiol* **7**: 460.

Chekabab, S.M., Daigle, F., Charette, S.J., Dozois, C.M., and Harel, J. (2013) Shiga toxins decrease
enterohaemorrhagic Escherichia coli survival within Acanthamoeba castellanii. *FEMS Microbiol*
Lett **344**: 86-93.

Contreras-Moreno, F.J., Munoz-Dorado, J., Garcia-Tomsig, N.I., Martinez-Navajas, G., Perez, J.,
and Moraleda-Munoz, A. (2020) Copper and Melanin Play a Role in Myxococcus xanthus
Predation on Sinorhizobium meliloti. *Front Microbiol* **11**: 94.

Damron, F.H., and Goldberg, J.B. (2012) Proteolytic regulation of alginate overproduction in
Pseudomonas aeruginosa. *Mol Microbiol* **84**: 595-607.

DePas, W.H., Syed, A.K., Sifuentes, M., Lee, J.S., Warshaw, D., Saggar, V. et al. (2014) Biofilm
formation protects Escherichia coli against killing by Caenorhabditis elegans and Myxococcus
xanthus. *Appl Environ Microbiol* **80**: 7079-7087.

Deretic, V., Martin, D.W., Schurr, M.J., Mudd, M.H., Hibler, N.S., Curcic, R., and Boucher, J.C.
(1993) Conversion to mucoidy in Pseudomonas aeruginosa. *Biotechnology (N Y)* **11**: 1133-1136.

Dillies, M.A., Rau, A., Aubert, J., Hennequet-Antier, C., Jeanmougin, M., Servant, N. et al. (2013)
A comprehensive evaluation of normalization methods for Illumina high-throughput RNA
sequencing data analysis. *Brief Bioinform* **14**: 671-683.

Drake, E.J., Cao, J., Qu, J., Shah, M.B., Straubinger, R.M., and Gulick, A.M. (2007) The 1.8 A crystal
structure of PA2412, an MbtH-like protein from the pyoverdine cluster of Pseudomonas
aeruginosa. *J Biol Chem* **282**: 20425-20434.

Ellis, B.M., Fischer, C.N., Martin, L.B., Bachmann, B.O., and McLean, J.A. (2019) Spatiochemically
Profiling Microbial Interactions with Membrane Scaffolded Desorption Electrospray Ionization-
Ion Mobility-Imaging Mass Spectrometry and Unsupervised Segmentation. *Anal Chem* **91**: 13703-
13711.

Erken, M., Lutz, C., and McDougald, D. (2013) The rise of pathogens: predation as a factor driving
the evolution of human pathogens in the environment. *Microb Ecol* **65**: 860-868.

Felnagle, E.A., Barkei, J.J., Park, H., Podevels, A.M., McMahon, M.D., Drott, D.W., and Thomas,
M.G. (2010) MbtH-like proteins as integral components of bacterial nonribosomal peptide
synthetases. *Biochemistry* **49**: 8815-8817.

Findlay, B.L. (2016) The Chemical Ecology of Predatory Soil Bacteria. *ACS Chem Biol* **11**: 1502-
1510.

Garneau-Tsodikova, S., and Labby, K.J. (2016) Mechanisms of Resistance to Aminoglycoside
Antibiotics: Overview and Perspectives. *Medchemcomm* **7**: 11-27.

Hassett, D.J., Howell, M.L., Sokol, P.A., Vasil, M.L., and Dean, G.E. (1997a) Fumarase C activity is
elevated in response to iron deprivation and in mucoid, alginate-producing Pseudomonas
aeruginosa: cloning and characterization of fumC and purification of native fumC. *J Bacteriol* **179**:
1442-1451.

Hassett, D.J., Howell, M.L., Ochsner, U.A., Vasil, M.L., Johnson, Z., and Dean, G.E. (1997b) An
operon containing fumC and sodA encoding fumarase C and manganese superoxide dismutase is
controlled by the ferric uptake regulator in Pseudomonas aeruginosa: fur mutants produce
elevated alginate levels. *J Bacteriol* **179**: 1452-1459.

Jeanson, S., Floury, J., Gagnaire, V., Lortal, S., and Thierry, A. (2015) Bacterial Colonies in Solid
Media and Foods: A Review on Their Growth and Interactions with the Micro-Environment. *Front
Microbiol* **6**: 1284.

Jones, C.J., Ryder, C.R., Mann, E.E., and Wozniak, D.J. (2013) AmrZ modulates Pseudomonas
aeruginosa biofilm architecture by directly repressing transcription of the psl operon. *J Bacteriol*
195: 1637-1644.

Justice, S.S., Hunstad, D.A., Cegelski, L., and Hultgren, S.J. (2008) Morphological plasticity as a
bacterial survival strategy. *Nat Rev Microbiol* **6**: 162-168.

Knutson, C.A., and Jeanes, A. (1968) A new modification of the carbazole analysis: application to
heteropolysaccharides. *Anal Biochem* **24**: 470-481.

Lainhart, W., Stolfa, G., and Koudelka, G.B. (2009) Shiga toxin as a bacterial defense against a
eukaryotic predator, Tetrahymena thermophila. *J Bacteriol* **191**: 5116-5122.

Leong, W., Lutz, C., Williams, J., Yan, H.P., Benny, Y.K.Y., Chua, C., Rice, S.A., Givskov, M.,
Sanderson-Smith, M., McDougald, D. (2020) *Pseudomonas aeruginosa* isolates co-incubated with
Acanthamoeba castellanii exhibit phenotypes similar to chronic cystic fibrosis isolates. *bioRxiv*.

Li, S., Lou, X., Xu, Y., Teng, X., Liu, R., Zhang, Q. et al. (2019) Structural basis for the recognition of
MucA by MucB and AlgU in Pseudomonas aeruginosa. *FEBS J* **286**: 4982-4994.

Li, X.Z., Plesiat, P., and Nikaido, H. (2015) The challenge of efflux-mediated antibiotic resistance
in Gram-negative bacteria. *Clin Microbiol Rev* **28**: 337-418.

Livingstone, P.G., Morphew, R.M., and Whitworth, D.E. (2017) Myxobacteria Are Able to Prey
Broadly upon Clinically-Relevant Pathogens, Exhibiting a Prey Range Which Cannot Be Explained
by Phylogeny. *Front Microbiol* **8**: 1593.

Livingstone, P.G., Millard, A.D., Swain, M.T., and Whitworth, D.E. (2018) Transcriptional changes
when Myxococcus xanthus preys on Escherichia coli suggest myxobacterial predators are
constitutively toxic but regulate their feeding. *Microb Genom* **4**.

Malone, J.G. (2015) Role of small colony variants in persistence of Pseudomonas aeruginosa
infections in cystic fibrosis lungs. *Infect Drug Resist* **8**: 237-247.

Martin, D.W., Schurr, M.J., Mudd, M.H., and Deretic, V. (1993) Differentiation of Pseudomonas
aeruginosa into the alginate-producing form: inactivation of mucB causes conversion to mucoidy.
Mol Microbiol **9**: 497-506.

Matz, C., and Kjelleberg, S. (2005) Off the hook--how bacteria survive protozoan grazing. *Trends*
Microbiol **13**: 302-307.

Matz, C., Deines, P., and Jurgens, K. (2002) Phenotypic variation in Pseudomonas sp. CM10
determines microcolony formation and survival under protozoan grazing. *FEMS Microbiol Ecol*
39: 57-65.

Matz, C., Bergfeld, T., Rice, S.A., and Kjelleberg, S. (2004) Microcolonies, quorum sensing and
cytotoxicity determine the survival of Pseudomonas aeruginosa biofilms exposed to protozoan
grazing. *Environ Microbiol* **6**: 218-226.

Matz, C., McDougald, D., Moreno, A.M., Yung, P.Y., Yildiz, F.H., and Kjelleberg, S. (2005) Biofilm
formation and phenotypic variation enhance predation-driven persistence of Vibrio cholerae.
Proc Natl Acad Sci U S A **102**: 16819-16824.

Matz, C., Moreno, A.M., Alhede, M., Manefield, M., Hauser, A.R., Givskov, M., and Kjelleberg, S.
(2008) Pseudomonas aeruginosa uses type III secretion system to kill biofilm-associated
amoebae. *ISME J* **2**: 843-852.

Maunders, E., and Welch, M. (2017) Matrix exopolysaccharides; the sticky side of biofilm
formation. *FEMS Microbiol Lett* **364**.

Miskinyte, M., Sousa, A., Ramiro, R.S., de Sousa, J.A., Kotlinowski, J., Caramalho, I. et al. (2013)
The genetic basis of Escherichia coli pathoadaptation to macrophages. *PLoS Pathog* **9**: e1003802.

Mohr, K.I. (2018) Diversity of Myxobacteria-We Only See the Tip of the Iceberg. *Microorganisms*
6.

Mulcahy, L.R., Burns, J.L., Lory, S., and Lewis, K. (2010) Emergence of Pseudomonas aeruginosa
strains producing high levels of persister cells in patients with cystic fibrosis. *J Bacteriol* **192**: 6191-
6199.

Muller, S., Strack, S.N., Ryan, S.E., Kearns, D.B., and Kirby, J.R. (2015) Predation by Myxococcus
xanthus induces Bacillus subtilis to form spore-filled megastructures. *Appl Environ Microbiol* **81**:
203-210.

Muller, S., Strack, S.N., Hoefler, B.C., Straight, P.D., Kearns, D.B., and Kirby, J.R. (2014) Bacillaene
and sporulation protect Bacillus subtilis from predation by Myxococcus xanthus. *Appl Environ*
Microbiol **80**: 5603-5610.

Muller, S., Strack, S.N., Ryan, S.E., Shawgo, M., Walling, A., Harris, S. et al. (2016) Identification of
Functions Affecting Predator-Prey Interactions between Myxococcus xanthus and Bacillus
subtilis. *J Bacteriol* **198**: 3335-3344.

Munoz-Dorado, J., Marcos-Torres, F.J., Garcia-Bravo, E., Moraleda-Munoz, A., and Perez, J. (2016)
Myxobacteria: Moving, Killing, Feeding, and Surviving Together. *Front Microbiol* **7**: 781.

Nair, R.R., Vasse, M., Wielgoss, S., Sun, L., Yu, Y.N., and Velicer, G.J. (2019) Bacterial predator-
prey coevolution accelerates genome evolution and selects on virulence-associated prey
defences. *Nat Commun* **10**: 4301.

Parker, D.L., Lee, S.W., Geszvain, K., Davis, R.E., Gruffaz, C., Meyer, J.M. et al. (2014) Pyoverdine
synthesis by the Mn(II)-oxidizing bacterium Pseudomonas putida GB-1. *Front Microbiol* **5**: 202.

Perez, J., Moraleda-Munoz, A., Marcos-Torres, F.J., and Munoz-Dorado, J. (2016) Bacterial
predation: 75 years and counting! *Environ Microbiol* **18**: 766-779.

Perez, J., Munoz-Dorado, J., Brana, A.F., Shimkets, L.J., Sevillano, L., and Santamaria, R.I. (2011)
Myxococcus xanthus induces actinorhodin overproduction and aerial mycelium formation by
Streptomyces coelicolor. *Microb Biotechnol* **4**: 175-183.

Perez, J., Jimenez-Zurdo, J.I., Martinez-Abarca, F., Millan, V., Shimkets, L.J., and Munoz-Dorado,
J. (2014) Rhizobial galactoglucan determines the predatory pattern of Myxococcus xanthus and
protects Sinorhizobium meliloti from predation. *Environ Microbiol* **16**: 2341-2350.

Pombinho, R., Camejo, A., Vieira, A., Reis, O., Carvalho, F., Almeida, M.T. et al. (2017) Listeria
monocytogenes CadC Regulates Cadmium Efflux and Fine-tunes Lipoprotein Localization to
Escape the Host Immune Response and Promote Infection. *J Infect Dis* **215**: 1468-1479.

Reyrolle, J., and Letellier, F. (1979) Autoradiographic study of the localization and evolution of
growth zones in bacterial colonies. *J Gen Microbiol* **111**: 399-406.

Ringel, M.T., and Bruser, T. (2018) The biosynthesis of pyoverdines. *Microb Cell* **5**: 424-437.

Riquelme, S., Varas, M., Valenzuela, C., Velozo, P., Chahin, N., Aguilera, P. et al. (2016) Relevant
Genes Linked to Virulence Are Required for Salmonella Typhimurium to Survive Intracellularly in
the Social Amoeba Dictyostelium discoideum. *Front Microbiol* **7**: 1305.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010) edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**: 139-140.

Scanlan, P.D., and Buckling, A. (2012) Co-evolution with lytic phage selects for the mucoid
phenotype of Pseudomonas fluorescens SBW25. *ISME J* **6**: 1148-1158.

Seccareccia, I., Kost, C., and Nett, M. (2015) Quantitative Analysis of Lysobacter Predation. *Appl*
Environ Microbiol **81**: 7098-7105.

Seiler, C., van Velzen, E., Neu, T.R., Gaedke, U., Berendonk, T.U., and Weitere, M. (2017) Grazing
resistance of bacterial biofilms: a matter of predators' feeding trait. *FEMS Microbiol Ecol* **93**.

Steinberg, K.M., and Levin, B.R. (2007) Grazing protozoa and the evolution of the Escherichia coli
O157:H7 Shiga toxin-encoding prophage. *Proc Biol Sci* **274**: 1921-1929.

Sun, S., Kjelleberg, S., and McDougald, D. (2013) Relative contributions of Vibrio polysaccharide
and quorum sensing to the resistance of Vibrio cholerae to predation by heterotrophic protists.
PLoS One **8**: e56338.

Sun, S., Noorian, P., and McDougald, D. (2018) Dual Role of Mechanisms Involved in Resistance
to Predation by Protozoa and Virulence to Humans. *Front Microbiol* **9**: 1017.

Sutton, D., Livingstone, P.G., Furness, E., Swain, M.T., and Whitworth, D.E. (2019) Genome-Wide
Identification of Myxobacterial Predation Genes and Demonstration of Formaldehyde Secretion
as a Potentially Predation-Resistant Trait of Pseudomonas aeruginosa. *Front Microbiol* **10**: 2650.

Vieira, A., Ramesh, A., Seddon, A.M., and Karlyshev, A.V. (2017) CmeABC Multidrug Efflux Pump
Contributes to Antibiotic Resistance and Promotes Campylobacter jejuni Survival and
Multiplication in Acanthamoeba polyphaga. *Appl Environ Microbiol* **83**.

Wang, C., Liu, X., Zhang, P., Wang, Y., Li, Z., Li, X. et al. (2019) Bacillus licheniformis escapes from
Myxococcus xanthus predation by deactivating myxovirescin A through enzymatic glucosylation.
Environ Microbiol **21**: 4755-4772.
Weitere, M., Bergfeld, T., Rice, S.A., Matz, C., and Kjelleberg, S. (2005) Grazing resistance of
Pseudomonas aeruginosa biofilms depends on type of protective mechanism, developmental
stage and protozoan feeding mode. *Environ Microbiol* **7**: 1593-1601.
Winstanley, C., O'Brien, S., and Brockhurst, M.A. (2016) Pseudomonas aeruginosa Evolutionary
Adaptation and Diversification in Cystic Fibrosis Chronic Lung Infections. *Trends Microbiol* **24**:
327-337.
Xiao, Y., Wei, X., Ebright, R., and Wall, D. (2011) Antibiotic production by myxobacteria plays a
role in predation. *J Bacteriol* **193**: 4626-4633.
Yang, F., Gu, J., Zou, J., Lei, L., Jing, H., Zhang, J. et al. (2018) PA0833 Is an OmpA C-Like Protein
That Confers Protection Against Pseudomonas aeruginosa Infection. *Front Microbiol* **9**: 1062.
628

Figure 1: A) 4 day sequence of *C. ferrugineus* predation of *P. putida* with complete swarming of prey (t= d3). B) 4 day sequence depicting predator avoidance observed from *P. putida* survivor phenotype including an additional image from day 9 to demonstrate length of avoidance. C) Comparison of colony sizes of predator unexposed *P. putida* (left) and *P. putida* survivor phenotype (right) depicting small colony variance of the survivor phenotype. D) Time to predation data for *P. putida* control and *P. putida* survivor phenotype (n=3; p≤0.002) with E) end point colony forming unit data (n=3; p≤0.006) to determine differences in cell viabilities post-swarming. Unpaired t-test with Welch's correction used for statistical analyses included in D and E; all scale bars depict 1mm.

Figure 2: A) Most abundantly over-expressed genes from survivor phenotype when compared to predator unexposed *P. putida*. B) Up-regulated features associated with alginate production and the associated genomic context for each (C). D) Up-regulated transport proteins including those associated with antibiotic efflux. All data depicted as an averages from 3 biological replicates (p≤0.05).

Figure 3: Growth curves for predator unexposed *P. putida* (A) and *P. putida* survivor phenotype (B) depicting antibiotic resistance unique to the survivor phenotype (n=12). LB supplemented with gentamicin (10 μg/ml), kanamycin (50 μg/ml), and tetracycline (10 μg/ml) where indicated.

Figure 4: Isolated alginate concentration determined from carbazole assays depicting

increased alginate production observed from the survivor phenotype (n=10; p≤0.02).

Relative units determined from average of control values with statistical significance

calculated from an unpaired t test with Welch's correction.

Figure 5: A) Direct contact predation assay as depicted in Figure 1 comparing control

(predator unexposed) *P. putida* with freezer stocks of the survivor phenotype (biological

replicates) (n=8; p≤0.0001). B) Modified indirect contact predation assay with prey

introduced 2 cm from predator (n=8; p≤0.01) and time to predator-prey contact recorded

(C). Swarming assays with *C. ferrugineus* plated on prey lawns and swarming diameters

recorded after 7 days (n=8). Statistical significance calculated using an unpaired t test

with Welch's correction.