The spatial-temporal dynamics of respiratory syncytial virus infections

across the east-west coasts of Australia during 2016-17

Mark Robertson[1#], John-Sebastian Eden[2,3#], Avram Levy[4,5], Ian Carter[1], Rachel L Tulloch[2,3],

Elena J Cutmore[2,3], Bethany A Horsburgh[2,3], Chisha T Sikazwe[4,5], Dominic E Dwyer[1,3], David W

Smith[4,6] and Jen Kok[1]

[1]NSW Health Pathology - Institute for Clinical Pathology and Medical Research, NSW Health

Pathology, Westmead Hospital, Westmead, NSW 2145, Australia

[2]Centre for Virus Research, Westmead Institute for Medical Research, Westmead, NSW 2145,

Australia

[3]Marie Bashir Institute for Infectious Diseases and Biosecurity, Sydney Medical School, The

University of Sydney, Sydney, NSW 2006, Australia

[4]PathWest Laboratory Medicine WA, Department of Microbiology, Nedlands, WA 6009,

Australia

[5]School of Biological Sciences, The University of Western Australia, Crawley, WA 6009,

Australia

[6]School of Medicine, The University of Western Australia, Crawley, WA 6009, Australia

[#]These authors contributed equally

Corresponding authors

Dr John-Sebastian Eden

Email: js.eden@sydney.edu.au

Phone: +61 2 8627 1817

Dr Jen Kok

Email: jen.kok@health.nsw.gov.au

Phone: +61 2 8890 6255

Abstract

Respiratory syncytial virus (RSV) is an important human respiratory pathogen. In temperate

regions a distinct seasonality is observed, where peaks of infections typically occur in early

winter, often preceding the annual influenza season. Infections are associated with high rates of

morbidity and mortality, and in some populations exceeds that of influenza. Two subtypes, RSV-

A and RSV-B, have been described, and molecular epidemiological studies have shown that

both viruses mostly co-circulate. This trend also appears to be the case for Australia, however

previous genomic studies have been limited to cases from one Eastern state - New South

Wales. As such, the broader spatial patterns and viral traffic networks across the continent are

not known. Here, we conducted a whole genome study of RSV comparing strains across

eastern and western Australia during the period January 2016 to June 2017. In total, 96 new

RSV genomes were sequenced, compiled with previously generated data, and examined using

a phylodynamic approach. This analysis revealed that both RSV-A and RSV-B strains were

circulating, and each subtype was dominated by a single genotype, RSV-A/ON1-like and RSV-

B/BA10-like viruses. Some geographical clustering was evident in strains from both states with

multiple distinct sub-lineages observed and relatively low mixing across jurisdictions suggesting

that endemic transmission was likely seeded from imported, unsampled locations. Overall, the

RSV phylogenies reflected a complex pattern of interactions across multiple epidemiological

scales from fluid virus traffic across global and regional networks to fine-scale local transmission

events.

Keywords

respiratory syncytial virus, molecular epidemiology, Australia, whole-genome sequencing,

phylogenetics

Introduction

Respiratory syncytial virus (RSV) is a major cause of acute respiratory tract infections in patients of all ages, producing significant morbidity and mortality [1]. The greatest burden of disease is in children under 1 year old, where it is the most common cause of acute respiratory tract infection, and in this age group is second only to malaria as a cause of death globally [2,3]. Importantly, RSV infection in young children may also lead to long-term sequelae such as asthma, chronic bronchitis and obstructive pulmonary disease [2,4]. Other populations particularly impacted by RSV include those over 65, and immunosuppressed patients, such as solid organ and bone marrow transplant recipients [2,4].

Despite being discovered in the 1950s, the burden of RSV disease has only recently been appreciated, which is in part due to more reliable methods of diagnosis and detection. The laboratory diagnosis of RSV was initially reliant on viral isolation and the visualisation of characteristic syncytial cytopathic effects, for which its name is derived [5]. These techniques were slow and required technical expertise. The most commonly used modalities now include rapid antigen (such as lateral flow immunochromatography and fluorescent immunoassays) and nucleic acid amplification tests (NATs). Molecular assays, including both commercial and in-house NATs, offer increased sensitivity and specificity, as well as the ability to be multiplexed to detect other respiratory pathogens such as influenza, parainfluenza, rhinovirus and human metapneumovirus [6]. Rapid test assays can also offer the additional benefit of early diagnosis, which allows for appropriate infection control interventions, rationalisation of unnecessary antibiotic therapy and shorter hospitalisation periods [5]. More generally, improved diagnostics and reporting have begun to shed light on the true incidence, seasonal patterns and peaks of activity of RSV. In temperate regions such as the southern major metropolitan areas of Australia, the seasonal peak typically occurs in the late autumn to early winter period (late-March to mid-August) in the months leading into the influenza season [5,7,8].

RSV can be divided into two antigenically and genetically distinct subtypes, RSV-A and RSV-B. These may be further divided into genetic groups based on the viral glycoprotein (G gene), termed genotypes, with at least 11 and 23 for RSV-A and RSV-B, respectively. However, recent work has proposed shifting to a genotype classification based on whole genome sequencing (WGS) in order to increase phylogenetic resolution [9]. Serological studies have shown that the majority of people are infected by age two, and whilst primary infections are typically more severe they are not protective against repeated infection [2]. The spectrum of

disease severity associated with RSV infection remains an important yet controversial topic.
Studies have made associations between increased severity and particular subtypes/genotypes
[10]; however these are complicated by both host [11] and viral factors [12], as well as their
interactions. Molecular epidemiological studies have shown that both RSV-A and RSV-B co-
circulate during a season, and often at similar levels [13–17]. Furthermore, for each of these
subtypes, a single genotype will tend to predominate, such as with the recent RSV-A ON1-like
and RSV-B BA10-like viruses [7,18,19]. Our understanding of the basic molecular epidemiology
of RSV has been strengthened by WGS, which, similar to other pathogens, is becoming
increasingly common in its application in infectious disease surveillance [7,20,21]. The added
resolution from WGS has been particularly useful for elucidating transmission networks at both
local [22] and epidemiological scales [7], as well as for identifying and classifying the
introduction and spread of new genotypes [9,20].
99
We recently performed the first genome-scale study of RSV molecular epidemiology in
Australia [7], demonstrating a wide diversity of co-circulating RSV lineages, with limited
evidence of strong age and geographical clustering. However, this study was limited to cases
obtained from eastern Australia in New South Wales (NSW) through the Western Sydney Local
Health District. Here, we expand on these initial investigations to perform a transcontinental
study of RSV genomic epidemiology in Australia. We compare RSV strains obtained from
Western Australia (WA) to those from NSW on the east coast over an equivalent time period to
examine the phylogenetic distribution of strains, and to describe viral traffic between these
regions.

Materials and Methods

Sample collection and processing

WGS was conducted on residual RSV-positive specimens collected and tested through routine
testing at diagnostic laboratories in two major Australian diagnostic laboratories. Specimens
were de-identified with basic demographic information collected including age, sex and
postcode, as per protocols approved by local ethics and governance committees
(LNR/17/WMEAD/128). One hundred and three RSV-A and RSV-B positive cases from WA with
a RT-PCR cycle threshold <30 in original screen were collected between January 2016 and
May 2017. Sample aliquots (typically, nasopharyngeal swabs in viral transport medium) were
transferred to the Institute of Clinical Pathology and Medical Research (ICPMR), NSW, and

subsequently extracted using the Qiagen EZ1 Advance XL extractor with the EZ1 Virus Mini Kit

v2.0 (Qiagen, Germany). For the eastern Australian RSV cases, 45 previously untyped RSV-

positive viral extracts from cases collected during early 2017 at the ICPMR were obtained from

storage at -80°C archive. Additional cases from 2016 in NSW were sequenced in a previous

study [7].

Whole genome sequencing

A previously published approach was used to amplify both RSV-A & RSV-B genomes [7]. In

short, RT-PCR was used to amplify four overlapping amplicons (each ~4kb) that together span

the RSV genome. The size and yield of each RT-PCR was determined by agarose gel

electrophoresis, and the four targets pooled equally. The pooled RSV amplicons were then

purified with Agencourt AMpure XP beads (Beckman Coulter, USA) and quantified using the

Quant-iT PicoGreen dsDNA Assay (Invitrogen, USA). The purified DNA was then diluted to 0.25

ng/µL and prepared for sequencing with the Nextera XT DNA library prep kit (Illumina, USA).

Libraries were sequenced on an Illumina MiSeq using a 300 cycle v2 kit (150 nt paired end

reads). Raw paired sequence reads were trimmed using Trim Galore

(https://github.com/FelixKrueger/TrimGalore) and then *de novo* assembled using Trinity [23].

RSV contigs were identified by a local Blastn [24] using a database of RSV reference genomes

from NCBI RefSeq. The trimmed reads were remapped to draft genome contigs using BBMap

(https://sourceforge.net/projects/bbmap/) to check the assembly, and the final majority

consensus was extracted for each sample.

Phylodynamic analysis

The aim of this study was to compare the spatial and temporal dynamics of RSV infection in

Australia. To do this, we used a phylogenetic approach to compare the distribution of strains

circulating in two major, geographically distinct regions – NSW and WA – representing eastern

and western Australia, respectively, during the period January 2016 to June 2017. All WA data,

as well as NSW data from the year 2017, was generated within this study. To ensure even

sampling across sites, these data were combined with NSW data from 2016 that was obtained

from a previous study [7]. To provide additional context, RSV genome data was also sourced

from NCBI GenBank where location (country) and collection date (year) was known. The

Australian and global RSV genomes were first aligned with MAFFT [25], using the FFT-NS-i

algorithm followed by manual inspection of gapped regions particularly in the G gene and non-

coding regions. The alignments were then trimmed to include only the coding regions, and

screened for potential recombinants using RDP4 [26]. The RSV-A alignment included 930

sequences with a length of 13,690 nt, whilst the RSV-B alignment included 602 sequences with

a length of 13,677 nt. The best-fit DNA substitution model for these data was determined with

jModelTest [27]. Maximum likelihood (ML) trees were then estimated for both RSV-A and RSV-

B alignments using RAxML [28] employing the best-fit model, which in both cases was found to

be the General-Time Reversible model with a gamma distribution of rates (GTR+G). Support for

individual nodes was determined by 1000 bootstrap replications.

Phylogeographic clustering

From our overall ML trees, it was found that all sequences from 2016 and 2017 were either

RSV-A ON1-like or RSV-B BA10-like viruses, therefore we limited further analyses to these

specific clusters. To evaluate the geographic structure in the RSV-A and RSV-B trees we used

the Bayesian Tip-association Significance (BaTS) program [29] with 1000 replicates. Using

BaTS we determined the parsimony score (PS), association index (AI), and maximum clade

size (MC) statistics for the location associated with each sequence, specifically focusing on the

comparison between eastern and western Australia. To account for other jurisdictions

(countries) present in the data, such sequences were assigned to their continent of sampling.

This analysis required a posterior distribution of trees, which were obtained using the Bayesian

Markov chain Monte Carlo (MCMC) method implemented in BEAST v1.8.2 [30]. Here, we used

the best-fit DNA model (GTR+G) with a strict clock and a constant population size, as shown to

be appropriate previously [7]. All analyses were run for 50 to 100 million steps with sampling

every 500 steps and 20% burn-in. To ensure convergence, two independent runs were

conducted and merged to obtain the final set of trees used in the clustering analysis.

Local transmission events

In order to identify potential local transmission events (for instance, an outbreak at an

institution), we examined both ML and BEAST trees, for monophyletic groups containing near

identical sequences (>99.9% nucleotide identity) sampled up to two weeks apart. Patient

demographics and sampling location were then mapped to determine traits associated with

these fine-scale phylogenetic groups.

Data availability

All sequences generated in this study have been submitted to NCBI GenBank (MW160744 -

MW160839). Furthermore, data and material relevant to this study are available from

https://github.com/jsede/RSV_NSW_WA.

Results and Discussion

Whole genome sequencing of RSV strains from NSW and WA in 2016 and 2017

In order to compare the spatial distribution of RSV strains across eastern (NSW) and western

(WA) Australia during the period January 2016 to June 2017, we performed WGS on stored

samples collected from community and hospitalised persons presenting with an influenza-like

illness. For the NSW cases, no specific sample selection criteria were used except for sample

availability from an archived collection of RSV positive residual diagnostic specimens. Most

samples from NSW were obtained from children ≤5 years age (52%, n=65/126) with a further

21% of samples collected from patients 65 years (n=26/126). In contrast, samples from WA

were selected to represent all age groups. Despite this, similar to NSW, most available samples

were collected from the young (62%, n=64/103 from patients ≤ 5 years age) and the elderly

(17%, n=18/103 from patients ≥ 65 years), reflecting the distribution of infection at both

extremes of age. The distribution between sexes was approximately equal for both study sites.

Virus genomes were successfully extracted and sequenced from 63 of the 103 respiratory

samples collected in WA during 2016 and 2017, and 32 of the 45 NSW cases collected in 2017

(total new genomes n=96, where one case from NSW was an RSV-A/B mixed infection). These

data were combined with existing genomes from NSW during 2016 (n=93) generated previously

[7], bringing the total number of Australian genomes for analysis across both states in 2016 and

2017 in the current study to 189. The breakdown between sampling location and RSV subtypes

determined by WGS shows that both RSV-A and RSV-B subtypes were present in both NSW

and WA populations (Figure 1). In NSW, there was a higher proportion of RSV-B strains (63%,

n=80/126), which was consistent across the entire study period. Furthemore, in NSW, the peak

in RSV activity typically occurs in May-June each year (NSW Health influenza surveillance

data), however most of our genome sequences in 2016 were from isolates in August to October

that year; that is, they were from the later part of the RSV season. In WA, there was even

representation of RSV-A and RSV-B in the WGS data across 2016 with an increase in the

relative proportion of RSV-A during 2017. While our sampling for WA was intentionally even with

regards to time, subtype, age and setting, the predominance of RSV-A during 2017 was

confirmed by the initial diagnostic screening using an assay that resolves subtype. Specifically,

a transition from RSV-B to RSV-A was observed between the 2016 and 2017 seasons in WA
(data not shown). These results are consistent with other molecular epidemiological studies
globally that show both RSV-A and RSV-B subtypes co-circulate with shifting predominance
across seasons [7,31–33]

Figure 1. Whole-genome sequencing of respiratory syncytial virus (RSV) in Western Australia and New South Wales between January 2016 and June 2017 by month. The y-axis represents counts of genomes sequenced. The x-axis represents months when the samples were collected, with year shown underneath. RSV-A and RSV-B subtypes have been coloured blue and red respectively, as per key provided. In total, over the period of investigation, 189 RSV genomes were compared including 63 from WA and 126 from NSW. WA samples were selected pre-sequencing to cover both subtypes during seasonal and inter-seasonal periods and therefore do not necessarily reflect the subtype or temporal distribution of RSV in WA. The asterisk indicates that of the 126 NSW genomes, 93 genomes from 2016 were generated in a previous study [7].

Phylodynamics of RSV infections in NSW and WA in a global context

The WGS data from NSW and WA were aligned and compared to global genome references to

provide genetic context for local strains. Phylogenetic analysis using a maximum likelihood

approach was then employed to examine the diversity of RSV-A and RSV-B strains (Fig 2). This

analysis revealed that for each subtype, regardless of location, a single genotype was

predominant. For RSV-A, recent viruses from both NSW and WA were derived from the ON1

lineage [18], which has been the predominant RSV-A genotype globally since it first emerged in

2011 [7,18,20,34]. Similarly, the RSV-B phylogeny showed that circulating viruses were mostly

of the BA10 lineage (Fig 2). From the limited available sequence data on GenBank, it appears

that the BA10-like viruses are circulating globally, however, there are few published molecular

epidemiology studies to support the suggestion they are predominant. It is clear however, that in

Australia, in both NSW and WA, that this has been the major RSV-B genotype since at least

2014 [7].

Across both RSV-A and RSV-B phylogenies there does not appear to be strict spatial

clustering when comparing viruses from NSW and WA (Fig 2). That is, the viruses from WA and

NSW do not form monophyletic groups. Such monophyletic clustering would not be expected

based on what is known for other respiratory pathogens - such as influenza - which are

characterised by high levels of gene flow and viral traffic at global scales [35,36]. As such, we

observed multiple co-circulating sub-lineages distributed across the entire diversity of RSV-

A/ON1-like and RSV-B/BA10-like viruses (Fig 2). However, within these sub-lineages the

viruses often formed clusters based on sampling location, therefore, some spatial structure was

apparent in our analysis. To explore this formally, we conducted a clustering analysis using

BaTS, and a posterior set of trees estimated using a time-scaled phylogenetic analysis in

BEAST. We limited the analysis to the RSV-A/ON1 and RSV-B/BA10-like viruses and measured

the degree of clustering based on sampling location (Table 1).

Figure 2. Maximum likelihood phylogeny of respiratory syncytial virus (RSV) strains circulating in Western Australia and New South Wales between January 2016 and June 2017. The phylogenies were first estimated using alignments of complete genome sequences of RSV-A (n=930) and RSV-B (n=602) strain circulating since the late 1970's, however for clarity, only the recently circulating lineages are shown (RSV-A and RSV-B on the left and right panels, respectively). This includes the RSV-A ON1 and RSV-B BA10 lineages, which are defined by specific branches marked with arrows. The tree branches have been coloured by sampling continents as per the key provided. Tip circles coloured grey and white indicate specific sequences sampled in NSW and WA, respectively. Scale bars represent the number of substitutions per site.

262 **Table 1.** Phylogeny-trait association test for RSV in Australia and other regions globally.

Location	RSV-A		RSV-B	
	Seqs (n)	p-value	Seqs (n)	p-value
Asia	13	0.001	8	0.052
Oceania[1]	27	0.001	2	1.000
Europe	0	~	4	1.000
America	137	0.001	3	0.006
NSW	60	0.035	84	0.011
WA	31	0.003	32	0.027
Overall clustering[2]	268	<0.001	133	<0.001

[1]New Zealand only
[2]Determined by Association Index (AI) and Parsimony Score (PS)

When sequences were grouped by broad geographical regions, specifically, continents
and the two Australian states NSW & WA, phylogeny-trait association tests indicated a
significant pattern of overall geographical clustering as measured by AI and PS scores (Table
1). For individual locations, many were found to cluster with significant scores. This includes
Asia, Oceania, America, NSW and WA for RSV-A viruses (all p-values < 0.035) and America,
NSW and WA for RSV-B. For RSV-A, each location was relatively well sampled (number of
sequences between 13 and 137), except that no European sequence was present in the
dataset. For RSV-B, some locations without significant evidence of clustering had low numbers
of sequences (between 2 to 8). This analysis demonstrates that, at larger epidemiological
scales, viral lineages are often imported and established locally. We note however, that this
analysis is biased for regions that have been highly sampled, as highlighted by the American
RSV-A viruses, which are mostly from one specific city and hospital (Supplementary Fig 1). Our
sampling of NSW and WA strains is similarly constrained, except that our sampling sites are
both major diagnostics hubs which perform testing across wide geographic regions that are
representative of large and diverse populations [7], and in our comparison the period of
sampling was equivalent. Despite this and any apparent geographical clustering, the
phylogenies both reflect a lack of data from many jurisdictions globally, and our insights into
global and local viral traffic will remain hampered until these genome sequences become
available.

Lineages and local clusters

A detailed examination of individual lineages (Supplementary Figs 1 & 2) confirmed that most
only contained viruses from either state. There were some instances where individual viruses
from NSW were nested within diversity solely comprised of WA strains, and vice-versa, and
which may represent viral migration events between the jurisdictions. The overall pattern
suggests low levels of mixing between NSW and WA, perhaps due to the large geographic area
of each state and the distance that separates them. Moreover, we had suggested previously
that some evidence of viral persistence across seasons was present in NSW. Here, we found
no direct link between the 2016 and 2017 seasons in both jurisdictions and that viral diversity is
likely maintained and seeded from imported yet unsampled locations. This is consistent with
other studies that have shown that local persistence makes a minimal contribution to the
diversity of RSV strains in any given area [37,38]. Rather, viral importation and global mixing of
strains seem to be the major driver of RSV diversity and their sources. Despite the data
suggesting limited east-west viral traffic across the Australian continent, the Sydney-Perth air
route remains one of the busiest in the country, nor can we discount other regions within
Australia such as the tropical North, where persistence of viral lineages may occur as important
sources of viral diversity. Indeed, from our understanding of influenza virus in Australia, the
synchronized dissemination of viral strains across the country is driven by both multiple
introductions from the global population and strong domestic connectivity [39]. Increasing the
breadth of WGS data for RSV cases nationally would assist in confirming what domestic viral
traffic networks exist, particularly between tropical and temperate regions, as well as, between
major population centres on the east coast such as Sydney and Melbourne.

Next, we examined genetic clusters defined by high genetic identity (>99.9%) and similar
sampling periods (collection dates within two weeks) to consider what features such as patient
age and sampling location define them. For RSV-A, we identified 8 clusters (Supp Figure 1 &
Table 1) and for RSV-B we identified 10 clusters (Supp Figure 2 & Table 2). In this study 75%
(n=6/8) and 80% (n=8/10) of RSV-A and RSV-B clusters, respectively could be linked to
common localities and/or institutions, including individual hospital wards or emergency
departments. While two of the RSV-B clusters were found to be repeat samples of the same
patient (Supp Table 2), overwhelmingly, these clusters of genetically and temporal related
sequences most likely represent fine-scale transmission events.

Conclusions

In summary, our analysis has identified a number of features of RSV epidemiology and patterns of spread including i) the co-circulation of both major subtypes: RSV-A and RSV-B; ii) a single genotype for each subtype predominates each season; iii) multiple distinct sub-lineages of each genotype will co-circulate and which are associated with regional and local clustering/outbreaks, iv) little viral mixing across the east-west coasts of Australia despite apparent overall geographic clustering and v) that genetically and temporal related sequences most likely represent fine-scale transmission events such as institutional outbreaks. Taken together, this presents a complex phylodynamic pattern with globally circulating diversity with viral mixing across different regions, yet, finer-scale patterns revealing multiple endemic sub-lineages and clusters consistent with local transmission events and outbreaks. This highlights the connections of genomic data across multiple epidemiological scales and further strengthens the need for much greater sampling of RSV, not just here in Australia, but globally.

Acknowledgments

We thank all the members of the Virology and Microbiology teams at NSW Health Pathology - ICPMR in Westmead, NSW and the PathWest QE2 Medical Centre laboratories in Perth, WA for all their contributions towards processing the diagnostics specimens used in this study.

Funding

Funding was provided through the ICPMR Private Practice Trust fund, the National Health and Medical Research Council Centre of Research Excellence in Emerging Infectious Diseases (1102962) and the Marie Bashir Institute for Infectious Diseases and Biosecurity at the University of Sydney.

Conflicts of Interest

None

References

1. Shi T, McAllister DA, O'Brien KL, Simoes EAF, Madhi SA, Gessner BD, et al. Global, regional, and national disease burden estimates of acute lower respiratory infections due to respiratory syncytial virus in young children in 2015: a systematic review and modelling study. Lancet. 2017;390: 946–958. doi:10.1016/S0140-6736(17)30938-8

2. Griffiths C, Drews SJ, Marchant DJ. Respiratory Syncytial Virus: Infection, Detection, and New Options for Prevention and Treatment. Clin Microbiol Rev. 2017;30: 277–319. doi:10.1128/CMR.00010-16

3. Hall CB, Weinberg GA, Iwane MK, Blumkin AK, Edwards KM, Staat MA, et al. The burden of respiratory syncytial virus infection in young children. N Engl J Med. 2009;360: 588–598. doi:10.1056/NEJMoa0804877

4. Beigelman A, Bacharier LB. Early-life respiratory infections and asthma development: role in disease pathogenesis and potential targets for disease prevention. Curr Opin Allergy Clin Immunol. 2016;16: 172–178. doi:10.1097/ACI.0000000000000244

5. Henrickson KJ, Hall CB. Diagnostic assays for respiratory syncytial virus disease. Pediatr Infect Dis J. 2007;26: S36–40. doi:10.1097/INF.0b013e318157da6f

6. Mahony J, Chong S, Merante F, Yaghoubian S, Sinha T, Lisle C, et al. Development of a respiratory virus panel test for detection of twenty human respiratory viruses by use of multiplex PCR and a fluid microbead-based assay. J Clin Microbiol. 2007;45: 2965–2970. doi:10.1128/JCM.02436-06

7. Di Giallonardo F, Kok J, Fernandez M, Carter I, Geoghegan JL, Dwyer DE, et al. Evolution of Human Respiratory Syncytial Virus (RSV) over Multiple Seasons in New South Wales, Australia. Viruses. 2018;10. doi:10.3390/v10090476

8. Yeoh DK, Foley DA, Minney-Smith CA, Martin AC, Mace AO, Sikazwe CT, et al. The impact of COVID-19 public health measures on detections of influenza and respiratory syncytial virus in children during the 2020 Australian winter. Clin Infect Dis. 2020. doi:10.1093/cid/ciaa1475

9. Ramaekers K, Rector A, Cuypers L, Lemey P, Keyaerts E, Van Ranst M. Towards a unified classification for human respiratory syncytial virus genotypes. Virus Evol. 2020;6: veaa052. doi:10.1093/ve/veaa052

10. Vandini S, Biagi C, Lanari M. Respiratory Syncytial Virus: The Influence of Serotype and Genotype Variability on Clinical Course of Infection. Int J Mol Sci. 2017;18. doi:10.3390/ijms18081717

11. Tal G, Mandelberg A, Dalal I, Cesar K, Somekh E, Tal A, et al. Association between common Toll-like receptor 4 mutations and severe respiratory syncytial virus disease. J Infect Dis. 2004;189: 2057–2063. doi:10.1086/420830

12. DeVincenzo JP, El Saleeby CM, Bush AJ. Respiratory syncytial virus load predicts disease severity in previously healthy infants. J Infect Dis. 2005;191: 1861–1868. doi:10.1086/430008

13. Cattoir L, Vankeerberghen A, Boel A, Van Vaerenbergh K, De Beenhouwer H.
Epidemiology of RSV and hMPV in Belgium: a 10-year follow-up. Acta Clin Belg. 2019;74:
229–235. doi:10.1080/17843286.2018.1492509

14. Otieno JR, Kamau EM, Agoti CN, Lewa C, Otieno G, Bett A, et al. Spread and Evolution of
Respiratory Syncytial Virus A Genotype ON1, Coastal Kenya, 2010-2015. Emerg Infect Dis.
2017;23: 264–271. doi:10.3201/eid2302.161149

15. Otieno JR, Kamau EM, Oketch JW, Ngoi JM, Gichuki AM, Binter Š, et al. Whole genome
analysis of local Kenyan and global sequences unravels the epidemiological and molecular
evolutionary dynamics of RSV genotype ON1 strains. Virus Evol. 2018;4: vey027.
doi:10.1093/ve/vey027

16. Pangesti KNA, Abd El Ghany M, Walsh MG, Kesson AM, Hill-Cawthorne GA. Molecular
epidemiology of respiratory syncytial virus. Rev Med Virol. 2018;28. doi:10.1002/rmv.1968

17. Park E, Park PH, Huh JW, Yun HJ, Lee HK, Yoon MH, et al. Molecular and clinical
characterization of human respiratory syncytial virus in South Korea between 2009 and
2014. Epidemiol Infect. 2017;145: 3226–3242. doi:10.1017/S0950268817002230

18. Eshaghi A, Duvvuri VR, Lai R, Nadarajah JT, Li A, Patel SN, et al. Genetic variability of
human respiratory syncytial virus A strains circulating in Ontario: a novel genotype with a
72 nucleotide G gene duplication. PLoS One. 2012;7: e32807.
doi:10.1371/journal.pone.0032807

19. Pretorius MA, van Niekerk S, Tempia S. Replacement and positive evolution of subtype A
and B respiratory syncytial virus G-protein genotypes from 1997–2012 in South Africa. The
Journal of. 2013. Available: https://academic.oup.com/jid/article-
abstract/208/suppl_3/S227/789685

20. Agoti CN, Otieno JR, Gitahi CW, Cane PA, Nokes DJ. Rapid spread and diversification of
respiratory syncytial virus genotype ON1, Kenya. Emerg Infect Dis. 2014;20: 950–959.
doi:10.3201/eid2006.131438

21. Dapat IC, Shobugawa Y, Sano Y, Saito R, Sasaki A, Suzuki Y, et al. New genotypes within
respiratory syncytial virus group B genotype BA in Niigata, Japan. J Clin Microbiol. 2010;48:
3423–3427. doi:10.1128/JCM.00646-10

22. Agoti CN, Otieno JR, Munywoki PK, Mwihuri AG, Cane PA, Nokes DJ, et al. Local
evolutionary patterns of human respiratory syncytial virus derived from whole-genome
sequencing. J Virol. 2015;89: 3444–3454. doi:10.1128/JVI.03391-14

23. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-length
transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol.
2011;29: 644–652. doi:10.1038/nbt.1883

24. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J
Mol Biol. 1990;215: 403–410. doi:10.1016/S0022-2836(05)80360-2

25. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol. 2013;30: 772–780.
doi:10.1093/molbev/mst010

26. Martin DP, Murrell B, Golden M, Khoosal A, Muhire B. RDP4: Detection and analysis of
recombination patterns in virus genomes. Virus Evol. 2015;1: vev003.
doi:10.1093/ve/vev003

27. Darriba D, Taboada GL, Doallo R, Posada D. jModelTest 2: more models, new heuristics
and parallel computing. Nat Methods. 2012;9: 772. doi:10.1038/nmeth.2109

28. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics. 2014;30: 1312–1313. doi:10.1093/bioinformatics/btu033

29. Parker J, Rambaut A, Pybus OG. Correlating viral phenotypes with phylogeny: accounting
for phylogenetic uncertainty. Infect Genet Evol. 2008;8: 239–246.
doi:10.1016/j.meegid.2007.08.001

30. Drummond AJ, Suchard MA, Xie D, Rambaut A. Bayesian phylogenetics with BEAUti and
the BEAST 1.7. Mol Biol Evol. 2012;29: 1969–1973. doi:10.1093/molbev/mss075

31. Yun KW, Choi EH, Lee HJ. Molecular epidemiology of respiratory syncytial virus for 28
consecutive seasons (1990-2018) and genetic variability of the duplication region in the G
gene of genotypes ON1 and BA in South Korea. Archives of Virology. 2020. pp. 1069–
1077. doi:10.1007/s00705-020-04580-z

32. Razanajatovo Rahombanjanahary NH, Rybkina K, Randriambolamanantsoa TH,
Razafimanjato H, Heraud J-M. Genetic diversity and molecular epidemiology of respiratory
syncytial virus circulated in Antananarivo, Madagascar, from 2011 to 2017: Predominance
of ON1 and BA9 genotypes. J Clin Virol. 2020;129: 104506. doi:10.1016/j.jcv.2020.104506

33. Luo H-J, Huang X-B, Zhong H-L, Ye C-X, Tan X, Zhou K, et al. Epidemiological
characteristics and phylogenic analysis of human respiratory syncytial virus in patients with
respiratory infections during 2011-2016 in southern China. Int J Infect Dis. 2020;90: 5–17.
doi:10.1016/j.ijid.2019.10.009

34. Tabatabai J, Prifert C, Pfeil J, Grulich-Henn J, Schnitzler P. Novel respiratory syncytial virus
(RSV) genotype ON1 predominates in Germany during winter season 2012-13. PLoS One.
2014;9: e109191. doi:10.1371/journal.pone.0109191

35. Rambaut A, Pybus OG, Nelson MI, Viboud C, Taubenberger JK, Holmes EC. The genomic
and epidemiological dynamics of human influenza A virus. Nature. 2008;453: 615–619.
doi:10.1038/nature06945

36. Vijaykrishna D, Holmes EC, Joseph U, Fourment M, Su YCF, Halpin R, et al. The
contrasting phylodynamics of human influenza B viruses. Elife. 2015;4: e05055.
doi:10.7554/eLife.05055

37. Agoti CN, Otieno JR, Ngama M, Mwihuri AG, Medley GF, Cane PA, et al. Successive
Respiratory Syncytial Virus Epidemics in Local Populations Arise from Multiple Variant
Introductions, Providing Insights into Virus Persistence. J Virol. 2015;89: 11630–11642.
doi:10.1128/JVI.01972-15

38. Zou L, Yi L, Wu J, Song Y, Huang G, Zhang X, et al. Evolution and Transmission of
Respiratory Syncytial Group A (RSV-A) Viruses in Guangdong, China 2008–2015. Front
Microbiol. 2016;7: 1263. doi:10.3389/fmicb.2016.01263

39. Geoghegan JL, Saavedra AF, Duchêne S, Sullivan S, Barr I, Holmes EC. Continental
synchronicity of human influenza virus epidemics despite climactic variation. PLoS Pathog.
2018;14: e1006780. doi:10.1371/journal.ppat.1006780