Newly identified proviruses in Thermotogota suggest that viruses are

the vehicles on the highways of interphylum gene sharing.

Running title: Viruses and Proviruses of Thermotogota.

Thomas H. A. Haverkamp[a#], Julien Lossouarn [b], Olga Zhaxybayeva[c], Jie Lyu[d], Nadège

Bienvenu[d], Claire Geslin[d] and Camilla L. Nesbø[e,f]

[a] Centre for Ecological and Evolutionary Synthesis, Department of Biosciences,

University of Oslo, Norway.

[b] Université Paris-Saclay, INRAE, AgroParisTech, Micalis Institute, 78350,

Jouy-en-Josas, France

[c] Department of Biological Sciences, Dartmouth College, Hanover, NH, USA.

[d] Université Brest, CNRS, IFREMER, Laboratoire de Microbiologie des Environnements

Extrêmes, F-29280 Plouzané, France

[e] Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada

[f] Department of Chemical Engineering and Applied Chemistry,

University of Toronto, Toronto, ON, Canada, M5S 3E5

[#]Present address: Department of Epidemiology, Norwegian Veterinary Institute, Oslo,

Norway.

ABSTRACT

Phylogenomic analyses of bacteria from the phylum Thermotogota have shown extensive

lateral gene transfer (LGT) with distantly related organisms, particularly with Firmicutes.

One likely mechanism of such DNA transfer is viruses. However, to date only three

temperate viruses have been characterized in this phylum, all infecting bacteria from the

Marinitoga genus. Here we report 17 proviruses integrated into genomes of eight

Thermotogota genera and induce viral particle production from one of the proviruses. The

proviruses fall into two groups based on sequence similarity, gene synteny and taxonomic

classification. Proviruses of one group are found in six genera and are similar to the

previously identified *Marinitoga* viruses, while proviruses from the second group are

only distantly related to the proviruses of the first group, have different genome

organization and are found in only two genera. Both groups are closely related to

Firmicutes in genomic and phylogenetic analyses, and one of the groups show evidence

of very recent LGT and are therefore likely capable of infecting cells from both phyla.

We conjecture that viruses are responsible for a large portion of the observed gene flow

between Firmicutes and Thermotogota.

Introduction

The phylum Thermotogota comprises anaerobic fermentative bacteria, most of which are thermophiles [1] . They are common in subsurface environments such as marine vents, terrestrial hot springs and deep subsurface oil reservoirs [2–5]. On phylogenetic trees of 16S rRNA gene, Thermotogota are usually a deep branching bacterial lineage, while ribosomal proteins and other markers do not always agree with that placement [6, 7]. Such discrepancies are likely due to lateral gene transfer (LGT), which has been an important force shaping the genomes of Thermotogota, with Firmicutes and Archaea being their most notable gene transfer partners [1, 7, 8]. The LGT between Firmicutes and Thermotogota is so extensive that the two phyla have been suggested to be linked by ''highways of gene sharing'' [7]. However, how these inter-phylum gene-sharing events occur is still unclear.

The subsurface constitutes the largest biosphere on Earth and is estimated to contain ~70% of all cells [9]. Viruses are likely to be particularly important in subsurface environments, since 97% of all viruses on earth being found in soil and sediments [10, 11]. Moreover, although both prokaryotic cell and virus numbers decrease with depth, the virus-to-cell ratio increases with depth [10, 12, 13]. Phylogeographic studies of hyperthermophilic *Thermotoga* and mesophilic *Mesotoga* have revealed genetic interaction between geographically distant populations, particularly among the hyperthermophilic *Thermotoga* [3, 5]. Viruses are one potential source of such long-distance dispersal of genetic material [14], especially for anaerobic organisms where surface dispersal is problematic.

Although viruses are likely candidates for transferring DNA both within and

between species, only three temperate siphoviruses (MCV1, MCV2, and MPV1), all

infecting one Thermotogota genus, *Marinitoga*, have been described [15, 16]. MCV1 and

MCV2 infect *Marinitoga camini* strains isolated from deep-sea hydrothermal vents [16].

MPV1 infects the deep-sea marine vent bacterium *Marinitoga piezophila*, where it is

highjacked by a plasmid co-occurring in the same host, illustrating the potential route of

gene mobilization in these ecosystems [15]. The three viruses are found as proviruses in

their host genomes and show similar genomic organization and virion morphology.

Phylogenetic and protein sequence-similarity analyses of the viral ORFs revealed that

they often group either with Firmicutes or Firmicutes' viruses, which suggests that

viruses infecting members of Firmicutes and Thermotogota phyla share a common gene

pool [15, 16].

Here we report 17 additional proviruses in Thermotogota genomes from eight

Thermotogota genera, and a successful induction of one of these proviruses. The

identified proviruses fall into two distinct groups. Both groups are closely related to

Firmicutes viruses, and the proviruses from one of these groups are likely able to infect

cells from both phyla. We hypothesize that membrane transport proteins, such as ABC

transporters, serve as receptors for Thermotogota viruses. We propose a mechanism that

could account for the highways of gene sharing observed between Thermotogota and

Firmicutes, where LGT of viral genes encoding transmembrane proteins may make the

host vulnerable to new viruses.

Material and Methods

Prediction and taxonomic classification of proviruses and functional annotation of their ORFs

One hundred eleven Thermotogota genomes were downloaded from either Genbank or IMG [17] prior to June 2018. For draft genomes, the contigs were combined into 'artificially closed' genome using the "union" command from the EMBOSS package (version 6.6.0) [18]. Each genome was screened for the presence of proviruses using the Prophinder web server [19], PHAST web server [20], and PhiSpy (version 2.3) [21] between October 2014 and July 2018 (**Supplementary Table S1**). For artificially closed genomes, the proviral regions that crossed contig borders were discarded. Putative provirus regions were inspected to identify the most likely provirus sequence by 1: looking at annotations, 2: identifying possible flanking tRNA genes and 3: comparing the region to genomes from the same genus and defining the boundaries, to ensure that flanking genomic regions present in closely related genomes without provirus were not included. Proviruses were considered complete if they contained modules for lysogeny, replication, packaging, head/tail morphogenesis and lysis. If one of these modules were missing, the provirus was scored as incomplete.

To see if there were close relatives of the predicted proviruses in other genomes, provirus ORFs were used as queries in BLASTX (ver.2.2.26) [22] searches of the NCBI non-redundant (nr) database [23] (accessed between July 2018 and March 2020). When homologs for multiple ORFs from the predicted provirus were found in the same distantly related subject genome (usually a Firmicutes genome), the identified genome was downloaded and aligned to the Thermotogota genome carrying the provirus using Progressive Mauve [24]. This resulted in the identification of a local alignment covering

similar proviruses in two otherwise distantly related genomes. The aligned region was

used to determine the boundaries of the provirus, limiting the provirus ends to the ends of

the alignment.

Proviral-ORF annotations were obtained from their respective Genbank entries

and supplemented by results from BLASTP searches [22] of the *nr* database with an

expected-value cutoff of 10^{-1}, and from HHpred searches [25] of the PDB database [26]

with a probability cutoff of 99%. In addition, recombinase- and terminase-encoding

ORFs were annotated using InterProScan [26], as implemented in Geneious v.10

(Biomatters Ltd.).

The sequences of the predicted proviruses were compared to each other using

BLASTN and TBLASTX (ver.2.2.26) [22] and visualized using genoPlotR [27] and

Circos [28]. Taxonomic classification of the provirus genomes was carried out using

searches of NCBI's viral RefSeq database (v. 94) as implemented in VContact2 using

Diamond [29] to identify viral protein clusters and ClusterONE [30] to obtain virus

clusters [31, 32]. Taxonomic classification was also assessed with Virfam [33], VIRIDIC

[34] and VIPTree [35]. Morphological classification was obtained with Virfam [33].

Inference of potential host range of the putative Thermotogota viruses. A database

containing all proteins from 59 Thermotogota genomes without identified proviruses was

constructed in Geneious v.10. Translated Thermotogota provirus proteins were used as

queries in BLASTP searches of this database. The provirus genes were scored as present

in the Thermotogota genomes if the query protein had a match with > 50% amino acid

identity and > 60% coverage.

CRPISPR spacer sequences from 90 Thermotogota genomes were obtained from

IMG [17]. The spacers were mapped to the provirus genomes in Geneious v.10, allowing

upto 10% nucleotide mismatches.

Phylogenetic analyses of provirus genes and candidate receptor genes. Homologs of

provirus genes selected for phylogenetic analysis were obtained by searching each

translated proviral gene against *nr* database (accessed between December 2019 and

March 2020), as well as a local database of all the *Thermotogota* virus proteins identified,

using BLASTP (version 2.2.26), with E-value cutoff of 10^{-1}. The 20 top-scoring matches

from each database were retrieved and aligned using MAFFT v. 7.450 with the G-INS-I

option [36]. Identical sequences and highly similar sequences from the same genus were

removed. Alignment positions with > 50% gaps were trimmed. Phylogenetic trees were

reconstructed using RAxML [37] with WAG+G substitution model with four rate

categories and 100 bootstrap replicates, as implemented in Geneious v.10.

Candidate receptor proteins in genomes of *Petrotoga* sp. 8T1HF07.NaAc.6.1,

Petrotoga olearia, *Petrotoga mobilis*, *Petrotoga* sp. 9T1HF07.CasAA.8.2, *Defluviitoga*

tunisiensis, *Lacticigenium naphtae*, and *Mahella australiensis*, which had proviruses

assigned to Group 2 (see the Results section for definition), were identified in IMG using

an amino acid identity cut-off of 50%. Homologs in *Geosporobacter ferrireducens*

genome, which was not available in IMG, were identified using BLASTP search with E-

value cutoff of 10^{-10}. Collection of additional homologs and phylogenetic analyses were

carried out as described above.

Phylogenetic analysis of single copy gene in Thermotogota genomes available in

Genbank (accessed May 27 2020) were done using the GToTree pipeline [38] with the

Bacterial hmm-set of 74 target genes. The resulting alignment was imported into

Geneious Prime 2020.1.2 where sites with more than 50% gaps were removed, giving an

alignment of 11,003 amino acid positions. The phylogenetic tree was reconstructed using

FastTree with the JTT model with optimized Gamma20 likelihood [39].

Virus induction and electron microscopy. *T. africanus* H17ap6033 and two *Petrotoga*

isolates, *P.olearia* and *Petrotoga* sp. 8T1HF07.NaAc.6.1, were cultivated in a modified

Ravot medium as previously described [15] at 65°C and 55°C, respectively. Attempts

were made to increase the viral production of the strains by using mitomycin C, as

reported previously [15, 16]. A final concentration of 5 µg/mL of mitomycin C was

added to 300 mL bacterial culture at early to mid-log growth phase. After 3 hours of

incubation with mitomycin C, cultures were centrifuged at 7500 rpm and 4°C for 15 min,

and supernatants were ultracentrifuged at 37 000 rpm (~100 000 g) and 10°C for 1h

(Beckman Optima LE-80 K; rotor 70.1.Ti). Pellets were resuspended in 100µL of buffer

(10 mM Tris-HCL, 100 mM NaCl, 5 mM $CaCl_2$, 20 mM $MgCl_2$) and suspensions were

prepared for negative staining electron microscopy as previously described [40]. Briefly,

5 µL of the suspensions were directly spotted onto a Formwar carbon coated copper grid.

Putative virus-like particles were allowed to adsorb to the carbon layer for 2 min and

excess of liquid was removed. 5 µL of a staining uranyl acetate solution (2%) was then

spotted to the grid for 45 s and excess of liquid was removed again. The grid was imaged

at 120 kV in a JEOL JEM 100 CXIIVR transmission electron microscope.

Results

Newly identified Thermotogota proviruses come from two distinct viral

lineages. Analysis of 111 Thermotogota genomes identified 20 proviruses, including the

three already characterized viruses from *Marinitoga* [15, 16] and four likely partial

proviruses (**Supplementary Table S1** and **Supplementary Table S2**). One of the 20

proviruses is present with 100% nucleotide sequence identity in all six available

Thermosipho melanesiensis genomes [41], and therefore is counted as just one novel

provirus. An additional provirus (MLaV1) was reported in a *Marinitoga lauensis* genome

after we completed the screening [42]. Due to its similarity to proviruses identified in

other *Marinitoga* genomes, it was not included in our further analyses.

The predicted proviruses can be divided into two distinct groups, hereafter

denoted as Group 1 (15 proviruses: 13 complete and 2 incomplete) and Group 2 (5

proviruses: 3 complete and 2 incomplete). First, the genome organization differs between

the proviruses in two groups (**Fig. 1** and **Fig. 2**). Second, the genes within each group are

more similar than the genes between groups (**Fig. 3, Supplementary Table S2**). Third,

the two groups form separate clusters in the VContact2 network (**Supplementary Fig.**

S1, panel A). Finally, the two groups show up as different clades on the Viral Proteomic

Tree (**Supplementary Fig. S2**). Group 1 proviruses are found in the genera *Marinitoga*,

Thermosipho, *Kosmotoga*, *Mesotoga*, *Geotoga* and *Mesoaciditoga*, while Group 2

proviruses are limited to the genera *Petrotoga* and *Defluviitoga* (**Supplementary Fig. S1,**

panel B). However, presence of 29 protein families shared between the two groups

(Supplementary Table S3) suggests that LGT may occur between the viruses of the two

groups.

Classification and genomic features of Group 1 proviruses.

All complete Group 1 proviruses are likely to encode siphoviruses based on their head,

neck and tail gene sequences [33], and the morphology observed for the earlier

characterized MPV1, MCV1, MCV2 viruses and the TAV1 virus induced in the current

study (see below).

None of the proviruses have significant nucleotide identity with viral genomes in

the NCBI nr/nt database. Following taxonomic criteria, where viruses with > 50-70%

nucleotide identity over the full genome belong to the same genus and viruses with >

95% nucleotide identity belong to the same species [43, 44], the complete proviruses are

assigned to 13 new species and at least 11 new genera (**Supplementary Table S4**). The

sequence similarity suggests that the closest relatives of the Group 1 proviruses are

Firmicutes' viruses (**Supplementary Fig. S1, panel A**), since 15% of the provirus genes

have Firmicutes as the top-scoring match, if members of the proviruses' host genus are

excluded (**Supplementary Table S5**).

The proviruses have the same modular structure as genomes of the earlier

described MPV1, MCV1 and MCV2 viruses [15, 16] (**Fig. 1**). The 5' module contains

genes involved in lysogeny and is encoded on the opposite strand compared to the rest of

the virus genes. The lysogeny module is followed by modules for replication, packaging,

morphogenesis and host lysis. Similar to the described *Marinitoga* viruses [15, 16], the

gene content of lysogeny module of all examined proviruses is very variable, with only

the recombinase gene conserved (**Fig. 1**).

All Group 1 proviruses are inserted next to a tRNA gene. Eight of them (the

Marinitoga proviruses MPV1, MCV1, MCV2, MHV1 and M1137V2; the *Kosmotoga*

pacifica provirus KPV1, *Geotoga petrae* provirus GPV1, and *Mesoaciditoga lauensis*

provirus MLV1), are inserted next to the tRNA-Glu gene, and carry similar site-specific

DNA serine recombinases (KEGG Orthology; K06400, homologs of Marpi_0291 in

MPV1 from *M. piezophila*) (**Fig. 4**). The *Thermosipho* proviruses TMV1 and T1074V1

carry more distant homologs of this recombinase (**Fig. 4**) and are inserted next to the

tRNA-Phe gene. The most divergent homolog of the serine recombinase is present in

TAV1, which is inserted next to the tRNA-Pro gene. MTOLDCV1 is also inserted next to

a tRNA-Pro gene, but it is located at the end of a contig in an incomplete single cell

genome and its 5' end (where recombinase would be found) is missing (the *Mesotoga* sp.

SC_TOLDC recombinase included in **Fig. 4** is located on a separate contig). We

hypothesize that these recombinases are integrases that specifically recognize the tRNA

genes next to which the provirus is inserted (i.e., the tRNA-Glu, tRNA-Phe, or tRNA-Pro

genes). The remaining three proviruses (M1135V1, M1138V1, and M1137V1) are

inserted next to tRNA-Cys gene and may also use similar integration mechanism, but

these proviruses do not have the detectable serine recombinase homologs. M1135V1 and

M1138V1 have homologous ORFs of unknown function in the recombinase gene

position (**Fig. 1**), which may or may not provide this function.

Another typical viral protein that shows variation across the Group 1 proviruses is

the large subunit of the terminase protein involved in the packaging of viral DNA into the

virus particle [45]. The proviruses carry three types of these proteins (BLASTP E-value

cut-off < 0.01, identity < 25%, **Supplementary Fig. S3**). The first type, exemplified by

the protein in MPV1 (Marpi_0320), contains a PBSX family domain and has homologs in

MCV1 and MCV2 and the proviruses in *Marinitoga* sp. 1135, *Marinitoga* sp. 1138,

Thermosipho sp. 1074 and *M. lauensis*. The second, exemplified by the terminase in

TAV1 (H17ap60334_04902), contains a 'Terminase_lsu_T4-like' domain and has

homologs in the proviruses from *K. pacifica*, *T. melanesiensis*, *Mesotoga* sp. TolDC and

G. petrae. The third type is found in the proviruses in *M. hydrogenitolerans* and

Marinitoga sp. 1137 (BUA62_RS02495, LN42_01905 and LN42_00550). These

terminases also contain a 'Terminase_lsu_T4-like' domain and are distant homologs of

the second terminase type.

In addition to the recombinase and terminase, other typical viral proteins such as

tail tape measure, capsid and portal proteins were identified, but did not always show

detectable similarity among the proviruses (**Fig. 1**). Two transcription regulators

(Marpi_0297 and Marpi_0298 in MPV1), a DNA repair exonuclease (Marpi_0340 in

MPV1) and a single stranded DNA-binding protein (Marpi_0306 in MPV1), show

relatively high (32-100%) identity across most Group 1 proviruses (**Fig. 1,**

Supplementary Table S4). Genes encoding two hypothetical proteins (homologs of

Marpi_0299 and Marpi_0338 in MPV1) are shared among 10 of the proviruses (36-96%

identity), suggesting these genes may provide important viral functions.

Broad host range of Group 1 proviruses.

Detection of Group 1 proviruses in the genera *Marinitoga*, *Thermosipho*, *Kosmotoga*,

Mesotoga, *Geotoga* and *Mesoaciditoga* (**Supplementary Table S1**), suggests that the

Group 1 viruses are widespread among Thermotogota, particularly among organisms

inhabiting hydrothermal vents. Such wide distribution and relatively high sequence

identity among the proviral genomes (**Fig. 1, Supplementary Table S4**) suggest that the

Group 1 temperate viruses might have broad host ranges. Experiments showing that

MPV1 from *M. piezophila* can infect and transfer a plasmid to a *Thermosipho* isolate is

consistent with this hypothesis [15].

Further support comes from mapping of CRISPR spacer sequences from 90

Thermotogota genomes to the Group 1 proviruses. Five of the 17 proviruses matched

CRISPR spacers in the genomes from a different genus (**Table 1**). For example, the

Thermosipho provirus TAV1 had 35 matches to spacers in the genomes of

Pseudothermotoga and *Thermotoga* spp. Anecdotal evidence corroborates an ability of

TAV1 to infect *Thermotoga* spp. Back in 2005, when the sample from the Hibernia oil

reservoir containing TAV1 and its host, *T. africanus* H17ap60334, was being processed

by one of us (Camilla L. Nesbø) in the laboratory, *Thermotoga* isolates from Troll oil

reservoir in the North Sea, which were at the same time being transferred to fresh media,

experienced a mass death. Analysis of the genomes of the surviving *Thermotoga* isolates

[3] revealed presence of three CRISPR spacer matching the TAV1 genome

(**Supplementary Fig. S4**). These spacers were located in the middle of the CRISPR

arrays, indicating that they were not new acquisitions [46]. Therefore the only surviving

isolates of *Thermotoga* must have had already experienced and survived TAV1 or related

virus infections in the oil reservoir.

Classification, genomic features and distribution of Group 2 proviruses.

Group 2 consists of three complete proviruses in the genomes of *Petrotoga* sp. 8T1HF07

(P8T1HF07V1), *Petrotoga olearia* (POV1) and *Defluviitoga tunesiensis* (DTV1) (**Fig. 2**),

and two incomplete proviruses in the genomes of *Petrotoga mobilis* SJ95 and *Petrotoga*

sp. 9T1HF07 (**Supplementary Fig. S5**). Due to the short length of the incomplete

proviruses, they were not included in the remaining analyses of this section.

Following the taxonomic classifications criteria described above, the three

complete proviruses P8T1HF07V1, DTV1 and POV1 are assigned to three new viral

species (**Supplementary Table S4**). Based on the head-neck-tail module classification

[33], these proviruses likely encode siphoviruses of Type1 - Cluster 2. All hosts of the

previously described members of this *Siphoviridae* lineage belong to Firmicutes. In

agreement with this, similarity searches revealed that these proviruses show very high

similarity to proviruses of three Firmicutes genomes: *Lacticigenium naphta*e (LNV1),

Geosporobacter ferrireducens (GFV1) and *Mahella australiensis* (MAV1) (**Fig. 2,**

Supplementary Table S4).

The sequences and genome organization of the three complete Group 2 proviruses

differ considerably from that of Group 1 (**Fig. 1** and **Fig. 2**). These proviruses are also not

located next to tRNA genes. The 5' module encodes genes involved in virus replication

and transcription, and the comparative genomic analysis shows high level of diversity in

this region (**Fig. 2**). This module is followed by highly conserved packaging,

morphogenesis and lysis modules. The lysogeny module is located at the 3' end of the

virus. The site-specific serine recombinases carried by the Group 2 proviruses in this

module are distant homologs of the earlier discussed Group 1 recombinases (**Fig. 4**).

When comparing the Group 2 provirus genomes from Thermotogota and

Firmicutes, each Thermotogota provirus is more similar to a Firmicutes provirus than to

other Thermotogota proviruses **(Fig. 2, Supplemental Table S4).** Alignments of the two

Thermotogota-Firmicutes provirus pairs, P8T1HF07V1 and GFV1, and DTV1 and

MAV1, have 66.7 % and 56.7 % intergenomic similarity values, respectively

(**Supplementary Table S4**), suggesting they may be assigned to the same genus.

Moreover, P8T1HF07V1 has 97% nucleotide identity to the GFV1 over specific

subregions that encode structural genes and the genes for DNA packaging and genome

integration **(Fig. 3)**. Similarly, in the same regions in DTV1 and MAV1 have 95-97%

identity. In contrast, the same regions in P8T1HF07V1 and POV1, and P8T1HF07V1 and

DTV1 have 53 and 75% nucleotide identity, respectively. Such similarity patterns suggest

that these viruses likely can infect hosts from both Thermotogota and Firmicutes phyla.

In contrast to the Group 1, none of the Group 2 proviruses had matches to

CRISPR spacers in 90 Thermotogota genomes, suggesting that the Group 2 viruses have

a more restricted host range within the Thermotogota or started to infect members of this

phylum recently.

Successful induction of TAV1 from *T*. *africanus* H17ap60333.

Induction assays were performed on three of the putatively lysogenized *Thermotogota*: *T.*

africanus H17ap6033 (Group 1), *Petrotoga* sp. P8T1HF07 (Group 2) and *P. olearia*

(Group 2). Only the provirus in *T. africanus* H17ap6033 (TAV1) was successfully

induced using mitomycin C. TAV1 was shown to produce viral particles with a

polyhedral head of ~50 nm in diameter and a flexible non-contractile tail of ~160 nm in

length and ~10 nm in width (**Fig. 5a**). Based on tail morphology, TAV1 was classified to

the order *Caudovirales* and the family *Siphoviridae*, confirming the sequence-based

classification. TAV1 morphology is similar to the three previously characterized

temperate *Marinitoga* viruses, whose virion tails were just slightly longer [15, 16]. In

addition to viral particles, a release of membrane vesicles or toga fragments was regularly

observed (**Fig. 5b**).

While the induction of the proviruses in *Petrotoga* sp. 8T1HF07 (P8T1HF07V1)

and *P. olearia* (POV1) using mitomycin C was unsuccessful, membrane vesicles of

various sizes and shapes (20 - 100nm) were produced by the cells, and in particular by the

induced *Petrotoga* sp. 8T1HF07 cells. Analysis of the supernatant of the latter culture

revealed similarly-sized round-shaped vesicles connected together in long chains by

hooking onto the flagella, like a "pearl necklace", while free vesicles showed more

diversity in size and shape (**Fig. 5c**). Some "sunflower-like" structures were also

observed inside a remaining cell (**Fig. 5d**). It is unknown if the provirus or stressors

influence the production of these vesicles and structures, or if they are produced

spontaneously.

A potential receptor for the Group 2 viruses

Several types of structures on the surface of bacteria, such as membrane proteins,

flagella, pili, or carbohydrate moieties, can act as virus receptors [47]. Most siphoviruses

of Gram-negative bacteria, and some of Gram-positive bacteria, use proteinaceous

receptors for adsorption [48, 49]. If the Group 2 viruses use the same protein receptor to

attach to both Thermotogota and Firmicutes cells, the large phylogenetic distance

between these hosts offers an opportunity to identify possible membrane protein receptors

bioinformatically, since the receptor proteins would be expected to be conserved across

the genomes from both phyla. It should be noted that this approach would only identify

possible protein receptors, while potential shared carbohydrate receptors would not be

detected.

Four predicted membrane proteins with transmembrane helices were identified in

all genomes carrying a Group 2 provirus. One of these was the viral holin gene, leaving

three receptor candidates: a ComEA family DNA-binding protein, an oxaloacetate

decarboxylase beta subunit, and an ABC transporter ATP-binding protein. Phylogenetic

analyses revealed that the ComEA and the oxaloacetate decarboxylase homologs are

widely distributed among Thermotogota (**Supplementary Fig. S6**). In contrast, the ABC

transporter is, among the Thermotogota, restricted to *Petrotoga* and *Defluviitoga*, the two

genera where the Group 2 proviruses are observed (**Supplementary Fig. S6, panel C**).

Moreover, the phylogenetic analysis suggests the homologs in *Petrotoga* and

Defluviitoga originated from an LGT event with a Firmicute (**Supplementary Fig. S6**).

These proteins show particularly high amino acid sequence similarity in the C-terminal

domain of both Thermotogota and Firmicutes homologs, which is facing the exterior of

the cell and could serve as a virus target (**Supplementary Fig. S7**). Although

experiments are needed to demonstrate if any of these proteins functions as receptor for

these viruses, we suggest that the ABC-transporter ATP-binding protein is a strong

candidate for a Group 2 virus receptor.

Moron genes are abundant in the identified proviruses.

Many temperate viruses are known to carry moron genes, which are genes that do not

have a direct viral function [50, 51]. The detected proviruses of Thermotogota are no

exception: the Group 1 proviruses carry up to 6 morons (**Fig. 1**), while the Group 2

proviruses have between 4 and 13 morons (**Fig. 2**). However, it should be noted that

because the 5' ends were hard to define for Group 2 proviruses, some of the moron genes

at the 5' ends might not be part of the proviruses. Sequencing virus DNA isolated from

capsids will help resolve this issue in the future.

Among the morons are several proteins that may confer a selective advantage to

the host (**Fig. 1** and **Fig. 2**, **Supplementary Table S2**). For instance, M1138V1 carry two

genes involved in sulfur metabolism. The Group 2 proviruses encode several transporters,

peptidases and hydrolases, likely to be beneficial for these heterotrophic bacteria. In

addition, all the viruses carry several hypothetical proteins that may also have non-viral

functions.

Evidence for the viruses' impact on lateral gene transfer

Eight hundred seventy homologs of 106 proviral genes were detected in 54 out of 59

Thermotogota genomes with no detectable proviruses (**Supplementary Table S6**). It

should be noted that some provirus genes, e.g. the Group 1 recombinases and terminases

(**Fig. 4** and Supplementary **Fig. S3**), did not pass our stringent screening criteria (see

Material and Methods), thus these represent minimum estimates of matches to proviral

genes in these genomes. Notably, 370 of 870 were homologs of 28 moron genes,

suggesting that the viruses may facilitate exchange of "host" genes among Thermotogota.

Moron genes also had the highest number of homologs across 54 genomes, with most

abundant being a queuine tRNA-ribosyltransferase in the Group 2 provirus DTV1 (found

in 48 genomes) and an aldo/keto reductase in Group 1 provirus GPV1 (found in 41

genomes).

Among phylogenetically informative datasets, 10 proviral genes group within

Thermotogota and 17 group within Firmicutes, suggesting that many of the proviral genes

originated either in Thermotogota and Firmicutes (**Supplementary Table S6**). For

instance, the above-described abundant moron gene queuine tRNA-ribosyltransferase is

of Thermotogota origin, while the aldo/keto reductase appears to be of Firmicutes origin

(**Supplementary Table S6**). In the phylogeny of another moron gene, a cadmium or

heavy metal transporter found in Firmicutes provirus MAV1 and Thermotogota Group 2

proviruses DTV1 and POV1, the provirus genes group closely with Firmicutes' homologs

(**Supplementary Fig. S8**). Notably, the other closely related Thermotogota homologs are

found in three *Fervidobacterium* and *Pseudothermotoga* genomes, genera where no

proviruses have yet been identified. Inspecting the genomic region surrounding these

genes in *Fervidobacterium* and *Pseudothermotoga*, revealed that the homolog of the

proviral recombinase (**Fig. 4**) is located immediately upstream of the transporter gene. No

other typical virus genes were observed in these regions, suggesting these genes are likely

remnants of proviruses. This also indicates that viruses related to Group 2 proviruses may

have broader host range that we presently detect.

Taken together the above analyses suggest that the viruses of both Group 1 and

Group 2 may facilitate exchange of genes not only among Thermotogota, but also

between Thermotogota and Firmicutes.

Discussion

In our search for proviruses in genomes of Thermotogota, we discovered two distinct

groups of temperate siphoviruses that have lysogenized this bacterial phylum. These

proviruses may represent multiple new viral species and genera. Our analyses suggest

that these viruses likely have broad host range that spans at least multiple genera. We also

found that the identified proviruses lineages are closely related to Firmicutes' viruses.

One of the bioinformatically identified Group 1 proviruses (TAV1) was induced

and shown to produce virus particles. The provirus resides in a genome of a *T. africanus*

isolate from the Hibernia oil reservoir off the Canadian east coast. The analysis of

CRISPR spacers suggested that this virus may have a particularly wide host range, with

the highest number of spacer-matches in genomes from outside its genus. For instance, a

virus very similar to TAV1 had likely infected *Thermotoga* spp. isolates from the North

Sea Troll oil reservoir. Similar predatory virus pressure in geographically and

geologically remote subsurface environments have been observed for *Methanohalophilus*

isolates from reservoirs in the USA and Russia [52].

We were not able to induce virus production from the selected Group 2 proviruses

This could be due to these proviruses currently being inactive, or we may not have

applied the right conditions to induce the expression of these proviruses. Nevertheless,

the high level of sequence identity between Group 2 provirus sequences from

Thermotogota and Firmicutes phyla uggests that they have been active very recently.

449   Many of the genes carried by both Group 1 and Group 2 proviruses are found in

genomes of Thermotogota that do not have detectable proviruses. These genes often

group with Firmicutes or viruses that infect Firmicutes, and many can be classified as

morons. This suggests that both Group 1 and Group 2 viruses transfer genes within

Thermotogota and between Thermotogota and Firmicutes, and may serve as a major

mechanism for the earlier reported large amounts of lateral gene transfer between

Thermotogota and Firmicutes [7, 8].

456   Based on our bioinformatic and phylogenetic analyses, we propose that an ABC

457   transporter may serve as a receptor for at least some of these proviruses. ABC transporter

458   proteins are, to our knowledge, not commonly identified as bacterioviral receptors.

459   However, *Lactoccocus* viruses from the siphoviral c2 group have been shown to use

460   membrane proteins Pip or YjaE, both with sequence similarity to ABC-transporter

461   domains, as secondary receptors [47, 53].

462   Intriguingly, transporters, and ABC transporters in particular, are among the most

frequently transferred genes both within the Thermotogota and between Thermotogota

and Firmicutes [7, 8, 54]. Transporter genes were also detected in the provirus genomes.

The possibility that transporters can function as viral receptors in the Thermotogota

therefore suggests that acquiring a new transporter, perhaps via a viral infection, might

result in the cell not only acquiring a new function but also becoming susceptible to a

new virus. This virus might carry another transporter gene, which can introduce yet

another virus, resulting in a ratchet-like process. Using transporters as receptors will

therefore not only provide the virus with the wide host range but could also make viruses

the vehicles on the highways of gene sharing observed between the Thermotogota and

Firmicutes.

Genes encoding proteins for membrane transport, including ABC transporters,

have been observed in several other viruses [55], and thus the proposed process could

operate widely among bacteria. This is contrary to a role commonly assigned to morons

where they often confer resistance to infections by other viruses [51]. Further studies and

experiments are needed to investigate if such ratchet processes are indeed occurring in

natural systems. However, regardless of the functions of the morons in the *Thermotogota*

proviruses, the observation of viruses potentially infecting organisms from different phyla

further demonstrates that viruses are key actors in the evolution of microbial diversity.

Acknowledgements

This work is supported by a Research Council of Norway award (project no.

180444/V40) to C.L.N., by the Sino-French LIA/PRC 1211 MicrobSea to J.L. and by the

Simons Foundation Investigator in Mathematical Modeling of Living Systems award

327936 to O.Z. Strains were obtained from the Université de Bretagne Occidentale

Culture Collection (UBOCC, Plouzané, France, www.univ-brest.fr/ubocc).

Competing Interests:

The authors declare no conflict of interest.

References

1.   Pollo SMJ, Zhaxybayeva O, Nesbø CL. Insights into thermoadaptation and the evolution of mesophily from the bacterial phylum *Thermotogae*. *Can J Microbiol* 2015; **61**: 655–670.

2.   Bhandari V, Gupta RS. The Phylum Thermotogae. In: Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F (eds). *The Prokaryotes*. 2014. Springer Berlin Heidelberg, Berlin, Heidelberg, pp 989–1015.

3.   Nesbø CL, S Swithers K, Dahle H, Haverkamp TH, Birkeland N-K, Sokolova T, et al. Evidence for extensive gene flow and Thermotoga subpopulations in subsurface and marine environments. *ISME J* 2015; **9**: 1532–1542.

4.   Foght JM, Gieg LM, Siddique T. The microbiology of oil sands tailings: past, present, future. *FEMS Microbiol Ecol* 2017; **93**.

5.   Nesbø CL, Charchuk R, Pollo SMJ, Budwill K, Kublanov IV, Haverkamp THA, et al. Genomic analysis of the mesophilic Thermotogae genus *Mesotoga* reveals phylogeographic structure and genomic determinants of its distinct metabolism: Comparative genomic analysis of *Mesotoga*. *Environ Microbiol* 2019; **21**: 456–470.

6.   Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, et al. A new view of the tree of life. *Nat Microbiol* 2016; **1**: 16048.

7.   Zhaxybayeva O, Swithers KS, Lapierre P, Fournier GP, Bickhart DM, DeBoy RT, et al. On the chimeric nature, thermophilic origin, and phylogenetic placement of the Thermotogales. *Proc Natl Acad Sci* 2009; **106**: 5865–5870.

8.   Nelson KE, Clayton RA, Gill SR, Gwinn ML, Dodson RJ, Haft DH, et al. Evidence

for lateral gene transfer between Archaea and Bacteria from genome sequence of

Thermotoga maritima. *Nature* 1999; **399**: 323–329.

9.   Magnabosco C, Biddle JF, Cockell CS, Jungbluth SP, Twing KI. Biogeography,

Ecology, and Evolution of Deep Life. In: Orcutt BN, Daniel I, Dasgupta R (eds).

Deep Carbon, 1st ed. 2019. Cambridge University Press, pp 524–555.

10. Anderson RE, Brazelton WJ, Baross JA. The Deep Viriosphere: Assessing the Viral

Impact on Microbial Community Dynamics in the Deep Subsurface. *Rev Mineral*

Geochem 2013; **75**: 649–675.

11. Cobián Güemes AG, Youle M, Cantú VA, Felts B, Nulton J, Rohwer F. Viruses as

Winners in the Game of Life. *Annu Rev Virol* 2016; **3**: 197–214.

12. Engelhardt T, Kallmeyer J, Cypionka H, Engelen B. High virus-to-cell ratios

indicate ongoing production of viruses in deep subsurface sediments. *ISME J* 2014;

8: 1503–1509.

13. Walsh EA, Kirkpatrick JB, Pockalny R, Sauvage J, Spivack AJ, Murray RW, et al.

Relationship of Bacterial Richness to Organic Degradation Rate and Sediment Age

in Subseafloor Sediment. *Appl Environ Microbiol* 2016; **82**: 4994–4999.

14. Ochman H, Lawrence JG, Groisman EA. Lateral gene transfer and the nature of

bacterial innovation. 2000; **405**: 6.

15. Lossouarn J, Nesbø CL, Mercier C, Zhaxybayeva O, Johnson MS, Charchuck R, et

al. 'Ménage à trois': a selfish genetic element uses a virus to propagate within

Thermotogales. *Environ Microbiol* 2015; **17**: 3278–3288.

16. Mercier C, Lossouarn J, Nesbø CL, Haverkamp THA, Baudoux AC, Jebbar M, et al.

Two viruses, MCV1 and MCV2, which infect Marinitoga bacteria isolated from

deep-sea hydrothermal vents: functional and genomic analysis. *Environ Microbiol*

2018; **20**: 577–587.

17. Markowitz VM, Chen I-MA, Chu K, Szeto E, Palaniappan K, Pillay M, et al.

IMG/M 4 version of the integrated metagenome comparative analysis system.

Nucleic Acids Res 2014; **42**: D568–D573.

18. Rice P, Longden I, Bleasby A. EMBOSS: The European Molecular Biology Open

Software Suite. *Trends Genet* 2000; **16,**: 276–277.

19. Lima-Mendez G, Van Helden J, Toussaint A, Leplae R. Prophinder: a

computational tool for prophage prediction in prokaryotic genomes. *Bioinformatics*

2008; **24**: 863–865.

20. Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. PHAST: A Fast Phage Search

Tool. *Nucleic Acids Res* 2011; **39**: W347–W352.

21. Akhter S, Aziz RK, Edwards RA. PhiSpy: a novel algorithm for finding prophages

in bacterial genomes that combines similarity- and composition-based strategies.

Nucleic Acids Res 2012; **40**: e126–e126.

22. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al.

BLAST+: architecture and applications. *BMC Bioinformatics* 2009; **10**: 421.

23. O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al.

Reference sequence (RefSeq) database at NCBI: current status, taxonomic

expansion, and functional annotation. *Nucleic Acids Res* 2016; **44**: D733–D745.

24. Darling AE, Mau B, Perna NT. progressiveMauve: Multiple Genome Alignment

with Gene Gain, Loss and Rearrangement. *PLoS ONE* 2010; **5**: e11147.

25. Zimmermann L, Stephens A, Nam S-Z, Rau D, Kübler J, Lozajic M, et al. A

Completely Reimplemented MPI Bioinformatics Toolkit with a New HHpred Server

at its Core. *J Mol Biol* 2018; **430**: 2237–2243.

26. Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan 5:

genome-scale protein function classification. 5.

27. Guy L, Roat Kultima J, Andersson SGE. genoPlotR: comparative gene and genome

visualization in R. *Bioinformatics* 2010; **26**: 2334–2335.

28. Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, et al. Circos:

An information aesthetic for comparative genomics. *Genome Res* 2009; **19**: 1639–

1645.

29. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using

DIAMOND. *Nat Methods* 2015; **12**: 59–60.

30. Nepusz T, Yu H, Paccanaro A. Detecting overlapping protein complexes in protein-

protein interaction networks. *Nat Methods* 2012; **9**: 471–472.

31. Bolduc B, Jang HB, Doulcier G, You Z-Q, Roux S, Sullivan MB. vConTACT: an

iVirus tool to classify double-stranded DNA viruses that infect *Archaea* and

Bacteria. *PeerJ* 2017; **5**: e3243.

32. Bin Jang H, Bolduc B, Zablocki O, Kuhn JH, Roux S, Adriaenssens EM, et al.

Taxonomic assignment of uncultivated prokaryotic virus genomes is enabled by

gene-sharing networks. *Nat Biotechnol* 2019; **37**: 632–639.

33. Lopes A, Tavares P, Petit M-A, Guérois R, Zinn-Justin S. Automated classification

of tailed bacteriophages according to their neck organization. *BMC Genomics* 2014;

15: 1027.

34. Moraru C, Varsani A, Kropinski AM. VIRIDIC – a novel tool to calculate the

intergenomic similarities of prokaryote-infecting viruses. 2020. Microbiology.

35. Nishimura Y, Yoshida T, Kuronishi M, Uehara H, Ogata H, Goto S. ViPTree: the

viral proteomic tree server. *Bioinformatics* 2017; **33**: 2379–2380.

36. Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Mol Biol Evol* 2013; **30**: 772–780.

37. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis

of large phylogenies. *Bioinformatics* 2014; **30**: 1312–1313.

38. Lee MD. GToTree: a user-friendly workflow for phylogenomics. *Bioinformatics*

2019; **35**: 4162–4164.

39. Price MN, Dehal PS, Arkin AP. FastTree 2 – Approximately Maximum-Likelihood

Trees for Large Alignments. *PLoS ONE* 2010; **5**: e9490.

40. Geslin C, Le Romancer M, Erauso G, Gaillard M, Perrot G, Prieur D. PAV1, the

First Virus-Like Particle Isolated from a Hyperthermophilic Euryarchaeote,

"Pyrococcus abyssi". *J Bacteriol* 2003; **185**: 3888–3894.

41. Haverkamp THA, Geslin C, Lossouarn J, Podosokorskaya OA, Kublanov I, Nesbø

CL. Thermosipho spp. immune system differences affect variation in genome size

and geographical distributions. *Genome Biol Evol* 2018.

42. L'Haridon S, Gouhier L, John ESt, Reysenbach A-L. Marinitoga lauensis sp. nov., a

novel deep-sea hydrothermal vent thermophilic anaerobic heterotroph with a

prophage. *Syst Appl Microbiol* 2019; **42**: 343–347.

43. Walker PJ, Siddell SG, Lefkowitz EJ, Mushegian AR, Adriaenssens EM, Dempsey

DM, et al. Changes to virus taxonomy and the Statutes ratified by the International

Committee on Taxonomy of Viruses (2020). *Arch Virol* 2020; s00705-020-04752–x.

44. Adriaenssens E, Brister JR. How to Name and Classify Your Phage: An Informal

Guide. *Viruses* 2017; **9**: 70.

45. Catalano CE. The terminase enzyme from bacteriophage lambda: a DNA-packaging

machine. *Cell Mol Life Sci CMLS* 2000; **57**: 128–148.

46. Rath D, Amlinger L, Rath A, Lundgren M. The CRISPR-Cas immune system:

Biology, mechanisms and applications. *Biochimie* 2015; **117**: 119–128.

47. Stone E, Campbell K, Grant I, McAuliffe O. Understanding and Exploiting Phage–

Host Interactions. *Viruses* 2019; **11**: 567.

48. Zhang Z, Yu F, Zou Y, Qiu Y, Wu A, Jiang T, et al. Phage protein receptors have

multiple interaction partners and high expressions. *Bioinformatics* 2020; **36**: 2975–

2979.

49. Bertozzi Silva J, Storms Z, Sauvageau D. Host receptors for bacteriophage

adsorption. *FEMS Microbiol Lett* 2016; **363**: fnw002.

50. Cumby N, Davidson AR, Maxwell KL. The moron comes of age. *Bacteriophage*

2012; **2**: e23146.

51. Taylor VL, Fitzpatrick AD, Islam Z, Maxwell KL. The Diverse Impacts of Phage

Morons on Bacterial Fitness and Virulence. *Adv Virus Res* 2019; **103**: 1–31.

52. Borton MA, Daly RA, O'Banion B, Hoyt DW, Marcus DN, Welch S, et al.

Comparative genomics and physiology of the genus *Methanohalophilus* , a

prevalent methanogen in hydraulically fractured shale. *Environ Microbiol* 2018; **20**:

4596–4611.

53. Millen AM, Romero DA. Genetic determinants of lactococcal C2viruses for host

infection and their role in phage evolution. *J Gen Virol* 2016; **97**: 1998–2007.

54. Nesbø CL, Nelson KE, Doolittle WF. Suppressive Subtractive Hybridization

Detects Extensive Genomic Diversity in Thermotoga maritima. *J Bacteriol* 2002;

184: 4475–4488.

55. Greiner T, Moroni A, Van Etten J, Thiel G. Genes for Membrane Transport

Proteins: Not So Rare in Viruses. *Viruses* 2018; **10**: 456.

56. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new

developments. *Nucleic Acids Res* 2019; **47**: W256–W259.

Table 1. CRISPR spacer matches to provirus genomes in Thermotogota genomes.

Matches to spacers from the provirus' host genome are labeled as a self-match.

provirus name	Genome with a CRISPR spacer match (number of spacers)
KPV1	*Kosmotoga olearia* (1)
MHV1	*Pseudothermotoga elfii* NBRC107921 (1), *Marinitoga* sp. 1154 (1)
MCV1*	*Marinitoga* sp. 1154 (2)
MCV2*	*Marinitoga* sp. 1155 (1) self-match, *Marinitoga* sp. 1154 (2)
MPV1*	*Marinitoga* sp. 1137 (1)
M1135	*Marinitoga* sp. 1155 (1)

M1137V1	*Pseudothermotoga elfii* NBRC107921 (1), *Marinitoga* sp. 1135 (2)
M1137V2	*Marinitoga piezophila* (1)
M1138	*Thermosipho africanus* TCF52B (1), *Thermosipho melanesiensis* (1), *Pseudothermotoga elfii* NBRC107921 (1), *Marinitoga* sp. 1137 (1)
TAV1*	*Thermosipho africanus* H17ap60334 (3) self -match, *Thermosipho africanus* TCF52B (2), *Thermosipho africanus* Ob7 (1), *Thermosipho melanesiensis* (2), *Thermotoga maritima* 2812B (1), *Thermotoga* sp. EMP (1), *Thermotoga* sp. XYL54 (3), *Thermotoga* sp. CELL2 (3), *Thermotoga* sp. TBGT1766 (3), *Thermotoga* sp. TBGT1765 (4), *Thermotoga* sp. A7A (1), *Thermotoga* sp. MC24 (2), *Pseudothermotoga elfii lettingae* (4), *Pseudothermotoga elfii* NBRC107921 (9), *Pseudothermotoga elfii* DSM9442 (2)
TMV1	*Thermosipho melanesiensis* (1) self-match, *Thermosipho africanus* TCF52B (1), *Thermosipho africanus* Ob7 (1), *Thermotoga* sp. TBGT1765 (1), *Thermotoga* sp. Mc24 (1), *Pseudothermotoga elfii* NBRC107921 (1), *Pseudothermotoga elfii lettingae* (2)
T1074V1	*Thermosipho affectus* Bl1070 (1), *Thermosipho affectus* 1223 (3), *Thermosipho affectus* Bl1063 (1)

641 *Proviruses that have been induced and shown to produce virus particles.
642

Fig. 1. Comparison of sequences from all detected Group 1 proviruses. Provirus

name and the species of its host are shown to the left of the nucleotide sequence, in which

predicted ORFs are depicted as arrows. Proviruses that have been induced and shown to

produce virus particles are marked with an asterisk. The lines connect regions of adjacent

viruses that have TBLASTX similarity of more than 30% over 100bp. Lines are colored

in red or blue indicate that the matching sequences encoded in the same or opposite

strand, respectively. The predicted ORFs are color-coded based on their function and

should be considered approximate, because it relies only on gene annotations. Selected

gene annotations are included and abbreviated as follows. Ser recomb: serine

recombinase, LexA: LexA repressor, ParB: ParB-like nuclease, RecT: RecT family

recombinase, dUTP hydrolase: deoxyuridine 5'-triphosphate nucleotidohydrolase, ssb:

single stranded DNA-binding protein, tss: terminase small subunit, tls: terminase large

subunit, mcp: major capsid protein, tail tape: tail tape measure protein, rRNA lsm:

ribosomal RNA large subunit methyltransferase, flagella bbp: flagella basal-body protein,

DnaC: DnaC replication protein, DnaD: DnaD replication protein, DNA pol sc: DNA

polymerase sliding clamp, SecB: SecB protein-export protein, rep organizer :replisome

organizer, RusA: RusA family crossover junction endodeoxyribonuclease, Cys peptidase:

cysteine peptidase, sulfate AT: sulfate adenylyltransferase subunit 2, PAPS reductase:

phosphoadenosine phosphosulfate reductase, CW hydrolase: cell wall-associated

hydrolase, MazF: MazF endoribonuclease, dsbr: DNA double-strand break repair protein,

metal bp: metal-binding protein, CMP hydrolase: cytidine 5′-monophosphate hydrolase.

The figure was produced using genoPlotR [27].

Fig. 2. Comparison of sequences from three complete Thermotogota Group 2

proviruses and their Firmicutes' homologs. Provirus name (in red for Thermotogota

and blue for Firmicutes) and the species of its host are shown to the left of the nucleotide

sequence, in which predicted ORFs are depicted as arrows. The lines connect regions of

adjacent viruses that have TBLASTX similarity of more than 30% over 100bp. Lines are

colored in red or blue indicate that the matching sequences encoded in the same or

opposite strand, respectively. The predicted ORFs are color-coded based on their function

and should be considered approximate, because it relies only on gene annotations.

Selected gene annotations are included and abbreviated; HMT ATPase: heavy metal

translocating ATPase, FMN reductase: flavine mono nucleotide reductase, HAD family

phosphatase: haloacid dehalogenase superfamily of hydrolase). The figure was produced

using genoPlotR [27].

Fig. 3. Comparison of representative Thermotogota proviruses. Due to sequence

similarity, only one provirus per Thermotogota genus is shown. The nucleotide sequences

of the proviruses are arranged around the circle and color-coded. Numbers indicate

kilobases (kb) and grey boxes outline locations of predicted genes. Lines connecting

different proviral sequences represent TBLASTX matches between the proviral regions,

with the percent identity shown in histograms at the ends of each line. The plot was

created using Circos [28].

Fig. 4. Maximum likelihood tree of recombinases found in Thermotogota proviruses

and of their homologs in Firmicutes proviruses, and Thermotogota and Firmicutes

genomes. Host names of Thermotogota and Firmicutes proviruses are colored in red and

blue, respectively. The names of their proviruses are added next to the host name. Names

of Thermotogota homologs that either resided outside of proviral regions or come from a

genome without detected proviruses are shown in black. Branches without labels

represent Firmicutes without an identified Group 2 provirus. Homologs from incomplete

proviruses are labeled with "(in)". Circles on the branches represent bootstrap support,

and only values above 70% are shown. Some proteins have identical amino acid

sequences in more than one organism. The protein labelled 'Bacteria inc. POLV1

Fervidobacterium spp.'corresponds to accession number WP_011994748.1 and is found

in *Fervidobacterium nodosum* Rt17-B1 (NC_009718.1), *Fervidobacterium pennivorans*

DSM 9078 (NC_017095.1), *Fervidobacterium islandicum* (NZ_CP014334.1),

*Fervidobacterium gondwane*nse DSM 13020 (FRDJ01), *Petrotoga olearia* (PNR98053)

and *Coprothermobacter proteolyticus* (PXJB01). The protein labelled 'Bacteria inc.

Mahella australiensis MAV1, *Pseudothermotoga elfii*' corresponds to accession number

WP_013782344.1 and is also found in *Clostridium* sp. SYSU GA15002T

(NZ_CP040924.1), *Thermoanaerobacter thermocopriae* JCM 7501 (NZ_KI912455.1),

Pseudothermotoga elfii and MAV1 from *Mahella australiensis*. The tree was rooted by

mid-point rooting and visualized using iTOL [56]. Tree scale, substitutions per site.

Fig. 5. Electron micrographs of the induced virus and vesicles, stained with 2%

uranyl acetate. Panel a. The TAV1 virus particle, which shows a typical Siphoviridae

morphology. **Panel b.** Vesicles and toga fragments produced by *Thermosipho africanus*

H17ap60334. **Panel c.** Vesicles produced by *Petrotoga* sp. 8T1HF07.NaAc.6.1, some of

which are attached to a flagellum. **Panel d.** Sunflower-like structures inside *Petrotoga* sp.

8T1HF07.NaAc.6.1 cells. The structures are highlighted by arrows.

Supplementary Figures:

Supplementary Fig. S1. Panel A. Gene-sharing network of proviruses calculated in

VContact2. The network is based on shared protein clusters between viral genomes.

Only proviruses at most three nodes away from MPV1 and P8T1HF07V1 are shown. The

Thermotogota proviruses are colored in red, viruses from Firmicutes are blue and viruses

infecting other taxa are colored orange. The quality scores calculated by ClusterOne are

0.94 (p=0.00004) for the Group 1 cluster and 0.83 (p=0.006) for the Group-2 cluster.

Panel B. Placement of proviruses on the phylogenetic tree of Thermotogota genomes

reconstructed from 74 single copy protein-coding genes. Closely related genomes

(distance > 0.1), monophyletic genomes from the same genus, and clades consisting of

only metagenome assembled genomes were collapsed. Identified proviruses are indicated

next to their respective host genera. The tree was visualized in iTOL [56]. Tree scale,

substitutions per site.

Supplementary Fig. S2. **Placement of complete Thermotogota and Firmicutes**

proviruses on the viral proteomic tree. The viral proteomic tree is from ViPTree v. 1.9

[35], and only the relevant region of the tree is shown. The Thermotogota and Firmicutes

proviruses are labeled with red stars. Taxonomy of the related viruses and their hosts is

indicated as color bars next to a terminal leaf on the tree.

Supplementary Fig. S3. Maximum likelihood trees of three families of terminase

large subunit genes. The phylogenetic trees displayed were constructed using RAxML

as implemented in Geneious v. 10 with a GAMMA-WAG substitution model and 100

bootstrap replicates. The trees should be considered unrooted. Bootstrap support > 70% is

shown on branches as circles, with the size corresponding to the strength of support.

Taxonomic labels of Thermotogota with proviruses are shown in red bold font, with the

provirus name listed after the host name. Thermotogota homologs from genomes with no

detected provirus are listed in bold font. Numbers in front of each taxon name represent

database accession numbers. The tree was visualized in iTOL and rooted by midpoint

rooting and should be considered unrooted [56].

Supplementary Fig. S4. Overview of CRISPR spacer sequences from Thermotoga

isolates from the Troll oil reservoir mapped on to the TAV1 genome. Alignment

position of each CRISPR spacer is indicated as black bars. Mapping and visualization

was performed in Geneious v. 10 and maximum of one mismatch was allowed.

Supplementary Fig. S5. Comparison of the three complete and two incomplete

Thermotogota Group 2 provirus sequences. Virus name and the genus the host belongs

to is indicated. The regions with significant pairwise BLASTX similarity scores are

connected, red indicates that sequence is in the same direction while blue indicates that

the similar sequences are on opposite strands. The predicted ORFs are color-coded based

on their function and should be considered approximate, because it relies only on gene

annotations. Selected gene annotations are included and abbreviated; HMT ATPase:

heavy metal translocating ATPase, FMN reductase: flavine mono nucleotide reductase,

HAD family phosphatase: haloacid dehalogenase superfamily of hydrolase),

dimethyladenosine trf: dimethyladenosine transferase, 2Fe-2S bp: 2Fe-2S binding p

rotein, MFS transporter: multi facilitator superfamily transporter. The figure was

produced using genoPlotR [27].

Supplementary Fig. S6. Maximum likelihood trees of three potential virus receptor

genes. Panel A: Competence protein ComEA, Panel B: oxaloacetate decarboxylase

and Panel C: ATP-binding cassette, subfamily B. Bootstrap support > 70% is shown

on branches as circles, with the size corresponding to the strength of support. The names

of Thermotogota taxa that contain Group 2 proviruses are displayed in red font and

Firmicutes with Group 2-like proviruses are displayed in blue font. Clades containing

sequences from the same genus are collapsed into wedges. The trees were rooted using

midpoint rooting, and should be considered unrooted. The trees were visualized in iTOL

[56].

Supplementary Fig. S7. Overview of the alignment of the ABC transporter ATP-

binding protein in Thermotogota and Firmicutes genomes with Group 2 proviruses.

Sites 100% conserved in all sequences sites are highlighted in color, while variable sites

are shown in grey. Transmembrane regions, predicted using the TMHMM Server v. 2.0,

are shown in red above the alignment (http://www.cbs.dtu.dk/services/TMHMM-2.0/).

Supplementary Fig. S8. Maximum likelihood tree of the moron gene annotated as a

cadmium transporter. Bootstrap support > 70% is shown on branches as circles, with

the size corresponding to the strength of support. Taxon names of Thermotogota with a

provirus are given in red and taxon name of Firmicutes with a Group 2-like provirus are

given in blue. Provirus name is also indicated. Thermotogota homologs from genomes

with no detected provirus, or where the homolog is found outside the provirus region, are

given in bold font. Database accession numbers are shown in front of taxonomic names.

The tree was rooted by midpoint rooting and visualized in iTOL [56].

Mesotoga sp. SC_TOLDC MTOLDCV1 (incomplete)

Mesoaciditoga lauensis MLV1

Kosmotoga pacifica KPV1

Thermosipho sp. 1074 T1074V1

Thermosipho africanus TAV1*

Thermosipho melanesiensis TMV1

Geotoga petraea GPV1

Marinitoga sp. 1138 M1138V1

Marinitoga sp. 1135 M1135V1

Marinitoga piezophila MPV1*

Marinitoga camini MCV2*

MCV1*

Marinitoga hydrogenitolerans MHV1

Marinitoga sp. 1137 M1137V2

M1137V1 (incomplete)

5 kb

integration ▶, transcriptional regulation ▶, DNA metabolism ▶, DNA packaging and head ▶, head to tail ▶, tail ▶, HNH endonuclease ▶, lysis ▶, morons ▶, hypothetical ▶