**Viruses of the Thermotogota; the vehicles on the highways of gene**

**sharing.**

Running title: Proviruses from Thermotogota.

Thomas Haverkamp[a#], Julien Lossouarn [b], Olga Zhaxybayeva[c], Jie Lyu[d], Nadège

Bienvenu[d], Claire Geslin[d], Camilla L. Nesbø[e,f]

[a] Centre for Ecological and Evolutionary Synthesis, Department of Biosciences,

University of Oslo, Norway.

[b] Université Paris-Saclay, INRAE, AgroParisTech, Micalis Institute, 78350,

Jouy-en-Josas, France

[c] Department of Biological Sciences, Dartmouth College, Hanover, NH, USA.

[d] Univ Brest, CNRS, IFREMER, Laboratoire de Microbiologie des Environnements

Extrêmes, F-29280 Plouzané, France

[e] Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada

[f] Biozone, Department of Chemical Engineering and Applied Chemistry and BioZone,

University of Toronto, 200 College Street, Toronto, Ontario, Canada, M5S 3E5

[#] present address: Department of Epidemiology, Norwegian Veterinary Institute, Oslo,

Norway.

**Abstract**

Viruses are drivers of microbial ecology and evolution controlling populations and

disseminating DNA laterally among the cells they infect. Phylogenetic and comparative

genomic analyses of Thermotogota bacteria have shown high levels of lateral gene

transfer with distantly related organisms, particularly with Firmicutes. One likely source

of such DNA transfers is viruses, however, to date only three temperate viruses infecting

*Marinitoga* bacteria have been characterized in this phylum. Here we use a bioinformatic

approach to identify an additional 17 proviruses integrated into genomes of eight

Thermotogota genera including *Marinitoga*. We also induce viral particle production

from one of the newly identified proviruses. The proviruses fall into two groups based on

sequence similarities, gene synteny and virus classification tools. Group-1 shows similar

genome structure to the three previously identified *Marinitoga* viruses while Group-2 are

distantly related to these viruses and have different genome organization. Both groups

show close connections to Firmicutes in genomic- and phylogenetic analyses, and the

Group-2 viruses show evidence of very recent transfer between these lineages and are

likely capable of infecting cells from both phyla. We suggest that viruses are responsible

for a large portion of the laterally transferred DNA between these distantly related

lineages.

**Introduction**

The phylum Thermotogota comprises anaerobic fermentative bacteria, most of which are

thermophiles [1]. They are found as a deep branching lineage in some universal

phylogenetic trees, for example those made from 16S rRNA genes [2], and are common

in subsurface environments such as marine vents, terrestrial hot springs and deep

subsurface oil reservoirs [3][4–6].

Lateral gene transfer (LGT) has been an important force shaping the genomes of

Thermotogota, with Firmicutes and Archaea being their most notable gene transfer

partners [1, 7, 8]. In fact Firmicutes and Thermotogota have been suggested to be linked

by ''highways of gene sharing'' [8]. How these gene-sharing connections between phyla

occur, is however, still unclear. Homologous recombination among closely related

genomes is another important evolutionary force acting on Thermotogota genomes [4, 9].

For both the *Thermotoga* and *Mesotoga* genera, comparative genomic analyses and

analyses of homologous recombination have shown that long fragments of DNA (>

10,000 nucleotides) are transferred between closely related strains [6]. Both types of

gene-sharing processes might be driven by viruses.

The subsurface constitutes the largest biosphere on earth and is estimated to

contain ~70% of all cells [10]. Viruses are likely to be particularly important in

subsurface environments with 97% of all viruses on earth being found in soil and

sediments [11, 12]. Moreover, although both prokaryotic cell and virus numbers

decreases with depth, the virus to cell ratio increases with depth [11, 13, 14].

Phylogeographic studies of hyperthermophilic *Thermotoga* and mesophilic *Mesotoga*

bacteria have revealed genetic interaction between geographically distant populations,

particularly among the hyperthermophilic Thermotoga [4, 6]. Viruses are one potential

source of such long-distance dispersal of genetic material, particularly for anaerobic

organisms where surface dispersal is problematic.

However, although viruses are likely candidates for transferring DNA, evidence

of viral activity is so far limited to one Thermotogota genus. Three temperate

*Siphoviridae* viruses have been described that infect *Marinitoga* species [15, 16]. MPV1

infects the deep-sea marine vent bacterium *Marinitoga piezophila* and displays an

intricate relationship with a plasmid co-occurring in the same host, illustrating the

importance of mobile elements in these ecosystems, while MCV1 and MCV2 infect

*Marinitoga camini*. The three viruses show similar genomic organization and virion

morphology. Phylogenetic analyses of the virus ORFs revealed close connections to

Firmicutes viruses, which prompted the conclusion that viruses belonging to these phyla

sample genes from a common pool.

Here we identify 17 provirus sequences (including 13 likely complete provirus

elements) in Thermotogota genomes using a bioinformatic approach that included several

phage-detection tools as well as manual inspection of the results. We successfully

induced one of these novel proviruses. These new proviral sequences are found in

genomes from eight Thermotogota genera, including *Marinitoga*. We report that the

Thermotogota proviruses fall into two distinct groups, and that the proviruses from one of

these groups are very similar to Firmicutes viruses. We also hypothesize that membrane

transport proteins (e.g. ABC-transporters) may serve as receptors for Thermotogota

viruses. We propose a ratchet-like process, where moron genes encoding transmembrane

proteins may make the host vulnerable to new viruses, which could account for the

highways of gene-sharing observed between Thermotogota and Firmicutes.

**Material and Methods**

**Data collection and genome analyses of viruses.**

111 Thermotogota genomes available in Genbank or IMG [17] were accessed until June

2018. For draft genomes we combined the contigs using the "union" command from the

EMBOSS package (version 6.6.0) [18]. This created 'artificially' closed-genomes needed

for Prohinder (http://aclame.ulb.ac.be/Tools/Prophinder/) [19]. For consistency, we also

used these 'artificially' closed genomes with the other provirus detection tools. We

screened each genome for the presence of prophages using three tools. The genomes were

uploaded to webservers for Prophinder and PHAST (http://phast.wishartlab.com/) [20]

and processed using default settings at both websites. In addition, we screened all the

genomes using PhiSpy (version 2.3) [21]. From the output we recorded: start and stop

position of the provirus sequence, the size of the provirus, the number of open reading

frames (ORFs), the presence of tRNA sequences. The annotation of ORFs in the

predicted regions were manually screened for the presence of prophage /mobile element

related genes by searching for the terms: "Viral", "Phage". Putative provirus regions with

few provirus-like annotations, were discarded. This might lead to an underestimate of

proviruses. In the case of the genomes composed of concatenated contigs, we discarded

provirus identifications that crossed contig borders. The provirus calls for each of the

three tools were compared and when proviruses were predicted by multiple tools in the

same region of the chromosomes, the annotations were inspected to identify the most

likely regions. When only one tool identified a putative virus sequence, we manually

checked if the spanned region had to be corrected. This included the following steps: 1)

PHAST can predict attR and attL integration sequences for the identified provirus, and

when predicted, the region between the sites was manually inspected to make sure the

region contained virus genes. 2) We screened for the presence of a tRNA gene close to

the attR/L site that could function as an integration site. 3) We determined if genes from

the host genome were incorrectly included in the predicted prophage region by

comparisons to closely related genomes (from the same genus) that did not have a

provirus. The predicted provirus sequences were compared to publicly available genomes

by searching the NCBI non-redundant (nr) database [22] with provirus ORFs as queries

using BlastX ver.2.2.26 [23]. When homologs for multiple ORFs from the predicted

provirus were found in the same distantly related reference genome (usually a Firmicutes

genome), the identified reference genome was downloaded for a more detailed

comparison. The reference genome was aligned to the host genome of the putative

provirus using Progressive Mauve [24]. In most cases this visualized only the aligned

proviruses in two otherwise distantly related microbial genomes and this information was

included when determining the genomic coordinates of the provirus.

Proviruses were considered complete if they contain modules for lysogeny,

replication, packaging, head/tail morphogenesis and lysis. If one or more of these

modules were missing, the provirus was scored as partial. It should be noted that for two

of the proviruses considered complete (M1135V1, M1138V1) the lysogeny module was

not clearly identified and for two others (MLV1, T1074V1) the lysis module was not

easily detected.

The predicted provirus sequences were compared to each other using blastn and

tblastx [23] and visualized in GenoplotR [25] and Circos [26]. Provirus-ORF annotations

were obtained from their respective Genbank entries, from blastp searches [23] of the

NCBI non redundant (nr) database with an expected-value cutoff of $10^{-1}$, and to the PDB

database using HHpred [27] with a probability cutoff of 99%. In addition, recombinase

and terminase ORFs were annotated using InterProScan as implemented in Geneious

v.10.

The provirus genomes were compared to NCBI's viral ref seq v. 94 database

using VContact2 [28][29], and their morphological and  taxonomic classification was

assessed using Virfam [30], VIRIDIC [31] and VIPTree [32].

Possible receptor proteins for the Group 2 viruses were identified in IMG using a

similarity cut-off at 50%. Homologs in *G. ferrireducens*, not available in IMG, were

identified using blastp.

**Phylogenetic analyses of virus genes.** Each translated virus gene from the gene-family

analyzed were queried using blastp against *nr* (accesses December 2019 – March 2020),

as well as a local database of *Thermotogota* virus proteins, and the 20 top scoring

matches were retrieved and aligned using MAFFT (expected value cut-off at $10^{-5}$).

Identical sequences were removed, as well as highly similar sequences from the same

genus, and alignment positions with > 50% gaps were trimmed. Phylogenetic trees were

constructed using RAxML [33] as implemented in Geneious v.10 with GAMMA+WAG

model of substitution (with 4 rate categories) and 100 bootstrap replicates.

**Virus induction and electron microscopy.**

*T. africanus* H17ap6033 and *Petrotoga* sp. 8T1HF07.NaAc.6.1 were cultivated in a

modified Ravot medium as previously described in reference [15] at 65°C or 55°C,

respectively. Attempts were made to increase the viral production of the strains by using

mitomycin C, as reported previously [15, 16]. A final concentration of 5 µg/mL of

mitomycin C was added to 300 mL bacterial culture at early to mid-log growth phase.

After 3 hours of incubation with mitomycin C, cultures were centrifuged at 7500 rpm and

4°C for 15 min, and supernatants were ultracentrifuged at 37 000 rpm (~100 000 g) and

10°C for 1h (Beckman Optima LE-80 K; rotor 70.1.Ti). Pellets were resuspended in

100µL of buffer (10 mM Tris-HCL, 100 mM NaCl, 5 mM $CaCl_2$, 20 mM $MgCl_2$) and

suspensions were prepared for negative staining electron microscopy as previously

described [34]. Briefly, 5 µL of the suspensions were directly spotted onto a Formwar

carbon coated copper grid. Putative virus-like particles were allowed to adsorb to the

carbon layer for 2 min and excess of liquid was removed. 5 µL of a staining uranyl

acetate solution (2%) was then spotted to the grid for 45 s and excess of liquid was

removed again. The grid was imaged at 120 kV in a JEOL JEM 100 CXIIVR transmission

electron microscope.

**Results and Discussion**

**Genomic characterization of Thermotogota proviruses identify two distinct virus**

**lineages.**

Analysis of 111 Thermotogota genomes identified 20 proviruses, including the three

already characterized viruses from *Marinitoga* and four partial proviruses

(Supplementary Table S1). All available *Thermosipho melanesiensis* genomes have

identical provirus sequences [35] and only one representative of these genomes were

included in our analyses below. Annotation of coding sequences (CDS) in the proviruses

are reported in Supplementary Table S2.

Comparisons of the predicted provirus genomes using several different tools

outlined below revealed two distinct groups of Thermotogota proviruses, here denoted as

Group-1 and Group-2 (Fig. 1 - Fig. 3). For these comparative genomic analyses, we also

included a provirus reported in *Marinitoga lauensis* [27] after our bioinformatic screening

was performed. The genome organization differs between the two groups and the

majority of top scoring blast-matches when comparing the Thermotogota proviruses

occur within the two groups (Fig. 1, Supplementary Table S2). The distinction of these

groups was supported by network analysis comparing the predicted provirus proteins to

the NCBI virus refseq database (Supplementary Fig. S1), where the Thermotogota

proviruses form two distinct clusters connected to a larger cluster mainly composed of

Firmicutes viruses. This clustering was also observed by the VIPTree tool

(Supplementary Fig. S2). Group-1 (N=15) is most widespread, found in the genera

*Marinitoga*, *Thermosipho*, *Kosmotoga*, *Mesotoga*, *Geotoga* and *Mesoaciditoga*, while

Group-2 (N=5) is limited to the genera *Petrotoga* and *Defluviitoga* (Supplementary Fig.

S1). Some of the proviruses appear to be incomplete, (M1137V1, MTOLDCV1, PMV1,

P9T1HF07V1), and may be in the process of being lost from the genomes or

domestication.

*Genomic characterization of Group-1 proviruses:* None of the 15 Group-1

proviruses identified here (including the three characterized temperate viruses and the

incomplete proviruses), exhibits significant nucleotide similarity with viral genomes in

the NCBI virus database. Following taxonomic criteria where viruses belonging to the

same genus display > 50-70% nucleotide identity over the full genome while species

display > 95% nucleotide identity [36][37], the VIRIDIC tool suggested that the complete

Group-1 proviruses identified here, correspond to 13 new species and at least 11 new

genera (Supplementary Table S2). Two provirus pairs, MCV1- MCV2 and M1135V1 -

M1138V1, may be assigned to the same genus having intergenomic similarity scores of

56% and 60% respectively (Supplementary Table S2).

The newly identified Group-1 proviruses show the same modular structure as

MPV1, MCV1 and MCV2 [15, 16] (Fig. 2). The 5' module in the provirus contains genes

involved in lysogeny and is encoded on the opposite strand compared to the rest of the

virus genes. This module is followed by modules for replication, packaging,

morphogenesis and host lysis. As noted for the described *Marintoga* viruses [15, 16], the

most variable module is the lysogeny module, where only the recombinase is conserved

between different proviruses (Fig. 2). Higher level of conservation is observed for the

other modules, and some of the conserved genes are discussed in more detail below. The

novel proviruses identified here likely correspond to siphovirus based on their head-neck-

tail gene sequences [30], which is consistent with the morphology observed for the

characterized viruses in this group (MPV1, MCV1, MCV2), and TAV1 induced in the

current study (see below).

All the complete Group-1 proviruses are inserted next to a tRNA gene. Eight of

them (the *Marinitoga* proviruses, MPV1, MCV1, MCV2, MHV1 and M1137V2, KPV1

(*Kosmotoga pacifica*), GPV1 (*Geotoga petrae*), MLV1 (*Mesoaciditoga lauensis*)), are

inserted next to tRNA-Glu, and share a very similar site-specific DNA serine

recombinase (KEGG Orthology; K06400, homologs of Marpi_0291 in MPV1 from *M.*

*piezophila*) (Fig. 4). The *Thermosipho* proviruses also carry homologs of this

recombinase but are inserted next to different tRNA genes. TMV1 and T1074V1, which

have recombinases that cluster together in the tree in Fig. 4, are inserted next to tRNA-

Phe. The most divergent homolog of this recombinase is found in TAV1 which is inserted

next to tRNA-Pro. We suggest that these recombinases are integrases that specifically

recognizes the tRNA genes the provirus is inserted next to (i.e. tRNA-Glu, tRNA-Phe or

tRNA-Pro). M1135V1 and M1138V1 are inserted next to tRNA-Cys and have two

unrelated ORFs with no suggested function in this position.

Terminase proteins are biological machines involved in the packaging of virus

DNA into the virus particle [38]. The Group-1 virus genomes have three families of

terminases (BLASTP cut-off exp. $> e^{-1}$, ident. $> 25\%$, Supplementary Fig. S3). The first

family, exemplified by Marpi_0320 in MPV1, has a PBSX family domain and has

homologs in MCV1 and MCV2 and the proviruses in *Marinitoga* sp. 1135, *Marinitoga*

sp. 1138, *Thermosipho* sp. 1074 and *M. lauensis*. The second type, exemplified by the

terminase protein in TAV1 (H17ap60334_04902) has a 'Terminase_lsu_T4-like' domain

and close homologs in the proviruses from *K. pacifica*, *T. melanesiensis*, *Mesotoga* sp.

TolDC and *G. petrae*. Finally, the proviruses in *M. hydrogenitolerans* and *Marinitoga* sp.

1137 have a third group of similar terminases (BUA62_RS02495, LN42_01905 and

LN42_00550, a pseudogene) in the same position, also with a 'Terminase_lsu_T4-like'

domain, thus these are distant homologs of the second terminase family (Supplementary

Fig. S3).

In addition to the recombinase and terminase, other typical virus proteins such as

capsid and portal proteins were also identified (Fig. 2). Moreover, two transcription

regulators (Marpi_0297 and Marpi_0298 in MPV1), a DNA repair exonuclease

(Marpi_0340 in MPV1), a single stranded DNA-binding (Marpi_0306 in MPV1) and

the tail tape measure protein (Marpi_0334 in MPV1) show relatively high identity across

most Group 1 viruses (Fig. 2, Supplementary Table S2). Interestingly, some hypothetical

proteins (Marpi_0299, Marpi_0338 in MPV1) are also shared among several of the

Group-1 provirus genomes, suggesting these genes provide important functions for these

viruses.

*Genomic characterization of Group-2 proviruses:* The sequences and genome

organization of the three likely complete Group-2 proviruses differs considerably from

that of Group-1 (Fig. 1, Fig. 3). In addition to the Group-2 provirus genomes found in

*Petrotoga* sp. P8T1HF07, *Petrotoga olearia* and *Defluviitoga tunesiensis,* we identified

partial proviruses in the genomes of *Petrotoga mobilis* SJ95 and *Petrotoga* sp. 9T1HF07

(Supplementary Fig. S4).

The classification based on the head-neck-tail module [30] assigned them to

*Siphoviridae* of Type1 - Cluster 2. All hosts of previously described members of this

virus lineage belong to Firmicutes. In agreement with this, similarity searches in Genbank

revealed that these proviruses show remarkably high similarity to proviruses found in

three Firmicutes genomes (Fig. 3); *Lacticigenium_naphta*e (LNV1), *Geosporobacter*

*ferrireducens* (GFV1) and *Mahella australiensis* (MAV1).

The VIRIDIC analysis of the three full-length Group-2 proviruses (P8T1HF07V1,

DTV1 and POV1), suggested they represent three new viral species (Supplementary

Table S2). Interestingly, two proviruses-pairs from different phyla, P8T1HF07V1-GFV1,

and DTV1-MAV1, have intergenomic similarity scores of 66.7 % and 56.7 %,

respectively, suggesting they may be assigned to the same genus (Supplementary Table

S2).

The 5' module of these proviruses, encoding genes involved in virus replication

and transcription, shows high level of diversity across the three Thermotogota viruses

analyzed and contains several morons (i.e. proviral genes that do not have a direct viral

function [39, 40]). This module is followed by highly conserved packaging,

morphogenesis and lysis modules. The lysogeny module is located in the 3' end of the

provirus, and also contains variable morons. The observation of many morons in Group-2

suggests that these proviruses might affect host physiology and metabolism, as well as

facilitating lateral transfer of these genes between hosts. However, it should be noted that

some of these morons, particularly at the 5'end, which was hard to define, might not be

part of the provirus. Sequencing virus DNA isolated from capsids will resolve this issue.

The site-specific serine recombinases carried by the Group-2 proviruses are

homologous, although distantly related (> 25% identity), to the ones found in the

majority of the Group-1 proviruses (Fig. 4). We identified a total of 28 additional protein

families shared between the two groups (Table 2, Supplementary Table S2), among these

are the two methylases and the regulatory proteins discussed above that are widely

distributed among the Group-1 viruses. This suggests that these distinct virus-lineages are

sampling genes from a common pool.

**Distribution and broad host range of Thermotogota Group-1 (pro)viruses**

The Group-1 proviruses are widespread across the Thermotogota phylum (Supplementary

Fig. 1B). In addition to the identified proviruses, several Thermotogota genomes with no

provirus carry close homologs of Group-1 provirus genes, e.g. the serine recombinase

discussed above, suggesting a history of infection of a Group-1 virus (Fig. 4). Blast

searches of hallmark virus genes [41] against a database of 59 Thermotogota genomes,

with no provirus identified, identified homologs (> 50% identity and 60% coverage, see

below) of typical virus genes in 50 genomes. Among these were a terminase homolog in

the genome of *Thermotoga* sp. RQ7 and the recombinase in the genome of *Thermotoga*

sp. SG1 (Fig. 4, Supplementary Fig. S2), suggesting Group-1 viruses also have infected

bacteria from the genus *Thermotoga* where no provirus have been identified yet. Several

*Thermosipho*, *Marinitoga* and *Mesotoga* genomes with no provirus also carry homologs

of this recombinase (Fig. 4).

The relatively high similarity among the Group-1 proviral genomes (Fig. 2)

suggest that they display broad host ranges, as observed for MPV1 from *M. piezophila*,

which was shown to be able infect and transfer a plasmid to a *Thermosipho* isolate [15].

To explore the host range of these predicted or active temperate viruses, we mapped

CRISPR spacer sequences from 90 Thermotogota genomes to the Group-1 genomes

(Table 1). This revealed that 12 of the provirus genomes have one or more matches to

CRISPR spacers in Thermotogota genomes. Three proviruses had no matches; MLV1

from *Mesoaciditoga*, the incomplete MTOLDCV1 from *Mesotoga*, GPV1 from *Geotoga*.

Two viruses had matches to spacers from its host genome: MCV2, TAV1. Most of the

viruses had matches to CRISPRs from genomes from the same genus as the host genome.

Interestingly, five viruses had matches to spacers from a genus different than the host,

supporting the notion that the Group-1 viruses have broad host ranges (Table 1). TAV1

had the most of such matches with 43 spacers from *Pseudothermotoga elfii, Thermosipho*

spp. and *Thermotoga* spp. mapping to the genome, suggesting that this virus has a

particularly broad host range.

Anecdotal evidence also supports a broad host range of TAV1. When the sample

from the Hibernia oil reservoir containing TAV1 and its host, *T. africanus* H17ap60334,

was processed in the lab, we observed a mass death of *Thermotoga* isolates, originating

from Troll oil reservoir in the North Sea, that were in the process of being transferred to

fresh media. When sequencing the genomes of the surviving *Thermotoga* isolates [4], all

survivors had at least one CRISPR spacer matching the TAV1 genome (Supplementary

Fig. S5). Interestingly, these spacers are located in the middle of the CRISPR array,

indicating that the spacers are not new acquisitions [42], but that the surviving isolates

have encountered a very similar virus, probably while inhabiting the Troll oil reservoir.

These observations suggest that TAV1 can infect both *Thermotoga* and *Thermosipho*

strains. It also suggests that the Troll and the Hibernia oil reservoir harbor closely related

virus populations. Similar predatory stress in geographically and geologically remote

subsurface environments have been observed for *Methanohalophilus* genomes originating

from reservoirs from USA and Russia [43].

**Close relationship of Thermotogota Group-2 proviruses and *Firmicutes***

For the Group-2 proviruses, highly similar (65-100% amino acid identity for individual

proteins) provirus were detected in three Firmicutes genomes; *Geosporobacter*

*ferrireducens* IRF9 (NZ_CP017269), *Lacticigenium naphtae* DSM 19658 and *Mahella*

*australiensis* 50-1 BON (NC_015520). When comparing the Thermotogota and

Firmicutes proviruses it was apparent that the levels of similarity does not correspond to

phylogeny of the host (Fig. 3). Genome alignments revealed that the provirus in

*Petrotoga* sp. P8T1HF07 shows highest similarity to the provirus in *G. ferrireducens*

with 97% DNA identity over 11,229bp and 8,389bp aligned by MAUVE [24] covering

structural proteins and the packing machinery. In comparison, for the same regions,

P8T1HF07V1 show 53-75% DNA identity to the proviruses from *P. olearia* and *D.*

*tunesiensis*.  Similarly, the provirus in *D. tunesiensis* is most similar to the one from *M.*

*australiensis* (95% and 97% identity in the same regions). These patterns suggest that

these viruses are frequently transferred between cells belonging to different phyla.

In contrast to the Group-1 proviruses, no Thermotogota CRISPR spacers mapped

to the Group-2 proviruses, suggesting that the Group-2 proviruses either have a more

restricted host range within the Thermotogota or a more recent introduction to this

phylum. With the exception of *D. tunesiensis*, the genomes carrying Group-2 proviruses

originated from hydrocarbon impacted environments (Supplementary Table S1). The

three Firmicutes isolates also originated from hydrocarbon impacted sites  (*M.*

*australiensis*; Riverslea oil field, Bowen-Surat basin, Australia [44], *L. naphtae*; crude oil

from oil-water well in Akita, Japan [45] or oil contaminated soil (*G. ferrireducens*;

Daejeon, South Korea [46]). These putative temperate viruses therefore appear to be

particularly abundant in oil reservoirs and hydrocarbon impacted sites.

**Importance of Thermotogota viruses in lateral gene transfer.**

The Group-2 proviruses highlights the close connection between Thermotogota and

Firmicutes viruses. However, the phylogenetic analysis of Group-1 provirus ORFs also

demonstrates a close connection to Firmicutes or viruses that infect Firmicutes (Fig. 4,

Supplementary Fig. S2, Supplementary Fig. S3). Blast searches of all Group-1 provirus

genes against nr supported this, and Firmicutes is the top scoring match of 15% of the

virus ORFs outside their genus (8-30% for individual Group-1 proviruses, Supplementary

Table S2). This is in agreement with previous analyses reporting large amounts of gene

transfer between the Thermotogota and Firmicutes e.g. reference [7, 8].

In order to further assess the importance of these viruses in LGT, we screened

genomes with no provirus for the presence of virus genes (Fig. 6, Supplementary Table

S2). The Group-2 viruses had the most Blast matches, with 76 Group-2 proteins having a

total of 613 matches, while 31 Group-1 proteins had a total of 262 matches. This is likely

due to the Group-2 proviruses having more morons. Interestingly, Group-2 genes were

abundant in *Pseudothermotoga* and *Fervidobacterium*, where no proviruses have been

detected yet.

Phylogenetic analyses revealed that some of the genes are of Thermotogota

origin, while for other genes, such as a cadmium transporter found in proviruses DTV1

and POV1, the virus homolog was distantly related to the homologs in most

Thermotogota genomes and was likely acquired from Firmicutes (Supplementary Fig.

S6). The phylogeny of the cadmium transporter also revealed that the virus gene has been

transferred into *Fervidobacterium* and *Pseudothermotoga* genomes, supporting the notion

that Group-2 viruses have interacted with these genera. For the Group-2 proviruses, the

highest number of matches was to a moron in DTV1 annotated as Queuine tRNA-

ribosyltransferase of Thermotogota origin (Supplementary Table 2S). The Group-1 gene

with the highest number of matches was a moron in GPV1 from *G. petraea* annotated as

an aldo/keto reductase, and also this gene appears to be of Firmicutes origin. These

analyses suggest that both Group-1 and Group-2 viruses are transferring genes both

within the Thermotogota, but also between Thermotogota and Firmicutes.

**Induction of TAV1 from *T. africanus* H17ap60333**

Induction assays were performed on three of the putatively lysogenised *Thermotogota*

isolated from petroleum reservoirs: one from Group-1 (*T. africanus* H17ap6033) and two

from Group-2 (*Petrotoga* sp. P8T1HF07 and *P. olearia*). We successfully induced the

provirus in *T. africanus* H17ap6033 using mitomycin C [15, 16]. TAV1 was shown to

produce viral particles with a polyhedral head of ~50 nm in diameter and a flexible non-

contractile tail of ~160 nm in length and ~10 nm in width (Fig. 5a). Based on tail

morphology, TAV1 was classified to the order *Caudovirales* and the family *Siphoviridae*,

confirming the classification using bioinformatic tools. This is similar to the three

previously characterized temperate *Marinitoga* viruses, whose virion tails were just

slightly longer [15, 16]. TAV1 induction produced low amounts of virions upon

mitomycin C induction (data not shown). In addition to viral particles, we regularly

observed membrane vesicles or toga fragments released from *T. africanus* H17ap6033

(Fig. 5b).

We also attempted, but were not successful in, inducing the provirus in *Petrotoga*

sp. P8T1HF07 and *Petrotoga olearia*, using mitomycin C. However, as observed for *T.*

*africanus* H17ap6033, large numbers of membrane vesicles of various sizes and shapes

(20 - 100nm) were produced by the cells (Fig. 5c). For *Petrotoga* sp. P8T1HF07, we

observed vesicles of similar round shape, connected together and extended as long chains

by hooking on to the flagella, like a "pearl necklace", while free vesicles showed more

diversity in size and shape (Fig. 5c). Some "sunflower-like" structures were also

observed inside the cell (Fig. 5d). It is unknown if the provirus or environmental stressors

influence the production of these vesicles and structures, or if they are produced

spontaneously.

Thus to date, only Group-1 proviruses from *Marinitoga* [15, 16] and *Thermosipho*

have been induced leading to the production of viral particles. The fact that we have not

been able to induce the Group-2 proviruses tested, could suggest they are inactive in

Thermotogota genomes or that they are in the process of domestication or deletion from

the genomes. Alternatively, we may not have provided the optimal conditions for

induction.

**Identification of a potential receptor of the Group-2 viruses**

Several types of structures on the surface of bacteria can act as virus receptors, e.g.

membrane proteins, flagella, pili, carbohydrate moieties on the cell surface [47]. Most

siphoviruses of Gram-negative bacteria, and some of Gram-positive bacteria, use

proteinaceous receptors for adsorption [48][49]. If the Group-2 viruses uses the same

protein receptor to attach to both Thermotogota and Firmicutes cells, the large

phylogenetic distance between these hosts offer an opportunity to identify possible

membrane protein receptors bioinformatically. It should be noted that this approach only

identifies possible protein receptors, while potential shared carbohydrate receptors will

not be identified. Four predicted membrane proteins with transmembrane helices where

identified in all the genomes carrying a Group-2 provirus (Supplementary Fig. S7). One

of these was the virus holin gene, leaving three receptors candidates; a ComEA family

DNA-binding protein, an oxaloacetate decarboxylase beta subunit and an ABC

transporter ATP-binding protein. Phylogenetic analyses revealed that the ComEA

homolog and the oxaloacetate decarboxylase are widely distributed among the

Thermotogota (Supplementary Fig. S7). In contrast, the ABC transporter is restricted to

*Petrotoga* and *Defluviitoga* (Supplementary Fig. S7), which could explain why the

Group-2 proviruses are only observed in these Thermotogota genera. Inspection of the

alignment of the proteins, revealed that these proteins show particularly high similarity in

the last C-terminal domain which is facing the exterior of the cell and could serve as a

virus target (Supplementary Fig. S8). Although experiments are needed to explore if any

of these proteins functions as receptor for these viruses, we suggest that the ABC-

transporter protein is the strongest candidate for a Group-2 virus receptor.

ABC-transporter proteins are, to our knowledge, not commonly identified as

bacterioviral receptors. However, membrane proteins with a variety of functions have

been shown to act as receptors [47]. Moreover,  *Lactoccocus* viruses from the siphoviral

c2 group have been shown to use membrane proteins Pip or YjaE as secondary receptors

[47][50], which both show similarity to ABC2 transporter domains.

**A role of membrane transport proteins and viruses in connecting Thermotogota and**

**Firmicutes?**

ABC-transporters are among the most frequently transferred genes both within the

Thermotogota and also between Thermotogota and other phyla, particularly Firmicutes

[51][7, 8]. The possibility that ABC transporters, and other transporters, can function as

virus receptors in the Thermotogota is intriguing, especially since transporters also are

among the morons discussed above. This suggests that acquiring a new transporter,

perhaps via a virus infection, might result in the cell not only acquiring a new function

but also becoming susceptible to a new virus. This virus might carry another transporter

gene, which can introduce yet another virus, resulting in a ratchet like process. Using

transporters as receptors will therefore not only provide the virus with the wide host

range inferred for both Thermotogota virus groups but could also make viruses the

vehicles on the highways of gene sharing observed between the Thermotogota and

Firmicutes. Membrane transport protein encoding genes, including ABC transporters,

have been observed in several other viruses [52], thus this process could also operate on

other genomes. This is contrary to a role commonly assigned to morons where they often

confer resistance to infections by other viruses [40]. Further studies and experiments are

needed to investigate if such ratchet processes are indeed occurring in natural systems.

However, regardless of the functions of the morons in the *Thermotogota* proviruses, the

observation of shared viruses among organisms from different phyla demonstrates that

viruses are key actors in the biosphere acting as engines of evolution.

**Acknowledgements**

This work is supported by a Norwegian research council award (project no. 180444/V40) to C.L.N, and by the Sino-French LIA/PRC 1211 MicrobSea for J.L.  Strains were obtained from the Université de Bretagne Occidentale Culture Collection (UBOCC, Plouzané, France, www.univ-brest.fr/ubocc). We thank Emily St John and Stéphane l'Haridon who provided the *Marinitoga lauensis* proviral sequence.

**Competing Interests:**

The authors declare no conflict of interest.

**References**

1.  Pollo SMJ, Zhaxybayeva O, Nesbø CL. Insights into thermoadaptation and the evolution of mesophily from the bacterial phylum *Thermotogae*. *Can J Microbiol* 2015; **61**: 655–670.

2.  Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, et al. A new view of the tree of life. *Nat Microbiol* 2016; **1**: 16048.

3.  Bhandari V, Gupta RS. The Phylum Thermotogae. In: Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F (eds). *The Prokaryotes*. 2014. Springer Berlin Heidelberg, Berlin, Heidelberg, pp 989–1015.

4.    Nesbø CL, S Swithers K, Dahle H, Haverkamp TH, Birkeland N-K, Sokolova T, et

al. Evidence for extensive gene flow and Thermotoga subpopulations in subsurface

and marine environments. *ISME J* 2015; **9**: 1532–1542.

5.    Foght JM, Gieg LM, Siddique T. The microbiology of oil sands tailings: past,

present, future. *FEMS Microbiol Ecol* 2017; **93**.

6.    Nesbø CL, Charchuk R, Pollo SMJ, Budwill K, Kublanov IV, Haverkamp THA, et

al. Genomic analysis of the mesophilic Thermotogae genus *Mesotoga* reveals

phylogeographic structure and genomic determinants of its distinct metabolism:

Comparative genomic analysis of *Mesotoga*. *Environ Microbiol* 2019; **21**: 456–470.

7.    Nelson KE, Clayton RA, Gill SR, Gwinn ML, Dodson RJ, Haft DH, et al. Evidence

for lateral gene transfer between Archaea and Bacteria from genome sequence of

Thermotoga maritima. *Nature* 1999; **399**: 323–329.

8.    Zhaxybayeva O, Swithers KS, Lapierre P, Fournier GP, Bickhart DM, DeBoy RT, et

al. On the chimeric nature, thermophilic origin, and phylogenetic placement of the

Thermotogales. *Proc Natl Acad Sci* 2009; **106**: 5865–5870.

9.    Nesbø CL, Dlutek M, Doolittle WF. Recombination in Thermotoga: Implications

for Species Concepts and Biogeography. *Genetics* 2006; **172**: 759–769.

10.   Magnabosco C, Biddle JF, Cockell CS, Jungbluth SP, Twing KI. Biogeography,

Ecology, and Evolution of Deep Life. In: Orcutt BN, Daniel I, Dasgupta R (eds).

*Deep Carbon*, 1st ed. 2019. Cambridge University Press, pp 524–555.

11.   Anderson RE, Brazelton WJ, Baross JA. The Deep Viriosphere: Assessing the Viral

Impact on Microbial Community Dynamics in the Deep Subsurface. *Rev Mineral*

*Geochem* 2013; **75**: 649–675.

12. Cobián Güemes AG, Youle M, Cantú VA, Felts B, Nulton J, Rohwer F. Viruses as

Winners in the Game of Life. *Annu Rev Virol* 2016; **3**: 197–214.

13. Engelhardt T, Kallmeyer J, Cypionka H, Engelen B. High virus-to-cell ratios

indicate ongoing production of viruses in deep subsurface sediments. *ISME J* 2014;

**8**: 1503–1509.

14. Walsh EA, Kirkpatrick JB, Pockalny R, Sauvage J, Spivack AJ, Murray RW, et al.

Relationship of Bacterial Richness to Organic Degradation Rate and Sediment Age

in Subseafloor Sediment. *Appl Environ Microbiol* 2016; **82**: 4994–4999.

15. Lossouarn J, Nesbø CL, Mercier C, Zhaxybayeva O, Johnson MS, Charchuck R, et

al. 'Ménage à trois': a selfish genetic element uses a virus to propagate within

Thermotogales. *Environ Microbiol* 2015; **17**: 3278–3288.

16. Mercier C, Lossouarn J, Nesbø CL, Haverkamp THA, Baudoux AC, Jebbar M, et al.

Two viruses, MCV1 and MCV2, which infect Marinitoga bacteria isolated from

deep-sea hydrothermal vents: functional and genomic analysis. *Environ Microbiol*

2018; **20**: 577–587.

17. Markowitz VM, Chen I-MA, Chu K, Szeto E, Palaniappan K, Pillay M, et al.

IMG/M 4 version of the integrated metagenome comparative analysis system.

*Nucleic Acids Res* 2014; **42**: D568–D573.

18. Rice P, Longden I, Bleasby A. EMBOSS: The European Molecular Biology Open

Software Suite. *Trends Genet* 2000; **16,**: 276–277.

19. Lima-Mendez G, Van Helden J, Toussaint A, Leplae R. Prophinder: a

computational tool for prophage prediction in prokaryotic genomes. *Bioinformatics*

2008; **24**: 863–865.

20.   Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. PHAST: A Fast Phage Search

Tool. *Nucleic Acids Res* 2011; **39**: W347–W352.

21.   Akhter S, Aziz RK, Edwards RA. PhiSpy: a novel algorithm for finding prophages

in bacterial genomes that combines similarity- and composition-based strategies.

*Nucleic Acids Res* 2012; **40**: e126–e126.

22.   O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al.

Reference sequence (RefSeq) database at NCBI: current status, taxonomic

expansion, and functional annotation. *Nucleic Acids Res* 2016; **44**: D733–D745.

23.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al.

BLAST+: architecture and applications. *BMC Bioinformatics* 2009; **10**: 421.

24.   Darling AE, Mau B, Perna NT. progressiveMauve: Multiple Genome Alignment

with Gene Gain, Loss and Rearrangement. *PLoS ONE* 2010; **5**: e11147.

25.   Guy L, Roat Kultima J, Andersson SGE. genoPlotR: comparative gene and genome

visualization in R. *Bioinformatics* 2010; **26**: 2334–2335.

26.   Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, et al. Circos:

An information aesthetic for comparative genomics. *Genome Res* 2009; **19**: 1639–

1645.

27.   Zimmermann L, Stephens A, Nam S-Z, Rau D, Kübler J, Lozajic M, et al. A

Completely Reimplemented MPI Bioinformatics Toolkit with a New HHpred Server

at its Core. *J Mol Biol* 2018; **430**: 2237–2243.

28.   Bolduc B, Jang HB, Doulcier G, You Z-Q, Roux S, Sullivan MB. vConTACT: an

iVirus tool to classify double-stranded DNA viruses that infect *Archaea* and

*Bacteria*. *PeerJ* 2017; **5**: e3243.

29.  Bin Jang H, Bolduc B, Zablocki O, Kuhn JH, Roux S, Adriaenssens EM, et al.

Taxonomic assignment of uncultivated prokaryotic virus genomes is enabled by

gene-sharing networks. *Nat Biotechnol* 2019; **37**: 632–639.

30.  Lopes A, Tavares P, Petit M-A, Guérois R, Zinn-Justin S. Automated classification

of tailed bacteriophages according to their neck organization. *BMC Genomics* 2014;

**15**: 1027.

31.  Moraru C, Varsani A, Kropinski AM. VIRIDIC – a novel tool to calculate the

intergenomic similarities of prokaryote-infecting viruses. 2020. Microbiology.

32.  Nishimura Y, Yoshida T, Kuronishi M, Uehara H, Ogata H, Goto S. ViPTree: the

viral proteomic tree server. *Bioinformatics* 2017; **33**: 2379–2380.

33.  Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis

of large phylogenies. *Bioinformatics* 2014; **30**: 1312–1313.

34.  Geslin C, Le Romancer M, Erauso G, Gaillard M, Perrot G, Prieur D. PAV1, the

First Virus-Like Particle Isolated from a Hyperthermophilic Euryarchaeote,

"Pyrococcus abyssi". *J Bacteriol* 2003; **185**: 3888–3894.

35.  Haverkamp THA, Geslin C, Lossouarn J, Podosokorskaya OA, Kublanov I, Nesbø

CL. Thermosipho spp. immune system differences affect variation in genome size

and geographical distributions. *Genome Biol Evol* 2018.

36.  Walker PJ, Siddell SG, Lefkowitz EJ, Mushegian AR, Adriaenssens EM, Dempsey

DM, et al. Changes to virus taxonomy and the Statutes ratified by the International

Committee on Taxonomy of Viruses (2020). *Arch Virol* 2020; s00705-020-04752–x.

37.  Adriaenssens E, Brister JR. How to Name and Classify Your Phage: An Informal

Guide. *Viruses* 2017; **9**: 70.

38.   Catalano CE. The terminase enzyme from bacteriophage lambda: a DNA-packaging

machine. *Cell Mol Life Sci CMLS* 2000; **57**: 128–148.

39.   Cumby N, Davidson AR, Maxwell KL. The moron comes of age. *Bacteriophage*

2012; **2**: e23146.

40.   Taylor VL, Fitzpatrick AD, Islam Z, Maxwell KL. The Diverse Impacts of Phage

Morons on Bacterial Fitness and Virulence. *Adv Virus Res* 2019; **103**: 1–31.

41.   Roux S, Enault F, Hurwitz BL, Sullivan MB. VirSorter: mining viral signal from

microbial genomic data. *PeerJ* 2015; **3**: e985.

42.   Rath D, Amlinger L, Rath A, Lundgren M. The CRISPR-Cas immune system:

Biology, mechanisms and applications. *Biochimie* 2015; **117**: 119–128.

43.   Borton MA, Daly RA, O'Banion B, Hoyt DW, Marcus DN, Welch S, et al.

Comparative genomics and physiology of the genus *Methanohalophilus* , a

prevalent methanogen in hydraulically fractured shale. *Environ Microbiol* 2018; **20**:

4596–4611.

44.   Bonilla Salinas M. Mahella australiensis gen. nov., sp. nov., a moderately

thermophilic anaerobic bacterium isolated from an Australian oil well. *Int J Syst*

*Evol Microbiol* 2004; **54**: 2169–2173.

45.   Iino T, Suzuki K -i., Harayama S. Lacticigenium naphtae gen. nov., sp. nov., a

halotolerant and motile lactic acid bacterium isolated from crude oil. *Int J Syst Evol*

*Microbiol* 2009; **59**: 775–780.

46.   Hong H, Kim S-J, Min U-G, Lee Y-J, Kim S-G, Jung M-Y, et al. Geosporobacter

ferrireducens sp. nov., an anaerobic iron-reducing bacterium isolated from an oil-

contaminated site. *Antonie Van Leeuwenhoek* 2015; **107**: 971–977.

47.   Stone E, Campbell K, Grant I, McAuliffe O. Understanding and Exploiting Phage–

Host Interactions. *Viruses* 2019; **11**: 567.

48.   Zhang Z, Yu F, Zou Y, Qiu Y, Wu A, Jiang T, et al. Phage protein receptors have

multiple interaction partners and high expressions. *Bioinformatics* 2020; **36**: 2975–

2979.

49.   Bertozzi Silva J, Storms Z, Sauvageau D. Host receptors for bacteriophage

adsorption. *FEMS Microbiol Lett* 2016; **363**: fnw002.

50.   Millen AM, Romero DA. Genetic determinants of lactococcal C2viruses for host

infection and their role in phage evolution. *J Gen Virol* 2016; **97**: 1998–2007.

51.   Nesbø CL, Nelson KE, Doolittle WF. Suppressive Subtractive Hybridization

Detects Extensive Genomic Diversity in Thermotoga maritima. *J Bacteriol* 2002;

**184**: 4475–4488.

52.   Greiner T, Moroni A, Van Etten J, Thiel G. Genes for Membrane Transport

Proteins: Not So Rare in Viruses. *Viruses* 2018; **10**: 456.

53.   L'Haridon S, Gouhier L, John ESt, Reysenbach A-L. Marinitoga lauensis sp. nov., a

novel deep-sea hydrothermal vent thermophilic anaerobic heterotroph with a

prophage. *Syst Appl Microbiol* 2019; **42**: 343–347.

54.   Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new

developments. *Nucleic Acids Res* 2019; **47**: W256–W259.

55.   Lee MD. GToTree: a user-friendly workflow for phylogenomics. *Bioinformatics*

2019; **35**: 4162–4164.

**Table 1.** CRISPR matches to provirus genomes. Proviruses that have been induced and

shown to produce virus particles are indicated by a *. Spacers were mapped to virus

sequences in Geneious v.10 allowing up to 10% mismatches. Numbers of spacers from

each genome mapping to the provirus is given in parentheses. Matches to spacers from

the host genome are noted as a self-match.

| provirus name | CRISPR spacer match from (number of spacers) |
|---|---|
| **KPV1** | *Kosmotoga olearia* (1) |
| **MHV1** | *Pseudothermotoga elfii* NBRC107921 (1), *Marinitoga* sp. 1154 (1) |
| **MCV1*** | *Marinitoga* sp. 1154 (2) |
| **MCV2*** | *Marinitoga* sp. 1155 (1) self-match, *Marinitoga* sp. 1154 (2) |
| **MPV1*** | *Marinitoga* sp. 1137 (1) |
| **M1135** | *Marinitoga* sp. 1155 (1) |
| **M1137V1** | *Pseudothermotoga elfii* NBRC107921 (1), *Marinitoga* sp. 1135 (2) |
| **M1137V2** | *M. piezophila* (1) |
| **M1138** | *T. africanus* TCF52B (1), *T. melanesiensis* (1), *Pseudothermotoga elfii* NBRC107921 (1), *Marinitoga* sp. 1137 (1) |
| **TAV1*** | *T. africanus* H17ap60334 (3) self -match, *T. africanus* TCF52B (2), *T. africanus* Ob7 (1), *T. melanesiensis* (2), *Thermotoga maritima* 2812B (1), *Thermotoga* sp. EMP (1), *Thermotoga* sp. XYL54 (3), *Thermotoga* sp. CELL2 (3), *Thermotoga* sp. TBGT1766 (3), *Thermotoga* sp. TBGT1765 (4), *Thermotoga* sp. A7A (1), *Pseudothermotoga elfii lettingae* (4), *Pseudothermotoga elfii* NBRC107921 (9), *Pseudothermotoga elfii* DSM9442 (2) |

| | |
|---|---|
| **TM1** | *T. melanesiensis* (1) self-match, *T. africanus* TCF52B (1), *T. africanus* Ob7 (1), *Pseudothermotoga elfii lettingae* (2), *Thermotoga* sp. TBGT1765 (1), *Thermotoga* sp. Mc24 (1), *Pseudothermotoga elfii* NBRC107921 (1) |
| **T1074V1** | *Thermosipho affectus* Bl1070 (1), *Thermosipho affectus* 1223 (3), *Thermosipho affectus* Bl1063 (1) |

643 **Table 2**. Homologous genes detected among Group-1 and Group-2 provirus proteins

644 using blastp and expected value cut-off at $1e^{-01}$. For Recombinase homologs see Fig. 5.

| Annotation | Group 1 homologs | Group 2 homologs |
|---|---|---|
| **Aldo/keto reductase** | BLS00_RS05685 (GPV1) | X929_01195 (POV1) |
| **ATP-binding protein** | BLS00_RS05500 (GPV1), IX53_RS00630 (KPV1) | DTL3_1118 (DTV1) |
| **Capsid protein** | LN42_00565 (M1137V1), LH53_05775 (MTOLDC1) | NAAC61_01950 (P8T1HF07V1), X929_01335 (POV1), DTL3_1133 (DTV1) |
| **Cro/Cl family transcriptional regulator** | EK18DRAFT_00606 (MLV1), H17ap60334_04807 (TAV1), BUA62_RS02415 (MHV1), LN42_00465 (M1137V1) | NAAC61_02080 (P8T1HF07V1) |
| **DNA methylase** | BLS00_RS05440 (GPV1), BUA62_RS02350 (MHV1), | NAAC61_02100 (P8T1HF07V1) |

| | | |
|---|---|---|
| | EK18DRAFT_00547 (MLV1) | |
| **DNA methylase** | Tmel_1480 (TMV1), Marpi_0296 (MPV1), XO11_00615 (M1138V1), EK18DRAFT_00605 (MLV1), LH53_05700 (MTOLDCV1) | DTL3_1116, DTL3_1121, DTL3_1122 (DTV1), NAAC61_01990 (P8T1HF07V1), X929_01285 (POV1) |
| **Endonuclease-like protein** | Tmel_1473 (TMV1) | NAAC61_02075 (P8T1HF07V1) |
| **HNH endonuclease** | EK18DRAFT_00580 (MLV1) | NAAC61_02005 (P8T1HF07V1), X929_01255 (POV1) |
| **Hypothetical protein/ Helicase** | XO11_00565 (M1138V1), XO10_00565 (M1135V1) | X929_01210 (POV1) |
| **Hypothetical protein** | UF08_38 (MCV1) | X925_04785 (P9T1HF07V1) |
| **Hypothetical protein** | XO08_07915 (T1074V1), LH53_05685 (MTOLDCV1) | DTL3_1144 (DTV1), NAAC61_01815* (P8T1HF07V1) |
| **Hypothetical protein** | Marpi_0334 (MPV1), LN42_00625 (M1137V1), XO11_00700 (M1138V1), H17ap60334_04967 (TAV1), Tmel_1449 (TMV1), XO08_07890 (T1074V1), LH53_05725 (MTOLDCV1), | DTL3_1141 (DTV1) |

| | EK18DRAFT_00564 (MAV1) | |
|---|---|---|
| **Hypothetical protein** | XO08_07730 (T1074V1) | DTL3_1114 (DTV1) |
| **Hypothetical protein** | LH53_05765 (MTOLDCV1) | DTL3_1134 (DTV1) |
| **Hypothetical protein** | EK18DRAFT_00608 (MAV1) | DTL3_1111 (DTV1) |
| **Hypothetical protein** | EK18DRAFT_00582 (MAV1) | X929_01400 (POV1) |
| **Hypothetical protein** | IX53_RS00750 (KPV1) | NAAC61_02025 (P8T1HF07V1) |
| **Hypothetical protein** | IX53_RS00690 (KPV1) | Pmob_0430* (PMV1) |
| **Hypothetical protein** | LN42_01810 (M1137V2) | NAAC61_02035 (P8T1HF07V1) |
| **Hypothetical protein** | Marpi_0313 (MPV1) | DTL3_1106 (DTV1) |
| **Hypothetical protein/ transporter** | Marpi_0315 (MPV1), XO10_00580 (1135V1) | DTL3_1155 (DTV1), X929_01440 (POV1), X925_04770 (P9T1HF07V1), Pmob_0433 (PMV1) |
| **Phage minor structural protein** | BUA62_RS02360 (MHV1) | X929_01385 (POV1), DTL3_1143 (DTV1) |
| **Phage portal protein** | LH53_05785 (MTOLDCV1) | NAAC61_01960 (P8T1HF07V1), X929_01325 (POV1),  DTL3_1131 (DTV1) |
| **Primase / Terminase** | EK18DRAFT_00577 (MAV1), IX53_RS00695 (KPV1) | X929_01235 (POV1) |
| **Tail protein** | UF08_32 (MCV1), BUA62_RS02555 (MHV1), | X929_01375 (POV1) |

| | Tmel_1449 (TMV1), H17ap60334_04967 (TAV1) | |
|---|---|---|
| **Transcription regulator** | Marpi_0298 Marpi_0297 (MPV1),  UF08_76 UF08_59 UF08_65 (MCV1), X274_02275 (MCV2), XO08_07670 (T1074V1), BUA62_RS02435 BUA62_RS02375 BUA62_RS02385 (MHV1), XO11_00475 (M1138), XO10_00515 XO10_00685 (M1135V1),  LN42_01800 LN42_01805 LN42_01820 (M1137V2), H17ap60334_04807 (TAV1), Tmel_1478 Tmel_1479 (TMV1) | DTL3_1158 (DTV1) |
| **Transposase** | Tmel_1465 (TMV1) | DTL3_1117 (DTV1) |
| **XRE family transcriptional regulator** | XO11_00475 (M1138V1), UF08_9 (MCV1), XO10_00475 (M1135) | NAAC61_02010 (P8T1HF07V1) |

645    *recombinase

**Fig. 1**. **Comparison of proviruses identified in eight different Thermotogota genera**.

For genera where multiple different viruses were detected, only one virus is represented

as viruses from the same genus were highly similar (Fig. 2 and 3). For this figure all

tblastx matches between the viruses were drawn as links with the percent identity shown

in histograms attached to each link. Blast matches were drawn in the following order;

DTV1 (dark blue), GPV1 (grey), TV1 (orange), MPV1(red), KPV1 (dark green),

MTOLDCV1 (light green), MLV1 (purple) and P8T1HF07V1 (light blue). Reciprocal

best matches are therefore hidden in many cases (e.g. the dark blue links between DTV1

and P8T1HF07V1), however, these are still visible in the histogram connecting to these

links. The figure was drawn in Circos [26].

**Fig. 2. Comparison of the Thermotogota Group-1 provirus sequences.** (Pro)virus

name and the genus of the host it belongs to are shown on the left side. Proviruses that

have been induced and shown to produce virus particles are indicated by a *. Proviruses

determined to be incomplete are indicated by (in). The origin of the sequences are

(accession number); GPV1 - *Geotoga petraea* WG14 (FMYV01), MPV1 - *Marinitoga*

*piezophila* (NC_016751), MCV1 – *Marinitoga camini* 1197 (AZAY01), MCV2 -

*Marinitoga camini* 1155 (AZAX01), MHV1 - *Marinitoga hydrogenitolerans* DSM 16785

(FQUI01), M1137V2 and M1137V2 - *Marinitoga* sp. 1137 (CP009471), M1135V1 -

*Marinitoga* sp. 1135 (LBEZ01), M1138V1 - *Marinitoga* sp. 1138 (LBFA01), MLaV1 -

*Marinitoga lauensis* (QHLX01), TMV1 - *Thermosipho melanesiensis* BI429

(CP000716), TAV1 – *Thermosipho africanus* H17ap60334 (AJIP01), T1074V1 –

*Thermosipho* sp. 1074 (LBEX01), KPV1 – *Kosmotoga pacifica* SLHLJ1 (CP011232),

MAV1 - *Mesoaciditoga lauensis* cd-1655R (JQJI01) and MTOLDCV1 – *Mesotoga* sp.

SC_TOLDC (AYSI01). We also included in this figure, MLaV1 - *Marinitoga lauensis*

(QHLX01) reported in reference [53] after our bioinformatic screening was performed.

The regions with significant pairwise tblastx similarity scores are connected (minimum

identity: 30 %, minimum lenght : 100 nucleotides), red indicates that sequence is in the

same direction while blue indicates that the similar sequences are on opposite strands.

The division of the provirus genomes into functional modules is approximate, as it is

based only on gene annotations. Gene functions are color-coded as follows; red:

integration, yellow: transcriptional regulation, orange: DNA metabolism, green: DNA

packaging and head, light blue: head to tail, dark blue: tail, beige: HNH endonuclease,

pink: lysis, black: other (morons), grey: hypothetical proteins). Some gene annotations

are included and abbreviated; Ser recomb: serine recombinase, LexA: LexA repressor,

ParB: ParB-like nuclease, RecT: RecT family recombinase, dUTP hydrolase:

deoxyuridine 5'-triphosphate nucleotidohydrolase,  ssb: single stranded DNA-binding

protein, tss: terminase small subunit, tls: terminase large subunit, mcp: major capsid

protein, tail tape: tail tape measure protein, rRNA lsm: ribosomal RNA large subunit

methyltransferase, flagella bbp: flagella basal-body protein, DnaC: DnaC replication

protein, DnaD: DnaD replication protein, DNA pol sc: DNA polymerase sliding clamp,

SecB: SecB protein-export protein, rep organizer :replisome organizer, RusA: RusA

family crossover junction endodeoxyribonuclease, Cys peptidase: cysteine peptidase,

sulfate AT: sulfate adenylyltransferase subunit 2, PAPS reductase: phosphoadenosine

phosphosulfate reductase, CW hydrolase: cell wall-associated hydrolase, MazF: MazF

692 endoribonuclease, dsbr: DNA double-strand break repair protein, metal bp: metal-binding

693 protein, CMP hydrolase: cytidine 5′-monophosphate hydrolase. The figure was produced

694 using genoPlotR (Guy *et al.*, 2010).

696 **Fig. 3. Comparison of the putative Thermotogota and Group-2 provirus sequences**

697 **and their Firmicutes closest relatives.** Virus name and the genus the host belongs to are

698 included. The origin of the sequences are (accession number); DTV1 - *Defluviitoga*

699 *tunisiensis* DTL3 (LN824141), POV1 – *Petrotoga olearia* DSM 13574 (AZRL01),

700 LNV1 – *Lacticigenium naphtae* DSM 19658 (AUEG01), P8T1HF07V1 - *Petrotoga* sp.

701 8T1HF07 (AUPG01), GSV1 - *Geosporobacter ferrireducens* IRF9 (CP017269), MAUV1

702 - *Mahella australiensis* 50-1 BON (NC_015520). The regions with significant pairwise

703 tblastx similarity scores are connected (mininum identity: 30 %, minimum lenght: 100

704 nucleotides), red indicates that sequence is in the same direction while blue indicates that

705 the similar sequences are on opposite strands. The division of the provirus genomes into

706 functional modules is approximate, as it is based only on gene annotations. Selected

707 informative ORF annotations are shown. Gene functions are color-coded: red:

708 integration, yellow: transcriptional regulation, orange: DNA metabolism, green: DNA

709 packaging and head, light blue: head to tail, dark blue: tail, white: HNH endonuclease,

710 pink: lysis, black: other (morons), purple: transposase, grey: hypothetical proteins). Some

711 gene annotations are included and abbreviated; HMT ATPase: heavy metal translocating

712 ATPase, FMN reductase: flavine mono nucleotide reductase, HAD family phosphatase:

713 haloacid dehalogenase superfamily of hydrolase). The figure was produced using

714 genoPlotR (Guy *et al.*, 2010).

**Fig. 4. Maximum likelihood tree of recombinase genes.** Sequences homologous to the

recombinase from all Thermotogota viruses and the three Firmicutes viruses in Fig. 4

were identified by blastp searches against Genbank and the twenty top matches of each

virus homolog were retrived. The resulting dataset was filtered keeping only unique

sequences and the sequences were aligned using MAFFT as implemented in Geneious v.

10.  A FastTree analysis was performed first in Genenious v. 10 and clades containing

numerous almost identical Firmicute sequences were trimmed. The phylogenetic tree

displayed was constructed using RAxML as implemented in Geneious v. 10 with a

GAMMA WAG model and 100 bootstrap replicates. Bootstrap support > 70% is shown

on branches as circles, with the size corresponding to the strength of support. (Pro)virus

name is added in to the host name for proteins from the Thermotogota proviruses (e.g.

***Thermosipho africanus* TAV1**), as well as from the Firmicute proviruses in Fig. 3.

Incomplete viruses are indicated by (in). Thermotogota homologs from genomes with no

detected provirus are indicated by a star. Numbers attached to each sequence represent

database accession numbers. Some proteins have identical sequences in more than one

organisms. The protein labelled 'Bacteria inc. Petrotoga olearia POLV1

*Fervidobacterium* spp.' corresponds to accession number WP_011994748.1 and is found

in *Fervidobacterium nodosum* Rt17-B1 (NC_009718.1), *Fervidobacterium pennivorans*

DSM 9078 (NC_017095.1), *Fervidobacterium islandicum* (NZ_CP014334.1) and

*Fervidobacterium gondwane*nse DSM 13020 (FRDJ01), *Petrotoga olearia* (PNR98053)

and *Coprothermobacter proteolyticus* (PXJB01). The protein labelled 'Bacteria inc.

*Mahella australiensis* MAV1, *Pseudothermotoga elfii*' corresponds to accession number

WP_013782344.1 found in *Clostridium* sp. SYSU GA15002T (NZ_CP040924.1) and

*Thermoanaerobacter thermocopriae* JCM 7501 (NZ_KI912455.1), *Pseudothermotoga*

*elfii* and MAV1 from *Mahella australiensis.* All branches without labels represent

Firmicutes sequences. The tree was visualized in iTOL [54].

**Fig. 5. Electron micrographs, stained with 2% uranyl acetate, of the TAV1 virus**

**particle (panel a) and vesicles produced by *Thermosipho africanus* H17ap60334**

**(panel b) and *Petrotoga* sp. 8T1HF07.NaAc.6.1 (panel c, d)**. The TAV1 virus particle

shows a typical Siphoviridae morphology (panel a). The virus particles were observed

after adding 5µg/ml mitomycin C at the beginning of the exponential growth phase of

*Thermosipho africanus* H17ap60334 growing at 65°C. Induction of the provirus in

*Petrotoga* sp. 8T1HF07.NaAc.6.1 was attempted by adding 5µg/ml mitomycin C at the

beginning of the exponential growth phase at 55°C. No virus particles were observed,

however, large amounts of vesicles were observed (panel b and d), as well as sunflower-

like structures (arrows in panel c). Panel b shows the vesicles attached to the flagellum.

**Fig. 6. Virus genes in genomes with no provirus.** Predicted proteins from proviruses

were compared to a database of protein from 59 Thermotogota genomes (from eight

genera) with no proviruses using blastp. This resulted in a total 875 matches for 107

provirus proteins to proteins from 50 genomes (> 50% identity and 60% coverage). The

Thermotogota genera represented in this analysis is shown on the x-axis, and the number

of provirus matches pr genome searched for each genus is shown on y-axis.

**Supplementary Figures:**

**Supplementary Fig. S1. A: Network of provirus genomes calculated in VContact2.**

Proviruses a maximum of three nodes away from MPV1 and P8T1HF07V1 are show.

The Thermotogota proviruses are colored blue, viruses from Firmicutes are green and

viruses infecting other taxa are colored orange. The quality score calculated by

ClusterOne for the Group-1 cluster was 0.94 and p=0.00004, the quality score for the

Group-2 cluster was 0.83 and p=0.00557971. **B: Phylogenetic tree of Thermotogota**

**genomes**. Identified proviruses are indicated next to their respective host genera.

Available Thermotogota genomes in Genbank, and a list of assembly accession numbers

were used as input for the GToTree pipeline, which downloads the genomes, extracts

single copy gene (SCGs) sets and aligns them based on a given hmm-gene set [55]. The

Bacterial hmm-set with 74 target genes was used, and the resulting alignment was

imported into Geneious Prime 2020.1.2 where sites with more than 50% gaps were

removed, resulting in an alignment of 11,003 amino acid positions. The phylogenetic tree

was constructed using FastTree and visualized in iTOL [54].

**Supplementary Fig. S2**. The region of the phylogenetic tree, calculated from virus

proteins by Vip-tree [32], that include the Group-1 and Group-2 Thermotogota and

Firmicutes proviruses that were determined to be likely complete (indicated by red stars).

Taxonomy of the most closely related viruses and their host is indicated as color bars in

front of the virus name for each terminal leaf on the tree.

**Supplementary Fig. S3. Maximum likelihood trees of Terminase genes.** Sequences

homologous to the Terminases from all Thermotogota Group 1 viruses were identified by

blastp searches against Genbank retrieving the twenty top matches of each virus

homolog. The resulting dataset was filtered keeping only unique sequences and the

sequences were aligned using MAFFT as implemented in Geneious v. 10.  A FastTree

analysis was performed first in Genenious v. 10 and clades containing numerous almost

identical sequences were trimmed.  The phylogenetic trees displayed were constructed

using RAxML as implemented in Geneious v. 10 with a GAMMA-WAG substitution

model and 100 bootstrap replicates. Bootstrap support > 70% is shown on branches as

circles, with the size corresponding to the strength of support. Virus name is added in

bold to the host name in the tree for the Thermotogota viruses (e.g. *Thermosipho*

*africanus* **TAV1**). Thermotogota homologs from genomes with no detected provirus are

also given in bold (e.g. *Thermotoga* **sp. RQ7**). Numbers attached to each sequence

represent database accession numbers. The tree was visualized in iTOL [54].

**Supplementary Fig. S4. Comparison of the Thermotogota Group 2 provirus**

**sequences including two incomplete ones.** Virus name and the genus the host belongs to

is indicated. Proviruses determined to be incomplete are indicated by (in). The origin of

the sequences are (accession number in parenthesis); DTV1 - *Defluviitoga tunisiensis*

DTL3 (LN824141), POV1 – *Petrotoga olearia* DSM 13574 (AZRL01), P8T1HF07V1 -

*Petrotoga* sp. 8T1HF07 (AUPG01), PMOV1 - *Petrotoga olearia* (NZ_AZRL01),

P9T1HF07V1 - *Petrotoga* sp. P9T1HF07 (AZRI01). The regions with significant

pairwise BLASTN similarity scores are connected, red indicates that sequence is in the

same direction while blue indicates that the similar sequences are on opposite strands.

Selected informative ORF annotations are shown. The division of the virus into

functional modules is approximate, as it is based only on gene annotations. The Fig. was

produced using genoPlotR (Guy *et al*., 2010).

**Supplementary Fig. S5. Overview of CRISPR spacer sequences from Thermotoga**

**isolates from the Troll oil reservoir mapped on to the TAV1 genome.** Mapping and

visualization was performed in Geneious v. 10 and maximum of one mismatch was

allowed.

**Supplementary Fig. S6. Maximum likelihood trees of the moron annotated as a**

**cadmium transporter**. Sequences homologous to the gene from all Group 2 viruses were

identified by blastp searches against Genbank retrieving the twenty top matches of each

virus homolog as well as searching a custom database containing genes from both

Thermotogota genomes with or without a provirus. The resulting dataset was filtered

keeping only unique sequences and the sequences were aligned using MAFFT as

implemented in Geneious v. 10. A FastTree analysis was performed first in Genenious v.

10 and clades containing numerous almost identical sequences were trimmed.  The

phylogenetic trees displayed were constructed using RAxML as implemented in

Geneious v. 10 with a GAMMA WAG model and 100 bootstrap replicates. Bootstrap

support > 70% is shown on branches as circles, with the size corresponding to the

strength of support. Virus name is added in bold to the host name in the tree for the

Thermotogota viruses (e.g. ***Defluviitoga tunisiensis* DTV1**). Thermotogota homologs

from genomes with no detected provirus are also given in bold (e.g. ***Fervidobacterium***

**sp. SC NGM5 G05**). Numbers attached to each sequence represent database accession

numbers. The tree was visualized in iTOL [54].

**Supplementary Fig. S7. Maximum likelihood trees of potential virus receptor genes;**

**Panel A: ComEA, Panel B: oxaloacetate decarboxylase and Panel C: ATP-binding**

**cassette, subfamily B.** Sequences homologous to the genes from the *Petrotoga*,

*Defluviitoga* and the three Firmicute genomes carrying Group-2 proviruses were

identified by blastp searches against Genbank, retrieving the twenty top matches of each

homolog, as well as the hundred top matches from *Petrotoga* sp. 8T1HF07 to ensure as

many Thermotogota homologes as possible were included. The resulting dataset was

filtered keeping only unique sequences and the sequences were aligned using MAFFT as

implemented in Geneious v. 10.  The phylogenetic trees displayed were constructed using

RAxML as implemented in Geneious v. 10 with a GAMMA WAG model and 100

bootstrap replicates. Bootstrap support > 70% is shown on branches as circles, with the

size corresponding to the strength of support. The genomes with Group-2 viruses are

displayed in bold and clades containing sequences from the same genus or with identical

taxonomic assignment are collapsed. The trees were visualized in iTOL [54].

**Supplementary Fig. S8. Alignment of ATP-binding cassette, subfamily B homologs.**

Sites identical among all the sequences are highlighted. Transmembrane regions were

predicted by TMHMM v. 2 at the TMHMM Server v. 2.0

(http://www.cbs.dtu.dk/services/TMHMM-2.0/). Identical sites are highlighted with the

same color and variable sites are in grey.

Matches to proirus genes pr. genome

■ All □ Group-1 ■ Group-2