Decreased homotopic interhemispheric functional connectivity in children with

autism spectrum disorder

Running title: Homotopic connectivity in autistic children

Shuxia Yao*, Menghan Zhou, Yuan Zhang, Qianqian Zhang, Zhongbo Zhao, Xi Jiang,

Xiaolei Xu, Benjamin Becker, Keith M. Kendrick*

The Clinical Hospital of Chengdu Brain Science Institute, MOE Key Laboratory for

NeuroInformation, Center for Information in Medicine, University of Electronic Science

and Technology of China, Chengdu, Sichuan 611731, China

*Corresponding authors

Correspondence: Shuxia Yao or Keith M. Kendrick, No.2006, Xiyuan Ave., West

Hi-Tech Zone, Chengdu, Sichuan 611731, China. Fax: +86-28-61830811. Email:

yaoshuxia@uestc.edu.cn or k.kendrick.uestc@gmail.com

Acknowledgments

This study was supported by the National Natural Science Foundation of China (NSFC)

grants (grant number: 31700998, 31530032) and by the Department of Science and

Technology of Guangdong Province (grant number: 2018B030335001). The primary

funding source for ABIDE-II is from NIMH 5R21MH107045.

Declaration of interest: The authors declare no conflict of interest.

Lay Summary: Homotopic interhemispheric functional connectivity plays an important

role in synchronizing activity between the two hemispheres and is altered in adults and

adolescents with autism spectrum disorder (ASD). In the present study focused on

children with ASD, we have observed similar decreased homotopic connectivity in

extensive brain networks, suggesting that alterations in homotopic interhemispheric

connectivity may occur early in ASD and be a useful general biomarker across ages.

Abstract

While a number of functional and structural changes occur in large-scale brain

networks in autism spectrum disorder (ASD), reduced interhemispheric resting state

functional connectivity (rsFC) between homotopic regions may be of particular

importance as a biomarker. ASD is an early-onset developmental disorder and neural

changes are often age-dependent, reflecting dysregulated developmental trajectories,

although no studies have investigated whether homotopic interhemispheric rsFC

alterations occur in ASD children. The present study conducted a voxel-based homotopic

interhemispheric rsFC analysis in 152 ASD and 185 typically developing children under

age 10 and examined associations with symptom severity in the Autism Brain Imaging

Data Exchange datasets. Given the role of corpus callosum (CC) in interhemispheric

connectivity and reported CC volume changes in ASD we additionally examined whether

there were parallel volume changes in ASD children. Results demonstrated decreased

homotopic rsFC in ASD children in the medial prefrontal cortex, precuneus and posterior

cingulate cortex of the default mode network (DMN), the dorsal anterior cingulate cortex

of the salience network, the precentral gyrus and inferior parietal lobule of the mirror

neuron system, the lingual, fusiform and inferior occipital gyri of the visual processing

network and thalamus. Symptom severity was associated with homotopic rsFC in regions

in the DMN and visual processing network. There were no significant CC volume

changes in ASD children. The present study shows that reduced homotopic

interhemispheric rsFC in brain networks in ASD adults/adolescents is already present in

children of 5-10 years old and further supports their potential use as a general ASD

biomarker.

Keywords: Autism Spectrum Disorder; Children; Resting State; Homotopic

Interhemispheric Functional Connectivity; Corpus Callosum

Introduction

Autism spectrum disorder (ASD) is a heterogeneous neurodevelopmental disorder with core symptomatology generally characterized by deficits in social interaction, verbal and nonverbal communication and restrictive or stereotyped behavior (Kanner, 1943). Behavioral studies have revealed a variety of impairments in social-cognitive and affective domains in ASD patients, including deficits in joint attention (Charman, 2003), weak central coherence (Happé & Frith, 2006), avoidance of eye contact (Madipakkam, Rothkirch, Dziobek, & Sterzer, 2017), difficulties in face recognition (Fedor et al., 2018) and diminished interest and motivation in social interactions (Chevallier, Grèzes, Molesworth, Berthoz, & Happé, 2012; Morrison et al., 2017). Task-based neuroimaging studies have demonstrated that these deficits are mediated by dysfunctions in large-scale brain systems sub-serving social-cognitive and affective processes including the default mode network (DMN), mirror neuron system (MNS), theory of mind (ToM), and executive control networks (Dapretto et al., 2006; Herringshaw, Kumar, Rody, & Kana, 2018; Oberwelland et al., 2017; von dem Hagen, Stoyanova, Rowe, Baron-Cohen, & Calder, 2014; Wang, Dapretto, Hariri, Sigman, & Bookheimer, 2004).

The behavioral deficits in ASD and the need for diagnostics at an early age limit the application of task-based fMRI assessments in ASD. Consequently, resting state fMRI (rs-fMRI), which has the advantage of being task-free and providing comparable measures across different studies and sites, has been considered as a promising approach

for investigating neural abnormalities and for providing possible promising biomarkers in

ASD. In line with the task-based fMRI findings, ASD-associated alterations in the

intrinsic organization of the brain as assessed by resting state functional connectivity

(rsFC) have been found in the DMN, MNS, ToM and executive control networks (Chen,

Yang, Wu, Chuang, & Huang, 2019; Fishman, Keown, Lincoln, Pineda, & Müller, 2014;

Funakoshi et al., 2016; Monk et al., 2009), suggesting that large-scale aberrant functional

organization occurs both in a task-free and task-based conditions in ASD. Additional

brain networks showing rsFC abnormities in autistic patients have been reported in

salience, mesolimbic reward and language networks (Lee, Park, James, Kim, & Park,

2017; Supekar et al., 2018; Uddin et al., 2013). Furthermore, atypical developmental

trajectories in the functional and structural organization of these networks have also been

found in ASD (Ecker, Bookheimer, & Murphy, 2015; Geschwind & Levitt, 2007;

Washington et al., 2014).

Based on altered functional connectivity patterns, ASD has been proposed as a

developmental dysfunctional connectivity syndrome with local overconnectivity but

long-distance underconnectivity (Courchesne & Pierce, 2005; Geschwind & Levitt, 2007;

Lau, Leung, & Lau, 2019). Recent studies have focused on interhemispheric functional

connections particularly the homotopic ones, which may be of great importance in ASD.

Homotopic interhemispheric functional connectivity is a measurement of the rsFC

between each voxel in one hemisphere and its mirrored counterpart in the opposite

hemisphere (Stark et al., 2008; Zuo et al., 2010). As a key characteristic of the intrinsic

functional architecture of the brain, homotopic interhemispheric functional connections in

the brain are exceptionally strong and may therefore play an important role in

synchronizing activity between the two hemispheres (Stark et al., 2008). Anderson et al.

(2011) first investigated the homotopic interhemispheric rsFC in ASD and found

decreased homotopic interhemispheric rsFC between sensorimotor cortex, insula,

fusiform gyrus, superior temporal gyrus, and superior parietal lobule in ASD compared to

typically developing (TD) controls. Two other studies using the large Autism Brain

Imaging Data Exchange (ABIDE) multisite datasets (ABIDE-I or -II), which have the

advantage of greater statistical power, confirmed homotopic interhemispheric rsFC

alterations in ASD, particularly in the interhemispheric communication of the posterior

cingulate cortex (PCC), insula and thalamus (Di Martino et al., 2014 – ABIDE-I) and

across a variety of large-scale networks (King et al., 2019; ABIDE-I and -II). Including

gender as an additional covariate, Li et al. (2019) using the ABIDE-I dataset further

confirmed decreased homotopic interhemispheric connectivity in ASD subjects and

reported associations between interhemispheric functional connectivity in the PCC, insula

and superior temporal gyrus and social communication impairment severity using the

Autism Diagnostic Observation Schedule (ADOS). Thus, interhemispheric functional

connectivity changes may represent a particular promising biomarker for discriminating

ASD from TD individuals (Li et al., 2019). Note that some ASD-related rsFC changes are

age-dependent (Hull, Jacokes, Torgerson, Irimia, & van Horn, 2017) and there is also

evidence that homotopic interhemispheric connectivity shows different developmental

trajectories in ASD compared with TD individuals (Kozhemiako et al., 2018). It is

therefore possible that interhemispheric communication dysfunctions may also represent

a result of dysregulated developmental trajectories. Although interhemispheric

connectivity alterations have been investigated in ASD patients, subjects used in previous

studies, including the ones using ABIDE datasets, are primarily high functioning

adults/adolescents with ASD (e.g., mean = 22.4 years old in Anderson et al., 2011 and

17.39 years old for ABIDE-I). However, despite the neurodevelopmental nature of the

disorder (Hull et al., 2017; Kozhemiako et al., 2018), no study to date has specifically

examined homotopic interhemispheric rsFC changes in ASD children.

To shed new light on the utility of using homotopic interhemispheric rsFC

alterations as biomarkers for ASD in general and as early biomarkers to determine ASD

in childhood in particular, the present study has therefore conducted a voxel-based

functional connectivity analysis to investigate whether alterations in homotopic

interhemispheric rsFC are already present in ASD children and whether the changes are

associated with ASD symptom severity. Furthermore, as the primary structure connecting

the left and right brain hemispheres, corpus callosum (CC) volumes have been considered

as an index of interhemispheric connectivity (Vidal et al., 2006; Wegiel et al., 2018) and

volumetric CC alterations have also been associated with ASD (Anderson et al., 2011;

Frazier & Hardan, 2009). The present study thus additionally examined whether there

were parallel CC volume changes in ASD children and, if so, associations with symptom

severity.

Methods

Participants

Resting-state fMRI and T1-weighted anatomical MRI images of 476 children under the

age of 10 years (mean = 8.57, SD = 0.947; range: 5.3 - 10) were obtained from the

ABIDE-I and -II datasets (http://fcon_1000.projects.nitrc.org/indi/abide/ for more

information). The threshold of 10 years old was selected based on definition of age range

for life phases from the American Academy of Pediatrics (Hagan, Shaw, & Duncan, 2008)

and the World Health Organization

(https://www.who.int/southeastasia/health-topics/adolescent-health). Consistent with Li

et al. (2019), subjects were excluded in cases where preprocessing was not successful and

head motion was excessive. Subjects were excluded in cases where: (a) fMRI images

could not be successfully preprocessed, as determined by visual inspection of the

normalization success; (b) mean framewise displacement (FD) was > 0.5 mm; (c) the

percentage of FD > 0.5 mm was over 25%. To further control for head motion, we also

matched head motion (mean FD) in the 2 groups (ASD group: mean = 0.20, SD = 0.096;

control group: mean = 0.18, SD = 0.096; t(335) = 1.62, p = 0.105). To further control for

head motion, we further matched the mean framewise displacement (FD) between the 2

groups (details see supplementary information). These steps led to 152 subjects (50 from

ABIDE-I) in the ASD group and 185 subjects (62 from ABIDE-I) in the TD control

group being included in the final interhemispheric functional connectivity analysis, with

the majority of subjects in the two groups (102 ASD vs. 123 control) being from the

ABIDE-II dataset. Subject demographic information is reported in Supplementary Table

S1 and clinical information of subjects with available scores on symptom severity as

measured by ADOS were reported in Supplementary Table S2 (N = 76). For transparency

and replicability, a full list of subject IDs was also reported in Supplementary Table S3

and Table S4 for ABIDE-I and -II respectively. The data from the ABIDE datasets was

fully anonymized. All contributions in this project were based on studies approved by

local Institutional Review Boards. Written informed consent was obtained from all

subjects or carers.

Data Analyses

Interhemispheric rsFC Analyses

Interhemispheric rsFC analyses were similar to Li et al. (2019). Resting-state fMRI

images were preprocessed using the Data Processing Assistant for Resting-State fMRI

(DPARSF 2.3) software (Yan & Zang, 2010; http://www.restfmri.net) for each site

separately following a standardized procedure (Yan et al., 2013). After discarding the

first 10 images, pre-processing included slice timing correction, realignment, segment

and spatial normalization. For spatial normalization, a pediatric template was created

using the Template-O-Matic Toolbox based on the age and sex of children used in the

present study (Wilke, Holland, Altaye, & Gaser, 2008). Head-motion parameters (using

the Friston 24-parameter model; Friston, Williams, Howard, Frackowiak, & Turner, 1996;

Satterthwaite et al., 2013), global signal, white matter and cerebrospinal fluid were

regressed out to control for potential confounding artifacts. To further exclude residual

effects of head motion, we included the mean FD as a nuisance covariate in the group

analysis (Power et al., 2014) and matched the mean FD between the 2 groups as

mentioned above. Following resampling with a 3 mm $\times$ 3 mm $\times$ 3 mm resolution, the

pre-processed images were spatially smoothed with a Gaussian kernel (8-mm full-width

at half-maximum) and filtered using a band pass filter (0.01-0.08 Hz). The homotopic

interhemispheric functional connectivity map was then generated based on a

voxel-mirrored homotopic connectivity (VMHC) analysis (Zuo et al., 2010) implemented

in the DPARSF toolbox (Yan & Zang, 2010). As a voxel-based functional connectivity

analysis approach, the VMHC reflects rsFC between each voxel in one hemisphere and

its mirrored counterpart in the opposite hemisphere (Stark et al., 2008; Zuo et al., 2010).

The homotopic interhemispheric functional connectivity maps were then converted into

Z-maps using Fisher's z transformation for subsequent group level analyses.

For group level analyses, a two-sample t-test was used to test group functional

connectivity differences between the ASD and TD children. To control for potential

confounding effects, gender, age, total brain volume (grey matter + white matter) and

head motion (mean FD) were included as nuisance covariates in the group analysis. A

threshold of p < 0.05 false discovery rate (FDR) corrected for multiple comparisons at the

voxel level was used in the whole-brain analysis and only clusters larger than 20 voxels

were reported. Given that ADOS scores were only available for 76 ASD children in the

final sample, we only conducted an exploratory correlation analysis between the

homotopic interhemispheric connectivity strength and symptom severity using Pearson

correlation coefficients. The homotopic interhemispheric connectivity strength was

extracted from a 6-mm sphere centered on the peak voxel for regions showing altered

homotopic interhemispheric connectivity in the ASD relative to the control group. ADOS

consisted of social, communication, and stereotyped behavior subscales and total ADOS

scores and scores for each subscale were provided for these 76 ASD children in the

ABIDE datasets.

Corpus Callosum Volume Analyses

Volume analyses of the CC were conducted on T1 data using the VBM8 toolbox

(http://dbm.neuro.unijena.de/vbm8/; Ashburner & Friston, 2000) and kept consistent with

Li et al. (2019). Volume analyses of the CC were conducted on T1 data using the VBM8

toolbox (http://dbm.neuro.unijena.de/vbm8/; Ashburner and Friston, 2000) and kept

consistent with Li et al. (2019). T1 images were first normalized to the Montreal

Neurological Institute (MNI) template and then segmented into GM, WM and

cerebrospinal fluid. Data quality check was performed via the modules of "display one

slice for all images" and "check sample homogeneity using covariance". 10 subjects were

further excluded due to their overall covariance homogeneity being below 2 standard

deviations from the mean. The remaining 145 ASD subjects and 182 controls were

entered into the final analysis. Segmented WM images from these subjects were then

smoothed with a Gaussian kernel of 8-mm full-width at half-maximum.

Group differences in CC volume were tested using a two-sample t-test on smoothed

WM images. Gender, age and the total brain volume (TBV) were included as nuisance

covariates to control for potential confounding effects. The structural CC mask was

derived from the Talairach Daemon database atlas (Lancaster, 1997). Within the CC ROI,

a threshold of $p < 0.05$ FDR corrected at peak-level was set for multiple comparisons and

only clusters larger than 10 voxels reported.

Results

Interhemispheric Functional Connectivity

A two-sample t-test was used to examine homotopic interhemispheric functional

connectivity differences between the ASD and TD children. Results revealed that ASD

compared to TD children exhibited decreased homotopic interhemispheric functional

connectivity in regions of the default mode network (DMN), including the PCC, the

dorsal medial prefrontal cortex (mPFC), ventral mPFC, precuneus and the angular gyrus;

the dorsal anterior cingulate cortex (dACC) of the salience network; the precentral gyrus

and inferior parietal lobule (IPL) of the mirror neuron system; the visual processing

network, including the lingual, fusiform, inferior occipital gyri, inferior and middle

temporal gyri; and thalamus (extended from PCC) ($p_{FDR} < 0.05$; Figure 1 and Table 1).

Categorization of regions into large-scale brain networks was based on previous reviews

and studies on the default mode network (Raichle, 2015; Raichle et al., 2001; Seghier,

2013), the salience network (Menon & Uddin, 2010), the mirror neuron system (Iacoboni

& Dapretto, 2006; Rizzolatti & Craighero, 2004), and the visual processing network

(Bogousslavsky, Miklossy, Deruaz, Assal, & Regli, 2017; Eickenberg, Gramfort,

Varoquaux, & Thirion, 2017; Machielsen, Rombouts, Barkhof, Scheltens, & Witter,

2000). There were no regions showing significantly stronger interhemispheric functional

connectivity in the ASD relative to TD groups using the same threshold ($p_{FDR} < 0.05$).

These findings were consistent with the patterns found in our previous study (Li et

al., 2019). Given that subjects from the ABIDE-I dataset used in the present study were a

small part (~ 12.96%) of the subject sample used in our previous study (Li et al., 2019),

we aimed at further controlling the possibility that subject overlap may have contributed

to the consistent findings between the two studies. To this end, we further conducted an

identical analysis using only ASD and TD children from the ABIDE-II dataset and found

a highly similar but less robust pattern (See Figure S1A and B), which may be due to the

smaller sample size when focusing only on the ABIDE-II dataset. Additionally, we also

confirmed that removing data from children in the ABIDE-I dataset did not influence the

pattern of results we had previously reported using data predominantly from adolescents

and adults (Li et al., 2019) (see Figure S1C and D).

253   ****Insert Figure 1 here****

254   ****Insert Table 1 here****

Correlations between Interhemispheric Functional Connectivity and Symptom

Severity

The exploratory correlation analysis between homotopic interhemispheric

connectivity strength and symptom severity was first conducted for the total ADOS

scores. Results revealed a significant positive correlation between the homotopic

interhemispheric functional connectivity strength of the PCC (Pearson r = 0.301, p =

0.008; Figure 2A), the vmPFC (Pearson r = 0.238, p = 0.039; Figure 2B), the inferior

(Pearson r = 0.277, p = 0.015; Figure 2C) and middle temporal gyri (Pearson r = 0.229, p

= 0.047; Figure 2D) and total ASD symptom load (total ADOS scores). Further

explorations using sub-scale scores revealed significant positive correlations between the

homotopic connectivity strengths of the PCC and social sub-scale scores (Pearson r =

0.227, p = 0.049; Figure 2E), and of the vmPFC (Pearson r = 0.227, p = 0.049; Figure 2F)

and the inferior temporal gyrus (Pearson r = 0.239, p = 0.037; Figure 2G) for the

communication sub-scale scores. The strength of the inferior temporal gyrus connectivity

was also positively associated with stereotyped behavior sub-scale scores (Pearson r =

0.332, p = 0.003; Figure 2H). However, none of these correlations can survive

family-wise error correction for multiple comparisons ($p < 0.05/21 = 0.002$, where 21 is

the total number of regions showing significant altered homotopic connectivity in ASD as

presented in Table 1).

****Insert Figure 2 here****

Corpus Callosum Volume

The ROI-based analysis of CC volume did not reveal any significant volume changes in

the ASD compared to the TD group at a threshold of $p_{FDR} < 0.05$.

Discussion

The present study investigated homotopic interhemispheric rsFC alterations in ASD

children under 10 years of age and their association with symptom severity. Given that

the CC is the key fiber tract connecting the two hemispheres, changes in CC volumes

were also examined. Results showed decreased homotopic interhemispheric functional

connectivity in regions of the DMN, the salience network, the mirror neuron system, the

visual processing network and thalamus in the ASD relative to the TD group, with

interhemispheric functional connectivity of regions in the DMN and the visual processing

network being correlated with symptom severity. No significant CC volume changes

were found in ASD children relative to TD controls. Altered homotopic functional

connectivity may therefore be a potential biomarker for ASD in both children and adults.

Previous studies investigating homotopic interhemispheric rsFC changes in ASD

mainly used subjects ranging from children to adults (Anderson et al., 2011; Di Martino

et al., 2014; King et al., 2019; Li et al., 2019). For example, subjects from the ABIDE-I

dataset used in most of these studies have an average age of 17.39 years with only 11.92%

under the age of 10. However, only focusing on ASD children aged under 10, the present

study using combined ABIDE-I and -II datasets found similar widespread decreased

homotopic interhemispheric rsFC patterns as reported in these previous predominantly

adult/adolescent-based studies. More specifically, the present study replicated the

majority of brain networks exhibiting decreased homotopic interhemispheric connectivity

in ASD patients reported by Li et al. (2019). These included the PCC, the dorsal and

ventral mPFC, precuneus and the angular gyrus of the DMN, the dACC of the salience

network, the precentral gyrus and IPL of the mirror neuron system, the lingual, fusiform

and inferior occipital gyri of the visual processing network and thalamus. These networks

are associated with a variety of cognitive and social/affective domains including

self-referential/consciousness processing and theory of mind (the DMN; Davey, Pujol, &

Harrison, 2016; Raichle, 2015; Saxe, Moran, Scholz, & Gabrieli, 2006), cognitive control

and emotion regulation (dACC; MacDonald, Cohen, Stenger, & Carter, 2000; Stevens,

Hurley, & Taber, 2011), action/intentions understanding and imitation (the mirror neuron

system; Cattaneo & Rizzolatti, 2009; Oberman & Ramachandran, 2007) and the

processing and integration of afferent sensory information (thalamus; Halassa & Kastner,

2017; Hwang, Bertolero, Liu, & D'Esposito, 2017). All of these systems can be

dysfunctional in ASD (Burrows, Laird, & Uddin, 2016; Cheng, Rolls, Gu, Zhang, & Feng,

2015; Dichter, Felder, & Bodfish, 2009; Hamilton, 2013; Linke, Keehn, Pueschel,

Fishman, & Müller, 2018; Padmanabhan, Lynch, Schaer, & Menon, 2017). The present

study thus extends previous ones by providing the first evidence that, at least for children

included in the ABIDE datasets, alterations in homotopic interhemispheric rsFC in

large-scale brain networks have already occurred in individuals aged from 5 to 10 years

of age.

Furthermore, an exploratory analysis between the homotopic interhemispheric

functional connectivity and symptom severity revealed significant positive correlations

between interhemispheric functional connectivity of the PCC and the vmPFC in the

DMN and the inferior and middle temporal gyri in the visual processing network and

total ADOS scores. This may indicate that stronger activity synchronizing between the

left and right parts of these regions, as reflected by stronger interhemispheric functional

connectivity, corresponds to greater overall symptom load in ASD. Similar correlation

patterns were also observed between the homotopic interhemispheric connectivity of the

inferior temporal gyrus and the vmPFC and communication subscale scores and between

the homotopic interhemispheric connectivity of the inferior temporal gyrus and the

stereotyped behavior subscale scores. These findings may indicate a closer clinical

relevance of the homotopic interhemispheric connectivity of these regions specifically in

ASD children. However, these positive correlations, particularly the ones with regions

within the DMN, contrast to some extent with our previous study involving primarily

adults/adolescents from the ABIDE-I dataset where we observed a negative association

between homotopic interhemispheric PCC functional connectivity and the ADOS

communication sub-scale (Li et al., 2019). A study on the developmental trajectories of

homotopic functional connectivity strength in ASD children has reported that some DMN

regions show an increase in connectivity during the first decade of life, followed

subsequently by a maintained decrease during adolescence and adulthood (Kozhemiako

et al., 2018). It is possible that initially a greater reduction in the homotopic functional

connectivity of the PCC and vmPFC may offer some compensatory benefit and reduced

symptoms, but that subsequently in a more fully developed brain in

adolescence/adulthood there is benefit in increasing the homotopic functional

connectivity strength. This might explain the positive correlation with symptom severity

during childhood but negative correlation during adolescence/adulthood. Interestingly,

the study on homotopic developmental trajectories also reported a positive association

with ADOS scores as a function of developmental curvature coefficient (Kozhemiako et

al., 2018). However, some caution is required when interpreting this finding since ADOS

scores were only available for a relatively small number of ASD children in our current

study.

Reduced homotopic functional connectivity has also been consistently reported in

another developmental psychiatric disorder, schizophrenia (Guo et al., 2013; Li, Xu,

Zhang, Hoptman, & Zuo, 2015) suggesting that impaired interhemispheric connectivity

may be a hallmark of these disorders and contribute to some of the cognitive, social,

motor and self-processing dysfunctions common to both. Decreased homotopic

functional connectivity has been observed in early onset schizophrenia patients (mean

age 14.5 years – Li et al., 2015) as well as adults (Guo et al., 2013), although specific

regions associated with symptom severity differ from those associated with symptom

severity in ASD.

We did not find significant differences between the ASD and TD children in CC

volume. Previous findings on CC volume changes in ASD patients have been

inconsistent, with some previous studies reporting reductions in both ASD adults and

children (Haar, Berman, Behrmann, & Dinstein, 2016; Hardan et al., 2009; Keary et al.,

2019; Li et al., 2019) and others in contrast observing no significant alterations (Herbert

et al., 2014; Lefebvre, Beggiato, Bourgeron, & Toro, 2015; Rice et al., 2005). The

inconsistency between these findings may well be as a result of many of the previous

studies being statistical underpowered (Lefebvre et al., 2015) and CC volume changes

derived from simple VBM-derived measures may also be too coarse an index to

characterize subtle changes in ASD (Lefebvre et al., 2015; Li et al., 2019).

There are some limitations in the present study. Firstly, the ages of the ASD and TD

children included in the present study ranged from 5 to 10 years old and therefore it is

possible that younger children might not show similar reduced homotopic functional

connectivity. This needs to be taken into consideration when making general inferences

based on findings concerning age-independency of homotopic connectivity changes.

Secondly, the ABIDE dataset only includes high-functioning ASD patients, therefore the

current findings related to ASD children may not necessarily apply to low-functioning

ASD populations. Finally, ADOS measurements were only available for a relatively

small number of ASD children included in the study.

In conclusion, the present study using the ABIDE dataset has revealed decreased

homotopic interhemispheric functional connectivity in diffuse brain networks including

the DMN, the salience network, the mirror neuron system, the visual processing network,

and thalamus in the ASD relative to the TD children between the ages of 5 and 10. Homo

topic functional connectivity changes between regions in the DMN and the visual

processing network were associated with the symptom severity. Thus, reduced homotopic

interhemispheric functional organization appears to be present in both high functioning

ASD children and adults/adolescents and may provide a useful general biomarker across

both early and late developmental stages of the disorder.

References

Anderson, J. S., Druzgal, T. J., Froehlich, A., DuBray, M. B., Lange, N., Alexander, A.

L., … Lainhart, J. E. (2011). Decreased interhemispheric functional connectivity in

autism. *Cerebral Cortex, 21,* 1134-1146. https://doi.org/10.1093/cercor/bhq190

Ashburner, J., & Friston, K. J. (2000). Voxel-based morphometry-the methods.

Neuroimage, 11, 805-821. https://doi.org/10.1006/nimg.2000.0582

Bogousslavsky, J., Miklossy, J., Deruaz, J. P., Assal, G., & Regli, F. (1987). Lingual and

fusiform gyri in visual processing: a clinico-pathologic study of superior altitudinal

hemianopia. Journal of Neurology, *Neurosurgery & Psychiatry, 50,* 607-614.

http://dx.doi.org/10.1136/jnnp.50.5.607

Burrows, C. A., Laird, A. R., & Uddin, L. Q. (2016). Functional connectivity of brain

regions for self- and other-evaluation in children, adolescents and adults with autism.

Developmental Science, 19, 564-580. https://doi.org/10.1111/desc.12400

Cattaneo, L., & Rizzolatti, G. (2009). The mirror neuron system. *Archives of Neurology,*

66, 557-560. https://doi.org/10.1146/annurev.neuro.27.070203.144230

Charman, T. (2003). Why is joint attention a pivotal skill in autism? *Philosophical*

transactions of the Royal Society of London. Series B, Biological sciences, 358,

315-324. https://doi.org/10.1098/rstb.2002.1199

Chen, C. M., Yang, P., Wu, M. T., Chuang, T. C., & Huang, T. Y. (2019). Deriving and

validating biomarkers associated with autism spectrum disorders from a large-scale

resting-state database. *Scientific Reports, 9,* 9043.

https://doi.org/10.1038/s41598-019-45465-9

Cheng, W., Rolls, E. T., Gu, H., Zhang, J. & Feng, J. (2015). Autism: reduced

connectivity between cortical areas involved in face expression, theory of mind and

the sense of self. *Brain, 138,* 1382-1393. https://doi.org/10.1093/brain/awv051

Chevallier, C., Grèzes, J., Molesworth, C., Berthoz, S., & Happé, F. (2012). Brief report:

Selective social anhedonia in high functioning autism. *Journal of Autism and*

Developmental Disorders, 42, 1504-1509.

https://doi.org/10.1007/s10803-011-1364-0

Courchesne, E., & Pierce, K. (2005). Why the frontal cortex in autism might be talking

only to itself: local over-connectivity but long-distance disconnection. *Current*

Opinion in Neurobiology, 15, 225-230. https://doi.org/10.1016/j.conb.2005.03.001

Dapretto, M., Davies, M. S., Pfeifer, J. H., Scott, A. A., Sigman, M., Bookheimer, S. Y.,

& Iacoboni, M. (2006). Understanding emotions in others: mirror neuron

dysfunction in children with autism spectrum disorders. *Nature Neuroscience, 9,*

28-30. https://doi.org/10.1038/nn1611

Davey, C. G., Pujol, J., & Harrison, B. J. (2016). Mapping the self in the brain's default

mode network. *Neuroimage, 132,* 390-397.

https://doi.org/10.1016/j.neuroimage.2016.02.022

Dichter, G. S., Felder, J. N., & Bodfish, J. W. (2009). Autism is characterized by dorsal

anterior cingulate hyperactivation during social target detection. *Social Cognitive*

and Affective Neuroscience, 4, 215-226. https://doi.org/10.1093/scan/nsp017

Eickenberg, M., Gramfort, A., Varoquaux, G., & Thirion, B. (2017). Seeing it all:

Convolutional network layers map the function of the human visual system.

NeuroImage, 152, 184-194. https://doi.org/10.1016/j.neuroimage.2016.10.001

Martino, A. D., Yan, C. G., Li, Q., Denio, E., Castellanos, F. X., Alaerts, K., …

Milham, M. P. (2014). The autism brain imaging data exchange: towards large-scale

evaluation of the intrinsic brain architecture in autism. *Molecular Psychiatry, 19,*

659-667. https://doi.org/10.1038/mp.2013.78

Ecker, C., Bookheimer, S. Y., & Murphy, D. G. M. (2015). Neuroimaging in autism

spectrum disorder: brain structure and function across the lifespan. *Lancet*

Neurology, 14, 1121-1134. https://doi.org/10.1016/S1474-4422(15)00050-2

Fedor, J., Lynn, A., Foran, W., DiCicco-Bloom, J., Luna, B., & O'Hearn, K. (2018).

Patterns of fixation during face recognition: Differences in autism across age.

Autism, 22, 866-880. https://doi.org/10.1177/1362361317714989

Fishman, I., Keown, C. L., Lincoln, A. J., Pineda, J. A., & Müller, R. A. (2014). Atypical

cross talk between mentalizing and mirror neuron networks in autism spectrum

disorder. *JAMA Psychiatry, 71,* 751-760. doi:10.1001/jamapsychiatry.2014.83 %J

JAMA Psychiatry

Frazier, T. W., & Hardan, A. Y. (2009). A meta-analysis of the corpus callosum in autism.

Biological Psychiatry, 66, 935-941. https://doi.org/10.1016/j.biopsych.2009.07.022

Friston, K. J., Williams, S., Howard, R., Frackowiak, R. S., & Turner, R. (1996).

Movement-related effects in fMRI time-series. *Magnetic Resonance in Medicine, 35,*

346-355. https://doi.org/10.1002/mrm.1910350312

Funakoshi, Y., Harada, M., Otsuka, H., Mori, K., Ito, H., & Iwanaga, T. (2016). Default

mode network abnormalities in children with autism spectrum disorder detected by

resting-state functional magnetic resonance imaging. *The Journal of Medical*

Investigation, 63, 204-208. https://doi.org/10.2152/jmi.63.204

Geschwind, D. H., & Levitt, P. (2007). Autism spectrum disorders: developmental

disconnection syndromes. *Current Opinion in Neurobiology, 17,* 103-111.

https://doi.org/10.1016/j.conb.2007.01.009

Guo, S., Kendrick, K. M., Zhang, J., Broome, M., Yu, R., Liu, Z., & Feng, J. (2013).

Brain-wide functional inter-hemispheric disconnection is a potential biomarker for

schizophrenia and distinguishes it from depression. *Neuroimage: Clinical, 2,*

818-826. https://doi.org/10.1016/j.nicl.2013.06.008

Haar, S., Berman, S., Behrmann, M., & Dinstein, I. (2016). Anatomical abnormalities in

autism. *Cerebral Cortex, 26,* 1440-1452. https://doi.org/10.1093/cercor/bhu242

Hagan, J. F. Jr, Shaw, J. S., & Duncan, P., eds. (2008). *Bright Futures: Guidelines for the*

Health Supervision of Infants, Children, and Adolescents (3rd ed.). Elk Grove

Village, IL: American Academy of Pediatrics.

Halassa, M. M., & Kastner, S. (2017). Thalamic functions in distributed cognitive control.

Nature Neuroscience, 20, 1669-1679. https://doi.org/10.1038/s41593-017-0020-1

Hamilton, A. F. D. C. (2013). Reflecting on the mirror neuron system in autism: a

systematic review of current theories. *Developmental Cognitive Neuroscience, 3,*

91-105. https://doi.org/10.1016/j.dcn.2012.09.008

Happé, F., & Frith, U. (2006). The weak coherence account: detail-focused cognitive

style in autism spectrum disorders. *Journal of Autism and Developmental Disorders,*

36, 5-25. https://doi.org/10.1007/s10803-005-0039-0

Hardan, A. Y., Pabalan, M., Gupta, N., Bansal, R., Melhem, N. M., Fedorov, S., …

Minshew, N.J. (2009). Corpus callosum volume in children with autism. *Psychiatry*

Research, 174, 57-61. https://doi.org/10.1016/j.rasd.2012.09.007

Herbert, M. R., Ziegler, D. A., Makris, N., Filipek, P. A., Kemper, T. L., Normandin, J.

J., … Caviness Jr, V. S. (2004). Localization of white matter volume increase in

autism and developmental language disorder. *Annals of Neurology, 55,* 530-540.

https://doi.org/10.1002/ana.20032

Herringshaw, A. J., Kumar, S. L., Rody, K. N., & Kana, R. K. (2018). Neural Correlates

of Social Perception in Children with Autism: Local versus Global Preferences.

Neuroscience, 395, 49-59. https://doi.org/10.1016/j.neuroscience.2018.10.044

Hull, J. V., Jacokes, Z. J., Torgerson, C. M., Irimia, A., & van Horn, J. D. (2017).

Resting-state functional connectivity in autism spectrum disorders: a review.

Frontiers in Psychiatry, 7, 205. https://doi.org/10.3389/fpsyt.2016.00205

Hwang, K., Bertolero, M. A., Liu, W. B., & D'Esposito, M. (2017). The human thalamus

is an integrative hub for functional brain networks. *Journal of Neuroscience, 37,*

5594-5607. https://doi.org/10.1523/JNEUROSCI.0067-17.2017

Iacoboni, M., & Dapretto, M. (2006). The mirror neuron system and the consequences of

its dysfunction. *Nature Reviews Neuroscience, 7,* 942-951.

https://doi.org/10.1038/nrn2024

Kanner, L. (1943). Autistic disturbances of affective contact. *Nervous Child, 2,* 217-250.

Keary, C. J., Minshew, N. J., Bansal, R., Goradia, D., Fedorov, S., Keshavan, M. S., &

Hardan, A. Y. (2009). Corpus callosum volume and neurocognition in autism.

Journal of Autism and Developmental Disorders, 39, 834-841.

https://doi.org/10.1007/s10803-009-0689-4

King, J. B., Prigge, M. B. D., King, C. K., Morgan, J., Weathersby, F., Chancellor Fox,

J., … Anderson, J.S. (2019). Generalizability and reproducibility of functional

connectivity in autism. *Molecular Autism,10,* 27.

https://doi.org/10.1186/s13229-019-0273-5

Kozhemiako, N., Vakorin, V., Nunes, A. S., Iarocci, G., Ribary, U., & Doesburg, S. M.

(2018). Extreme male developmental trajectories of homotopic brain connectivity.

Human Brain Mapping, 40, 987-1000. https://doi.org/10.1002/hbm.24427

Lancaster, J. L. (1997). The Talairach Daemon, a database server for Talairach atlas

labels. *Neuroimage, 5,* S633.

Lau, W., Leung, M. K., & Lau, B. (2019). Resting-state abnormalities in Autism

Spectrum Disorders: A meta-analysis. *Scientific Reports, 9,* 3892.

https://doi.org/10.1038/s41598-019-40427-7

Lee, Y., Park, B. Y., James, O., Kim, S. G., & Park, H. (2017). Autism Spectrum

Disorder Related Functional Connectivity Changes in the Language Network in

Children, Adolescents and Adults. *Frontiers in Human Neuroscience, 11,* 418.

https://doi.org/10.3389/fnhum.2017.00418

Lefebvre, A., Beggiato, A., Bourgeron, T., & Toro, R. (2015). Neuroanatomical diversity

of corpus callosum and brain volume in autism: meta-analysis, analysis of the

autism brain imaging data exchange project, and simulation. *Biological Psychiatry,*

78, 126-134. https://doi.org/10.1016/j.biopsych.2015.02.010

Li, H. J., Xu, Y., Zhang, K. R., Hoptman, M. J., & Zuo, X. N. (2015). Homotopic

connectivity in drug□naïve, first□episode, early□onset schizophrenia. *Journal of*

Child Psychology and Psychiatry, 56, 432-443. https://doi.org/10.1111/jcpp.12307

Li, Q., Becker, B., Jiang, X., Zhao, Z., Zhang, Q., Yao, S., & Kendrick, K. M. (2019).

Decreased interhemispheric functional connectivity rather than corpus callosum

volume as a potential biomarker for autism spectrum disorder. *Cortex, 119,* 258-266.

https://doi.org/10.1016/j.cortex.2019.05.003

Linke, A. C., Keehn, R. J. J., Pueschel, E. B., Fishman, I., & Müller, R. A. (2018).

Children with ASD show links between aberrant sound processing, social symptoms,

and atypical auditory interhemispheric and thalamocortical functional connectivity.

Developmental Cognitive Neuroscience, 29, 117-126.

https://doi.org/10.1016/j.dcn.2017.01.007

Machielsen, W. C., Rombouts, S. A., Barkhof, F., Scheltens, P., & Witter, M. P. (2000).

FMRI of visual encoding: reproducibility of activation. *Human Brain Mapping, 9,*

156-164.

https://doi.org/10.1002/(SICI)1097-0193(200003)9:3<156::AID-HBM4>3.0.CO;2-

Q

MacDonald, A. W., Cohen, J. D., Stenger, V. A., & Carter, C. S. (2000). Dissociating the

role of the dorsolateral prefrontal and anterior cingulate cortex in cognitive control.

Science, 288, 1835-1838. https://doi.org/10.1126/science.288.5472.1835

Madipakkam, A. R., Rothkirch, M., Dziobek, I., & Sterzer, P. (2017). Unconscious

avoidance of eye contact in autism spectrum disorder. *Scientific Reports, 7,* 13378.

https://doi.org/10.1038/s41598-017-13945-5

Menon, V., & Uddin, L. Q. (2010). Saliency, switching, attention and control: a network

model of insula function. *Brain Structure and Function, 214,* 655-667.

https://doi.org/10.1007/s00429-010-0262-0

Monk, C. S., Peltier, S. J., Wiggins, J. L., Weng, S. J., Carrasco, M., Risi, S., & Lord, C.

(2009). Abnormalities of intrinsic functional connectivity in autism spectrum

disorders. *Neuroimage, 47,* 764-772.

https://doi.org/10.1016/j.neuroimage.2009.04.069

Morrison, K. E., Pinkham, A. E., Penn, D. L., Kelsven, S., Ludwig, K., & Sasson, N. J.

(2017). Distinct profiles of social skill in adults with autism spectrum disorder and

schizophrenia. *Autism Research, 10,* 878-887. https://doi.org/10.1002/aur.1734

Oberman, L. M., & Ramachandran, V. S. (2007). The simulating social mind: the role of

the mirror neuron system and simulation in the social and communicative deficits of

autism spectrum disorders. *Psychological Bulletin, 133,* 310.

https://doi.org/10.1037/0033-2909.133.2.310

Oberwelland, E., Schilbach, L., Barisic, I., Krall, S. C., Vogeley, K., Fink, G. R., …

Schulte-Rüther, M. (2017). Young adolescents with autism show abnormal joint

attention network: A gaze contingent fMRI study. *Neuroimage: Clinical, 14,*

112-121. https://doi.org/10.1016/j.nicl.2017.01.006

Padmanabhan, A., Lynch, C. J., Schaer, M., & Menon, V. (2017). The default mode

network in autism. *Biological Psychiatry, 2,* 476-486.

https://doi.org/10.1016/j.bpsc.2017.04.004

Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E.

(2014). Methods to detect, characterize, and remove motion artifact in resting state

fMRI. *Neuroimage, 84,* 320-341. https://doi.org/10.1016/j.neuroimage.2013.08.048

Raichle, M. E. (2015). The brain's default mode network. Annual Review of

Neuroscience, 38, 433-447. https://doi.org/10.1146/annurev-neuro-071013-014030

Raichle, M. E., MacLeod, A. M., Snyder, A. Z., Powers, W. J., Gusnard, D. A., &

Shulman, G. L. (2001). A default mode of brain function. *Proceedings of the*

National Academy of Sciences, 98, 676-682. doi:10.1073/pnas.98.2.676

Rice, S. A., Bigler, E. D., Cleavinger, H. B., Tate, D.F., Sayer, J., McMahon, W.,

Ozonoff, S., Lu, J., & Lainhart, J. E. (2005). Macrocephaly, corpus callosum

morphology, and autism. *Journal of Child Neurology, 20,* 34-41.

https://doi.org/10.1177/08830738050200010601

Rizzolatti, G., & Craighero, L. (2004). The mirror-neuron system. *Annual Review of*

Neuroscience, 27, 169-192.

https://doi.org/10.1146/annurev.neuro.27.070203.144230

Satterthwaite, T. D., Elliott, M. A., Gerraty, R. T., Ruparel, K., Loughead, J., Calkins, M.

E., … Wolf, D.H. (2013). An improved framework for confound regression and

filtering for control of motion artifact in the preprocessing of resting-state functional

connectivity data. *Neuroimage, 64,* 240-256.

https://doi.org/10.1016/j.neuroimage.2012.08.052

Seghier, M. L. (2013). The angular gyrus: multiple functions and multiple subdivisions.

The Neuroscientist, 19, 43-61. https://doi.org/10.1177/1073858412440596

Stark, D. E., Marguiles, D. S., Shehzad, Z. E, Reiss, P., Clare Kelly, A. M. C., … Milham,

M.P. (2008). Regional variation in interhemispheric coordination of intrinsic

hemodynamic fluctuations. *Journal of Neuroscience, 28,* 13754-13764.

https://doi.org/10.1523/JNEUROSCI.4544-08.2008

Stevens, F. L., Hurley, R. A., & Taber, K. H. (2011). Anterior cingulate cortex: unique

role in cognition and emotion. *The Journal of Neuropsychiatry and Clinical*

Neurosciences, 23, 121-125. doi:10.1001/jamapsychiatry.2013.104

Supekar, K., Kochalka, J., Schaer, M., Wakeman, H., Qin, S., Padmanabhan, A., &

Menon, V. (2018). Deficits in mesolimbic reward pathway underlie social

interaction impairments in children with autism. *Brain, 141,* 2795-2805.

https://doi.org/10.1093/brain/awy191

Uddin, L. Q., Supekar, K., Lynch, C. J., Khouzam, A., Phillips, J., Feinstein, C., …

Menon, V. (2013). Salience network-based classification and prediction of symptom

severity in children with autism. *JAMA Psychiatry, 70,* 869-879.

doi:10.1001/jamapsychiatry.2013.104

Vidal, C. N., Nicolson, R., DeVito, T. J., Hayashi, K. M., Geaga, J. A., Drost, D. J., …

Thompson, P.M. (2006). Mapping corpus callosum deficits in autism: an index of

aberrant cortical connectivity. *Biological Psychiatry, 60,* 218-225.

https://doi.org/10.1016/j.biopsych.2005.11.011

von dem Hagen, E. A., Stoyanova, R. S., Rowe, J. B., Baron-Cohen, S., & Calder, A. J.

(2014). Direct gaze elicits atypical activation of the theory-of-mind network in

autism spectrum conditions. *Cerebral Cortex, 24,* 1485-1492.

https://doi.org/10.1093/cercor/bht003

Wang, A. T., Dapretto, M., Hariri, A. R., Sigman, M., & Bookheimer, S. Y. (2004).

Neural correlates of facial affect processing in children and adolescents with autism

spectrum disorder. *Journal of the American Academy of Child and Adolescent*

Psychiatry, 43, 481-490. https://doi.org/10.1097/00004583-200404000-00015

Washington, S. D., Gordon, E. M., Brar, J., Warburton, S., Sawyer, A. T., Wolfe, A., …

VanMeter, J.W. (2014). Dysmaturation of the default mode network in autism.

Human Brain Mapping, 35, 1284-1296. https://doi.org/10.1002/hbm.22252

Wegiel, J., Kaczmarski, W., Flory, M., Martinez-Cerdeno, V., Wisniewski, T., Nowicki,

K., Kuchna, I., & Wegiel, J. (2018). Deficit of corpus callosum axons, reduced axon

diameter and decreased area are markers of abnormal development of

interhemispheric connections in autistic subjects. *Acta Neuropathologica*

Communications, 6, 143. https://doi.org/10.1186/s40478-018-0645-7

Wilke, M., Holland, S. K., Altaye, M., & Gaser, C. (2008). Template-O-Matic: a toolbox

for creating customized pediatric templates. *Neuroimage, 41,* 903-913.

https://doi.org/10.1016/j.neuroimage.2008.02.056

Yan, C., & Zang, Y. (2010). DPARSF: a MATLAB toolbox for "pipeline" data analysis

of resting-state fMRI. *Frontiers in Systems Neuroscience, 4,* 13.

https://doi.org/10.3389/fnsys.2010.00013

Yan, C. G., Craddock, R. C., Zuo, X. N., Zang, Y. F., & Milham, M. P. (2013).

Standardizing the intrinsic brain: towards robust measurement of inter-individual

variation in 1000 functional connectomes. *Neuroimage, 80,* 246-262.

https://doi.org/10.1016/j.neuroimage.2013.04.081

Zuo, X. N., Kelly, C., Di Martino, A., Mennes, M., Margulies, D. S., Bangaru, S., …

Milham, M. P. (2010). Growing together and growing apart: regional and sex

differences in the lifespan developmental trajectories of functional homotopy.

Journal of Neuroscience, 30, 15034-15043.

https://doi.org/10.1523/JNEUROSCI.2612-10.2010

Table 1. Regions showing decreased interhemispheric connectivity in the autism relative to the control groups.

Brain Region	BA	No. Voxels	Peak t-value	x	y	z
Posterior Cingulate Cortex	7/30/31	855	5.18	3	-48	12
Precuneus			4.43	3	-51	33
Lingual Gyrus			4.08	27	-54	-6
Fusiform Gyrus			4.01	30	-69	-3
Thalamus			3.43	12	-30	3
Middle Temporal Gyrus	21	70	4.78	66	-18	-9
Supramarginal/Angular Gyrus	40	37	4.24	60	-51	33
Inferior Parietal Lobule			3.16	60	-51	45
Inferior Temporal Gyrus	20/37	34	4.24	66	-54	-12
Middle Temporal Gyrus			3.59	54	-48	-12
Dorsal Medial Prefrontal Cortex	9/10	58	4.17	3	63	9
Dorsal Anterior Cingulate Cortex			3.32	3	45	24
Rostral Anterior Cingulate Cortex	24/32	48	4.15	6	36	3
Ventral Medial Prefrontal Cortex			2.88	6	36	-9
Middle Occipital Gyrus	18/19	110	4.08	30	-99	3
Inferior Occipital Gyrus			3.24	36	-90	-6
Caudate		63	3.74	9	3	15
Middle Frontal Gyrus	6/8	89	3.69	27	18	51
Superior Parietal Lobule	7/40	72	3.58	30	-60	48
Postcentral Gyrus	5/7	45	3.58	24	-42	63
Precentral Gyrus	6	23	3.54	36	-15	42

635 All with a p < 0.05 FDR corrected threshold and voxels larger than 20. BA: Brodmann Area. Brain coordinates
636 are reported in the Montreal Neurological Institute (MNI) space.
637

Figure Legends

Figure 1. Regions showing significantly ($p_{FDR} < 0.05$) decreased homotopic interhemispheric

functional connectivity in the ASD compared to the typically developing groups. PCC =

posterior cingulate cortex; dmPFC = dorsomedial prefrontal cortex; vmPFC = ventromedial

prefrontal cortex; PreCG = precentral gyrus; IOG = inferior occipital gyrus; LING = lingual

gyrus; FFG = fusiform gyrus; PostCG = postcentral gyrus; IPL = inferior parietal lobule. Color

bar indicates t values of the statistical map.

Figure 2. Correlations between the homotopic interhemispheric functional connectivity of the

posterior cingulate cortex (PCC) (A), the ventromedial prefrontal cortex (vmPFC) (B), the inferior

(ITG) (C) and middle temporal gyri (MTG) (D) and ADOS total scores. Correlations between the

homotopic connectivity strengths of the PCC and social sub-scale scores (E), and of the vmPFC (F)

and the ITG (G) for the communication sub-scale scores. The strength of the ITG connectivity was

also positively associated with stereotyped behavior sub-scale scores (H).