Mechanical worrying drives cell migration in crowded environments

Erik S. Welf,[1,2,*] Meghan K. Driscoll,[1,2,*] Etai Sapoznik,[2] Vasanth S. Murali,[1,2] Andrew Weems,[1,2] Juan Manuel Garcia-Arcos,[3] Minna Roh-Johnson,[4] Kevin M. Dean,[2] Matthieu Piel,[3] Reto Fiolka,[2,§] Gaudenz Danuser[1,2,§]

[1]Department of Bioinformatics, University of Texas Southwestern Medical Center, Dallas, TX
[2]Department of Cell Biology, University of Texas Southwestern Medical Center, Dallas, TX
[3]Institut Curie, PSL Research University, CNRS, Paris, France.
[4]Department of Biochemistry, School of Medicine, University of Utah, Salt Lake City, UT
*contributed equally
§Corresponding author, Gaudenz.Danuser@UTSouthwestern.edu, Reto.Fiolka@UTSouthwestern.edu

Migratory cells navigate through crowded 3D microenvironments *in vivo.* Amoeboid cells, such as immune cells and some cancer cells, are thought to do so by deforming their bodies to squeeze through tight spaces.[1] Yet large populations of nearly spherical amoeboid cells migrate[2-4] in microenvironments too dense[5,6] to move through without extensive shape deformations. How they do so is unknown. We used high-resolution light-sheet microscopy to visualize metastatic melanoma cells in dense environments, finding that cells maintain a round morphology as they migrate and create a path through which to move via bleb-driven mechanical degradation and subsequent macropinocytosis of extracellular matrix components. Proteolytic degradation of the extracellular matrix via matrix metalloproteinases is not required. Membrane blebs are short-lived relative to the timescale of migration, and thus persistence in their polarization is critical for productive ablation of the extracellular matrix. Interactions between small but long-lived cortical adhesions and collagen at the cell front induce PI-3 Kinase signaling that drive bleb enlargement via branched actin polymerization. Large blebs in turn abrade collagen, creating a feedback between extracellular matrix structure, cell morphology, and cell polarization that results in both path generation and persistent cell movement.

Introduction

Cell migration is critical to processes ranging from embryogenesis and wound healing to cancer metastasis.[1] When spatially confined, animal and non-animal cells alike exhibit bleb-based motility, a type of amoeboid migration characterized by weak adhesion and minimal proteolytic destruction of the surrounding matrix.[7,8] Amoeboid cells can migrate through tight spaces by deforming their body and nucleus, even to the point of nucleus rupture.[6,9,10] Studies of invasive cancer cells, particularly of melanoma, have reported that rounded amoeboid cells are enriched at the tumor edge,[11] and can move through Matrigel and tumor xenografts.[2,12,13] Indeed, the pro-migratory effect of intracellular contractility, which is associated with rounding and surface blebbing, is well known.[11,13-17] The pronounced rounded morphology of these cells seems to be at odds with a migration mode that relies on shape deformation. Round, amoeboid cells have been observed to move in tunnels.[4] Although under some conditions amoeboid cells secrete matrix metalloproteinases (MMPs),[18] amoeboid migration is not usually considered a proteolytic migration mode, and so it has been generally assumed that these tunnels were pre-formed by 'helper' cells moving in a mesenchymal manner, a migration mode in which the extracellular matrix (ECM) is degraded via MMPs.[4] Despite the notion that amoeboid cells cannot remodel their environment to generate their own paths and thus would be forced to immobility in a very dense microenvironment, blebbing cells in soft extracellular matrices have been observed to physically

manipulate fibers, pushing them out of their way[19] and tugging on them with adhesions,[20] suggesting the
possibility of an amoeboid migration mode enabled by matrix remodeling.

To investigate the mechanism of this potentially new path generating migration strategy, we created
dense, yet pliable *in vivo* mimetics by encapsulating cells in fibrous Collagen 1 gels with bulk stiffnesses
on the order of 1 kPa (Advanced biomatrix.com).[21,22] We imaged cells in 3D with high-resolution light-
sheet microscopy at near isotropic ~350 nm resolution, using specially designed sample chambers that do
not interfere with the mechanical properties of the mimetics (Fig. 1a, Movie 1).[23,24] Adsorption of collagen
to hard surfaces increases collagen stiffness near the surface,[25-28] which shifts and eventually diminishes
the rounded, blebbing morphotype. To observe migration of these cells without mechanical interference,
our chambers enabled cell imaging at greater than 1 mm away from any stiff surfaces.

To determine whether our *in vivo* mimetics prohibited deformation-based migration through pores, we
fluorescently labeled collagen and then measured collagen pore size.[24,29,30] Although blebs were small
enough to fit inside the pores in the collagen network, the nucleus and cell body of amoeboid melanoma
cells were too large to fit through the existing pores (Fig. 1a,b, Movie 2), thus rendering a deformation-
based mode of migration unlikely. Long-term time-lapse imaging of melanoma cells confirmed that
amoeboid cells were nevertheless able to move through the mimetics while maintaining their largely
spherical shape (Fig. 1c, Extended Data Fig. 1a,b).

As a model cell for bleb-based migration through soft crowded environments, we chose metastatic
melanoma cells. *In vivo*, melanoma metastasizes to soft environment such as the brain.[31] Consistent with
these clinical observations, we noted that melanoma in the soft environment of the zebrafish hindbrain
not only have an amoeboid morphology but bleb extensively (Extended Data Fig. 1c). We next tested
whether the amoeboid morphology was associated with melanoma metastatic potential within our *in vivo*
mimetic. We imaged populations of primary melanoma cells that were harvested from patients and then
passaged in a mouse xenotransplantation system.[32,33] Comparing three mechanically distinct collagen
microenvironments, including the mimetic, we found that samples with higher metastatic efficiency were
enriched in the amoeboid morphology compared to the stretched mesenchymal morphology (Figure 1d,
Extended Data Figure 1d,e). Furthermore, using unbiased cell shape motif detection,[34] we discovered that
a parental melanoma cell line (A375P) had a lower average bleb count than a subpopulation of the cell
line that had been enriched for metastatic potential in mouse xenografts (A375M2) (Fig. 1e,f).[35] Together,
these results establish the significance of our experimental system as a model of metastatic cell migration
in soft, ECM-dense tissues.

Migrating amoeboid cells can carve a path without the need for extracellular proteolytic degradation.
To determine how blebbing melanoma cells migrate through soft, dense collagen, we imaged them 24
hours after seeding. Over this time frame, many cells had created tunnels (Fig. 1g). We observed a similar
tunneling phenomenon with a different melanoma cell line (Extended Data Fig. 2a), as well as with
pediatric Ewing sarcoma cells (Extended Data Fig. 2b). Tunnel creation in dense matrices is usually ascribed
to matrix metalloproteinase (MMP)-dependent mesenchymal migration.[3,36] Thus, our finding of an
amoeboid cell morphology associated with a clearly demarcated, cell-generated path seemed paradoxical
in view of the current paradigms of cancer cell motility. To begin to solve this puzzle, we applied a broad
spectrum MMP inhibitor, GM6001,[37,38] and found no effect on the ability of melanoma cells to tunnel (Fig.
1h). The MMP-independence of the migration mode also held when collagen was not first solubilized
using pepsin, which removes a collagen crosslinking site potentially rendering the collagen easier to
digest.[39] Although tunneling is somewhat less frequent in unpepsinized collagen, the pore size is greater

Figure 1. Cells can generate a path through dense collagen without matrix metalloproteinases. (a) Surface rendering of a 3D light-sheet microscopy image of a melanoma cell in 3D collagen with collagen fibers behind it shown yellow. (b) Pore size analysis of 2.0 mg/ml bovine collagen gels showing the fraction of space occupied by a specific pore size (n = 6 gels). Error bars indicate the standard error of the mean. In comparison, the mean diameter of blebby melanoma cells in these types of gels is 20 ± 1 μm (n = 44 cells). (c) Time-lapse phase-contrast microscopy images showing the movement of a round, blebby melanoma cell in 3D collagen. (d) Fraction of melanoma cells with rounded morphology in cell populations extracted from patient-derived xenografts of high vs low metastatic efficiency under three different microenvironmental conditions. (e) Surface renderings of parental, P, and metastatic, MA2, A375 melanoma cells. Computationally identified blebs are shown randomly colored. (f) Number of blebs per cell in the parental, P (n = 33 cells), and metastatic, MA2 (n = 9 cells), cells. Cells with the median number of blebs are indicated in green and shown in panel e. (g) Maximum intensity projection across 4.8 μm of a light-sheet microscope image of an A375MA2 melanoma cell expressing GFP-F-tractin (red) in 3D collagen processed with a steerable filter to enhance edges (cyan). The white dashed line indicates the location of a tunnel. (h) Percent of cells in tunnels after 18-24 hours in pepsinized or unpepsinized bovine collagen. Samples were either untreated (un.; n = 176 cells in pepsinized collagen, 67 cells in unpepsinized collagen), or treated with DSMO (n = 52 cells in pepsinized collagen, 105 cells in unpepsinized collagen), or 40 mM GM6001 (gm; n = 82 cells in pepsinized collagen, 67 cells in unpepsinized collagen). Error bars indicate 95% confidence intervals calculated using the Normal approximation for the confidence interval of a binomial distribution. (i) Single optical sections of 3D light-sheet images of collagen samples containing MV3 cells treated with either DMSO or 40 μM GM6001 for ~24 hours. The cyan channel shows collagen with fibers enhanced by a steerable line filter and the red channel shows ¾ collagen antibody fluorescence near tunnels formed by cells. (j) Quantification of ¾ collagen antibody intensity inside a tunnel divided by the intensity outside the tunnel in samples treated with either DMSO or 40 μm GM6001. Error bars show 95% confidence intervals (n = 4 tunnels for each condition). All scale bars show 10 μm.

as is the extent of directional collagen realignment by the cells, suggesting a reduced need to tunnel (Extended Data Fig. 3). We confirmed that GM6001 inhibited MMP activity by measuring the abundance of an antibody that recognizes the MMP-cleaved collagen site in tunnels (Fig. 1i,j) and by direct measurement of MMP enzymatic activity on a synthetic substrate (Extended Data Fig. 2c). Thus, tunneling is not mediated by the enzymatic activity of MMPs.

Path generation is mediated by bleb-driven ablation of the extracellular matrix (ECM).
To address alternative path generation mechanisms to protease-activity, we analyzed the interactions of cells with collagen in greater detail. Cells inside tunnels were often highly polarized, with many large blebs at the cell front facing the enclosed end of the tunnel (Fig. 2a, Movies 3,4). Measuring the difference in frequencies of protrusive and retractive motion on and off blebs, we found that blebs were on average protrusive and non-blebs retractive (Fig. 2b), suggesting that blebs are responsible for cell protrusion through collagen. We next measured the colocalization of blebs with collagen, finding that collagen was enriched in regions near blebs, but not directly on blebs (Fig. 2c). This is explained by bleb interdigitation

Figure 2. Persistent agitation and internalization of collagen enables cells to dig tunnels. (a) Surface rendering of a light-sheet microscope image of an MV3 melanoma cell, colored by surface curvature, in 3D collagen. The scale for surface curvature is shown in Figure 1A. (b) Frequency of protrusive motion minus the frequency of retractive motion on and off blebs (aggregated over n = 9 cells). (c) Collagen intensity on and off blebs as a function of distance to the nearest bleb neck (aggregated over n= 23 cells). (d) XZ and XY views of maximum intensity projections of light-sheet images of a melanoma cell expressing GFP-AktPH (red) interacting with collagen fibers enhanced by steerable line filtering (cyan) and projected over 5.6 µm and 2.4 µm, respectively.

The region shown in the XY view is indicated by the dashed box in the XZ view. Orange arrows indicate blebs interdigitating
between collagen fibers. (e) Maximum intensity projection across 3.2 μm of a light-sheet image of a melanoma cell expressing
GFP-AktPH (red) in 3D collagen enhanced by steerable line filtering (cyan). The dashed box indicates the region magnified in panel
F. (f) Overlay of two different time points (green and magenta, separated by 120 seconds) of a maximum intensity projection
over 3.2 μm of a light-sheet microscope image of collagen fibers enhanced by steerable line filtering. Orange arrows indicate the
motion of individual collagen fibers. (g) Collagen motion (red), as measured by 3D optical flow, near a melanoma cell expressing
GFP-AktPH (white), imaged using light-sheet microscopy and shown as a maximum intensity projection over the entire cell. (h)
Collagen motion near the cell surface of blebs associated with either protrusive (blue) or retractive (red) motion (aggregated over
n = 5 cells). (i) Maximum intensity projection across 3.2 μm of a light-sheet image of a cell expressing GFP-AktPH (red) in 3D
collagen (cyan). The orange arrow indicates internalized collagen at the front of the cell. (j) Surface rendering of a light-sheet
microscopy image of a melanoma cell expressing GFP-AktPH (red) in 3D collagen (cyan). A quadrant of the cell and collagen is cut
away to show the collagen internalized at the cell periphery. (k) Maximum intensity projection across 3.2 μm of a light-sheet
microscope image of a cell expressing GFP-AktPH (red) in 3D collagen (cyan). The dashed box indicates the region magnified in
the time-lapse panels to the right and the orange arrow indicates a piece of collagen that is broken off and brought in towards
the center of the cell. (L&M) Maximum intensity projection across 3.2 μm of a light-sheet microscope image of MV3 cells
(unlabeled) in 3D collagen (cyan) treated with 70 kDa FITC-dextran (red), as well as either DMSO (l) or 50 mM EIPA (m). The orange
arrows indicates internalized collagen and dextran in intracellular vesicles. (n) Number of internalized collagen fragments in either
DMSO (n = 20 cells) or EIPA-treated (n = 23 cells) cells (difference between conditions per one-sided t-test, p = 0.04). (o) Number
of internalized collagen fragments per cell vs. the number of internalized dextran dots in EIPA-treated cells. (p) The p value,
calculated for each cell, corresponding to the likelihood that collagen fluorescence intensity is not elevated at the location of
dextran dots. All scale bars show 10 μm.

into pores in the collagen network, resulting in high collagen fiber density at the base of blebs (Fig. 2d).
Then we examined collagen motion, which showed movement of individual collagen fibers at the front of
tunneling cells (Fig. 2e,f). Using a 3D optical flow algorithm designed to capture multi-scale motion both
near and away from cells (Fig. 2g),[40] we compared the collagen speed near blebs with the bleb speed for
both protruding and retracting blebs (Fig. 2h). For protrusive blebs, at low bleb speeds collagen speed
increased linearly with bleb speed, whereas at high bleb speeds collagen motion plateaued, consistent
with bleb interdigitation into collagen pores. For retracting blebs we found that even at high bleb speed,
collagen was pulled in concert with the blebs, meaning that retracting blebs pull collagen towards the cell
surface. Indeed, at the fronts of highly polarized cells, collagen was often enriched into a shell at the cell
periphery alongside extensive internalization of labeled collagen (Fig. 2i,j). Over long periods of time, cells
slowly agitated the collagen shell, breaking off fragments of the collagen and pulling them into the cell
(Fig. 2k).

To determine the mechanism of collagen internalization, we incubated cells with high molecular weight
dextran, finding that it was ingested alongside labeled collagen (Fig. 2l). Internalization of large liquid-
phase molecules is indicative of macropinocytosis, since such molecules are excluded from smaller
endocytic vesicles.[41] Treatment with the sodium hydrogen exchange inhibitor 5-(N-ethyl-N-
isopropyl)amiloride (EIPA), an inhibitor of macropinocytosis,[42] decreased the number of dextran-labeled
vesicles and internalized collagen fragments, with the number of dextran vesicles and collagen fragments
highly correlated across cells (Fig. 2m-o). Testing the hypothesis that collagen localization at detected
dextran vesicles was random, we found that the distribution of p-values was heavily tilted towards small
values, indicating likely collagen enrichment within dextran vesicles (Fig. 2p). Furthermore, we did not
observe intracellular vesicles containing fluorescently-labeled clathrin light chain (CLC) that were
associated with internalized collagen (Extended Data Fig. 4), indicating that clathrin-mediated endocytosis
does not contribute to internalization. Thus, we conclude that macropinocytosis is the dominant form of
collagen internalization in this form of path generation for cell migration. The centrality of
macropinocytosis to this form of amoeboid cancer cell migration, combined with its known role in

enabling nutrient uptake in depleted cancer microenvironments,[43] highlights the importance of
macropinocytosis to metastasis.

Phosphoinositide 3-kinase (PI3K) establishes bleb polarity in feedback with collagen remodeling.
For productive path generation, the slow destruction of dense extracellular matrix at the cell front
critically depends on persistence factors that promote the highly polarized and continuous formation of
large blebs, which abrade and internalize matrix material in a directed fashion. To measure polarization
on the 3D cell surface, we used an approximation of a spherical normal distribution, which has fit
parameters that intuitively correspond to the direction of the peak and the peak's inverse width, here
termed the polarization magnitude (Fig. 3a). Using these statistics, we first confirmed that the distribution
of large blebs was more polarized than the overall bleb distribution (Fig. 3b). Measuring the directional
correlation of large bleb polarization and collagen localization, we next found that large blebs were
systematically biased towards areas of high collagen density (Fig. 3c). Hypothesizing that adhesions might
couple collagen and bleb localization, we found that paxillin-containing adhesion complexes indeed
formed (Fig. 3d, Extended Data Fig. 5) in the direction of the high collagen density at the closed end of the
tunnel (Fig. 3e). A canonical cell polarity factor that is organized by nascent adhesions via focal adhesion
kinase (FAK) is PI3K.[44-46] Similar to the distribution of adhesions, we observed a striking polarization of
PI3K near the closed end of the tunnel (Fig. 3f, Movie 5). This was surprising given that fibroblasts
exhibiting pseudopodial motility in 3D microenvironments do not seem to require polarized PI3K
signaling.[47] Moreover, PI3K signaling was more directionally aligned with large blebs than with blebs of all
sizes (Fig. 3g), suggesting that PI3K signaling is involved specifically in the polarization of large blebs.
Despite their small size, adhesions in the cortical area persisted for several minutes (Fig. 3h), in contrast
to the ~1 min lifetime of similarly-sized nascent adhesions formed in cells on a coverslip (Fig. 3i,j).[48] The
localization and persistence of cortical adhesions at the front of tunneling cells may enable the
recruitment of PI3K to the cell front.

To test this hypothesis, we acutely inhibited FAK signaling using a small molecule inhibitor of FAK-kinase
activity. FAK inhibition resulted in a decrease in bleb volume, even though bleb polarization and number
were unaffected (Fig. 3k). PI3K polarization and mean cell surface motion were also decreased (Fig. 3k,
Extended Data Fig. 6). Measuring the full-width half-maxima of the FAK inhibition response times, we
found that PI3K polarization fell first, followed by bleb volume and then cell surface motion. This led us to
conclude that PI3K polarity is upstream of large bleb formation at the cell front. Indeed, stratification of
blebs by volume revealed that large blebs in particular were enriched for high PI3K signaling and also
associated with increased collagen motion (Extended Data Fig. 7).

To determine if the relationship between PI3K and bleb size was causative, we used photoactivation to
increase PI3K signaling locally in blebbing cells, resulting in a striking increase in proximal but not distal
bleb size (Fig. 3l, Extended Data Fig. 8, Movie 6). We also pharmacologically inhibited PI3K signaling by
acute addition of a low dose of an inhibitor specific for PI3Kα. In the region of former high PI3K activity,
PI3K biosensor intensity and bleb size rapidly decreased, even though de novo bleb formation was not
inhibited (Extended Data Fig. 9, Movie 7). Aggregating over multiple cells, we found that both PI3K
polarization and bleb size were decreased by PI3K inhibition, whereas the number of blebs and bleb
polarization were not affected (Fig. 3m). Altogether, these results indicate that PI3K is responsible for
generating large blebs but does not govern the frequency or location of bleb initiation.

Figure 3. Large blebs are polarized to the cell front via a feedback between collagen remodeling, adhesion formation, and PI3K localization. (a) Simulated species concentrations illustrating example polarization magnitudes and directional correlations. (b) Distributions of polarization magnitudes for all blebs (blue) and for the largest decile of blebs by volume (green). (n = 34 cells) (c) Directional correlation of collagen polarization near the cell surface with polarization of the largest decile of blebs. (d) A 3D surface rendering of a melanoma cell in 3D collagen with the cell surface shown transparent in gray and paxillin-marked adhesions in red.

(e) The directional correlation of collagen polarization with adhesion polarization. (f) A surface rendering of a light-sheet
microscope image of a melanoma cell in collagen, colored by the localization of GFP-AktPH. (g) The directional correlation of bleb
polarization with PI3K polarization for all blebs and for the largest decile of blebs by volume. (h) Time-lapse images of GFP-paxillin
adhesions localized to the cortical region in a worrying cell. (i) Time-lapse images of GFP-paxillin adhesions within a dynamic
protrusion, indicated by the dashed blue box. (j) Adhesion lifetimes for adhesions near the cell cortex (red) and within dynamic
protrusions (blue) in three different cells for each condition. Each data point represents the persistence of a single adhesion. (k)
Representative temporal response of blebs and PI3K signaling in an MV3 cell treated with FAK inhibitor 14. From top to bottom,
shown are mean bleb volume, bleb polarization magnitude, number of blebs, PI3K polarization magnitude, and mean surface
motion magnitude. Dashed lines indicate the approximate full-width half-maximum decay times of measures that are reduced
by FAK inhibition. (l) Change in bleb size due to photoactivation, calculated as mean maximum area per bleb during activation
divided by mean maximum area per bleb in the same region before activation (p=0.0053, two sided t-test, n=6 regions from 6
cells expressing mCherry-CRY2-iSH2 along with CIBN-CAAX and n=8 regions in 6 cells not expressing cry2-mRuby2-PRL3) (m) Effect
of PI3Kα inhibitor IV compared to DMSO control on bleb and PI3K properties in MV3 cells. PI3K polarization (p = 0.001) and bleb
volume (p = 9x10^{-8}) show statistically significant differences across treatments, whereas bleb polarization (p = 0.4) and number
of blebs (p = 0.12) do not. Statistical testing was performed with a one-sided t-test.

PI3K enlarges blebs by enhancing branched actin polymerization.

237 The fact that PI3K activity was responsive to FAK manipulation left open the possibility of a mechanical
238 feedback from large blebs to PI3K, which would also explain the observed persistence in polarization of
239 blebs and signaling. To test if the presence of large blebs affected PI3K activity, we reduced cell blebbing
240 via addition of wheat germ agglutinin (WGA), which binds to sialic acid and N-acetylglucosaminyl residues
241

on the extracellular surface of the cell membrane, thereby increasing membrane stiffness.[49,50] We found
that decreasing bleb abundance via WGA decreased the polarity and strength of PI3K signaling (Fig. 4a,
Extended Data Fig. 10), supporting the notion of a sustained feedback between blebbing and local PI3K
activity.

We therefore sought to uncover the mechanism that couples PI3K activity and bleb size. Bleb growth is
thought to be driven by pressure-based cytoplasmic flows[51] with actin filament formation only at very late
growth stages to reform the actin cortex.[52] In contrast, PI3K is usually associated with actin-driven
protrusion.[53] Thus, the involvement of PI3K in bleb expansion seemed paradoxical. We noted, however,
that PI3K and actin were directionally correlated (Fig. 4b,c) and that regions of the cell with higher PI3K
activity protruded faster (Fig. 4d). Indeed, in agreement with previous findings,[51] we found that blebs that
ultimately reached a larger size did so by growing faster (Fig. 4e). Inhibition of PI3K activity dramatically
decreased the growth rate and final bleb size (Fig. 4e), confirming that fast bleb growth and large size are
due to PI3K signaling.

Given the known association of PI3K with actin-based migration, we wondered if actin polymerization
could be playing a role in bleb growth despite previous reports otherwise. We found that the F-tractin
construct, which localizes to filamentous actin,[54] was absent during bleb expansion and localized only to
the bleb cortex, as previously reported.[52] However, expressing low levels of HALO-tagged actin showed
that actin is present during bleb expansion (Extended Data Fig. 11a). Calculating the ratio of actin to F-
tractin revealed that actin was enriched in blebs relative to filamentous actin (Extended Data Fig. 11b),
raising the possibility that nascent actin filaments formed in blebs, but are not recognized by the F-tractin
probe. To further explore this possibility, we used a TIRF microscope with a high numerical aperture (NA)
objective to image actin in the growing blebs of HeLa cells gene-edited to express a copy of the beta actin
gene fused to GFP. We found that actin filaments are indeed formed in the bleb during expansion (Fig. 4f,
Movie 8). Furthermore, stratification of blebs by their final size revealed that large blebs contained a
significantly higher concentration of actin than small blebs, in particular later in their life (Fig. 4g).

Figure 4. Bleb size is locally controlled by branched actin polymerization. (a) The number of blebs and the PI3K polarization in individual cells as a function of Wheat Germ Agglutinin (WGA) dose. In each category, the value of the median cell is colored red. (b) Surface renderings of 3D light-sheet microscopy images showing GFP-AktPH and HALO-actin intensity on the surface of an MV3 cell. (c) Directional correlation of PI3K and actin polarization in individual MV3 cells (n= 11 cells). (d) Local cell surface motion as a function of local PI3K intensity at the cell surface (n= 9 cells). (e) Protrusion speed of individual blebs as a function of final bleb size in either unperturbed or PI3K inhibited cells. (f) Time-lapse sequence of TIRF microscopy images of a single bleb in a HeLa cell expressing GFP-actin. The bleb edge is indicated by the dark blue line. (g) Mean HALO-actin intensity within large and small blebs in a light-sheet image of MV3 cells as a function of time after bleb initiation. Dashed lines represent 95% confidence intervals (n= 12 blebs from 3 different cells). (h) Effect of Arp2/3 inhibition via CK666 on PI3K and bleb properties, compared to DMSO control. PI3K polarization (p = 0.004) and bleb volume (p = 0.001) show statistically significant differences across samples, whereas the number of blebs do not (p = 0.48). Statistical testing was performed with a one-sided t-test. (i) Spinning disk confocal microscope images showing a single optical slice of mCherry-PA-Rac1 in a mouse embryonic fibroblast before, during and after photoactivation of PI3K in the area indicated by a red circle. (j) Change in bleb size due to photoactivation, calculated as the mean of the maximum bleb area during activation divided by the mean of the maximum bleb area in the same region before activation (p=1.6 x10[-5], two sided t-test; n=6 regions expressing mCherry-PA-Rac1, n=11 regions in cells not expressing mCherry-PA-Rac1).

The putative link between PI3K signaling and actin polymerization may rest on the Rac1 – WAVE – Arp2/3 pathway,[44,55] which promotes lamellipodia expansion. To test the involvement of this pathway also in bleb expansion, we first blocked Arp2/3 activity by the small molecule inhibitor CK666,[56,57] which decreased bleb volume but did not alter bleb number (Fig. 4h). This is consistent with previous findings that both genetic and acute inhibition of the Arp2/3 complex reduces bleb size.[58] Coupled with previous findings that CK666 treatment does not decrease intracellular pressure[59] and thus is not expected to globally decrease the driving force for bleb expansion, our results suggest that active actin polymerization contributes to bleb growth, especially in large blebs. To add to this conclusion, we employed a photoactivatable Rac1 construct,[60] which allowed us to increase Rac1 activity locally and acutely, avoiding the more global and pleiotropic effects Arp2/3 inhibition may exert. Local photoactivation resulted in a dramatic, reproducible increase in local bleb size that was immediately reversible upon light cessation

294 (Fig. 4i,j, Movie 9). Our results thus show that the mechanism by which PI3K localization promotes bleb
295 growth at the front of tunneling cells is via Rac1-mediated actin polymerization.

Discussion.

Our data uncover a mode of cell migration that is effective in dense, soft environments. We refer to this
mode as worrying, which means wearing down or tearing repeatedly, like a dog worrying a bone, a
meaning that predates the more figurative use as a term related to anxiety.[61] In the context of cell
migration worrying denotes that the core element of the mechanism is the sustained agitation and tearing
of the extracellular matrix at the cell front by persistently polarized and dynamic cell surface blebs. The
persistence is mounted by a mechanochemical feedback between actin-enforced large bleb formation,
matrix ablation, adhesion signaling, and PI3K/Rac1 triggered activation of actin filament assembly inside
the bleb. The discovery of this self-reinforcing machinery depended on the development of 3D imaging
assays to capture cell dynamic behaviors without mechanical interference from the microscope optics,
computer vision to extract the relations between cell blebbing and signaling, and optogenetic approaches
to acutely interfere with the feedback loop.

Our custom-built technology enabled the study of migration in dense, yet soft environments. Not only are
such tissues common *in vivo*, with melanoma in particular known to prefer soft environments,[62] but many
tissues throughout the body are likely mechanical composites with pockets of mechanically soft
microenvironments.[63,64] It is well established that mesenchymal migration, which is especially common in
stiff environments, is facilitated by ECM remodeling leading to tunnel generation and directional fiber
alignment.[65] From the recent discovery that amoeboid immune cells build specialized actin structures to
push fibers out of the way,[66] combined with the results of this study, we now conclude that all major
modes of migration are remodeling modes, in which the environment is at least transiently reorganized.
Hence, alongside the core processes of cell migration established by Abercrombie[67] decades ago, i.e. of
protrusion, adhesion and contraction, we add environmental remodeling as the fourth process. Moving
forward, the use of advanced technologies to dissect migration in ever more complex environments will
be critical to enhancing our understanding of the mechanisms of environmental remodeling and its
integration with the other core processes.

Our results additionally underscore how much remains to be understood about blebs. Blebs are known to
play a role in diverse processes beyond cancer cell migration, including apoptosis, cytokinesis, cell
spreading, and virus entry.[68] Although the role of blebs as spatial compartments has been previously
understood, we show here as an additional function for blebs the pushing and pulling of the environment
in order to remodel it. In so doing, we found that bleb expansion is locally controlled by PI3K signaling and
mediated by branched actin polymerization. This raises the possibility that the branched actin machinery
governing bleb expansion in cell migration, may also be critical to other processes involving blebs, for
example organelle segregation into blebs in apoptosis, demanding the need for future studies to
understand how cellular control of blebbing enables critical cell functions.

Acknowledgments

We thank Allan Zhang for manually tracking the cells imaged via phase-contrast microscopy, Ugur
Eskiocak, Elena Piskounova, Arin Aurora, and Sean Morrison for providing primary melanoma cells. We
thank Stefan Wieser and Verena Ruprecht for use of their custom high NA TIRF microscope and Philippe
Roudot for advice using the optical flow algorithm. We also thank Alba Diz-Muñoz and Jörg Renkawitz for
kindly commenting on the manuscript. Cell and collagen rendering was performed with UCSF ChimeraX,
developed by the Resource for Biocomputing, Visualization, and Informatics at the University of California,

San Francisco, with support from National Institutes of Health R01-GM129325 and the Office of Cyber
Infrastructure and Computational Biology, National Institute of Allergy and Infectious Diseases. This work
was supported by the following grants: K25CA204526 to ESW, K99GM123221 to MKD, INSERM ITMO
Cancer Plan Cancer 2015-2020 to MP, Fondation ARC no. DOC20190508743 PhD fellowship to WMG-A,
NIH R33CA235254 and R35GM133522 to RF, Cancer Prevention Research Institute of Texas grant
RR160057 to RF, and R35-GM136428 to GD.

Competing Interests
The authors declare no competing interests.
345

346 **Data Availability**
347 The data that support the findings of this study are available from the corresponding authors upon
348 reasonable request.
349

Code Availability
Much of the code used in this study was associated with previously published methods papers and is
available at https://github.com/DanuserLab. All remaining code will be made available at that repository
upon publication.

References

1   Yamada, K. M. & Sixt, M. Mechanisms of 3D cell migration. *Nat Rev Mol Cell Biol* **20**, 738-752,
(2019).
2   Sahai, E. & Marshall, C. J. Differing modes of tumour cell invasion have distinct requirements for
Rho/ROCK signalling and extracellular proteolysis. *Nature Cell Biology* **5**, 711-719, (2003).
3   Wolf, K., Mazo, I., Leung, H., Engelke, K., von Andrian, U. H., Deryugina, E. I., Strongin, A. Y.,
Brocker, E. B. & Friedl, P. Compensation mechanism in tumor cell migration: Mesenchymal-
amoeboid transition after blocking of pericellular proteolysis. *Journal of Cell Biology* **160**, 267-277,
(2003).
4   Paul, C. D., Mistriotis, P. & Konstantopoulos, K. Cancer cell motility: lessons from migration in
confined spaces. *Nat Rev Cancer* **17**, 131-140, (2017).
5   Zaman, M. H., Trapani, L. M., Sieminski, A. L., Mackellar, D., Gong, H., Kamm, R. D., Wells, A.,
Lauffenburger, D. A. & Matsudaira, P. Migration of tumor cells in 3D matrices is governed by
matrix stiffness along with cell-matrix adhesion and proteolysis. *Proc Natl Acad Sci U S A* **103**,
10889-10894, (2006).
6   Wolf, K., te Lindert, M., Krause, M., Alexander, S., te Riet, J., Willis, A. L., Hoffman, R. M., Figdor,
C. G., Weiss, S. J. & Friedl, P. Physical limits of cell migration: Control by ECM space and nuclear
deformation and tuning by proteolysis and traction force. *Journal of Cell Biology* **201**, 1069-1084,
(2013).
7   Ruprecht, V., Wieser, S., Callan-Jones, A., Smutny, M., Morita, H., Sako, K., Barone, V., Ritsch-
Marte, M., Sixt, M., Voituriez, R. & Heisenberg, C. P. Cortical contractility triggers a stochastic
switch to fast amoeboid cell motility. *Cell* **160**, 673-685, (2015).
8   Brunet, T., Albert, M., Roman, W., Coyle, M. C., Spitzer, D. C. & King, N. A flagellate-to-amoeboid
switch in the closest living relatives of animals. *Elife* **10**, (2021).
9   Lammermann, T., Bader, B. L., Monkley, S. J., Worbs, T., Wedlich-Soldner, R., Hirsch, K., Keller, M.,
Forster, R., Critchley, D. R., Fassler, R. & Sixt, M. Rapid leukocyte migration by integrin-
independent flowing and squeezing. *Nature* **453**, 51-55, (2008).

10   Raab, M., Gentili, M., de Belly, H., Thiam, H. R., Vargas, P., Jimenez, A. J., Lautenschlaeger, F.,
382   Voituriez, R., Lennon-Dumenil, A. M., Manel, N. & Piel, M. ESCRT III repairs nuclear envelope
383   ruptures during cell migration to limit DNA damage and cell death. *Science* **352**, 359-362, (2016).

384   11   Georgouli, M., Herraiz, C., Crosas-Molist, E., Fanshawe, B., Maiques, O., Perdrix, A., Pandya, P.,
385   Rodriguez-Hernandez, I., Ilieva, K. M., Cantelli, G., Karagiannis, P., Mele, S., Lam, H., Josephs, D.
386   H., Matias-Guiu, X., Marti, R. M., Nestle, F. O., Orgaz, J. L., Malanchi, I., Fruhwirth, G. O.,
387   Karagiannis, S. N. & Sanz-Moreno, V. Regional Activation of Myosin II in Cancer Cells Drives Tumor
388   Progression via a Secretory Cross-Talk with the Immune Microenvironment. *Cell* **176**, 757-774
389   e723, (2019).

390   12   Pinner, S. & Sahai, E. PDK1 regulates cancer cell motility by antagonising inhibition of ROCK1 by
391   RhoE. *Nat Cell Biol* **10**, 127-137, (2008).

392   13   Sanz-Moreno, V., Gaggioli, C., Yeo, M., Albrengues, J., Wallberg, F., Viros, A., Hooper, S., Mitter,
393   R., Feral, C. C., Cook, M., Larkin, J., Marais, R., Meneguzzi, G., Sahai, E. & Marshall, C. J. ROCK and
394   JAK1 signaling cooperate to control actomyosin contractility in tumor cells and stroma. *Cancer*
395   *Cell* **20**, 229-245, (2011).

396   14   Gadea, G., de Toledo, M., Anguille, C. & Roux, P. Loss of p53 promotes RhoA-ROCK-dependent cell
397   migration and invasion in 3D matrices. *Journal of Cell Biology* **178**, 23-30, (2007).

398   15   Madsen, C. D., Hooper, S., Tozluoglu, M., Bruckbauer, A., Fletcher, G., Erler, J. T., Bates, P. A.,
399   Thompson, B. & Sahai, E. STRIPAK components determine mode of cancer cell migration and
400   metastasis. *Nat Cell Biol* **17**, 68-80, (2015).

401   16   Sadok, A., McCarthy, A., Caldwell, J., Collins, I., Garrett, M. D., Yeo, M., Hooper, S., Sahai, E.,
402   Kuemper, S., Mardakheh, F. K. & Marshall, C. J. Rho kinase inhibitors block melanoma cell
403   migration and inhibit metastasis. *Cancer Res* **75**, 2272-2284, (2015).

404   17   Rodriguez-Hernandez, I., Maiques, O., Kohlhammer, L., Cantelli, G., Perdrix-Rosell, A., Monger, J.,
405   Fanshawe, B., Bridgeman, V. L., Karagiannis, S. N., Penin, R. M., Marcolval, J., Marti, R. M., Matias-
406   Guiu, X., Fruhwirth, G. O., Orgaz, J. L., Malanchi, I. & Sanz-Moreno, V. WNT11-FZD7-DAAM1
407   signalling supports tumour initiating abilities and melanoma amoeboid invasion. *Nat Commun* **11**,
408   5315, (2020).

409   18   Orgaz, J. L., Pandya, P., Dalmeida, R., Karagiannis, P., Sanchez-Laorden, B., Viros, A., Albrengues,
410   J., Nestle, F. O., Ridley, A. J., Gaggioli, C., Marais, R., Karagiannis, S. N. & Sanz-Moreno, V. Diverse
411   matrix metalloproteinase functions regulate cancer amoeboid migration. *Nat Commun* **5**, 4255,
412   (2014).

413   19   Wyckoff, J. B., Pinner, S. E., Gschmeissner, S., Condeelis, J. S. & Sahai, E. ROCK- and myosin-
414   dependent matrix deformation enables protease-independent tumor-cell invasion in vivo. *Curr*
415   *Biol* **16**, 1515-1523, (2006).

416   20   Guzman, A., Avard, R. C., Devanny, A. J., Kweon, O. S. & Kaufman, L. J. Delineating the role of
417   membrane blebs in a hybrid mode of cancer cell invasion in three-dimensional environments. *J*
418   *Cell Sci* **133**, (2020).

419   21   Bordeleau, F., Mason, B. N., Lollis, E. M., Mazzola, M., Zanotelli, M. R., Somasegar, S., Califano, J.
420   P., Montague, C., LaValley, D. J., Huynh, J., Mencia-Trinchant, N., Negron Abril, Y. L., Hassane, D.
421   C., Bonassar, L. J., Butcher, J. T., Weiss, R. S. & Reinhart-King, C. A. Matrix stiffening promotes a
422   tumor vasculature phenotype. *Proc Natl Acad Sci U S A* **114**, 492-497, (2017).

423   22   Burla, F., Dussi, S., Martinez-Torres, C., Tauber, J., van der Gucht, J. & Koenderink, G. H.
424   Connectivity and plasticity determine collagen network fracture. *Proc Natl Acad Sci U S A* **117**,
425   8326-8334, (2020).

426   23   Dean, K. M., Roudot, P., Welf, E. S., Danuser, G. & Fiolka, R. Deconvolution-free Subcellular
427   Imaging with Axially Swept Light Sheet Microscopy. *Biophys. J.* **108**, 2807-2815, (2015).

24 Welf, E. S., Driscoll, M. K., Dean, K. M., Schafer, C., Chu, J., Davidson, M. W., Lin, M. Z., Danuser,
G. & Fiolka, R. Quantitative Multiscale Cell Imaging in Controlled 3D Microenvironments. *Dev. Cell*
36, 462-475, (2016).
25 Ma, X., Schickel, M. E., Stevenson, M. D., Sarang-Sieminski, A. L., Gooch, K. J., Ghadiali, S. N. &
Hart, R. T. Fibers in the extracellular matrix enable long-range stress transmission between cells.
Biophys J **104**, 1410-1418, (2013).
26 Provenzano, P. P., Inman, D. R., Eliceiri, K. W. & Keely, P. J. Matrix density-induced
mechanoregulation of breast cell phenotype, signaling and gene expression through a FAK-ERK
linkage. *Oncogene* **28**, 4326-4343, (2009).
27 Rao, S. S., Bentil, S., DeJesus, J., Larison, J., Hissong, A., Dupaix, R., Sarkar, A. & Winter, J. O.
Inherent interfacial mechanical gradients in 3D hydrogels influence tumor cell behaviors. *PLoS*
One **7**, e35852, (2012).
28 Wang, H., Abhilash, A. S., Chen, C. S., Wells, R. G. & Shenoy, V. B. Long-range force transmission
in fibrous matrices enabled by tension-driven alignment of fibers. *Biophys J* **107**, 2592-2603,
(2014).
29 Aguet, F., Jacob, M. & Unser, M. Three-dimensional feature detection using optimal steerable
filters. *IEEE Image. Proc.* **2**, 1158-1161, (2005).
30 Lomakin, A. J., Cattin, C. J., Cuvelier, D., Alraies, Z., Molina, M., Nader, G. P. F., Srivastava, N., Saez,
P. J., Garcia-Arcos, J. M., Zhitnyak, I. Y., Bhargava, A., Driscoll, M. K., Welf, E. S., Fiolka, R., Petrie,
R. J., De Silva, N. S., Gonzalez-Granado, J. M., Manel, N., Lennon-Dumenil, A. M., Muller, D. J. &
Piel, M. The nucleus acts as a ruler tailoring cell responses to spatial constraints. *Science* **370**,
(2020).
31 Achrol, A. S., Rennert, R. C., Anders, C., Soffietti, R., Ahluwalia, M. S., Nayak, L., Peters, S., Arvold,
N. D., Harsh, G. R., Steeg, P. S. & Chang, S. D. Brain metastases. *Nat Rev Dis Primers* **5**, 5, (2019).
32 Quintana, E., Piskounova, E., Shackleton, M., Weinberg, D., Eskiocak, U., Fullen, D. R., Johnson, T.
M. & Morrison, S. J. Human melanoma metastasis in NSG mice correlates with clinical outcome in
patients. *Sci Transl Med* **4**, 159ra149, (2012).
33 Zaritsky, A., Jamieson, A. R., Welf, E. S., Nevarez, A., Cillay, J., Eskiocak, U., Cantarel, B. L. &
Danuser, G. Interpretable deep learning of label-free live cell images uncovers functional
hallmarks of highly-metastatic melanoma. *BioRxiv*, (2020).
34 Driscoll, M. K., Welf, E. S., Jamieson, A. R., Dean, K. M., Isogai, T., Fiolka, R. & Danuser, G. Robust
and automated detection of subcellular morphological motifs in 3D microscopy images. *Nat*
Methods, (2019).
35 Clark, E. A., Golub, T. R., Lander, E. S. & Hynes, R. O. Genomic analysis of metastasis reveals an
essential role for RhoC. *Nature* **406**, 532-535, (2000).
36 Kessenbrock, K., Plaks, V. & Werb, Z. Matrix metalloproteinases: regulators of the tumor
microenvironment. *Cell* **141**, 52-67, (2010).
37 Grobelny, D., Poncz, L. & Galardy, R. E. Inhibition of human skin fibroblast collagenase,
thermolysin, and Pseudomonas aeruginosa elastase by peptide hydroxamic acids. *Biochemistry*
31, 7152-7154, (1992).
38 Winer, A., Adams, S. & Mignatti, P. Matrix Metalloproteinase Inhibitors in Cancer Therapy: Turning
Past Failures Into Future Successes. *Mol Cancer Ther* **17**, 1147-1155, (2018).
39 Sabeh, F., Shimizu-Hirota, R. & Weiss, S. J. Protease-dependent versus -independent cancer cell
invasion programs: three-dimensional amoeboid movement revisited. *Journal of Cell Biology* **185**,
11-19, (2009).
40 Manandhar, S., Bouthemy, P., Welf, E., Danuser, G., Roudot, P. & Kervrann, C. 3D flow field
estimation and assessment for live cell fluorescence microscopy. *Bioinformatics* **36**, 1317-1325,
(2020).
41 Kerr, M. C. & Teasdale, R. D. Defining macropinocytosis. *Traffic* **10**, 364-371, (2009).

42 Masereel, B., Pochet, L. & Laeckmann, D. An overview of inhibitors of Na(+)/H(+) exchanger. *Eur J Med Chem* **38**, 547-554, (2003).

43 Stow, J. L., Hung, Y. & Wall, A. A. Macropinocytosis: Insights from immunology and cancer. *Curr Opin Cell Biol* **65**, 131-140, (2020).

44 Chen, H. C. & Guan, J. L. Association of focal adhesion kinase with its potential substrate phosphatidylinositol 3-kinase. *Proc Natl Acad Sci U S A* **91**, 10148-10152, (1994).

45 Johnson, H. E., King, S. J., Asokan, S. B., Rotty, J. D., Bear, J. E. & Haugh, J. M. F-actin bundles direct the initiation and orientation of lamellipodia through adhesion-based signaling. *Journal of Cell Biology* **208**, 443-455, (2015).

46 Welf, E. S., Ahmed, S., Johnson, H. E., Melvin, A. T. & Haugh, J. M. Migrating fibroblasts reorient directionality by a metastable, PI3K-dependent mechanism. *Journal of Cell Biology* **197**, 105-114, (2012).

47 Petrie, R. J., Gavara, N., Chadwick, R. S. & Yamada, K. M. Nonpolarized signaling reveals two distinct modes of 3D cell migration. *Journal of Cell Biology* **197**, 439-455, (2012).

48 Choi, C. K., Vicente-Manzanares, M., Zareno, J., Whitmore, L. A., Mogilner, A. & Horwitz, A. R. Actin and alpha-actinin orchestrate the assembly and maturation of nascent adhesions in a myosin II motor-independent manner. *Nat Cell Biol* **10**, 1039-1050, (2008).

49 Charras, G. T., Yarrow, J. C., Horton, M. A., Mahadevan, L. & Mitchison, T. J. Non-equilibration of hydrostatic pressure in blebbing cells. *Nature* **435**, 365-369, (2005).

50 Evans, E. & Leung, A. Adhesivity and rigidity of erythrocyte membrane in relation to wheat germ agglutinin binding. *Journal of Cell Biology* **98**, 1201-1208, (1984).

51 Cunningham, C. C. Actin polymerization and intracellular solvent flow in cell surface blebbing. *Journal of Cell Biology* **129**, 1589-1599, (1995).

52 Charras, G. T., Hu, C. K., Coughlin, M. & Mitchison, T. J. Reassembly of contractile actin cortex in cell blebs. *J. Cell Biol.* **175**, 477-490, (2006).

53 Funamoto, S., Meili, R., Lee, S., Parry, L. & Firtel, R. A. Spatial and temporal regulation of 3-phosphoinositides by PI 3-kinase and PTEN mediates chemotaxis. *Cell* **109**, 611-623, (2002).

54 Schell, M. J., Erneux, C. & Irvine, R. F. Inositol 1,4,5-trisphosphate 3-kinase A associates with F-actin and dendritic spines via its N terminus. *J Biol Chem* **276**, 37537-37546, (2001).

55 Welch, H. C., Coadwell, W. J., Stephens, L. R. & Hawkins, P. T. Phosphoinositide 3-kinase-dependent activation of Rac. *FEBS Lett* **546**, 93-97, (2003).

56 Bisi, S., Disanza, A., Malinverno, C., Frittoli, E., Palamidessi, A. & Scita, G. Membrane and actin dynamics interplay at lamellipodia leading edge. *Curr Opin Cell Biol* **25**, 565-573, (2013).

57 Nolen, B. J., Tomasevic, N., Russell, A., Pierce, D. W., Jia, Z., McCormick, C. D., Hartman, J., Sakowicz, R. & Pollard, T. D. Characterization of two classes of small molecule inhibitors of Arp2/3 complex. *Nature* **460**, 1031-1034, (2009).

58 Bergert, M., Chandradoss, S. D., Desai, R. A. & Paluch, E. Cell mechanics control rapid transitions between blebs and lamellipodia during migration. *Proc Natl Acad Sci U S A* **109**, 14434-14439, (2012).

59 Cartagena-Rivera, A. X., Logue, J. S., Waterman, C. M. & Chadwick, R. S. Actomyosin Cortical Mechanical Properties in Nonadherent Cells Determined by Atomic Force Microscopy. *Biophys J* **110**, 2528-2539, (2016).

60 Wu, Y. I., Frey, D., Lungu, O. I., Jaehrig, A., Schlichting, I., Kuhlman, B. & Hahn, K. M. A genetically encoded photoactivatable Rac controls the motility of living cells. *Nature* **461**, 104-U111, (2009).

61 Worry. *Oxford English Dictionary.*

62 Bras, M. M., Radmacher, M., Sousa, S. R. & Granja, P. L. Melanoma in the Eyes of Mechanobiology. *Front Cell Dev Biol* **8**, 54, (2020).

63 Bonnans, C., Chou, J. & Werb, Z. Remodelling the extracellular matrix in development and disease. *Nat. Rev. Mol. Cell Biol.* **15**, 786-801, (2014).

64   Handorf, A. M., Zhou, Y., Halanski, M. A. & Li, W. J. Tissue stiffness dictates development,
homeostasis, and disease progression. *Organogenesis* **11**, 1-15, (2015).
65   Winkler, J., Abisoye-Ogunniyan, A., Metcalf, K. J. & Werb, Z. Concepts of extracellular matrix
remodelling in tumour progression and metastasis. *Nat Commun* **11**, 5120, (2020).
66   Gaertner, F., Reis-Rodrigues, P., Vries, I. d., Hons, M., Aguilera, J., Riedl, M., Leithner, A., Merrin,
J., Zheden, V., Kaufmann, W. A., Hauschild, R. & Sixt, M. Mechanosensitivity of amoeboid cells
crawling in 3D. *BioRxiv*, (2021).
67   Abercrombie, M. Croonian lecture, 1978: the crawling movement of metazoan cells. *Proceedings*
of the Royal Society Series B-Biological Sciences **207**, 129-147, (1980).
68   Charras, G. & Paluch, E. Blebs lead the way: How to migrate without lamellipodia. *Nat Rev Mol*
Cell Biol **9**, 730-736, (2008).

Methods
Cell culture and reagents
MV3 cells were obtained from Peter Friedl (MD Anderson Cancer Center, Houston TX). A375 (ATCC® CRL-
1619) and A375MA2 (ATCC® CRL-3223) cells were acquired from ATCC. SKNMC Ewing sarcoma cells were
obtained from the Whitehurst lab at UT Southwestern. MV3, A375, and SKNMC cells were cultured in
DMEM (Gibco) supplemented with 10% fetal bovine serum (FBS; ThermoFisher) at 37 °C and 5% CO_2.

Populations of primary melanoma cells were created from tumors grown in murine xenograft models as
described previously,[1] and provided as a gift by the Laboratory of Sean Morrison (UT Southwestern
Medical Center, Dallas, TX). Briefly, cells were suspended in Leibovitz's L-15 Medium (ThermoFisher)
containing mg/ml bovine serum albumin, 1% penicillin/streptomycin, 10 mM HEPES and 25% high protein
Matrigel (product 354248; BD Biosciences). Subcutaneous injections of human melanoma cells were
performed in the flank of NOD.CB17-Prkdcscid Il2rgtm1Wjl/SzJ (NSG) mice (Jackson Laboratory). These
experiments were performed according to protocols approved by the animal use committees at the
University of Texas Southwestern Medical Center (protocol 2011-0118). After surgical removal, tumors
were mechanically dissociated and subjected to enzymatic digestion for 20 min with 200 U ml−1
collagenase IV (Worthington), 5 mM CaCl2, and 50 U ml−1 DNase at 37oC. Cells were filtered through a
40 μm cell strainer to break up cell aggregates and washed through the strainer to remove cells from large
tissue pieces. The cells were then cultured in medium containing the Melanocyte Growth Kit (ATCC PCS-
200-042) and Dermal Cell Basal Medium (ATCC PCS-200-030).

Inhibitors
FAK inhibitor 14 was purchased from Tocris (3414). PI3K alpha inhibitor IV was purchased from Santa Cruz
(sc-222170). CK666 was purchased from Millipore Sigma (SML0006). The ¾-collagen antibody was
purchased from Adipogen (AG-25T-011). NEIPA was purchased from Sigma (A3085). WGA was purchased
from VWR (101098-084). 70 kDa Fluorescein isothiocyanate–dextran was purchased from Sigma (46945).
GM6001 was purchased from Sigma (CC1010).

MMP activity assay
We used the MMP activity assay kit by Abcam (ab112146) according to their instructions in order to test
the efficiency of MMP inhibitors. MV3 melanoma cells were cultured in a 24 well plate with (a) media with
no cells, (b) cells treated with vehicle (DMSO) control, and (c) cells treated with GM6001 for 24h. Two
positive controls of recombinant human MMP1 and MMP8 from RnD systems (901-MP & 908-MP) were
used for the assay. The control MMPs were dissolved in assay buffer and a 2mM AMPA working solution
was prepared with assay buffer. The MMP and the test samples were mixed 1:1 vol/vol with the AMPA

working solution and incubated for 1h at 37 °C. The MMP green substrate working solution was prepared
in assay buffer and then mixed 1:1 vol/vol in the black walled 96 well plate and further incubated for 1h.
The samples were then read on a Biotek, Synergy H1 hybrid plate reader at Ex/Em = 490/525 nm.

Recombinant DNA Constructs
The GFP-AktPH construct was obtained from the laboratory of Jason Haugh (North Carolina State
University, Raleigh NC)[2] and cloned into the pLVX-IRES-puro vector (Clontech). The GFP-tractin construct
was a gift from Dyche Mullins (Addgene plasmid # 58473; http://n2t.net/addgene:58473;
RRID:Addgene_58473)[3] and was cloned into the pLVX-IRES-puro vector (Clontech). Paxillin-pEGFP was a
gift from Rick Horwitz (Addgene plasmid # 15233 ; http://n2t.net/addgene:15233;
RRID:Addgene_15233)[4]. mRuby2-CLC was a gift from the laboratory of Sandra Schmid (UT Southwestern
Medical Center). Cells expressing lentiviral vectors were created by following the manufacturer's
instructions for virus preparation and cell infection (Clontech). Cells were selected for expression by
treatment with puromycin, G418, or by fluorescence activated cell sorting.

The photoactivatable PI3K construct (Idevall-Hagren et al., 2012) was created by cloning mCherry-CRY2-
iSH2 (Addgene Plasmid #66839) into the pLVX-neo vector (Clontech). The CIBN-CAAX plasmid was
obtained from Addgene (Plasmid #79574) and cloned into the pLVX-puro vector. Cells expressing both the
mCherry-CRY2-iSH2 and the CIBN-CAAX constructs were selected by treatment with 10 mg/mL puromycin
and fluorescence activated cell sorting. It is critical for the two part cry2 photoactivation system that cells
express sufficient concentration of the CIBN-CAAX construct or the cry2 construct will aggregate in the
cytosol instead of being recruited to the membrane. Thus, the optimal ratio of CIBN:cry2 is greater than
one; cells expressing insufficient CIBN-CAAX will not respond to light. We also noted through the course
of our experiments that cells will stop expressing one or both of these constructs if not kept constantly
under selective pressure. Such a loss of expression will result in non-responsive cells. The PA-Rac1
construct was obtained from Yi I. Wu (University of Connecticut Health Center, Farmington, CT).

Overexpression of fluorescently tagged monomeric actin can perturb cell cytoskeletal dynamics. To avoid
this artifact while imaging tagged actin, we expressed HALO-tagged actin under the control of a truncated
CMV promotor, which results in lower expression of tagged actin than the full length promoter. The
original actin construct features an 18 amino acid linker between mNeonGreen and actin in a pLVX-
shRNA2 vector and was obtained from Allele Biotech. We truncated the CMV promoter, and replaced the
mNeonGreen fluorophore with the HALO tag sequence. The sequence of the CMV100 promoter region is
as follows, with the CMV sequence highlighted and the start codon in bold:

AGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCCCATATATGGAGTTCCGCGTTACATAAC
TTACGGTAAATGGCCCGCCTGGCTGACCGCCGCTAGCGCTAACTAGT**GCCACCATG**

Phase-contrast imaging
Live-cell phase-contrast imaging was performed on a Nikon Ti microscope equipped with an
environmental chamber held at 37°C and 5% CO_2 and imaged with 20x magnification.

Cells on top of gels
Collagen slabs were made from rat tail collagen Type 1 (Corning; 354249) at a final concentration of 3
mg/mL, created by mixing with the appropriate volume of 10x PBS and water and neutralized with 1N
NaOH. A total of 200 μL of collagen solution was added to the glass bottom portion of a gamma irradiated
35 mm glass bottom culture dish (MatTek P35G-0-20-C). The dish was then placed in an incubator at 37°C
for 15 minutes to allow for polymerization.

Cells were seeded on top of the collagen slab at a final cell count of 5000 cells in 400 uL of medium per
dish. The dish was then placed in a 37°C incubator for 4 hours. Following incubation, 1 mL of medium was
gently added to the dish. The medium was gently stirred to suspend debris and unattached cells. The
medium was then drawn off and gently replaced with 2 mL of fresh medium.

Cells embedded in 3D collagen
Collagen gels were created by mixing bovine collagen I (Advanced Biomatrix 5005 and 5026) with
concentrated phosphate buffered saline (PBS) and water for a final concentration of 2 mg/mL collagen.
This collagen solution was then brought to pH 7 with 1N NaOH and mixed with cells just prior to incubation
at 37°C to induce collagen polymerization. Cells were suspended using trypsin/EDTA (Gibco), centrifuged
to remove media, and then mixed with collagen just prior to incubation at 37°C to initiate collagen
polymerization. To image collagen fibers, a small amount of collagen was conjugated directly to AlexaFluor
568 dye and mixed with the collagen sample just prior to polymerization. FITC-conjugated collagen was
purchased from Sigma (C4361).

3D confocal imaging
The cell/collagen mixture described in the previous section was added to Nunc Lab-Tek II Chambered
Coverglass samples holders with a No. 1.5 borosilicate glass bottom (Thermo Scientific). Cells were fixed
with paraformaldehyde and stained with Hoechst and FITC-phalloidin. Images were acquired on a Zeiss
LSM 880 using a Plan-Apochromat 63x/1.4 Oil objective.

Classification of cell morphology

For Figure 1d, we classified the fraction of cells as "rounded" as follows. For cells embedded in 3D collagen,
we labeled as rounded cells with extensive blebbing as well as round cells with few protrusions of any
sort. For cells placed on top of collagen and imaged using phase contrast microscopy, we manually scored
each cell as either rounded or stretched, with the rounded morphology indicating the amoeboid
phenotype.

Zebrafish injection and imaging
B16F10 melanoma cells expressing Lifeact-eGFP were injected into the hindbrain ventricle of 2 days post-
fertilization wildtype zebrafish larvae using previously described protocols.[5] Briefly, B16F10 melanoma
cells were suspended in HBSS. 25-50 cancer cells were transplanted into the hindbrain ventricle of
anesthetized larvae. Injected zebrafish larvae were incubated at 31°C with 0.2 mM PTU to prevent
pigment formation. Live-cell *in vivo* imaging was performed using a Zeiss spinning disc microscope with a
QuantEM EMCCD camera.

High NA TIRF microscopy
Human cervical adenocarcinoma cells HeLa-Kyoto with TALEN-edited ActB fused with GFP (Cellectis,
France) were maintained in DMEM/F12 supplemented with 10% FBS (Invitrogen) at 37°C and 5% CO2, and
imaged using CO2-independent medium (Invitrogen) supplemented with 10% FBS. Cells were confined by
a PDMS stamp in a non-adhesive, PLL-g-PEG (0.5mg/ml) coated chamber of ~3μm height, as described
previously,[6] and imaged 20 minutes after initiating the confinement to observe actin dynamics within
blebs. The high NA TIRF consisted of a standard setup equipped with a 473nm laser 500mW
(Laserquantum), an objective TIRF NA=1.49 (Olympus), and a camera (Andor Zyla 4.2). A single notch filter
was used in the emission light path to block the laser line at 473 nm (Chroma). Acquisition was controlled
by the Andor SOLIS software.

3D cell tracking from phase-contrast movies

Cells were embedded in 2.0 mg/mL pepsinized bovine collagen in Nunc Lab-Tek II Chambered Coverglass
samples holders as described above. Live-cell phase-contrast imaging was performed on a Nikon Ti
microscope as described above. Cells were outlined manually using ImageJ, and position and shape data
were exported for analysis using Matlab. Cell shape was calculated using roundness, given by
$4*area/(\pi*major\_axis^2)$, and cells were classified as either round (roundness > 0.8) or stretched
(roundness < 0.8). Autocorrelation was calculated using the Matlab function *xcorr*. Cell velocity was
calculated from cell centroid positions.

3D light-sheet imaging
3D samples were imaged using either an axially-swept light-sheet microscope[7] or a meSPIM microscope,[8]
both of which provide nearly isotropic, diffraction-limited 3D images. Samples were imaged in phenol red
free DMEM containing 25mM HEPES (ThermoFisher) with 10% FBS and antibiotic-antimycotic (Gibco),
held at 37°C during imaging. Images were collected using sCMOS cameras (Orca Flash4.0 v2, Hamamatsu)
and microscopes were operated using custom Labview software. All software was developed using a 64-
bit version of LabView 2016 equipped with the LabView Run-Time Engine, Vision Development Module,
Vision Run-Time Module and all appropriate device drivers, including NI-RIO Drivers (National
Instruments). Software communicated with the camera via the DCAM-API for the Active Silicon Firebird
frame-grabber and delivered a series of deterministic TTL triggers with a field programmable gate array
(PCIe 7852R, National Instruments). These triggers included analog outputs for control of mirror
galvanometers, piezoelectric actuators, laser modulation and blanking, camera fire and external trigger.
All images were saved in the OME-TIFF format (https://docs.openmicroscopy.org/ome-model/5.6.3/ome-
tiff/). Some of the core functions and routines in the microscope control software are licensed under a
material transfer agreement from Howard Hughes Medical Institute, Janelia Research Campus.

3D cell image analysis
3D light-sheet images of cells were first deconvolved using the Richardson-Lucy algorithm built-in to
Matlab (Mathworks). To reduce deconvolution artifacts, images were apodized, as previously described.[8]
Following deconvolution, we used our previously published u-shape3D analysis framework.[9] to segment
cells, detect blebs, map fluorescence intensity to the cell surface, measure surface motion, and calculate
polarization statistics. Briefly, images of cells were segmented to create a cell surface represented as a 3D
triangle mesh. We used u-shape3D's *twoLevelSurface* segmentation mode, which combines a blurred
image of the cell interior with an automatically thresholded image of the cell surface. Blebs were detected
by decomposing the surface into convex patches, and using a machine learning algorithm to classify the
patches as a bleb or not a bleb. For each patch classified as a bleb, the bleb neck was defined as the
boundary between that patch and neighboring patches. Distance from a bleb neck was calculated at every
face on the mesh as the geodesic distance to the closest bleb neck. To determine the fluorescence
intensity at each mesh face, we used the raw, non-deconvolved, fluorescence image. At each mesh face,
a kd-tree was used to identify the cell-interior voxels within a sampling radius of 1 or 2 μm of the mesh
face. Before averaging the intensity values in these voxels, the intensity values were depth-normalized to
correct for surface-curvature dependent artifacts.[10] The u-shape3D software, as well as the trained
machine learning models used here, are available with the previously published manuscript.[9]

Polarization statistics were calculated by mapping data defined on the cell surface to a sphere, and fitting
the mapped data to a 3D von Mises distribution, which is akin to a spherical normal distribution. We
calculated bleb polarization by representing each bleb by the location on the bleb surface farthest from
the bleb neck, with distances measured on the cell surface. Additionally, since the adhesion images had

substantial fluorescence background, to measure adhesion polarization, we bandpass filtered the raw
images via a difference of Gaussians procedure, selecting for objects between 1 and 6 pixels in radius.

3D collagen image analysis

To enhance linear image features, such as collagen fibers, the 3D collagen images were processed with a
steerable filter of width 2 pixels, as previously described.[8] To emphasize collagen fiber location, some
figure panels, as indicated in the figure legends, show steerable-filter enhanced collagen. Other collagen
images, especially those related to endocytosis, were neither filtered nor deconvolved to avoid the
creation of artifacts. Collagen polarization near the cell surface was measured after mapping image
intensity values from steerable-filtered images onto the cell surface. Following steerable filtering and
automatic thresholding, the nematic order parameter of collagen networks was calculated as described
previously[8], except that the average fiber directionality in each 3D image was used as the reference
direction. The fiber directionality was calculated at each voxel via a steerable filter. Collagen pore size
analysis was also performed as described previously.[11] Images were filtered and then thresholded at 2.5
times the intensity threshold calculated by Otsu's algorithm.[12] To measure pore sizes, for each image, we
first fitted the largest possible sphere into the collagen pores. We then iteratively fitted the next largest
sphere into the pores space minus the volume of previously fitted spheres until no remaining spheres
above a size threshold would fit. We defined the distribution of collagen pore sizes as the distribution of
fitted sphere diameters. Collagen motion was measured using a previously published 3D optical flow
algorithm.[13] This algorithm combines a matching framework for large displacements across frames with a
variational framework for small displacements. We mapped the magnitude of the collagen motion
calculated via optical flow onto the cell surface using the framework for mapping fluorescence intensity
onto mesh faces described above. To create panel 3H, we separated cell surface motion, aggregated
across multiple cells, into bins by magnitude and found the mean collagen motion magnitude associated
with each bin. The collagen sample moves during imaging, and although the average collagen motion in
each frame was subtracted from the measured collagen velocities, residual local motions contribute to
create a non-zero background collagen motion.

3D dextran assay image analysis

To measure the uptake of collagen fragments alongside 70 kDa dextran, we first segmented the cell using
the dextran channel. To do so, we inverted each 3D image, subtracted the median intensity, normalized
by the 99th intensity percentile, subtracted the image background, thresholded, morphologically dilated
by 1 pixel, morphologically eroded by 8 pixels, filled holes, and finally selected the largest image
component. Since the cell is morphologically eroded to a greater extent than it is dilated, the cell
segmentation is effectively shrunk, reducing the effect of segmentation errors on later analysis. To detect
dots of endocytosed collagen and dextran, we employed a previously published multiscale stochastic
filter.[14] For this filter, we used scales of 1.5 to 4 pixels, an α = 0.01, and detected dots only inside the
segmented cell. The p-value distribution shown in Fig. 2o results from testing, for each cell, the hypothesis
that collagen fluorescence intensity is greater at the location of detected dextran dots than elsewhere in
the cell. To calculate the p value for each cell, we randomly picked n collagen intensity values within the
cell 100,000 times, where n is the number of detected dextran dots, and calculated the probability that
the mean of the randomly picked values was greater than the mean of the collagen intensity values at the
true detected dextran dots.

Photoactivation

739 Photoactivation of subcellular regions was performed using a 488 nm laser at 10% power via the FRAP
740 module of a Zeiss LSM780 outfitted with temperature and CO_2 control. Cells for the PI3K optogenetics

were treated with 200uM of PI3K inhibitor IV just prior to photo activation. To assess bleb size change in phase contrast movies, we analyzed multiple blebs within the stimulated region by manually outlining individual blebs at their largest size using ImageJ. Bleb size was measured prior to activation and during activation in the same sub-region of the cell.

Visualization and Statistics

3D surface renderings were made in ChimeraX.[15] Colored triangle meshes representing the cell surface were imported into ChimeraX from u-shape3D as Collada dae files, as previously described.[9] To render collagen, steerable-filtered images were opened directly in ChimeraX and thresholded. To create the rendering of adhesions shown in Figure 3d, the raw paxillin images were bandpassed, admitting objects between 0.5 and 3 pixels in radius, and then median filtered.

Figure 2h has histograms with varied bin sizes. To avoid the existence of bins with very little data, each bin in this panel contains a decile of data. Furthermore, to ease visual interpretation, the time series data in Figure 3k were smoothed using a moving average filter with a span of 5 frames.

All statistical comparisons shown in figures were calculated using a one-sided or two-sided t-test with α = 0.05. Error bars in figures show either 95% confidence intervals or the standard error of the mean, as stated in the figure legends. Number of cells and/or number of different experiments analyzed are given in the figure legends.

Methods References

Quintana, E., Piskounova, E., Shackleton, M., Weinberg, D., Eskiocak, U., Fullen, D. R., Johnson, T. M. & Morrison, S. J. Human melanoma metastasis in NSG mice correlates with clinical outcome in patients. *Sci Transl Med* **4**, 159ra149, (2012).

Haugh, J. M., Codazzi, F., Teruel, M. & Meyer, T. Spatial sensing in fibroblasts mediated by 3' phosphoinositides. *Journal of Cell Biology* **151**, 1269-1280, (2000).

Belin, B. J., Goins, L. M. & Mullins, R. D. Comparative analysis of tools for live cell imaging of actin network architecture. *Bioarchitecture* **4**, 189-202, (2014).

Laukaitis, C. M., Webb, D. J., Donais, K. & Horwitz, A. F. Differential dynamics of alpha 5 integrin, paxillin, and alpha-actinin during formation and disassembly of adhesions in migrating cells. *Journal of Cell Biology* **153**, 1427-1440, (2001).

Roh-Johnson, M., Shah, A. N., Stonick, J. A., Poudel, K. R., Kargl, J., Yang, G. H., di Martino, J., Hernandez, R. E., Gast, C. E., Zarour, L. R., Antoku, S., Houghton, A. M., Bravo-Cordero, J. J., Wong, M. H., Condeelis, J. & Moens, C. B. Macrophage-Dependent Cytoplasmic Transfer during Melanoma Invasion In Vivo. *Dev. Cell* **43**, 549-562, (2017).

Le Berre, M., Zlotek-Zlotkiewicz, E., Bonazzi, D., Lautenschlaeger, F. & Piel, M. Methods for two-dimensional cell confinement. *Methods Cell Biol* **121**, 213-229, (2014).

Dean, K. M., Roudot, P., Welf, E. S., Danuser, G. & Fiolka, R. Deconvolution-free Subcellular Imaging with Axially Swept Light Sheet Microscopy. *Biophys. J.* **108**, 2807-2815, (2015).

Welf, E. S., Driscoll, M. K., Dean, K. M., Schafer, C., Chu, J., Davidson, M. W., Lin, M. Z., Danuser, G. & Fiolka, R. Quantitative Multiscale Cell Imaging in Controlled 3D Microenvironments. *Dev. Cell* **36**, 462-475, (2016).

Driscoll, M. K., Welf, E. S., Jamieson, A. R., Dean, K. M., Isogai, T., Fiolka, R. & Danuser, G. Robust and automated detection of subcellular morphological motifs in 3D microscopy images. *Nat Methods*, (2019).

Elliott, H., Fischer, R. S., Myers, K. A., Desai, R. A., Gao, L., Chen, C. S., Adelstein, R. S., Waterman, C. M. & Danuser, G. Myosin II controls cellular branching morphogenesis and migration in three dimensions by minimizing cell-surface curvature. *Nat. Cell Biol.* **17**, 137-147, (2015).

11 Lomakin, A. J., Cattin, C. J., Cuvelier, D., Alraies, Z., Molina, M., Nader, G. P. F., Srivastava, N., Saez,
P. J., Garcia-Arcos, J. M., Zhitnyak, I. Y., Bhargava, A., Driscoll, M. K., Welf, E. S., Fiolka, R., Petrie,
R. J., De Silva, N. S., Gonzalez-Granado, J. M., Manel, N., Lennon-Dumenil, A. M., Muller, D. J. &
Piel, M. The nucleus acts as a ruler tailoring cell responses to spatial constraints. *Science* **370**,
(2020).
12 Otsu, N. A threshold selection method from gray-level histograms. *IEEE Trans. Syst. Man, Cybern.*
9, 62–66, (1979).
13 Manandhar, S., Bouthemy, P., Welf, E., Danuser, G., Roudot, P. & Kervrann, C. 3D flow field
estimation and assessment for live cell fluorescence microscopy. *Bioinformatics* **36**, 1317-1325,
(2020).
14 Roudot, P., Legant, W. R., Zou, Q., Dean, K. M., Welf, E. S., David, A. F., Gerlich, D. W., Fiolka, R.,
Betzig, E. & Danuser, G. u-track 3D: measuring and interrogating intracellular dynamics in three
dimensions. *BioRxiv*, (2020).
15 Goddard, T. D., Huang, C. C., Meng, E. C., Pettersen, E. F., Couch, G. S., Morris, J. H. & Ferrin, T. E.
UCSF ChimeraX: Meeting modern challenges in visualization and analysis. *Protein Sci* **27**, 14-25,
(2018).

    Extended Data Figures and Legends

Extended Data Figure 1. Melanoma cell morphology and migration. (a) Mean instantaneous velocity of
melanoma cells in 3D collagen, categorized by cell shape (n = 68 cells). (b) Directional persistence as
measured by the directional autocorrelation of single cell trajectories, categorized by cell shape (n = 68
cells). Greater lag times indicate more persistent migration. (c) Maximum intensity projection of a 3D
confocal image of a B16 melanoma cell xenografted into a zebrafish embryo. The imaging site is indicated
on the zebrafish schematic. Surface renderings of 3D light-sheet microscopy images of (d) amoeboid and
(e) mesenchymal melanoma cells in 3D collagen. Scale bar is 10 μm.

Extended Data Figure 2. Digging tunnels through 3D collagen. (a) Maximum intensity projection across
3.2 μm (20 slices) of a light-sheet microscope image of two MV3 melanoma cells expressing GFP-AktPH
(red) in 3D collagen (cyan). The white dashed line indicates the location of a tunnel. (b) Maximum intensity
projection across 18 μm of a light-sheet microscope image of an Ewing sarcoma cell expressing GFP-F-
tractin (red) in 3D collagen (cyan). The white dashed line indicates the location of a tunnel. The collagen
images in a & b were computationally enhanced by a steerable line filter amplifying collinear structures,
and in both panels the scale bars show 10 μm. (c) MMP activity is evaluated by measuring fluorescence
resonance energy transfer (FRET) fluorescence of a generic MMP peptide, measured via plate reader. In
the intact FRET peptide, the fluorescence of one part is quenched by another. After cleavage into two
separate fragments by MMPs, the fluorescence is recovered. 'Medium' shows MMP activity of DMEM cell
culture medium with FBS but no cell contact. 'DMSO' shows activity of cell culture medium incubated with
cells and DMSO vehicle as control for 24 hours. 'MMP8' shows the activity of purified MMP8 protein
dissolved in sample measurement buffer. 'GM6001' shows the activity of cell culture medium incubated
with cells and 40 μM GM6001 for 24 hours. P value via two sample t-test for the comparison between
DMSO and GM6001 is 0.0068, n = 4 separate culture wells for each condition. Error bars show 95%
confidence interval.

Extended Data Figure 3. Comparing path generation in pepsinized and unpepsinized collagen. (a) Maximum intensity projections of a single light-sheet microscope image of an MV3 melanoma cell expressing GFP-AktPH (red) in pepsinized 3D collagen with fibers enhanced (cyan), projected across 1.1 μm (left) and 8.2 μm (right). In the left panel, the white dashed line indicates the location of a tunnel behind the cell. In the right panel, the orange arrow indicates a collagen fiber being dragged behind the cell, and the white arrows indicate collagen fibers being dragged from the sides in the cell's wake. Scale bar show 10 μm. (b) Pore size analysis of pepsinized and unpepsinized 3D collagen samples. Error bars indicate the standard error of the mean (n=6 gels per condition). The pepsinized data was shown in 1b. (c) Nematic order parameter quantifying the extent of collagen fiber alignment in images of pepsinized and unpepsinized 3D collagen containing cells (difference between conditions per a one-sided t-test, p = 0.009). A mean nematic order parameter of 1 indicates high fiber alignment and 0 indicates no overall alignment.

Extended Data Figure 4. Spatial location of internalized collagen and clathrin-containing vesicles. Images show a single optical slice of a 3D light sheet microscope image of an MV3 melanoma cell expressing mRuby2-CLC in FITC-labeled 3D collagen. Orange arrows indicate internalized collagen fragments and white arrows indicate clathrin-containing vesicles.

Extended Data Figure 5. Adhesion localization. (E) Maximum intensity projections across 0.8 μm of a
light-sheet microscope image of a melanoma cell expressing GFP-paxillin (red in left image, black in right
image) in 3D collagen (white in left image). Scale bar indicates 10 μm.

Extended Data Figure 6. Surface motion of a melanoma cell treated with FAK inhibitor 14. Surface
renderings colored by local surface motion of the cell analyzed in 3k. Purple indicates protrusive regions,
whereas green indicates retractive regions. Panel 3f shows PI3K activity localization on the same cell prior
to treatment.

Extended Data Figure 7. PI3K localization is associated with large blebs. (a) A surface rendering of a light-
sheet microscope image of a melanoma cell in collagen, colored by the localization of GFP-AktPH. A
timelapse of the same cell is shown in Movie 5. (b) Relative frequency distributions of GFP-AktPH near the
cell surface reporting PI3K products in large blebs (top decile by volume) vs small blebs (bottom decile by
volume) (n = 34 cells). (c) Frequency distributions of collagen motion near cell surfaces of large blebs (top
decile by volume) vs small blebs (bottom decile by volume) (n = 6 cells).

Extended Data Figure 8. PI3K photoactivation. Spinning disk confocal microscope images showing a single
optical slice of GFP-AktPH biosensor localization in an MV3 cell before, during and after photoactivation
of PI3K in the area indicated by a red circle.

Extended Data Figure 9. PI3K inhibition. (a) Maximum intensity projections across 1.6 μm of a light-sheet
microscope image of an MV3 cell expressing GFP-AktPH, before and after PI3K inhibition. Scale bar shows
10 μm. (b) Temporal response of a different MV3 cell treated with PI3Kα inhibitor IV. From left to right,
shown are mean bleb volume, bleb polarization magnitude, number of blebs, PI3K polarization
magnitude, and mean surface motion magnitude.

Extended Data Figure 10. Wheat Germ Agglutinin (WGA) treatment. (a) Maximum intensity projections of light-sheet microscope images of representative melanoma cells treated with 0 μg/mL or 40 μg/mL WGA. Scale bar shows 10 μm. (b) PI3K intensity as a function of time after WGA treatment in different regions of a cell.

Extended Data Figure 11. HALO-actin and tractin localization within blebs. (a) Time-lapse maximum intensity projections across 3.2 μm of HALO-actin (total actin) and GFP-F-tractin in an MV3 cell. Volumes were acquired every 4.3 sec. (b) Actin/tractin ratio on and off blebs as a function of distance from the bleb neck (n = 5 cells).

Movie Legends.

Movie 1. Collagen movement in soft collagen. Single optical slice of a light sheet fluorescence time-lapse series showing motion of collagen labeled with Alexafluor 568 in the proximity of a single melanoma cell (unlabeled).

Movie 2. Collagen network surrounding a melanoma cell. Movie animates stepping through single slices of a light sheet fluorescence image of collagen labeled with Alexafluor 568 along with a single melanoma cell showing local surface curvature mapped to the surface of the cell.

Movie 3. A cell tunneling through collagen. Movie shows a time-lapse sequence of a single slice of a light sheet microscope image of a melanoma cell expressing GFP-AktPH (green) and collagen (red).

Movie 4. Animation showing blebs identified by u-shape 3D. Movie shows rotation of a 3D rendering of the surface of a single melanoma cell, imaged using light sheet microscopy. Blebs identified by u-shape 3D are shown in different colors.

Movie 5. PI3K polarity in a migrating cell. Movie shows a time-lapse sequence of the surface of light sheet microscope images of a melanoma cell expressing GFP-AktPH. The local intensity of GFP-AktPH is projected onto the surface of the cell, according to the colormap shown in Extended Data Figure 7a.

Movie 6. Photoactivation of PI3K in a blebbing cell. Time-lapse sequence of a single optical section of a confocal microscope acquisition of an MV3 cell expressing GFP-AktPH. Photoactivation of Cry2-iSH2/CIBN-CAAX occurred via activation of the FRAP module in the location and time shown by the green circle.