1

4   A tractable bottom-up model of the yeast polarity genotype-phenotype map for evolutionary

5   relevant predictions

9   Werner Karl-Gustav Daalman[1], Els Sweep[1], Liedewij Laan[1,2,*]

13   [*] Corresponding author

14   E-mail: L.Laan@tudelft.nl

[1] Department of Bionanoscience, TU Delft, 2629 HZ Delft, the Netherlands
[2] Lead contact

Abstract

Accurate phenotype prediction based on genetic information has numerous societal applications, such as crop design or cellular factories. Epistasis, when biological components interact, complicates modelling phenotypes from genotypes. Here we show an approach to mitigate this complication for polarity establishment in budding yeast, where mechanistic information is abundant. We coarse-grain molecular interactions into a so-called mesotype, which we combine with gene expression noise into a physical cell cycle model. Firstly, we show with computer simulations that the mesotype allows validation of the most current biochemical polarity models by quantitatively matching doubling times. Secondly, the mesotype elucidates epistasis emergence as exemplified by evaluating the predicted mutational effect of key polarity protein Bem1p when combined with known interactors or under different growth conditions. This example also illustrates how unlikely evolutionary trajectories can become more accessible. The tractability of our biophysically justifiable approach inspires a road-map towards bottom-up phenotype modelling beyond statistical inferences.

Introduction

Many fields, such as personalized medicine (Li, 2011), agriculture (Cobb et al., 2013), chemical production (Oud et al., 2012) and forensics (Kayser and Schneider, 2009), will greatly benefit from better understanding of how traits connect to genes, the so-called genotype-phenotype (GP-) map. However, resolving this connection is generally not straightforward even for known heritable traits (Eichler et al., 2010). For example, multiple genes can contribute to a single trait (polygenic inheritance) while multiple traits can emerge from a single gene (pleiotropy) (Wagner and Zhang, 2011). Frequently, mutational effects have been shown to be non-additive in (model)

species as humans (Cordell, 2009; Zuk et al., 2012), *Escherichia coli* (Wünsche et al., 2017) and

Saccharomyces cerevisiae (budding yeast) (van Leeuwen et al., 2016). This phenomenon is known

as epistasis. Theoretically, epistasis surfaces easily based on metabolic network analysis

(Kryazhimskiy, 2021), and some molecular origins are known (Lehner, 2011). Although

sometimes inconsequential for fitness evolution (Kryazhimskiy et al., 2014), epistasis complicates

the predictions of phenotype and consequently gene evolution (Papp et al., 2011; Miton and

Tokuriki, 2016). Therefore, epistasis predictions pose an important challenge for GP-map models.

In order to unravel the complication epistasis imposes, intermediate levels in the GP-map

are commonly employed (Blanco-Gómez et al., 2016). An intermediate level is any quantity more

complex than individual proteins, but less complex than the phenotype level. This level addition is

meant to produce a more modular and hence more tractable GP-map (Figure 1A). Ideally, a level

that clarifies trait generation also elucidates the handles for evolution, the reverse path in the GP-

map. While phenomenological or statistical level formulations may suffice for predictions, they are

not designed to provide insight into the GP-map. We therefore formulate a biophysically sound

intermediate level from the bottom-up, which we refer to as the "mesotype".

To test our mesotype approach, we model polarity establishment in *S. cerevisiae*, the cell

cycle step where budding yeast breaks its spherical symmetry to direct bud growth (see Figure 1B).

The challenge is reproducing the ample epistasis exhibited in the polarity network in e.g., doubling

times (Laan et al., 2015). Conveniently, the molecular interaction network for polarity

establishment (simplified in Figure 1C) was recently modelled in broad agreement with literature

(Brauns et al., 2020). In short, polarity is essential for cell proliferation and relies on the small

GTPase Cdc42p (Adams et al., 1990). To cluster the active form of Cdc42p, bound to a GTP

molecule, on one plasma membrane patch, mutual recruitment of Cdc42p and Bem1p complexes

containing Cdc24p occurs (Goryachev and Pokhilko, 2008; Kozubowski et al., 2008; Klünder et

al., 2013; Witte et al., 2017). Cdc24p activates Cdc42p locally (Woods et al., 2015), while Cdc42p

is inactivated by GTPase activating proteins (GAPs), namely Bem2p, Bem3p (Zheng et al., 1994),

Rga1p (Stevenson et al., 1995) and Rga2p (Smith et al., 2002). In absence of Bem1p, transport

presumably occurs by actin (Freisinger et al., 2013) and/or Cla4p (Tiedje et al., 2008). Clusters of

active Cdc42p arise by locally saturating the GAPs (Brauns et al., 2020). In the Δ*bem1* background,

polarity establishment benefits from deleting the gene *NRP1*, possibly by influencing cell cycle

timing (Laan et al., 2015). However, Nrp1p's molecular mechanism is currently unknown.

Based on these molecular details and associated rigorous analysis of reaction-diffusion

equations, we construct the genotype-phenotype map for yeast cell polarity by coarse-graining.

One particular result in (Brauns et al., 2020) motivates simplifying polarity success to certain

GAP/Cdc42p dosage ratios (Figure 1D). More specifically, a minimum Cdc42p concentration

exists below which polarization is not possible. This simple rule on protein dosage defines the

mesotype in this context and facilitates the understanding of epistasis emergence while sharply

reducing computational costs.

To complete our GP-map model, the mesotype is incorporated into a physical cell cycle

model, which is further composed of simple volume growth and stochastic protein production

(Figure 1E), to reproduce phenotypes as epistasis. In this paper we assess the quality of our

predictions by comparing these to documented interactions, validating the underlying molecular

polarity model quantitatively. Our tractable approach also illustrates how epistasis and therefore

feasible evolutionary trajectories depend on growth conditions. Finally, we show our framework

relies on biofunctional (and ideally mechanistic information) of the key proteins to yield

informative predictions, which delineates the appropriate conditions for applying our method.

Results

Cell cycle model design with the mesotype as level between genotype and phenotype

We modelled the yeast cell cycle as a process involving three modules, namely (I) polarity

mesotype, (II) stochastic protein turnover and (III) cell size expansion (Figure 1E). We integrated

these modules into a simulated population of cells, with properties size and protein content. In a

Gillespie-style algorithm, the properties are updated per cell (Figure 2A) to yield population

dynamics (Figure 2B). We sketch the essence of each module, and further details are found in the

Materials and Methods. Moreover, a summary of model parameter values is given in Table S1.

For module I, we reduced the biochemical network for polarity to the mesotype of a minimal

[Cdc42p] threshold (dotted line Figure 2A). GAP mutations shift this threshold downward, which

in part constitutes the polarity mesotype checkpoint (see module III). Upon checkpoint passage,

cell expansion continues for a time t_{pol}, which increases exponentially with the excess [Cdc42p].

This time period simplifies the functional dependency uncovered with the aforementioned analysis

of the underlying reaction-diffusion equations (Brauns et al., 2020).

For module II, we only consider Cdc42p to induce cell-to-cell variability, as GAP dosage

population noise (Chong et al., 2015) is much smaller than the noise of Cdc42p (coefficient of

variation 0.83, this study). As mRNA of Cdc42 lives much shorter than the protein (Christiano et

al., 2014; Grigull et al., 2004), we assume bursty Cdc42p production (Figure 2A). By contrast, we

assume deterministic Cdc42p degradation due to its high abundance (Kulak et al., 2014).

For module III, we assume two stages of constant outer membrane area growth in G1 and

S/G2/M phase respectively. In the first stage, spherical mother cells of radius $r_m(t)$ grow until the

polarity mesotype checkpoint is passed and polarization is completed, or until cell death when $r_m(t)$

exceeds maximum radius r_{max} (black arrow Figure 2A). The polarity checkpoint entails next to the

$[Cdc42p]_{min}$ of module I, exceeding a minimal radius r_{min}, and a minimal time since last division

$t_{G1,min}$,. The last two criteria represent e.g., cell size dependent control of Cln3p nuclear arrival by

Ydj1p (Vergés et al., 2007). The *nrp1* deletion is phenomenologically incorporated by reducing

this minimal time as suggested by Laan et al., 2015). In the next stage, a bud with radius $r_b(t)$ grows

to 70% of the mother volume (the second checkpoint). Then, mother and bud restart the cell cycle

independently. Resulting cell expansion rates are in decent agreement with literature (Ferrezuelo

et al., 2012; Goranov et al., 2009), see Materials and Methods: size expansion.

Finally, we validated the convergence of the phenotypes dosage distribution, size and

doubling times (Figure 2B). In particular, the size and doubling time change 0.6% and 0.9%

respectively across the last 200 minutes, well within the typical experimental error of (Laan et al.,

2015). Confirmation of doubling time convergence is also depicted in Figure S1.

The mesotype enables detailed validation of the underlying biochemical model and bottom-

up interpretation of phenotype emergence

Our physical cell cycle model relies on the validity of the underlying biochemical network

model, which was coarse-grained to the mesotype. While the biochemical network model was

tested in (Brauns et al., 2020), our cell cycle model allows otherwise unfeasible quantitative

phenotype descriptions, improving the validation precision of the biochemical network model.

To this end, we considered 20 previously documented experimental values from (Laan et

al., 2015). These constitute doubling times, whose reciprocals are defined as fitness, and cell cycle

times. The three most prominent, non-trivial phenotypes are (i) strong epistasis in growth rates

between GAP mutants only in the Δ*bem1* background, (ii) strong epistasis between *BEM1* and

NRP1, and (iii) non-monotonous optimization of G1 times for (reconstructed) experimentally

evolved mutants starting from Δ*bem1*. For the latter phenotype, the acceleration of G1 speed of the

Δ*bem1* cells, despite their poor fitness, compared to WT cells is particularly noteworthy.

To model the mutants in our target observable set, we required five fitting parameters.

Firstly, the *BEM1* and Δ*bem1* have two different values for the mesotype. Secondly, because the

mesotype scales with GAP concentration (Figure 1D), any combination of *bem2* and *bem3* null

mutations is described by two additional parameters. Lastly, the *nrp1* mutant is

phenomenologically incorporated through a (25%) decrease in $t_{G1,min}$.

The first phenotype of interest, GAP epistasis, is quantitatively well described (Figure 3A,

left). This description is robust to several model assumption modifications (Figure S2). Particularly

in presence of *NRP1*, fitness values were modelled in accurate accordance with experiments of

(Laan et al., 2015). As expected, less cells produce sufficient Cdc42p for high mesotypes, leading

to increased cell death and lower fitness (Figure 3B). Therefore, we see a diffuse viability threshold

despite a sharp mesotype, as in previous experiments with inducible Cdc42p (Brauns et al., 2020).

The four fitted mesotype thresholds are consistent with a $\Delta bem3$ effect that is twice as large

as for $\Delta bem2$. Given the GAP abundancies (Kulak et al., 2014), this sets the *in vivo* Bem3p effective

GAP activity to be almost four times that of Bem2p. This robust result (Figure S2) suggests that

the GAPs are functionally more similar *in vivo* than measured *in vitro,* where Bem2p activity was

negligible compared to Bemp3 activity. (Zheng et al., 1994).

If we consider the second phenotype of interest, *BEM1-NRP1* epistasis, we focus attention

to the $\Delta nrp1$ background. There, fitness values (Figure 3A, right) are not always well fitted (5/8

correct within experimental error), although these mutants have relatively large experimental

uncertainties. Nevertheless, the strongest feature, the *BEM1-NRP1* epistasis, is at least qualitatively

described. This description confirms the intuition offered by the mesotype. If the *nrp1* deletion

reduces the mandatory G1 waiting time $t_{G1,min}$, $\Delta bem1$ cells have more chance to exploit temporary

Cdc42p overproduction before excessive dilution, thereby improving fitness (Figure 3C).

Quantitatively, we explain one-third of the epistasis following the definition of (da Silva et al.,

2010). An alternative G1 time formulation cannot strongly improve this result (see Materials and

Methods, G1 timing). In the next section, we confirm that incorporating mutants

phenomenologically rather than based on molecular information limits the accuracy of phenotype

description.

Lastly, we turned to the third phenotype of interest concerning non-monotonous G1-time

shifts during adaptation. These shifts are an example of the more detailed traits that can be

modelled. To mimic the poorer content of synthetic medium in (Laan et al., 2015), we performed

the simulations with half the normal membrane area rates. The observed trends in G1 times along

the evolutionary trajectory from WT to the fully evolved mutant in that paper were qualitatively

matched, including the unusual G1 time decrease for the Δ*bem1* (Figure 3D). The mesotype

clarifies this subtle phenotype. The Δ*bem1* cells are relatively larger and less limited by the

minimum size criterion r_{min}. This eliminates a potential waiting step in G1 for this background,

which may allow the cells to pass G1 faster if these have lucky Cdc42p overproduction. Other

phases are on average extended by the relatively high minimum Cdc42 concentration threshold.

In summary, the three phenotype examples illustrate two main points. Firstly, comparison

of experimental data with simulation of our physical cell cycle model further confirms the current

biochemical model view of polarity. Secondly, the mesotype framework exhibits the tractability

needed for understanding phenotype emergence, as demonstrated with several examples.

The mesotype generates model predictions for genetic interactions.

Poorer medium quality reduces fitness differences.

After establishing the value of the mesotype in trait generation, we turn to the reverse

transition in the GP-map. This transition is embodied by evolution (Figure 1A), since phenotypes

such as fitness determine the selective pressure to shape genotypes. Environmental factors are

important for growth, with even subtle changes noticeable under highly controlled laboratory

settings (Atolia et al., 2020). Given that historical evolution has occurred in the wild, where

conditions are expected to be much more variable and more often difficult than not, adaptive

trajectories can be very different than for laboratory evolution. For example, step-wise evolution

of Bem1p has already been hypothesized (Brauns et al., 2020) given fitness in lab conditions. Yet,

more difficult conditions, such as slower growth and concordant ease of exceeding the mesotype,

suggests that the evolution of cells lacking Bem1p may be more likely than initially anticipated.

To further substantiate our mesotype intuition on condition-dependence for Bem1p

evolution, we first simulate the effect of changes in growth media quality through a change in

membrane area growth rates. We considered a roughly three-fold area growth rate range that caused

WT fitness to span between 0.5 and 1 (normalized to maximum growth). We further assume

Cdc42p expression remains the same across media, which is at least true upon switching from

dextrose to ethanol, an inferior carbon source (Smith and Kruglyak, 2008).

As a result of our model simulations, Figure 4A displays the fitness ratio between the

Δ*bem1* and *BEM1* background, as a function of GAP concentration and medium quality.

Intuitively, we expected Δ*bem1* cells to benefit greatly from less Cdc42p dilution and the extra

time to exceed the mesotype threshold. We indeed observe the trend of smaller fitness differences

for decreasing GAP concentrations and decreasing medium quality in our simulations. This

observation suggests poorer media acts as a fitness equalizer, facilitating the evolution of Bem1p.

To experimentally test our hypothesized environmental effect, Figure 4B demonstrates the

effect of poorer medium on fitness with and without Bem1p. In this case, we modulated medium

quality through sugar content (from 2% to 0.1%). To avoid lengthy exposure of the Δ*bem1*

background to high selective pressure, this genotype is mimicked by auxin inducible degradation

(Nishimura et al., 2009) of Bem1p. As it is difficult to exactly integrate the media conditions into

the simulations, the match cannot be expected to be quantitative. However, the nullifying effect of

poor media on the fitness differences between the Δ*bem1* and *BEM1* backgrounds is visible. The

relative fitness compared to WT for the effective Δ*bem1* background in the 'poorest' media (0.1%

dextrose) is significantly better compared to the 'richest' and intermediate medium conditions at

2% and 0.5% dextrose (one-sided Welch's t-test, p-value $1.3 \cdot 10^{-3}$ and $3.3 \cdot 10^{-3}$ respectively), also

when considering the Holm-Bonferroni correction (Holm, 1979). Thus, the mesotype framework

can provide a new, intuitive explanation of otherwise non-trivial environmental interactions that

are relevant for evolution.

The polarity mesotype predictions on epistasis become useful when functions of

mutated genes are known. While we showed that our cell cycle model produced accurate epistasis

predictions, the mechanistic information we used is not always available. We therefore determined

the required mutant information content for qualitative epistasis predictions with modules where

no mechanistic information is assumed. For this purpose, we considered high-throughput data on

BEM1 interactors, assuming many are documented given the strong *bem1* phenotype. Intersecting

BEM1 interactors and mutants with various phenotype specificities, i.e. information content, yields

three sets. (Figure 4C). From low to high information content, the coarse phenotype set covers

(competitive/fermentative) fitness mutants, the mid-detail set G1 mutants (in size/speed), and the

functional set covers proteasomal, phospholipid or ribosomal mutants. Our model subdivides each

set by sign of the predicted epistasis, positive (X_{n+}) or negative (X_{n-}). These predictions can be

intuitively constructed as explained in the next paragraphs or derived from simulations (Table S3).

We then determined for which set, and thus information content, the observed epistasis sign

prevalence qualitatively matched predictions (see Materials and Methods, quantification and

statistical analysis).

Firstly, the coarse phenotype mutants are incorporated in the membrane area growth rate

C_2. Similar to the media effect in Figure 4A, smaller rates mitigate the $\Delta bem1$ effect. The model

prediction is hence that epistasis with $\Delta bem1$ for deleterious mutants (ε_{1-}), which slow down

growth, is more often positive than the epistasis for beneficial mutants (ε_{1+}), We require a Bayesian

odds ratio $Pr(\varepsilon_{1+} > 0)/Pr(\varepsilon_{1-} > 0) > 10$ (Kass and Raftery, 1995) or equivalently, a probability that

our epistasis statement is true of $> 90\%$. However, in the literature data only 14% of the deleterious

mutants have positive epistasis with $\Delta bem1$, compared to 19% of the beneficial mutants.

Consequently, the data shows no support for our statement on epistasis prevalence given coarse

mutant information, with the 18% chance (Figure 4D) worse than a coin-flip (50%).

Secondly, the mid-detail mutants are incorporated by changing G1 waiting time $t_{G1,min}$ (for

G1 speed mutants) and additionally minimal radius before Start r_{min} (for G1 size mutants). As was

the case for to Nrp1p (Figure 3C), mutations which shorten Start disproportionally benefit the

$\Delta bem1$ background. Therefore, the model prediction is that mutants fast or small in G1 have more

positive epistasis with $\Delta bem1$ than mutants that are slow or large in G1. While positive epistasis is

more abundant for fast/small in G1 mutants (27% against 20%), the experimental evidence is not

yet compelling, with 71% chance that this model claim is true.

Lastly, the functional mutant set is incorporated by increasing the Cdc42p half-life for proteasomal

mutants, decreasing membrane growth rate C_2 for phospholipid mutants and decreasing average

Cdc42p burst size for ribosomal mutants. The proteasomal and phospholipid mutants mitigate the

problematic lack of Cdc42p in the $\Delta bem1$ cells. These two mutant types should therefore exhibit

more positive epistasis than the ribosomal mutants, which lowers [Cdc42p]. From the literature

data, 29% of the proteasomal and phospholipid mutants have positive epistasis, much more than

the 11% of the ribosomal mutants. There is hence strong positive evidence for our epistasis

prevalence statement, which is true with around 95% certainty. This shows we minimally require

functional information on mutants for meaningful epistasis predictions, once we have a core where

mechanical information is known.

Discussion

Epistasis forms a general hurdle for reverse-engineering the GP-map, complicating

modelling of protein networks and in turn limiting the predictability of phenotypes and evolution.

To alleviate the GP-map complexity, we tested the mesotype as an intermediate level between

genotype and phenotype. This level emerges from coarse-graining the biophysics of the underlying

biochemical network, which in our case corresponded to budding yeast polarity (Figure 1). We

employed the mesotype, which in our context is the minimum [Cdc42p] to polarize, in a tractable

cell cycle model with simple volume growth and stochastic protein production, making bottom-up

reconstruction of phenotypes and epistasis feasible and insightful.

Firstly, we validated the polarity network model to unprecedented detail by our cell cycle

model. Good quantitative agreement with literature was found for 13 out of 16 polarity mutant

doubling times, and qualitative agreement for documented unintuitive G1 times (Figure 3).

Phenomenological linkage of minimum G1 time to Nrp1p, a protein normally not included in

polarity models, also provided fruitful model predictions. Thus, our cell cycle model also provides

direction to deciphering the remaining mechanistic unknowns in the polarity network.

Secondly, we showed how the mesotype elucidates epistasis. We hypothesize that lower

quality growth medium extends protein production time, alleviating minimum protein

concentration thresholds. We verified this for yeast polarity, where the [Cdc42p] mesotype

threshold proves less problematic for the Δ*bem1* background in poorer media in simulations and

experiments (Figure 4A and 4B). The Δ*bem1* case fits the more general picture that

haploinsufficiency in YPD is typically lifted in poorer medium (Deutschbauer et al., 2005). By the

same token, evolvability of (near-)essential genes may be enhanced under difficult circumstances,

where fitness values across genetic backgrounds converge. In our example, step-wise evolution of

Bem1p is still feasible given laboratory conditions (Brauns et al., 2020). Yet, for other proteins,

medium quality change may be the only manner to circumvent fitness valleys.

Finally, we determined the key for successful epistasis predictions. As exemplified by the

robustness of results to many alternative growth detail formulations, the key here is combining the

mesotype with dosage noise. In the case of a sharp mesotype threshold as for polarity, acting on a

noisy protein like Cdc42p (about four times the typical coefficient of variation in (Chong et al.,

2015)), epistasis with a strong perturbation as with Δ*bem1* makes other mechanistic details

irrelevant. Indeed, when examining high-throughput literature data, functional information proved

necessary and sufficient to generate meaningful predictions for epistasis (Figure 4D), in line with

the ontotype strategy (Yu et al., 2016). As the sharp mesotype implies essentiality (around 19% of

yeast genes (Giaever et al., 2002)) or toxicity, generally other mesotypes are more appropriate. Yet,

given the simple functions with which fitness landscapes as function of single genes can be fitted

(Keren et al., 2016), the mesotype approach still seems feasible to model many biological networks.

Using our findings on yeast polarity as a template, we envision a road-map to apply to

general genotype-phenotype maps (Figure 5). The core functional component, in this case polarity,

is to be modelled by justifiable coarse-graining, which results in the mesotype of the system. This

mesotype in turn emerges from functional subunits (Brauns et al., 2020), identifiable from the

rigorous analysis of the underlying biophysics. Currently, yeast polarity stands alone at the ideal

intersection between complexity and required mechanistic knowledge. Mesotype generation for

other systems would again require detailed numerical analysis of reaction-diffusion systems, or

results more simply from dose-reponse curves when spatial information is not essential for the

chemical reactions. But once multiple model systems (such as the PAR protein system in

Caenorhabditis elegans (Goldstein and Macara, 2007)) have been described in this manner, it may

be possible to construct a limited library of recurring subunits. Based on this library, construction

of the corresponding mesotype may be much simpler. In combination with a coarse-grained view

of cell growth and noisy protein production, this paves the general path to bottom-up (population)

phenotype prediction.

Limitations of the study

Yeast polarity, our system of interest, is an example of a self-organization network, where we

cannot by default assume tractable Michaelis-Menten kinetics, because we cannot neglect the

spatiotemporal interplay of protein species. In other model systems where proteins can be assumed

to be well-mixed, other modelling approaches have been shown to be highly effective. For

example, in budding yeast there is the iMeGroCy model (Pelillo et al., 2018), more than 100 protein

species were covered in a mechanistic model of macrophage polarization (Zhao et al., 2021), and

modelling metabolism in *S. cerevisiae* can even encompass thousands of protein species (Sánchez

et al., 2017).

Although mechanistic information is generally not widely available, this is not always a problem.

As we demonstrated in Figure 3A and more specifically in Figure 4D, we extended predictions

beyond polarity, where much is known, with mutants that affect the cell in other modules, where

less details are known, as for example the case with Nrp1p. This extension is still useful for epistatic

predictions if there is at least functional information about the added mutant gene, but becomes

dubious once the mutant information is more generic.

Another consideration is to what extent our approach will work for other model systems where a

core module depends on the intricate interplay of spatiotemporal interactions. The mesotype relies

on coarse-graining abundant, detailed mechanistic information concerning the core biochemical

network, and this property currently prohibits application to many other self-organizing systems.

Concerns are that some protein networks, particularly for complex organisms, become too complex

to coarse-grain, or that including the diverse environmental dependencies required to fully reveal

biological functions (Bergelson et al., 2021), also sharply increases complexity. After all, other

systems that share the necessary biochemical network knowledge, for example the Min system in

E. coli (Ramm et al., 2019), exhibit less complexity, while for systems with higher complexity

insufficient mechanistic information is available.

Presumably, advances in the near future in systems with insufficient information will follow. For

example, , recent theoretical advances provide a promising tractable framework for the PAR-

system of *C. elegans* (Geßele et al., 2020). But for more complex networks, our method may take

several decades to fully roll-out. For example, the time span between discovery of the first Cdc42p

GAP and revealing its mechanism in yeast polarity spans three decades (Brauns et al., 2020; Zheng

et al., 1994). The key question on whether the definition of a tractable mesotype following rigorous

analysis of the biochemical network remains possible for more difficult networks, is therefore still

awaiting a definitive answer.

Author Contributions:

Conceptualization, WD and LL; Methodology, WD; Software, WD; Validation, WD and ES; Formal Analysis; WD and ES; Investigation, WD, ES and LL; Data Curation WD and ES; Writing - Original Draft, WD; Writing - Review & editing, WD, ES and LL; Visualization, WD and ES; Supervision, LL; Project Administration, LL.

Acknowledgements

We thank Marit Smeets for flow cytometry measurements of two strains. We gratefully acknowledge Leila Iñigo de la Cruz for growth data on RWS1421 and constructing strains yLIC132 and yLIC133. We express our gratitude to Enzo Kingma and Wessel Teunisse for constructing strain yWT3. Additionally, we thank Fridtjof Brauns and Jos Zwanikken for careful reading of the manuscript and Sophie Tschirpke for help in optimizing figure formatting. We acknowledge (Smith et al., 2015) for making the viridis colormap generously available. LL and WD gratefully acknowledge support from the Netherlands Organization for Scientific Research (NWO/OCW), as part of the Gravitation Program: Frontiers of Nanoscience. LL and ES gratefully acknowledge funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant agreement No. [758132]).

Declarations of interest

The authors declare no competing interests.

398   **Figure titles and legends**

Figure 1. Decomposition of the genotype-phenotype map with the mesotype as an intermediate, showcased in yeast polarity.

Figure 1 Legend

(A) General conceptualization of a genotype-phenotype (GP-) map containing epistatic interactions. Introduction of an intermediate level (right) ideally simplifies the connections between genes to phenotype, improving tractability for trait generation (upwards trajectory) and evolution (coupling downwards from the phenotype). (B) Visualization of yeast polarity as the function template for decomposing a GP-map. Brightfield (left) and widefield fluorescence (right) example images of a polarizing budding yeast cell (yLL129), scale bar 5 µm. Key for polarity is clustering to one zone on the membrane of active Cdc42p, of which a binding partner (Spa2p) is fluorescently labelled in the images. This zone marks the site of polarized growth. (C) Schematic model overview of core polarity protein network (proteins not to scale). A positive feedback for (active) Cdc42p-GTP is mediated by the Bem1p-Cdc24p complex, and to lesser extent by Cla4p. Nrp1p represents a mechanistic unknown. (D) Phase diagram summarizing the GP-map, depicting the phenotype viability (green) as function of genotype (purple) through Cdc42p (active and inactive) and GAP concentration in the cell with or without Bem1p. The GP-map contains epistatic interactions, for example the same increase in GAP concentration can yield inviability in the $\Delta bem1$ background but not in the *BEM1* background. An intermediate, the 'mesotype', can be identified here as the limiting Cdc42p concentration (blue). (E) Implementation of the mesotype into a physical cell cycle model, to tractably decompose the polarity GP-map. Starting from G1, every cell aims to divide once the polarity mesotype checkpoint has passed. This implies mother radius r_m exceeding minimum radius r_{min}, time thus far in G1 t_1 exceeding minimum G1 time $t_{G1,min}$, and a Cdc42 concentration (abbreviated as $[P]$) exceeding mesotype $[P]_{min}$. If maximum size r_{max} is exceeded in the process, the cell dies. If the mesotype checkpoint is passed, the mother continues to grow

isotropically for polarization time t_{pol}, which is at least $t_{pol,min}$. Then, only the bud grows until the

second, coarse-grained growth checkpoint, which involves bud size must be a certain fraction of

mother size before division. Throughout all phases, Cdc42p is subject to stochastic protein

production and deterministic degradation. See also Table S1.

Figure 2. Verification of our physical cell cycle model.

Figure 2 Legend

(A) Example trace of the Cdc42 copy number (purple) and concentration (emerald) of a single cell,

which are subject to protein production and degradation (and dilution for concentrations). The cell

must exceed the mesotype threshold ([Cdc42p]$_{min}$) before division can take place. When this is

delayed for too long, the cell expands beyond the predefined maximal size r_{max} and dies (after

almost 15h). (B) Convergence of Cdc42 copy number distribution during simulations. Simulated

time since ancestor is approximate as birth times of the cells in the starting population are

distributed across an 83 min. bandwidth. The inset shows how the estimates of the population

doubling time and the average effective cell size equilibrate as a function of time. See also Figure

S1 concerning the doubling time convergence, and Figure S3 concerning measured stochasticity of

Cdc42p expression.

Figure 3. Our physical cell cycle model including the mesotype permits detailed and tractable

validation of the underlying biochemical polarity model.

Figure 3 Legend

(A) Experimental fitness values relative to WT (phenotypes) for 16 different polarity genotypes

(Laan et al., 2015) denoted by diamonds, which are fitted by our physical cell cycle model as

depicted by the dark and light lines for Δ*nrp1* and *NRP1* background respectively. GAP genotypes

can be linearly linked to the minimum Cdc42 concentration to polarize, the mesotype, as displayed

on the top horizontal axis. Error of the *BEM2 NRP1* was not available and conservatively guessed.

See also Figure S2. (B) Graphical representation of the effects of GAP and *bem1* deletions from

the mesotype perspective. The stochasticity of exceeding the minimal [Cdc42p], the mesotype,

only becomes apparent in absence of Bem1p and with high GAP concentrations. (C) Graphical

representation of the effects of the *nrp1* deletion from the mesotype perspective. When cells are

less fit and become larger, the minimal G1 time reduction associated with the *nrp1* deletion

becomes most apparent. Otherwise, Start is more frequently set by the minimum size criterion. The

added time for lucky overproduction of Cdc42p is beneficial for the backgrounds that suffer from

Cdc42p underproduction, like the Δ*bem1*. (D) The detailed phenotype of minimum G1 time as

displayed by WT and four polarity mutants, comprising a single evolutionary trajectory.

Experimental values are from (Laan et al., 2015) in light emerald (defined there as tie to first

polarity spot), model values are in dark emerald (defined here as the time in G1 until both the size

and time criteria are met). Both cases are normalized to their respective WT values, such that each

column denoted the relative change in G1 time compared to the previous step in the trajectory. See

also Table S2.

Figure 4. The mesotype allows evolutionary relevant predictions on the effect of medium

quality and epistasis.

Figure 4 Legend

(A) Simulated fitness differences between *BEM1* and Δ*bem1* backgrounds as a function of medium

quality, which is integrated in the cell cycle model through varying cell membrane area growth

rates. The medium quality translates to an associated WT fitness, which sets the horizontal axis.

Different GAP concentrations are denoted by different colors. Generally, poorer medium reduces

differences in fitness and genetic interactions between GAPs when comparing the *BEM1* and

Δ*bem1* backgrounds. (B) Experimental assay comparing the effect of the *bem1* deletion in rich and

poor medium. Bars denote the fitness is plotted of the effective Δ*bem1* background (*BEM1-AID*

with added auxin) relative to the *BEM1* background (all with *BEM2* and *BEM3*), for three medium

conditions. From left to right, quality of the YPD medium increases, with increasing dextrose

concentration as indicated on the axis. Asterisks denote significant differences. (C) Workflow for

model prediction on epistasis ε of various mutants with $\Delta bem1$. Mutants are divided into three sets

by the intersection of Bem1 interactors with mutants of various phenotype specificity (and hence

varying model implementation). Each set consists of two subsets, depending on the subsequent

model prediction on epistasis sign. For each subset, the beta posterior density of the observed

positive epistasis fraction can be constructed (from a binomial likelihood and uniform prior). The

model prediction given the information per set is that + subset has more positive epistasis than -

subset. The probability of a true prediction is then defined as the area below the posterior density

of the difference in positive epistasis abundance in both subsets. See also Table S3. (D) Probability

bars reflecting Bayes factors for the model hypothesis; the ratio between the odds that the model

prediction on positive epistasis abundance differences in subsets is true and false.

Figure 5. Proposed flow chart for phenotype predictions through intermediate levels.

Figure 5 Legend

Bottom-up approach for phenotype prediction from genotype through mesotypes, which result

from selecting the appropriate functional subunits. The mesotype will typically result from dose-

response curves, which are relatively simple to derive from Michaelis-Menten kinetics when there

is no spatial dependence. Otherwise, either rigorous numerical analysis of reaction-diffusion

equations is required, which may in the future be bypassed when more model systems are analyzed

(e.g., polarization in *S. cerevisiae*, the min-system in *E. coli*, PAR-system in *C. elegans*). From a

sufficiently large library of functional subunits, the applicable mesotype for the system of interest

may be extrapolated. Bottom-half in each box exemplifies the flow chart with the yeast polarity

case.

Materials and Methods

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, Liedewij Laan (L.Laan@tudelft.nl).

Materials availability

Strains generated in this study are available from the Lead Contact on request, we will require a

completed Materials Transfer Agreement due to TU Delft policy.

Data and code availability

The datasets and code generated during this study are available as Supplemental Information.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Microbe strains

All strains used in this study are *S. cerevisiae* strains in the W303 background. A full list of strains used in this study can be found in the Materials and strains table. Cells are kept for long-term storage in a frozen glycerol stock at -80°C, and are cultured for experiments at 30°C by inoculating in liquid fresh media. Strains RWS116 and RWS1421 require absence of methionine in the media for growth, so we used YNB (0.69% w/v, Sigma-Aldrich) with CSM -MET (0.77% w/v, Formedium) amino acid mix and 2% dextrose (Sigma-Aldrich). For fluorescence microscopy of yLL129a (Figure 1B), we used low-fluorescence media, namely non-fluorescent nitrogen base (0.69% w/v, Formedium), CSM amino acid mix (0.79% w/v, Formedium) and 2% dextrose. For the growth rate assay in Figure 4B, we used YPD (10 g/L Yeast extract, 20 g/L peptone, 0.1/0.5/2% dextrose, 20 μg/ml adenine) and 0.25 mM auxin for strains yLIC133 and yWT3.

Method details

Extended explanation cell cycle model

Size expansion

From (Goranov et al., 2009), two growth phases are distinguished, with seemingly linear growth of volume with respect to time for the mother and sublinear volume growth for the bud, possibly consistent with constant growth of membrane area across time. Two phases are similarly visible in (Ferrezuelo et al., 2012), but also indicating sublinear dependency between volume and time. In light of absence of consensus, we accommodated both the constant area and constant volume expansion across time implementation in our model, but used the former by default. However, for the description of GAP epistasis this choice is inconsequential, see Figure S2. We follow the

542   observation in (Ferrezuelo et al., 2012) that mother expansion stops during bud growth. We hence

543   describe the size expansion as:

$$
\begin{cases}
\dfrac{dr_m}{dt} = \dfrac{C_1}{r_m}, r_b = 0 & G1\ phase \\[2mm]
\dfrac{dr_m}{dt} = 0, \dfrac{dr_b}{dt} = \dfrac{C_2}{r_b} & S/G2/M\ phase
\end{cases}
\tag{1}
$$

where r_m and r_b are mother and bud radius respectively, and C_1 and C_2 are constants.

To set C_1, we assume an optimized WT such that at checkpoint 1, the minimum size requirement

$r_m \geq r_{min}$ is typically met at the same time that the minimum G1 time requirement (time before

polarization starts must be $\geq t_{G1,min}$) is met, and that subsequent polarization time is minimal. After

G1, the (squared) mother radius is then (integrating equation 1 from 0 to $t_{pol,min}$, the duration needed

to polarize for a fast WT):

$$
\frac{dr_m}{dt} = \frac{C_1}{r_m} \Longrightarrow r_m^2 = r_{min}^2 + \frac{C_1 t_{pol,min}}{2}
\tag{2}
$$

551   From (Ferrezuelo et al., 2012; Goranov et al., 2009), we infer final bud volumes to be around 70%

552   of mother volumes. This means the bud radius after the next M-phase $r_b = 0.7^{1/3}\ r_m \approx 0.89\ r_m$. For

553   self-consistency, this new cell must then expand to size r_{min} again at checkpoint 1, such that:

554
$$
r_{min}^2 = r_b^2 + \frac{C_1 t_{G1,min}}{2} = 0.79 r_{min}^2 + 0.79 \frac{C_1 t_{pol,min}}{2} + \frac{C_1 t_{G1,min}}{2}
$$

$$
\Longrightarrow C_1 = \frac{0.42 r_{min}^2}{0.79\ t_{pol,min} + t_{G1,min}}
\tag{3}
$$

From Table S6 of (Ferrezuelo et al., 2012), we see the volume at Start in glucose of WT is 36.1

µm³, which assuming a spherical cell implies $r_{min} \leq 2.05$ µm, which we round to 2 µm. From Table

S7 of the same paper, we see Time at Start for mother WT cells to be 15.6 min., so we set $t_{G1,min}$ to

this value. Note it is this value that alters when we incorporate the *nrp1* deletion. Finally, we

estimate $t_{pol,min}$ from (Howell et al., 2012), where it is noted that polarization of Bem1p typically

occurs in 2 minutes starting from the point of first detection of significant fluorescence. To estimate

the full minimal polarization time including the time when Bem1p accumulation is still too faint,

we turned to the power spectrum of the oscillation time in Figure 1F of that paper, which peaks

around 0.2 1/min., corresponding to a time to build a spot of 5 min., so we set $t_{pol,min}$ to this value.

From these parameter values, we calculate C_1=0.086 µm²/min from equation 3. This value is

consistent with the aforementioned papers of (Ferrezuelo et al., 2012; Goranov et al., 2009). For

example, reading out from the plot in Fig. 7C of (Goranov et al., 2009) we see for the mother

growth phase a shift from 1500 voxels to 1700 voxels in -30 to 0 min. 1 voxel is $0.26 \ \mu m^3$, so

$$\frac{200 \ voxels}{30 \ min.} = 4\pi r \ C_1 \implies C_1 = \frac{0.14}{r \ [\mu m]} \ [\mu m^2/min] \tag{4}$$

With a typical size of $2 \ \mu m$, we have $C_1 \approx 0.07$ µm²/min. For (Ferrezuelo et al., 2012), we see

volume growth of 0.15 µm³/min, and given $\frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt} = \frac{r}{2}\frac{dA}{dt}$, we expect $C_1 \approx 0.15$ µm²/min .

Therefore, our value of C_1=0.086 µm²/min is within the expected range.

The value of C_2 (equation 1) follows from calibration, by running the model implementation (see

further on this Materials and Methods) and varying C_2 such that under WT parameters, the doubling

time is close to 83 minutes, as expected from literature (Laan et al., 2015). This calibration yields

C_2=2.1324 µm²/min. Again, this is in line with literature, as the rate during bud growth phase is

roughly twice as fast as the isotropic growth phase in (Ferrezuelo et al., 2012; Goranov et al.,

2009). To accommodate different growth media into the calculation as for the growth rates of

Figure 4A, both membrane growth rate can be multiplied by the same factor. For Figure 4A, this

factor was between 0.3 and 1 (steps of 0.1).

We also set a maximum size value, r_{max}, which plays a role when Cdc42 turnover has been

unfavorable to prevent polarization. The value is set at 6 µm, which prevents excessive calculations

on cells with very poor forecasts, and accommodates the smallest and thus healthiest >90% of the

cells of the ill Δ*bem1* background, assuming a normal distribution for the cell diameter with mean

6 µm and standard deviation 4.3 µm, with values estimated from Figures 1E and 1F of (Laan et al.,

2015). Setting the maximum radius to i.e. 20 µm has a negligible influence (<0.25%) on simulated

doubling times of WT and Δ*bem1* cells.

Cdc42p turnover

We expect Cdc42p expression bursts at intervals and with sizes which are exponentially distributed

(Friedman et al., 2006). In this paper, it was also explained how the average burst interval time and

size can be inferred from flow cytometry data. The former, $t_{b,WT}$, follows from dividing the average

cycle time by the shape parameter of a gamma fit on the endogenous Cdc42p distribution across

the population. The latter, $p_{b,WT}$, in principle follows from dividing the average Cdc42p copy

number which can be estimated from (Kulak et al., 2014) to be 8700, by the same shape parameter

of the aforementioned gamma fit. In this way, the mean of the gamma distribution should equal

8700. However, as we explicitly wish to include degradation into our model to better mimic the

critical part of the turnover concerning passage of the mesotype threshold, we must adjust the

scaling parameter slightly. Following calibration as in the previous section, we adjusted $p_{b,WT}$ (see

Table S1) such that the average copy number of Cdc42p in WT is close to the 8700 from (Kulak et

al., 2014). This required an 18,67% reduction of the original $p_{b,WT}$.

The flow cytometry for setting $t_{b,WT}$ and $p_{b,WT}$ was performed using strains RWS116 and RWS1421,

from (Freisinger et al., 2013; Gulli et al., 2000)). RWS1421 contains C*DC42pr-GFP-CDC42* in

addition to its endogenous *CDC42*, as this GFP tagged Cdc42p is not fully functional. However,

we do not expect this tag to be of major influence on the inferred copy number from fluorescence,

as the main issue is mislocalization of this Cdc42-variant (Watson et al., 2014). RWS116 has the

same strain background as RWS1421 but without fluorescence, to serve as a background

measurement. Fluorescence measurements were acquired using FlowJo CE software and

performed on a BD FACScan flow cytometer. Cells were pregrown in YNB (Sigma) + CSM -Met

(Formedium) + 2% dextrose (Sigma-Aldrich), diluted to an OD_{600} of 0.1 and measured after 15h.

The flow cytometry data was then analyzed (for the script see SI experimental code) by manually

imposing a polygon in the Forward/Side scattering plot to remove data points that likely correspond

to debris or dead cells (gating). Using Matlab's R2016a built-in *mle* function, we fit a gamma

distribution using maximum likelihood, to the background and endogenous Cdc42p distribution.

This yields the shape parameters $k = 2.36 \pm 0.05$ and 4.74 ± 0.07 and the scale parameters $\theta = 20.2$

± 0.5 and 59.6 ± 0.9 for the background and the endogenous distribution respectively, where the $\pm$

sign reports the 95% confidence interval.

We then approximate the analytical deconvolution of the endogenous distribution by setting the

deconvolved distribution as a gamma distribution (Stewart et al., 2007), where parameters follow

from subtracting the mean and variance of the background from the mean and variance respectively

of the endogenous signal. In that paper, the gamma approximation for deconvolution is shown to

hold as long as the scaling parameters of the background and endogenous signal are within an order

of magnitude, as is the case here. We retrieve for the background corrected endogenous signal the

shape parameter $k = 3.48 \pm 0.07$ and the scale parameter $\theta = 67.6 \pm 0.9$. Here the 95% confidence

interval indicated by the $\pm$ sign follows from 2.5% and 97.5% quantiles of Monte Carlo simulations

(10000 runs) of calculations of the shape and scaling parameters assuming normal errors on the

background and uncorrected endogenous parameters, also assuming independence between a draw

of a shape parameter and a draw of a scaling parameter.

The growth rate assay for RWS1421 was performed in the same media type as the flow cytometry,

in an Infinite M-200 Pro Tecan plate reader at 30°C.OD_{600} measurements (9 nm bandwidth) were

conducted with an interval of 12 minutes for a total duration of 24 hours, using a Thermo Fisher

Scientific Nunclon 96 Flat Bottom Plate input template. After an initial 1000s of orbital shaking (1

mm amplitude), linear shaking between measurements (25 flashes, 5 ms settle time) lasted 330s

each time (amplitude 1mm).

Only three wells (E2, E3, E4) were designated for RWS1421 in the appropriate medium. These

were analyzed using a home-written Matlab GUI already used in (Brauns et al., 2020) and made

available as Supplemental Information. The GUI allows the user to import the plate reader output

file in excel format, after which the user can specify a time bandwidth within which to perform an

ordinary (OLS) or weighted least squares (WLS) fit (OLS/WLS regressions described in (Heij et

al., 2004)). This fit follows from a linear regression on the log (base 2) of the (OD_{600} – background

OD_{600} value), where the background is set as the mean of the first ten OD_{600} values in that well.

The weights for WLS are set by the reciprocal of the difference between log of the OD_{600} values

with the instrument error added and subtracted. The instrument error is approximated by taking the

standard deviation of the first ten OD_{600} or by $10^{-digit}/2$ where digit is the value of the exponent with

base 10 of the last significant digit. The latter case applies when the data output from the plate

reader has rounded the data to an extent that the first ten point are identical. WLS is performed for

every fit window of a user-defined size that fits within the time bandwidth that the user also defines,

and that only contains data corresponding to the longest time span above a user defined signal-to-

noise ratio (with noise from the instrument error). For every fit window, the reciprocal of the

doubling time is the slope of the WLS regression, which follows a Student's t-distribution with the

number of data points in the fitting window minus 2 as the degrees of freedom (Heij et al., 2004).

This distribution sets the standard error. The fastest doubling time which pertains to a fit with a R^2

squared above a user defined value, is chosen to be the doubling time.

Since the flow cytometry measurement protocol suggests the protein distribution reflects more

accurately as the distribution is during late log phase (measuring 15h after dilution), we

intentionally fit above an OD above 0.1, roughly between 10 and 14h into the growth experiment.

We set our fitting window size to 21 points, weighted least squares option to on, minimum R^2 to

0.9 and mininmal signal-to-noise ratio to 2. The doubling times following the analysis in the GUI

were 129, 199 and 199 minutes. We take the median value and round to 200 minutes to use as the

cycle time for the calculation of $t_{b,WT}$.

Table S4 contains the flow cytometry (raw, processed and fitted) of RWS116 and RWS1421 used

in this study, see also Figure S3. Table S5 contains OD_{600} growth assay measurements of RWS1421

(with fits).

Polarity mesotype

The rigorous analysis of the reaction-diffusion equations and numerical simulations done in

(Brauns et al., 2020) provided the support to simplify the a priori complex dependency of

polarization success on polarity protein concentrations and reaction rates to a simple cone-like

phase space (see Figure 1D). More concretely, this also resulted in the support to justify a simple

approximation for the polarization time, see the qualitative plots in Figures 23 and 24 in (Daalman,

2020) (courtesy of Fridtjof Brauns). We approximate the polarization time t_{pol} as an exponential:

682
$$t_{pol} = t_{pol,min} e^{\frac{[Cdc42]-[Cdc42]_{min}}{[Cdc42]_{min}\phi_{pol}}}$$

where $[Cdc42p]_{min}$ is the polarity mesotype and φ_{pol} is a scaling parameter to govern the

theoretically predicted exponential increase in polarization time when there is excessive Cdc42p.

As the aforementioned qualitative plots indicate similar behavior for low GAPs concentrations

(relative to Cdc42p), we assume $[Cdc42p]_{min}$ φ_{pol} for both the *BEM1* and Δ*bem1* background. As

no clear deleterious effects are apparent for high induction of Cdc42p under the Gal1 promoter in

(Brauns et al., 2020), we set

ϕ_{pol} at a sufficiently high value of 25 for the Δ*bem1* background. For the *BEM1* background, we

notice that setting a value of 500 yields the approximately self-consistent result $[Cdc42p]_{min,\Delta bem1}$

$\varphi_{pol\ \Delta bem1} = 116 \cdot 25 \approx [Cdc42p]_{min,BEM1}\ \varphi_{pol\ BEM1} = 4 \cdot 500$.

Because of the shape of the phase space, the mesotypes are simple linear functions of GAP

concentrations, such that we only require to fit the polarity mesotype for the WT and Δ*bem1*

background and fit relative shift in mesotype for a *bem3* deletion and for the *bem2* deletion (the

same in both backgrounds). The relative shift in mesotype with both GAPs deleted is then simply

the sum of the two relative shifts, and is not modelled to alter upon *nrp1* deletion. Thus, four

mesotypes (and a change in $t_{G1,min}$ for the *nrp1* deletion) describe all polarity mutants from (Laan

et al., 2015) considered in this study.

As a final note, we impose for practical purposes a maximum polarization time of 600 min., to

truncate excessive long polarization times. Extending this time to 2000 min. has no influence on

doubling times of WT and Δ*bem1* cells.

G1 timing

We also experimented with a more realistic (and less coarse-grained) modification of the modelled

cell cycle progression to improve the quantitative match of the *BEM1-NRP1* epistasis in Figure

3A. Suppose for example the *Δbem1* cells if the assumed minimum G1 time set is not a constant

but results from a distribution. This is more realistic, as times for symmetry breaking in daughter

cells can exhibit notable stochasticity (Moran et al., 2019). Some Δ*bem1* cells have an early

opportunity to fulfil the mesotype threshold concentration requirement, with which they usually

struggle, while others are delayed more. This increases the cell-to-cell variation in fates in G1,

since cells with fast G1 times are most likely to generate a first spot, while slow cells never generate

this spot and thus do not show up in the statistics. This is how less coarse-graining can lead to a

larger decrease in G1 times from WT to Δ*bem1* than is the case with constant G1 waiting time

$t_{G1,min}$. Implementing a minimum G1 time distribution yields a slight quantitative improvement for

the G1 times in the genotype trajectory of Figure 3D (see Table S2). Thus, other unmodelled parts

of the cell cycle are likely responsible for the remaining quantitative discrepancy in describing G1

times.

Fitting parameters cell cycle model

Doubling times of (Laan et al., 2015) in Figure 3A were fitted using the built-in *fminsearch* on a

normalized score objective for varying [Cdc42]$_{min}$ and manual inspection for setting t_{mut} (to 0.75)

for the *nrp1* deletion. The objective involves the sum of squares of the fitting errors divided by the

standard errors of the experimental literature values. The experimental error for the Δ*bem2* Δ*nrp1*

is not available, and conservatively set at 30 minutes.

Computational implementation cell cycle model

Overview algorithm

Cell cycle model simulations were performed in MATLAB R2016a following a partial leap-

like Gillespie algorithm (Gillespie, 2001) implementation (the G1 time until $r<r_{min}$ and $t_1<t_{G1,min}$,

t_{pol} and the time through S/G2/M are one leap each). Cell cycle model parameters are summarized

in Table S1. The core function Mesotype_model.m, function to generate the data for the figures

and an example script to demonstrate the functionality are found as Supplemental Information.

Information considering the input parameters for each function are included in the code, and can

be called from Matlab's command window by e.g., the command "help Mesotype_model". A

pseudocode representation of the core Mesotype_model.m function follows the descriptions in the

next paragraphs.

An initial population (ancestor seed) asynchronized across a bandwidth of 83 minutes (all

cells with equal radii of 2.2 µm and without proteins) is defined. For every iteration in the simulated

growth loop, the lowest 50% of the population in terms of simulation time since the ancestor seed

is considered for growth, to avoid excessive asynchrony for the cells. Each of those cells aim to

progress through the cell cycle, where the step to fulfill all polarity conditions ($r < r_{min}$ and $t_1 < t_{G1,min}$

and $[Cdc42p] > [Cdc42p]_{min}$) is the only step where permanent cell death can take place. The 50%

of the cells not considered in the iteration are temporary scored as death, and this is reset in every

iteration.

The population grows until a user-defined population size placed as an input argument in

the core function, after which the colony is diluted to 1000 cells. Throughout this study, we use a

population size value of 5 million except for the single cell time trace where this was set at 2e3 to

avoid diluting too many of the initial cells. After dilution, the population grows again until the same

user-defined population size, or the highest user-defined time for copy number and volume

distribution probes if this requires the simulation to continue. All cells left alive must have a

simulated time since the ancestor seed to consider a copy number or volume probe complete.

If the colony fails to complete the simulation due to all cells dying prematurely, the

simulation is restarted. If the colony fails five times, the simulation returns a default doubling time

of 1e9 minutes. Otherwise, doubling times are the average of the last hundred moving window

(size 201 min.) linear regressions on the log number of cells.

Pseudocode representation of our physical cell cycle model

% Initialization

num_cells = 1000, radius=2.2, copy_number =0, simulation_time = -83 to 0

num_tries = 0

while num_cells < 5e6 & num_tries < 5

for all alive cells

769   **% Isotropic growth until $r = r_{min}$ and at least time_step $> t_{G1,min}$**

770   time_step = max($4\pi(r_{min}^2 - r^2) / C_1$, $t_{G1,min}$)

771   num_bursts ~ poiss(time_step / $t_{b,WT}$) % draw form poisson distribution

772   distribute burst_times uniformly across time step

773   for all bursts

774   burst_size ~ exp($p_{b,WT}$) % draw form exponential distribution

775   burst_size = burst_size * exp(burst_time / τ_h) % correction for dilution

776   mother copy number = mother copy number * exp(time_step / τ_h)

777   % correction for dilution

778   mother copy number = mother copy number + sum of all burst sizes

779   mother radius = mother radius + C_1 time_step % update mother radius

781   **% Isotropic growth until mother copy number $> P_{min}$ or $r > r_{max}$**

782   while mother copy number $< P_{min}$ and $r < r_{max}$

783   time_step ~ exp($t_{b,WT}$) % draw form exponential distribution

784   burst_size = ~ exp($p_{b,WT}$) % draw form exponential distribution

785   mother copy number = mother copy number * exp(time_step / τ_h)

786   % correction for dilution

787   mother copy number = mother copy number + burst_size

mother radius = mother radius + C_1 time_step % update mother radius

end

for all cells with **mother copy number** $< P_{min}$ and $r > r_{max}$

mark cells as dead

for all alive cells

% Isotropic growth until polarization is achieved

time_step   = $t_{pol,min}$ exp((mother copy number - P_{min}))/(P_{min} ϕ_{pol}))

num_bursts ~ poiss(time_step / $t_{b,WT}$) % draw form poisson distribution

distribute burst_times uniformly across time step

for all bursts

800   burst_size ~ exp($p_{b,WT}$) % draw form exponential distribution

801   burst_size = burst_size * exp(burst_time / τ_h) % correction for dilution

mother copy number = mother copy number * exp(time_step / τ_h)

% correction for dilution

mother copy number = mother copy number + sum of all burst sizes

mother radius = mother radius + C_1 time_step % update mother radius

update mother simulation_time

% Polarized growth of bud

time_step   = 4π ($(0.7)^{1/3}$ mother radius)^2 / C_2

num_bursts ~ poiss(time_step / $t_{b,WT}$) % draw form poisson distribution

distribute burst_times uniformly across time step

for all bursts

813   burst_size ~ exp($p_{b,WT}$) % draw form exponential distribution

814   burst_size = burst_size * exp(burst_time / τ_h) % correction for dilution

mother copy number = mother copy number * exp(time_step / τ_h)

% correction for dilution

mother copy number = mother copy number + sum of all burst sizes

bud radius = bud radius + C_2 * time_step   % update bud radius

bud copy_number = copy_number * bud volume / total volume % update bud

copy_number

Define buds as new cells

if no cells left

num_tries = num_tries + 1

% Re-initialize

num_cells = 1000, radius=2.2, copy_number =0, simulation_time = -83 to 0

continue with next iteration while-loop

if num_cells > 5e6 & dilution did not occur yet

Dilute: Randomly select 1000 cells and restart while loop

end

Note: checks for simulated measurements of volumes and copy numbers occur in each time_step.

Required intermediate radii follow deterministically and are stored. Intermediate copy numbers

follow from uniformly distributing the bursts across the time_step and store the intermediate copy

number update of the bursts that fall in the time until the simulated measurment.

Growth assay varying medium quality

Strains for the growth assay of Figure 4B were inoculated in YP2%D + 20 µg/ml adenine from

glycerol stock in glass tubes. These were then grown in a turning wheel for at least 24h at 30 °C.

After this pre-growth the cultures were diluted (at least 100x) to an OD_{600} of approx. 0.05 into a

96-well plate. Each well contains 100 µL of one of the three medium types with differing glucose

concentrations (YP + 0.1%, 0.5%, and 2% dextrose). Auxin was added to a final concentration of

0.25mM this medium to induce the Bem1p degradation. 0.25mM auxin was also added to the wells

with WT yeast as a control. Then the OD_{600} was measured for 48h using a Biotek Epoch™ 2

Microplate Spectrophotometer using linear and orbital shaking at 30C, as described in (Brauns et

al., 2020)

Predictions using literature data

Interaction and phenotype data for Figure 4D were obtained from BioGRID (Stark, 2006) and SGD

(Cherry et al., 2012) respectively (date of access March 2018), see also Table S6 for detailed

references   per   interaction.   Concretely,   interactions   with   Bem1   are   from

https://thebiogrid.org/32897/summary/saccharomyces-cerevisiae/bem1.html. Data for competitive

fitness   mutants   are   from   the   SGD   Project,

https://www.yeastgenome.org/observable/APO:0000110 [6 March 2018]. Data for fermentative

growth   mutants   are   from   the   SGD   Project.

859 https://www.yeastgenome.org/observable/APO:0000308 [6 March 2018]. Data for critical cell size

860 at START (G1 cell-size checkpoint) mutants are from the SGD Project.

861 https://www.yeastgenome.org/observable/APO:0000141 [8 March 2018]. Data for cell cycle

862 progression in G1 phase mutants are from the SGD Project.

863 https://www.yeastgenome.org/observable/APO:0000255 [8 March 2018]. Data for genes

864 annotated for "proteasome" are from the SGD Project.

865 http://yeastgenome.org/search?category=locus&geneMode=wrap&page=0&q=proteasome [8

866 March 2018]. Data for genes annotated for "ribosomal" are from the SGD Project.

867 http://yeastgenome.org/search?category=locus&geneMode=wrap&page=0&q=ribosomal [8

868 March 2018]. Data for Genes annotated for "phospholipid" are from the SGD Project.

869 http://yeastgenome.org/search?category=locus&geneMode=wrap&page=0&q=phospholipid [8

870 March 2018].

872 We consider all 'Positive Genetic' and 'Synthetic Rescue' annotations as positive interactions, and

873 'Negative Genetic' and 'Synthetic Lethality' as negative interactions. For the competitive fitness

874 mutants, we only considered those with phenotype competitive fitness: decreased (negative

875 interactions) or increased (positive interactions) in YPD. For the fermentative fitness mutants, we

876 only considered those with phenotype competitive fitness: decreased rate (negative interactions) or

877 increased rate (positive interactions) in YPD. For the mutants with altered critical cell size at

878 START (G1 cell-size checkpoint), we only considered the null, reduced and overexpression

879 mutants. For the overexpression mutants, we assumed the null/reduced mutants have the opposite

880 effect (e.g., decreased size to increased size). This was done similarly for the cell cycle progression

881 in G1 phase mutants, where we considered the same mutant types but for the phenotypes cell cycle

882 progression in G1 phase: delayed/increased duration/increased rate/decreased duration. When

interactions in all previously mentioned categories led to a conflict between effects (sources have

opposing opinions), we either choose the effect that is most commonly reported or discard the gene

if there is no single most commonly reported effect. We overwrite the genetic interaction with

Bem2 as positive (Laan et al., 2015).

We assume for simplicity that all genes annotated for "proteasome" have a negative effect on

degradation when mutated. We assume for simplicity that all genes annotated for "ribosomal" have

a negative effect on translation and hence protein production when mutated. We assume for

simplicity that all genes annotated for "phospholipid" have a negative effect on phospholipid

synthesis and hence membrane area growth when mutated. Similar to the fitness mutants in the

coarse phenotype mutant set, membrane area growth rate mutations are incorporated in C_2 alone,

as this modification is more effective in changing fitness for the WT background than changing

both C_1 and C_2.

Strain construction

Firstly, the *ade2* deletion was performed through transformation by homologous recombination.

The repair fragment was generated by overlap extension PCR using primers olic24, olic15, olic18

and olic26, and encompasses the *URA3* (from a template derived from the pRL368 plasmid

(Wedlich-Soldner, 2004)) gene with homology regions with *ADE2* (from a yLL3a genomic

template). Selection on uracil dropout media completed the transformation of yLL3a to yLIC132.

Similarly, overlap extension PCR using primers olic24, olic20, olic26 and olic21 yielded the repair

fragment to remove the URA3 marker by transformation, yielding yLIC133. This strain then

formed the basis for yWT3 used in this study, which was generated as follows.

Firstly, the PCR product to integrate *osTIR1* in the HO locus was amplified from plasmid

pOsTir1w/oGFP. Plasmid pOsTIR1w/oGFP was a gift from Matthias Heinemann (Addgene

plasmid # 102883 ; http://n2t.net/addgene:102883 ; RRID:Addgene_102883. Secondly, using

Gibson assembly. a *BEM1* homology region was added to a pG23A plasmid backbone, upstream

and in frame of the *mCherry-AID* sequence. Plasmid pG23A was a gift from Matthias Heinemann

(Addgene plasmid # 102884 ; http://n2t.net/addgene:102884 ; RRID:Addgene_102884). Further

downstream, the *HPHMX6* cassette and another *BEM1* homology region was added, to allow

generation of a PCR fragment to integrate the *mCherry-AID* C-terminally on *BEM1* by selecting

with Hygromycin at transformation (for primers see Table S8). Starting from strain yLIC133,

successive transformation with the PCR fragment from the *osTIR1-KANMX4* plasmid (selecting

with G418) and the *BEM1-mCherryAID-HPHMX6* plasmid (selection with hygromycin) generates

strain yWT3 (deposition of latter plasmid at Addgene in progress).

Quantification and statistical analysis

Pooling and testing growth data

Three experiments with two plates each were done for the glucose sensitivity assay. In each plate

we used one and two biological replicates for the *BEM1* and *BEM1-AID* background respectively,

with at least two technical replicates per medium. The doubling time of each well was calculated

using the Matlab GUI as explained above (section Cdc42p turnover). Per run, we averaged the

fitness (defined by the reciprocal of the doubling time) of *BEM1* replicates. We then divided the

fitness of each *BEM1-AID* strain by the mean *BEM1* fitness of that run. We then pooled all *BEM1-*

AID replicates per medium for data analysis. Wells where no growth were observed were excluded.

We neglected the error from the doubling time fits as the largest error source is the variation across

replicates. We used a one-sided Welch t-test (Matlab R2016a's native ttest2, for unequal variances)

to test for significant differences, and applied a Holm-Bonferroni correction factor (Holm, 1979)

to the significance values for the three t-tests performed.

We disregarded the 0.25% and 1% medium type, to only plot the media containing the most runs.

We excluded the other *BEM1* background strain measured in some plates as it lacked the *ade2*

mutation present in the *BEM1-AID* strains. Finally, we excluded the biological replicate yWT3a,

as its behavior was less comparable to the other two *BEM1-AID* biological replicates. The latter

two exclusions do not affect the conclusions on which fitness differences were significant.

Epistasis predictions with Δ*bem1*

From the interaction sets described in the previous section, we calculated posterior distributions

for the probability θ of a positive interaction with *BEM1*, using the frequency of such an interaction

encountered in literature. We describe the probability of finding a positive interaction given a total

number of interactions as following from a binomial likelihood. If we then assume a beta

distribution beta(α,β) as our prior (more specifically, the uninformed prior with beta(1,1)), the

resulting posterior distribution for the probability of having a positive interaction θ is also a beta

distribution, but with parameters $\alpha+k$ and $\beta+N$, where k is the encountered positive interactions and

N the number of encountered negative interactions with *BEM1* (Hobbs and Hooten, 2015). The

positive and negative interaction counts for this study can be found in Table S7. This gives us the

distributions for θ for the pooled deleterious fitness mutants (competitive and fermentative), pooled

beneficial fitness mutants, mutants that are small at G1 or fast through G1, mutants that are large

in G1 or slow through G1, mutants that have less degradation or phospholipid production, and

finally mutants that have reduced protein production.

From 10000 random draws of every θ, we can simulate the distributions of the differences between

pairs of θ. In particular, we consider the difference distribution of θ between deleterious and

beneficial mutants, the difference distribution of the small/fast in G1 mutants and large/slow in G1

mutants, and the difference distribution of the low degradation/lipid and low production mutants.

The probability of a difference above zero corresponds to the probability that positive epistasis is

more prevalent in deleterious than in beneficial mutants, more prevalent in small/fast in G1 mutants

than in large/slow mutants, and more prevalent in low degradation/lipid than in low production

mutants. This probability is plotted in Figure 4D. Alternative, we can rewrite this probability as a

Bayes factor for the model hypotheses that the positive epistasis is indeed more prevalent in each

of the three aforementioned cases. The Bayes factors are these probabilities divided by 1 minus

these probabilities, and we turn to (Kass and Raftery, 1995) to interpret these factors.

Materials and strains table

Materials used	
Indole-3-acetic-acid	Merck Millipore
Adenine	Sigma-Aldrich
Yeast extract	Sigma-Aldrich
Peptone	Sigma-Aldrich
Dextrose	Sigma-Aldrich
CSM amino acid mix	Formedium

CSM -Methionine amino acid mix	Formedium
Yeast nitrogen base	Sigma-Aldrich
Yeast nitrogen base low fluorescence	Formedium
G418 sulphate	Thermo Fisher
Hygromycin B	Formedium
Strains used	
S. cerevisiae: Strain RWS116. Strain background: W303. MATa, *his3-11,15, ade2-1, can1-100, ura3, cln1::HisG, Δcln2, cln3::HisG, YipLac204-MET-CLN2::TRP1*	(Gulli et al., 2000)
S. cerevisiae: Strain RWS1421. Strain background: W303. MATa, *his3-11,15, ade2-1, can1-100, ura3, cln1::HisG, Δcln2, cln3::HisG, YipLac204-MET-CLN2::TRP1, CDC42p-GFP-CDC42::URA3*	(Freisinger et al., 2013)
S. cerevisiae: Strain yLL3a. Strain background: W303. *MATα can1-100, leu2-3, 112, his3-11,15, ura3Δ, BUD4-S288C*	(Laan et al., 2015)
S. cerevisiae: Strain yLL129a. Strain background: W303. *MATa can1-100::P_{MFA}-HIS3, leu2-3, 112, his3-11,15, ura3Δ, bem1Δ::KANMX6, bem3Δ::NATMX4, BUD4-S288C, SPA2-yeCitrine-URA3*	(Laan et al., 2015)

S. cerevisiae: Strain yLIC132. Strain background: W303. *MATα, can1-100, leu2-3, 112, his3-11,15, ura3Δ, ade2Δ::URA3, BUD4-S288C*	This work
S. cerevisiae: Strain yLIC133. Strain background: W303. *MATα can1-100, leu2-3, 112, his3-11,15, ura3Δ, ade2Δ, BUD4-S288C*	This work
S. cerevisiae: Strain yWT3. Strain background: W303. *MATα can1-100, leu2-3, 112, his3-11,15, ura3Δ, ade2Δ, BEM1-mCherry-AID-HPHMX6, BUD4-S288C, HO::osTIR1-3xMyc-KANMX4*	This work
Plasmids used	
pOsTir1w/oGFP pllasmid	(Papagiannakis et al., 2017) Addgene plasmid #102883
Hygro-Bem1-mCherry-AID plasmid	Deposition at Addgene in progress
pG23A	(Papagiannakis et al., 2017) Addgene plasmid # 102884
pFA6-HPHMX6	(Hentges et al., 2005) Euroscarf plasmid 30438
Software used	
Matlab 2016a	www.mathworks.com

NIS Elements AR 4.3	www.microscope.healthcare.nikon.com/ products/software/nis-elements/nis-elements-advanced-research
ImageJ	(Schneider et al., 2012), https://imagej.nih.gov/ij

Supplementary information titles and legends

Figure S1 Convergence of simulated doubling times.

Figure S1 Legend

Doubling times as function of simulation time for a fast (WT), medium (Δ*bem1* Δ*bem2*), and slow

(Δ*bem1*) growing strain background. Doubling times and simulation times normalized to their

respective final value. The dilution step midway temporarily causes unreliable estimates and these

are omitted from display.

Figure S2 Cell cycle model details effect on relative and absolute quantities

Figure S2 Legend

(A) Calculated GAP activity given cell cycle model fits and abundancies from (Kulak et al., 2014),

(B) relative multiplicative epistasis (definition of (da Silva et al., 2010)) for the GAPs in the Δ*bem1*

NRP1 background and (C) fitted mesotype threshold (minimum Cdc42 concentration to polarize)

in the Δ*bem1* background for the default parameter set in the cell cycle model and four variations;

doubling of membrane area growth rate C_2, 15% reduction of r_{min}, change of second checkpoint to

a fixed bud size threshold of 1.8 µm, and constant cell volume instead of area expansion. WT

membrane growth rates are recalibrated in each case to match 83 minutes. Fits are performed on

the doubling times of *NRP1* strains in (Laan et al., 2015) alone, in slight contrast with fits in Figure

3A, to focus on variation due to GAP deletions. Quantities in panels (A) and (B) are relative,

dimensionless quantities, whereas the quantity in (C) is absolute.

Figure S3 Measured Cdc42p distributions

Figure S3 Legend

(A) Flow cytometry data (histogram) of RWS116, which acts as a measure of the background count

distribution without fluorescence. The solid green line shows the maximum likelihood gamma

distribution fit. (B) Flow cytometry data (histogram) of RWS1421, which shows the WT

distribution of Cdc42p. The solid green line shows the maximum likelihood gamma distribution

fit, the dashed green line the fit corrected for the background fit in panel (A). Underlying data is

found in Table S4.

1004

1005 **Table titles and legends**

1006 **Table S1. Growth model parameter list.**

1007 **Table S1 Legend**

1008 Parameters used in the physical cell cycle model in this study, with associated symbols, values and

1009 justification.

Parameter	Symbol	Value (background)	Source
Cdc42 concentration threshold (mesotype)	$[Cdc42]_{min}$	116 # proteins/μm^3 ($\Delta bem1$) 4 # proteins /μm^3 (*BEM1*) -61% ($\Delta bem3$) -35% ($\Delta bem2$)	This study, fitted
Minimum G1 time multiplier	t_{mut}	1 (*NRP1*) 0.75 ($\Delta nrp1$)	This study, fitted
Minimum polarization time	$t_{pol,min}$	5 min.	(Howell et al., 2012)
Maximum polarization time	$t_{pol,max}$	600 min.	To truncate computations for cells with extremely low GAP content

Polarization time scaling parameter	ϕ_{pol}	25 ($\Delta bem1$) 500 (*BEM1*)	$\phi_{pol,BEM1} \gg 1$ and $\phi_{pol,BEM1}/\phi_{pol,\Delta bem1} \cong$ $[Cdc42]_{min,\Delta bem1}$ $/[Cdc42]_{min,BEM1}$ (see Materials and Methods)
Minimum G1 time	$t_{G1,min}$	15.6 min.	(Ferrezuelo et al., 2012)
Minimum radius to polarize	r_{min}	2 µm	(Ferrezuelo et al., 2012)
Maximum radius in G1	r_{max}	6 µm	To truncate computations for cells with very low Cdc42 content
Average Cdc42 expression burst interval time	$t_{b,WT}$	57 min.	This study, assuming theory from (Friedman et al., 2006)
Average Cdc42 expression burst size	$p_{b,WT}$	4900	This study, assuming theory from (Friedman et al., 2006) and with calibration
Cdc42p half-life	τ_h	474 min.	(Christiano et al., 2014)
Bud/mother volume ratio checkpoint 2	c_r	0.89	Consistent with (Ferrezuelo et al., 2012)
Ratio polarized/ isotropic membrane	c_p	2.13	Calibration

area growth rates (C_2/C_1)			
Isotropic membrane area growth rate	C_1	0.086 µm^2/min.	Analytical considerations for optimized WT (see S3 Text)

1010

Table S2. Additional G1 times shifts explained by stochastic minimum G1 time.

Table S2 Legend.

Percentage of the mean shift in observed G1 times in the adaptive trajectory of (Laan et al., 2015)

that is left unexplained by the standard cell cycle model implementation as in Figure 3D with

minimum G1 time $t_{mut} t_{G1,min}$, and is explained when the minimum G1 time is stochastic. In this

alternative implementation, the minimum G1 time is on average the same, but results from a gamma

distribution $\Gamma(2, t_{mut} t_{G1,min}/2)$.

Genotype transition	% mean G1 time shift mismatch described
WT→ $\Delta bem1$	5.8
$\Delta bem1 \rightarrow \Delta bem1\ \Delta bem3$	1.7
$\Delta bem1\ \Delta bem3 \rightarrow \Delta bem1\ \Delta bem3\ \Delta nrp1$	2.7
$\Delta bem1\ \Delta bem3\ \Delta nrp1\rightarrow$ $\Delta bem1\ \Delta bem3\ \Delta nrp1\ \Delta bem2$	5.2

1018

1019 **Table S3. Example calculation of quantitative epistasis of mutants with Δ*bem1*.**

1020 **Table S3 Legend**

1021 Epistasis values for interactions with Δ*bem1* assuming several mutations with certain impact on

1022 model parameters. Epistasis is defined as relative multiplicative epistasis as in (da Silva et al.,

1023 2010). Default doubling time values are resulting from the fit in Figure 3A.

Case	Parameter	Shift rel. to default [%]	Simulated doubling time [min.]	Simulated doubling time [min.]	Epistasis
Default	-	-	82	406	-
Fast membrane growth	C_2	10	77	460	-0.08
Slow membrane growth	C_2	-10	89	368	0.07
Small min. Start radius	r_{min}, $t_{G1,min}$	-15,-20	70	234	0.17
Large min. Start radius	r_{min}, $t_{G1,min}$	15,20	95	1018	-0.34
Slow in G1	$t_{G1,min}$	20	94	1016	-0.34
Fast in G1	$t_{G1,min}$	-20	71	237	0.17
Slow degradation	τ_h	40	82	324	0.10

Fast degradation	τ_h	-40	83	1038	-0.41
Small bursts	$p_{b,WT}$	-5	82	555	-0.14
Large bursts	$p_{b,WT}$	5	82	327	0.09

1024

1025 **Table S4. Flow cytometry data.**

1026 **Table S4 Legend**

1027 Flow cytometry data underlying Fig S3 of RWS116 and RWS1421. The raw data, gated data and

1028 fitted data are in separate tabs. The fit of RWS1421 corrected for the background distribution fit of

1029 RWS116 is also given in the fitted data tab. The data in Table S1 underlies Figure S3.

1030

1031 **Table S5. Growth assay data**

1032 **Table S5 Legend**

1033 Growth assay data of RWS1421 (OD_{600} values) with a weighted least squares fit on the log of the

1034 background corrected OD_{600} values, applied through our growth curve GUI.

1035

1036 **Table S6. Growth assay data**

1037 **Table S6 Legend**

1038 Interaction data with *BEM1* from (Stark, 2006) and mutant data associating genes with various

1039 phenotypes and gene functions from (Cherry et al., 2012), SGD Project.

1040 https://www.yeastgenome.org/ [6 and 8 March 2018].

1041

1042 **Table S7. Epistasis prevalence**

1043 **Table S7 Legend**

1044 Number of positive and negative interactions with *BEM1*, subdivided by mutant type.

	Number of negative interactors with BEM1	Number of positive interactors with BEM1
Beneficial mutants (competitive fitness)	21	5
Deleterious mutants (competitive fitness)	192	32
Beneficial mutants (fermentative growth)	0	0
Deleterious mutants (fermentative growth)	47	6
Slow through G1	22	5
Fast through G1	6	2
Small at Start	2	1
Large at Start	1	1
Proteasomal	5	2
Ribosomal	41	5
Phospholipid	5	2

1045

1046 **Table S8. Oligonucleotides**

1047 **Table S8 Legend**

1048 Oligonucleotides used in this study.

1049

1050 **Data file S1. Flow cytometry processing code**

1051 Matlab code to process flow cytometry data as found in Table S4.

1052

1053 **Data file S2. Yeast polarity GP map simulation code**

1054 Zip-file containing the Matlab codes for the generation of the model simulation data.

1055

1056 **Data file S3. Plate reader GUI**

1057 Zip-file containing the Matlab codes for analyzing growth assay data from excel sheets using a

1058 graphical user interface.

1059

1060 **Data file S4. Growth data effect of media quality**

1061 Zip-file containing the excel sheets with measured OD_{600} values of the growth assays underlying

1062 Figure 4B.

1063

References

Adams, A.E., Johnson, D.I., Longnecker, R.M., Sloat, B.F., and Pringle, J.R. (1990). CDC42 and
CDC43, two additional genes involved in budding and the establishment of cell polarity in the yeast
Saccharomyces cerevisiae. J. Cell Biol. *111*, 131–142.

Atolia, E., Cesar, S., Arjes, H.A., Rajendram, M., Shi, H., Knapp, B.D., Khare, S., Aranda-Díaz,
A., Lenski, R.E., and Huang, K.C. (2020). Environmental and physiological factors affecting high-
throughput measurements of bacterial growth. Mbio *11*.

Bergelson, J., Kreitman, M., Petrov, D.A., Sanchez, A., and Tikhonov, M. (2021). Functional
biology in its natural context: A search for emergent simplicity. eLife *10*.

Blanco-Gómez, A., Castillo-Lluva, S., del Mar Sáez-Freire, M., Hontecillas-Prieto, L., Mao, J.H.,
Castellanos-Martín, A., and Pérez-Losada, J. (2016). Missing heritability of complex diseases:
Enlightenment by genetic variants from intermediate phenotypes. BioEssays *38*, 664–673.

Brauns, F., de la Cruz, L.M.I., Daalman, W.K.-G., de Bruin, I., Halatek, J., Laan, L., and Frey, E.
(2020). Adaptability and evolution of the cell polarization machinery in budding yeast. bioRxiv.

Cherry, J.M., Hong, E.L., Amundsen, C., Balakrishnan, R., Binkley, G., Chan, E.T., Christie, K.R.,
Costanzo, M.C., Dwight, S.S., Engel, S.R., et al. (2012). Saccharomyces Genome Database: the
genomics resource of budding yeast. Nucleic Acids Res. *40*, D700–D705.

Chong, Y.T., Koh, J.L.Y., Friesen, H., Kaluarachchi Duffy, S., Cox, M.J., Moses, A., Moffat, J.,
Boone, C., and Andrews, B.J. (2015). Yeast Proteome Dynamics from Single Cell Imaging and
Automated Analysis. Cell *161*, 1413–1424.

Christiano, R., Nagaraj, N., Fröhlich, F., and Walther, T.C. (2014). Global Proteome Turnover
Analyses of the Yeasts S. cerevisiae and S. pombe. Cell Rep. *9*, 1959–1965.

Cobb, J.N., DeClerck, G., Greenberg, A., Clark, R., and McCouch, S. (2013). Next-generation
phenotyping: requirements and strategies for enhancing our understanding of genotype–phenotype
relationships and its relevance to crop improvement. Theor. Appl. Genet. *126*, 867–887.

Cordell, H.J. (2009). Detecting gene–gene interactions that underlie human diseases. Nat. Rev.
Genet. *10*, 392–404.

Daalman, W.K.-G. (2020). Coupling of genotype-phenotype maps to noise-driven adaptation,
showcased in yeast polarity. TU Delft.

Deutschbauer, A.M., Jaramillo, D.F., Proctor, M., Kumm, J., Hillenmeyer, M.E., Davis, R.W.,
Nislow, C., and Giaever, G. (2005). Mechanisms of Haploinsufficiency Revealed by Genome-
Wide Profiling in Yeast. Genetics *169*, 1915–1925.

Eichler, E.E., Flint, J., Gibson, G., Kong, A., Leal, S.M., Moore, J.H., and Nadeau, J.H. (2010).
Missing heritability and strategies for finding the underlying causes of complex disease. Nat. Rev.
Genet. *11*, 446–450.

Ferrezuelo, F., Colomina, N., Palmisano, A., Garí, E., Gallego, C., Csikász-Nagy, A., and Aldea,
M. (2012). The critical size is set at a single-cell level by growth rate to attain homeostasis and
adaptation. Nat. Commun. *3*, 1012.

Freisinger, T., Klünder, B., Johnson, J., Müller, N., Pichler, G., Beck, G., Costanzo, M., Boone,
C., Cerione, R.A., Frey, E., et al. (2013). Establishment of a robust single axis of cell polarity by
coupling multiple positive feedback loops. Nat. Commun. *4*, 1807.

Friedman, N., Cai, L., and Xie, X.S. (2006). Linking stochastic dynamics to population distribution: an analytical framework of gene expression. Phys. Rev. Lett. *97*, 168302.

Geßele, R., Halatek, J., Würthner, L., and Frey, E. (2020). Geometric cues stabilise long-axis polarisation of PAR protein patterns in C. elegans. Nat. Commun. *11*.

Giaever, G., Chu, A.M., Ni, L., Connelly, C., Riles, L., Véronneau, S., Dow, S., Lucau-Danila, A., Anderson, K., and Andre, B. (2002). Functional profiling of the Saccharomyces cerevisiae genome. Nature *418*, 387–391.

Gillespie, D.T. (2001). Approximate accelerated stochastic simulation of chemically reacting systems. J. Chem. Phys. *115*, 1716–1733.

Goldstein, B., and Macara, I.G. (2007). The PAR Proteins: Fundamental Players in Animal Cell Polarization. Dev. Cell *13*, 609–622.

Goranov, A.I., Cook, M., Ricicova, M., Ben-Ari, G., Gonzalez, C., Hansen, C., Tyers, M., and Amon, A. (2009). The rate of cell growth is governed by cell cycle stage. Genes Dev. *23*, 1408–1422.

Goryachev, A.B., and Pokhilko, A.V. (2008). Dynamics of Cdc42 network embodies a Turing-type mechanism of yeast cell polarity. FEBS Lett. *582*, 1437–1443.

Grigull, J., Mnaimneh, S., Pootoolal, J., Robinson, M.D., and Hughes, T.R. (2004). Genome-Wide Analysis of mRNA Stability Using Transcription Inhibitors and Microarrays Reveals Posttranscriptional Control of Ribosome Biogenesis Factors. Mol. Cell. Biol. *24*, 5534–5547.

Gulli, M.-P., Jaquenoud, M., Shimada, Y., Niederhäuser, G., Wiget, P., and Peter, M. (2000). Phosphorylation of the Cdc42 exchange factor Cdc24 by the PAK-like kinase Cla4 may regulate polarized growth in yeast. Mol. Cell *6*, 1155–1167.

Heij, C., de Boer, P., Franses, P.H., Kloek, T., and van Dijk, H.K. (2004). Econometric methods with applications in business and economics (Oxford University Press).

Hentges, P., Van Driessche, B., Tafforeau, L., Vandenhaute, J., and Carr, A.M. (2005). Three novel antibiotic marker cassettes for gene disruption and marker switching in Schizosaccharomyces pombe. Yeast *22*, 1013–1019.

Hobbs, N.T., and Hooten, M.B. (2015). Bayesian models: a statistical primer for ecologists (Princeton University Press).

Holm, S. (1979). A simple sequentially rejective multiple test procedure. Scand. J. Stat. 65–70.

Howell, A.S., Jin, M., Wu, C.-F., Zyla, T.R., Elston, T.C., and Lew, D.J. (2012). Negative Feedback Enhances Robustness in the Yeast Polarity Establishment Circuit. Cell *149*, 322–333.

Kass, R.E., and Raftery, A.E. (1995). Bayes Factors. J. Am. Stat. Assoc. *90*, 773–795.

Kayser, M., and Schneider, P.M. (2009). DNA-based prediction of human externally visible characteristics in forensics: Motivations, scientific challenges, and ethical considerations. Forensic Sci. Int. Genet. *3*, 154–161.

Keren, L., Hausser, J., Lotan-Pompan, M., Vainberg Slutskin, I., Alisar, H., Kaminski, S., Weinberger, A., Alon, U., Milo, R., and Segal, E. (2016). Massively Parallel Interrogation of the Effects of Gene Expression Levels on Fitness. Cell *166*, 1282–1294.e18.

Klünder, B., Freisinger, T., Wedlich-Söldner, R., and Frey, E. (2013). GDI-Mediated Cell Polarization in Yeast Provides Precise Spatial and Temporal Control of Cdc42 Signaling. PLoS Comput. Biol. *9*, e1003396.

Kozubowski, L., Saito, K., Johnson, J.M., Howell, A.S., Zyla, T.R., and Lew, D.J. (2008). Symmetry-Breaking Polarization Driven by a Cdc42p GEF-PAK Complex. Curr. Biol. *18*, 1719–1726.

Kryazhimskiy, S. (2021). Emergence and propagation of epistasis in metabolic networks. Elife *10*, e60200.

Kryazhimskiy, S., Rice, D.P., Jerison, E.R., and Desai, M.M. (2014). Global epistasis makes adaptation predictable despite sequence-level stochasticity. Science *344*, 1519–1522.

Kulak, N.A., Pichler, G., Paron, I., Nagaraj, N., and Mann, M. (2014). Minimal, encapsulated proteomic-sample processing applied to copy-number estimation in eukaryotic cells. Nat. Methods *11*, 319–324.

Laan, L., Koschwanez, J.H., and Murray, A.W. (2015). Evolutionary adaptation after crippling cell polarization follows reproducible trajectories. eLife *4*, e09638.

van Leeuwen, J., Pons, C., Mellor, J.C., Yamaguchi, T.N., Friesen, H., Koschwanez, J., U aj, M.M., Pechlaner, M., Takar, M., U aj, M., et al. (2016). Exploring genetic suppression interactions on a global scale. Science *354*, aag0839-aag0839.

Lehner, B. (2011). Molecular mechanisms of epistasis within and between genes. Trends Genet. *27*, 323–331.

Li, C. (2011). Personalized medicine–the promised land: are we there yet? Clin. Genet. *79*, 403–412.

Miton, C.M., and Tokuriki, N. (2016). How mutational epistasis impairs predictability in protein evolution and design. Protein Sci. *25*, 1260–1272.

Moran, K.D., Kang, H., Araujo, A.V., Zyla, T.R., Saito, K., Tsygankov, D., and Lew, D.J. (2019). Cell-cycle control of cell polarity in yeast. J. Cell Biol. *218*, 171–189.

Nishimura, K., Fukagawa, T., Takisawa, H., Kakimoto, T., and Kanemaki, M. (2009). An auxin-based degron system for the rapid depletion of proteins in nonplant cells. Nat. Methods *6*, 917–922.

Oud, B., Maris, A.J.A., Daran, J.-M., and Pronk, J.T. (2012). Genome-wide analytical approaches
for reverse metabolic engineering of industrially relevant phenotypes in yeast. FEMS Yeast Res.
12, 183–196.

Papagiannakis, A., Niebel, B., Wit, E.C., and Heinemann, M. (2017). Autonomous Metabolic
Oscillations Robustly Gate the Early and Late Cell Cycle. Mol. Cell *65*, 285–295.

Papp, B., Notebaart, R.A., and Pál, C. (2011). Systems-biology approaches for predicting genomic
evolution. Nat. Rev. Genet. *12*, 591–602.

Pelillo, M., Poli, I., Roli, A., Serra, R., Slanzi, D., and Villani, M. (2018). Artificial Life and
Evolutionary Computation (Springer).

Ramm, B., Heermann, T., and Schwille, P. (2019). The E. coli MinCDE system in the regulation
of protein patterns and gradients. Cell. Mol. Life Sci. *76*, 4245–4273.

Sánchez, B.J., Zhang, C., Nilsson, A., Lahtvee, P., Kerkhoven, E.J., and Nielsen, J. (2017).
Improving the phenotype predictions of a yeast genome-scale metabolic model by incorporating
enzymatic constraints. Mol. Syst. Biol. *13*, 935.

Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of
image analysis. Nat. Methods *9*, 671–675.

da Silva, J., Coetzer, M., Nedellec, R., Pastore, C., and Mosier, D.E. (2010). Fitness Epistasis and
Constraints on Adaptation in a Human Immunodeficiency Virus Type 1 Protein Region. Genetics
185, 293–303.

Smith, E.N., and Kruglyak, L. (2008). Gene–Environment Interaction in Yeast Gene Expression.
PLoS Biol. *6*, e83.

Smith, G.R., Givan, S.A., Cullen, P., and Sprague, G.F. (2002). GTPase-Activating Proteins for
Cdc42. Eukaryot. Cell *1*, 469–480.

Smith, N.J., Walt, S. van der, and Firing, E. (2015). Magma, inferno, plasma and viridis colormaps,
https://github.com/BIDS/colormap/blob/master/colormaps.py.

Stark, C. (2006). BioGRID: a general repository for interaction datasets. Nucleic Acids Res. *34*,
D535–D539.

Stevenson, B.J., Ferguson, B., De Virgilio, C., Bi, E., Pringle, J.R., Ammerer, G., and Sprague,
G.F. (1995). Mutation of RGA1, which encodes a putative GTPase-activating protein for the
polarity-establishment protein Cdc42p, activates the pheromone-response pathway in the yeast
Saccharomyces cerevisiae. Genes Dev. *9*, 2949–2963.

Stewart, T., Strijbosch, L.W.G., Moors, H., and Batenburg, P. van (2007). A simple approximation
to the convolution of gamma distributions.

Tiedje, C., Sakwa, I., Just, U., and Höfken, T. (2008). The rho gdi rdi1 regulates rho gtpases by
distinct mechanisms. Mol. Biol. Cell *19*, 2885–2896.

Vergés, E., Colomina, N., Garí, E., Gallego, C., and Aldea, M. (2007). Cyclin Cln3 Is Retained at
the ER and Released by the J Chaperone Ydj1 in Late G1 to Trigger Cell Cycle Entry. Mol. Cell
26, 649–662.

Wagner, G.P., and Zhang, J. (2011). The pleiotropic structure of the genotype–phenotype map: the
evolvability of complex organisms. Nat. Rev. Genet. *12*, 204–213.

Watson, L.J., Rossi, G., and Brennwald, P. (2014). Quantitative Analysis of Membrane Trafficking
in Regulation of Cdc42 Polarity. Traffic *15*, 1330–1343.

Wedlich-Soldner, R. (2004). Robust cell polarity is a dynamic state established by coupling
transport and GTPase signaling. J. Cell Biol. *166*, 889–900.

Witte, K., Strickland, D., and Glotzer, M. (2017). Cell cycle entry triggers a switch between two
modes of Cdc42 activation during yeast polarization. Elife *6*.

Woods, B., Kuo, C.-C., Wu, C.-F., Zyla, T.R., and Lew, D.J. (2015). Polarity establishment
requires localized activation of Cdc42. J. Cell Biol. *211*, 19–26.

Wünsche, A., Dinh, D.M., Satterwhite, R.S., Arenas, C.D., Stoebel, D.M., and Cooper, T.F. (2017).
Diminishing-returns epistasis decreases adaptability along an evolutionary trajectory. Nat. Ecol.
Evol. *1*, 1–6.

Yu, M.K., Kramer, M., Dutkowski, J., Srivas, R., Licon, K., Kreisberg, J.F., Ng, C.T., Krogan, N.,
Sharan, R., and Ideker, T. (2016). Translation of Genotype to Phenotype by a Hierarchy of Cell
Subsystems. Cell Syst. *2*, 77–88.

Zhao, C., Medeiros, T.X., Sové, R.J., Annex, B.H., and Popel, A.S. (2021). A data-driven
computational model enables integrative and mechanistic characterization of dynamic macrophage
polarization. Iscience *24*, 102112.

Zheng, Y., Cerione, R., and Bender, A. (1994). Control of the yeast bud-site assembly GTPase
Cdc42. Catalysis of guanine nucleotide exchange by Cdc24 and stimulation of GTPase activity by
Bem3. J. Biol. Chem. *269*, 2369–2372.

Zuk, O., Hechter, E., Sunyaev, S.R., and Lander, E.S. (2012). The mystery of missing heritability:
Genetic interactions create phantom heritability. Proc. Natl. Acad. Sci. *109*, 1193–1198.

1235