Bulk flow of cerebrospinal fluid observed in periarterial spaces is not an artifact of injection

Aditya Raghunandan[1,+], **Antonio Ladrón-de-Guevara**[2,+], **Jeffrey Tithof**[1,3], **Humberto Mestre**[2], **Maiken Nedergaard**[2,4], **John H. Thomas**[1], and **Douglas H. Kelley**[1,*]

[1]Department of Mechanical Engineering, University of Rochester, Rochester, NY 14620, USA
[2]Center for Translational Neuromedicine, University of Rochester Medical Center, Rochester, NY 14642, USA
[3]Department of Mechanical Engineering, University of Minnesota, Minneapolis, MN 55455, USA
[4]Center for Translational Neuromedicine, University of Copenhagen, Copenhagen N, Denmark, 2200
[*]d.h.kelley@rochester.edu
[+]these authors contributed equally to this work

ABSTRACT

The flow of cerebrospinal fluid (CSF) through perivascular spaces surrounding the vasculature is thought to be an important part of the brain's mechanism for clearing metabolic waste products, including those associated with disorders such as Alzheimer's disease. Experiments that track tracer particles injected into the cisterna magna of mouse brains have shown evidence of pulsatile CSF flow in perivascular spaces around pial (surface) arteries, with a bulk (average) flow in the same direction as the blood flow. Although measurements are consistent with the idea that this flow is driven primarily by arterial pulsations, the driving mechanism is still not completely understood, and several published articles have suggested that the bulk flow might be an artifact, driven by the injection itself. Here we address this hypothesis with new *in vivo* experiments in which the injection of suspended tracer particles into the cisterna magna is done with a dual-syringe system, with simultaneous injection and withdrawal of equal amounts of fluid. This method produces no net increase in CSF volume and no significant increase in intracranial pressure, and yet particle-tracking reveals flows in the pial periarterial spaces that are completely consistent with the flows observed in earlier experiments with single-syringe injection.

Introduction

Cerebrospinal fluid (CSF) flowing in perivascular spaces (PVS) – annular tunnels that surround the brain's vasculature — plays a crucial role in clearing metabolic waste products from the brain[1,2]. The failure to remove such waste products, including toxic protein species, has been implicated in the etiology of several neurological disorders, including Alzheimer's disease[1,3]. Recently, *in vivo* experiments that combine two-photon microscopy and flow visualization in live mice have used the motion of fluorescent microspheres injected into the cisterna magna (CM) to measure the flow of CSF through the periarterial spaces surrounding pial (surface) arteries. The results show pulsatile flow, in lock-step synchrony with the cardiac cycle and with an average (bulk) flow in the same direction as that of the arterial blood flow[4,5]. This flow is driven primarily by "perivascular pumping"[6,7] due to motions of the arterial walls that drive the movement of the fluid[5]. Earlier, the existence of such a bulk flow of CSF into the brain was questioned[8,9], and while arterial pulsation has long been considered as a possible driving mechanism[10–12], that notion remains controversial[13–15].

One issue is whether the injection of tracers into the CM might cause a pressure gradient that drives a flow in the surface PVS[8,14–19]. Injection of CSF tracers is known to raise the intracranial pressure (ICP) by 1 - 3 mm Hg[12,20]. Reports contend that this addition of fluid volume leads to non-natural increases in ICP[21,22] that drive fluid into low-resistance pathways such as surface periarterial spaces[4,23]. In other words, these studies suggest that the bulk flow observed in detail by Mestre et al.[5] is an artifact of the injection. Mestre et al. showed that the flows did not decay over time, as would be expected if they were injection artifacts, but given that this criticism has since arisen in several publications, we decided to test the hypothesis with additional *in vivo* experiments, essentially identical to the earlier experiments[5], but employing a new particle-injection method.

Figure 1. Schematic representation of the cisterna magna injection using (a) the single-syringe protocol for injection of $10\,\mu\text{L}$ at $2\,\mu\text{L/min}$ and (b) the dual-syringe protocol for simultaneous injection and withdrawal of $20\,\mu\text{L}$ at $2\,\mu\text{L/min}$.

The new injection protocol, illustrated in Figure 1b, employs a dual-syringe system to infuse the tracer particles. In this system, two cannulae connected to synchronized syringe pumps are inserted into the CM; one line injects fluid in which the tracer particles are suspended, while the other line simultaneously withdraws an identical amount of fluid at the same volumetric flow rate. Thus, no net volume of fluid is added to the intracranial compartment, and hence we expect no significant change in ICP. We use two-photon microscopy to visualize the motion of the fluorescent tracer particles and measure the flow in the surface periarterial spaces using particle tracking velocimetry. We also simultaneously measure changes to ICP while monitoring heart and respiration rates. We compare the flow characteristics measured under the new protocol with those measured previously using the traditional single-syringe injection[4,5] (depicted in Fig. 1a). (For this comparison, the data from Mestre et al.[5] analyzed here are from the control mice, not the hypertension mice.) Our new results are completely consistent with the previous results. With the new infusion protocol, the flow is again pulsatile in nature, in step with the cardiac cycle, with a net (bulk) flow in the direction of arterial blood flow. We find nearly identical mean flow speeds and other flow characteristics with the new infusion protocol. Our new experiments confirm that the flows we observed in periarterial spaces in our earlier experiments are natural, not artifacts of the tracer infusion, and provide additional statistical information about these flows.

Results

Changes in intracranial pressure

In a group of mice, we evaluated the effect of tracer infusion upon ICP. A 30-gauge needle was inserted stereotactically into the right lateral ventricle and connected to a pressure transducer to monitor ICP during CSF tracer injection into the CM, using both the single-injection (n = 6) and dual-syringe (n=5) protocols (Fig. 2). In agreement with prior studies using similar single-injection protocols[2, 12, 20], we found that the injection of 10 μL of CSF tracer into the CM at a rate of 2 μL/min resulted in a mild elevation of ICP ($\sim$ 2.5 mmHg) that relaxed rapidly upon cessation of the injection (Fig. 2). On the other hand, when ICP was measured during the dual-syringe infusion, we observed that the simultaneous injection of the tracer and withdrawal of CSF did not significantly alter ICP (Fig. 2), as to be expected given the absence of any net change in the volume of fluid in the intracranial CSF compartment. Based upon these findings, we conducted intracisternal infusion of fluorescent microspheres into the CM using the dual-syringe protocol to perform particle-tracking studies and determine the characteristics of CSF flow in the absence of any transient elevation of ICP caused by the infusion protocol.

Figure 2. The effect of single-injection and dual-syringe tracer infusion upon intracranial pressure (ICP). The ICP was monitored continuously during injection of CSF tracers into the CM of mice. Injection begins at 60 s, indicated by the vertical dashed line. Single-injection infusion of 10 μL at a rate of 2 μL/min resulted in a mild change of $\sim$ 2.5 mmHg in ICP, whereas little or no change in ICP was observed during the simultaneous injection and withdrawal in the dual-syringe protocol. Repeated measures two-way analysis of variance (ANOVA) was performed; interaction P value < 0.0001; n = 5 mice for single-injection and n = 6 mice for dual-syringe. The shaded regions above and below the plot lines indicate the standard error of the mean (SEM).

Even though the increase in ICP observed during a standard single-injection infusion is within the physiological range, and comparable to variations in ICP in response to respiration or changes in body posture[20], this artificial elevation of ICP is one of the most commonly criticized aspects of tracer studies in which an exogenous fluorescent tracer is injected intracisternally. Although we should not ignore the effect of this transient ICP elevation on the initial distribution of the tracer, especially at the ventral portion of the CSF compartment, the ICP returns to baseline within 5 minutes of cessation of the infusion, while the CSF influx continues for the duration of the experiment ($\sim$ 30 min)[20]. Moreover, if the induced ICP elevation were responsible for the tracer penetration into the brain, then the differences in tracer influx between different arousal states, anesthetics, or experimental groups that receive the same infusion protocols[2, 24, 25] would not occur.

Flow measurements in perivascular spaces

We studied the motion of tracer particles infused with the new dual-syringe protocol (lower panels in Fig. 3) and compared it with the motion of tracer particles observed by Mestre et al.[5] using the single-injection protocol (upper panels in Fig. 3), using particle tracking to examine flow of CSF in pial periarterial spaces. The periarterial spaces of the cortical branches of the middle cerebral artery (MCA) were chosen for imaging. In the new protocol, particles appeared in the visualized spaces at $\sim$ 300 s after infusion was complete. This time scale is similar to that in our previous report[5] of 292 ± 26s, but particle counts were lower than those observed using the single-injection technique, likely because some of the injected particles were siphoned into the withdrawal line of the dual-syringe setup. However, a sufficient number of particles made their way into the PVSs to enable rigorous flow measurements (see Supplemental Video S1). Results obtained from the particle tracking analysis are shown in Figure 3. In experiments that lasted at least 10 minutes, we tracked more than 6200 particles in each of the six

mice that were infused using the new protocol. An example of the superimposed particle tracks imaged in an experiment is shown in Figure. 3e. The particle tracks are mostly confined to the perivascular spaces surrounding the artery, with occasional cross-overs from one side of the artery to the other. The distribution of particle tracks is spatially continuous across the width of the imaged PVSs under both infusion methods (Figs. 3a[5] & 3e), reaffirming that surface periarterial spaces are open, rather than porous, spaces[26]. The direction of the observed fluid flow in the different branches is indicated by the arrows in Figures 3b and 3f. If injection were driving the flow, we should observe dominant directional transport of tracer particles only when they are infused using the single-injection method, and little or no transport with the dual-syringe method. The time-averaged (bulk) flow for both infusion methods is in the same direction as that of the blood flow, providing evidence that net CSF transport into the brain from the subarachnoid space is not caused by the injection. For both infusion methods, we observed no net flow in the direction opposite to that of blood flow, as some recent reports have suggested[15,27]. Figure 3g shows that the average flow speed in pial (surface) periarterial spaces varies across the PVS, consistent with prior reports[5] shown in Figure 3c. The velocity profile is parabolic-like (Figs. 3d and 3h); the flow is fastest ($\sim 50\ \mu$m/s) at the center of the PVS and slows to zero at the walls. This parabolic-like shape of the velocity profile is consistent with laminar, viscous-dominated flow of CSF through an open annular space, and not through a porous medium, indicating that pial periarterial spaces are open[26].

Figure 3. Particle tracking velocimetry in surface periarterial spaces using the single-injection method (panels in first row[5]) and the new dual-syringe method (second row). The superimposed particle tracks shown in panels (a) and (e) have similar, continuous spatial distributions and show similar sizes of the perivascular spaces. The time-averaged velocity fields shown in panels (b) and (f) both show net downstream flow of fluid in the same direction as that of the blood flow. The flow-speed distributions plotted in panels (c) and (g) show comparable speeds, with the fastest flow at the center of the imaged periarterial space and the slowest flow near the boundaries. Panels (d) and (h) show average flow-speed profiles across the corresponding colored lines spanning the PVS in panels (c) and (g), smoothed by interpolation. The parabolic-like nature of these velocity profiles is what is expected for viscous flow in an open channel. Scale bars indicate 50 μm.

Further analysis of the data obtained from particle tracking demonstrates the close similarity between the flows observed in the two protocols, as shown in Figure 4. A time-history of the measured flows — quantified by the spatial root-mean-square velocity computed at each instant of time (V_{rms}) — portrays very similar behavior over times much longer than the time it takes for the ICP to return to normal after the infusion (Fig. 4a). (The times shown here begin when particles were first seen or when the imaging was started: these times differ by less than 1 minute and so do not affect the results significantly.) The pulsatile nature of the flow at small time scales is depicted in Figures 4b and 4c. If injection-induced elevated ICP were propelling the flow, we would observe large V_{rms} values early in the single-injection experiments, followed by an exponential decay, and we would observe little or no flow in the dual-syringe experiments, as the ICP remains unchanged. Since we observe very similar trends in the time-history profiles in both infusion protocols, the pressure differences driving the flow must be the same and independent of the infusion method.

Figure 4. Measured flow characteristics. Panel (a) shows that V_{rms} remains much the same over the course of the velocity measurements for both infusion methods. Repeated measures two-way ANOVA was performed; ns, not significant; n = 5 mice for single-injection and n = 6 mice for dual-syringe. The solid lines represent the mean value of V_{rms} and the shaded area represents the standard error of the mean within each time bin. The pulsatility of typical measured flows is depicted in panels (b) and (c). Panel (d) shows mean downstream flow speeds and panel (e) shows backflow fractions for the individual experiments, with overall mean values shown as open circles (and bars showing the standard error of the mean). The nearly identical values for the two protocols demonstrate that the flow is independent of the injection method employed. Unpaired Student's t-test was performed; n = 5 or 6 mice per group; ns, not significant; mean $\pm$ SEM.

Figure 4d shows mean flow speeds computed by averaging the downstream velocity component over space and time for each experiment. The overall mean flow speed (open circles) is 15.71 ± 6.2 μm/s for all the single-injection experiments and 17.67 ± 4.42 μm/s for all the dual-syringe experiments, *nearly identical* values. These values are also nearly identical to the mean speed of 17 ± 2 μm/s reported by Bedussi et al.[4], from experiments that also used a single-injection protocol. The mean flow speeds represent the speeds at which tracer particles (or cerebrospinal fluid) are transported in the direction of arterial blood flow (downstream), and presumably into the brain. If the observed flows were injection-induced, we would expect vastly different transport speeds with faster mean flows associated with the single-syringe injection method. We also computed a 'backflow fraction' for each experiment, as the fraction of the downstream velocity measurements showing motion in the retrograde direction (opposite that of the blood flow): the results are shown in Figure 4e. An injection-driven flow would be dominantly unidirectional and would exhibit a much smaller backflow fraction. However, the backflow fraction is nearly identical ($\sim$ 0.25) for single-injection and dual-syringe infusions. The nearly identical mean flow speeds and backflow fractions further demonstrate that the observed flows are natural, and not artifacts of the infusion.

Flow is strongly regulated by the cardiac cycle

It has been variously suggested that CSF bulk flow might be driven by the cardiac cycle, the respiratory cycle, or perhaps both[4, 6, 28, 29], with evidence indicating much stronger correlation with the cardiac cycle[5, 12]. Having shown here that the tracer infusion is not driving the observed flows, we wanted to use our new data to examine whether arterial pulsations are indeed the dominant driving mechanism, using methods similar to those in our earlier report[5]. We use the simultaneous measurements of the electrocardiogram (ECG) and respiration in conjunction with particle tracking to determine the relative importance of the

cardiac and respiratory cycles[30], and also to see if there is any difference between the two infusion methods (Fig. 5).

Figure 5. CSF velocity variations over the cardiac and respiratory cycles. Panels (a) and (b) show the measured V_{rms} conditionally averaged over the cardiac and respiratory cycles, based on the synchronized measurements of ECG, respiration, and velocity, for the single-injection (a) and the dual-syringe (b) protocols. Panel (c) for single injection and panel (e) for dual syringe both show that the peaks in the ECG are immediately followed by peaks in V_{rms}, indicating a strong correlation between heart rate and fluid motion in both injection protocols. No consistent trends are seen when V_{rms} is averaged over the respiratory cycle, as shown in panels (d) and (f). Panels (g) and (h) show the mean and the standard error of the mean of probability density functions of the delay time Δt between the peak in the cardiac (cyan) or respiration (green) cycle and the subsequent peak in V_{rms}, for single-injection ($n = 5$) and dual-syringe ($n = 6$) methods respectively. Panel (i) shows that the average Δt between peaks in the cardiac cycle and V_{rms} is nearly identical for both protocols, and that the peak in V_{rms} typically occurs ~ 0.05 s after the peak in the cardiac cycle. Unpaired Student's t-test was performed; n = 5 or 6 mice per group; ns, not significant; mean $\pm$ SEM.

We find that the measured flow quantities, such as V_{rms}, are strongly modulated by the cardiac cycle but only very weakly by respiration (Figs. 5a[5] & 5b). This strong correlation of the flow with the heart rate is exhibited equally under both the single-syringe and dual-syringe protocols, as shown in Figures 5c and 5e, where the peak in V_{rms} occurs soon after the peak in the cardiac cycle. Probability density functions of Δt, the delay time between peaks in V_{rms} and cardiac/respiratory cycles, also predict a much greater likelihood of peaks in V_{rms} following the peak in the cardiac cycle (Figs. 5g & 5h). We also observe nearly identical average delay times of ~ 0.05 s between peaks in V_{rms} and the cardiac cycle for both protocols (Fig. 5i). No such correlation is observed when the V_{rms} is conditionally averaged over respiration cycles (Figs. 5d and 5f). These observations corroborate prior reports that the cardiac cycle, through the pulsation of arterial walls, is the strongest mechanism for pumping CSF through surface periarterial spaces.

Discussion and conclusions

The net transport of CSF from the subarachnoid space deep into the brain tissue through periarterial spaces surrounding the pial cerebral arteries is a critical component by which metabolic waste products are removed to keep the brain healthy[31–33]. While experiments in live mice have shown that fluid is pumped in the direction of blood flow and into brain by forces linked to the pulsation of arterial walls, several published papers have speculated that the observed flows might instead be artifacts of non-natural elevation of ICP levels caused by tracer infusion into the cisterna magna. In this study, we designed a new infusion protocol that enabled tracer-particle infusion with no net addition of fluid and near-zero changes in ICP. We used two-photon microscopy and particle-tracking velocimetry, and found flows of CSF in the surface periarterial spaces that are essentially identical to the flows found earlier using the traditional single-syringe injection protocol. The measured flows are pulsatile, viscous-dominated, laminar flows, with mean flow in the direction of blood flow in the cerebral arteries. Our flow visualization techniques and synchronized measurements of ICP and heart and respiration rates enabled us to show that the observed flows are not driven by pressure differences induced by the tracer infusion. Furthermore, we confirmed that forces linked to the cardiac cycle remain the strongest driver of the flow in pial periarterial spaces.

Our new experiments provide several lines of evidence to prove that the observed bulk flow is not induced by tracer infusion methods currently used. Tracer infusion at rates of $1 - 2$ μL/min have typically been employed[4,5] to add 10 μL of fluid to the subarachnoid space. Although this addition of fluid is greater than the natural CSF production rate (0.38 μl/min[34]), and it induces a small increase in ICP, our results show that the ICP returns to its baseline value within 5 minutes after injection is completed. If infusion were propelling the mean flow, particle transport would cease after the ICP reverted to normal, or would not occur at all with dual-syringe injection, where the ICP is not affected. We continue to observe particles being transported along the periarterial spaces long after the return of ICP to its baseline value in the single-syringe experiments and at comparable times in the dual-syringe experiments. This demonstrates that the bulk flow we observe is not caused by the infusion.

Elevated ICP levels create large pressure differences across the brain, but these pressure differences will undergo exponential decay because of the brain's compliance and proclivity to achieve stasis. If this exponential relaxation of ICP were to drive fluid flow, the measurements from particle tracking would reflect this decay. exhibiting fast flows at early times which then gradually subside. However, our measurements show that the mean flow remains nearly constant and similar over periods that are 2 to 3 times longer than the infusion time, for several healthy mice and both infusion protocols. Any variablity is probably due to physiological differences between different mice. Moreover, if infusion-driven flows were prevalent, we would expect to see much faster mean flow speeds. Yet, with the new dual-syringe method, with no net infusion, we observe flow speeds that are nearly identical to those observed in our earlier study[5], and very close to those in a study that used injection rates many orders of magnitude smaller[4]. An infusion-driven flow would also be unidirectional, with no retrograde motion, but we observed consistent backflows of particles for both infusion protocols.

Our results show that arterial pulsations are the strongest driver of the observed periarterial flow. The peaks of V_{rms} that we measured across specimens, for both infusion protocols, appear shortly after the peaks in the cardiac cycle, but are not correlated with the respiratory cycle. Probability density functions show that the delay times between the peaks in the cardiac cycle and the peaks in V_{rms} are nearly identical for the two infusion methods. While we can not rule out other physiological mechanisms that might drive CSF flow, such as functional hyperemia[35] or vasomotion[15,36], we conclude that the currently employed methods of tracer infusion are not responsible for the observed flows.

Methods

Animals and surgical preparation

All experiments were approved and conducted in accordance with the relevant guidelines and regulations stipulated by the University Committee on Animal Resources of the University of Rochester Medical Center (Protocol No. 2011-023), certified by Association for Assessment and Accreditation of Laboratory Animal Care. An effort was made to minimize the number of animals used. We used 8- to 12-week-old male C57BL/6 mice acquired from Charles River Laboratories (Wilmington, MA, USA). In all experiments, animals were anesthetized with a combination of ketamine (100 mg/kg) and xylazine (10 mg/kg) administered intraperitoneally. Depth of anesthesia was determined by the pedal reflex test. Once reflexes had ceased, anesthetized mice were fixed in a stereotaxic frame for the surgical procedure and body temperature was kept at 37°C with a temperature-controlled warming pad.

Dual-syringe protocol

For *in vivo* imaging, anesthetized mice were fixed in a stereotaxic frame and body temperature was maintained at 37.5°C with a rectal probe-controlled heated platform (Harvard Apparatus). Two 30-gauge needles were inserted into the cisterna magna, as previously described[37]. Briefly, the dura mater of mice was exposed after blunt dissection of the neck muscles so that a

cannula could be implanted into the cisterna magna (CM), which is continuous with the subarachnoid space. A cranial window was prepared over the right middle cerebral artery (MCA) distribution. The dura was left intact, and the craniotomy ($\simeq$4 mm in diameter) was filled with aCSF, covered with a modified glass coverslip, and sealed with dental acrylic. Afterwards, two 30-gauge needles were inserted into the cisterna magna, as described above. Using a syringe pump (Harvard Apparatus Pump 11 Elite), red fluorescent polystyrene microspheres (FluoSpheres^{TM} 1.0 μm, 580/605 nm, 0.25% solids in aCSF, Invitrogen) were infused up to a total volume of 20 μL via one of the cisterna magna cannulae while CSF was simultaneously withdrawn through the other cannula at an equal rate of 2 μL/min with a coupled syringe pump.

Intracranial pressure measurements

Anesthetized mice were fixed in a stereotaxic frame, and two 30-gauge needles were inserted into the cisterna magna, as described above. A third cannula was inserted via a small burr hole into the right lateral ventricle (0.85 mm lateral, 2.10 mm ventral and 0.22 mm caudal to bregma). Mice were then placed in a prone position. In the first set of experiments, 10 μL of artificial CSF (aCSF) was injected into the CM at a rate of 2 μL/min via one of the CM cannulae using a syringe pump (Harvard Apparatus Pump 11 Elite). In the second set of experiments, aCSF was injected at the same rate while withdrawing CSF from the cisterna magna via the other CM cannula at an equal rate using a coupled syringe pump (Harvard Apparatus Pump 11 Elite). In both experiments, intracranial pressure (ICP) was monitored via the ventricle cannulation connected to a transducer and a pressure monitor (BP-1, World Precision Instruments). ICP was acquired at 1 kHz, digitized, and monitored continuously for the duration of the infusion experiments with a DigiData 1550B digitizer and AxoScope software (Axon Instruments).

In vivo two-photon laser-scanning microscopy

Two-photon imaging was performed using a resonant scanner B scope (Thorlabs) with a Chameleon Ultra II laser (Coherent) and a 20$\times$ water immersion objective (1.0 NA, Olympus). Intravascular FITC-dextran and red microspheres were excited at a 820 nm wavelength and images were acquired at 30 Hz (ThorSync software) simultaneously with physiological recordings (3 kHz, ThorSync software), as previously described[5]. To visualize the vasculature, fluorescein isothiocyanate–dextran (FITC–dextran, 2,000 kDa) was injected intravenously via the femoral vein immediately before imaging. Segments of the middle cerebral artery were distinguished on the basis of morphology: surface arteries passing superficially to surface veins and exhibiting less branching at superficial cortical depths. ECG and respiratory rate were acquired at 1 kHz and 250 Hz, respectively, using a small-animal physiological monitoring device (Harvard Apparatus). The signals were digitized and recorded with a DigiData 1550A digitizer and AxoScope software (Axon Instruments).

Image processing

Images with spatial dimensions 512 x 512 were obtained from two-photon microscopy. Each image is 16-bit with two channels, red and green. The FITC-dextran injected in the vasculature is captured via the green channel while the red channel is used to image the fluorescent microspheres flowing in the perivascular spaces. Image registration via rigid translation is performed on each image in the time series to account for movement by the mouse in the background. The image registration is implemented using an efficient algorithm in Matlab[38] to an accuracy of 0.2 pixels. Erroneous correlations in the translation are manually corrected by linear interpolation. The translations obtained are sequentially applied to images that are padded with zero-value pixels. This ensures spatial dimension homogeneity across all images without modifying the image resolution. Particles are then detected by applying a minimum intensity threshold to each image. Typically, particles were resolved across 3-4 pixels in the image with spatial resolution of 1.29 μm.

Particle-tracking velocimetry

The particles detected in each image were tracked using an automated PTV routine in MATLAB[39,40]. Briefly, the algorithm locates each particle with a sub-pixel accuracy and obtains a series of particle locations (particle tracks) for the entire duration of the recorded video. Particle velocities were calculated by convolution with a Gaussian smoothing and differentiation kernel. Stagnant particles that have adhered to the wall of the artery or the outer wall of the PVS, and hence no longer track the CSF flow, were masked in each image by subtracting a dynamic background image. This image was different for each frame and was computed by taking the average of 100 frames before and 100 frames after the given image. This method of masking was applied only to the dual-syringe data; the single-syringe data used a simpler masking approach with a single background image[5]. Time-averaged flow velocities were obtained by segregating the imaged domain into a 70 x 70 grid, with a resolution of 7.5 x 7.5 pixels in each direction. All velocity measurements for a chosen time interval were binned based on their grid position. Average flow speeds were computed using bins with at least 15 measurements. The downstream velocity component was calculated as the dot product $\boldsymbol{u} \cdot \hat{\boldsymbol{u}}_{avg}$, where $\boldsymbol{u}$ is the instantaneous particle velocity and $\hat{\boldsymbol{u}}_{avg}$ is the field of unit vectors computed from the time-averaged flow field, in the direction of arterial blood flow.

Statistical analysis

All statistical analyses were performed on GraphPad Prism 8 (GraphPad Software). Data in all graphs are plotted as mean $\pm$ standard error of the mean (SEM) over the individual data points and lines from each mouse. Statistical tests were selected after evaluating normality (D'Agostino Pearson omnibus test). When the sample size did not allow for normality testing, both parametric and nonparametric tests were performed and, in all cases, yielded the same result. Sphericity was not assumed; in all repeated measures, two-way ANOVAs and a Geisser-Greenhouse correction were performed. All hypothesis testing was two-tailed and exact P values were calculated at a 0.05 level of significance and stated in the figure legends.

References

1. Iliff, J. J. *et al.* A paravascular pathway facilitates CSF flow through the brain parenchyma and the clearance of interstitial solutes, including amyloid β. *Sci Transl Med* **4**, 4 (2012).

2. Xie, L. *et al.* Sleep drives metabolite clearance from the adult brain. *Science* **342**, 373–377 (2013).

3. Peng, W. *et al.* Suppression of glymphatic fluid transport in a mouse model of Alzheimer's disease. *Neurobiol. disease* **93**, 215–225 (2016).

4. Bedussi, B., Almasian, M., de Vos, J., VanBavel, E. & Bakker, E. N. T. P. Paravascular spaces at the brain surface: Low resistance pathways for cerebrospinal fluid flow. *J Cerebr Blood F Met* 0271678X1773798–8 (2017).

5. Mestre, H. *et al.* Flow of cerebrospinal fluid is driven by arterial pulsations and is reduced in hypertension. *Nat Commun* **9**, 4878 (2018).

6. Hadaczek, P. *et al.* The "perivascular pump" driven by arterial pulsation is a powerful mechanism for the distribution of therapeutic molecules within the brain. *Mol Ther* **14**, 69–78 (2006).

7. Thomas, J. H. Fluid dynamics of cerebrospinal fluid flow in perivascular spaces. *J. R. Soc. Interface* **16**, 52–57 (2019).

8. Smith, A. J., Yao, X., Dix, J. A., Jin, B. J. & Verkman, A. S. Test of the 'glymphatic' hypothesis demonstrates diffusive and aquaporin-4-independent solute transport in rodent brain parenchyma. *eLife* **6**, e27679 (2017).

9. Asgari, M., de Zélicourt, D. & Kurtcuoglu, V. Glymphatic solute transport does not require bulk flow. *Sci Rep* 1–11 (2016).

10. Bilston, L. E., Fletcher, D. F., Brodbelt, A. R. & Stoodley, M. A. Arterial pulsation-driven cerebrospinal fluid flow in the perivascular space: A computational model. *Comput. Method. Biomech.* **6**, 235–241 (2003).

11. Wang, P. & Olbricht, W. L. Fluid mechanics in the perivascular space. *J Theor Biol* **274**, 52–57 (2011).

12. Iliff, J. J. *et al.* Cerebral arterial pulsation drives paravascular CSF-interstitial fluid exchange in the murine brain. *J Neurosci* **33**, 18190–18199 (2013).

13. Diem, A. K. *et al.* Arterial Pulsations cannot Drive Intramural Periarterial Drainage: Significance for Aβ Drainage. *Front Neurosci* **11**, 353–9 (2017).

14. Kedarasetti, R. T., Drew P., T. & Costanzo, F. Arterial pulsations drive oscillatory flow of CSF but not directional pumping. *Sci. Rep.* **10** (2020).

15. van Veluw, S. J. *et al.* Vasomotion as a driving force for paravascular clearance in the awake mouse brain. *Neuron* **105**, 549–561 (2020).

16. Smith, A. J. & Verkman, A. S. The "glymphatic" mechanism for solute clearance in Alzheimer's disease: game changer or unproven speculation? *The FASEB J.* **32**, 543–551 (2018).

17. Croci, M., Vinje, V. & Rognes, M. E. Uncertainty quantification of parenchymal tracer distribution using random diffusion and convective velocity fields. *Fluids Barriers CNS* **16**, 32 (2019).

18. Sharp, M. K., Carare, R. O. & Martin, B. A. Dispersion in porous media in oscillatory flow between flat plates: applications to intrathecal, periarterial and paraarterial solute transport in the central nervous system. *Fluids Barriers CNS* **16**, 13 (2019).

19. Vinje, V., Eklund, A., Mardal, K.-A., Rognes, M. E. & Støverud, K.-H. Intracranial pressure elevation alters CSF clearance pathways. *Fluids Barriers CNS* **17**, 1–19 (2020).

20. Mestre, H., Mori, Y. & Nedergaard, M. The brain's glymphatic system: Current controversies. *Trends Neurosci.* (2020).

21. Hladky, S. B. & Barrand, M. A. Elimination of substances from the brain parenchyma: efflux via perivascular pathways and via the blood–brain barrier. *Fluids Barriers CNS* **15**, 30 (2018).

22. Bakker, E. N. T. P., Naessens, D. M. P. & VanBavel, E. Paravascular spaces: entry to or exit from the brain? *Exp. physiology* **104**, 1013–1017 (2019).

23. Faghih, M. M. & Sharp, M. K. Is bulk flow plausible in perivascular, paravascular and paravenous channels? *Fluids Barriers CNS* **15**, 1–10 (2018).

24. Mestre, H. *et al.* Aquaporin-4-dependent glymphatic solute transport in the rodent brain. *eLife* **7**, e40070 (2018).

25. Hablitz, L. M. *et al.* Increased glymphatic influx is correlated with high EEG delta power and low heart rate in mice under anesthesia. *Sci. Adv.* **5**, eaav5447 (2019).

26. Min-Rivas, F. *et al.* Surface periarterial spaces in the mouse brain are open, not porous. *J Roy Soc Interface* in press (2020).

27. Aldea, R., Weller, R. O., Wilcock, D. M., Carare, R. O. & Richardson, G. Cerebrovascular smooth muscle cells as the drivers of intramural periarterial drainage of the brain. *Front. Aging Neurosci.* **11** (2019).

28. Rennels, M. L., Gregory, T. F., Blaumanis, O. R., Fujimoto, K. & Grady, P. A. Evidence for a 'paravascular' fluid circulation in the mammalian central nervous system, provided by the rapid distribution of tracer protein throughout the brain from the subarachnoid space. *Brain Res* **326**, 47–63 (1985).

29. Yamada, S. *et al.* Influence of respiration on cerebrospinal fluid movement using magnetic resonance spin labeling. *Fluids Barriers CNS* **10**, 36 (2013).

30. Santisakultarm, T. P. *et al.* In vivo two-photon excited fluorescence microscopy reveals cardiac-and respiration-dependent pulsatile blood flow in cortical blood vessels in mice. *Am. J. Physiol. Circ. Physiol.* **302**, H1367–H1377 (2012).

31. Rasmussen, M. K., Mestre, H. & Nedergaard, M. The glymphatic pathway in neurological disorders. *The Lancet Neurol.* **17**, 1016–1024 (2018).

32. Reeves, B. C. *et al.* Glymphatic system impairment in Alzheimer's disease and idiopathic normal pressure hydrocephalus. *Trends Mol. Medicine* **26**, 285–295 (2020).

33. Nedergaard, M. & Goldman, S. A. Glymphatic failure as a final common pathway to dementia. *Science* **370**, 50–56 (2020).

34. Oshio, K., Watanabe, H., Song, Y., Verkman, A. S. & Manley, G. T. Reduced cerebrospinal fluid production and intracranial pressure in mice lacking choroid plexus water channel Aquaporin-1. *The FASEB J.* **19**, 76–78 (2005).

35. Kedarasetti, R. T. *et al.* Functional hyperemia drives fluid exchange in the paravascular space. *Fluids Barriers CNS* **17**, 1–25 (2020).

36. Kiviniemi, V. *et al.* Ultra-fast magnetic resonance encephalography of physiological brain activity–Glymphatic pulsation mechanisms? *J. Cereb. Blood Flow & Metab.* **36**, 1033–1045 (2016).

37. Xavier, A. L. *et al.* Cannula implantation into the cisterna magna of rodents. *J. Vis. Exp.* e57378 (2018).

38. Guizar-Sicairos, M., Thurman, S. T. & Fienup, J. R. Efficient subpixel image registration algorithms. *Opt. Lett.* **33**, 156–158 (2008).

39. Kelley, D. H. & Ouellette, N. T. Using particle tracking to measure flow instabilities in an undergraduate laboratory experiment. *Am J Phys* **79**, 267–273 (2011).

40. Ouellette, N. T., Xu, H. & Bodenschatz, E. A quantitative study of three-dimensional Lagrangian particle tracking algorithms. *Exp. Fluids* **40**, 301–313 (2006).

Acknowledgements

This work was supported by the NIH/National Institute of Aging (grant RF1AG057575) and by the U. S. Army Research Office (grant MURI W911NF1910280). J.T. was supported by a Career Award at the Scientific Interface by the Burroughs Wellcome Fund. We thank Keith Sharp for recommending something like our dual-syringe experiments to us in July 2019.

Author contributions

A.R. and A.L. contributed equally to this work. J.H.T. suggested the experiments, which were then carried out by A.L. and A.R., following the methods developed by M.N., H.M, D.H.K, J.T., and J.H.T. Data analysis was performed by A.R. and A.L. All of the authors contributed to writing and editing the paper.

Competing interests

The authors declare that they have no competing interests.