Evaluating and Improving SSU rRNA PCR Primer Coverage via Metagenomes from Global

Ocean Surveys

Jesse McNichol[1], Paul M. Berube[2], Steven J. Biller[2,3], Jed A. Fuhrman[1]

1 - Department of Biological Sciences, University of Southern California, Los Angeles,

California, USA

2 - Department of Civil and Environmental Engineering, Massachusetts Institute of

Technology, Cambridge, MA, USA

3 - Department of Biological Sciences, Wellesley College, Wellesley, MA, USA

Abstract:

Small subunit ribosomal RNA (SSU rRNA) amplicon sequencing comprehensively profiles

microbiomes, but results will only be accurate if PCR primers perfectly match environmental

sequences. To evaluate whether primers commonly used in microbial oceanography match naturally-

occurring organisms, we compared primers with > 300 million rRNA sequences retrieved from

globally-distributed metagenomes. The best-performing 16S primers were 515Y/926R and

515Y/806RB which perfectly matched most sequences (~0.95). Considering Cyanobacteria/Chloroplast

16S, 515Y/926R had the highest coverage (0.99), making it ideal for quantifying phytoplankton. For

18S sequences, 515Y/926R performed best (0.88), followed by V4R/V4RB (18S-specific; 0.82). Using

Atlantic and Pacific BioGEOTRACES field samples, we demonstrate high correspondence between

515Y/926R amplicons (generated as part of this study) and metagenomic 16S rRNA (median R^2=0.98,

n=272), indicating amplicons can produce equally accurate community composition data versus

shotgun metagenomics. Since our pipeline identifies missed taxa, we suggest modifications to improve

coverage of biogeochemically-important oceanic microorganisms - a strategy applicable to any

environment with metagenomic data.

Significance Statement:

Quantification of taxonomically-informative marker genes using PCR amplification and

high-throughput sequencing is a low-cost technique for monitoring distributions and changes of

microbial communities across space and time. In order to maximize this procedure's effectiveness, it is

essential that environmental organisms match PCR primer sequences exactly. In this study, we

developed a software pipeline to evaluate how well commonly-used primers match primer-binding

regions from globally-distributed short-read oceanic metagenomes. Our results demonstrate common

primer sets vary widely in performance, and that including additional degenerate bases is a simple

strategy to maximize environmental coverage. Written in the reproducible snakemake workflow

language and publicly accessible, our pipeline provides a general-purpose tool to guide rational design

of PCR primers for any environment with metagenomic data.

Introduction:

Amplicon sequencing is a powerful tool for understanding microbial community composition

and dynamics in the oceans and other ecosystems (1), but the PCR amplification step is potentially

biased due to both technical issues during amplification and mismatches to organisms found in natural

ecosystems (2–5). Despite these potential issues, PCR amplicon sequencing retains several advantages

that make it desirable to investigate and correct biases. First of all, it is a high-throughput and low cost

technique, making it suitable for large numbers of samples e.g. for global surveys of sediment, water,

animal-associated, and other microbial communities (1). Secondly, the targeted nature of the assay

means relatively small numbers of sequences are sufficient for detecting rare organisms even when we

have no genomes from any of their relatives, due to the conserved nature of the molecule and

comprehensive SSU rRNA sequence database. Thirdly, modern "denoising" methods produce exact

amplicon sequence variants (ASVs) that can be used as stable and intercomparable biogeographic

markers (6). Finally, amplicon datasets are simpler to analyze since they consist of a single gene.

Because of this relative simplicity, the complete dataset can be classified with databases such as SILVA

(7). In contrast, the complexity of metagenomic samples and current lack of reference genomes for

many organisms makes it less possible to classify all sequences.

Recent studies have shown that PCR amplicon sequencing of mock microbial communities

can recover known abundances extremely well (3, 8). Conversely, these studies have also underscored

the key weakness of PCR-based methods - i.e. that a single mismatch between the primer and template

sequences can have a dramatic effect on the measured community composition (e.g. ~10-fold bias

against the taxon in question, from a single internal mismatch). In addition, the extent of this bias is not

easily predicted if the mismatched base is not at the 3' terminus of the primer, where biases are known

to be severe (3, 8). Therefore, PCR primers should ideally incorporate degenerate bases to match all

targeted organisms without sacrificing specificity or PCR efficiency. This is typically accomplished by

comparing primer sequences to a SSU rRNA database such as SILVA and then checking for

mismatches to organisms known to be abundant in the environment of interest. This general approach

has led to marked improvements in primer design, for example by Apprill et al (2015) and Parada et al

(2015) who reported that small modifications to existing primers could better quantify the dominant

marine taxa SAR11 and *Thaumarchaea* (2, 3).

However, this approach remains limited because it relies on expert knowledge that may be

incomplete and does not take into account the potential heterogeneity of natural systems. Wear et al.

(2018) addressed this knowledge gap by testing 4 commonly-used primer sets on a 16S mock microbial

community derived from natural seawater communities near Santa Barbara (9). These authors tested

the effect of their analysis pipeline on recovery of mock community sequences for different primers,

and found their primer-pipeline combination had several sequence-specific biases, in some cases due to

a primer mismatch. While an important step towards cross-comparing primers in an oceanographic

context, Wear et al. note that results are specific to their mock microbial community, analyzed with

their particular pipeline, and thus it remains unknown how representative their results are for other

ecosystems and other pipelines. More specifically, it is currently unknown how many primer-

mismatched rRNA sequences occur across diverse oceanographic environments, a key piece of

information for designing field measurements and interpreting results.

To more fully evaluate the extent to which commonly-used primers perfectly match naturally

occurring sequences, we developed a workflow to conduct *in silico* primer evaluations with

metagenomic SSU rRNA fragments, under the broad assumption that metagenomes have fewer

methodological biases than PCR. We use the results from this pipeline to carry out a more objective

evaluation of commonly used primers for pelagic marine ecosystems, suggest improvements, and also

compare the fidelity of SSU rRNA amplicon sequencing versus shotgun metagenomic sequencing on

these globally-distributed samples. Because this pipeline is broadly applicable for any environment

where metagenomic data exists, it will allow investigators to design environment-specific PCR probes

that optimally recover targeted organisms. Our approach has the added advantage of providing

evidence-based data to correct datasets generated with sub-optimal primers in cases where new data

cannot be regenerated due to a lack of archived sample material.

Results and Discussion:

Overall primer coverage for pelagic ocean datasets (BiHOBA, TARA, MSPINA,

MBARI, SPOT):

To evaluate whether commonly used primer pairs are likely to return biased views of marine

microbial communities, we applied our workflow to globally-distributed metagenomic datasets that

reflect a range of depths and nutrient regimes (Table 1; Fig 1), testing primers that are commonly used

in environmental microbiology (Table 2; (2–4, 10–15)). Metagenomic samples included worldwide

sampling expeditions such as Tara oceans (TARA (16)), the Malaspina expedition (MSPINA (17)), and

BioGEOTRACES (18). Among these three campaigns, the MSPINA metagenomes focus largely on the

deep sea, whereas TARA and BioGEOTRACES sample primarily the sun-lit ocean. In addition, we

used time-series data from the Hawaii Ocean Time-Series and the Bermuda Atlantic Time-Series (19,

20) (HOT/BATS; both analyzed with BioGEOTRACES as BiHOBA [BioGEOTRACES-HOT-BATS]),

a bloom time-series in Monterey Bay (21) (MBARI), and the San Pedro Ocean Time-Series (22)

(SPOT). We also included in the analysis two sites representing suboxic/anoxic systems - the Saanich

inlet time-series (Saanich) (23), and samples collected from hydrothermally-altered sediments in

Guaymas basin (24) (GuaySed). We also generated new 16S/18S universal amplicon sequences for two

BioGEOTRACES cruise transects (GA03/GP13; noted with open circles on Fig 1) which were used for

making intercomparisons between metagenomes and amplicons (discussed further below).

We chose the percentage of perfect matches as a metric to evaluate potential PCR

performance since previous studies have shown that even a single mismatch can significantly bias

results. To summarize data from pelagic ocean ecosystems, we plotted the fraction of metagenomic

reads that perfectly matched primer pairs across all pelagic oceanic datasets (excluding GuaySed and

Saanich) with violin plots/boxplots to visualize the distribution of metagenomic primer coverage values

(Fig 2). In the following discussion, we mostly use "median coverage" as a summary statistic to

compare primer set performance.

Median coverage varied widely among primer pairs, with 515Y/806RB and 515Y/926R

showing the most consistently high coverage (Fig 2). The ability of all primers to capture the

underlying diversity in the metagenomic samples can be improved to some extent by increasing

degeneracies, though this varied significantly between primer sets. Adding a single additional

degeneracy fixes the majority of mismatches for some primers (e.g. 785R; discussed further below),

whereas others have greater than 2 mismatches to metagenomic sequences and thus would require

extensive modifications (Figures S1-S28). Below, we discuss the performance of these primers across 4

broad taxonomic groups (*Archaea, Bacteria, Cyanobacteria+*Plastidal 16S, and *Eukarya*). We

separated *Cyanobacteria* + phytoplankton plastidal 16S from bacterial 16S since these organisms are

responsible for the vast majority of marine primary productivity and thus are very important to quantify

accurately.

For *Archaea*, the primer with the best median coverage was the EMP 515Y/806RB

combination (coverage=0.996), followed by 515Y/926R (0.982), and 926wF/1392R (0.952). We also

note that our results suggest the 341F/785R primer combination could amplify most *Archaea* with the

addition of several degeneracies in 341F (Fig S1-S7).

For *Bacteria* (excluding *Cyanobacteria* and plastidal 16S sequences), the primer with the

best median coverage was the 515Y/926R combination (0.961), followed by 515Y/806RB (0.955),

27F/338R (0.888), 926wF/1392R (0.869), and 341F/785R (0.525). The relatively poor performance of

341F/785R is mainly due to the reverse primer (median individual primer coverages: 341F=0.887,

785R=0.624), whereas the other two less optimal primer pairs have more even individual performances

(27F=0.968, 338R=0.913; 926wF=0.916, 1392R=0.943).

For *Cyanobacteria* and chloroplast 16S (derived from eukaryotic phytoplankton), the primer

pair with the highest median coverage was 515Y/926R (0.992), followed by 515Y/806RB (0.881), 27F/

338R (0.826), 926wF/1392R (0.714), and 341F/785R (0.702). We note that the vast majority of these

mismatched sequences come from chloroplast 16S sequences - the common cyanobacterial groups

Prochlorococcus and *Synechococcus* have high coverage with all primers listed above, so this is only a

concern for those wishing to use chloroplast 16S data to quantify eukaryotic phytoplankton.

For *Eukarya*, the primer pair with the highest median coverage was 515Y/926R (0.883),

followed by V4F/V4RB (0.822), 926wF/1392R (0.792), and 1389F/1510R (0.732). The distributions of

coverage were considerably broader for *Eukarya* versus all other 16S categories, which may be due to a

higher sequence heterogeneity among eukaryotic genomes. As above, all primer sets could be improved

with the addition of some degeneracy (Fig S22-S28), but approaching perfect coverage seems less

practically achievable for eukaryotic SSU rRNA sequences. In addition, our analysis indicates that

improvements in eukaryotic coverage are most likely to come from targeting universally-conserved

rRNA regions, such as those containing the 515Y/V4F and the 1389F/1392R primers respectively (Fig

S22-S28).

In summary, the primer pairs with the highest environmental coverage were 515Y/926R

(universal), 515Y/806RB (prokaryote-only), and V4F/V4RB (eukaryote-only). It is worth emphasizing

that both universal primer sets have the potential to be equally or more comprehensive for *Eukarya*

versus the *Eukarya*-specific primer sets tested here. In addition, for those wishing to quantify the

abundance of eukaryotic phytoplankton using chloroplast data, the 515Y/926R primer set nearly

perfectly matches all environmental sequences. We note, however, that recognizable dinoflagellate

chloroplast rRNA sequences are almost completely missing from the metagenomic sequences (for all

primer regions, not just 515Y/926R), and thus will likely not appear in the resulting amplicons

regardless of primer choice. This is possibly due to the unusual genomic organization/subcellular

localization of chloroplast genes in dinoflagellates (25). However, dinoflagellate 18S sequences can be

recovered from the 515Y/926R primer pair (26), making it possible to reconstruct their abundance *in*

situ in combination with the chloroplast 16S. Overall, these results confirm that the careful primer

design previously reported for the 515Y/806RB/926R primers has resulted in very high overall

coverage in oceanic ecosystems. On the other hand, certain primer sets (e.g. 341F/785R) have very low

coverage due to mismatches to dominant organisms. Resulting data may thus be distorted, though we

note that removing taxa with mismatches to primers from processed data will recover accurate relative

abundances from the remaining taxa (8). This relative abundance correction approach could also be

combined with ecosystem-specific full-length 16S databases (27) to generate accurate intercomparisons

between datasets derived from different primer sets. This approach will be important to "translate" data

from different primer sets, both from legacy samples that cannot be reanalyzed and in the future as

sequencing technology / primer design inevitably evolves.

A Quantitative Intercomparison between SSU rRNA Amplicon Sequencing and

Metagenomics:

To determine whether amplicon-based strategies recover similar patterns as metagenomes,

we generated PCR amplicon sequences for a subset of the BioGEOTRACES samples (cruises GA03

and GP13, using the same DNA), and thus were able to make an "apples to apples" quantitative

comparisons between the relative abundances of organisms determined with amplicons vs.

metagenomic reads.

To do so, we directly compared the abundance of exact amplicon sequence variants (ASVs)

from the 515Y/926R primer pair with metagenomic reads from the same region of the SSU rRNA

molecule. ASVs were clustered as necessary to account for the fact that short (150bp) metagenomic

reads could in some cases only be attributed to a broader phylogenetic group (e.g. *Prochlorococcus* and

SAR11). The summed abundance of all taxonomic groups, as determined by ASVs and metagenomic

reads, was then compared using a linear least-squares regression. For 16S sequences, the strength of the

linear correlation was uniformly high across both cruise datasets (median R^2=0.98; range=0.55-1.0; Fig

3a). These patterns were generally consistent across depth for the GA03 cruise which sampled into the

mesopelagic where community composition is markedly different from the surface. We do however

note that at depths > 150m about 38% of samples had R^2 values below 0.95 (Fig 3a), vs. 9.2% for

samples at depths < 150m. These lower correspondences are unlikely to be due to mismatches to the

primer region, since predicted primer coverage is uniformly high across depth for the 515Y/926R

primer pair across depth (Fig 3b). Other factors such as DNA quality or quantity may have been

responsible for this, but we cannot rule out the possibility it involves characteristics of the sequences

between the priming regions. Regardless, these results show that amplicon sequencing and

metagenomic profiling typically have a very high quantitative correspondence with the primers tested

here (515Y/926R).

Improving Existing Primers:

In addition to quantifying mismatches, our pipeline also identifies the specific primer region

variants found in particular taxa and their coverage across datasets. This information can guide

improvements for any of the primers tested here. In the sections below, we provide three case studies

for improving primers using these metagenomic data.

To illustrate how a primer with poor predicted performance could be improved, we use the

case of 785R which has a mismatch to the dominant organism SAR11. This could be corrected by

adding a single additional degeneracy at position 7 in the primer GACTAC**N**VGGGTATCTAATCC.

However, 785R also has mismatches to a number of abundant chloroplast sequences (Fig 2), which

could be corrected with two additional degeneracies **R**A**Y**TACNVGGGTATCTAATCC. Compared to

the original primer (GACTACHVGGGTATCTAATCC), these primers would require the synthesis of a

larger number of oligonucleotide combinations (12 and 48 vs. 9 originally) but are well within the

number of combinations currently in use for other environmental amplification strategies (28).

Interestingly, despite being the worst-performing primer for marine ecosystems due to mismatches to

the dominant taxon SAR11, we predict that by adding the above degeneracies this primer would

achieve nearly perfect coverage for *Bacteria* (Fig S15-S21). This suggests the primer-binding region of

785R is highly-conserved among *Bacteria,* consistent with the initial publication describing this primer

as the best overall for prokaryotic taxa across diverse environments (10). This observation also

underscores the value of comparing primer sequences with environmental data since it uncovered a

significant, unexpected mismatch to an abundant taxon that can now be corrected.

We also investigated how primer sets that already perform well (515Y/806RB/926R) could

be further improved. To do so, we identified datasets that had significant potential for coverage

improvement (Figures S1-S28) and certain rare order-level taxa that had many mismatches or are

known to be biogeochemically-important. The 515Y primer was relatively easy to improve, given that

its performance was already very robust across diverse datasets (Fig S1-S28). An increase in

degeneracies at four positions (GTG**B**CAGCM**SY**CGCGGT**M**A) was sufficient to resolve the vast

majority of mismatches (including to rare taxa such as certain representatives of the *Patescibacteria*

a.k.a. candidate phyla radiation (CPR) bacteria) which were previously identified as taxonomic "blind

spots" in PCR amplicon analysis (5). These changes led to a primer with a degeneracy of 48, which we

term 515Yp ("p" for "pelagic").

Our data showed that the 806RB primer performed extremely well overall for 16S sequences,

but did miss a number of 16S chloroplast sequences mostly from the *Mamielliales* clade, which

includes small picoeukaryotes such as *Ostreococcus* and *Bathycoccus*. As shown in Fig S8-S14, these

taxa were still missed when allowing for up to 2 mismatches, meaning they are unlikely to be amplified

with the current primer design. Alterations would be possible, but we note that this may be an

intentional design choice, since chloroplast amplification may be undesirable in certain environments

(29) but is often very useful in euphotic zone marine water samples where it can be used to quantify

primary producer abundance. Related to this, we note that a context-dependent advantage of 806RB is

that it lacks an 18S binding site. This means that in combination with 515Y(p), nuclear 18S will not be

amplified with 806RB - a potential advantage for host-associated microbiome studies where 18S could

overwhelm targeted 16S sequences. This is, however, unlikely to occur in general for pelagic ocean

samples that are generally dominated by prokaryotic microorganisms (Fig S29), making the 926R

primer a feasible choice.

The 926R primer had mismatches to more taxa, and thus required more extensive

modifications. For example, Table S1 shows that 926R has mismatches to both *Ectothiorhodospirales*

(an uncultivated group related to purple sulfur bacteria (30)), and the *Rickettsiales* (an order that

includes mitochondrial sequences in addition to free-living organisms (31)). While less abundant

overall, we also noted mismatches in the Tara Oceans dataset to *Brocadiales*, the order containing

anammox bacteria (32). Resolving mismatches for all these three taxa as well as those having > 2%

overall relative abundance, we arrive at a new primer CCGYCAATT**HH**TTTR**R**GTT**Y**

(degeneracy=144), which could be further simplified by removing the trailing wobble base to

CCGYCAATT**HH**TTTR**R**GTT (degeneracy=72). Removing the trailing base would allow more

efficient amplification of *Rickettsiales* templates, which have a single mismatch at the 3' end of the

primer which was shown to completely prevent amplification of such sequences (9). For example, in

the BiHOBA dataset, these modifications increase *Rickettsiales* coverage from ~73% to ~90%. Since

the order *Rickettsiales* contains sequences from mitochondrial SSU rRNA, this modification will

potentially allow for more effective quantification of protist abundances, similar to how chloroplasts

quantify marine protistan phytoplankton.

266   While this new version of 926R would increase coverage for all abundant groups sampled in

the Tara/BioGEOTRACES expeditions (including eukaryotes), it would still miss some taxa important

in low-oxygen regions. For example, the *Campylobacteria* are dominant chemoautotrophs in many

sulfidic systems (33), and could be targeted by adding a single further degeneracy

(CCGYC**W**ATTHHTTTRRGTT; degeneracy = 144). Since *Campylobacteria* are known to be

abundant in certain rare low-oxygen microenvironments such as in sediment trap particles (34), they

would be important to include for a primer set used across diverse marine pelagic systems.

273   Since it is beyond the scope of this paper to provide exhaustive recommendations for each

primer set, we direct interested readers to an Open Science Foundation repository which contains all

necessary output files, scripts, and a suggested workflow for improving primers (osf.io/gr4nc/). This

should be possible for most taxa since a few mismatches typically dominated, with a long tail of rare

variants that may derive from sequencing errors or pseudogenized SSU rRNA. This information would

allow an interested reader to optimize one of the primers investigated here for particular taxa or

environments of interest.

280   Moving forward, we recommend oceanographers consider applying these two new

degenerate primers for pelagic water column surveys to maximize organismal coverage (515Yp:

GTGBCAGCMSYCGCGGTMA; 926Rp: CCGYCWATTHHTTTRRGTT [5'-3']). If proven to be as

quantitative as the original 515Y/926R primers (3, 8), they would provide an affordable,

comprehensive (all organisms from Archaea - to Zooplankton), and scaleable method to measure whole

community compositions across time and space in the world's oceans. In turn, this would allow routine

measurement of microbial community compositions alongside physical and chemical parameters on

global oceanographic surveys, helping expand our understanding of how oceanic microbes interact

with global biogeochemical cycles and respond to global climate change.

Finally, we believe a similar approach could be applied elsewhere to generate ecosystem-

specific primer sets. For example, while our revised primer set has high coverage for the vast majority

of *Archaea* and *Bacteria* in the Guaymas Basin samples, it remains unclear whether this is broadly true

for other marine sediment ecosystems. Our reproducible workflow, combined with expert curation and

broader environmental sampling could ensure that sediment taxa of interest are accurately quantified by

oligonucleotide primers, or indeed taxa from any environment where deeply-sampled metagenomes

exist.

Materials and Methods:

Evaluation of primer coverage/biases using metagenomic reads:

In order to evaluate the performance of our primer set compared to others, we used

metagenomic reads to estimate the fraction of SSU rRNA fragments that could be successfully targeted

by (i.e. perfectly match) each primer set. In brief, our pipeline retrieves SSU rRNA from fastq-

formatted shotgun metagenomic samples, removes sequences containing uninformative repeats, sorts

into four organismal categories (*Archaea, Bacteria, Cyanobacteria* (including plastidal 16S), and

Eukarya), and aligns these sequences to a group-specific reference SSU rRNA sequence (e.g.

Saccharomyces cerevisiae for *Eukarya*). Aligned sequences are then subsetted by primer region,

quality-filtered, and compared with primer sequences (Table 2) at 0, 1 and 2-mismatch thresholds (Fig

S1-S28). Retrieved reads were then classified using the SILVA 132 database (*Archaea, Bacteria,*

Eukarya) and PhytoRef (8, 35) (*Cyanobacteria* + plastidal 16S). The specific software implementation

is described fully in the supplementary material. For those wishing to inspect results further, complete

plaintext summaries of output are available online at https://osf.io/gr4nc/ in addition to summary

graphs.

These computational steps were implemented with the snakemake workflow engine (36) to

create a documented and reproducible pipeline which is freely available

(github.com/jcmcnch/MGPrimerEval). It automatically produces tabular output / summary plots and

can be applied to new samples and primers by modifying template configuration files. The specific

functioning of the three workflows are described in more detail in the supplemental material.

PCR amplification and ASV generation:

DNA used for PCR amplicons was the same DNA material used to produce metagenomes

described by Biller et al (2018) (18) except that it was diluted to 0.5 ng/µL in low-EDTA TE buffer

prior to amplification. DNA was amplified with the 515Y/926R primers (515Y:

GTGYCAGCMGCCGCGGTAA / 926R: CCGYCAATTYMTTTRAGTTT) (3, 8) using 0.5ng of DNA

template in a 25 µL reaction with a final primer concentration of 0.3mM. Primers were part of a larger

construct with Illumina adapter constructs/barcodes already included, allowing for single-step library

preparation. Thermocycling was as follows: an initial denaturation at 95°C for 120s followed by 30

amplification cycles of 95°C × 45s; 50°C × 45s; 68°C × 90s and a final elongation step at 68°C ×

300s. Laboratory methods are described in more detail at

https://dx.doi.org/10.17504/protocols.io.vb7e2rn. Both 16S and 18S mock communities (staggered and

even versions of both (3, 8)) were included in the sequencing run as quality control. Amplicons were

sequenced with HiSeq 2x250 RapidRun technology at the USC sequencing center in a run pooled with

shotgun metagenomic samples. Raw basecall data was demultiplexed with bcl2fastq (-r 20 -p 20 -w 32

--barcode-mismatches 0) using the 6 basepair reverse indices and subsequently with the forward 5

basepair barcode with cutadapt according to the templates provided at github.com/jcmcnch/demux-

notes. Raw data from this step of the pipeline is available on NCBI under the umbrella Bioproject

PRJNA659851 and three sub-projects PRJNA658608 (controls), PRJNA658384 (GA03), and

PRJNA658385 (GP13).

Sequence data was analyzed with a custom *in-silico* pipeline available at

github.com/jcmcnch/eASV-pipeline-for-515Y-926R and based on qiime2 2019.4 (35). Briefly, 16S and

18S sequences were first partitioned using a custom 16S/18S-specific databases derived from the

SILVA 132 and PR2 (7, 36) which is available at osf.io/e65rs/. After partitioning, primers were

removed from amplicons allowing up to 20% mismatches to the primer, and raw sequences were

imported into qiime2 and denoised with DADA2 (6). For 18S sequences that do not overlap, they were

denoised as single-end reads by trimming to a consistent length (forward=220bp, reverse=180bp) and

concatenating the two reads. Taxonomy was assigned using a naïve Bayesian classification algorithm

with the SILVA 132 database subsetted to the amplicon region as a reference (37). Sequences identified

as chloroplasts were re-annotated with PhytoRef (38), and 18S sequences were additionally annotated

with PR2 (36). A step-by-step protocol for these informatic steps is available at

https://dx.doi.org/10.17504/protocols.io.vi9e4h6. After denoising and annotation, biom tables were

exported with taxonomy as tsv files and converted to relative abundances which were used as input for

the metagenome-amplicon comparison.

Acknolwedgements: This work was supported by the Simons Collaboration on

Computational Biogeochemical Modeling of Marine Ecosystems (CBIOMES) grant 549943 to JAF, the

Gordon and Betty Moore Foundation Marine Microbiology Initiative grant 3779 and NSF grant

OCE1737409. We gratefully acknowledge the effort of the scientists who collected the

BioGEOTRACES field samples analyzed here: Christel Hassler, Debbie Hulston, and Elizabeth Maas

(GP13) and Jeremy Jacquot (GA03). We also acknowledge the efforts of Keven Dooley and Madeline

Williams who conducted DNA extractions, which was funded in part by grants to Sallie W. Chisholm at

MIT that enabled the collection and processing of DNA samples from GEOTRACES cruises. These

include the Simons Foundation (Life Sciences Project Award ID 337262, SCOPE Award ID 329108),

the Moore Foundation (Grant IDs GBMF495 and GBMF4511), and the National Science Foundation

(OCE-1153588, OCE-1356460, and DBI-0424599). We thank S.W.C. for donating the samples and

supporting parts of the study.

References Cited:

1.   L. R. Thompson, *et al.*, A communal catalogue reveals Earth's multiscale microbial diversity. *Nature* **551**, 457–463 (2017).

2.   A. Apprill, S. McNally, R. Parsons, L. Weber, Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. *Aquatic Microbial Ecology* **75**, 129–137 (2015).

3.   A. E. Parada, D. M. Needham, J. A. Fuhrman, Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. *Environ Microbiol* **18**, 1403–1414 (2016).

4.   W. Walters, *et al.*, Improved Bacterial 16S rRNA Gene (V4 and V4-5) and Fungal Internal Transcribed Spacer Marker Gene Primers for Microbial Community Surveys. *mSystems* **1** (2016).

5.   E. A. Eloe-Fadrosh, N. N. Ivanova, T. Woyke, N. C. Kyrpides, Metagenomics uncovers gaps in amplicon-based detection of microbial diversity. *Nature Microbiology* **1**, 15032 (2016).

6. B. J. Callahan, *et al.*, DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods* **13**, 581–583 (2016).

7. C. Quast, *et al.*, The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucl. Acids Res.* **41**, D590–D596 (2013).

8. Y.-C. Yeh, J. C. McNichol, D. M. Needham, E. B. Fichot, J. A. Fuhrman, Comprehensive single-PCR 16S and 18S rRNA community analysis validated with mock communities and denoising algorithms. *bioRxiv*, 866731 (2019).

9. E. K. Wear, E. G. Wilbanks, C. E. Nelson, C. A. Carlson, Primer selection impacts specific population abundances but not community dynamics in a monthly time-series 16S rRNA gene amplicon analysis of coastal marine bacterioplankton. *Environmental Microbiology* **20**, 2709–2726 (2018).

10. A. Klindworth, *et al.*, Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucl. Acids Res.*, gks808 (2012).

11. C. S. Fortunato, L. Herfort, P. Zuber, A. M. Baptista, B. C. Crump, Spatial variability overwhelms seasonal patterns in bacterioplankton communities across a river to ocean gradient. *The ISME Journal* **6**, 554–563 (2012).

12. T. Stoeck, *et al.*, Multiple marker parallel tag environmental DNA sequencing reveals a highly complex eukaryotic community in marine anoxic water. *Molecular Ecology* **19**, 21–31 (2010).

13. S. Balzano, E. Abs, S. Leterme, Protist diversity along a salinity gradient in a coastal lagoon. *Aquatic Microbial Ecology* **74**, 263–277 (2015).

14. D. Wilkins, E. van Sebille, S. R. Rintoul, F. M. Lauro, R. Cavicchioli, Advection shapes Southern Ocean microbial assemblages independent of distance and environment effects. *Nature Communications* **4**, 2457 (2013).

15. L. A. Amaral-Zettler, E. A. McCliment, H. W. Ducklow, S. M. Huse, A Method for Studying Protistan Diversity Using Massively Parallel Sequencing of V9 Hypervariable Regions of Small-Subunit Ribosomal RNA Genes. *PLOS ONE* **4**, e6372 (2009).

16. S. Pesant, *et al.*, Open science resources for the discovery and analysis of Tara Oceans data. *Sci Data* **2**, 1–16 (2015).

17. S. G. Acinas, *et al.*, Metabolic Architecture of the Deep Ocean Microbiome. *bioRxiv*, 635680 (2019).

18. S. J. Biller, *et al.*, Marine microbial metagenomes sampled across space and time. *Scientific Data* **5**, 180176 (2018).

19. D. M. Karl, M. J. Church, Microbial oceanography and the Hawaii Ocean Time-series programme. *Nat Rev Micro* **12**, 699–713 (2014).

20. D. K. Steinberg, *et al.*, Overview of the US JGOFS Bermuda Atlantic Time-series Study (BATS): a decade-scale look at ocean biology and biogeochemistry. *Deep Sea Research Part II: Topical Studies in Oceanography* **48**, 1405–1447 (2001).

21. B. Nowinski, *et al.*, Microbial metagenomes and metatranscriptomes during a coastal phytoplankton bloom. *Sci Data* **6**, 1–7 (2019).

22. E. T. Sieradzki, J. C. Ignacio-Espinoza, D. M. Needham, E. B. Fichot, J. A. Fuhrman, Dynamic marine viral infections and major contribution to photosynthetic processes shown by spatiotemporal picoplankton metatranscriptomes. *Nat Commun* **10**, 1–9 (2019).

23. A. K. Hawley, *et al.*, A compendium of multi-omic sequence information from the Saanich Inlet water column. *Sci Data* **4**, 1–11 (2017).

24. N. Dombrowski, K. W. Seitz, A. P. Teske, B. J. Baker, Genomic insights into potential interdependencies in microbial hydrocarbon and nutrient cycling in hydrothermal sediments. *Microbiome* **5**, 106 (2017).

25. T. Laatsch, S. Zauner, B. Stoebe-Maier, K. V. Kowallik, U.-G. Maier, Plastid-Derived Single Gene Minicircles of the Dinoflagellate *Ceratium horridum* Are Localized in the Nucleus. *Mol Biol Evol* **21**, 1318–1322 (2004).

26. D. M. Needham, J. A. Fuhrman, Pronounced daily succession of phytoplankton, archaea and bacteria following a spring bloom. *Nature Microbiology* **1**, 16005 (2016).

27. M. S. Dueholm, *et al.*, Generation of Comprehensive Ecosystem-Specific Reference Databases with Species-Level Resolution by High-Throughput Full-Length 16S rRNA Gene Sequencing and Automated Taxonomy Assignment (AutoTax). *mBio* **11** (2020).

28. J. C. Gaby, D. H. Buckley, A Comprehensive Evaluation of PCR Primers to Amplify the *nifH* Gene of Nitrogenase. *PLoS One* **7** (2012).

29. S. Thijs, *et al.*, Comparative Evaluation of Four Bacteria-Specific Primer Pairs for 16S rRNA Gene Surveys. *Front. Microbiol.* **8** (2017).

30. A. Oren, "The Family *Ectothiorhodospiraceae*" in *The Prokaryotes*, (Springer, Berlin, Heidelberg, 2014), pp. 199–222.

31. D. N. Fredricks, "Introduction to the *Rickettsiales* and Other Intracellular Prokaryotes" in *The Prokaryotes*, M. D. P. Dr, S. Falkow, E. Rosenberg, K.-H. Schleifer, E. Stackebrandt, Eds. (Springer New York, 2006), pp. 457–466.

32. L. Medina Faull, P. Mara, G. T. Taylor, V. P. Edgcomb, Imprint of Trace Dissolved Oxygen on Prokaryoplankton Community Structure in an Oxygen Minimum Zone. *Front. Mar. Sci.* **7** (2020).

33. B. J. Campbell, A. S. Engel, M. L. Porter, K. Takai, The versatile ε-proteobacteria: key players in sulphidic habitats. *Nature Reviews Microbiology* **4**, 458–468 (2006).

34. D. Boeuf, *et al.*, Biological composition and microbial dynamics of sinking particulate organic matter at abyssal depths in the oligotrophic open ocean. *PNAS* **116**, 11824–11832 (2019).

35.   E. Bolyen, *et al.,* Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nature Biotechnology* **37**, 852–857 (2019).

36.   L. Guillou, *et al.,* The Protist Ribosomal Reference database (PR2): a catalog of unicellular eukaryote Small Sub-Unit rRNA sequences with curated taxonomy. *Nucleic Acids Res* **41**, D597–D604 (2013).

37.   N. A. Bokulich, *et al.,* Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome* **6**, 90 (2018).

38.   J. Decelle, *et al.,* PhytoREF: a reference database of the plastidial 16S rRNA gene of photosynthetic eukaryotes with curated taxonomy. *Molecular Ecology Resources* **15**, 1435–1445 (2015).

Figure 1: Distribution of metagenomic samples used in this study. BioGEOTRACES samples that were used in the metagenome/amplicon intercomparison are noted with open circles (GA03 - N. Atlantic; GP13 - S. Pacific).

Dataset name (**abbreviation**)	Sample and environment description	Sample size fraction(s)	Sample volume (L)	Reference(s)	# samples tested	Total # SSU rRNA sequences	Median filtered sequences per sample/group/primer/direction**			
							Archaea	Bacteria	Cyanobacteria & Chloroplast	Eukarya
BioGEOTRACES / HOT / BATS (**BiHOBA**)	Mostly euphotic and upper mesopelagic ocean waters; primarily oligotrophic	>0.2µm	0.1	18-20	610	3.16E+07	36.9	256.9	46.3	55.5
Tara Oceans (**TARA**)	Euphotic, deep chlorophyll maximum, and mesopelagic samples	All with upper size fraction <3µm*	≥100	16	458	1.32E+08	109.1	1148	141.7	155.4
Malaspina Deep Metagenomes (**MSPINA**)	Deep ocean metagenomes from samples collected >2000m in depth	0.2 - 0.8 µm (FL), 0.8 - 20 µm (PA)	120	17	58	3.46E+06	26.3	158.4	NA	32.6
MBARI bloom metagenome (**MBARI**)	Coastal surface seawater sampled during phytoplankton bloom across 52 days	0.22–5.0 µm	up to 1	21	84	1.32E+08	25.7	667.7	52.4	68.3
SPOT / Catalina metagenome (**SPOT**)	Surface seawater collected from the San Pedro Ocean Time-Series and Catalina island across 4 seasons	0.22-1 µm	20	22	8	5.04E+05	16.9	434.2	186.2	201.7
Saanich Inlet Metagenomes (**Saanich**)	Depths spanning the oxycline (10, 100, 120, 135, 150, and 200 m)	0.22-2.7 µm	1 or 20	23	87	1.37E+07	96.8	662.6	88.9	21.5
Guaymas sediment metagenomes (**GuaySed**)	Total DNA from > 10g hydrothermally-influenced sediment from 0-15cm depth	NA, whole sediment	NA, 10g sediment	24	4	1.95E+06	1047.3	3065.6	93	433.7

*<0.2µm; 0.2-1.6µm; 0.2-3µm; 0.8-3µm; **groups={1: Archaea,2:Bacteria less Cyanobacteria+Plastids,3:Chloroplast+Plastid,4:Eukarya}, direction=forward and reverse paired-end reads (analyzed separately)

374   Table 1: Data used in this study. SRA/ENA Accessions available on github.com/jcmcnch/MGPrimerEval in config files.

Primer	SSU rRNA region targeted	Organismal coverage	Primer sequence (5' - 3')	Reference(s)
27F	V1-V2	16S only	AGRGTTTGATYMTGGCTCAG	11
338R	V1-V2	16S only	TGCWGCCWCCCGTAGGWGT	11
341F	V3-V4	16S (Bacteria-only)	CCTACGGGNGGCWGCAG	10
785R	V3-V4	16S only	GACTACHVGGGTATCTAATCC	10
V4F	V4	18S only	CCAGCASCYGCGGTAATTCC	12
V4R	V4	18S only	ACTTTCGTTCTTGATYRA	12
V4RB	V4	18S only	ACTTTCGTTCTTGATYRR	13
515Y	V4/V4-V5	16S and 18S	GTGYCAGCMGCCGCGGTAA	2-4
806RB	V4	16S only	GGACTACNVGGGTWTCTAAT	2-4
926R	V4-V5	16S and 18S	CCGYCAATTYMTTTRAGTTT	2-4
926wF	V6-V8	16S and 18S	AAACTYAAAKGAATTGRCGG	14
1392R	V6-V8	16S and 18S	ACGGGCGGTGTGTRC	14
1389F	V9	18S only	TTGTACACACCGCCC	15
1510R	V9	18S only	CCTTCYGCAGGTTCACCTAC	15

377   Table 2: Primers tested in this study.

Figure 2: Violin plot with inlaid boxplot showing coverage for oligonucleotide PCR primers based on comparisons with globally-distributed oceanic metagenomes. The shapes of the violin plots illustrate the distribution of primer coverage across all samples. Note that the "All SSU rRNA" panel represents all of the SSU rRNA targeted by a particular primer set (e.g. for 1389F/1510R this includes only Eukarya *while for 926wF/1392R it includes all 4 taxonomic groups). Data are corrected for predicted taxonomic overlap between forward and reverse primers. Coordinates refer to locations of primer alignments to the* Escherichia coli *16S rRNA gene (strain K-12; substrain MG1655).*

Fig 3: A) Metagenome-amplicon quantitative correspondence across all depths for the GA03 cruise (BioGEOTRACES), showing the R^2 value of the plot of the relative abundances of amplicon sequence variants (ASVs; 515Y/926R primers) vs relative abundances of metagenomic reads from the same organisms (see text for details of comparisons). B) The fraction of exact matches of the 515Y and 926R amplicon primers compared to metagenomic reads of the priming regions for each sample, summarized as the "worst-case" coverage for the combined primers (i.e. 1 - (missed$_{fwd}$ + missed$_{rev}$)).

Supplementary Information:

Supplementary Methods:

Software Implementation: Software dependencies are handled by conda implemented in

snakemake (1). Examples of how the pipeline can be run are available at

github.com/jcmcnch/MGPrimerEval in the "runscripts" folder. Both the "classify" and "compare"

workflows depend on the output of the "compute" workflow. The "classify" workflow is for detailed

exploration of the taxonomy of retrieved sequences and the "compare" workflow allows for

intercomparisons between SSU rRNA amplicon sequence variants (ASVs) and metagenomic SSU

rRNA from the same amplicon region.

Compute workflow ("Snakefile-compute.smk"; Fig S30): This portion of the analysis

calculates the coverage of primer sequences for metagenomic samples specified in a configuration file.

Results are calculated as the fraction of total SSU rRNA reads matching the primer at 0, 1, and 2-

mismatches for 4 taxonomic partitions as described below. A rule graph summarizing the steps is

shown in Figure S30. Briefly, fastq reads are first cleaned using fastp's (v 0.14.1) default parameters

and trimming 3 trailing and leading bases (2). To retrieve SSU rRNA sequences from metagenomes, we

used the phyloFlash (v 3.3b1) package (3) which uses bbmap (v 38.69) (4) and a curated version of

SILVA132 (5) to retrieve both 16S and 18S sequences. Since retrieved sequences contained a small

fraction of low-complexity repeats that were not homologous to the SSU rRNA, we used komplexity (v

0.3.6) (6) to remove these likely spurious hits.

SSU rRNA sequences thus identified were then split into 4 different pools (Eukaryota,

Archaea, Bacteria (non-cyanobacterial), and Cyanobacteria (including chloroplasts)) using bbsplit.sh

v38.22 (4) comparing against curated subsets of the SILVA/PR2 reference databases (5, 7) (sequences

and a template script for creating databases are available at osf.io/e65rs/). Partial SSU rRNA sequences

were then aligned using pyNAST v1.2.2 (8) to group-specific references available as part of the github

repository. Primer-specific subsets were then parsed from the alignment using python/biopython (9),

only outputting sequences that overlap the primer region plus 5 leading or trailing bases. The 1-based

coordinates of each primer region (for each organismal reference) must be supplied by the user; several

widely-used primers are already included in template configuration files.

To place all reads in the correct orientation, fastq files were reverse-complemented based on

information found in pyNAST alignment headers. cutadapt (version 1.18) (10) was used to identify

primer regions in fastq sequences with 0, 1, 2, and 6-mismatches to the primer. Only fastq sequences

with 6 or fewer mismatches to the primer were further analyzed to exclude any spurious matches

caused by pyNAST misalignments. Based on primer comparisons to references, this is likely to include

even highly divergent taxa so long as they have a true primer binding site (NB: this is not true for some

primers designed to specifically target a particular group - e.g. the 806RB primer does not contain a

binding site for eukaryotic sequences). Primer coverage at each mismatch cutoff was then calculated

using only sequences where all bases in the primer region plus 5 leading and trailing bases had a phred

score of > 30. Plots shown in the main and supplemental text were generated with python/cartopy (11),

ggplot2 (12), veusz (13), and snakemake (1) (Fig 1; Fig 2, Fig S1-S29; Fig 3; Fig S31-S32

respectively).

Classify workflow ("Snakefile-classify.smk"; Fig S31):

In order to determine which taxa were missed by a particular primer set and provide

recommendations about how to further improve existing primers' performance, we implemented an

optional subsequent workflow to taxonomically classify reads and provide summary statistics for each

group. Briefly, full-length fastq files from mismatched taxa are classified with VSEARCH 2.10.4 (14)

and the SINTAX algorithm (15) against the SILVA132 nr99 database (5) or PhytoRef (16) databases

(for cyanobacteria/plastids) that had been converted into ".udb" format. The same procedure was

followed for matches except a maximum of 5,000 reads were randomly subsampled using

"filterbyname.sh" (4) to reduce computation time.

Summary statistics and output files were then generated for each unique combination of

group, primer, and mismatch threshold (e.g. EUK, 926R, 0-mismatch) at order-level resolution since

informative taxonomic annotations could generally not be made at finer taxonomic levels. Summaries

contain the fraction of each taxon not matching the primer at a given mismatch threshold, as well as the

fraction of total sequences represented by these mismatches. In addition, multiple sequence alignments

of mismatches and their relative abundances are generated, with summary plots automatically

generated showing the relative abundances of particular mismatches both for the whole dataset and

subsetted by individual taxa. In addition, the same output files are generated for specific order-level

taxa, allowing an investigator to see whether a given primer covers a particular taxon of interest. For

the datasets analyzed in this study, raw output files/plots can be downloaded at https://osf.io/gr4nc/.

Our pipeline generates coverage information at the individual primer level since

metagenomic reads will rarely/never cover the entire amplicon region (depending on lengths of

sequence reads and the amplicon region). However, it is advantageous to present results in terms of

predicted coverage for the primer set. If the mismatches to the forward and reverse primer were

completely non-overlapping (i.e. the mismatches were to completely different organisms) then the

mismatched fractions would be simply additive ("worst case" as mentioned in main text and shown in

Fig 3). In contrast, if organisms overlapped between the forward and reverse primer, the largest

mismatched fraction for that organism could be used. We used the taxonomic assignments discussed

above to estimate this fraction of overlapping sequences ("rule normalize_summaries" and its

dependent rules/scripts) and thus correct the primer coverage values as shown in Fig 2.

Compare workflow ("Snakefile-compare2tags-16s.smk" and "Snakefile-compare2tags-

18s.smk"):

This workflow compares the abundance of taxa quantified by universal 16S and 18S

amplicons using the 515Y/926R primers with abundances of taxa from metagenomic SSU rRNA in the

same region of the molecule targeted by the PCR primers. The primer region is targeted by subsetting

pyNAST alignments by primer coordinates. Only reads with at least 100bp overlap with the amplicon

region are retained, and subsetted reads from the three prokaryotic categories are merged for the 16S

comparison. For 18S reads that do not overlap (since the amplicon region is longer than the 2x250

sequencing length employed here), they were subsetted to the trim length of the partial amplicons (e.g.

220bp for forward read, 180 bp for reverse read).

These metagenomic sequences were then compared using BLASTn (17) (minimum 97%

similarity, requiring 100% overlap) against a sample-specific database of exact amplicon sequence

variants (ASVs) generated with DADA2 (18) that had a nonzero abundance in that particular sample.

In order to make a direct comparison between the two data sources, the ASVs were clustered for each

sample based on BLAST results with networkx (19) to account for the resolution mismatch between

short metagenomic reads and longer, denoised amplicons. Total relative abundances of each cluster

were plotted using seaborn (https://seaborn.pydata.org/) and sample-sample correspondence was

evaluated using least squared linear regression with the stats.linregress function in scipy (20). The R^2

value of this regression was used to evaluate general trends in MG-ASV sample-sample

correspondence, and was plotted against both environmental covariates and the "worst-case" (simply

additive) predicted primer performance (Fig 3b).

Amplicon Analysis Pipeline Supplementary Information:

SSU rRNA partitioning was carried out with the database mentioned in the main text

(osf.io/e65rs/) using bbsplit.sh (4). These databases were constructed from SILVA132 (5) and PR2 (7)

by first clustering with CDHIT (21) at 95% identity to reduce total data volume and subsequently

curated to remove sequences that were extremely long or short relative to the average sequence length.

Demultiplexing and trimming steps were carried out with cutadapt (10), and trimming prior to

denoising (for 18S sequences) was carried out with bbduk.sh (4).

Supplementary Results:

Technical evaluation of pipeline:

After retrieval, quality-filtering, and subsetting to the primer region, dozens to hundreds of

rRNA sequences were recovered for each group, sample, read direction (fwd & rev reads), and primer

combination (Table 1). Considering the pelagic ocean datasets introduced above and discussed in detail

below (BiHoBa, TARA, MSPINA, MBARI, SPOT), we retrieved between 158-1148 primer-matching

regions for *Bacteria* for each combination (medians within datasets; Table 1), whereas lower numbers

were retrieved for the remaining three groups of *Archaea*, *Cyanobacteria*+Plastids and *Eukarya* (on

average 17-109, 46-186 and 32-202 respectively) most likely due to their lower relative abundances in

the environmental DNA. Among the primers, smaller number of sequences were identified for the two

primers located near the end of the SSU rRNA molecule. For example, for Tara samples, a median of

455 sequences were retrieved per sample for 27F (close to end of molecule) for *Bacteria* vs. 1299 per

sample for 338R (middle of molecule). Similarly, 58 eukaryotic sequences were retrieved per sample

with 1510R vs. 136 per sample for 1389F.

123   With the exception of regions near the end of the SSU rRNA molecule, sequence coverage

was mostly evenly distributed across the alignments to group-specific references. Some relatively small

regions had lower coverage, most likely due to their absence in dominant environmental taxa. For

example, the SAR11 isolate HTCC1062 is missing nucleotides (relative to the *E. coli* reference)

corresponding to small portions of loops in the V1, V2, V3, and V9 regions (22). We note also that

phyloFlash retrieved approximately double the total number of 18S sequences compared to a graftM

(23) package based on an alignment of the greengenes database (data.ace.uq.edu.au/public/graftm/7/)

which was employed in an earlier version of our pipeline. In addition, the coverage across the *S.*

cerevisae 18S reference was much more even with phyloFlash-retrieved 18S rRNA vs. graftM. No

such bias was noted for 16S sequences, suggesting this graftM package may be appropriate for 16S

retrieval but not 18S.

135   **Primer Limitations and Potential Improvements:**

137   Table S1 shows a summary of the top 3 most dominant mismatched orders for each primer

and group for the BiHoBa dataset. These samples mostly represent surface and upper mesopelagic

environments across all organismal size classes (Table 1), making it useful for exploring general

patterns of primer performance for oceanic ecosystems. Among the most prominent mismatches to

Archaea are those between *Nitrosopumilales* MGI *Archaea* and primers 341F or 1389F. Neither of

these primers were specifically designed for capturing archaeal sequences but our results show they

could be modified to include most *Archaea* as shown in Figs S1-S7 for 341F, and likely also for 1389F

given that a universal primer with high coverage exists for the same general region (1392R). For other

primers, the dominant mismatches include abundant groups such as MGII, and MGIII but we also note

a smaller number of mismatches from the poorly-described *Aigarchaeales* and *Woesearchaeia.*

For *Bacteria*, many of the abundant mismatches were already discussed in the main text.

Other patterns of particular note are mismatches to abundant members of the SAR86 clade and

Flavobacteriales for 341F/338R primers. Together, these groups represent over 4% of the total

sequences, meaning they would likely be desirable to correct in the future. Readers should note that the

dominant taxa in Table S1 are the most abundant sequences, and there are a number of rare bacterial

taxa that have very low coverage. These include *Poribacteria*, a number of members of the candidate

phyla radiation or CPR, *Rubrobacterales* (*Actinobacteria*), and *Chlamydiales*. Interestingly, some

members of these taxa (e.g. *Poribacteria*, CPR) had more than one mismatch to the 926R primer set,

meaning they are likely very evolutionarily-divergent from other organisms and may require a more

extensive primer redesign to quantify them fully. However, we note that these organisms are very rare

in these datasets, so their absences are unlikely to bias overall community profiles in oxic marine

environments and that the additional degeneracies proposed in the main text will increase coverage for

some of these taxa.

With respect to the cyanobacterial/chloroplast fraction, while the 515Y/926R primer pair

achieves nearly perfect coverage, the other primers generally have significant mismatches to one or

more abundant chloroplast groups, including the *Mamelliophyceae*, *Prasinophyceae*, and

Prymnesiophyceae. This means that care must be used in interpreting chloroplast results from some

popular primer sets which may miss important taxa.

Among eukaryotic (nuclear 18S) sequences, the most commonly mismatched taxa included

Dinoflagellata, Discicristata, Protalveolata, Ochrophyta, Charophyta, and *Metazoa.* As noted above,

the universal primer sets appear to perform nearly as well or better versus *Eukarya*-specific primers.

Improvements described in the main text do increase 18S coverage for the 515Y/926R primers, but a

fully comprehensive 18S primer may be less feasible since primer binding regions have a much higher

sequence diversity versus 16S. This may be due to large copy numbers of SSU rRNA, some of which

may become pseudogenized and undergo random evolutionary drift (24). In practical terms, this means

that some of these more rare variants will inevitably be missed by primers since creating an overly-

degenerate primer could cause problems with non-specific amplification during PCR. The same may be

true for some mitochondrial 16S genes that have become pseudogenized after being incorporated into

the host nuclear genome (25).

Similarly, one of the most challenging datasets for improving our primers came from the

Guaymas basin (Table 1). Anoxic sediment environments are known to host unusual *Archaea* that are

now recognized to be much more diverse than previously thought and are missed by some PCR primers

(25). For 926R, we identified another degeneracy that combined with an additional 3' trim would target

many of these rare *Archaea* (CCGYCWATTHH**Y**TTRRGT). However, this would increase total

degeneracy to 288, and it is not clear whether this level of degeneracy would be desirable. Instead, it

may be more effective to try to use our general approach to design an environment-specific primer for

sediment microbiomes.

A Practical Consideration for the use of the 515Y/926R primer set:

One concern about using truly universal primers is the amplification of non-microbial

sequences, e.g. in the case of host-associated microbiomes. This has led to the design of prokaryotic-

specific primer sets (e.g. 515Y/806RB), that intentionally do not amplify 18S sequences. However,

universal primers retain the advantage of quantifying a broad swathe of the entire biological

community together. Therefore, such primers could be potentially be very useful if 18S sequences do

not overwhelm the SSU rRNA sequences in typical samples.

To test this, we quantified the total fraction of 18S (out of the total 16S + 18S) for the

unfractionated (>0.2µM) BiHoBa metagenomes. For these data, we show 18S sequences rarely exceed

~1/3 of the total SSU rRNA sequences (Fig S29), indicating that universal primers will still be useful

even when whole seawater samples are analyzed, providing quantification of both prokaryotic and

eukaryotic communities simultaneously. The fraction of eukaryotic SSU rRNA relative to total 16S +

18S averaged 19.5 +/- 0.09 % (range = 0.024 - 0.7). Samples with large fractions of eukaryotes (>0.25)

were dominated by *Metazoa*, mainly composed of taxa such as *Arthropoda, Platyhelminthes,* and

Hydrozoa. These animal-derived sequences have the potential to overwhelm the signal from other

environmental eukaryotes, especially for unfractionated samples. This can be alleviated to some extent

by increasing overall sequence coverage such that sufficient non-metazoan 18S sequences can also be

recovered.

Supplemental References Cited:

1.   J. Köster, S. Rahmann, Snakemake—a scalable bioinformatics workflow engine. *Bioinformatics* **28**, 2520–2522 (2012).

2.   S. Chen, Y. Zhou, Y. Chen, J. Gu, fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

3.   H. R. Gruber-Vodicka, B. K. B. Seah, E. Pruesse, phyloFlash – Rapid SSU rRNA profiling and targeted assembly from metagenomes. *bioRxiv*, 521922 (2019).

4.   B. Bushnell, BBMap short read aligner. *https://sourceforge.net/projects/bbmap/*.

5.   C. Quast, *et al.*, The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucl. Acids Res.* **41**, D590–D596 (2013).

6.   E. Clarke, komplexity: A method of assessing sequence complexity based on kmer frequencies. *https://github.com/eclarke/komplexity*.

7.   L. Guillou, *et al.*, The Protist Ribosomal Reference database (PR2): a catalog of unicellular eukaryote Small Sub-Unit rRNA sequences with curated taxonomy. *Nucleic Acids Res* **41**, D597–D604 (2013).

8.   J. G. Caporaso, *et al.*, PyNAST: a flexible tool for aligning sequences to a template alignment. *Bioinformatics* **26**, 266–267 (2010).

9.   P. J. A. Cock, *et al.*, Biopython: freely available Python tools for computational molecular biology and bioinformatics. *Bioinformatics* **25**, 1422–1423 (2009).

10.   M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**, 10–12 (2011).

11.   Met Office, *Cartopy: a cartographic python library with a matplotlib interface* (2010).

12.   H. Wickham, *ggplot2: Elegant Graphics for Data Analysis* (Springer-Verlag, 2009) https:/doi.org/10.1007/978-0-387-98141-3.

13.   J. Sanders, Veusz: A Scientific Plotting Package. *https://github.com/veusz/veusz*.

14. T. Rognes, T. Flouri, B. Nichols, C. Quince, F. Mahé, VSEARCH: a versatile open source tool for metagenomics. *PeerJ* **4**, e2584 (2016).

15. R. C. Edgar, SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS sequences. *bioRxiv*, 074161 (2016).

16. J. Decelle, *et al.*, PhytoREF: a reference database of the plastidial 16S rRNA gene of photosynthetic eukaryotes with curated taxonomy. *Molecular Ecology Resources* **15**, 1435–1445 (2015).

17. S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment search tool. *J. Mol. Biol.* **215**, 403–410 (1990).

18. B. J. Callahan, *et al.*, DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods* **13**, 581–583 (2016).

19. A. A. Hagberg, D. A. Schult, P. J. Swart, Exploring Network Structure, Dynamics, and Function using NetworkX in *Proceedings of the 7th Python in Science Conference*, G. Varoquaux, T. Vaught, J. Millman, Eds. (2008), pp. 11–15.

20. P. Virtanen, *et al.*, SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nature Methods* **17**, 261–272 (2020).

21. W. Li, A. Godzik, CD-HIT: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

22. P. Yarza, *et al.*, Uniting the classification of cultured and uncultured bacteria and archaea using 16S rRNA gene sequences. *Nat Rev Micro* **12**, 635–645 (2014).

23.   J. A. Boyd, B. J. Woodcroft, G. W. Tyson, GraftM: a tool for scalable, phylogenetically informed classification of genes within metagenomes. *Nucleic Acids Res* **46**, e59–e59 (2018).

24.   S. R. Santos, R. A. Kinzie, K. Sakai, M. A. Coffroth, Molecular Characterization of Nuclear Small Subunit (ISS)-rDNA Pseudogenes in a Symbiotic Dinoflagellate (Symbiodinium, Dinophyta). *Journal of Eukaryotic Microbiology* **50**, 417–421 (2003).

25.   D. Bensasson, D.-X. Zhang, D. L. Hartl, G. M. Hewitt, Mitochondrial pseudogenes: evolution's misplaced witnesses. *Trends in Ecology & Evolution* **16**, 314–321 (2001).

Table S1: The fraction of total identified sequences mismatched for the top 3 most numerically abundant mismatched orders in the BioHOBA dataset (when allowing 0 mismatches). Primers that have "NA" for particular groups often represents an intentional design choice to avoid off-target amplification.

Primer	Archaea	Bacteria (without Cyanobacteria and plastid 16S)	Cyanobacteria (including plastid 16S)	Eukarya (18S nuclear SSU rRNA only)
27F	NA, no apparent primer binding site	Rickettsiales (0.008), Ectothiorhodospirales (0.008), SAR11 clade (0.002)	Prasinococcales (0.011), Cyanobactera;Subsection I (0.003), Chrysophyceae-Synurophyceae (0.002)	NA, no apparent primer binding site
338R	NA, no apparent primer binding site	SAR86 clade (0.027), Flavobacteriales (0.015), SAR11 clade (0.009)	Prymnesiophyceae (0.123), Bacillariophyta (0.006), Prasinococcales (0.005)	NA, no apparent primer binding site
341F	Nitrosopumilales (0.51), MG II (0.41), MG III (0.07)	SAR86 clade (0.028), Flavobacteriales (0.016), SAR11 clade (0.016)	Prymnesiophyceae (0.121), Bacillariophyta (0.006), Prasinococcales (0.005)	NA, 5-mismatch to reference
785R	MG II (0.003), Nitrosopumilales (0.002), MG III (0.0004)	SAR11 clade (0.40), Marinimicrobia (SAR406_clade); unclassified order/class (0.006), Verrucomicrobiae;Arctic97B-4 (0.003)	Mamiellophyceae (0.103), Prymnesiophyceae (0.006), Cyanobacteria;SubsectionI (0.001)	NA, 3-mismatch to reference
V4F	NA, 5-mismatch to reference	NA, 3-mismatch to reference	NA, 4-mismatch to reference	Dinoflagellata (0.022), Metazoa (Animalia) (0.010), Protalveolata (0.007)
V4R	NA, 8-mismatch to reference	NA, 7-mismatch to reference	NA, many mismatches to reference	Dinoflagellata (0.081), Discicristata (0.027), Protalveolata (0.026)
V4RB	NA, 8-mismatch to reference	NA, 7-mismatch to reference	NA, many mismatches to reference	Dinoflagellata, (0.040), Discicristata (0.013), Protalveolata (0.012)
515Y	Nitrosopumilales (0.003), MG II (0.002), MG III (0.0003)	Rickettsiales (0.002), SAR11 clade (0.001), Chlamydiales (0.0006)	Cyanobacteria;Subsection I (0.003), Prymnesiophyceae (0.001), Pelagophyceae (0.000)	Dinoflagellata (0.021), Discicristata (0.014), Metazoa (Animalia) (0.010)
806RB	Nitrosopumilales (0.002), MG II (0.002), Aigarchaeales (0.001)	Marinimicrobia (SAR406_clade); unclassified order/class (0.006), Flavobacteriales (0.005), SAR202 clade (0.002)	Mamiellophyceae (0.108), Prymnesiophyceae (0.014), Prasinophyceae (0.013)	NA, many mismatch to reference
926R	Nitrosopumilales (0.016), MG II (0.001), Woesearchaeia; unclassified order (0.0007)	Ectothiorhodospirales (0.007), Rickettsiales (0.004), SAR11 clade (0.002)	Cyanobacteria;Subsection I (0.003), Prymnesiophyceae (0.000) Dictyochophyceae (0.000)	Dinoflagellata (0.015), Protalveolata (0.012), Metazoa (Animalia) (0.010)
926wF	Nitrosopumilales (0.016), MG II (0.001), Woesearchaeia; unclassified order (0.0007)	Rhodospirillales (0.041), Ectothiorhodospirales (0.007), Rickettsiales (0.004)	Cyanobacteria;Subsection I (0.003), Prymnesiophyceae (0.000) Dictyochophyceae (0.000)	Metazoa (Animalia) (0.040), Ochrophyta (0.033), Dinoflagellata (0.016)
1392R	Nitrosopumilales (0.002), Woesearchaeia; unclassified order (0.002), MG II (0.001)	SAR11 clade (0.003), Clostridiales (0.001), Rickettsiales (0.001)	Mamiellophyceae (0.227), Dictyochophyceae (0.011), Bacillariophyta (0.002)	Dinoflagellata (0.014), Protalveolata (0.010), Metazoa (Animalia) (0.008)
1389F	Nitrosopumilales (0.53), MG II (0.37), MG III (0.05)	Dadabacteriales (0.002), Clostridiales (0.002), SAR11 clade (0.002)	Mamiellophyceae (0.167), Cyanobacteria;Subsection I (0.103), Trebouxiophyceae (0.055)	Dinoflagellata (0.013), Protalveolata (0.008), Metazoa (Animalia) (0.006)
1510R	NA, 4-mismatch to reference	NA, 4-mismatch to reference	Prasinophyceae (0.393), Pelagophyceae (0.375), Pavlovophyceae (0.054)	Metazoa (Animalia) (0.055), Charophyta (0.036), Protalveolata (0.012)

Fig S1: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Archaea*

using metagenomes derived from Guaymas Basin sediments.

Fig S2: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Archaea*

using metagenomes derived from Saanich Inlet water samples.

Fig S3: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Archaea*

using metagenomes derived from BioGEOTRACES/BATS/HOT water samples.

Fig S4: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Archaea*

using metagenomes derived from SPOT/Catalina water samples.

Fig S5: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Archaea*

using metagenomes derived from Malaspina deep-sea water samples.

Fig S6: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Archaea*

using metagenomes derived from MBARI bloom water samples.

Fig S7: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Archaea* using metagenomes derived from Tara Oceans water samples (<3µm).

Fig S8: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for

Cyanobacteria and plastids using metagenomes derived from Guaymas Basin sediments.

Fig S9: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for

Cyanobacteria and plastids using metagenomes derived from Saanich Inlet water samples.

Fig S10: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for

Cyanobacteria and plastids using metagenomes derived from SPOT/Catalina water samples.

242    Fig S11: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for

243  *Cyanobacteria* and plastids using metagenomes derived from BioGEOTRACES/BATS/HOT water

244  samples.

Fig S12: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for

Cyanobacteria and plastids using metagenomes derived from Malaspina deep-sea water samples.

Fig S13: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for

Cyanobacteria and plastids using metagenomes derived from MBARI bloom water samples.

Fig S14: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for

Cyanobacteria and plastids using metagenomes derived from Tara Oceans water samples (<3µm).

Fig S15: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Bacteria*

(not including *Cyanobacteria* and plastids) using metagenomes derived from Guaymas Basin sediment

samples.

Fig S16: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Bacteria*

(not including *Cyanobacteria* and plastids) using metagenomes derived from Saanich Inlet water

samples.

Fig S17: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Bacteria*

(not including *Cyanobacteria* and plastids) using metagenomes derived from BioGEOTRACES/BATS/

HOT water samples.

267   Fig S18: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Bacteria*

268 (not including *Cyanobacteria* and plastids) using metagenomes derived from SPOT/Catalina water

269 samples.

Fig S19: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Bacteria*

(not including *Cyanobacteria* and plastids) using metagenomes derived from Malaspina deep-sea water

samples.

Fig S20: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Bacteria*

(not including *Cyanobacteria* and plastids) using metagenomes derived from MBARI bloom water

samples.

Fig S21: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Bacteria*

(not including *Cyanobacteria* and plastids) using metagenomes derived from Tara Oceans water

samples (<3µm).

Fig S22: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Eukarya*

using metagenomes derived from Guaymas Basin sediment samples.

Fig S23: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Eukarya*

using metagenomes derived from Saanich Inlet water samples.

Fig S24: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Eukarya*

using metagenomes derived from BioGEOTRACES/BATS/HOT water samples.

Fig S25: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Eukarya*

using metagenomes derived from SPOT/Catalina water samples.

Fig S26: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Eukarya*

using metagenomes derived from Malaspina deep-sea water samples.

Fig S27: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Eukarya*

using metagenomes derived from MBARI bloom water samples.

301    Fig S28: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Eukarya*

302 using metagenomes derived from Tara Oceans water samples (<3µm).

Fraction of 18S SSU rRNA from BiHoBa Metagenomes

304   Figure S29: Violin plot with inlaid boxplot showing the fraction of eukaryotic (18S) SSU

305   rRNA as a fraction of total across BATS, HOT, and bioGEOTRACES metagenomic datasets. The

306   violin plot provides a visual description of the distribution of data - i.e. in this case showing that the

307   vast majority of samples had < 30% of SSU rRNA reads accounted for by 18S sequences.

Figure S30: **Rulegraph for "Snakefile-compute.smk" portion of workflow**. Briefly, SSU rRNA is identified using phyloFlash from fastp quality-filtered fastq reads, which are then sorted with bbsplit.sh into 4 categories, and aligned to SSU rRNA references with pyNAST after removing low-complexity sequences. For each group and primer, matches to primer regions are then identified using custom python scripts and cutadapt (at 0, 1, 2, and 6-mismatches), quality filtered to omit sequences where the primer region plus 5 leading and trailing bases has one individual base with a phred score below 30. Matches are then summarized in plain-text output and ggplot2 graphs.

Figure S31:

Rulegraph for "Snakefile-compare.smk" portion of workflow to intercompare metagenome and amplicon data.