Evaluating and Improving SSU rRNA PCR Primer Coverage for Bacteria, Archaea, and Eukaryotes

Using Metagenomes from Global Ocean Surveys

Jesse McNichol[1], Paul M. Berube[2], Steven J. Biller[2,3], Jed A. Fuhrman[1]

1 - Department of Biological Sciences, University of Southern California, Los Angeles, California,

USA

2 - Department of Civil and Environmental Engineering, Massachusetts Institute of Technology,

Cambridge, MA, USA

3 - Department of Biological Sciences, Wellesley College, Wellesley, MA, USA

Abstract:

Small subunit ribosomal RNA (SSU rRNA) amplicon sequencing can quantitatively and

comprehensively profile natural microbiomes, representing a critically important tool for studying

diverse global ecosystems. However, results will only be accurate if PCR primers perfectly match the

rRNA of all organisms present. To evaluate how well marine microorganisms across all 3 domains are

detected by this method, we compared commonly-used primers with > 300 million rRNA gene

sequences retrieved from globally-distributed marine metagenomes. The best-performing primers when

comparing to 16S rRNA of Bacteria and Archaea were 515Y/926R and 515Y/806RB, which perfectly

matched over 96% of all sequences. Considering Cyanobacteria and Chloroplast 16S rRNA,

515Y/926R had the highest coverage (99%), making this set ideal for quantifying marine primary

producers. For eukaryotic 18S rRNA sequences, 515Y/926R also performed best (88%), followed by

V4R/V4RB (18S rRNA-specific; 82%) – demonstrating that the 515Y/926R combination performs best

overall for all 3 domains. Using Atlantic and Pacific Ocean samples, we demonstrate high

correspondence between 515Y/926R amplicon abundances (generated for this study) and metagenomic

16S rRNA (median R^2=0.98, n=272), indicating amplicons can produce equally accurate community

composition data versus shotgun metagenomics. Our analysis also revealed that expected performance

of all primer sets could be improved with minor modifications, pointing toward a nearly-completely

universal primer set that could accurately quantify biogeochemically-important taxa in ecosystems

ranging from the deep-sea to the surface. In addition, our reproducible bioinformatic workflow can

guide microbiome researchers studying different ecosystems or human health to similarly improve

existing primers and generate more accurate quantitative amplicon data.

Significance Statement:

PCR amplification and sequencing of marker genes is a low-cost technique for monitoring prokaryotic

and eukaryotic microbial communities across space and time, but will only work optimally if

environmental organisms match PCR primer sequences exactly. In this study, we evaluated how well

primers match globally-distributed short-read oceanic metagenomes. Our results demonstrate primer

sets vary widely in performance, and that at least for marine systems, rRNA amplicon data from some

primers lack significant biases compared to metagenomes. We also show that it is possible to create a

nearly universal primer set for diverse saline environments by defining a specific mixture of a few

dozen oligonucleotides and present a software pipeline that can guide rational design of primers for any

environment with available meta'omic data.

Introduction:

       Amplicon sequencing is a powerful tool for understanding microbial community composition

and dynamics in the oceans and other ecosystems (1), but the PCR amplification step is potentially

biased due to both technical issues during amplification and mismatches to organisms found in natural

environments (2–6). Despite these concerns, PCR amplicon sequencing retains several advantages that

make it desirable to investigate and correct biases. Firstly, it is a high-throughput and low-cost

technique, making it suitable for large numbers of samples - e.g. for global surveys of sediment, water,

animal-associated, and other microbial communities (1).

       Secondly, the targeted nature of the PCR assay means relatively small numbers of sequences are

sufficient for detecting rare organisms even when we have no genomes from any of their relatives, due

to the conserved nature of the molecule and the existence of comprehensive SSU rRNA sequence

databases. Being able to quantify the abundance and dynamics of rare organisms is important for

understanding ecosystem function since many rare bacteria have impacts far greater than their low

abundances might imply (7, 8). While PCR assays are targeted, we note that this does not imply they

need to be taxonomically-restricted. For example, there are some primer binding sites in the SSU rRNA

molecule that are nearly universally conserved between Archaea, Bacteria, and Eukarya (discussed

further below).

       Thirdly, using untargeted metagenomic sequencing for taxonomic profiling still has a number of

significant disadvantages versus amplicon sequencing. Direct taxonomic assignment of randomly-

sheared metagenomic reads by recruitment to reference genomes has the potential to assign taxonomy

at a very high resolution. However, the best currently available genome databases still only recruit ~ 40

% of metagenomic reads for marine prokaryotes (9), meaning a significant fraction of untargeted

metagenomic datasets are taxonomically uncharted. This problem is even more acute for environmental

eukaryotes with large amounts of non-coding genomic DNA and fewer genomic references. It is also

possible to extract SSU rRNA (or other marker genes) from metagenomes to generate a taxonomic

profile. However, SSU rRNA fragments typically represent only a small fraction a given dataset (~ 0.1

%), so is a far more costly way to obtain a comprehensive community profile. In addition, because

metagenomically-retrieved marker genes are randomly sheared fragments covering both conserved and

hypervariable regions and are in most cases too short to provide a unique match to reference sequences,

they must be assembled or otherwise clustered using non-deterministic algorithms. Because of this, the

taxonomic resolution of short-read metagenomic marker gene data is limited (e.g. near family or genus

level for the best known conserved markers like rRNA). In contrast, with rRNA amplicons, modern

"denoising" algorithms produce exact amplicon sequence variants (ASVs) that are stable biogeographic

markers that can be intercompared without reanalysis (10). The resulting datasets are also relatively

simple to analyze since they consist of a single gene region and can be comprehensively classified with

databases such as SILVA or RDP (11, 12).

Recent studies have shown that PCR amplicon sequencing of mock microbial communities can

recover known relative and absolute abundances extremely well, and thus provides accurate

quantification of natural gene copy abundances (3, 13, 14). In turn, this accuracy allows amplicon data

to serve as a ground-truth for modeling / ecological studies by quantifying community members and

their dynamics. While well-designed primers have the potential to recover truly quantitative data, these

studies also underscored the critical fact that a single mismatch between the primer and template

sequences can have a dramatic effect on the measured community composition in complex natural

mixtures. For example, a single internal mismatch in the popular Earth Microbiome Project primer

caused a ~10-fold bias against the most common bacteria in seawater (SAR11 cluster), and terminal 3'

mismatches can completely prevent amplification (2, 3, 6, 13). Since the extent of this bias is not easily

predicted, PCR primers should incorporate degenerate bases or specific oligonucleotide variants so that

all targeted organisms are perfectly matched without overly diluting the common perfect matches in

primer mixtures. This is especially critical for abundant taxa such as SAR11 as distortions in their

relative abundances will skew the remainder of the community, but it is also important to consider for

rare taxa which might have essential biogeochemical or ecological roles.

Previous studies have shown high coverage of natural taxa is possible by designing moderately

degenerate primers without sacrificing specificity or PCR efficiency (3, 13). This primer design was

accomplished by comparing oligonucleotide sequences to a SSU rRNA database such as SILVA or

RDP (11, 12) and then checking for mismatches to organisms known to be abundant in the environment

of interest. This approach led to marked improvements in primer design, for example by Apprill et al

(2015) and Parada et al (2015) who reported that small modifications to existing primers could better

quantify the dominant marine taxa SAR11 and *Thaumarchaea* (2, 3). However, in these primer

evaluations, the reliance on full-length references and giving each sequence in a database equal weight

can lead to a distorted perspective of the actual extent of matches and mismatches expected in real

samples since they do not take into account the highly unequal abundances in nature. In addition, some

environments may have abundant taxa poorly represented or unrepresented in these curated reference

databases.

Wear et al. (2018) studied the effect of these potential biases empirically by testing 4 primer sets

currently in broad use by marine microbiologists on a 16S rRNA mock microbial community derived

from natural seawater communities near Santa Barbara (6). These authors tested the effect of their

analysis pipeline on recovery of mock community sequences for different primers, and found their

primer-pipeline combination had several sequence-specific biases, in some cases due to a primer

mismatch. While an important step towards cross-comparing primers in an oceanographic context,

Wear et al. note that results are specific to their mock microbial community, analyzed with their

particular pipeline, and thus it remains unknown how representative their results are for other

ecosystems and other pipelines. More specifically, it is currently unknown how many primer-

mismatched rRNA sequences occur across diverse oceanographic environments, a key piece of

information for designing field measurements and interpreting results.

To more fully evaluate the extent to which primers currently in broad use by the marine

microbiology community perfectly match naturally occurring sequences, we developed a workflow to

conduct *in silico* primer evaluations with metagenomic SSU rRNA fragments, under the general

assumption that metagenomes have fewer methodological biases than PCR. While it is already known

that some primers have better coverage for abundant marine organisms versus others (3, 13), a precise

quantitative comparison of these primers with others that are in broad use has not yet been conducted.

We developed a new software pipeline for analyzing globally-distributed marine metagenomic

datasets (Table S1) to make a quantitative and objective evaluation of PCR primers for pelagic marine

ecosystems and to suggest specific improvements based on this evidence. We further experimentally

compared the quantitative performance of SSU rRNA amplicon sequencing, performed for this study,

against the published shotgun metagenomic data from the same sample DNA from the

bioGEOTRACES study. Our pipeline also allowed us to make intercomparisons between primer sets

often not considered together. For example, we compared the performance of two "universal" (3-

domain) PCR primers that amplify both 16S and 18S rRNA with primers designed to specifically target

only 16S or 18S rRNA and exclude the other molecule.

Because our software pipeline is broadly applicable for any environment where metagenomic

data exists, it will allow investigators to design environment-specific PCR probes that recover as many

(or as few) targeted organisms as desired. It is also flexible, allowing design or improvement of primers

from different variable regions to either retain stable intercomparisons with existing data, or to target a

region of the SSU rRNA molecule that has better taxonomic resolution for taxa of interest (15). Our

approach has the added advantage of providing a quantitative, evidence-based framework to identify

which taxa may have been affected by biases in existing datasets generated with sub-optimal primers -

information that is often unknown or only recognized anecdotally by specific investigators.

Results and Discussion:

Overall primer coverage for pelagic ocean datasets:

To evaluate the real-world extent of primer biases, we compared PCR primers currently in

broad use by the marine microbiology community (Table S2) to SSU rRNA sequences extracted from a

geographically diverse set of pelagic ocean metagenomes (Fig 1; Table S1). These analyses were based

on the percentage of perfect matches between primers and metagenome sequences as a metric to

determine potential PCR performance, since previous studies have shown that even a single mismatch

can significantly bias results (3, 13).

We observed relatively even recruitment of metagenomic reads across the SSU rRNA molecule,

which indicates that each different primer region has broadly similar potential to generate accurate

taxonomic profiles of naturally-occurring organisms (Supplemental Results). However, when we

compared the precise sequences derived from the primer-binding regions to the oligonucleotide primer

sequences, we observed that predicted median coverage of perfect matches varied widely among

primer pairs (Table S2), with 515Y/806RB and 515Y/926R showing the most consistently high

incidence of perfect matches (Fig 2). The ability of all primers to capture the underlying diversity in the

metagenomic samples can be improved to some extent by increasing degeneracies, though this varied

significantly between primer sets. Adding a single additional degeneracy fixes the majority of

mismatches for some primers (e.g. 785R; discussed further below), whereas others have greater than 2

mismatches to metagenomic sequences and thus would require more extensive modifications (Fig S1-

S28). Below, we discuss the performance of these primers across 4 broad taxonomic groups (*Archaea,*

*Bacteria, Cyanobacteria+*Plastidal 16S rRNA, and *Eukarya*). We separated *Cyanobacteria* +

phytoplankton plastidal 16S rRNA from bacterial 16S rRNA since these organisms are responsible for

the vast majority of marine primary productivity and thus are important to quantify accurately.

For *Archaea,* the primer with the best median coverage was the EMP 515Y/806RB combination

(coverage=0.996), followed by 515Y/926R (0.982), and 926wF/1392R (0.952). We also note that our

results suggest the 341F/785R primer combination could amplify most *Archaea* with the addition of

several degeneracies in 341F (Fig S1-S7).

For *Bacteria* (excluding *Cyanobacteria* and plastidal 16S rRNA sequences), the primer with the

best median coverage was the 515Y/926R combination (0.961), followed by 515Y/806RB (0.955),

27F/338R (0.888), 926wF/1392R (0.869), and 341F/785R (0.525). The relatively poor performance of

341F/785R is mainly due to the reverse primer (median individual primer coverages: 341F=0.887,

785R=0.624), whereas the other two less optimal primer pairs have more even individual performances

(27F=0.968, 338R=0.913; 926wF=0.916, 1392R=0.943).

For *Cyanobacteria* and chloroplast 16S rRNA (derived from eukaryotic phytoplankton), the

primer pair with the highest median coverage was 515Y/926R (0.992), followed by 515Y/806RB

(0.881), 27F/338R (0.826), 926wF/1392R (0.714), and 341F/785R (0.702). We note that the vast

majority of these mismatched sequences come from chloroplast 16S rRNA sequences - the common

cyanobacterial groups *Prochlorococcus* and *Synechococcus* have high coverage with all primers listed

above, so this is only a concern for those wishing to use chloroplast 16S rRNA data to quantify

eukaryotic phytoplankton.

For *Eukarya*, the primer pair with the highest median coverage was 515Y/926R (0.883),

followed by V4F/V4RB (0.822), 926wF/1392R (0.792), and 1389F/1510R (0.732). The distributions of

coverage were considerably broader for *Eukarya* versus all other 16S rRNA categories, which may be

due to a higher sequence heterogeneity among eukaryotic genomes. As above, all primer sets could be

improved with the addition of some degeneracy (Fig S22-S28), but approaching perfect coverage

seems less practically achievable for eukaryotic SSU rRNA sequences. In addition, our analysis

indicates that improvements in eukaryotic coverage are most likely to come from targeting universally-

conserved rRNA regions, such as those containing the 515Y/V4F and the 1389F/1392R primers

respectively (Fig S22-S28).

We note, however, that recognizable dinoflagellate chloroplast rRNA sequences were almost

completely missing from the metagenomic sequences (for all primer regions, not just 515Y/926R), and

thus may not appear in the resulting amplicons regardless of primer choice. This is possibly due to the

unusual genomic organization/subcellular localization of chloroplast genes in dinoflagellates (16), or

alternatively because we lack insufficiently accurate or comprehensive databases for identifying

dinoflagellate chloroplast 16S rRNA (discussed further below). It is possible, however, to recover and

confidently identify dinoflagellate 18S rRNA sequences from the 515Y/926R primer pair (17), making

it feasible to reconstruct a holistic picture of phytoplankton community composition / abundance in

combination with the chloroplast 16S rRNA.

In summary, the primer pairs with the highest environmental coverage were 515Y/926R

(universal), 515Y/806RB (prokaryote-only), and V4F/V4RB (eukaryote-only). It is worth emphasizing

that both universal primer sets have the potential to be equally or more comprehensive for *Eukarya*

versus the *Eukarya*-specific primer sets tested here. In addition, for those wishing to quantify the

abundance of eukaryotic phytoplankton using chloroplast data, the 515Y/926R primer set nearly

perfectly matches all environmental sequences. These results are a quantitative confirmation that the

careful primer design previously reported for the 515Y/806RB/926R primers has resulted in very high

overall coverage in oceanic ecosystems. On the other hand, certain primer sets (e.g. 341F/785R) have

very low coverage due to mismatches to dominant organisms, despite being designed to maximize

environmental coverage (18). Resulting data may thus be distorted, though we note that removing taxa

with mismatches to primers from processed data will recover accurate relative abundances for the

remaining taxa (13), due to the fact that biases are thought to be taxon-specific (19). This relative

abundance correction approach depends on our pipeline's specific identification of mismatched taxa,

and provides a way to recover useful quantitative information from legacy datasets that may have been

biased during PCR. In combination with ecosystem-specific full-length 16S rRNA databases, this

correction approach could represent an important tool for integrating historical amplicon datasets and

modern long-read sequencing data into a single intercomparable framework for observing longer-term

changes in ecosystem structure (15). In addition, by eliminating or at least controlling for primer bias,

our approach could help better evaluate how well particular primer regions perform with respect to

taxonomic profiling and classification since it would tease apart the effect of variable region from

amplification artifacts.

A Quantitative Intercomparison between SSU rRNA Amplicon Sequencing and Metagenomics:

In order for amplicon sequencing to become useful as a quantitative tool for measuring natural

gene abundances, it is desirable to benchmark amplicon abundances against an external reference that

is not affected by potential PCR biases to gain a more objective determination of how well a primer

recovers true gene abundances. While shotgun metagenomics is not necessarily free of all biases, these

biases are distinct from those in PCR-based amplicon studies; thus, if both data types yield similar

community composition patterns, we can be more confident about our overall conclusions. We tested

whether amplicon-based strategies recover similar patterns as metagenomes by generating PCR

amplicon sequences for a subset of 272 BioGEOTRACES (20) samples (cruises GA03 and GP13). We

used the same DNA, eliminating extraction biases as a potential confounding factor in this

intercomparison. We were thus able to make a direct "apples to apples" quantitative comparison

between the relative abundances of organisms determined with amplicons vs. metagenomic reads.

To accomplish this, we compared exact amplicon sequence variants (ASVs) from the

515Y/926R primer pair with metagenomic reads from the same region of the SSU rRNA molecule. To

make a more robust quantitative comparison, ASVs were clustered as necessary to account for the fact

that short (150bp) metagenomic reads could in some cases only be attributed to a broader phylogenetic

group (e.g. *Prochlorococcus* and SAR11). The summed abundance of all taxonomic groups, as

determined by ASVs and metagenomic reads, was then compared using a linear least-squares

regression. For 16S rRNA sequences, we observed very similar community composition for both

techniques, as shown by very high R^2 values and coverage in Fig 3a (median R^2=0.98; range=0.55-1.0).

This indicated that the ASV sequences were recovering the same diversity of organisms present in

metagenomic reads from the primer region and is consistent with the aforementioned observation of

even coverage of metagenomic reads across the SSU rRNA molecule.

These patterns were generally consistent across the surface-mesopelagic transition where there

is a major change in community composition (Fig 3). We do however note that at depths > 150m about

38% of samples had R^2 values below 0.95 (Fig 3a), vs. 9.2% for samples at depths < 150m. These

lower correspondences are unlikely to be due to mismatches to the primer region, since predicted

primer coverage is uniformly high across depth for the 515Y/926R primer pair across depth (Fig 3b).

Other factors such as DNA quality or quantity may have been responsible for this, but we cannot rule

out the possibility it involves characteristics of the sequences between the priming regions such as

secondary structure or expanded loops known from some eukaryotes (21). Regardless, these results

show that amplicon sequencing and metagenomic profiling typically have a very high quantitative

correspondence with the primers tested here (515Y/926R). Additionally we provide a software toolbox

for making further intercomparisons between arbitrary primer sets and metagenomes that could be

useful for optimizing both PCR amplicon sequencing and metagenomics for diverse environments.

Improving Existing Primers:

In addition to quantifying mismatches, our pipeline also identifies the specific primer region

variants found in particular naturally-occurring taxa and their coverage across datasets. This

information can guide improvements for any of the primers tested here, and we identify two specific

use cases. Firstly, some of the primers identified as having major mismatches are dominated by a few

organisms, and thus small modifications are all that is necessary to make them more appropriate for

oceanic ecosystems. Secondly, we were interested to know whether for the better-performing primers it

would be possible to identify a combination of oligonucleotides that would perform well across all

marine environments tested here. Such a primer set could have the advantage of being able to monitor

rare but biogeochemically-important taxa that may become more abundant due to climate change, e.g.

as a result of expanding oxygen minimum zones.

To illustrate how a primer with poor predicted performance could be improved, we use the case

of 785R which has a mismatch to the dominant organism SAR11. Because the SAR11 mismatches

represent up to 40% of total 16S rRNA reads (Table S3), this one mismatch will have a major effect on

the resulting data but could be corrected by adding a single additional degeneracy at position 7 in the

primer (5'-GACTAC**N**VGGGTATCTAATCC). However, 785R also has mismatches to a number of

abundant chloroplast sequences (Fig 2), which could be corrected with two additional degeneracies (5'-

RA**Y**TACNVGGGTATCTAATCC). Compared to the original primer (5'-

GACTACHVGGGTATCTAATCC), these primers would require the synthesis of a larger number of

oligonucleotide combinations (12 and 48 vs. 9 originally) but are well within the number of

combinations currently in use for other environmental amplification strategies (22). Interestingly,

despite being the worst-performing primer for marine ecosystems due to mismatches to the dominant

taxon SAR11, we predict that by adding the above degeneracies this primer would achieve nearly

perfect coverage for *Bacteria* (Fig S15-S21). This suggests the primer-binding region of 785R is

highly-conserved among *Bacteria*, consistent with the initial publication describing this primer as the

best overall for prokaryotic taxa across diverse environments (18). This observation also underscores

the value of comparing primer sequences with environmental data since it uncovered a significant,

unexpected mismatch to an abundant taxon that can now be corrected.

We also investigated whether it is possible to improve primer sets that already perform well

(e.g. 515Y/806RB/926R) to the point where they offer near-universal coverage of all taxa identified in

this study. To do so, we identified datasets that had significant potential for coverage improvement

(Figures S1-S28) and certain rare order-level taxa that had many mismatches or are known to be

biogeochemically-important.

Our data showed that the 806RB primer performed extremely well overall for 16S rRNA

sequences, but did miss a number of 16S rRNA chloroplast sequences mostly from the *Mamielliales*

clade, which includes small picoeukaryotes such as *Ostreococcus* and *Bathycoccus*. As shown in Fig

S8-S14, these taxa were still missed when allowing for up to 2 mismatches, meaning they are unlikely

to be accurately quantified with the current primer design. Alterations would be possible, but we note

that this may be an intentional design choice, since chloroplast amplification may be undesirable in

certain environments (23) but is often very useful in euphotic zone marine water samples where it can

be used to quantify primary producer abundance. Related to this, we note that a context-dependent

advantage of 806RB is that it lacks an 18S rRNA binding site. This means that in combination with

515Y, nuclear 18S rRNA will not be amplified with 806RB. In other words, even though the variable

region covered by 515Y/806RB contains eukaryotic sequences, this primer set will not recover them as

amplicons – demonstrating that primer design places a first-order constraint on how well a given

amplicon product reflects the natural community. In contrast, the 51Y/926R combination will amplify

these eukaryotic sequences. In a practical sense, amplifying 18S rRNA sequences with 16S rRNA can

present problems, for example in host-associated microbiome studies where 18S rRNA could

overwhelm targeted 16S rRNA. This overwhelming of libraries by 18S rRNA sequences is however

unlikely to be a major issue for the 515Y/926R primer pair for pelagic ocean samples. This is due to the

fact that 18S rRNA sequences will generally be selected against in the sequencing process (thus

reducing their overall abundances (13)), and the fact that marine environments are generally dominated

by prokaryotic microorganisms (13; Fig S29).

The 515Y primer was relatively easy to improve, given that its performance was already very

robust across diverse datasets (Fig S1-S28). We identified four positions where further degeneracies

could be added (5'-GTG**B**CAGCM**SY**CGCGGT**M**A), and were sufficient to resolve the vast majority

of mismatches including to rare taxa such as certain representatives of the *Patescibacteria* (a.k.a.

candidate phyla radiation (CPR) bacteria) which were previously identified as taxonomic "blind spots"

in PCR amplicon analysis (5). However, incorporating these new degeneracies produced a relatively

large number of oligonucleotide combinations compared to the original primer (48 versus 4 originally)

and most of these new variants were not observed in the metagenomic data. We thus decided to

redesign the primer based on a different approach, starting from a non-degenerate primer and only

adding in variants that could be detected in the natural environment or in the SILVA database. This

resulted in a specific mixture of 13 specific oligonucleotides that we term 515Yp-min ("p" for pelagic,

"min" for minimal). This mixture of oligonucleotides is tailored specifically to the environmental

sequences present in natural marine environments. It both maximizes organismal coverage, while

keeping the overall degeneracy at a relatively low level compared to other degenerate primers used in

environmental microbiology (22). Another key advantage of this tailored approach is that it permits

iterative improvements to a primer set over time. For example, we added a variant matching

dinoflagellate chloroplast sequences from the PhytoRef database (24) that may potentially improve

quantification of these organisms' plastidal 16S rRNA in combination with other modifications to 926R

(discussed below). In the future, if we were able to identify other variants explaining why

dinoflagellate chloroplast 16S sequences are conspicuously absent from 515Y/926R amplicon data, it

would then be feasible to add these new olignucleotides to the mixture without creating excessive

sequence redundancy.

The 926R primer had mismatches to more taxa, and thus required more extensive

modifications. For example, Table S3 shows that 926R has mismatches to both *Ectothiorhodospirales*

(an uncultivated group related to purple sulfur bacteria (25)), and the *Rickettsiales* (an order that

includes mitochondrial sequences in addition to free-living organisms (26)). While less abundant

overall, we also noted mismatches in the Tara Oceans dataset to *Brocadiales*, the order containing

anammox bacteria (27). In the Guaymas basin sediment samples, some of the dominant mismatches

were to the *Campylobacteria*, dominant chemoautotrophs in many sulfidic systems (28), as well as

unusual *Archaea* that are now recognized to be much more diverse than previously thought and missed

by some PCR primers (29). Accurate quantification of these rare taxa would allow the monitoring of

unusual but potentially significant changes in biogeochemistry. For example, the *Campylobacteria* are

known to be abundant in certain low-oxygen microenvironments such as in sediment trap particles

(30), and the *Brocadiales* are abundant in oxygen minimum zones (27). Being able to better quantify

these "sentinel taxa" could allow us to monitor the dynamics of these low-oxygen environments that

are likely to expand under global climate change.

Following the same approach discussed above for 515Yp-min, we developed a mixture of 38

oligonucleotides ("926Rp-min") that contains perfect matches for aforementioned taxa, primer variants

that have > 2% overall relative abundance across all environments, as well as 2 sequences from

dinoflagellate chloroplasts that were not previously covered. This set of oligonucleotides is

comprehensive for all the environments tested in our study and would likely result in major

improvements in quantification all of the aforementioned taxa. To achieve the same organismal

coverage by adding more degenerate sites would increase the total number of primer variants to at least

144 (compared to 16 in the original primer). Not only would this dilute perfectly-matching

oligonucleotides and be unlikely to result in efficient PCR amplification, it would also include many

redundant oligonucleotides unnecessary with an explicit oligonucleotide mixture. The improvements in

coverage for this new mixture may be particularly apparent for *Rickettsiales* templates, many of which

have a single mismatch at the 3' end of the primer that is known to completely prevent PCR

amplification (6). Since the order *Rickettsiales* contains sequences from mitochondrial SSU rRNA, this

modification will potentially allow for more effective quantification of protist abundances, similar to

how chloroplasts quantify marine protistan phytoplankton.

Since it is beyond the scope of this paper to provide exhaustive recommendations for each

primer set, we direct interested readers to an Open Science Foundation repository which contains all

necessary output files, scripts, and a suggested workflow for improving primers using metagenomic

data (osf.io/gr4nc/). Improvements should be possible for most taxa/primer combinations since a few

mismatches typically dominated, with a long tail of rare variants that may derive from sequencing

errors or pseudogenized SSU rRNA. This information would allow an interested reader to optimize one

of the primers investigated here for particular taxa or environments of interest.

Moving forward, we recommend oceanographers consider applying these two new primer

mixtures for pelagic water column surveys to maximize organismal coverage (515Yp-min, 926Rp-min;

Table S4). By correcting for 3' mismatches that lead to taxonomic "blind spots" and including exact

matches to taxa found in low-oxygen regions, we should be able to better quantify the abundance and

dynamics of organisms critical to marine carbon, nitrogen, and sulfur cycling. In addition, by

representing primers as a specific mixture of oligonucleotides rather than a single sequence with

degeneracies, it leaves open the possibility of small modifications in the future to improve coverage;

the same is not true for fully degenerate primers where each new degeneracy will multiply the total

number of combinations by at least 2-fold and further dilute the other perfectly matching

oligonucleotides. If proven to be as quantitative as the original 515Y/926R primers (3, 13), these new

mixtures would provide an affordable, comprehensive (all organisms from Archaea - to Zooplankton),

and scalable method to measure whole community compositions across time and space in the world's

oceans. While further validation would be necessary, we believe our results and those of previous

studies (3, 13–15) demonstrate the potential for PCR amplicon methods to accurately quantify natural

marker gene abundances in a robust and accurate manner, either by accurately recovering true relative

abundances, or in combination with internal standards to obtain absolute quantification (14). This

would allow confident measurement of microbial community composition alongside well-developed

methods such as those for quantifying phytoplankton photosynthetic pigments. In turn, this would help

expand our understanding of how oceanic microbes interact with biogeochemical cycles and respond to

global climate change.

Finally, we believe our bioinformatic approach could be applied elsewhere to generate

ecosystem-specific primer sets. For example, it remains unknown whether the patterns we observed for

seawater would be true in other systems, such as terrestrial sediment, soil, freshwater, or animal/plant-

associated microbiomes. Our reproducible workflow, combined with expert curation and broad

environmental sampling could ensure that taxa of interest are accurately quantified by oligonucleotide

primers for any environment where deeply-sampled metagenomes could be used as a database. In

addition, our approach could allow for the rational evaluation and improvement of new primers for next

generation long-read amplicon sequencing, or to improve the performance of primers used to identify

animal species of economic or conservation importance. By evaluating primer coverage objectively and

providing simple ways to iteratively improve existing primer sets, it would ensure that the potential of

these techniques for monitoring microbiomes relevant to ecosystem functioning and human health is

fully realized.

Materials and Methods:

Sample scope / biogeographic distribution and evaluated primers:

We analyzed globally-distributed metagenomic datasets that reflect a range of depths and

nutrient regimes (Table S1; Fig 1), and tested primers that are commonly used in environmental

microbiology (Table S2; (2–4, 18, 31–35)). Metagenomic samples included worldwide sampling

expeditions such as Tara oceans (36), the Malaspina expedition (37), and BioGEOTRACES (20).

Among these three campaigns, the Malaspina metagenomes focus largely on the deep sea, whereas

TARA and BioGEOTRACES sample primarily the sun-lit ocean. In addition, we used time-series data

from the Hawaii Ocean Time-Series and the Bermuda Atlantic Time-Series (38, 39) (HOT/BATS; both

analyzed with BioGEOTRACES as BiHOBA [BioGEOTRACES-HOT-BATS]), a bloom time-series in

Monterey Bay (40), and the San Pedro Ocean Time-Series (41) (SPOT). We also included in the

analysis two sites representing suboxic/anoxic systems - the Saanich inlet time-series (42), and samples

collected from hydrothermally-altered sediments in Guaymas basin (43). We also generated new

16S/18S rRNA universal amplicon sequences for two BioGEOTRACES cruise transects (GA03/GP13;

noted with open circles on Fig 1) which were used for making intercomparisons between metagenomes

and amplicons.

Evaluation of primer coverage/biases using metagenomic reads:

In order to evaluate the performance of our primer set compared to others, we used

metagenomic reads to estimate the fraction of SSU rRNA fragments that could be successfully targeted

by (i.e. perfectly match) each primer set. In brief, our pipeline retrieves SSU rRNA from fastq-

formatted shotgun metagenomic samples, removes sequences containing uninformative repeats, sorts

into four organismal categories (*Archaea, Bacteria, Cyanobacteria* (including plastidal 16S rRNA), and

Eukarya), and aligns these sequences to a group-specific reference SSU rRNA sequence (e.g.

Saccharomyces cerevisiae for *Eukarya*). To obtain sequences overlapping with each oligonucleotide

primer and to exclude non-primer reads, coordinates provided in the config file were used to extract

reads from alignments that overlapped with the primer-binding region plus 5 leading or trailing bases.

These reads, which represent SSU rRNA fragments with a primer-binding region, were then quality-

filtered and compared with primer sequences (Table 2) at 0, 1 and 2-mismatch thresholds (Fig S1-S28)

to identify matched and mismatched sequences. As an additional quality-control step, we excluded any

aligned read falling in the primer region that did not have a match to the primer at a 6-mismatch

threshold (e.g. misalignments or distantly homologous sequences). Both mismatched and matched

reads were then classified using the SILVA 132 database (*Archaea, Bacteria, Eukarya*) and PhytoRef

(11, 24) (*Cyanobacteria* + plastidal 16S rRNA). The specific software implementation is described

fully in the supplementary material. For those wishing to inspect results further, complete plaintext

summaries of output and summary plots are available online at https://osf.io/gr4nc/.

These computational steps were implemented with the snakemake workflow engine (44) to

create a documented and reproducible pipeline which is freely available

(github.com/jcmcnch/MGPrimerEval). It automatically produces tabular output / summary plots and

can be applied to new samples and primers by modifying template configuration files. The specific

functioning of the three workflows are described in more detail in the supplemental material and on the

github page.

PCR amplification and ASV generation:

272 DNA samples from GEOTRACES cruises GA03 and GP13 were used to generate PCR

amplicons, and represent either surface (GP13) or surface plus upper mesopelagic water samples

(GA03). GP13 is a longitudinal transect in the southern Pacific, whereas GA03 is a longitudinal

transect in the north Atlantic. DNA used for PCR amplicons was the exact same DNA material used to

produce metagenomes described by Biller et al (2018) (20) except that it was diluted to 0.5 ng/µL in

low-EDTA TE buffer prior to amplification. DNA was amplified with the 515Y/926R primers (515Y:

5'-GTGYCAGCMGCCGCGGTAA / 926R: 5'-CCGYCAATTYMTTTRAGTTT) (3) using 0.5ng of

DNA template in a 25 µL reaction with a final primer concentration of 0.3mM. Primers were part of a

larger construct with Illumina adapter constructs/barcodes already included, allowing for single-step

library preparation. Thermocycling was as follows: an initial denaturation at 95°C for 120s followed by

30 amplification cycles of 95°C × 45s; 50°C × 45s; 68°C × 90s and a final elongation step at 68°C ×

300s. Laboratory methods are described in more detail at

https://dx.doi.org/10.17504/protocols.io.vb7e2rn. Both 16S and 18S rRNA mock communities

(staggered and even versions of both (3, 13)) were included in the sequencing run as quality control.

Amplicons were sequenced with HiSeq 2x250 RapidRun technology at the USC sequencing center in a

run pooled with shotgun metagenomic samples. Raw basecall data was demultiplexed with bcl2fastq (-

r 20 -p 20 -w 32 --barcode-mismatches 0) using the 6 basepair reverse indices and subsequently with

the forward 5 basepair barcode with cutadapt according to the templates provided at

github.com/jcmcnch/demux-notes. Raw data from this step of the pipeline is available on NCBI under

the umbrella Bioproject PRJNA659851 and three sub-projects PRJNA658608 (controls),

PRJNA658384 (GA03), and PRJNA658385 (GP13)**.**

Sequence data was analyzed with a custom *in-silico* pipeline available at

github.com/jcmcnch/eASV-pipeline-for-515Y-926R and based on qiime2 2019.4 (45). Briefly, 16S and

18S rRNA sequences were first partitioned using a custom 16S/18S rRNA-specific databases derived

from the SILVA 132 and PR2 (11, 46) which is available at osf.io/e65rs/. After partitioning, primers

were removed from amplicons allowing up to 20% mismatches to the primer, and raw sequences were

imported into qiime2 and denoised with DADA2 (10). Taxonomy was assigned using a naïve Bayesian

classification algorithm with the SILVA 132 database subsetted to the amplicon region as a reference

(47). Sequences identified as chloroplasts were re-annotated with PhytoRef (24), and 18S rRNA

sequences were additionally annotated with PR2 (46). A step-by-step protocol for these informatic steps

is available at https://dx.doi.org/10.17504/protocols.io.vi9e4h6. After denoising and annotation, biom

tables were exported with taxonomy as tsv files and converted to relative abundances which were used

as input for the metagenome-amplicon comparison.

Acknowledgements: This work was supported by the Simons Collaboration on Computational

Biogeochemical Modeling of Marine Ecosystems (CBIOMES) grant 549943 to JAF, the Gordon and

Betty Moore Foundation Marine Microbiology Initiative grant 3779 and NSF grant OCE-1737409. We

gratefully acknowledge the effort of the scientists who collected the BioGEOTRACES field samples

analyzed here: Christel Hassler, Debbie Hulston, and Elizabeth Maas (GP13) and Jeremy Jacquot

(GA03). We also acknowledge the efforts of Keven Dooley and Madeline Williams who conducted

DNA extractions, which was funded in part by grants to Sallie W. Chisholm at MIT that enabled the

collection and processing of DNA samples from GEOTRACES cruises. These include the Simons

Foundation (Life Sciences Project Award ID 337262, S.W.C; SCOPE Award ID 329108, S.W.C), the

Moore Foundation (Grant IDs GBMF495 and GBMF4511), and the National Science Foundation

(OCE-1153588, OCE-1356460, and DBI-0424599). We thank S.W.C. for donating the samples and

supporting parts of the study.

References Cited:

1. L. R. Thompson, *et al.*, A communal catalogue reveals Earth's multiscale microbial diversity. *Nature* **551**, 457–463 (2017).

2. A. Apprill, S. McNally, R. Parsons, L. Weber, Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. *Aquatic Microbial Ecology* **75**, 129–137 (2015).

3. A. E. Parada, D. M. Needham, J. A. Fuhrman, Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. *Environ Microbiol* **18**, 1403–1414 (2016).

4. W. Walters, *et al.*, Improved Bacterial 16S rRNA Gene (V4 and V4-5) and Fungal Internal Transcribed Spacer Marker Gene Primers for Microbial Community Surveys. *mSystems* **1** (2016).

5. E. A. Eloe-Fadrosh, N. N. Ivanova, T. Woyke, N. C. Kyrpides, Metagenomics uncovers gaps in amplicon-based detection of microbial diversity. *Nature Microbiology* **1**, 15032 (2016).

6. E. K. Wear, E. G. Wilbanks, C. E. Nelson, C. A. Carlson, Primer selection impacts specific population abundances but not community dynamics in a monthly time-series 16S rRNA gene amplicon analysis of coastal marine bacterioplankton. *Environmental Microbiology* **20**, 2709–2726 (2018).

7. B. J. Campbell, L. Yu, J. F. Heidelberg, D. L. Kirchman, Activity of abundant and rare bacteria in a coastal ocean. *PNAS* **108**, 12776–12781 (2011).

8. A. Jousset, *et al.*, Where less may be more: how the rare biosphere pulls ecosystems strings. *The ISME Journal* **11**, 853–862 (2017).

9. M. G. Pachiadaki, *et al.*, Charting the Complexity of the Marine Microbiome through Single-Cell Genomics. *Cell* **179**, 1623-1635.e11 (2019).

10. B. J. Callahan, *et al.*, DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods* **13**, 581–583 (2016).

11. C. Quast, *et al.*, The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucl. Acids Res.* **41**, D590–D596 (2013).

12. J. R. Cole, Q. Wang, B. Chai, J. M. Tiedje, "The Ribosomal Database Project: Sequences and Software for High-Throughput rRNA Analysis" in *Handbook of Molecular Microbial Ecology I*, F. J. de Bruijn, Ed. (John Wiley & Sons, Inc., 2011), pp. 313–324.

13. Y.-C. Yeh, J. C. McNichol, D. M. Needham, E. B. Fichot, J. A. Fuhrman, Comprehensive single-PCR 16S and 18S rRNA community analysis validated with mock communities, and estimation of sequencing bias against 18S. *Environ. Microbiol.* In Press. (2021). doi: 10.1111/1462-2920.15553.

14. Y. Lin, S. Gifford, H. Ducklow, O. Schofield, N. Cassar, Towards quantitative microbiome community profiling using internal standards. *Appl. Environ. Microbiol.,* AEM.02634-18 (2018).

15. M. S. Dueholm, *et al.,* Generation of Comprehensive Ecosystem-Specific Reference Databases with Species-Level Resolution by High-Throughput Full-Length 16S rRNA Gene Sequencing and Automated Taxonomy Assignment (AutoTax). *mBio* **11** (2020).

16. T. Laatsch, S. Zauner, B. Stoebe-Maier, K. V. Kowallik, U.-G. Maier, Plastid-Derived Single Gene Minicircles of the Dinoflagellate Ceratium horridum Are Localized in the Nucleus. *Mol Biol Evol* **21**, 1318–1322 (2004).

17. D. M. Needham, J. A. Fuhrman, Pronounced daily succession of phytoplankton, archaea and bacteria following a spring bloom. *Nature Microbiology* **1**, 16005 (2016).

18. A. Klindworth, *et al.,* Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucl. Acids Res.,* gks808 (2012).

19. M. R. McLaren, A. D. Willis, B. J. Callahan, Consistent and correctable bias in metagenomic sequencing experiments. *eLife* **8**, e46923 (2019).

20. S. J. Biller, *et al.,* Marine microbial metagenomes sampled across space and time. *Scientific Data* **5**, 180176 (2018).

21. A. Obiol, *et al.,* A metagenomic assessment of microbial eukaryotic diversity in the global ocean. *Molecular Ecology Resources* **20**, 718–731 (2020).

22. J. C. Gaby, D. H. Buckley, A Comprehensive Evaluation of PCR Primers to Amplify the nifH Gene of Nitrogenase. *PLoS One* **7** (2012).

23. S. Thijs, *et al.,* Comparative Evaluation of Four Bacteria-Specific Primer Pairs for 16S rRNA Gene Surveys. *Front. Microbiol.* **8** (2017).

24. J. Decelle, *et al.,* PhytoREF: a reference database of the plastidial 16S rRNA gene of photosynthetic eukaryotes with curated taxonomy. *Molecular Ecology Resources* **15**, 1435–1445 (2015).

25. A. Oren, "The Family Ectothiorhodospiraceae" in *The Prokaryotes*, (Springer, Berlin, Heidelberg, 2014), pp. 199–222.

26. D. N. Fredricks, "Introduction to the Rickettsiales and Other Intracellular Prokaryotes" in *The Prokaryotes*, M. D. P. Dr, S. Falkow, E. Rosenberg, K.-H. Schleifer, E. Stackebrandt, Eds. (Springer New York, 2006), pp. 457–466.

27. L. Medina Faull, P. Mara, G. T. Taylor, V. P. Edgcomb, Imprint of Trace Dissolved Oxygen on Prokaryoplankton Community Structure in an Oxygen Minimum Zone. *Front. Mar. Sci.* **7** (2020).

28. B. J. Campbell, A. S. Engel, M. L. Porter, K. Takai, The versatile ε-proteobacteria: key players in sulphidic habitats. *Nature Reviews Microbiology* **4**, 458–468 (2006).

29. B. J. Baker, *et al.*, Diversity, ecology and evolution of Archaea. *Nat Microbiol*, 1–14 (2020).

30. D. Boeuf, *et al.*, Biological composition and microbial dynamics of sinking particulate organic matter at abyssal depths in the oligotrophic open ocean. *PNAS* **116**, 11824–11832 (2019).

31. L. A. Amaral-Zettler, E. A. McCliment, H. W. Ducklow, S. M. Huse, A Method for Studying Protistan Diversity Using Massively Parallel Sequencing of V9 Hypervariable Regions of Small-Subunit Ribosomal RNA Genes. *PLOS ONE* **4**, e6372 (2009).

32. D. Wilkins, E. van Sebille, S. R. Rintoul, F. M. Lauro, R. Cavicchioli, Advection shapes Southern Ocean microbial assemblages independent of distance and environment effects. *Nature Communications* **4**, 2457 (2013).

33. S. Balzano, E. Abs, S. Leterme, Protist diversity along a salinity gradient in a coastal lagoon. *Aquatic Microbial Ecology* **74**, 263–277 (2015).

34. T. Stoeck, *et al.*, Multiple marker parallel tag environmental DNA sequencing reveals a highly complex eukaryotic community in marine anoxic water. *Molecular Ecology* **19**, 21–31 (2010).

35. C. S. Fortunato, L. Herfort, P. Zuber, A. M. Baptista, B. C. Crump, Spatial variability overwhelms seasonal patterns in bacterioplankton communities across a river to ocean gradient. *The ISME Journal* **6**, 554–563 (2012).

36. S. Pesant, *et al.*, Open science resources for the discovery and analysis of Tara Oceans data. *Sci Data* **2**, 1–16 (2015).

37. S. G. Acinas, *et al.*, Metabolic Architecture of the Deep Ocean Microbiome. *bioRxiv*, 635680 (2019).

38. D. M. Karl, M. J. Church, Microbial oceanography and the Hawaii Ocean Time-series programme. *Nat Rev Micro* **12**, 699–713 (2014).

39. D. K. Steinberg, *et al.*, Overview of the US JGOFS Bermuda Atlantic Time-series Study (BATS): a decade-scale look at ocean biology and biogeochemistry. *Deep Sea Research Part II: Topical Studies in Oceanography* **48**, 1405–1447 (2001).

40. B. Nowinski, *et al.*, Microbial metagenomes and metatranscriptomes during a coastal phytoplankton bloom. *Sci Data* **6**, 1–7 (2019).

41. E. T. Sieradzki, J. C. Ignacio-Espinoza, D. M. Needham, E. B. Fichot, J. A. Fuhrman, Dynamic marine viral infections and major contribution to photosynthetic processes shown by spatiotemporal picoplankton metatranscriptomes. *Nat Commun* **10**, 1–9 (2019).

42. A. K. Hawley, *et al.*, A compendium of multi-omic sequence information from the Saanich Inlet water column. *Sci Data* **4**, 1–11 (2017).

43. N. Dombrowski, K. W. Seitz, A. P. Teske, B. J. Baker, Genomic insights into potential interdependencies in microbial hydrocarbon and nutrient cycling in hydrothermal sediments. *Microbiome* **5**, 106 (2017).

44. J. Köster, S. Rahmann, Snakemake—a scalable bioinformatics workflow engine. *Bioinformatics* **28**, 2520–2522 (2012).

45. E. Bolyen, *et al.*, Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nature Biotechnology* **37**, 852–857 (2019).

46. L. Guillou, *et al.*, The Protist Ribosomal Reference database (PR2): a catalog of unicellular eukaryote Small Sub-Unit rRNA sequences with curated taxonomy. *Nucleic Acids Res* **41**, D597–D604 (2013).

47. N. A. Bokulich, *et al.*, Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome* **6**, 90 (2018).

Figure 1: Distribution of metagenomic samples used in this study. BioGEOTRACES samples that were used in the metagenome/amplicon intercomparison are noted with open circles (GA03 - N. Atlantic; GP13 - S. Pacific).

Figure 2: Oligonucleotide PCR primer coverage based on comparisons with globally-distributed oceanic metagenomes. The shapes of the plots illustrate the distribution of primer coverage across all samples, where perfect correspondence of primers and metagenome sequences is represented by data piled up against "1.0" on the x axis, a situation approached by the UNIV V4-V5 primer for Cyanobacteria+ Plastid 16S rRNA, for example. Note that the "All SSU rRNA" panel represents all of the SSU rRNA targeted by a particular primer set (e.g. for 1389F/1510R this includes only Eukarya *while for 926wF/1392R it includes all 4 taxonomic groups). Data are corrected for predicted taxonomic overlap between forward and reverse primers. Coordinates refer to locations of primer alignments to the* Escherichia coli *16S rRNA gene (strain K-12; substrain MG1655).*

Fig 3: Metagenome-amplicon/oligonucleotide primer comparisons across the marine water column. A) Metagenome-amplicon quantitative correspondence across all depths for the GA03 cruise (BioGEOTRACES), showing the R^2 value of the plot of the relative abundances of amplicon sequence variants (ASVs; 515Y/926R primers) vs. relative abundances of metagenomic reads from the same organisms (see text for details of comparisons). B) The fraction of exact matches of the 515Y and 926R amplicon primers compared to metagenomic reads of the priming regions for each sample, summarized as the "worst-case" coverage for the combined primers (i.e. 1 - ($missed_{fwd}$ + $missed_{rev}$)).

Supplementary Information:

Supplementary Methods:

Software Implementation: Software dependencies are handled by conda implemented in snakemake

(1). Examples of how the pipeline can be run are available at github.com/jcmcnch/MGPrimerEval in

the "runscripts" folder. Both the "classify" and "compare" workflows depend on the output of the

"compute" workflow. The "classify" workflow is for detailed exploration of the taxonomy of retrieved

sequences and the "compare" workflow allows for intercomparisons between SSU rRNA amplicon

sequence variants (ASVs) and metagenomic SSU rRNA from the same amplicon region.

Compute workflow ("Snakefile-compute.smk"; Fig S30): This portion of the analysis calculates the

coverage of primer sequences for metagenomic samples specified in a configuration file. Results are

calculated as the fraction of total SSU rRNA reads matching the primer at 0, 1, and 2-mismatches for 4

taxonomic partitions as described below. A rule graph summarizing the steps is shown in Figure S30.

Briefly, fastq reads are first cleaned using fastp's (v 0.14.1) default parameters and trimming 3 trailing

and leading bases (2). To retrieve SSU rRNA sequences from metagenomes, we used the phyloFlash (v

3.3b1) package (3) which uses bbmap (v 38.69) (4) and a curated version of SILVA132 (5) to retrieve

both 16S and 18S rRNA sequences. Since retrieved sequences contained a small fraction of low-

complexity repeats that were not homologous to the SSU rRNA, we used komplexity (v 0.3.6) (6) to

remove these likely spurious hits.

SSU rRNA sequences thus identified were then split into 4 different pools (Eukaryota, Archaea,

Bacteria (non-cyanobacterial), and Cyanobacteria (including chloroplasts)) using bbsplit.sh v38.22 (4)

comparing against curated subsets of the SILVA/PR2 reference databases (5, 7) (sequences and a

template script for creating databases are available at osf.io/e65rs/). Partial SSU rRNA sequences were

then aligned using pyNAST v1.2.2 (8) to group-specific references available as part of the github

repository. Primer-specific subsets were then parsed from the alignment using python/biopython (9),

only outputting sequences that overlap the primer region plus 5 leading or trailing bases. The 1-based

coordinates of each primer region (for each organismal reference) must be supplied by the user; several

widely-used primers are already included in template configuration files.

To place all reads in the correct orientation, fastq files were reverse-complemented based on

information found in pyNAST alignment headers. cutadapt (version 1.18) (10) was used to identify

primer regions in fastq sequences with 0, 1, 2, and 6-mismatches to the primer. Only the 0, 1 and 2-

mismatch data were considered for plots/results following the established conventions from tools such

as SILVA *testprime.* The 6-mismatch threshold was used to exclude any spurious matches caused by

pyNAST misalignments or distantly homologous sequences; any sequence with greater than 6-

mismatches to the primer was not further considered. Based on primer comparisons to references, this

cutoff is likely to include even highly divergent taxa so long as they have a true primer binding site

(NB: this is not true for some primers designed to specifically target a particular group - e.g. the 806RB

primer does not contain a binding site for eukaryotic sequences). Primer coverage at each mismatch

cutoff was then calculated using only sequences where all bases in the primer region plus 5 leading and

trailing bases had a phred score of > 30. Plots shown in the main and supplemental text were generated

with python/cartopy (11), ggplot2 (12), veusz (13), and snakemake (1) (Fig 1; Fig 2, Fig S1-S29; Fig 3;

Fig S31-S32 respectively).

Classify workflow ("Snakefile-classify.smk"; Fig S31):

In order to determine which taxa were missed by a particular primer set and provide recommendations

about how to further improve existing primers' performance, we implemented an optional subsequent

workflow to taxonomically classify reads and provide summary statistics for each group. Briefly, full-

length fastq files from mismatched taxa (that is, SSU rRNA sequences with a primer-binding region but

not matching primers at a given threshold) are classified with VSEARCH 2.10.4 (14) and the SINTAX

algorithm (15) against the SILVA132 nr99 database (5) or PhytoRef (16) databases (for

cyanobacteria/plastids) that had been converted into ".udb" format. The same procedure was followed

for matches except a maximum of 5,000 reads were randomly subsampled using "filterbyname.sh" (4)

to reduce computation time.

Summary statistics and output files were then generated for each unique combination of group, primer,

and mismatch threshold (e.g. EUK, 926R, 0-mismatch) at order-level resolution since informative

taxonomic annotations could generally not be made at finer taxonomic levels. Summaries contain the

fraction of each taxon not matching the primer at a given mismatch threshold, as well as the fraction of

total sequences represented by these mismatches. In addition, multiple sequence alignments of

mismatches and their relative abundances are generated, with summary plots automatically generated

showing the relative abundances of particular mismatches both for the whole dataset and subsetted by

individual taxa. In addition, the same output files are generated for specific order-level taxa, allowing

an investigator to see whether a given primer covers a particular taxon of interest. For the datasets

analyzed in this study, raw output files/plots can be downloaded at https://osf.io/gr4nc/.

Our pipeline generates coverage information at the individual primer level since metagenomic reads

will rarely/never cover the entire amplicon region (depending on lengths of sequence reads and the

amplicon region). However, it is advantageous to present results in terms of predicted coverage for the

primer set. If the mismatches to the forward and reverse primer were completely non-overlapping (i.e.

the mismatches were to completely different organisms) then the mismatched fractions would be

simply additive ("worst case" as mentioned in main text and shown in Fig 3). In contrast, if organisms

overlapped between the forward and reverse primer, the largest mismatched fraction for that organism

could be used. We used the taxonomic assignments discussed above to estimate this fraction of

overlapping sequences ("rule normalize_summaries" and its dependent rules/scripts) and thus correct

the primer coverage values as shown in Fig 2.

Compare workflow ("Snakefile-compare2tags-16s.smk" and "Snakefile-compare2tags-18s.smk"):

This workflow compares the abundance of taxa quantified by universal 16S and 18S rRNA amplicons

using the 515Y/926R primers with abundances of taxa from metagenomic SSU rRNA in the same

region of the molecule targeted by the PCR primers. The primer region is targeted by subsetting

pyNAST alignments by primer coordinates. Only reads with at least 100bp overlap with the amplicon

region are retained, and subsetted reads from the three prokaryotic categories are merged for the 16S

rRNA comparison. For 18S rRNA reads that do not overlap (since the amplicon region is longer than

the 2x250 sequencing length employed here), they were subsetted to the trim length of the partial

amplicons (e.g. 220bp for forward read, 180 bp for reverse read).

These metagenomic sequences were then compared using BLASTn (17) (minimum 97% similarity,

requiring 100% overlap) against a sample-specific database of exact amplicon sequence variants

(ASVs) generated with DADA2 (18) that had a nonzero abundance in that particular sample. In order

to make a direct comparison between the two data sources, the ASVs were clustered for each sample

based on BLAST results with networkx (19) to account for the resolution mismatch between short

metagenomic reads and longer, denoised amplicons. Total relative abundances of each cluster were

plotted using seaborn (https://seaborn.pydata.org/) and sample-sample correspondence was evaluated

using least squared linear regression with the stats.linregress function in scipy (20). The R^2 value of this

regression was used to evaluate general trends in MG-ASV sample-sample correspondence, and was

plotted against both environmental covariates and the "worst-case" (simply additive) predicted primer

performance (Fig 3b).

Amplicon Analysis Pipeline Supplementary Information:

SSU rRNA partitioning was carried out with the database mentioned in the main text (osf.io/e65rs/)

using bbsplit.sh (4). These databases were constructed from SILVA132 (5) and PR2 (7) by first

clustering with CDHIT (21) at 95% identity to reduce total data volume and subsequently curated to

remove sequences that were extremely long or short relative to the average sequence length.

Demultiplexing and trimming steps were carried out with cutadapt (10), and trimming prior to

denoising (for 18S rRNA sequences) was carried out with bbduk.sh (4).

Supplementary Results:

Technical evaluation of pipeline:

After retrieval, quality-filtering, and subsetting to the primer region, dozens to hundreds of rRNA

sequences were recovered for each group, sample, read direction (fwd & rev reads), and primer

combination (Table 1). Considering the pelagic ocean datasets introduced above and discussed in detail

below (BiHoBa, TARA, MSPINA, MBARI, SPOT), we retrieved between 158-1148 primer-matching

regions for *Bacteria* for each combination (medians within datasets; Table 1), whereas lower numbers

were retrieved for the remaining three groups of *Archaea*, *Cyanobacteria*+Plastids and *Eukarya* (on

average 17-109, 46-186 and 32-202 respectively) most likely due to their lower relative abundances in

the environmental DNA. Among the primers, smaller number of sequences were identified for the two

primers located near the end of the SSU rRNA molecule. For example, for Tara samples, a median of

455 sequences were retrieved per sample for 27F (close to end of molecule) for *Bacteria* vs. 1299 per

sample for 338R (middle of molecule). Similarly, 58 eukaryotic sequences were retrieved per sample

with 1510R vs. 136 per sample for 1389F.

With the exception of regions near the end of the SSU rRNA molecule, sequence coverage was mostly

evenly distributed across the alignments to group-specific references. Some relatively small regions

had lower coverage, most likely due to their absence in dominant environmental taxa. For example, the

SAR11 isolate HTCC1062 is missing nucleotides (relative to the *E. coli* reference) corresponding to

small portions of loops in the V1, V2, V3, and V9 regions (22). We note also that phyloFlash retrieved

approximately double the total number of 18S rRNA sequences compared to a graftM (23) package

based on an alignment of the greengenes database (data.ace.uq.edu.au/public/graftm/7/) which was

employed in an earlier version of our pipeline. In addition, the coverage across the *S. cerevisae* 18S

rRNA reference was much more even with phyloFlash-retrieved 18S rRNA vs. graftM. No such bias

was noted for 16S rRNA sequences, suggesting this graftM package may be appropriate for 16S rRNA

retrieval but not 18S rRNA.

Primer Limitations and Potential Improvements:

Table S1 shows a summary of the top 3 most dominant mismatched orders for each primer and group

for the BiHoBa dataset. These samples mostly represent surface and upper mesopelagic environments

across all organismal size classes (Table 1), making it useful for exploring general patterns of primer

performance for oceanic ecosystems. Among the most prominent mismatches to *Archaea* are those

between *Nitrosopumilales* MGI *Archaea* and primers 341F or 1389F. Neither of these primers were

specifically designed for capturing archaeal sequences but our results show they could be modified to

include most *Archaea* as shown in Figs S1-S7 for 341F, and likely also for 1389F given that a universal

primer with high coverage exists for the same general region (1392R). For other primers, the dominant

mismatches include abundant groups such as MGII, and MGIII but we also note a smaller number of

mismatches from the poorly-described *Aigarchaeales* and *Woesearchaeia.*

For *Bacteria,* many of the abundant mismatches were already discussed in the main text. Other patterns

of particular note are mismatches to abundant members of the SAR86 clade and *Flavobacteriales* for

341F/338R primers. Together, these groups represent over 4% of the total sequences, meaning they

would likely be desirable to correct in the future. Readers should note that the dominant taxa in Table

S1 are the most abundant sequences, and there are a number of rare bacterial taxa that have very low

coverage. These include *Poribacteria,* a number of members of the candidate phyla radiation or CPR,

Rubrobacterales (*Actinobacteria*), and *Chlamydiales*. Interestingly, some members of these taxa (e.g.

Poribacteria, CPR) had more than one mismatch to the 926R primer set, meaning they are likely very

evolutionarily-divergent from other organisms and may require a more extensive primer redesign to

quantify them fully. However, we note that these organisms are very rare in these datasets, so their

absences are unlikely to bias overall community profiles in oxic marine environments and that the

additional degeneracies proposed in the main text will increase coverage for some of these taxa.

With respect to the cyanobacterial/chloroplast fraction, while the 515Y/926R primer pair achieves

nearly perfect coverage, the other primers generally have significant mismatches to one or more

abundant chloroplast groups, including the *Mamelliophyceae, Prasinophyceae,* and *Prymnesiophyceae.*

This means that care must be used in interpreting chloroplast results from some popular primer sets

which may miss important taxa.

Among eukaryotic (nuclear 18S rRNA) sequences, the most commonly mismatched taxa included

Dinoflagellata, Discicristata, Protalveolata, Ochrophyta, Charophyta, and *Metazoa.* As noted above,

the universal primer sets appear to perform nearly as well or better versus *Eukarya*-specific primers.

Improvements described in the main text do increase 18S rRNA coverage for the 515Y/926R primers,

but a fully comprehensive 18S rRNA primer may be less feasible since primer binding regions have a

much higher sequence diversity versus 16S rRNA. This may be due to large copy numbers of SSU

rRNA, some of which may become pseudogenized and undergo random evolutionary drift (24). In

practical terms, this means that some of these more rare variants will inevitably be missed by primers

since creating an overly-degenerate primer could cause problems with non-specific amplification

during PCR. The same may be true for some mitochondrial 16S rRNA genes that have become

pseudogenized after being incorporated into the host nuclear genome (25).

A Practical Consideration for the use of the 515Y/926R primer set:

One concern about using truly universal primers is the amplification of non-microbial sequences, e.g.

in the case of host-associated microbiomes. This has led to the design of prokaryotic-specific primer

sets (e.g. 515Y/806RB), that intentionally do not amplify 18S rRNA sequences. However, universal

primers retain the advantage of quantifying a broad swathe of the entire biological community together.

Therefore, such primers could be potentially be very useful if 18S rRNA sequences do not overwhelm

the SSU rRNA sequences in typical samples.

To test this, we quantified the total fraction of 18S rRNA (out of the total 16S + 18S rRNA) for the

unfractionated (>0.2µM) BiHoBa metagenomes. For these data, we show 18S rRNA sequences rarely

exceed ~1/3 of the total SSU rRNA sequences (Fig S29), indicating that universal primers will still be

useful even when whole seawater samples are analyzed, providing quantification of both prokaryotic

and eukaryotic communities simultaneously. The fraction of eukaryotic SSU rRNA relative to total

16S + 18S rRNA averaged 19.5 +/- 0.09 % (range = 0.024 - 0.7). Samples with large fractions of

eukaryotes (>0.25) were dominated by *Metazoa*, mainly composed of taxa such as *Arthropoda,*

Platyhelminthes, and *Hydrozoa.* These animal-derived sequences have the potential to overwhelm the

signal from other environmental eukaryotes, especially for unfractionated samples. This can be

alleviated to some extent by increasing overall sequence coverage such that sufficient non-metazoan

18S rRNA sequences can also be recovered.

Supplemental References Cited:

1. J. Köster, S. Rahmann, Snakemake—a scalable bioinformatics workflow engine. *Bioinformatics* **28**, 2520–2522 (2012).

2. S. Chen, Y. Zhou, Y. Chen, J. Gu, fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

3. H. R. Gruber-Vodicka, B. K. B. Seah, E. Pruesse, phyloFlash – Rapid SSU rRNA profiling and targeted assembly from metagenomes. *bioRxiv,* 521922 (2019).

4. B. Bushnell, BBMap short read aligner. *https://sourceforge.net/projects/bbmap/*.

5. C. Quast, *et al.*, The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucl. Acids Res.* **41**, D590–D596 (2013).

6. E. Clarke, komplexity: A method of assessing sequence complexity based on kmer frequencies. *https://github.com/eclarke/komplexity*.

7.   L. Guillou, *et al.*, The Protist Ribosomal Reference database (PR2): a catalog of unicellular eukaryote Small Sub-Unit rRNA sequences with curated taxonomy. *Nucleic Acids Res* **41**, D597–D604 (2013).

8.   J. G. Caporaso, *et al.*, PyNAST: a flexible tool for aligning sequences to a template alignment. *Bioinformatics* **26**, 266–267 (2010).

9.   P. J. A. Cock, *et al.*, Biopython: freely available Python tools for computational molecular biology and bioinformatics. *Bioinformatics* **25**, 1422–1423 (2009).

10.   M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**, 10–12 (2011).

11.   Met Office, *Cartopy: a cartographic python library with a matplotlib interface* (2010).

12.   H. Wickham, *ggplot2: Elegant Graphics for Data Analysis* (Springer-Verlag, 2009) https:/doi.org/10.1007/978-0-387-98141-3.

13.   J. Sanders, Veusz: A Scientific Plotting Package. *https://github.com/veusz/veusz*.

14.   T. Rognes, T. Flouri, B. Nichols, C. Quince, F. Mahé, VSEARCH: a versatile open source tool for metagenomics. *PeerJ* **4**, e2584 (2016).

15.   R. C. Edgar, SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS sequences. *bioRxiv*, 074161 (2016).

16.   J. Decelle, *et al.*, PhytoREF: a reference database of the plastidial 16S rRNA gene of photosynthetic eukaryotes with curated taxonomy. *Molecular Ecology Resources* **15**, 1435–1445 (2015).

17. S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment search tool. *J. Mol. Biol.* **215**, 403–410 (1990).

18. B. J. Callahan, *et al.*, DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods* **13**, 581–583 (2016).

19. A. A. Hagberg, D. A. Schult, P. J. Swart, Exploring Network Structure, Dynamics, and Function using NetworkX in *Proceedings of the 7th Python in Science Conference*, G. Varoquaux, T. Vaught, J. Millman, Eds. (2008), pp. 11–15.

20. P. Virtanen, *et al.*, SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nature Methods* **17**, 261–272 (2020).

21. W. Li, A. Godzik, CD-HIT: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

22. P. Yarza, *et al.*, Uniting the classification of cultured and uncultured bacteria and archaea using 16S rRNA gene sequences. *Nat Rev Micro* **12**, 635–645 (2014).

23. J. A. Boyd, B. J. Woodcroft, G. W. Tyson, GraftM: a tool for scalable, phylogenetically informed classification of genes within metagenomes. *Nucleic Acids Res* **46**, e59–e59 (2018).

24. S. R. Santos, R. A. Kinzie, K. Sakai, M. A. Coffroth, Molecular Characterization of Nuclear Small Subunit (ISS)-rDNA Pseudogenes in a Symbiotic Dinoflagellate (Symbiodinium, Dinophyta). *Journal of Eukaryotic Microbiology* **50**, 417–421 (2003).

25. D. Bensasson, D.-X. Zhang, D. L. Hartl, G. M. Hewitt, Mitochondrial pseudogenes: evolution's misplaced witnesses. *Trends in Ecology & Evolution* **16**, 314–321 (2001).

								Median filtered sequences per sample/group/primer/direction**			
Dataset name (**abbreviation**)	Sample and environment description	Sample size fraction(s)	Sample volume (L)	Reference(s)	SRA/ENA Accession(s)	# samples tested	Total # SSU rRNA sequences	Archaea	Bacteria	Cyanobacteria & Chloroplast	Eukarya
BioGEOTRACES / HOT / BATS (**BiHOBA**)	Mostly euphotic and upper mesopelagic ocean waters; primarily oligotrophic	>0.2µm	0.1	18-20	PRJNA385854	610	3.16E+07	36.9	256.9	46.3	55.5
Tara Oceans (**TARA**)	Euphotic, deep chlorophyll maximum, and mesopelagic samples	All with upper size fraction <3µm*	≥100	16	PRJEB1787, PRJEB1788, PRJEB4352, PRJEB4419	458	1.32E+08	109.1	1148	141.7	155.4
Malaspina Deep Metagenomes (**MSPINA**)	Deep ocean metagenomes from samples collected >2000m in depth	0.2 - 0.8 µm (FL), 0.8 - 20 µm (PA)	120	17	PRJNA329986 - PRJNA330161	58	3.46E+06	26.3	158.4	NA	32.6
MBARI bloom metagenome (**MBARI**)	Coastal surface seawater sampled during phytoplankton bloom across 52 days	0.22–5.0 µm	up to 1	21	PRJNA533622	84	1.32E+08	25.7	667.7	52.4	68.3
SPOT / Catalina metagenome (**SPOT**)	Surface seawater collected from the San Pedro Ocean Time-Series and Catalina island across 4 seasons	0.22-1 µm	20	22	PRJEB12234	8	5.04E+05	16.9	434.2	186.2	201.7
Saanich Inlet Metagenomes (**Saanich**)	Depths spanning the oxycline (10, 100, 120, 135, 150, and 200 m)	0.22-2.7 µm	1 or 20	23	PRJNA247822	87	1.37E+07	96.8	662.6	88.9	21.5
Guaymas sediment metagenomes (**GuaySed**)	Total DNA from > 10g hydrothermally-influenced sediment from 0-15cm depth	NA, whole sediment	NA, 10g sediment	24	PRJNA362212	4	1.95E+06	1047.3	3065.6	93	433.7

*<0.2µm; 0.2-1.6µm; 0.2-3µm; 0.8-3µm; **groups={1: Archaea,2:Bacteria less Cyanobacteria+Plastids,3:Chloroplast+Plastid,4:Eukarya}, direction=forward and reverse paired-end reads (analyzed separately)

201   Table S1: Data used in this study. Exact SRA/ENA Accessions available on github.com/jcmcnch/MGPrimerEval in config files.

Table S2: Primers tested in this study

Primer	SSU rRNA region targeted	Organismal coverage	Primer sequence (5' - 3')	Reference(s)	Reference(s)
27F	V1-V2	16S only	AGRGTTTGATYMTGGCTCAG	11	Fortunato et al., 2012
338R	V1-V2	16S only	TGCWGCCWCCCGTAGGWGT	11	Fortunato et al., 2012
341F	V3-V4	16S (Bacteria-only)	CCTACGGGNGGCWGCAG	10	Klindworth et al., 2013
785R	V3-V4	16S only	GACTACHVGGGTATCTAATCC	10	Klindworth et al., 2013
V4F	V4	18S only	CCAGCASCYGCGGTAATTCC	12	Stoeck et al., 2010
V4R	V4	18S only	ACTTTCGTTCTTGATYRA	12	Stoeck et al., 2010
V4RB	V4	18S only	ACTTTCGTTCTTGATYRR	13	Balzano et al., 2015
515Y	V4/V4-V5	16S and 18S	GTGYCAGCMGCCGCGGTAA	2-4	Parada et al., 2016
806RB	V4	16S only	GGACTACNVGGGTWTCTAAT	2-4	Apprill et al., 2015
926R	V4-V5	16S and 18S	CCGYCAATTYMTTTRAGTTT	2-4	Walters et al., 2015
926wF	V6-V8	16S and 18S	AAACTYAAAKGAATTGRCGG	14	Wilkins et al., 2013
1392R	V6-V8	16S and 18S	ACGGGCGGTGTGTRC	14	Wilkins et al., 2013
1389F	V9	18S only	TTGTACACACCGCCC	15	Amaral-Zettler et al., 2009
1510R	V9	18S only	CCTTCYGCAGGTTCACCTAC	15	Amaral-Zettler et al., 2009

Table S3: Top 3 most abundant mismatched orders for each organismal category in the BioHOBA dataset. Values in parentheses represent the fraction of sequences that do not perfectly match the primer at a 0-mismatch threshold as a fraction of total SSU rRNA sequences (matched + mismatched). For example, for the primer 785R, mismatches to SAR11 primer-binding regions represent 40% of total identified bacterial sequences (excluding cyanobacteria + plastids), whereas the most abundant mismatched taxon for 515Y is Rickettsiales at 0.2% total bacterial SSU rRNA read abundance. Primers that have "NA" for particular organismal category did not have a recognizable primer-binding region. This often represents an intentional design choice to avoid what was considered off-target amplification. For example, the 806RB primer was designed not to amplify 18S (Eukaryote) sequences.

Primer	Archaea	Bacteria (without Cyanobacteria and plastid 16S)	Cyanobacteria (including plastid 16S)	Eukarya (18S nuclear SSU rRNA only)
27F	NA, no apparent primer binding site	Rickettsiales (0.008), Ectothiorhodospirales (0.008), SAR11 clade (0.002)	Prasinococcales (0.011), Cyanobactera;Subsection I (0.003), Chrysophyceae-Synurophyceae (0.002)	NA, no apparent primer binding site
338R	NA, no apparent primer binding site	SAR86 clade (0.027), Flavobacteriales (0.015), SAR11 clade (0.009)	Prymnesiophyceae (0.123), Bacillariophyta (0.006), Prasinococcales (0.005)	NA, no apparent primer binding site
341F	Nitrosopumilales (0.51), MG II (0.41), MG III (0.07)	SAR86 clade (0.028), Flavobacteriales (0.016), SAR11 clade (0.016)	Prymnesiophyceae (0.121), Bacillariophyta (0.006), Prasinococcales (0.005)	NA, 5-mismatch to reference
785R	MG II (0.003), Nitrosopumilales (0.002), MG III (0.0004)	SAR11 clade (0.40), Marinimicrobia (SAR406_clade); unclassified order/class (0.006), Verrucomicrobiae;Arctic97B-4 (0.003)	Mamiellophyceae (0.103), Prymnesiophyceae (0.006), Cyanobacteria;SubsectionI (0.001)	NA, 3-mismatch to reference
V4F	NA, 5-mismatch to reference	NA, 3-mismatch to reference	NA, 4-mismatch to reference	Dinoflagellata (0.022), Metazoa (Animalia) (0.010), Protalveolata (0.007)
V4R	NA, 8-mismatch to reference	NA, 7-mismatch to reference	NA, many mismatches to reference	Dinoflagellata (0.081), Discicristata (0.027), Protalveolata (0.026)
V4RB	NA, 8-mismatch to reference	NA, 7-mismatch to reference	NA, many mismatches to reference	Dinoflagellata, (0.040), Discicristata (0.013), Protalveolata (0.012)
515Y	Nitrosopumilales (0.003), MG II (0.002), MG III (0.0003)	Rickettsiales (0.002), SAR11 clade (0.001), Chlamydiales (0.0006)	Cyanobacteria;Subsection I (0.003), Prymnesiophyceae (0.001), Pelagophyceae (0.000)	Dinoflagellata (0.021), Discicristata (0.014), Metazoa (Animalia) (0.010)
806RB	Nitrosopumilales (0.002), MG II (0.002), Aigarchaeales (0.001)	Marinimicrobia (SAR406_clade); unclassified order/ class (0.006), Flavobacteriales (0.005), SAR202 clade (0.002)	Mamiellophyceae (0.108), Prymnesiophyceae (0.014), Prasinophyceae (0.013)	NA, many mismatch to reference
926R	Nitrosopumilales (0.016), MG II (0.001), Woesearchaeia; unclassified order (0.0007)	Ectothiorhodospirales (0.007), Rickettsiales (0.004), SAR11 clade (0.002)	Cyanobacteria;Subsection I (0.003), Prymnesiophyceae (0.000) Dictyochophyceae (0.000)	Dinoflagellata (0.015), Protalveolata (0.012), Metazoa (Animalia) (0.010)
926wF	Nitrosopumilales (0.016), MG II (0.001), Woesearchaeia; unclassified order (0.0007)	Rhodospirillales (0.041), Ectothiorhodospirales (0.007), Rickettsiales (0.004)	Cyanobacteria;Subsection I (0.003), Prymnesiophyceae (0.000) Dictyochophyceae (0.000)	Metazoa (Animalia) (0.040), Ochrophyta (0.033), Dinoflagellata (0.016)
1392R	Nitrosopumilales (0.002), Woesearchaeia; unclassified order (0.002), MG II (0.001)	SAR11 clade (0.003), Clostridiales (0.001), Rickettsiales (0.001)	Mamiellophyceae (0.227), Dictyochophyceae (0.011), Bacillariophyta (0.002)	Dinoflagellata (0.014), Protalveolata (0.010), Metazoa (Animalia) (0.008)
1389F	Nitrosopumilales (0.53), MG II (0.37), MG III (0.05)	Dadabacteriales (0.002), Clostridiales (0.002), SAR11 clade (0.002)	Mamiellophyceae (0.167), Cyanobacteria;Subsection I (0.103), Trebouxiophyceae (0.055)	Dinoflagellata (0.013), Protalveolata (0.008), Metazoa (Animalia) (0.006)
1510R	NA, 4-mismatch to reference	NA, 4-mismatch to reference	Prasinophyceae (0.393), Pelagophyceae (0.375), Pavlovophyceae (0.054)	Metazoa (Animalia) (0.055), Charophyta (0.036), Protalveolata (0.012)

Table S4: Oligonucleotide mixtures specified for the new primers 515Yp-min and 926Rp-min.

515Yp-min	926Rp-min
GTGCCAGCAGCAGCGGTAA	CCGTCAATTCGTTTAAGTTT
GTGCCAGCACCCGCGGTAA	CCACCAGAACATTTAAGTTT
GTGCCAGCAGACGCGGTAA	CCATCAATTCATTTGAGTTT
GTGCCAGCAGCCGCGGTAA	CCGCCAATTCATTTAAGTTT
GTGCCAGCAGCCGCGGTCA	CCGCCAATTCATTTGAGTTT
GTGCCAGCAGCCGCGGTGA	CCGCCAATTCCCTTAAGTCT
GTGCCAGCAGCTGCGGTAA	CCGCCAATTCCCTTAAGTTT
GTGCCAGCAGTCGCGGCAA	CCGCCAATTCCTATAAGTTT
GTGCCAGCAGTCGCGGTAA	CCGCCAATTCCTTCAAGTTT
GTGCCAGCCGCCGCGGTAA	CCGCCAATTCCTTTAAGTCG
GTGGCAGCCGCCGCGGTAA	CCGCCAATTCCTTTAAGTCT
GTGTCAGCAGCCGCGGTAA	CCGCCAATTCCTTTAAGTTA
GTGTCAGCCGCCGCGGTAA	CCGCCAATTCCTTTAAGTTT
	CCGCCAATTCCTTTGAGTTT
	CCGCCAATTGGTTTAAGTTT
	CCGCCAATTTCTTTAAGTTT
	CCGCCAATTTCTTTGAGTTT
	CCGTCAACTCATTTGAGTTT
	CCGTCAATCCCTTTAAGTTT
	CCGTCAATGCCTTTAAGTTT
	CCGTCAATTACTTTAAGTTT
	CCGTCAATTACTTTGAGTTT
	CCGTCAATTCATTTAAGTTT
	CCGTCAATTCATTTGAGTTT
	CCGTCAATTCCCTTGAGTTT
	CCGTCAATTCCTTCAAGTTT
	CCGTCAATTCCTTTAAATTT
	CCGTCAATTCCTTTAAGTTT
	CCGTCAATTCCTTTGAATTT
	CCGTCAATTCCTTTGAGTTC
	CCGTCAATTCCTTTGAGTTT
	CCGTCAATTCCTTTGGGTTC
	CCGTCAATTCCTTTGGGTTT
	CCGTCAATTCTTTTGAGTTT
	CCGTCAATTTATTTGAGTTT
	CCGTCAATTTCTTTAAGTTT
	CCGTCAATTTCTTTGAGTTT
	CCGTCTATTCCTTTGAGTTT

 Fig S1: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Archaea* using

 metagenomes derived from Guaymas Basin sediments.

Fig S2: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Archaea* using

metagenomes derived from Saanich Inlet water samples.

Fig S3: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Archaea* using

metagenomes derived from BioGEOTRACES/BATS/HOT water samples.

Fig S4: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Archaea* using

metagenomes derived from SPOT/Catalina water samples.

Fig S5: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Archaea* using

metagenomes derived from Malaspina deep-sea water samples.

Fig S6: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Archaea* using

metagenomes derived from MBARI bloom water samples.

Fig S7: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Archaea* using

metagenomes derived from Tara Oceans water samples (<3µm).

Fig S8: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Cyanobacteria* and

plastids using metagenomes derived from Guaymas Basin sediments.

Fig S9: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Cyanobacteria* and

plastids using metagenomes derived from Saanich Inlet water samples.

Fig S10: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Cyanobacteria* and

plastids using metagenomes derived from SPOT/Catalina water samples.

Fig S11: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Cyanobacteria* and plastids using metagenomes derived from BioGEOTRACES/BATS/HOT water samples.

Fig S12: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Cyanobacteria* and

plastids using metagenomes derived from Malaspina deep-sea water samples.

Fig S13: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Cyanobacteria* and

plastids using metagenomes derived from MBARI bloom water samples.

Fig S14: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Cyanobacteria* and

plastids using metagenomes derived from Tara Oceans water samples (<3µm).

Fig S15: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Bacteria* (not

including *Cyanobacteria* and plastids) using metagenomes derived from Guaymas Basin sediment

samples.

Fig S16: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Bacteria* (not

including *Cyanobacteria* and plastids) using metagenomes derived from Saanich Inlet water samples.

Fig S17: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Bacteria* (not

including *Cyanobacteria* and plastids) using metagenomes derived from BioGEOTRACES/BATS/HOT

water samples.

Fig S18: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Bacteria* (not

including *Cyanobacteria* and plastids) using metagenomes derived from SPOT/Catalina water samples.

Fig S19: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Bacteria* (not

including *Cyanobacteria* and plastids) using metagenomes derived from Malaspina deep-sea water

samples.

Fig S20: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Bacteria* (not

including *Cyanobacteria* and plastids) using metagenomes derived from MBARI bloom water samples.

Fig S21: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Bacteria* (not

including *Cyanobacteria* and plastids) using metagenomes derived from Tara Oceans water samples

(<3µm).

Fig S22: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Eukarya* using

metagenomes derived from Guaymas Basin sediment samples.

Fig S23: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Eukarya* using

metagenomes derived from Saanich Inlet water samples.

Fig S24: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Eukarya* using

metagenomes derived from BioGEOTRACES/BATS/HOT water samples.

Fig S25: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Eukarya* using

metagenomes derived from SPOT/Catalina water samples.

Fig S26: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Eukarya* using

metagenomes derived from Malaspina deep-sea water samples.

Fig S27: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Eukarya* using

metagenomes derived from MBARI bloom water samples.

Fig S28: Predicted individual primer coverage allowing 0, 1, and 2-mismatches for *Eukarya* using

metagenomes derived from Tara Oceans water samples (<3µm).

Fraction of 18S SSU rRNA from BiHoBa Metagenomes

293   Figure S29: Violin plot with inlaid boxplot showing the fraction of eukaryotic (18S) SSU rRNA as a

294   fraction of total across BATS, HOT, and bioGEOTRACES metagenomic datasets. The violin plot

295   provides a visual description of the distribution of data - i.e. in this case showing that the vast majority

296   of samples had < 30% of SSU rRNA reads accounted for by 18S rRNA sequences.

Figure S30: Rulegraph for "Snakefile-compute.smk" portion of workflow. Briefly, SSU rRNA is identified using phyloFlash from fastp quality-filtered fastq reads, which are then sorted with bbsplit.sh into 4 categories, and aligned to SSU rRNA references with pyNAST after removing low-complexity sequences. For each group and primer, matches to primer regions are then identified using custom python scripts and cutadapt (at 0, 1, 2, and 6-mismatches), quality filtered to omit sequences where the primer region plus 5 leading and trailing bases has one individual base with a phred score below 30. Matches are then summarized in plain-text output and ggplot2 graphs.

Figure S31: Rulegraph for "Snakefile-compare.smk" portion of workflow to intercompare metagenome and amplicon data.