**Notch signaling represses cone photoreceptor formation through the**

**regulation of retinal progenitor cell states**

Xueqing Chen[1,2], Mark M. Emerson[1,2,3*]

[1] Biology PhD Program, The Graduate Center, The City University of New

York, New York, NY, 10016

[2] Department of Biology, The City College of New York, The City University of

New York, New York, NY, 10031

[3] Biochemistry PhD Program, The Graduate Center, The City University of

New York, New York, NY, 10016

*Corresponding author: memerson@ccny.cuny.edu

Email addresses:

Xueqing Chen: xchen2@gradcenter.cuny.edu

Mark Emerson: memerson@ccny.cuny.edu

Keywords: retinal progenitor cells, multipotency, cis-regulation, cone

photoreceptors, Notch signaling, electroporation

**Abstract**

Vertebrate cone photoreceptor formation is a multistep process. First, multipotent retinal progenitor cells generate genetically-defined restricted/neurogenic progenitor cells and these cells then divide to preferentially produce cones and horizontal cells. Notch signaling represses cone formation and maintains the proliferative potential of retinal progenitor cells. However, the mechanisms through which it affects these processes are unknown. Here we use cell type specific inhibition of Notch signaling to localize the primary role of Notch signaling during cone genesis to the regulation of restricted retinal progenitor cells from multipotent retinal progenitor cells. Notch signaling inhibition in restricted progenitor cells does not alter the number of cones derived from these cells but does affect horizontal cell development. Cell cycle promotion is not a primary effect of Notch signaling but an indirect effect on progenitor cell state transitions that leads to depletion of the multipotent progenitor cell population. Taken together, this suggests that the roles of Notch in cone photoreceptor formation and cell cycle promotion are both mediated by a localized function in multipotent retinal progenitor cells to repress the formation of restricted progenitor cells.

**Introduction**

Notch signaling is an evolutionarily conserved signaling pathway that functions in diverse cellular contexts to drive both cell cycle and cell fate decisions. In canonical Notch signaling, extracellular binding of a member of the Delta/Serrate/Lag family of proteins to the extracellular portion of the transmembrane Notch receptor leads to a proteolytic cleavage that releases the Notch intracellular domain (NICD) from the membrane (Bray, 2006; Kopan and Ilagan, 2009). The NICD then translocates to the nucleus, and through a primary interaction with the MAML protein, forms a tripartite complex with the Rbpj transcription factor that affects transcription of target genes (Mumm and

Kopan, 2000; Wu et al., 2000). Notch signaling can act iteratively and

participate in several distinct decisions that occur during the process of

formation of a particular cell type. For example, during the formation of

Drosophila sensory neurons from ectoderm, Notch signaling is critical in three

successive cell fate decisions (Artavanis-Tsakonas et al., 1991; Bray, 1998).

Though the role of Notch signaling in cell fate decisions in Drosophila and

C. elegans has been implicated in binary cell fate choices (Lai, 2004), the

specific role of Notch signaling in the vertebrate nervous system has not been

as clear. During the formation of the vertebrate retina, a functional role for

Notch signaling in both proliferation and in cell fate choice has been identified.

Induction of conditional loss-of-function alleles for the Notch1 receptor during

early retinogenesis leads to cell cycle exit and increased cone photoreceptor

formation (Jadhav et al., 2006; Yaron et al., 2006). In one of these studies, a

loss of horizontal cells (HCs) was also described (Yaron et al., 2006).

Similarly, conditional Rbpj loss-of-function increases the formation of cone

photoreceptors while also promoting the formation of retinal ganglion cells

(RGCs) (Riesenberg et al., 2009). In both cases, Notch1 and Rbpj loss-of-

function leads to a decrease in proliferation. Conditional loss of Notch1 in the

later postnatal period, after the window of cone genesis, reveals that rod

photoreceptors are also increased, which suggests that Notch signaling plays

a more general role in repression of the photoreceptor fate (Jadhav et al.,

2006). In addition, it has been suggested in this postnatal context that Notch

signaling can function in postmitotic cell fates to repress the rod fate

(Mizeracka et al., 2013). Despite these significant advances, the mechanisms

through which Notch affects these processes are still unknown and has

implications for both visual system evolution and retinal therapy development.

While previous studies have implicated Notch signaling in the generation

of cone and rod photoreceptors, it is not yet clear at which step in their

formation that it acts. Recently, the formation of cones has been identified as

a multistep process. During the early stages of retinal formation when cone

photoreceptors are preferentially generated, multipotent retinal progenitor

cells (RPCs) divide to form restricted (also known as neurogenic RPCs) RPCs

that can be identified by the activity of the ThrbCRM1 cis-regulatory element,

the expression of specific transcription factors (Otx2, OC1 and Olig2), and the

decreased expression of canonical RPC markers, such as Vsx2

(Buenaventura et al., 2018; Emerson and Cepko, 2011; Emerson et al., 2013;

Jean-Charles et al., 2018). These restricted RPCs lack the diverse cell fate

potential of the multipotent RPC and instead preferentially divide to form cone

and HCs as well as a minor number of RGCs (Cepko, 2014; Emerson et al.,

2013; Schick et al., 2019). Thus, Notch signaling could mediate cone genesis

through regulation of the multipotent to restricted RPC transition, the cell fate

choices of the restricted RPC daughter cells, or in the production of cones

directly from multipotent RPCs or other cell types.

In this study, we used defined cis-regulatory elements to both quantitate

the effects of Notch signaling inhibition and also to limit interference with

Notch signaling to specific cell types. We determined that during the formation

of cone photoreceptors, inhibition of Notch signaling specifically promoted the

formation of restricted RPCs from multipotent RPCs but did not promote the

cone fate in the daughter cells of restricted RPCs. As expected for such a shift

in cycling populations, there was minimal short-term effect on proliferation as

measured by EdU labeling. Taken together, this suggests that a primary role

of Notch signaling during cone photoreceptor genesis is to regulate the

formation of restricted RPC states from multipotent RPCs.

**Results**

**Evaluation of Notch reporter activity in response to MAML dominant-**

**negative**

To provide precise and rapid interference with Notch signaling, we used a

dominant-negative approach. We first created a chicken version of a

previously validated, and widely used, C-terminal truncation version of MAML

that retains the ability to interact with the Notch intracellular domain but lacks

the domains necessary to induce a transcriptional response (referred to as

MAML-DN) (Tan et al., 2016; Weng et al., 2003). The MAML1 gene was used,

as it was the MAML gene most highly expressed in the developing chicken

retina according to reported RNA-Seq data (Buenaventura et al., 2018). To

test the effectiveness of the MAML-DN construct, we used previously reported

Notch reporters Hes5::GFP (Vilas-Boas et al., 2011), Hes1::GFP (Mizutani et

al., 2007) and 12XCSL::EGFP (Hansson et al., 2006). Hes1 and Hes5 are two

previously used transcriptional reporters of Notch signaling derived from

characterized Notch target genes. The Hes1 gene is broadly expressed in

RPCs while the Hes5 gene is dynamic and expressed in RPCs and

differentiating retinal ganglion neurons (Riesenberg et al., 2018). The 12XCSL

reporter construct is a synthetic element composed of 12 multimerized Rbpj

motifs.

E5 chicken retinas were co-electroporated ex vivo with one of the Notch

GFP reporters, a ubiquitous electroporation control (CAG::Nucßgal), and with

or without the MAML-DN construct. After two days of culture, retinas were

imaged by confocal microscopy or analyzed by flow cytometry (Fig. 1A). In

control retinas, the Hes1::GFP reporter was broadly distributed across the

retina and appeared to be in the majority of targeted cells (Fig. S1). This

pattern was unexpected for a Notch reporter given the enriched expression of

Notch in RPCs and the restricted expression of Hes1 expression. In addition,

the Hes1::GFP reporter activity levels did not appear to be affected by MAML-

DN expression (Fig. S1). In contrast, the Hes5::GFP reporter was active

primarily in the neuroblast layer of the retina, where multipotent RPCs are

located, and the reporter activity was drastically abrogated in response to

MAML-DN introduction (Fig. 1B). The 12XCSL::EGFP reporter showed low

expression levels in targeted cells, however, MAML-DN also decreased this

reporter (Fig. S1). Strikingly, we found in all the experimental groups that the

retinal distribution of the electroporated population was altered, with cells

found predominantly in the scleral region where photoreceptor cells and

Otx2/OC1+ RPCs are enriched, as well as the RGC layer, but not in the

middle portion of the retina where most multipotent RPCs are localized (Fig.

1B) (Buenaventura et al., 2018). This suggested that there could be cell fate

changes in response to the MAML-DN mutant.

To quantify Notch reporter activity, we used a flow cytometry assay and

calculated the percentage of GFP-positive cells within the electroporated cell

population (Fig. 1C). The quantification supported the conclusions of the

confocal imaging results, as Hes5::GFP reporter activity showed a significant

decrease in response to MAML-DN introduction. We also confirmed the

effects of inhibited Notch signaling using dominant-negative forms of Rbpj

(Rbpj-EnR), in which the full-length Rbpj was fused with an engrailed

repressor at either the N-terminus or C-terminus and co-electroporated with

the same Notch reporters. Hes5 reporter activity significantly decreased in the

presence of both dominant-negative constructs (Fig. 1C and Fig. S2). In

summary, the Notch reporter Hes5::GFP displayed the most robust readout of

Notch signaling activity in this developmental context and the MAML-DN and

Rbpj-EnR constructs were all effective at inhibiting Notch signaling as gauged

by the Hes5 reporter.

**Inhibition of Notch promotes the formation of restricted progenitor cells**

A mouse knockout of the Notch1 receptor leads to an increase in cone

photoreceptors (Jadhav et al., 2006; Yaron et al., 2006), but the mechanism

behind this phenotype is unknown. One possibility is that in the knockout,

multipotent RPCs produce more of the RPC type that is restricted to cone and

HC fates, which in turn divide to form cones. The cis-regulatory element

ThrbCRM1 provides a robust tool to answer the question as it is active

specifically in this restricted RPC type (Emerson et al., 2013).

We first tested the effect of Notch inhibition in retinal development using

DNA reporters, including reporters with enrichment in multipotent RPCs

(VSX2ECR4; Buenaventura et al., 2018), cone/HC restricted RPCs

(ThrbCRM1; Emerson et al., 2013), subsets of cones (ThrbCRM2; Emerson et

al., 2013; Schick et al., 2019) and HCs/RGCs (OC1ECR22; Patoori et al.,

2020 and manuscript in preparation). E5 chicken retinas were electroporated

with a co-electroporation control, with or without the MAML-DN, and assessed

by confocal microscopy after two days culture. In retinas in which MAML-DN

was introduced, the number of ThrbCRM1 reporter-labeled restricted RPCs

was visibly increased, while the number of VSX2ECR4-labeled multipotent

RPCs was decreased and presented with altered morphology (Fig. 2A). The

number of ThrbCRM2 reporter-labeled cones and OC1ECR22 reporter-

labeled HCs/RGCs were qualitatively unchanged at this timepoint (Fig. S3).

The quantification of these cell types using flow cytometry statistically

confirmed these effects (Fig. 2B). This suggested that the broad inhibition of

Notch signaling via CAG::MAML-DN promotes the formation of ThrbCRM1-

active restricted RPCs. The down-regulation of VSX2ECR4 reporter-positive

RPCs, and significantly increased number of VSX2ECR4/ThrbCRM1 double

reporter-positive RPCs (Fig. 2B), suggests a transition from multipotent RPCs

to ThrbCRM1-positive RPCs in CAG::MAML-DN mutant retina. Intriguingly,

ThrbCRM2-labeled cones did not significantly increase after 2 days culture,

which could indicate a delay in cone genesis at this time point.

The increase in ThrbCRM1 reporter-positive cells could be due to an

increase in the ThrbCRM1 state, but it could also represent an increase in

their daughter cell production (cones, HCs, or RGCs) that could be missed by

the specific reporters used (Schick et al., 2019). We first examined Otx2 and

Visinin expression to confirm the reporter-observed effects, as these

endogenous markers are expressed in both photoreceptors and restricted

RPCs (Buenaventura et al., 2018; Emerson et al., 2013). In retinas with

introduction of the CAG::MAML-DN construct, there was a significant increase

of electroporated Visinin and Otx2-labeled cells, consistent with the reporter

results (Fig. 3A, B). Assessment of Olig2 expression, which is expected to be

expressed in ThrbCRM1-labeled RPCs and not cones or HCs, was

significantly increased, which suggests that the ThrbCRM1 reporter-positive

population observed at this 2 day period represents the restricted RPC state

(Fig. 3A and 3C) (Emerson et al., 2013; Ghinia Tegla et al.; Hafler et al.,

2012). In alignment with this observation and the ThrbCRM2 reporter result,

expression of the cone gene Lhx4 was not qualitatively changed (Fig. S3)

(Buenaventura et al., 2019). In contrast, the representation of the H1 HC

population (marked by Lim1 and AP2α; Fischer et al., 2007) that is normally

generated by ThrbCRM1 restricted RPCs is decreased in response to

CAG::MAML-DN (Figure 3B). No significant changes in the H2-4 HC/RGC

marker Isl1 or the Brn3 family of RGC markers were observed (Fig. S4)

(Fischer et al., 2007; Liu et al., 2000). Taken together, these data suggest that

the primary effect of Notch inhibition at this timepoint is to increase the

formation of the restricted RPCs that generate cones and HCs.

To further validate that these MAML-DN effects were related to Notch

signaling, we used flow cytometry to assess the effects of dominant-negative

forms of Rbpj on these same DNA reporters and cell markers. Similar effects

were observed upon introduction of both Rbpj-EnR constructs in support of a

basis in Notch signaling (Fig. S5). However, it was observed that the

reduction of HCs in response to the Rbpj-EnR constructs was more severe

than with MAML-DN, which is consistent with a Notch-independent role for

Rbpj to function in a complex with Ptf1a, a known regulator of HC genesis

(reviewed in Johnson and MacDonald, 2011).

**Notch inhibition does not affect the number of cones produced from**

**restricted RPCs**

To test whether MAML-DN also induces a shift from HCs to cones within

ThrbCRM1 reporter-positive RPCs, we replaced the ubiquitous CAG promoter

upstream of MAML-DN with the ThrbCRM1 specific enhancer, so that MAML-

DN would only be expressed in ThrbCRM1-active restricted RPCs. In contrast

to CAG::MAML-DN, introduction of this ThrbCRM1::MAML-DN into E5 retinas

did not alter the number of ThrbCRM1::GFP labeled RPCs. This suggests that

the ectopic ThrbCRM1 cells observed with the CAG promoter are not due to

induction of increased proliferation within the ThrbCRM1 population. Instead,

this supports a role for Notch in the multipotent RPC population, in which the

ThrbCRM1 element is not active (Fig. 4B).

To genetically maintain labeling in post-mitotic daughter cells, we used a

lineage tracing strategy to identify the retinal cells that have a ThrbCRM1

history (Fig. 4A) which were previously shown to be predominantly cones and

H1 HCs (Schick et al., 2019). A driver plasmid was used in which the PhiC31

recombinase is transcriptionally regulated by the cis-regulatory element

ThrbCRM1. In the responder plasmid, there is a STOP codon between a

broadly active CAG promoter and GFP, which is flanked by two recombinase

target sites. Excision of the STOP sequence upon ThrbCRM1-driven

expression of PhiC31 recombinase, will lead to GFP expression in cells with a

history of ThrbCRM1 activity (Schick et al., 2019).

The lineage tracing plasmids were introduced into the E5 retina, and then

the percentage of cones and HCs within the GFP-labeled retinal cells was

quantified using flow cytometry after two days culture. Intriguingly, the number

of cone photoreceptors did not change, as assessed by the expression of

Visinin and Otx2, while the percentage of Lim1 and AP2α-labeled HCs was

reduced (Fig. 4C and 4D). A small percentage of Isl1-labeled HCs and Brn3-

labeled RGCs are formed from ThrbCRM1 restricted RPCs, and no significant

changes in the populations were detected through cell counting in response to

introduction of the ThrbCRM1::MAML-DN (Fig. S6). Thus, the observed

reduction in HCs did not correspond to increased formation of cone

photoreceptors or Isl1-labeled HCs and RGCs in the daughter cell population

of ThrbCRM1 restricted RPCs. These unidentified cells potentially become

other cell types that cannot be identified by the cell markers used.

**Effects of MAML-DN on retinal cell proliferation and multipotency**

Previous analysis of mouse Notch1 conditional knockout retinas identified a reduction in BrdU-positive cells, which suggested that loss of Notch signaling promoted premature cell cycle exit and down-regulation of cell proliferation (Jadhav et al., 2006; Yaron et al., 2006). However, the difference in the percentages of BrdU-positive cells in wild-type and Notch1 CKO mutant is relatively small. To investigate the role of Notch in retinal cell proliferation, we co-electroporated CAG::MAML-DN and CAG::TdT into E5 chick retinas and after two days culture pulsed with EdU for 1 hour to label proliferating cells. We first quantified the percentage of EdU/CAG::TdT double-positive cells in all the electroporated cells using flow cytometry. Intriguingly, the population of proliferating cells does not significantly change compared to the control (Fig. 5A). Introduction of the Rbpj dominant-negative construct resulted in similar effects on electroporated EdU-positive cells (Fig. S7).

These data suggest that Notch signaling does not directly control cell proliferation in retina. Although inhibition of Notch does not drastically alter the overall number of RPCs within this timeframe, the experiments described above suggest that it causes a shift from multipotent RPCs to restricted RPCs with a significant increase in ThrbCRM1-positive RPCs, a reduction of VSX2ECR4-positive RPCs, and a large increase in the number of ThrbCRM1/VSX2ECR4 double reporter-positive cells (Fig. 2B). To confirm that these identified populations are indeed proliferating, we co-electroporated ThrbCRM1 and VSX2ECR4 reporters and pulsed with EdU just prior to harvest to label RPCs. Flow cytometry was used to quantify the EdU/ThrbCRM1 and EdU/VSX2ECR4 double-positive cells in the electroporated population. We also measured the percentage of EdU/ThrbCRM1/VSX2ECR4 triple-positive cells versus the combination of EdU/ThrbCRM1 and EdU/VSX2ECR4 double-positive cells (Fig. 5B and 5D). There was a significant up-regulation of ThrbCRM1+/EdU+ cells as well as ThrbCRM1+/VSX2ECR4+/EdU+ cells that indicated a transition from

VSX2ECR4-active RPCs to ThrbCRM1-active RPCs within the EdU+

population (Fig. 5C). In addition, immunostaining with the ThrbCRM1 marker

Otx2 and the multipotent RPC marker VSX2 showed a down-regulation of

VSX2+/EdU+ cells and an up-regulation of Otx2+/EdU+ cells, validating a shift

between VSX2+ RPCs and ThrbCRM1-active RPCs (Fig. S8).

To further determine if the induced ThrbCRM1 RPCs were derived from

multipotent RPCs, we used the VSX2ECR4 element that is active in this

population to drive expression of PhiC31 recombinase and conduct lineage

tracing of this population. The number of VSX2ECR4/ThrbCRM1 double

reporter-positive cells increased 5-fold in response to CAG::MAML-DN (Fig.

5E). This increase was consistent with the majority of new ThrbCRM1

reporter-positive cells being derived from multipotent RPCs, which suggests

that Notch inhibition specifically affects the transition from the multipotent to

restricted RPC state.

**Temporal parameters of the transition from multipotent RPCs to**

**restricted RPCs**

We were interested in determining the timeframe with which multipotent

RPCs transition to restricted RPCs. To test if the changes in multipotent RPC

numbers could be detected during their first cell cycle after electroporation, we

electroporated E5 chick retinas with VSX2ECR4::GFP, ThrbCRM1::TdT, the

co-electroporation reporter CAG::iRFP, and either CAG::MAML-DN or control

conditions. Retinas were harvested at 8 hours after electroporation, which is a

time before multipotent RPCs would be expected to complete one cell cycle.

There were no significant differences in the number of VSX2ECR4-marked

multipotent RPCs or ThrbCRM1-marked restricted RPCs (Fig. 6A). In

addition, the VSX2ECR4/ThrbCRM1 double reporter-positive cells were

present in a very low level as observed in wild type in both MAML-DN and

Rbpj-EnR mutants (Fig. 6B and Fig. S7). After 20 hours culture, the

ThrbCRM1-active population started to increase slightly but not nearly as

much as observed after two days (Fig. 6C). However, the transition from multipotent RPCs to restricted RPCs showed a small but significant up-regulation, as the VSX2ECR4/ThrbCRM1 double reporter-positive cells increased (Fig. 6C). Examination of Otx2 and VSX2 endogenous markers revealed a more robust response, with a decreased VSX2-positive population and an increased Otx2-positive one after 20 hours culture (Fig. 6D). Taken together, these data suggest that the Notch inhibition transition did not occur at 8 hours post-electroporation, but could be detected after 20 hours, and increased further at 2 days post-electroporation. This suggests that multipotent RPCs do not immediately transition to a restricted RPC state, but do so after a delay that could be accounted for by a required progression through a single, or multiple rounds, of cell division.

**Increased cone genesis in response to Notch inhibition is delayed relative to restricted RPC formation**

While multiple experiments support the conclusion that Notch inhibition leads to the formation of an increased restricted RPC population, we did not detect any increased cone genesis within a 2-day post-electroporation window (Fig. 2B and Fig. S3). To investigate the possibility that there is a delay in cone genesis that will occur from the induced cone/HC RPCs, we used a lineage tracing strategy to follow the fates of multipotent RPCs and assessed the cone fate within this population. Chick retinas were electroporated with a VSX2ECR4::PhiC31 lineage tracing plasmid, a GFP lineage tracing reporter, a ThrbCRM2::TdTomato reporter, and CAG::iRFP to mark all electroporated cells (Fig. 7A). Retinal samples were harvested after two to five days culture. The total number of ThrbCRM2 reporter-positive cones did not change at 2 or 3 days post-electroporation, however, the number of ThrbCRM2 reporter-positive cells within the VSX2ECR4 lineage-traced population showed a significant increase at 3 days, in line with the hypothesis that the effects of Notch inhibition originate from multipotent RPCs (Fig. 7B). Both the overall

ThrbCRM2 population and the population specifically lineage-traced from

multipotent RPCs showed a very significant enrichment at 4 and 5 days after

electroporation (Fig. 7C), which correlated in a delayed manner with the

changes of ThrbCRM1 restricted RPCs from 1 to 2 days (Fig. 6C and Fig.

2B). This supports our hypothesis that increased cone genesis has a delay of

2 days relative to the up-regulated ThrbCRM1 RPCs formed in response to

Notch inhibition and that this increased cone cell number is ultimately caused

by effects that originate in multipotent RPCs.

**Discussion**

While critical gene regulators of retinal cell type formation have been

identified in the last decades, the relationships of these factors within gene

regulatory networks and the cellular events in which they function are much

less clear. One of the primary reasons for this is that, unlike other

developmental systems, strict lineages of cell types originating from

multipotent RPCs have not been identified, leading to some hypotheses that

this system is driven primarily by stochastic choices (Gomes et al., 2011).

However, identification of restricted RPC states with limited cell fate potential

suggests that there are at least some deterministic processes involved and

these are additional, discrete cellular steps in the formation of specific cell

types during which genes could act. Here we attempted to integrate the well-

documented role for the Notch pathway in the repression of cone

photoreceptor formation and promotion of RPC proliferation, within a

framework for retinal development that includes restricted RPCs.

We initially hypothesized that, similar to other systems, the effect of

Notch inhibition on cone photoreceptors would be mediated through sister cell

fate decisions. This would be consistent with the fact that there is a restricted

RPC type that primarily forms cones and HCs and that several reports that

examined the Notch1 mouse mutant identified an increase in cones and one

also reported a decrease in HCs. However, though we also detected reduced

HCs when we interfered with Notch signaling, we did not detect a concomitant

increase in photoreceptors within the ThrbCRM1 restricted RPC daughter cell

population. This suggests that sister cell fate choice is not the primary site of

functional Notch signaling during cone photoreceptor formation. The most

prominent and consistent effects of Notch signaling we observed were on the

formation of the ThrbCRM1-active restricted RPC population and a

concomitant depletion of the multipotent RPC population, which suggests that

the primary role for Notch signaling in cone formation is at the step of

restricted RPC derivation from multipotent RPCs. Our interpretation of these

results is consistent with another recent developmental study that undertook a

comprehensive transcriptome analysis of DAPT-treated embryonic mouse

retinas (Kaufman et al., 2019). Here, it was reported that the earliest effects of

Notch inhibition through DAPT treatment include the loss of multipotent RPC

gene expression (observed as well by earlier studies of Notch1 and Rbpj), but

also the early increase of genes that have identified primary correlations with

restricted RPCs and with reduced to no expression in postmitotic cones

(Jadhav et al., 2006; Riesenberg et al., 2009; Yaron et al., 2006). These gene

expression changes include Olig2 (Hafler et al., 2012), Onecut1 (Emerson et

al., 2013), and Otx2, all of which have been previously shown to be expressed

in restricted RPCs that give rise to cones (Emerson et al., 2013; Hafler et al.,

2012). Additional genes such as Dll4 (Rocha et al., 2009), and Atoh7

(Buenaventura et al., 2018), are expressed in cycling cells and are enriched in

ThrbCRM1 cells as assessed by bulk RNA-Seq. All of these genes displayed

a temporal dynamic consistent with increased production of restricted RPCs in

this mouse model. In addition, several studies (Kaufman et al., 2019; Nelson

et al., 2007) have identified an upregulation of several members of the bHLH

family (NeuroD1, Ngn2, etc) which recent single cell analysis has associated

with neurogenic RPCs and not multipotent RPCs (Lu et al., 2020).

A role for Notch signaling in maintaining a self-renewing multipotent

RPC population is also consistent with its observed effects on proliferation.

Many studies (Jadhav et al., 2006; Kaufman et al., 2019; Yaron et al., 2006)

have identified a decrease in cycling cells when Notch signaling is inhibited.

We also detected a significant decrease within two days of electroporation,

but consistent with other analysis at early time points in the mouse, there was

only a partial loss of cycling cells. Given the identified shift from multipotent to

restricted RPCs identified here, we would predict that the majority of the

remaining cycling cells in mouse Notch loss-of-function paradigms represent

restricted RPCs and that the loss of multipotent RPCs is actually much larger

than was assumed to be the case.

The evidence presented in the current study, and interpreted from

previous ones, suggests a primary role in Notch signaling for mediating the

multipotent to restricted RPC transition, but there are critical questions that

remain to be addressed. One question is how cell division patterns of

multipotent RPCs change with inhibition of Notch signaling? Instead of a

multipotent RPC dividing asymmetrically to form another multipotent RPC and

a ThrbCRM1 RPC, does Notch inhibition cause symmetrical divisions to form

two ThrbCRM1 RPCs? Previous studies have also suggested that there may

exist an additional RPC state located between multipotent RPCs and

ThrbCRM1 RPCs (Schick et al., 2019; Suzuki et al., 2013). This RPC divides

to form an RGC and a ThrbCRM1 RPC on division. The formation of these

cells could be the first change elicited by Notch inhibition which would suggest

that RGCs should also increase. We do observe a number of cells in the RGC

layer, however, we have not detected an increase in RGCs using early RGC

markers such as Islet1 and Brn3. It is possible that the continued inhibition of

the Notch signaling pathway in these cells interferes with their differentiation.

This study focused on the formation of cone photoreceptors, but it will be

interesting to determine if Notch functions at other temporal points in

retinogenesis to control the formation of other restricted RPC types. Lastly, it

will be informative to determine the direct functional targets of Notch signaling

that drive RPC transitions. Identification of these pathways will have

implications for our understanding of visual system evolution and for the

development of strategies to control the proliferation dynamics and cell fate

choices of stem cells for therapeutic uses.

**Methods**

**Animals**

All experimental procedures involving animals were carried out in consultation

with the City College of New York, CUNY Institutional Animal Care protocols.

Fertilized chick eggs were acquired from Charles River, stored in a 16 °C

room for less than 10 days and incubated in a 38 °C humidified incubator for 5

days. All experiments that used animals were not sex-biased.

**DNA Plasmids**

The Hes1 reporter (CBFRE-EGFP) was obtained from Nicholas Gaiano

(Addgene plasmid #17705; http://n2t.net/addgene:17705;

RRID:Addgene_17705) and described in (Mizutani et al., 2007). The

Hes5::GFP reporter was described in (Vilas-Boas et al., 2011). 12XCSL-

EGFP reporter (12XCSL-d1EGFP) was obtained from Urban Lendahl

(Addgene plasmid # 47684; http://n2t.net/addgene:47684;

RRID:Addgene_47684) and described in (Hansson et al., 2006). CAG::βgal

was obtained from the Cepko lab (Cepko Lab, Harvard Medical School).

CAG::TdT was described in (Schick et al., 2019). OC1ECR22::GFP was

described in (Patoori et al., 2020 and manuscript in preparation).

VSX2ECR4::GFP reporter and CAG::iRFP were described in (Buenaventura

et al., 2018). ThrbCRM1::GFP and ThrbCRM2::GFP reporters were previously

reported (Emerson et al., 2013). The lineage trace driver ThrbCRM1::PhiC31

and its responder plasmid (CaNa::GFP) were described in (Schick et al.,

2019).

To make the CAG::MAML-DN plasmid (Addgene plasmid # 160923, this

study), an N-terminal MAML coding fragment (Weng et al., 2003) was

amplified from chicken cDNA and fused with a mouse Gapdh coding

sequence through overlap extension PCR. The PCR product was purified and

digested with AgeⅠ/NheⅠ before ligation into an AgeⅠ/NheⅠ digested

CAG::AU1 vector. ThrbCRM1::MAML-DN was made by replacing the CAG

promoter with the ThrbCRM1 element. The CAG::Rbpj-EnR ( Addgene

plasmid # 160924, this study) and CAG::EnR-Rbpj (Addgene plasmid #

160925, this study) were made by fusing mouse Rbpj (GenBank ID

BC051387.1 obtained from Transomics) with an engrailed repressor domain

(Vickers and Sharrocks, 2002) and amplification through overlap extension

PCR. The amplicon was digested by EcoRⅠ and ligated into an EcoRⅠ

digested CAG vector. All the plasmids described above were validated by

restriction digest and Sanger sequencing.

**Electroporation and Culture**

To deliver plasmid DNA into dissected retinas, *ex vivo* experiments were

carried out as described in (Emerson and Cepko, 2011) using a Nepagene

NEPA21 Type II Super Electroporator. DNA mixes for *ex vivo* electroporation

were made in 1X PBS with a final concentration of 0.1 µg/µl for the co-

electroporation control plasmids with CAG promoters and 0.16 µg/µl for

reporter and dominant-negative plasmids. Retinas were dissected at E5 and

cultured between 8 hours and 2 days after electroporation.

**Immunohistochemistry**

After harvest, retinas were fixed in 4% paraformaldehyde for 30 minutes,

washed 3 times in 1XPBS, and sunk in 30% sucrose/0.5XPBS overnight.

Retinas were frozen in OCT (Sakura Tissue-Tek, 4583). 20 µm vertical

sections were acquired using a Leica Cryostat and collected on 5-7 slides

(FisherBrand, 12-550-15).

Sections were blocked in 5% serum (Jackson ImmunoResearch, Donkey -

017-000121, Goat - 005-000-121) in 1XPBT (0.01% Tween-20 in 1XPBS) for

hour at room temperature and incubated in primary antibody at 4℃ overnight. The primary antibodies used were: chicken anti-GFP (ab13970, Abcam, 1:1000), rabbit anti-GFP (A-6455, Invitrogen, 1:1000), chicken anti-β-galactosidase (ab9361, Abcam, 1:1000), mouse IgG1 anti-β-galactosidase (40-1a-s, DSHB 1:20), rabbit anti-RFP (600-401-379, Rockland, 1:250), mouse IgG1 anti-Visinin (7G4-s, DSHB, 1: 500), mouse IgG1 anti-Lim1 (4F2-c, DSHB, 1:30), mouse anti-AP2 α (3B5, DSHB, 1:200), mouse IgG1 anti-Islet1 (39.3F7, DSHB, 1:50), mouse IgG1 anti-Brn3a (MAB1585, EMD Millipore, 1:800), mouse anti-Pan-Brn3 (Sc-390781, Santa Cruz, 1:100), sheep anti-Vsx2 (x1180p, Ex Alpha, 1:200), goat anti-Otx2 (AF1979, R&D, 1:500), rabbit anti-Otx2 (AB21990, Abcam, 1:500) and rabbit anti-Olig2 (4B9610-AF488, Millipore, 1:500). Sections were washed 3 times with 1XPBT, blocked at room temperature for 30 min and incubated in secondary antibody at 4℃ overnight. The secondary antibodies used were from Jackson Immunoresearch and appropriate for multiple labeling. Alexa 488- and 647-conjugated secondary antibodies (resuspended in 50% glycerol) were used at 1:400 and cy3-conjugated antibodies were used at 1:250 as described (Schick et al., 2019). 4'6-diamidino-2-phenylindole (DAPI) was added at 1 µg/ml after 3X wash with 1XPBT. Sections were mounted using Fluoromount-G (Southern Biotech, 0100-01) and 34x60 mm coverslips (VWR, 48393-106).

**EdU Labeling**

Before fixation, retinas were incubated at 37℃ for 1 hour with 50 µM EdU in culture media and harvested as described above. Sections were permeabilized for 15 minutes in 0.1% Triton in 1X PBS before incubating with the EdU Reaction Cocktail for 30 minutes in the dark. Sections were washed 3 times with 1X PBT before blocking and antibody staining. EdU was detected using a Click-iT EdU Alexa Fluor 647 imaging kit (Invitrogen, C10340).

**Confocal Microscopy and Cell Counting**

Confocal images were acquired using a Zeiss LSM710 inverted confocal

microscope with an EC Plan-Neofluar 40x/1.30 Oil DIC M27 immersion

objective, 488nm laser, 561 nm laser, 633 nm laser, 405 nm laser, and ZEN

Black 2015 21 SP2 software resolution of 1024 x 1024, acquisition speed of 6.

Fiji (Schindelin et al., 2012) was used to analyze images and convert into

picture format. Cell counting was performed using the Cell Counter plugin-in

for ImageJ developed by Kurt De Vos. Cell counts used at least 3 retinas for

each condition, 3 technical replicates per retina. Figures were assembled

using Affinity Designer vector editor (Serif [Europe] Ltd). Brightness and

contrast were adjusted uniformly in all the figures. All retinal sections are

oriented with the developing photoreceptor side at the top of the image. All

images are presented as maximum projections, with the exception of Olig2-

related images in Figure 3A, which are single z-planes.

**Retinal Cells Dissociation and Flow Cytometry**

After ex vivo culture, the remaining retinal pigment epithelium (RPE) and

condensed vitreal materials were removed in HBSS (GIBCO, 14170112) and

dissociated following a papain-based (Worthington, L5003126) protocol

(Jean-Charles et al., 2018). Dissociated retinal cells were fixed using 4%

paraformaldehyde for 15 minutes and washed 3 times in 1X PBS. Samples for

quantification were filtered into 4ml tubes (BD Falcon, 352054) through 40um

cell strainers (Biologix, 15-1040). Dissociated cells were analyzed with a BD

LSR II flow cytometer using the 488nm, 561nm, and 633nm lasers. All flow

cytometry data was analyzed using FlowJo Version 10.4.2.

**Statistical Analysis**

Graphs and statistical tests were conducted using Microsoft Office Excel and

GraphPad Prism 8. The error bars represent S.E.M. ANOVA with a post hoc

Dunn test was used in Fig. 4B and Fig. S2B. Mann-Whitney test was used in

Fig. 4D, Fig. 5B, Fig 6B, Fig. S4B and Fig. S7B. Student's t-test was used in

the rest of qualifications. All the tests were performed with independent samples after the Shapiro-Wilk normality test using GraphPad Prism8, and the results were significant as noted. Quantifications for flow cytometry were based on 4 biological replicates per condition and at least 2 technical replicates. Quantifications for cell counting were based on 3 biological replicates per condition and at least 3 technical replicates.

**Acknowledgements**

Funding for this project was provided by NSF CAREER 1453044 (to M.E.) and supported by NIMHD 3G12MD007603-30S2 (to CCNY). Jeffrey Walker and Jorge Morales provided technical support with flow cytometry and confocal microcopy experiments. We thank Diego Buenaventura for providing the VSX2ECR4::PhiC31 plasmid and for guidance with flow cytometry analysis. We thank Miruna Ghinia-Tegla for advice on cell counting and Estie Schick for comments on the manuscript. Shirley Chan, Denice Moran and Boguslava Dimitrova provided technical assistance. We thank all the members of the Emerson lab for support throughout the project.

**Author Contributions**

XC and ME designed the experiments and analyzed the data. XC and ME conducted the experiments. XC and ME wrote the manuscript.

**Competing Interests**

The author(s) declare no competing interests.

## References

Artavanis-Tsakonas, S., Delidakis, C., and Fehon, R.G. (1991). The Notch locus and the cell biology of neuroblast segregation. Annu. Rev. Cell Biol. 7, 427–452.

Bray, S. (1998). Notch signalling in Drosophila: three ways to use a pathway. Semin. Cell Dev. Biol. 9, 591–597.

Bray, S.J. (2006). Notch signalling: a simple pathway becomes complex. Nat. Rev. Mol. Cell Biol. 7, 678–689.

Buenaventura, D.F., Ghinia-Tegla, M.G., and Emerson, M.M. (2018). Fate-restricted retinal progenitor cells adopt a molecular profile and spatial position distinct from multipotent progenitor cells. Dev. Biol. 443, 35–49.

Buenaventura, D.F., Corseri, A., and Emerson, M.M. (2019). Identification of Genes With Enriched Expression in Early Developing Mouse Cone Photoreceptors. Invest. Ophthalmol. Vis. Sci. 60, 2787–2799.

Cepko, C. (2014). Intrinsically different retinal progenitor cells produce specific types of progeny. Nat. Rev. Neurosci. 15, 615–627.

Emerson, M.M., and Cepko, C.L. (2011). Identification of a retina-specific Otx2 enhancer element active in immature developing photoreceptors. Dev. Biol. 360, 241–255.

Emerson, M.M., Surzenko, N., Goetz, J.J., Trimarchi, J., and Cepko, C.L. (2013). Otx2 and Onecut1 promote the fates of cone photoreceptors and horizontal cells and repress rod photoreceptors. Dev. Cell 26, 59–72.

Fischer, A.J., Stanke, J.J., Aloisio, G., Hoy, H., and Stell, W.K. (2007). Heterogeneity of horizontal cells in the chicken retina. J. Comp. Neurol. 500, 1154–1171.

Ghinia Tegla, M.G., Buenaventura, D.F., Kim, D.Y., Thakurdin, C., Gonzalez, K.C., and Emerson, M.M. OTX2 represses sister cell fate choices in the developing retina to promote photoreceptor specification. ELife 9.

Gomes, F.L.A.F., Zhang, G., Carbonell, F., Correa, J.A., Harris, W.A., Simons, B.D., and Cayouette, M. (2011). Reconstruction of rat retinal progenitor cell lineages in vitro reveals a surprising degree of stochasticity in cell fate decisions. Dev. Camb. Engl. 138, 227–235.

Hafler, B.P., Surzenko, N., Beier, K.T., Punzo, C., Trimarchi, J.M., Kong, J.H., and Cepko, C.L. (2012). Transcription factor Olig2 defines subpopulations of retinal progenitor cells biased toward specific cell fates. Proc. Natl. Acad. Sci. 109, 7882–7887.

Hansson, E.M., Teixeira, A.I., Gustafsson, M.V., Dohda, T., Chapman, G., Meletis, K., Muhr, J., and Lendahl, U. (2006). Recording Notch signaling in real time. Dev. Neurosci. *28*, 118–127.

Jadhav, A.P., Mason, H.A., and Cepko, C.L. (2006). Notch 1 inhibits photoreceptor production in the developing mammalian retina. Dev. Camb. Engl. *133*, 913–923.

Jean-Charles, N., Buenaventura, D.F., and Emerson, M.M. (2018). Identification and characterization of early photoreceptor cis-regulatory elements and their relation to Onecut1. Neural Develop. *13*, 26.

Johnson, J.E., and MacDonald, R.J. (2011). Chapter three - Notch-Independent Functions of CSL. In Current Topics in Developmental Biology, C. Birchmeier, ed. (Academic Press), pp. 55–74.

Kaufman, M.L., Park, K.U., Goodson, N.B., Chew, S., Bersie, S., Jones, K.L., Lamba, D.A., and Brzezinski, J.A. (2019). Transcriptional profiling of murine retinas undergoing semi-synchronous cone photoreceptor differentiation. Dev. Biol. *453*, 155–167.

Kopan, R., and Ilagan, M.X.G. (2009). The canonical Notch signaling pathway: unfolding the activation mechanism. Cell *137*, 216–233.

Lai, E.C. (2004). Notch signaling: control of cell communication and cell fate. Dev. Camb. Engl. *131*, 965–973.

Liu, W., Khare, S.L., Liang, X., Peters, M.A., Liu, X., Cepko, C.L., and Xiang, M. (2000). All Brn3 genes can promote retinal ganglion cell differentiation in the chick. Dev. Camb. Engl. *127*, 3237–3247.

Lu, Y., Shiau, F., Yi, W., Lu, S., Wu, Q., Pearson, J.D., Kallman, A., Zhong, S., Hoang, T., Zuo, Z., et al. (2020). Single-Cell Analysis of Human Retina Identifies Evolutionarily Conserved and Species-Specific Mechanisms Controlling Development. Dev. Cell *53*, 473-491.e9.

Mizeracka, K., DeMaso, C.R., and Cepko, C.L. (2013). Notch1 is required in newly postmitotic cells to inhibit the rod photoreceptor fate. Dev. Camb. Engl. *140*, 3188–3197.

Mizutani, K., Yoon, K., Dang, L., Tokunaga, A., and Gaiano, N. (2007). Differential Notch signalling distinguishes neural stem cells from intermediate progenitors. Nature *449*, 351–355.

Mumm, J.S., and Kopan, R. (2000). Notch signaling: from the outside in. Dev. Biol. *228*, 151–165.

Nelson, B.R., Hartman, B.H., Georgi, S.A., Lan, M.S., and Reh, T.A. (2007).
Transient inactivation of Notch signaling synchronizes differentiation of neural
progenitor cells. Dev. Biol. *304*, 479–498.

Patoori, S., Jean-Charles, N., Gopal, A., Sulaiman, S., Gopal, S., Wang, B.,
Souferi, B., and Emerson, M.M. (2020). Cis-regulatory analysis of Onecut1
expression in fate-restricted retinal progenitor cells. Neural Develop. *15*, 5.

Riesenberg, A.N., Liu, Z., Kopan, R., and Brown, N.L. (2009). Rbpj cell
autonomous regulation of retinal ganglion cell and cone photoreceptor fates in
the mouse retina. J. Neurosci. Off. J. Soc. Neurosci. *29*, 12865–12877.

Riesenberg, A.N., Conley, K.W., Le, T.T., and Brown, N.L. (2018). Separate and
coincident expression of Hes1 and Hes5 in the developing mouse eye. Dev.
Dyn. Off. Publ. Am. Assoc. Anat. *247*, 212–221.

Rocha, S.F., Lopes, S.S., Gossler, A., and Henrique, D. (2009). Dll1 and Dll4
function sequentially in the retina and pV2 domain of the spinal cord to regulate
neurogenesis and create cell diversity. Dev. Biol. *328*, 54–65.

Schick, E., McCaffery, S.D., Keblish, E.E., Thakurdin, C., and Emerson, M.M.
(2019). Lineage tracing analysis of cone photoreceptor associated cis-
regulatory elements in the developing chicken retina. Sci. Rep. *9*, 1–14.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch,
T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an
open-source platform for biological-image analysis. Nat. Methods *9*, 676–682.

Suzuki, S.C., Bleckert, A., Williams, P.R., Takechi, M., Kawamura, S., and Wong,
R.O.L. (2013). Cone photoreceptor types in zebrafish are generated by
symmetric terminal divisions of dedicated precursors. Proc. Natl. Acad. Sci. U.
S. A. *110*, 15109–15114.

Tan, G.C., Mazzoni, E.O., and Wichterle, H. (2016). Iterative Role of Notch
Signaling in Spinal Motor Neuron Diversification. Cell Rep. *16*, 907–916.

Vickers, E.R., and Sharrocks, A.D. (2002). The use of inducible engrailed fusion
proteins to study the cellular functions of eukaryotic transcription factors.
Methods San Diego Calif *26*, 270–280.

Vilas-Boas, F., Fior, R., Swedlow, J.R., Storey, K.G., and Henrique, D. (2011).
A novel reporter of notch signalling indicates regulated and random Notch
activation during vertebrate neurogenesis. BMC Biol. *9*, 58.

Weng, A.P., Nam, Y., Wolfe, M.S., Pear, W.S., Griffin, J.D., Blacklow, S.C., and
Aster, J.C. (2003). Growth suppression of pre-T acute lymphoblastic leukemia
cells by inhibition of notch signaling. Mol. Cell. Biol. *23*, 655–664.

Wu, L., Aster, J.C., Blacklow, S.C., Lake, R., Artavanis-Tsakonas, S., and Griffin,
J.D. (2000). MAML1, a human homologue of Drosophila mastermind, is a
transcriptional co-activator for NOTCH receptors. Nat. Genet. *26*, 484–489.

Yaron, O., Farhy, C., Marquardt, T., Applebury, M., and Ashery-Padan, R. (2006).
Notch1 functions to suppress cone-photoreceptor fate specification in the
developing mouse retina. Dev. Camb. Engl. *133*, 1367–1378.

**Figure Legends**

**Fig. 1.** CAG::MAML-DN inhibits the expression of the Hes5 Notch reporter

**A.** Schematic representation of the electroporation paradigm with plasmids introduced ex vivo into E5 chick retinas, cultured for two days, dissociated into single cells and analyzed by flow cytometry or confocal micrsocopy. A schematic of the MAML dominant-negative construct is shown. **B.** Confocal images of vertically sectioned E5 chick retinas co-electroporated with CAG::Nucßgal, Hes5::GFP Notch reporter, and with or without CAG::MAML-DN and then cultured for two days. The scale bar in the bottom right picture denotes 40 µm and applies to all images. **C.** Flow cytometry dot plots of dissociated chick retinal cells electroporated *ex vivo* at E5 with a co-electroporation control (CAG::iRFP), Hes5::GFP, and with or without CAG::MAML-DN. The bar graph on the right shows the percentage of Hes5::GFP reporter-positive cells within the electroporated population. Each point represents one biological replicate. The Shapiro-Wilk normality test was used to confirm the normal distribution. ** signifies p<0.01 with a student's t-test.

**Fig. 2.** CAG::MAML-DN induced Notch inhibition promotes ThrbCRM1 restricted RPC formation

**A.** Confocal images of vertically sectioned E5 chick retinas co-electroporated with CAG::Nucßgal, either ThrbCRM1 or VSX2ECR4 GFP reporters, and with or without CAG::MAML-DN and then cultured for two days. The scale bar shown in the bottom right picture denotes 40 µm and applies to all images. **B.** Flow cytometry quantification of the percentage of ThrbCRM1, ThrbCRM2, VSX2ECR4 and OC1ECR22 reporter-positive cells within the CAG::iRFP electroporated population in E5 retinas cultured for two days. Green bars represent control conditions and yellow bars inclusion of the CAG::MAML-DN construct. Each point represents one biological replicate. The Shapiro-Wilk

normality test was used to confirm the normal distribution. * signifies p<0.05,

** signifies p<0.01, *** signifies p<0.001 with a student's t-test.

**Fig. 3.** Effects of CAG::MAML-DN on endogenous retinal markers

**A.** Confocal images of vertically sectioned E5 chick retinas co-electroporated

with CAG::Nucßgal, and with or without CAG::MAML-DN and then cultured for

two days. Sections were immunostained with Otx2 or Olig2 markers

(magenta), CAG::Nucßgal (yellow), and nuclei visualized with DAPI. The

scale bar in the bottom right picture denotes 40 μm and applies to all images.

**B.** Flow cytometry quantification of the percentage of Visinin, Otx2, Lim1 and

AP2 α -positive cells within the electroporated cells. Dissociated chick retinal

cells electroporated *ex vivo* at E5 with the co-electroporation control

CAG::TdTomato, under control (green bars) or CAG::MAML-DN (yellow bars)

conditions. The retinas were cultured for two days. Each point represents one

biological replicate. The Shapiro-Wilk normality test was used to confirm the

normal distribution. ** signifies p<0.01, *** signifies p<0.001 with a student's t-

test.   **C.** Quantification of the percentage of Olig2-positive cells within the

electroporated cell population from cell counting. Sectioned chick retinas

electroporated *ex vivo* at E5 with the co-electroporation control

CAG::Nucßgal. The retinas were cultured for two days. Each point represents

one biological replicate. The Shapiro-Wilk normality test was used to confirm

the normal distribution. *** signifies p<0.001 with a student's t-test.

**Fig. 4.** Targeted expression of MAML-DN to cone/HC restricted RPCs does

not alter cone formation

**A.** Schematic representation of PhiC31 recombinase-based lineage tracing

strategy (Schick et al., 2019).   **B.** Flow cytometry quantification of the

percentage of ThrbCRM1 reporter-positive cells within all the electroporated

cells. Dissociated chick retinal cells electroporated *ex vivo* at E5 with the

CAG::iRFP co-electroporation control, and with or without CAG::MAML-DN or

ThrbCRM1::MAML-DN and cultured for two days. Each point represents one

biological replicate. The Shapiro-Wilk normality test was used to confirm the

normal distribution. ANOVA with a post hoc Dunn test was used to test

significance. ** signifies $p<0.01$, ns signifies $p>0.05$. **C.** Confocal images of

vertically sectioned E5 chick retinas co-electroporated with CAG::Nucßgal

using ThrbCRM1::MAML-DN lineage trace with a PhiC31 responder plasmid

shown in A. The retinas were cultured for two days post-electroporation.

Sections were immunostained with Visinin (magenta), CAG::Nucßgal

(orange), and nuclei visualized with DAPI. The scale bar in the bottom right

picture denotes 40 μm and applies to all images. **D.** Flow cytometry

quantification of the percentage of Otx2, Lim1, AP2 α , OC1ECR22 and

ThrbCRM2 reporter-positive cells within the ThrbCRM1 lineage traced cell

population. Dissociated chick retinal cells electroporated *ex vivo* at E5 with the

co-electroporation control CAG::TdTomato, ThrbCRM1::PhiC31 and its

responder plasmid, and with or without CAG::MAML-DN. The retinas were

cultured for two days post-electroporation. Each point represents one

biological replicate. The Shapiro-Wilk normality test was used to confirm the

normal distribution. Mann-Whitney test was used to test significance in

ThrbCRM2 quantification. A student's t-test was used to test significance in

the rest of the quantifications. * signifies $p<0.05$, ** signifies $p<0.01$.

**Fig. 5.** CAG::MAML-DN induced Notch inhibition leads to a shift from

multipotent to cone/HC restricted RPCs. **A-E.** Retinas were cultured for two

days post-electroporation. **A.** Flow cytometry quantification of the percentage

of EdU-positive cells within the electroporated cell population. Dissociated

chick retinal cells electroporated *ex vivo* at E5 with the co-electroporation

control CAG::TdTomato with or without CAG::MAML-DN. Retinas were pulsed

with EdU for 1 hour before harvest. Each point represents one biological

replicate. The Shapiro-Wilk normality test was used to confirm the normal

distribution. A student's t-test was used to test significance. **B.** Flow cytometry

quantification of EdU+ cells positive for VSX2ECR4 or ThrbCRM1 reporters. Dissociated chick retinal cells electroporated *ex vivo* at E5 with VSX2ECR4::GFP and ThrbCRM1::TdTomato. Retinas were pulsed with EdU for 1 hour before harvest. Each point represents one biological replicate. The Shapiro-Wilk normality test was used to confirm the normal distribution. Mann-Whitney test was used to test significance in VSX2ECR4/EdU+ quantification. A student's t-test was used to test significance in ThrbCRM1/EdU+ quantification. ** signifies $p<0.01$. **C.** Flow cytometry quantification of ThrbCRM1 reporter-positive cells that also express VSX2ECR4 reporter and incorporate EdU from dissociated chick retinal cells electroporated *ex vivo* at E5 with VSX2ECR4::GFP and ThrbCRM1::TdTomato. Retinas were pulsed with EdU for 1 hour prior to harvest. Each point represents one biological replicate. The Shapiro-Wilk normality test was used to confirm the normal distribution. *** signifies $p<0.001$ with a student's t-test. **D.** Confocal images of vertically sectioned E5 chick retinas co-electroporated with CAG::Nucßgal, ThrbCRM1::GFP or VSX2ECR4::GFP, and with or without CAG::MAML-DN. Retinas were pulsed with EdU for 1 hour before harvest. The scale bar in the bottom right picture denotes 40 μm and applies to all images. **E.** Flow cytometry quantification of the percentage of VSX2ECR4/ThrbCRM1 double reporter-positive cells within the VSX2ECR4 lineage traced population. Dissociated chick retinal cells electroporated *ex vivo* at E5 with the co-electroporation control CAG::iRFP, VSX2ECR4::PhiC31 and its responder plasmid, and with or without CAG::MAML-DN. Each point represents one biological replicate. The Shapiro-Wilk normality test was used to confirm the normal distribution. ** signifies $p<0.01$ with a student's t-test.

**Fig. 6.** The short-term effects of Notch inhibition on RPC reporters

**A.** Confocal images of vertically sectioned E5 chick retinas co-electroporated with CAG::Nucßgal, ThrbCRM1::GFP or VSX2ECR4::GFP, and with or without CAG::MAML-DN. The retinas were cultured for 8 hours post-

electroporation. The scale bar in the bottom right picture denotes 40 µm and applies to all images.   **B.** Flow cytometry quantification of the percentage of VSX2ECR4, ThrbCRM1 and VSX2ECR4/ThrbCRM1 reporter-positive cells within all the electroporated cells. Dissociated chick retinal cells electroporated *ex vivo* at E5 with the co-electroporation control CAG::iRFP with or without CAG::MAML-DN. Retinas were cultured for 8 hours post-electroporation. Each point represents one biological replicate. The Shapiro-Wilk normality test was used to confirm the normal distribution. Mann-Whitney test was used to test significance in VSX2ECR4 quantification. A student's t-test was used to test significance in ThrbCRM1 and double-positive quantifications. **C.** Flow cytometry quantification of the percentage of VSX2ECR4, ThrbCRM1 and VSX2ECR4/ThrbCRM1 reporter-positive cells within all the electroporated cells. Dissociated chick retinal cells electroporated with CAG::iRFP *ex vivo* at E5 and cultured for 20 hours. Each point represents one biological replicate. The Shapiro-Wilk normality test was used to confirm the normal distribution. *** signifies $p<0.001$ with a student's t-test. **D.** Quantification of the percentage of VSX2, Otx2 and double marker-positive cells within the electroporated cell population from cell counting. Sectioned chick retinas electroporated *ex vivo* at E5 with the co-electroporation control CAG::Nucßgal with or without CAG::MAML-DN. The retinas were cultured for 20 hours and immunostained with VSX2 or Otx2 antibodies. Each point represents one biological replicate. The Shapiro-Wilk normality test was used to confirm the normal distribution. * signifies $p<0.05$, *** signifies $p<0.001$ with a student's t-test.

**Fig. 7.** Notch inhibition leads to a delayed increase in cones relative to cone/HC RPCs

**A.** Schematic of experimental set-up. Retinas were electroporated with VSX2ECR4::PhiC31 and CAG::stopGFP responder plasmid to label multipotent RPCs (mpRPC)s and their progeny. In addition, ThrbCRM2::TdT

was co-electroporated to identify a subset of cones. CAG::MAML-DN was included in experimental conditions and omitted in control conditions. CAG::iRFP was included as a co-electroporation control (not shown). Retinas were assessed 2 to 5 days later by flow cytometry to measure GFP, TdT, and iRFP activity. Non-electroporated cells are shown in blue, lineage-traced cells in green, non-lineage traced ThrbCRM2+ cells in magenta and lineage traced ThrbCRM2+ cells in orange. **B.** Flow cytometry quantification of the percentage of ThrbCRM2 reporter-positive cells within all the electroporated cells using VSX2ECR4 lineage trace strategy with the PhiC31 responder plasmid shown in 4A. Dissociated chick retinal cells electroporated *ex vivo* at E5 with the co-electroporation control CAG::iRFP, VSX2ECR4::PhiC31 and its responder plasmid, ThrbCRM2::TdTomato, and with or without CAG::MAML-DN. Retinas were cultured for 2 or 3 days post-electroporation. A student's t-test was used to test significance. Each point represents one biological replicate. The Shapiro-Wilk normality test was used to confirm the normal distribution. * signifies $p<0.05$, ** signifies $p<0.01$, *** signifies $p<0.001$ with a student's t-test. **C.** Same as A but with 4 or 5 days culture.

**Fig. 8.** Model of Notch inhibition effects on cone/HC regulatory network Schematic of the general relationship between multipotent RPCs (mpRPC, beige), cone/HC RPCs (green), cones (blue), and HCs (purple). Grey shading denotes cell-type limits of Notch inhibition. CAG::MAML-DN is broadly active in all cells (denoted by grey shading) and leads to reduction in mpRPCs, increase in cone/HC RPCs and cones. Bottom graphs represent relative numbers of cell type populations under control and Notch inhibition conditions. Dotted lines depict the control levels of cones and HCs.

**Fig. S1.** CAG::MAML-DN effects on Hes1 and 12XCSL-EGFP reporters.
**A.** Confocal images of vertically sectioned E5 chick retinas co-electroporated with CAG::Nucßgal, Hes1::GFP or 12XCSL-EGFP, and with or without

CAG::MAML-DN and cultured for two days. The scale bar in the bottom right

picture denotes 40 µm and applies to all images. **B.** Bar graphs of flow

cytometry analysis of dissociated chick retinal cells electroporated *ex vivo* at

E5 with the CAG::iRFP co-electroporation control, Hes1::GFP reporter (left

graph) or 12XCSL-EGFP (right graph), and with or without CAG::MAML-DN

and cultured for two days. The number of EGFP reporter-positive cells within

the electroporated population is plotted. Each point represents one biological

replicate. The Shapiro-Wilk normality test was used to confirm the normal

distribution. * signifies $p < 0.05$ with a student's t-test.

**Fig. S2.** Rbpj-EnR has similar effects as CAG::MAML-DN on Hes5 Notch

reporter and ThrbCRM1-marked RPCs

**A.** Confocal images of vertically sectioned E5 chick retinas co-electroporated

with CAG::Nucßgal, Hes5::GFP Notch reporter, and with or without Rbpj-EnR

and cultured for two days before Visinin immunostaining. The scale bar in the

bottom right picture denotes 40 µm and applies to all images. **B.** Flow

cytometry quantification of the percentage of Hes5 and ThrbCRM1-positive

cells within all the electroporated cells. Dissociated chick retinal cells

electroporated *ex vivo* at E5 with the CAG::iRFP co-electroporation control,

Hes5 or ThrbCRM1 reporters, and with or without EnR-Rbpj or Rbpj-EnR and

cultured for two days. Each point represents one biological replicate. The

Shapiro-Wilk normality test was used to confirm the normal distribution.

ANOVA with a post hoc Dunn test was used to test significance. *** signifies

$p < 0.001$.

**Fig. S3.** CAG::MAML-DN does not promote cone formation at two days post-

electroporation

Confocal images of vertically sectioned E5 chick retinas co-electroporated

with CAG::Nucßgal, ThrbCRM2::GFP, OC1ECR22::GFP or ThrbCRM1::GFP ,

and with or without CAG::MAML-DN and cultured for two days. LHX4 antibody

was used to label cones. The scale bar in the bottom right picture denotes 40 µm and applies to all images.

**Fig. S4.** CAG::MAML-DN does not shift the number of H2, H3 and H4 HCs and RGCs. **A.** Confocal images of vertically sectioned E5 chick retinas co-electroporated with CAG::Nucßgal, and with or without CAG::MAML-DN and cultured for two days. Sections were immunostained with Islet1 and Pan-Brn3 cell specific markers (magenta), CAG::Nucßgal (orange), and nuclei visualized with DAPI. The scale bar in the bottom right panel denotes 40 µm and applies to all images. **B.** Flow cytometry quantification of the percentage of Islet1, Brn3a and Pan-Brn3-positive cells within all electroporated cells. Dissociated chick retinal cells electroporated *ex vivo* at E5 with the CAG::TdTomato co-electroporation control and cultured for two days. Each point represents one biological replicate. The Shapiro-Wilk normality test was used to confirm the normal distribution. A student's t-test was used to test significance in Islet1 and Brn3a quantifications. Mann-Whitney test was used to test significance in Pan-Brn3 quantification.

**Fig. S5.** Rbpj-EnR promotes ThrbCRM1 RPCs and represses HCs
Flow cytometry quantification of the percentage of VSX2ECR4, ThrbCRM1, VSX2ECR4/ThrbCRM1 double reporter-positive, Visinin, ThrbCRM2, OC-ECR22, Lim1 and AP2 $\alpha$ -positive cells within all the electroporated cells. Dissociated chick retinal cells electroporated *ex vivo* at E5 with the CAG::iRFP or CAG::TdTomato co-electroporation control, and with or without Rbpj-EnR and cultured for two days. Each point represents one biological replicate. The Shapiro-Wilk normality test was used to confirm the normal distribution. ** signifies p<0.01, *** signifies p<0.001 with a student's t-test.

**Fig. S6.** ThrbCRM1::MAML-DN does not have effects on H2, H3 and H4 HCs and RGCs within ThrbCRM1 lineage traced population

**A.** Confocal images of vertically sectioned E5 chick retinas co-electroporated with CAG::Nucßgal, ThrbCRM1::PhiC31 and its responder plasmid, and with or without ThrbCRM1::MAML-DN. The retinas were cultured for two days post-electroporation. The sections were immunostained with Otx2, Lim1 and AP2$\alpha$ markers (magenta), CAG::Nucßgal (orange), and nuclei visualized with DAPI. The scale bar in the bottom right picture denotes 40 µm and applies to all images. **B.** Quantification of the percentage of Islet1 and Brn3 marker-positive cells within the ThrbCRM1 lineage traced cell population from cell counting. Sectioned chick retinas electroporated *ex vivo* at E5 with the co-electroporation control CAG::Nucßgal, ThrbCRM1::PhiC31 and its responder plasmid, and with or without ThrbCRM1::MAML-DN. The retinas were cultured for two days after electroporation. Each point represents one biological replicate. The Shapiro-Wilk normality test was used to confirm the normal distribution. A student's t-test was used to test significance.

**Fig. S7.** Rbpj-EnR induced Notch inhibition has similar effects as CAG::MAML-DN on cell proliferation

**A.** Flow cytometry quantification of the percentage of EdU-positive cells within all the electroporated cells. Dissociated chick retinal cells electroporated *ex vivo* at E5 with the co-electroporation control CAG::TdTomato with or without Rbpj-EnR. The retinas were cultured for two days after electroporation and pulsed with EdU for 1 hour before harvest. Each point represents one biological replicate. The Shapiro-Wilk normality test was used to confirm the normal distribution. A student's t-test was used to test significance. **B.** Flow cytometry quantification of the percentage of VSX2ECR4, ThrbCRM1 or VSX2ECR4/ThrbCRM1-positive cells within the electroporated cell population. Dissociated chick retinal cells electroporated *ex vivo* at E5 with the co-electroporation control CAG::iRFP, VSX2ECR4::GFP or ThrbCRM1::GFP, and with or without Rbpj-EnR. The retinas were cultured for 8 hours post-electroporation. Each point represents one biological replicate. The Shapiro-

Wilk normality test was used to confirm the normal distribution. A student's t-

test was used to test significance in ThrbCRM1 quantification. Mann-Whitney

test was used to test significance in VSX2ECR4 and double reporter-positive

quantifications. * signifies p<0.05.

**Fig. S8.** CAG::MAML-DN induced Notch inhibition led to a transition from

VSX2+/EdU+ cells to Otx2+/EdU+ cells

Confocal images of vertically sectioned E5 chick retinas co-electroporated

with CAG::Nucßgal and with or without CAG::MAML-DN. The retinas were

cultured for two days after electroporation and pulsed with EdU for 1 hour

before harvest. The sections were immunostained with VSX2 and Otx2

makers (magenta), CAG::Nucßgal (orange), and nuclei visualized with DAPI.

The scale bar in the bottom right picture denotes 40 µm and applies to all

images.

# Fig. 1

**A**

electroporation

E5 chick retina

2X AU1 Tag    N-term cMAML1    GAPDH

8 hours to 5 days culture

Confocal Microscopy

Dissociation

Flow cytometry

X%

Y%

**B**

|  | DAPI | CAG::Nucβgal | Hes5::GFP | Merge |
|---|---|---|---|---|
| Control | | | | |
| CAG::MAML-DN | | | | |

**C**

Hes5 reporter labeled cells (GFP+)

Control

Q1 0.020    Q2 0.77
Q4 86.9    Q3 12.3

CAG::MAML-DN

Q1 0.016    Q2 0.13
Q4 85.8    Q3 14.1

Electroporated cells (iRFP+)

% Hes5 reporter-positive cells

∗∗

Control    CAG::MAML-DN

# Fig. 2

# Fig. 3

# Fig. 4

# Fig. 5

# Fig. 6

# Fig. 7

## A

## B

## C

# Fig.8

| Wildtype | CAG::MAML-DN | ThrbCRM1::MAML-DN |
|---|---|---|

Cones:  —  ↑  —