Engineering Radioprotective Human Cells Using the Tardigrade Damage Suppressor Protein, DSUP

Authors: Craig Westover[1], Deena Najjar[1], Cem Meydan[1], Kirill Grigorev[1], Mike T. Veling[3,4], Sonia Iosim[1], Rafael Colon[1], Sherry Yang[1], Uriel Restrepo[1] Christopher Chin[1], Daniel Butler[1], Chris Moszary[1], Savlatjaton Rahmatulloev[1], Ebrahim Afshinnekoo[1,2,5], Roger L Chang[3,4], Pamela A Silver[3,4], Christopher E. Mason[1,2,5,6]*

[1]Department of Physiology and Biophysics, Weill Cornell Medicine, New York, NY, USA
[2]The HRH Prince Alwaleed Bin Talal Bin Abdulaziz Alsaud Institute for Computational Biomedicine, Weill Cornell Medicine, New York, NY, USA
[3]Department of Systems Biology, Blavatnik Institute at Harvard Medical School, Boston, MA 02115, USA.
[4]Wyss Institute for Biologically Inspired Engineering, Harvard University, Boston, MA 02115, USA.
[5]The WorldQuant Initiative for Quantitative Prediction, Weill Cornell Medicine, New York, NY, USA
[6]The Feil Family Brain and Mind Research Institute, Weill Cornell Medicine, New York, NY, USA

*Corresponding Author
Christopher E. Mason
Weill Cornell Medicine
1305 York Ave., Y13-05
New York, NY 10021
Tel: 203-668-1448
E-mail: chm2042@med.cornell.edu

Abstract

Spaceflight has been documented to produce a number of detrimental effects to physiology and genomic stability, partly a result of Galactic Cosmic Radiation (GCR). In recent years, extensive research into extremotolerant organisms has begun to reveal how they survive harsh conditions, such as ionizing radiation. One such organism is the tardigrade (*Ramazzottius varieornatus*) which can survive up to 5kGy of ionizing radiation and also survive the vacuum of space. In addition to their extensive network of DNA damage and response mechanisms, the tardigrade also possesses a unique damage suppressor protein (Dsup) that co-localizes with chromatin in both tardigrade and transduced human cells and protects against damage from reactive oxygen species via ionizing radiation. While Dsup has been shown to confer human cells with radioresistance; much of the mechanism of how it does this in the context of human cells remains to be elucidated. In addition, there is no knowledge yet of how introduction of Dsup into human cells can perturb cellular networks and if there are any systemic risks associated. Here, we created a stable HEK293 cell line expressing Dsup via lentiviral transduction and confirmed its presence and its integration site. We show that Dsup confers human cells with a reduction of apoptotic signals. Through measuring these biomarkers of DNA damage in response to

irradiation longitudinally along with gene expression analysis, we were able to demonstrate a
potential role for Dsup as DNA damage response and repair enhancer much in the same way its
human homologous counterpart HMGN1 functions. Our methods and tools provide evidence that
the effects of the Dsup protein can be potentially utilized to mitigate such damage during
spaceflight.

Introduction
As more public and private enterprises plan to send humans on long-term spaceflight missions,
the challenges associated with adapting to the harsh conditions of space become more
pronounced. One of the major risks to humans during spaceflight is the exposure to Galactic
Cosmic Rays (GCRs), which are made up of mostly high-energy protons, and to a lesser extent,
alpha particles, electrons, and highly damaging HZE nuclei [1,2]. High linear energy transfer (LET)
radiation in general produce more carcinogenic and complex breaks leading to genomic
instability than low LET radiation[1]. Low LET radiation such as ϒ-rays and x-rays deposit their
energy uniformly but still produce double stranded breaks mainly through the indirect method of
increasing reactive oxygen species (ROS) through the radiolysis of water[1-4]. Low LET-induced
oxidative stress is then defined as an imbalance between a lack of anti-oxidative defenses against
an excess of ROS[5-7].
64

As of now there is little data on the associated risks of exposure to space radiation for a 3-year
Mars mission[8], but it has been estimated that a return trip to Mars could expose astronauts to
600-1000mSv, which is near the NASA astronaut career limit of 800-1200mSv[1,2]. Based on
these estimates, the predicted attributable risk for GCR exposure would suggest a high likelihood
of returning astronauts facing higher risk of leukemia, stomach, colon, lung, bladder, ovarian,
and esophageal cancers[2]. Selection of radioresistant individuals for space travel has been
proposed as one option to circumvent the effects of these massive doses of radiation exposure.
Candidates could be selected based on their rate of DNA damage accumulation and repair, as
measured by comprehensive multi-omic analyses[1], or prioritizing those with lower rate of
mutations measured with clonal hematopoiesis (CH).
75

However, additional clues and protective mechanisms can be gleaned from studies on bacteria
and multicellular extremophiles. Studies on various bacteria have shown increases in mutation
rates as well as increases in virulence, antibiotic resistance, metabolic activity, shorter lag phase
time, and a number of beneficial adaptations in response to short orbital flights[8]. Bacteria such as
Deinococcus radiodurans have increased radiotolerance via enhanced and efficient DNA repair
systems and protection of protiens[10,11]. Many radiotolerant species have the ability to enter
anhydrobiotic states for extended periods of time and so there is a selective pressure to withstand
endogenous reactive oxygen species generated during times of desiccation[3,5,12,13]. As such,
desiccated tardigrades of the species *R. coronifer* and *M. tardigradum* have been shown to
survive the combined effects of the vacuum of space, galactic cosmic radiation on the scale of
9.1Gy, and different spectra of UV radiation at a total dose of $7577kJ/m^2$ at low earth orbit[3,5,12,13].
87

Anhydrobiosis in tardigrade species *R. varieornatus* has been shown to provide better aid in the
prevention of DNA damage accumulation than in the hydrated stage in response to UV-C
radiation as measured by UV-induced thymine dimers[16]. However, hydrated tardigrades in

another study were shown to be more resistant to heavy ion radiation than in their anhydrobiotic
state as demonstrated as the LD_{50} for heavy ions being 4.4kGy to 5.2kGy for hydrated and
dehydrated animals respectively[1]. As the majority of damage of High-LET is deposited along
linear tracks to pass right through cells and induce DSBs as opposed to the indirect ROS effects
of Low-LET, there could be separate mechanisms for defending against varying types of
radiation. An overlap of functional redundancy between proteins of both systems as adult
tardigrades of various species have been shown to be just as tolerant to High-LET as they are to
Low-LET radiation[1,3,5,13,16,17]. This includes the recently-discovered protein, Dsup[18].
99
Dsup was found to co-localize with nuclear DNA through its highly basic c-terminal domain.
This region was later discovered to have a very similarly conserved protein sequence with the
vertebrate HMGN family of proteins in its RRSARLSA consensus found in the nucleosome
binding domain of both proteins[19]. The current proposed mechanism for which Dsup prevents
DNA damage is that it suppresses DNA breaks and acts as a physical protectant against damage
inducing agents[18,20]. It is also possible that Dsup may act functionally similar to HMGN in terms
of enhancing DNA repair mechanisms in addition to its shielding effect[19], but there is no
functional genomics data to yet verify this hypothesis. Indeed, the HMGN proteins colocalize
with epigenetic marks of active chromatin, but also promote chromatin decompaction via
competitively binding with H1, perhaps similar to how Dsup interacts with nucleosomes[19,24,25,26].
110
Understanding the molecular mechanisms underpinning extremophile tolerance could provide us
clues on how human survival in space could be improved through genetic engineering. However,
it is essential to understand how introducing a foreign gene from one species to another affects
the overall system. Here, we developed a lentiviral-transduced HEK293 cell line containing
Dsup (HEK293-Dsup) from which we created a clonal cell line to test functional assays that
measure biomarkers indicative of DNA damage. We then performed RNA-seq analysis on these
cell lines at various doses of radiation and time points in order to understand the dynamics of
how Dsup responds to radiation in the context of human cells. These data show overall high
functionality of the radioprotective effects, with some changes in overall gene expression, and
also provide a resource of data on human cell regulatory changes when utilizing the foreign Dsup
protein.
122
Results
Confirmation of presence of Dsup in HEK293 cells and integration into chromosome 5
After transduction of pLVX Puro Dsup into HEK293 cells, Dsup protein expression was
confirmed via western blot analysis using an HA tag antibody and visualized as a band at 43kDa
along with the larger 110kDa vinculin housekeeping gene **(Supplemental Figure S1a)**. Vinculin
was present in all cell types but only HA tagged antibodies were present in HEK293-Dsup. Dsup
expression was also later confirmed in Fgsea from DESeq2 results **(Supplemental Figure S1b)**.
Dsup remains stably expressed in terms of TPM across all time points and doses of radiation but
not in control HEK293 cells, further validating the expression of Dsup in our engineered cell
line. Next, long read PromethION Nanopore sequencing was done to determine integration site
of pLVX-Puro Dsup. The complete pLVX-Puro Dsup vector plasmid was used as a reference for
mapping of long reads to the reference HEK293 genome. 1,043,859 reads were obtained with a
median of 3,573 reads for HEK293 Dsup with 98% of reads maps and 1.1x genome coverage.
According to FindGenomics digital karyotyping service analysis, our cell lines matched to

HEK293 reference as outlined in Lin et. al[65]. Accordingly, part of the pLVX Puro Dsup vector
mapped to the right arm of chromosome 5 further confirming Dsup lentiviral integration
(**Supplemental Figure S2**).
140
Measuring apoptosis
We sought to look at apoptosis by luciferase fusion annexin V binding of phosphatidylserine. We
detected early apoptosis in live cells measured over the span of 48hrs as a measure of relative
luminescence units for different doses of radiation and 100μM of H_2O_2 (**Figure 1a**). For the 1hr
post treatment cells, there were already significant differences (p=0.01) between controls and
Dsup cells, with control HEK293 cells displaying significantly more apoptotic signals at 2Gys of
radiation and up. About twice as much apoptotic signal was detected up until 6hrs later for
control cells as compared to HEK293-Dsup, except the only significant difference between H_2O_2
treatment was at 1hr post treatment, indicating perhaps a different mechanism of ROS damage
response for Dsup-expressing cells. At 24hrs, this signal began to become saturated and only at
higher doses of radiation again were these signals significant. Interestingly, at 48hrs HEK293-
Dsup no longer had less apoptotic signal than HEK293 cells at 1Gy, 2Gy, and 4G. Notably, at
0Gy, apoptotic signaling continued to rise over time for both control and HEK293-Dsup cells,
indicating that perhaps natural baseline apoptotic signaling pathways are still functional in
response to a stress free environment[60], but nevertheless diminished in HEK293-Dsup after a
certain time post seeding of cells. In conjunction with PS annexin V binding, Dsup expressing
cells also showed significantly less (p = 0.009) apoptotic activity as measured by Casapase-3
activity over a time frame of 22.5 hours (**Figure 1b & 1c**).
159
Differential expression profiling
In order to probe how integration of Dsup affects gene expression and regulatory pathways, we
subjected HEK293 control cells and HEK293-Dsup cells to varying doses of X-ray radiation and
collected cells for RNA-sequencing (RNAseq) library prep across three time points
(**Supplemental Figure S3**), in triplicate. These time points were chosen based off of RNAseq
analyses involved in irradiation of human cell lines[66,67] and to observe differences in how these
cell lines react initially to radiation stress (0.5, 1, 2, 4 Gy) and overtime through the iterative
process of DNA damage repair[49,53]. After differential gene expression analysis was conducted
using DESeq2 normalization and GSEA analysis, PCA and t-SNE plots were generated to
observe any clustering of genotype, time, and radiation dose. We observed the stronger variance
and clustering of time followed by genotype and then radiation dose (**Figure 2**). Accordingly,
HEK293 Dsup at 0Gys baseline separated with higher variance from the other groups across the
three timepoints, indicating differences in baseline expression between Dsup expressing cells and
controls. When comparing genotypes at individual hours, clustering can be seen more clearly
between HEK293-Dsup and controls.
175
Similarly, differences in gene expression was observed in volcano plots for each timepoint. Here
we see that DSUP is the most upregulated gene across all time points in terms of adjusted p-
value and log_2 fold change. Overall we observed more upregulation of genes in Dsup-expressing
cells (63%) compared to controls when all time points are analyzed at once **Figure 3b.** Across
each time point including each dose of radiation however, there appears to be more differential
gene expression occurring later at 6hrs and 24hrs as compared to 1hr post irradiation
(**Supplemental Figure S5b,d,e**). Heatmaps in **Figure 4a.** reveal a similar pattern of differential
expression. Here we see differential signatures at 1hr post irradiation **(Figure 3a &**

Supplemental Figure S4a) at 0Gy, 0.5Gy, and 1Gy for HEK293 cells that remain consistent
until higher doses of radiation at 2Gy and 4Gy. The opposite pattern is seen in HEK293 Dsup
cells at 1hr, but begin to look more like control cells at higher doses of 2Gy and 4Gy of
radiation. At 6hrs (**Supplemental Figure S4c**) and 24hrs (**Supplemental Figure S4d**)
differences in expression remain the most consistently changed across all radiation doses but
differential expression at 0Gys in HEK293 Dsup displays the most dramatically enriched
changes in expression as compared to 0Gy baseline HEK293 cells.
191
Gene set enrichments related to stress response
To assess what baseline expression looks like in HEK293 Dsup cells in comparison to wild type,
enrichment analysis was done on just the intersection of genes shared among all three time
points. Enrichr was used to generate gene set libraries using a modification of Fishers exact test
to generate a combined z-score a p-value in order to rank genes. The Histone Modification gene-
set library from ENCODE was analyzed to obtain a gene set library based on histone
modifications (**Figure 4a & 4b**). We separated Running Enrichment Score plots by time for both
upregulated and downregulated gene sets. Gene sets were ranked by adjusted p-value correlated
with density of genes in each gene set. Downregulated pathways are shown in (**Supplemental
Figure S5a).** At 1hr UV damage responses, CD5 targets, and haptotaxis are downregulated while
at 6 and 24hrs, UV damage responses, homologous recombination, rRNA metabolic processes,
and inflammation responses are down. (**Supplemental Figure S5b).** At 1hr, Zinc Finger Protein
target genes, nucleic acid binding, and metastasis related gene sets were upregulated, while at 6
and 24hrs inflammation, growth regulation, and hypoxia responses were upregulated. The
intersection for all time points of the upregulated and downregulated gene sets were also
analyzed. **Figure 4c** shows mostly a downregulation of gene sets belonging to ubiquitin protein
ligases, nuclear protein import, and whole sets of certain chromosomal regions corresponding to
ch13q14 and chr2q33. **Figure 5d** shows a shared upregulation of chromatin modulators involved
with histone methylation and zinc finger protein targets.
211
Enriched KEGG pathways
Next we looked at overall differential expression between HEK293 Dsup cells and WT HEK293
cells. This included gene set enrichment analysis for all time points and doses of radiation
combined. Here, we ranked our multivariate DESeq2 results by adjusted p-value multiplied by
Log_2FC (<0.05 for p-adjusted and >1.5 for Log_2FC). For biological process we see an
upregulation of metabolism, response to stimuli, and growth and cell proliferation (**Figure 5a,
5b, 5c**). A few of the top cellular component categories included cell membrane, nucleus, and
membrane-enclosed lumen. For molecular function, we observed significant enrichment of
protein binding, ion binding, and nucleic acid binding gene sets, as well as chromatin binding,
antioxidant activity, and oxygen binding. Of note, the top upregulated pathways included many
that are common in cancerous phenotypes that are resistant to different types of therapy
including radiation therapy. We also see up regulation of lipolysis, a common stress response in
exposure to oxidation and ionizing radiation[14,84]. There is also a downregulation of oxidative
phosphorylation, indicative of radiation resistant hypoxic cells[69,70].
226
As Dsup shares amino acid sequence homology with the high mobility group proteins, in
particular HMGN1 and 2, we checked to see if expressing Dsup in HEK293 cells would
potentially lead to a perturbance of these expression levels, as they are hypothesized to bind to
the same nucleosome structures[19]. Using our DESeq2 input and running Fsgea, TPM was

counted for each of the HMGN1 and 2 proteins. While levels of transcripts appear to fluctuate
across different doses of radiation and time, both genes were still found to be stably expressed in
HEK293 Dsup cells as compared to controls **(Supplemental Figure S6)**.
234
Discussion
Chromosomal aberrations due to GCR-induced DSBs can be a major detriment to future long-
duration space flight missions. In the NASA Twins Study data, a greater rate of chromosomal
inversions were found during space flight, and after, indicating genomic instability and
rearrangement[8]. Since the tardigrade Dsup protein has been shown to suppress DNA damage we
tested if a stably-transfected human cell line could also sustain such protective phenotypes, and if
there would be other changes to human gene expression networks. Overall, the results validate
the functional radioprotective capacity of Dsup in human cells, to establish the stable line for use
by others, and provide new molecular details about the functional genomic impact of Dsup
integration into the human genome.
245
Interestingly, there were significant upregulated enrichments in DSUP expressing cells for
pathways associated with histone H3 acetylation and methylation. Specifically, an upregulation
of both the gene signatures of the repressive mark H3K27me3 and the active mark H3K4me3
indicative of bivalent gene targets associated with EZH2, the catalytic subunit of PRC2. These
findings support the model that Dsup binds to nucleosomes and remodels chromatin as an
architectural chromatin protein similar to its homologous counterparts the HMGN proteins[19].
This same expression pattern from our dataset was also observed in a study where HMGN1 was
overexpressed in Ts1Rhr mouse models where EZH2 targets and H3K27ME3 pathways were
significantly enriched in overexpressing HMGN1 cells as compared to wild type[27]. This could be
the result of overexpression of chromatin architectural proteins as TPM normalization of
HMGN1 appears stable in DSUP expressing cells.
257
Of note, as damage from radiation increased, so did many of our top pathways at 1hr post
irradiation involving HAT activity. It has been demonstrated in multiple *in vivo* and *in vitro*
models that cells lacking high mobility group proteins are far more sensitive to radiation and
other damaging agents as these chromatin architectural modifiers are shown to enhance the
recognition and processing of DNA damage by making DNA lesions more accessible to repair
machinery[36,37,38]. With the decompaction of chromatin, High Mobility Group proteins also induce
kinks in exposed DNA in order to make these sites more accessible as well. Given that Dsup is
about 4 times as large as HMGN1, it is possible that Dsup could confer human cells with greater
chromatin decompaction and accessibility to sites of damage[19,38], or work with other proteins to
accomplish such reorganization.
268
Strikingly, these patterns of chromatin remodeling pathways also move together with an
upregulation of DNA damage repair and response pathways within 1hr of radiation and become
more pronounced with greater doses. Here we see even at 0.5Gy and 1Gy of radiation an
upregulation of characteristic radiation damage response pathways[39]. At higher doses of 2Gy and
4Gy there is a more pronounced upregulation of multiple damage response pathways that include
upregulation of members of double strand break and repair pathways[40-46]. In addition, a number
of other Zinc Finger protein targets were upregulated, which could be indicative of an enhanced
DNA damage repair process[73]. GSEA of KEGG pathways also showed an upregulation of

lipolysis, a known protective metabolism stress response of both humans and tardigrades[14,84].
Long fatty acid chains can often serve as damage response signaling molecules and can regulate
inflammation as well as mediate recovery after injury[84]. Fatty acid metabolism was shown to be
affected dynamically by space flight in the NASA Twin study and as such, spikes in midflight
expression levels of genes in the lipolysis regulation pathways were upregulated[84].
282
Cell cycle checkpoint pathways, such as those belonging to the DNA damage response CHEK2
network, also became upregulated in response to higher doses at 1hr as well as the ANP32B
pathway necessary for cell cycle progression from G1 to S phase[47,48]. Dsup-expressing cells
could move through cell cycle progression phases faster than wild type cells because of an
increased rate of damage clearance that's necessary to enter the next cell cycle phase. DNA
damage repair is an iterative process and cells enter a state of senescence until the damage is
repaired sufficiently or if the accumulation of damage is beyond repair then apoptotic pathways
are triggered[49].
291
Given these broad results, one can imagine Dsup and related genes helping in the creation of a
temporal, radiation, or molecule controlled, inducible Dsup system that is activated in response
to intense periods of galactic cosmic radiation. Indeed, such work has already begun for atomic
energy workers and in other trials of engineered cancer therapies that leverage CRISPR[85]. Even if
not directly used as a therapeutic, the gene expression perturbation data here, as well as the
stably-expressing cell line, can serve as resources to guide future research on radioresistance and
baseline levels of transcriptional disruption after the stable integration of a radioresistance gene.
Together, such tools and data sets will help keep astronauts safer as missions move farther from
Earth, become riskier, and are laden with ever-increasing levels of GCR and irradiation.
301
Methods
Plasmid Design and Extraction
The complete coding sequence for DSUP from *Ramazzottius varieornatus* as described in
Hashimoto et. Al was obtained from The Kumashi Genome Project database. This sequence was
cloned into a custom pLVX Puro plasmid lentiviral vector with a constitutive CMV promoter.
Dsup was ligated into the 5' EcoR1 and 3' XbaI in the multiple cloning site of pLVX Puro.
GeneWiz® Gene synthesis services was used to codon optimize DSUP for expression in
mammalian systems. An influenza Hemagglutinin tag was synthesized on the N-terminal region
of DSUP to minimize interference of c-terminal binding of DSUP to nucleosomes and for
downstream antibody detection.
312
The plasmid was then transformed into DH5α Competent Cells. 10ng of DSUP pLVX Puro
plasmid was added to 100μl of competent cells and incubated on ice for 30mins followed by heat
shock for 45 seconds in a 42°C water bath and then incubated on ice for 2mins. 900μl of S.O.C.
medium was added to cells and shook at 225rpm at 37°C for 1hr before 100ul were spread using
glass beads on LB 100μg/ml ampicillin plates. Plates were incubated overnight at 37°C and
colonies were picked the following day for miniprep. Multiple colonies were picked and grown
in 4mls of LB broth containing 100μg/ml of ampicillin shaking overnight at 250rpm and 37°C.
DNA from cultures were extracted the following day using Promega™ Wizard® Plus SV
Miniprep kit. Plasmid DNA was eluted from spin columns in 30μl of nuclease free water.
322

500ng of plasmid DNA sample was restriction enzyme digested in NEB 10X CutSmart® Buffer
with 1μl of EcoRI and XbaI at 37°C for 3hrs before run on a gel using NEB 6X loading dye and
Invitrogen™ 1KB plus DNA ladder for 45mins at 130V. Samples cut at 1377 Bps corresponding
to DSUP and 8069 base pairs corresponding to pLVX Puro vector backbone were chosen for
maxiprep using Invitrogen™ PureLink™ HiPure Plasmid Filter Maxiprep Kit. The correctly
digested sample was grown in 500mls of LB with 100μg/ml ampicillin overnight shaking at
250rpm at 37°C. In parallel pLVX Puro empty vector, pMD2.G VSV-G envelope plasmid, and
psPAX2 lentiviral packaging plasmid transformed and extracted using the above protocols were
also grown overnight up for maxiprep. 300μl of nuclease free water was passed through the
precipitator to elute the plasmid DNA into 1.5ml Eppendorf microcentrifuge tubes. Plasmid
DNA was then quantified using Qubit™ 1X dsDNA HS assay. 1μg and above was quantified for
each plasmid DNA sample.
335
Generation of lentiviral transduced stable HEK293 DSUP Cell line
HEK293 cells and HEK293T cells at passage 13 and 17 respectively were donated by Dr. Levitz
at Weill Cornell were grown and incubated at standard incubation conditions at 37°C with 5%
CO_2 in complete DMEM media supplemented with 10% FBS, 1% Pen/Strep, and 1X Glutamax
on T-75 Nunc™ EasYFlasks™. HEK293T cells were split at 1:10 and $2.2x10^6$ cells were seeded
on two PLL coated 100mm plates for transfection of pLVX Puro DSUP and empty vector pLVX
Puro as an empty vector control for downstream experiments.
343
Mirus TransIt® Lentivirus System was used to transfect 80-95% confluent HEK293T cells
grown the previous day in Opti-MEM reduced serum media. TransIT® Lenti Reagent was
warmed to room temperature and vortexed gently before using. 200μl of Opti-Mem™ media was
added to sterile 1.5ml Eppendorf microcentrifuge tubes while 1.65 pmol of either pLVX Puro
DSUP or empty vector pLVX Puro was combined with 1.3pmol of psPAX2 plasmid and
0.72pmol of pMD2.G plasmid. 2μg of plasmid mixture was then transferred to the tube
containing Opti-MEM. 6μl of TransIT®-Lenti Reagent was added to the diluted plasmid DNA
mixture and incubated for 10mins at room temperature. The mixture was then added dropwise to
different areas of the plate and gently rocked back and forth to ensure even distribution. Plates
were stored at 37°C with 5% CO_2 for 48hrs before packaged lentivirus particles were harvested.
Viral particles were filtered through a 10ml syringe with a 0.45μm PVDF filter and separated
into 1ml aliquots to be flash frozen and stored in -80°C.
356
Mirus TransduceIT™ Reagent was used to efficiently transduce HEK293 cells plated on 100mm
plates at a density of $2.2x10^6$ cells seeded the previous day. TransduceIT™ Reagent was added to
the HEK293 cells along with 1ml of virus particles distributed evenly over the plate dropwise.
Cells were incubated for 48hrs before being selected with 2μg/ml of puromycin as determined by
puromycin kill curve analysis on WT HEK293 cells. After 1 week of selection HEK293 DSUP
and HEK293 empty vector cells were obtained and expanded on T-75 flasks.
363
Serial dilution cloning was performed on HEK293 Dsup cells in order to obtain clonal cell lines.
200μl of cell suspension at $2x10^4$ was added to the first well A1 of a 96 well plate. 100μl of
complete DMEM media was added to each well except A1 using a multichannel pipette. Using a
single channel pipette 100μl of cell suspension was added below to well B1 and mixed before

100μl of that cell suspension was added to the next well below it until dilutions were made to
well H1. Then using a multichannel pipette 100μl of those cell suspensions were added to the
next columns further creating serial dilutions that would span across the plate. Cells were
incubated at standard conditions and after 5 days individual clones were isolated in some wells
as detected by microscopy. Once these wells were confluent, they were treated with trypsin and
resuspended in complete DMEM media to be expanded on 24 well plates followed by 6 well
plates, T-25 flasks, and finally T-75 flasks.
375
Western Blot
Confirmation of Dsup was assessed by western blot. HEK293 Dsup, HEK293 empty vector, and
HEK293 cells were plated in triplicates on 100mm dishes at a seeding density of $2.2x10^6$ cells
per dish. The following day after incubation the dishes were placed in ice and washed with ice
cold Tris-buffered saline. The TBS was aspirated and 1ml of ice-cold RIPA buffer was added to
dissolve cell membranes. Cell lysates were collected in 2ml pre-cooled Eppendorf tubes.
Constant agitation was maintained at 1400rpm for 30mins and 4°C before being spun down at
16,000xg for 20mins at 4°C. The supernatant was transferred to a new pre-cooled
microcentrifuge tube and 10μl of lysate was used to determine protein concentration using
Qubit™ Protein Assay kit. 20ug of each sample was then added to an equal volume of 2x
Laemmli sample buffer before being boiled at 95°C for 5mins. Equal amounts of protein along
with a PageRuler™ Plus Pre-stained protein ladder were loaded into a Bio-Rad 10 well 4-20%
Mini-PROTEAN® TGX™ Precast Protein Gel. Protein separation by electrophoresis was then
performed in a Mini PROTEAN Tetra cell with 1liter of Tris/Glycine/SDS running buffer. The
gel was run a 50V for 5mins and then 100V for 1hr.
391
The gel was then placed in 1X transfer buffer for 15mins and the transfer sandwich was
assembled using a 0.20μm Nitrocellulose membrane and sponges. The assembled transfer
sandwich was placed in the transfer cassette and proteins were transferred for 1hr at 100Volts
and 350mA current in the Mini PROTEAN Tetra cell with 1L of transfer buffer. The blot was
briefly rinsed with Millipore water and stained with Ponceau S solution to confirm transfer of
proteins. The blot was then rinsed three times with 1X TBST and blocked in 5% milk in 1X
TBST at room temperature for 1hr. R&D Systems recombinant mouse HA Tag antibody
(MAB0601) and control Bio-Rad Mouse anti Human Vinculin antibody (V284) was diluted to a
working concentration of 0.1μg/ml and a 1:500 dilution in 1% milk in 1X TBST respectively and
the primary antibodies were incubated on the membrane overnight at 4°C. The following day the
blot was rinsed 5 times for 5mins each with 1X TBST before adding the Invitrogen Goat anti-
Rabbit IgG (H+L) Cross-Absorbed Secondary Antibody DyLight 800 (SAS-10036) and Rabbit
anti-Mouse IgG (H+L) Cross-adsorbed Secondary Antibody Alexa Fluor Plus 680 (A32729)
diluted to working concentrations of 1:5000 dilution and 0.1μg/ml respectively in 1% milk in 1X
TBST. The blot was incubated at room temperature for 1hr, rinsed 5 times for 5 mins in 1X
TBST and then imaged on a Li-Cor Odyssey® CLx imagine system using both 700nm and
800nm channels.
409
Confirmation of lentiviral insertion site using Oxford Nanopore long read sequencing
DNA from HEK293 Dsup cell line at passage 20 and 22 and the parental HEK293 cell line at
passage 18 were extracted using Trizol as mentioned above. Briefly, at phase separation ethanol
was added to the lower phenol-chloroform phase and interphase, centrifuged and pelleted, and

resuspended in sodium citrate. After 2 wash steps with 75% ethanol, the DNA pellet was allowed
to air dry and then resuspended in 8mM NaOH. DNA concentration was assessed via QuBit 1X
dsDNA assay and prepared for long read sequencing using the Ligation Sequencing Kit (LSK-
109) for Oxford Nanopore. The adapter ligated gDNA was sequenced using a PromethION flow
cell and base called in real time using Guppy 3.4. The data was analyzed using FindGenomics
digital karyotyping service.
420
Cell Irradiation
Cells were incubated overnight post seeding at 37°C with 5% CO_2 before irradiation. The
following day cells were taken out of the incubator and subjected to X-rays in a Rad Source
Technologies RS 2000 Biological research irradiator and irradiated at 1Gy, 2Gy, 4Gy, and 16Gy
while the corresponding 0Gy plate was kept outside the incubator as a no treatment control. For
the RNAseq experiment, cells were irradiated at 0.5Gy, 1Gy, 2Gy, and 4Gy. Dose time was
determined divided by the dose rate of 2Gy/min.
428
Annexin V Apoptosis Assay
Live cell real time monitoring of apoptosis was conducted using Promega RealTime-Glo™
Annexin V Apoptosis Assay. HEK293, HEK293 Dsup, and HEK293 empty vector cells were
seeded on 6 separate 96 Well White Polystyrene microplates at a density of 1,000 cells/well at a
volume of 100µl in triplicates and incubated at 37°C with 5% CO_2 overnight. 100µl of complete
medium were added to wells not containing cells as no cell and no compound controls. Each
plate was a separate treatment condition and each row contained all cell types in triplicates and
corresponded to a different time point from which apoptosis would be assayed.
437
The following day cells were taken out of the incubator and subjected to X-ray irradiation and
hydrogen peroxide treatment as indicated above. Apoptosis was assessed at 1hr, 6hrs, and 24hrs
post irradiation. To prepare the 2X detection reagent, 1000X Annexin NanoBiT® substrate was
added to prewarmed DMEM complete medium at a 500-fold dilution. 1000X $CaCl_2$ was then
added to combined DMEM complete medium with diluted NanoBiT® substrate at a 500-fold
dilution. 1000X Annexin V-SmBiT and 1,000X Annexin V-LgBiT was added at a 500-fold
dilution to the combined substrate, $CaCl_2$, complete medium and was inverted gently a few times
to mix. 100µl of the 2X detection reagent was then added to each 96 well plate using a
multichannel pipette and mixed at 700rpm for 30 seconds on a plate shaker. Luminescence and
fluorescence at 485nm$_{Ex}$/525-530nm$_{Em}$ data were measured on a Promega™ Glowmax® Plate
Reader.
449
Microscopy for Caspase-3 assay:
HEK293 cells containing the Empty Vector, DSUP, or no construct were cultured in DMEM
supplemented with 10% FBS. Cells were washed once with PBS before trypsinization. 5 mL of
Trypsinized cells were transferred to 10 mL of FluoroBrite DMEM supplemented with 10% FBS
and 1% 100x glutamax in a 15 mL conical tube. Cells were pelleted at 330 xg for 3 min. Media
trypsin mix was removed before resuspending the cell pellet in 5 mL of the FluoroBrite DMEM
described above. 10 µL of resuspended cells were added to 10 µL of trypan blue from the BioRad
Cell counting kit. 10 µL of this cell trypan mix was added to a cell counting slide from the same
kit. The slide was placed in a TC20™ Automated Cell Counter for cell counting. Based on the
count of live cells, each cell line was diluted to 75 cells/µL in the FluoroBrite DMEM described

above. 200 µL of this mix was added to each well of a Corning 3603 96 well microscopy grade
plate for a total of 15,000 cells per well. The Plate was spun at 330 xg for 3 min. The plate was put
in a humidified 37°C incubator with 5% CO_2 for 4 hours to allow cell adherence to the bottom of
the well.
464
After cell adherence, plates were removed from the incubator and inoculated with 10 µL of a mix
containing 105 µM Hoechst 33342 (diluted from Thermo H3570), 26.25 µM Caspase dye (Diluted
from Sartorius 4440), and 277.2 µM Camptothecin (diluted from Millipore Sigma C9911) in water.
This inoculation amounts to 5 µM Hoechst 33342, 1.25 µM Caspase dye, 13.2 µM Camptothecin
in the well. The plate was sealed with a Breathe-Easy® sealing membrane before incubating for
30 min in the same incubator described above.
After incubation, cells were placed in a humidified 37°C ImageXpress Micro Confocal High-
Content Imaging System with 5% CO_2. All wells of the plate were imaged with a 10x Plan Apo
Lambda lenses in wide field mode. 9 fields from each well were collected with a 4 ms DAPI
exposure and a 93 ms FITC exposure. This process was repeated every 45 min for 22.5 hours (30
time points).
476
Images were analyzed using MetaXpress. Briefly nuclei were segmented using the built-in find
round objects function applied to the DAPI channel. The local background of these objects was
defined by using the expand objects without touching function (with the ultimate option
checked). The difference between the expanded objects and the nuclei was used to define the
background area. The average FITC intensity was calculated for the background and for the
nuclear regions. The difference between the nuclear and background FITC intensity was
calculated and used as the Caspase signal for each cell at each time point. Upon inspection of the
distribution, a cutoff value of 650 caspase signal units was determined to be an appropriate cutoff
for cell death as there was a bimodal break around that signal intensity. Cells with lower than
650 units of caspase signal were called non-apoptotic, cells with greater than 650 units of
caspase signal were considered apoptotic. The percentage of apoptotic cells was calculated for
every well at every time point. These points were plotted for all wells tested for each cell line. A
bar graph was made with error bars at 1 standard deviation of the apoptotic rate of the cells at the
final time point (22.5 h post inoculation). A heterostatic two tailed student t-test was performed
to test the null hypothesis that the Empty Vector and DSUP containing cell lines have equal
apoptotic rates.
493
RNA sequencing and Differential Gene Expression Analysis
Each cell type was harvested from T-175 at around 75% confluency. Viability and cell count
were assessed using TrypanBlue stain on a Countess™ Automated Cell Counter. Each cell type
had over 90% viable cells and were then seeded onto 6 well round bottom plates in triplicates for
each cell type, treatment, and time point at a density of 30,000 cells/well. Cells were irradiated
the following day post seeding at 0.5Gy, 1Gy, 2Gy, and 4Gy of X-rays while 0gy of no treatment
controls were kept outside the incubator for the duration of radiation treatment. Cells were
incubated at standard cell culture conditions and harvested at 1hr, 6hrs, and 24hrs post treatment.
502
A total of 93 samples were selected for NEBNext® Ultra™ II RNA library prep. These include 3
HEK293 DSUP replicates x 3 time points x 4 radiation doses + 3 0gy Controls. The same was
done for HEK293 cells and 3 HEK293 empty vectors at 2Gy and 6hrs were included. Media was

aspirated, washed once with 1xPBS, and cells were detatched with trypsin before being
resuspended and collected into 2ml Eppendorf tubes to be spun down at 6,000xg and pelleted.
Media was aspirated and then the cell pellet was resuspended and dissolved in Trizol. Trizol
extractions were performed as mentioned above but this time RNA was isolated carefully from
the upper aqueous layer and pelleted before being eluted in 30μl of RNAse free water. RNA
concentration was measured using Qubit™ HS RNA assay and RIN and fragment distribution
was obtained using Agilent High Sensitivity RNA ScreenTape System for TapeStation 2200.
RIN was above 9 for all samples and 250ng of RNA sample was brought forward for rRNA
ribosomal depletion using NEBNext® rRNA Depletion Kit for human/mouse/rat samples.
Following depletion, RNA samples were library prepped and adapter ligated with single Index
Primer set 1 NEBNext® Multiplex Oligos for Illumina®. Libraries were then sequenced at 40
million single end 100bp reads on Novaseq 6000.
518
Read alignment
Libraries were single end 100bp sequenced on NovaSeq6000, targeting 40 million reads. Raw
data was base called using Illumina Basespace. GTF and FASTA files for the human reference
genome (GRCh38) were downloaded from ENSEMBL. Both GTF and FASTA files were edited
to contain the sequence for the protein coding tardigrade gene DSUP. After a reference index
was created using STAR (ver 2.7), the reads were then mapped to the human genome[76]. The
output BAM files were then sorted using Samtools[78]. Read counts were obtained using Feature
Counts in which multi-mapped reads were removed [77].
527
Normalization using DESeq2
The variance mean was estimated and differential expression was computed using a negative
binomial distribution using DESeq2 from raw read count data. First a count matrix was
generated and for each gene a general linearized model is fitted. Read counts are then modeled as
a negative binomial distribution and the mean is taken as a quantity that is proportional to the
concentration of DNA fragments from the gene in the sample and scaled by a normalization
factor. The corresponding size factors are then computed using a median on ratios method. When
comparing two groups the GLM fit returns coefficients that's indicates the expression strength of
a gene and its corresponding Log_2FC. Empirical Bayes shrinkage was used to estimate dispersion
parameters within groups to assess for variability between replicates and for shrinkage of fold
change estimation to then rank genes. To test for differential expression, DESeq2 uses a Wald
test which returns a z-statistic and p-values from a subset of genes that are adjusted using
multiple testing via Benjamini and Hochberg correction[79].
541
Fgsea analysis
Fgsea, a Bioconductor R-package, was used to obtain pre-ranked expression to identify classes
of overrepresented genes[64]. Predefined gene sets from MsigDB were grouped together by their
involvement in the same biological pathway or by location on chromosome[80]. Features were
ranked by differential expression analysis adjusted p-value with corresponding Log_2FC to get
normalized enrichment scores. First an enrichment score was calculated by running a sum
statistic on each gene in the set to assess the degree to which overrepresented gene sets at the top
and bottom of the ranked list deviates from zero using a weighted Kolmogorov-Smirnov like
statistic. Then the significance levels of the enrichment scores were calculated to get nominal p-
values by calculation of cumulative gene set enrichment values. Last normalized enrichment

scores (NES) are returned by normalizing each enrichment score by the size of the gene set and
then the proportion of false positives are controlled by calculating the FDR[81].
554
Enrichr analysis
Common genes among different time points for 0Gys of radiation were obtained as the
intersection of all genes across the three timepoints. Erichr was used to identify enriched gene
sets using a hypergeometric model. To compute enrichment, Enrichr uses the Fisher exact test
and then computes a mean rank and standard deviation from the expected rank for the terms in
the gene set library. A z-score for deviation can also then be computed using a lookup table for
expected ranks with associated variances, the output of which is then used as corrected score for
ranking terms. In addition, Enrichr also can combine the computed p-value from the Fisher exact
test with the z-score of the deviation from the expected rank by multiplying log(p) by z to get c
the combined score[74,75].
565
WebGestalt analysis
To observe overall effect of Dsup in human cells, KEGG pathways were visualized with the web
based integrative data mining system WebGestalt[83]. Input data set from DESeq2 was ranked by
adjusted p-value multiplied by Log_2FC with a p-value cutoff of 0.05 and Log_2FC cutoff of 1.5
and used for pathway enrichment analysis. This list of genes of interest was then compared to a
reference gene set in KEGG using a hypergeometric test to assess the significance of enrichment
for a category in the input gene set. The hypergeometric test then provides for an over and
underrepresentation of individual genes. Gene sets can then be explored and organized using
visualization interfaces through KEGG tables[82,83].
575
576

References
1. Cortese, F., Klokov, D., Osipov, A., Stefaniak, J., Moskalev, A., Schastnaya, J., Cantor,
C., Aliper, A., Mamoshina, P., Ushakov, I., Sapetsky, A., Vanhaelen, Q., Alchinova, I.,
Karganov, M., Kovalchuk, O., Wilkins, R., Shtemberg, A., Moreels, M., Baatout, S.,
Izumchenko, E., … Zhavoronkov, A. (2018). Vive la radiorésistance!: converging
research in radiobiology and biogerontology to enhance human radioresistance for deep
space exploration and colonization. *Oncotarget*, *9*(18), 14692–14722.
https://doi.org/10.18632/oncotarget.24461
2. Cucinotta, F. A., Kim, M. H., Chappell, L. J., & Huff, J. L. (2013). How safe is safe
enough? Radiation risk for a human mission to Mars. *PloS one*, *8*(10), e74988.
https://doi.org/10.1371/journal.pone.0074988
3. Jönsson K. I. (2019). Radiation Tolerance in Tardigrades: Current Knowledge and
Potential Applications in Medicine. *Cancers*, *11*(9), 1333.
https://doi.org/10.3390/cancers11091333
4. Gusev, O., Nakahara, Y., Vanyagina, V., Malutina, L., Cornette, R., Sakashita, T.,
Hamada, N., Kikawada, T., Kobayashi, Y., & Okuda, T. (2010). Anhydrobiosis-
associated nuclear DNA damage and repair in the sleeping chironomid: linkage with
radioresistance. *PloS one*, *5*(11), e14008. https://doi.org/10.1371/journal.pone.0014008
5. Rizzo, A. M., Altiero, T., Corsetto, P. A., Montorfano, G., Guidetti, R., & Rebecchi, L.
(2015). Space flight effects on antioxidant molecules in dry tardigrades: the TARDIKISS

experiment. *BioMed research international*, *2015*, 167642.
https://doi.org/10.1155/2015/167642

6. Daly, M. J., Gaidamakova, E. K., Matrosova, V. Y., Vasilenko, A., Zhai, M., Leapman,
R. D., Lai, B., Ravel, B., Li, S. M., Kemner, K. M., & Fredrickson, J. K. (2007). Protein
oxidation implicated as the primary determinant of bacterial radioresistance. *PLoS*
biology, *5*(4), e92. https://doi.org/10.1371/journal.pbio.0050092

7. Wang, L., He, L., Bao, G., He, X., Fan, S., & Wang, H. (2016). Ionizing Radiation
Induces HMGB1 Cytoplasmic Translocation and Extracellular Release. *Guo ji fang she yi*
xue he yi xue za zhi = International journal of radiation medicine and nuclear
medicine, *40*(2), 91–99.

8. Taylor, L., Vitaterna, M. H., Afkarian, M., Afshinnekoo, E., Ahadi, S., Ambati, A., …
Turek, F. W. (2019). The NASA Twins Study: A multidimensional analysis of a year-
long human spaceflight. *Science (New York, N.Y.)*, *364*(6436), eaau8650.
https://doi.org/10.1126/science.aau8650

9. Taylor P. W. (2015). Impact of space flight on bacterial virulence and antibiotic
susceptibility. *Infection and drug resistance*, *8*, 249–262.
https://doi.org/10.2147/IDR.S67275

10. Hashimoto, T., & Kunieda, T. (2017). DNA Protection Protein, a Novel Mechanism of
Radiation Tolerance: Lessons from Tardigrades. *Life (Basel, Switzerland)*, *7*(2), 26.
https://doi.org/10.3390/life7020026

11. Daly, M. J., Gaidamakova, E. K., Matrosova, V. Y., Vasilenko, A., Zhai, M., Leapman,
R. D., Lai, B., Ravel, B., Li, S. M., Kemner, K. M., & Fredrickson, J. K. (2007). Protein
oxidation implicated as the primary determinant of bacterial radioresistance. *PLoS*
biology, *5*(4), e92. https://doi.org/10.1371/journal.pbio.0050092

12. Gusev, O., Nakahara, Y., Vanyagina, V., Malutina, L., Cornette, R., Sakashita, T.,
Hamada, N., Kikawada, T., Kobayashi, Y., & Okuda, T. (2010). Anhydrobiosis-
associated nuclear DNA damage and repair in the sleeping chironomid: linkage with
radioresistance. *PLoS one*, *5*(11), e14008. https://doi.org/10.1371/journal.pone.0014008

13. Weronika, E., & Łukasz, K. (2017). Tardigrades in Space Research - Past and
Future. *Origins of life and evolution of the biosphere : the journal of the International*
Society for the Study of the Origin of Life, *47*(4), 545–553.
https://doi.org/10.1007/s11084-016-9522-1

14. Förster, F., Beisser, D., Grohme, M. A., Liang, C., Mali, B., Siegl, A. M., Engelmann, J.
C., Shkumatov, A. V., Schokraie, E., Müller, T., Schnölzer, M., Schill, R. O., Frohme,
M., & Dandekar, T. (2012). Transcriptome analysis in tardigrade species reveals specific
molecular pathways for stress adaptations. *Bioinformatics and biology insights*, *6*, 69–96.
https://doi.org/10.4137/BBI.S9150

15. Neumann, S., Reuner, A., Brümmer, F., & Schill, R. O. (2009). DNA damage in storage
cells of anhydrobiotic tardigrades. *Comparative biochemistry and physiology. Part A,*
Molecular & integrative physiology, *153*(4), 425–429.
https://doi.org/10.1016/j.cbpa.2009.04.611

16. Horikawa, D. D., Cumbers, J., Sakakibara, I., Rogoff, D., Leuko, S., Harnoto, R.,
Arakawa, K., Katayama, T., Kunieda, T., Toyoda, A., Fujiyama, A., & Rothschild, L. J.
(2013). Analysis of DNA repair and protection in the Tardigrade Ramazzottius
varieornatus and Hypsibius dujardini after exposure to UVC radiation. *PLoS one*, *8*(6),
e64793. https://doi.org/10.1371/journal.pone.0064793

17. Nilsson, E. J., Jönsson, K. I., & Pallon, J. (2010). Tolerance to proton irradiation in the
eutardigrade Richtersius coronifer--a nuclear microprobe study. *International journal of*
radiation biology, *86*(5), 420–427. https://doi.org/10.3109/09553000903568001
18. Hashimoto, T., Horikawa, D. D., Saito, Y., Kuwahara, H., Kozuka-Hata, H., Shin-I, T.,
Minakuchi, Y., Ohishi, K., Motoyama, A., Aizu, T., Enomoto, A., Kondo, K., Tanaka, S.,
Hara, Y., Koshikawa, S., Sagara, H., Miura, T., Yokobori, S. I., Miyagawa, K., Suzuki,
Y., … Kunieda, T. (2016). Extremotolerant tardigrade genome and improved
radiotolerance of human cultured cells by tardigrade-unique protein. *Nature*
communications, *7*, 12808. https://doi.org/10.1038/ncomms12808
19. Chavez, C., Cruz-Becerra, G., Fei, J., Kassavetis, G. A., & Kadonaga, J. T. (2019). The
tardigrade damage suppressor protein binds to nucleosomes and protects DNA from
hydroxyl radicals. *eLife*, *8*, e47682. https://doi.org/10.7554/eLife.47682
20. Mah, L., El-Osta, A. & Karagiannis, T. γH2AX: a sensitive molecular marker of DNA
damage and repair. *Leukemia* **24,** 679–686 (2010). https://doi.org/10.1038/leu.2010.6
21. Subramanian, M., Gonzalez, R. W., Patil, H., Ueda, T., Lim, J. H., Kraemer, K. H.,
Bustin, M., & Bergel, M. (2009). The nucleosome-binding protein HMGN2 modulates
global genome repair. *The FEBS journal*, *276*(22), 6646–6657.
https://doi.org/10.1111/j.1742-4658.2009.07375.x
22. Birger, Y., West, K. L., Postnikov, Y. V., Lim, J. H., Furusawa, T., Wagner, J. P., Laufer,
C. S., Kraemer, K. H., & Bustin, M. (2003). Chromosomal protein HMGN1 enhances the
rate of DNA repair in chromatin. *The EMBO journal*, *22*(7), 1665–1675.
https://doi.org/10.1093/emboj/cdg142
23. He, B., Deng, T., Zhu, I., Furusawa, T., Zhang, S., Tang, W., Postnikov, Y., Ambs, S., Li,
C. C., Livak, F., Landsman, D., & Bustin, M. (2018). Binding of HMGN proteins to cell
specific enhancers stabilizes cell identity. *Nature communications*, *9*(1), 5240.
https://doi.org/10.1038/s41467-018-07687-9
24. Apelt, K., Zoutendijk, I., Gout, D. Y., Wondergem, A. P., van den Heuvel, D., &
Luijsterburg, M. S. (2020). Human HMGN1 and HMGN2 are not required for
transcription-coupled DNA repair. *Scientific reports*, *10*(1), 4332.
https://doi.org/10.1038/s41598-020-61243-4
25. Ding, H. F., Bustin, M., & Hansen, U. (1997). Alleviation of histone H1-mediated
transcriptional repression and chromatin compaction by the acidic activation region in
chromosomal protein HMG-14. *Molecular and cellular biology*, *17*(10), 5843–5855.
https://doi.org/10.1128/mcb.17.10.5843
26. Catez, F., Yang, H., Tracey, K. J., Reeves, R., Misteli, T., & Bustin, M. (2004). Network
of dynamic interactions between histone H1 and high-mobility-group proteins in
chromatin. *Molecular and cellular biology*, *24*(10), 4321–4328.
https://doi.org/10.1128/mcb.24.10.4321-4328.2004
27. Mowery, C. T., Reyes, J. M., Cabal-Hierro, L., Higby, K. J., Karlin, K. L., Wang, J. H.,
Kimmerling, R. J., Cejas, P., Lim, K., Li, H., Furusawa, T., Long, H. W., Pellman, D.,
Chapuy, B., Bustin, M., Manalis, S. R., Westbrook, T. F., Lin, C. Y., & Lane, A. A.
(2018). Trisomy of a Down Syndrome Critical Region Globally Amplifies Transcription
via HMGN1 Overexpression. *Cell reports*, *25*(7), 1898–1911.e5.
https://doi.org/10.1016/j.celrep.2018.10.061
28. He, J., Liu, C., Wang, B., Li, N., Zuo, G., & Gao, D. (2015). HMGN5 blockade by
siRNA enhances apoptosis, suppresses invasion and increases chemosensitivity to

temozolomide in meningiomas. *International journal of oncology*, *47*(4), 1503–1511. https://doi.org/10.3892/ijo.2015.3131

29. Subramanian, M., Gonzalez, R. W., Patil, H., Ueda, T., Lim, J. H., Kraemer, K. H., Bustin, M., & Bergel, M. (2009). The nucleosome-binding protein HMGN2 modulates global genome repair. *The FEBS journal*, *276*(22), 6646–6657. https://doi.org/10.1111/j.1742-4658.2009.07375.x

30. Martens, J. H., Brinkman, A. B., Simmer, F., Francoijs, K. J., Nebbioso, A., Ferrara, F., Altucci, L., & Stunnenberg, H. G. (2010). PML-RARalpha/RXR Alters the Epigenetic Landscape in Acute Promyelocytic Leukemia. *Cancer cell*, *17*(2), 173–185. https://doi.org/10.1016/j.ccr.2009.12.042

31. Lu, C., Han, H. D., Mangala, L. S., Ali-Fehmi, R., Newton, C. S., Ozbun, L., Armaiz-Pena, G. N., Hu, W., Stone, R. L., Munkarah, A., Ravoori, M. K., Shahzad, M. M., Lee, J. W., Mora, E., Langley, R. R., Carroll, A. R., Matsuo, K., Spannuth, W. A., Schmandt, R., Jennings, N. B., … Sood, A. K. (2010). Regulation of tumor angiogenesis by EZH2. *Cancer cell*, *18*(2), 185–197. https://doi.org/10.1016/j.ccr.2010.06.016

32. McCabe, M. T., Graves, A. P., Ganji, G., Diaz, E., Halsey, W. S., Jiang, Y., Smitheman, K. N., Ott, H. M., Pappalardi, M. B., Allen, K. E., Chen, S. B., Della Pietra, A., 3rd, Dul, E., Hughes, A. M., Gilbert, S. A., Thrall, S. H., Tummino, P. J., Kruger, R. G., Brandt, M., Schwartz, B., … Creasy, C. L. (2012). Mutation of A677 in histone methyltransferase EZH2 in human B-cell lymphoma promotes hypertrimethylation of histone H3 on lysine 27 (H3K27). *Proceedings of the National Academy of Sciences of the United States of America*, *109*(8), 2989–2994. https://doi.org/10.1073/pnas.1116418109

33. Hübner, J. M., Müller, T., Papageorgiou, D. N., Mauermann, M., Krijgsveld, J., Russell, R. B., Ellison, D. W., Pfister, S. M., Pajtler, K. W., & Kool, M. (2019). EZHIP/CXorf67 mimics K27M mutated oncohistones and functions as an intrinsic inhibitor of PRC2 function in aggressive posterior fossa ependymoma. *Neuro-oncology*, *21*(7), 878–889. https://doi.org/10.1093/neuonc/noz058

34. Pasini, D., Bracken, A. P., Jensen, M. R., Lazzerini Denchi, E., & Helin, K. (2004). Suz12 is essential for mouse development and for EZH2 histone methyltransferase activity. *The EMBO journal*, *23*(20), 4061–4071. https://doi.org/10.1038/sj.emboj.7600402

35. Gerlitz G. (2010). HMGNs, DNA repair and cancer. *Biochimica et biophysica acta*, *1799*(1-2), 80–85. https://doi.org/10.1016/j.bbagrm.2009.10.007

36. Lange, S. S., Mitchell, D. L., & Vasquez, K. M. (2008). High mobility group protein B1 enhances DNA repair and chromatin modification after DNA damage. *Proceedings of the National Academy of Sciences of the United States of America*, *105*(30), 10320–10325. https://doi.org/10.1073/pnas.0803181105

37. Lange, S. S., & Vasquez, K. M. (2009). HMGB1: the jack-of-all-trades protein is a master DNA repair mechanic. *Molecular carcinogenesis*, *48*(7), 571–580. https://doi.org/10.1002/mc.20544

38. Birger, Y., West, K. L., Postnikov, Y. V., Lim, J. H., Furusawa, T., Wagner, J. P., Laufer, C. S., Kraemer, K. H., & Bustin, M. (2003). Chromosomal protein HMGN1 enhances the rate of DNA repair in chromatin. *The EMBO journal*, *22*(7), 1665–1675. https://doi.org/10.1093/emboj/cdg142

39. da Costa, R. M., Riou, L., Paquola, A., Menck, C. F., & Sarasin, A. (2005). Transcriptional profiles of unirradiated or UV-irradiated human cells expressing either

735   the cancer-prone XPB/CS allele or the noncancer-prone XPB/TTD allele. *Oncogene*,
736   *24*(8), 1359–1374. https://doi.org/10.1038/sj.onc.1208288

40. Deb, S., Xu, H., Tuynman, J. *et al.* RAD21 cohesin overexpression is a prognostic and predictive marker exacerbating poor prognosis in *KRAS* mutant colorectal carcinomas. *Br J Cancer* 110, 1606–1613 (2014). https://doi.org/10.1038/bjc.2014.3

41. Bordoni, M., Pansarasa, O., Dell'Orco, M., Crippa, V., Gagliardi, S., Sproviero, D., Bernuzzi, S., Diamanti, L., Ceroni, M., Tedeschi, G., Poletti, A., & Cereda, C. (2019). Nuclear Phospho-SOD1 Protects DNA from Oxidative Stress Damage in Amyotrophic Lateral Sclerosis. *Journal of clinical medicine*, *8*(5), 729. https://doi.org/10.3390/jcm8050729

42. Ng, J. M., Vermeulen, W., van der Horst, G. T., Bergink, S., Sugasawa, K., Vrieling, H., & Hoeijmakers, J. H. (2003). A novel regulation mechanism of DNA repair by damage-induced and RAD23-dependent stabilization of xeroderma pigmentosum group C protein. *Genes & development*, *17*(13), 1630–1645. https://doi.org/10.1101/gad.260003

43. Brace, L. E., Vose, S. C., Stanya, K., Gathungu, R. M., Marur, V. R., Longchamp, A., Treviño-Villarreal, H., Mejia, P., Vargas, D., Inouye, K., Bronson, R. T., Lee, C. H., Neilan, E., Kristal, B. S., & Mitchell, J. R. (2016). Increased oxidative phosphorylation in response to acute and chronic DNA damage. *NPJ aging and mechanisms of disease*, *2*, 16022. https://doi.org/10.1038/npjamd.2016.22

44. Nemoz, C., Ropars, V., Frit, P., Gontier, A., Drevet, P., Yu, J., Guerois, R., Pitois, A., Comte, A., Delteil, C., Barboule, N., Legrand, P., Baconnais, S., Yin, Y., Tadi, S., Barbet-Massin, E., Berger, I., Le Cam, E., Modesti, M., Rothenberg, E., … Charbonnier, J. B. (2018). XLF and APLF bind Ku80 at two remote sites to ensure DNA repair by non-homologous end joining. *Nature structural & molecular biology*, *25*(10), 971–980. https://doi.org/10.1038/s41594-018-0133-6

45. Boehm, E. M., Gildenberg, M. S., & Washington, M. T. (2016). The Many Roles of PCNA in Eukaryotic DNA Replication. The Enzymes, 39, 231–254. https://doi.org/10.1016/bs.enz.2016.03.003

46. Song, M., Wang, Y., Zhang, Z., & Wang, S. (2017). PSMC2 is up-regulated in osteosarcoma and regulates osteosarcoma cell proliferation, apoptosis and migration. *Oncotarget*, *8*(1), 933–953. https://doi.org/10.18632/oncotarget.13511

47. Pujana, M. A., Han, J. D., Starita, L. M., Stevens, K. N., Tewari, M., Ahn, J. S., Rennert, G., Moreno, V., Kirchhoff, T., Gold, B., Assmann, V., Elshamy, W. M., Rual, J. F., Levine, D., Rozek, L. S., Gelman, R. S., Gunsalus, K. C., Greenberg, R. A., Sobhian, B., Bertin, N., … Vidal, M. (2007). Network modeling links breast cancer susceptibility and centrosome dysfunction. *Nature genetics*, *39*(11), 1338–1349. https://doi.org/10.1038/ng.2007.2

48. Yang, S., Zhou, L., Reilly, P. T., Shen, S. M., He, P., Zhu, X. N., Li, C. X., Wang, L. S., Mak, T. W., Chen, G. Q., & Yu, Y. (2016). ANP32B deficiency impairs proliferation and suppresses tumor progression by regulating AKT phosphorylation. *Cell death & disease*, *7*(2), e2082. https://doi.org/10.1038/cddis.2016.8

49. Redza-Dutordoir, M., & Averill-Bates, D. A. (2016). Activation of apoptosis signalling pathways by reactive oxygen species. *Biochimica et biophysica acta*, *1863*(12), 2977–2992. https://doi.org/10.1016/j.bbamcr.2016.09.012

50. Rothkamm, K., & Löbrich, M. (2003). Evidence for a lack of DNA double-strand break repair in human cells exposed to very low x-ray doses. *Proceedings of the National*

Academy of Sciences of the United States of America, *100*(9), 5057–5062. https://doi.org/10.1073/pnas.0830918100

51. Marín, A., Martín, M., Liñán, O., Alvarenga, F., López, M., Fernández, L., Büchser, D., & Cerezo, L. (2014). Bystander effects and radiotherapy. *Reports of practical oncology and radiotherapy : journal of Greatpoland Cancer Center in Poznan and Polish Society of Radiation Oncology*, *20*(1), 12–21. https://doi.org/10.1016/j.rpor.2014.08.004

52. Cadet, J., & Wagner, J. R. (2013). DNA base damage by reactive oxygen species, oxidizing agents, and UV radiation. *Cold Spring Harbor perspectives in biology*, *5*(2), a012559. https://doi.org/10.1101/cshperspect.a012559

53. Kennedy A. R. (2014). Biological Effects of Space Radiation and Development of Effective Countermeasures. *Life sciences in space research*, *1*, 10–43. https://doi.org/10.1016/j.lssr.2014.02.004

54. Thannickal, V. J., & Fanburg, B. L. (2000). Reactive oxygen species in cell signaling. *American journal of physiology. Lung cellular and molecular physiology*, *279*(6), L1005–L1028. https://doi.org/10.1152/ajplung.2000.279.6.L1005

55. Stier, A., Schull, Q., Bize, P. *et al.* Oxidative stress and mitochondrial responses to stress exposure suggest that king penguins are naturally equipped to resist stress. *Sci Rep* 9, 8545 (2019). https://doi.org/10.1038/s41598-019-44990-x

56. Ock, C. Y., Kim, E. H., Choi, D. J., Lee, H. J., Hahm, K. B., & Chung, M. H. (2012). 8-Hydroxydeoxyguanosine: not mere biomarker for oxidative stress, but remedy for oxidative stress-implicated gastrointestinal diseases. *World journal of gastroenterology*, *18*(4), 302–308. https://doi.org/10.3748/wjg.v18.i4.302

57. Lee, S. H., Meng, X. W., Flatten, K. S., Loegering, D. A., & Kaufmann, S. H. (2013). Phosphatidylserine exposure during apoptosis reflects bidirectional trafficking between plasma membrane and cytoplasm. *Cell death and differentiation*, *20*(1), 64–76. https://doi.org/10.1038/cdd.2012.93

58. Wang, S., El-Deiry, W. TRAIL and apoptosis induction by TNF-family death receptors. *Oncogene* 22, 8628–8633 (2003). https://doi.org/10.1038/sj.onc.1207232

59. Duez, P., Dehon, G., Kumps, A., & Dubois, J. (2003). Statistics of the Comet assay: a key to discriminate between genotoxic effects. *Mutagenesis*, *18*(2), 159–166. https://doi.org/10.1093/mutage/18.2.159

60. Elmore S. (2007). Apoptosis: a review of programmed cell death. *Toxicologic pathology*, *35*(4), 495–516. https://doi.org/10.1080/01926230701320337

61. Bowden, R. D., Buckwalter, M. R., McBride, J. F., Johnson, D. A., Murray, B. K., & O'Neill, K. L. (2003). Tail profile: a more accurate system for analyzing DNA damage using the Comet assay. *Mutation research*, *537*(1), 1–9. https://doi.org/10.1016/s1383-5718(03)00056-1

62. Collins, A. R., Dobson, V. L., Dusinská, M., Kennedy, G., & Stětina, R. (1997). The comet assay: what can it really tell us?. *Mutation research*, *375*(2), 183–193. https://doi.org/10.1016/s0027-5107(97)00013-4

63. Mair, P., Wilcox, R. Robust statistical methods in R using the WRS2 package. *Behav Res* **52,** 464–488 (2020). https://doi.org/10.3758/s13428-019-01246-w

64. Korotkevich G, Sukhov V, Sergushichev A (2019). "Fast gene set enrichment analysis." *bioRxiv*. doi: 10.1101/060012, http://biorxiv.org/content/early/2016/06/20/060012.

65. Lin, Y., Boone, M., Meuris, L. *et al.* Genome dynamics of the human embryonic kidney
293 lineage in response to cell biology manipulations. *Nat Commun* **5,** 4767 (2014).
https://doi.org/10.1038/ncomms5767
66. Purbey, P. K., Scumpia, P. O., Kim, P. J., Tong, A. J., Iwamoto, K. S., McBride, W. H.,
& Smale, S. T. (2017). Defined Sensing Mechanisms and Signaling Pathways Contribute
to the Global Inflammatory Gene Expression Output Elicited by Ionizing
Radiation. *Immunity*, *47*(3), 421–434.e3. https://doi.org/10.1016/j.immuni.2017.08.017
67. Ma, H., Rao, L., Wang, H. *et al.* Transcriptome analysis of glioma cells for the dynamic
response to γ-irradiation and dual regulation of apoptosis genes: a new insight into
radiotherapy for glioblastomas. *Cell Death Dis* **4,** e895 (2013).
https://doi.org/10.1038/cddis.2013.412
68. Hurov, K. E., Cotta-Ramusino, C., & Elledge, S. J. (2010). A genetic screen identifies the
Triple T complex required for DNA damage signaling and ATM and ATR
stability. *Genes & development*, *24*(17), 1939–1950. https://doi.org/10.1101/gad.1934210
69. Samuni, A. M., Kasid, U., Chuang, E. Y., Suy, S., Degraff, W., Krishna, M. C., Russo,
A., & Mitchell, J. B. (2005). Effects of hypoxia on radiation-responsive stress-activated
protein kinase, p53, and caspase 3 signals in TK6 human lymphoblastoid cells. *Cancer
research*, *65*(2), 579–586.
70. Wang, H., Jiang, H., Van De Gucht, M., & De Ridder, M. (2019). Hypoxic
Radioresistance: Can ROS Be the Key to Overcome It?. *Cancers*, *11*(1), 112.
https://doi.org/10.3390/cancers11010112
71. Contiliani, D. F., de Araújo Ribeiro, Y., de Moraes, V. N., & Pereira, T. C.
(2020). *Panagrolaimus superbus* tolerates hypoxia within Gallium metal cage:
implications for the understanding of the phenomenon of anhydrobiosis. *Journal of
nematology*, *52*, 1–6. https://doi.org/10.21307/jofnem-2020-046
72. Wang, C., Grohme, M. A., Mali, B., Schill, R. O., & Frohme, M. (2014). Towards
decrypting cryptobiosis--analyzing anhydrobiosis in the tardigrade Milnesium
tardigradum using transcriptome sequencing. *PloS one*, *9*(3), e92663.
https://doi.org/10.1371/journal.pone.0092663
73. Vilas, C. K., Emery, L. E., Denchi, E. L., & Miller, K. M. (2018). Caught with One's
Zinc Fingers in the Genome Integrity Cookie Jar. *Trends in genetics : TIG*, *34*(4), 313–
325. https://doi.org/10.1016/j.tig.2017.12.011
74. Chen, E. Y., Tan, C. M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G. V., Clark, N. R., &
Ma'ayan, A. (2013). Enrichr: interactive and collaborative HTML5 gene list enrichment
analysis tool. *BMC bioinformatics*, *14*, 128. https://doi.org/10.1186/1471-2105-14-128
75. Kuleshov, M. V., Jones, M. R., Rouillard, A. D., Fernandez, N. F., Duan, Q., Wang, Z.,
Koplev, S., Jenkins, S. L., Jagodnik, K. M., Lachmann, A., McDermott, M. G., Monteiro,
C. D., Gundersen, G. W., & Ma'ayan, A. (2016). Enrichr: a comprehensive gene set
enrichment analysis web server 2016 update. *Nucleic acids research*, *44*(W1), W90–
W97. https://doi.org/10.1093/nar/gkw377
76. Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., & Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq
aligner. *Bioinformatics (Oxford, England)*, *29*(1), 15–21.
https://doi.org/10.1093/bioinformatics/bts635
77. Conesa, A., Madrigal, P., Tarazona, S., Gomez-Cabrero, D., Cervera, A., McPherson, A.,
Szcześniak, M. W., Gaffney, D. J., Elo, L. L., Zhang, X., & Mortazavi, A. (2016). A

survey of best practices for RNA-seq data analysis. *Genome biology*, *17*, 13.
https://doi.org/10.1186/s13059-016-0881-8

78. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., Durbin, R., & 1000 Genome Project Data Processing Subgroup (2009). The
Sequence Alignment/Map format and SAMtools. *Bioinformatics (Oxford,*
England), *25*(16), 2078–2079.

79. Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15,** 550 (2014).
https://doi.org/10.1186/s13059-014-0550-8

80. Liberzon, A., Birger, C., Thorvaldsdóttir, H., Ghandi, M., Mesirov, J. P., & Tamayo, P.
(2015). The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell*
systems, *1*(6), 417–425. https://doi.org/10.1016/j.cels.2015.12.004

81. Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M.
A., Paulovich, A., Pomeroy, S. L., Golub, T. R., Lander, E. S., & Mesirov, J. P. (2005).
Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide
expression profiles. *Proceedings of the National Academy of Sciences of the United*
States of America, *102*(43), 15545–15550. https://doi.org/10.1073/pnas.0506580102

82. Liao, Y., Wang, J., Jaehnig, E. J., Shi, Z., & Zhang, B. (2019). WebGestalt 2019: gene set
analysis toolkit with revamped UIs and APIs. *Nucleic acids research*, *47*(W1), W199–
W205. https://doi.org/10.1093/nar/gkz401

83. Zhang, B., Kirov, S., & Snoddy, J. (2005). WebGestalt: an integrated system for
exploring gene sets in various biological contexts. *Nucleic acids research*, *33*(Web
Server issue), W741–W748. https://doi.org/10.1093/nar/gki475

84. Schmidt, M. A., Meydan, C., Schmidt, C. M., Afshinnekoo, E., & Mason, C. E. (2020).
The NASA Twins Study: The Effect of One Year in Space on Long-Chain Fatty Acid
Desaturases and Elongases. *Lifestyle genomics*, *13*(3), 107–121.
https://doi.org/10.1159/000506769

85. MacKay M, Afshinnekoo E, Roboz G, Guzman M, Melnick A, Wu S, Mason CE. The
Therapeutic Landscape for Cells Engineered with Chimeric Antigen Receptors. *Nature*
Biotechnology. 2020 Jan;6(8):120-128.

908
Figure 1. Apoptosis Assay.
A) Apoptosis was measured by Phosphatidyl Serine exposure to luminescent annexin V fusion proteins (Annexin V-
LgBit and Annexin VsmBit). HEK293 Dsup and HEK293 controls were subjected to 1Gy, 2Gy, 4Gy, and 16Gy of
X-ray radiation and 100μM of H202 for the duration of the time points 1hr, 6hr, 24hrs, and 48hrs. At each time
point apoptosis was measured in terms of relative light units (RLUs). HEK293 control cells experience roughly
twice as much apoptotic signals than HEK293 DSUP up to 24hrs but then saturates at 48hrs. Anova analysis
followed by Tukeys Post Hoc test was used to determine p value significance, ns=p>0.05, p=0.5*, ** p<0.05, ***
p<0.01, **** p<0.001. Values represent average of triplicates and error bars represent standard error of the mean.
B) Apoptosis via caspase-3 activity was measured in HEK293, Empty vector, and Dsup expressing cells.
13.2uM of CPT was added to each cell line and Hoechst and Caspase staining was used to discriminate live
cells from apoptotic cells. Images were collected and FITC intensity was analyzed and plotted over time for
all wells.
C) Percentage of apoptotic cells at the end of the experiment was calculated at 22.5 hours. Error bars show 1
standard deviation of the apoptotic rate of the cells. A heteroscedastic two tailed student t-test was used to test
the null hypothesis that the empty vector and Dsup containing cells have equal rates of apoptosis (p = 0.005).
924

927
Fig... ...ratio...
A) ... ...ng of gene expression, princ... F...
dist... ...g (t-SNE) dimensionality re...
Pea... ...of gene expression, principal...
dist... ...g (t-SNE) dimensionality re...
Pea... ...of gene...
dist... ...g (t-S...
sho... ...UP vs...
clus... ...urs at 0...
can... ...-SNE...
rath...

939
Figure 3. HEK293...

A) Heatmap for HEK293 and Dsup cells across all radiation doses and time points. Gene expression changes were
seen across each time point and mostly corresponded to time and genotype. Reported here are genes expression
changes for any gene with a $q < 0.01$ and $logFC \geq 1$ in a multivariate model including mycoplasma correction, time,
radiation, and genotype. Orange indicates relative enrichment while purple indicates relative depletion. Differences
in gene clustering at 0Gys across all time points indicates gene expression differences for Dsup expressing cells at
baseline.
B) Volcano plot illustrating fold change log base 2 compared with adjusted p value (-log base 10) between HEK293
Dsup and HEK293 controls across all time points. Horizontal bar at $y=2$ represents a significance level of $p = 0.01$
significance level. Dsup is the most differentially and significantly expressed gene across all time points.
950

951
Figure 4. Shared pathways conserved across 0Gys at all timepoints.
ENCODE histone modifications were assessed using Enrichr with bars sorted by p-value rankings
A) The 10 genes that were found downregulated across all time points histone modification pathways were
analyzed. Gray bars are not significantly expressed and the top corresponding bar has a q-value of 1
B) The subset of the intersection of all genes upregulated at 0Gys was 148 genes and was analyzed for histone
modification pathways. Red bars indicate significantly expressed histone methylation pathways with the top bar
having an associated q-value of 7.941e-11.

959 C) Fgsea Running Enrichment Score plots corresponding to each time point at all doses of radiation for
960 downregulated pathways in Dsup expressing cells. Horizontal lines correspond to ranked leading edge subset. Fgsea
961 plots are ordered by p-adjusted values and correlated with density of genes of the curve.
962 D) Upregulated pathways across all time points for all doses of radiation.

963
Figure 5. Functional GSEA of Kegg pathways.
Genes were ranked by a p-value cutoff of .05 and Log$_2$FC cutoff of 1.5 was used to perform WebGesalt GSEA for
overall comparison of differentially expressed pathways in HEK293 Dsup vs. HEK293.
A) Biological process shows top enrichment of biological regulation, metabolic process, and response to stimulus.
B) Cellular component shows top enrichment of membrane, nucleus, and membrane enclosed lumen
C) Molecular function shows top enrichment of protein binding, ion binding, and nucleic acid binding.
D) Enrichment ratio. Displays top upregulated and top downregulated KEGG pathways for Dsup expressing human
cells compared to WT.

972
973 **Supplemental Figures**

974   A

976   B

977
978

Figure S1. Presence of HA-tagged DSUP.
A) Primary antibodies R&D Systems recombinant mouse HA Tag antibody (MAB0601) was diluted to 0.1ug/ml
and Anti Human Vinculin antibody (V284) was diluted to a working concentration of 1:500 dilution and used as a
housekeeping protein control. Invitrogen Goat anti-Rabbit IgG (H+L) Cross-Absorbed Secondary Antibody DyLight
800 (SAS-10036) and Rabbit anti-Mouse IgG (H+L) Cross-adsorbed Secondary Antibody Alexa Fluor Plus 680
(A32729) diluted to working concentrations of 1:5000 dilution and 0.1ug/ml respectively in 1% milk in 1X TBST.
The blot was imaged on a Li-Cor Odyssey® CLx imagine system using both 700nm and 800nm channels. Viniculin
was present in all samples at 110kDa and only HA tag at 43 kDa indicative of the MW of Dsup. Columns 1-3 are
HEK293, 4-6 are HEK293 Vector, and 7-9 are HEK293-Dsup expressing HA tag.
B) Boxplots shows presence of Dsup transcripts in HEK293-Dsup cells but not in control across all time points and
doses of radiation.

991
Figure S2. Identification of insertion site.
The left plot x-axis is the segment of the read that maps to the vector and the right plot x-axis is the segment of the
read that maps to the genome. The y-axis is the full length nanopore read. The first 500 bases of the nanopore read
map to the right arm of chromosome 5 while the remaining 1500 bases of the nanopore read map to the vector site
downstream of Dsup. 98% of reads were obtained for HEK293 parental cell line and HEK293 Dsup with a genome
coverage of ~2.8x and 1.1x respectively.
998
999
1000

1001
1002 **Figure S3. Experimental layout for RNAseq analysis.**
1003 HEK293 Dsup and Control cells were grown to 75% confluency on T-175 flasks before plated in triplicates on 6
1004 well plates. 1 day after seeding, cells from each set were irradiated with X-rays in a RS2000 Biological research
1005 irradiator. For 0Gy, each set was kept outside the incubator at the corresponding time frame it took to irradiate cells
1006 and were harvested along with irradiated cells at 1hr, 6hrs, and 24hrs, post irradiation.

1007
Figure S4. A) Heatmap for mycoplasma corrected HEK293 and DSUP cell lines across all radiation doses at 1hr
post radiation.
B) Volcano plot illustrating mycoplasma corrected gene fold change log base 2 compared with adjusted p value (-
log base 10) between HEK293 DSUP and HEK293 controls across all radiation doses at 1hr post radiation.
C) Heatmap for mycoplasma corrected HEK293 and DSUP cell lines across all radiation doses at 6hr post radiation.
D) Volcano plot illustrating mycoplasma corrected gene fold change log base 2 compared with adjusted p value (-
log base 10) between HEK293 DSUP and HEK293 controls across all radiation doses at 6hr post radiation.

E) Heatmap for mycoplasma corrected HEK293 and DSUP cell lines across all radiation doses at 24hr post
radiation.
F) Volcano plot illustrating mycoplasma corrected gene fold change log base 2 compared with adjusted p value (-log
base 10) between HEK293 DSUP and HEK293 controls across all radiation doses at 24hr post radiation.
1019
1020   A

1021
1022   B

1023
1024   **Figure S5. Differentially expressed pathways by introduction of Dsup at each timepoint.**

1025   (A) Fgsea Running Enrichment Score plots corresponding to each time point at all doses of radiation for
1026   downregulated pathways in Dsup expressing cells. Horizontal lines correspond to ranked leading edge subset. Fgsea
1027   plots are ordered by p.adjusted values and correlated with density of genes of the curve.
1028   (B) Upregulated pathways across all time points for all doses of radiation.
1029
1030

1031
1032   **Figure S6. Stable expression of HMGN1/2.** Fgsea Boxplots of TPM normalized gene expression of
1033   human homologous proteins HMGN1 and HMGN2 reveal relatively normal expression across time points for both
1034   HEK293 Dsup and HEK293 control cells.
1035