Integrative Transcriptomic Analysis Reveals the Immune Mechanism for A CyHV-3-Resistant Common Carp Strain

Zhiying Jia[1,2†], Nan Wu[3,4†], Xiaona Jiang[1], Heng Li[3,4], Jiaxin Sun[1], Mijuan Shi[3], Chitao
Li[1], Xuesong Hu[1], Yanlong Ge[1], Weidong Ye[3,4], Ying Tang[3], Yingyin Cheng[3], Xiao-Qin
Xia[3,4,5*], Lianyu Shi[1*]

[1]Heilongjiang River Fisheries Research Institute, Chinese Academy of Fishery
Sciences, Harbin, China

[2]Key Laboratory of Aquatic Genomics, Ministry of Agriculture, Chinese Academy of
Fishery Sciences, Beijing, China

[3]Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, China

[4]University of Chinese Academy of Sciences, Beijing, China

[5]The Innovative Academy of Seed Design, Chinese Academy of Sciences, Beijing,
China

[†] Co-first author.

* Correspondence:
Lianyu Shi
sly2552@aliyun.com

Xiao-Qin Xia
xqxia@ihb.ac.cn

Word count: 5652

Number of figures: 7

Keywords: transcriptome, lncRNA, CyHV-3, resistant immune mechanism, common
carp
22

ABSTRACT
Anti-disease breeding is becoming the most promising solution to cyprinid
herpesvirus-3 (CyHV-3) infection, the major threat to common carp aquaculture.
Mortality studies suggested that a breeding strain of common carp is resistant to this
disease. This study illustrates the immune mechanisms involved in anti-CyHV-3
ability. An integrative analysis of the protein-coding genes and long non-coding
RNAs (lncRNAs) using transcriptomic data was also performed. Tissues from the
head kidney of common carp were extracted at day 0 (the healthy control) and day 7
after CyHV-3 infection (the survivors), and used to analyze the transcriptome through
both Illumina and Pac-bio sequencing. Following analysis of the Kyoto Encyclopedia
of Genes and Genomes pathways and Gene Ontology terms involved, the immune-
related genes were merged. These genes were filtered using the current common carp
immune gene library, and information on the immune process was detailed. Immune
gene categories and their corresponding genes in different comparison groups were
revealed. Also, the immunological Gene Ontology terms for lncRNA modulation were
retained. The weighted gene co-expression network analysis was used for the
regulation of immune genes lncRNA. The results demonstrated that the breeding carp
strain develops marked resistance to CyHV-3 through a specific innate immune
mechanism. The featured biological processes were autophagy, phagocytosis,
cytotoxicity, and virus blockage by lectins and mucin 3 (MUC3). Moreover, the
immune suppressive signals, such as suppression of interleukin 21 receptor (IL21R)
on STAT3, PI3K mediated the inhibition of inflammation by dopamine upon
infection, as well as the inhibition of NLR family CARD domain containing 3
(NLRC3) on STING during a steady state. Possible susceptible factors for CyHV-3,
such as integrin beta 1 (ITGB1), toll-like receptor 18 (TLR18), and C-C motif
chemokine ligand 4 (CCL4), were also revealed from the common strain. The results
of this study suggested that the regulation of galectin 3 (LGALS3) and T cell
leukemia homeobox 3 (TLX3) by lncRNA may play a role in the resistance
mechanism. Therefore, immune factors that are immunogenetically insensitive or
susceptible to CyHV-3 infection have been revealed.

INTRODUCTION
Cyprinid herpesvirus-3 (CyHV-3) infection is a major threat to common carp
aquaculture (1), leading to widespread mortality and substantial economic loss.
CyHV-3 is thought to cause death by weakening the host's immune system, resulting
in susceptibility to pathogenic microbes (1). In common carp, clinical signs of the
disease develop rapidly and may induce morbidity and mortality within a period of 6
to 10 days following infection (2).
61
Carp that survive a primary infection with CyHV-3 can be resistant to future infection
with this virus. This immunity may be genetically passed onto offspring. Since

latency and life-long infections with CyHV-3 exist in carp (3, 4), genetic backgrounds
are crucial in developing an understanding resistance against the virus. Experimental
infections of carp from pure lines or crosses have indicated the existence of genetic
background of resistance by divergent survival rates (2). For example, a markedly
higher expression of immune-related genes involved in pathogen recognition,
complement activation, major histocompatibility complex class I (MHC I)-restricted
antigen presentation, and the development of adaptive mucosal immunity was noted
in the more resistant R3 line. Higher activation of CD8$^+$ T cells was also observed (5).
Additionally, carp strains of Asian origin, particularly Amur wild carp, currently being
treated as *Cyprinus rubrofuscus*, were shown to be more resistant to CyHV-3 than
strains originating from Europe and belonging to *Cyprinus carpio*, such as the Prerov
scale carp or koi carp from a breed in the Czech Republic (6).

The immune response of carp to CyHV-3 involves both innate and adaptive aspects
with the outcome of the disease largely depending on whether the balance is tipped in
favor of the host's immune response or virus's evasion strategy (2). In general,
transcriptomic analysis has revealed that three immune-related pathways, namely the
mitogen-activated protein kinase (MAPK) signaling pathway, the innate immune
response, and the cytokine-mediated signaling pathway, were highly involved in the
infection with CyHV-3 (7). In red common carp × koi, the expression of interleukin
12 (IL12) p35, interferon (IFN) αβ, and toll-like receptor 9 (TLR9) may provide
potential genes related to resistance against KHV (another term for CyHV-3) (8).
However, the magnitude of type I IFN response did not correlate with a higher
resistance in CyHV-3-infected carp, during the challenge test among different strains,
although CyHV-3 infection can induce type I IFNs (6, 9). Regarding the innate
resistance in carp, the mapped CyHV-3 survival quantitative trait loci have been
reported mainly in *IL10* and *MHC II* (10). Recently, by quantitative trait locus
mapping and genome-wide association study, *tumor necrosis factor-alpha* (*TNFA*),
hypoxia inducible factor 1 subunit alpha (*HIF1A*), *galectin-8* (*LGALS8*), *rootletin*,
and *paladin*, have also been related to resistance against CyHV-3 (11). Adaptive
immunity through both cytotoxicity and immunoglobulin (Ig) secretion may be
involved in resistance. Matthew et al. revealed that, in the anterior kidney, Ig secretion
plays an important role in the resistance during the persistent infection or reactivation
phases of CyHV-3 (1). In addition, CyHV-3 profoundly influences the expression of
host miRNA, although the regulation of immune processes by miRNA in the clinical
and latent phases differs (4). This suggests an important role of non-coding RNAs in
anti-CyHV-3 immunity.

Based on our previous study, a strain of common carp has already shown a higher
survival rate after breeding for four generations. In addition, it has displayed a better
immunological index as well as a reduced virus load in immune organs, such as the
kidneys and spleen (12, 13). However, there is a lack of systemic studies focusing on

a detailed network of the immune system for the anti-CyHV-3 immune mechanism in
this resistant strain. The third-generation sequencing method, such as Pac-bio, can
decode the genetic sequences that are markedly longer compared with the second-
generation method, such as Illumina (14). Therefore, a higher quality assembly of
transcripts, including both mRNA and non-coding RNAs, may provide a more
detailed understanding of the mechanism of genetic resistance.
112
In this study, integrative transcriptomic analysis was performed for both mRNA and
long non-coding RNA (lncRNA) during the CyHV-3 challenge. Immune-related
transcripts from survivors and healthy controls, from either resistant or normal strains,
were analyzed using the strategy shown in Figure 1. By comparing data from different
groups for different strains obtained through both Illumina (New England Biolabs,
Ipswich, MA, USA) and Pacific Biosciences (PacBio) sequencing, both immune-
related differentially expressed genes (DEG) and lncRNA have been revealed for the
Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways and/or Gene
Ontology (GO) terms involved. The immune gene-related lncRNA was also explored.
This study may shed some light on fish breeding for virus-resistant strains, and
facilitate disease control after CyHV-3 infection.

MATERIALS AND METHODS
Ethics Statement
All procedures involving animals in this study were conducted according to the
guidelines for the care and use of laboratory animals of Heilongjiang River Fisheries
Research Institute, Chinese Academy of Fishery Sciences (Harbin, China). The
studies involving animals were reviewed and approved by the Committee for the
Welfare and Ethics of Laboratory Animals of Heilongjiang River Fisheries Research
Institute, Chinese Academy of Fishery Sciences.
132
Animals and Virus Challenge
The fish used in this study were obtained from the Kuandian Research Base of
Heilongjiang River Research Institute (Liaoning, China). The breeding resistant strain
was the fourth generation after the selection of resistance to CyHV-3 (13). For the
common carp involved in this study, both the CyHV-3-resistant strain and common
strain were used for the virus challenge. Since CyHV-3 has a mucosal route of
infection mainly via the skin rather than the gut (2), this study induced infection with
CyHV-3 by adding the homogenate solution of internal organs from sick fish into the
water of the tanks. To separate the control and experimental groups from either the
resistant or common strains, four tanks (1.6 m × 1.2 m × 0.6 m) containing healthy
juvenile common carp (~70 g; N=300 fish per tank), were used. The water
temperature during the experiment was maintained at $25\pm1^{\circ}$C.
145
Sampling

A selection of common carp, from either the resistant or normal strains, were
sacrificed. Head kidney samples were obtained at days 0 and 7 after challenge,
representing the control and survivor carp. The samples (N=3) were collected
immediately and soaked in 10 volumes of RNAlater (Qiagen, Hilden, Germany), for
sequencing using Illumina (New England Biolabs). To better sequence and generate
the lncRNAs, PacBio sequencing was applied to analyze the mix of liver, spleen, and
kidney. The number of dead fish during the experiment was counted daily to calculate
the mortality rate.
155

RNA Extraction
RNA was isolated using the AllPrep DNA/RNA FFPE Kit (Qiagen, Hilden,
Germany), according to the instructions provided by the manufacturer. RNA
degradation and contamination were monitored on 1% agarose gels. RNA purity was
checked using the NanoPhotometer spectrophotometer (Implen Inc., Westlake Village,
CA, USA). RNA concentration was measured using the Qubit RNA Assay Kit in a
Qubit 2.0 Flurometer (Life Technologies, Carlsbad, CA, USA). RNA integrity was
assessed using the RNA Nano 6000 Assay Kit of the Agilent Bioanalyzer 2100 system
(Agilent Technologies, Santa Clara, CA, USA).
165

Library Preparation and Sequencing for Transcriptomic Analysis
Optimized RNA-Seq strategies, including both PacBio and Illumina (New England
Biolabs) sequencing (15), were used to more precisely resolve the sequence of
transcripts. Firstly, the total RNA isolated from the head kidney was used to construct
the cDNA library. Subsequently, the library was sequenced on a PacBio RS II
platform (Biomarker Technologies, Beijing, China). For the PacBio Long Read
Processing, raw reads were processed into error-corrected reads of insert using Iso-seq
pipeline with minFullPass = 0 and minPredictedAccuracy = 0.90. Next, full-length,
non-chemiric transcripts were determined by searching for the polyA tail signal and
the 5' and 3' cDNA primers in reads of insert. Iterative Clustering for Error Correction
was used to obtain consensus isoforms, and full-length consensus sequences from
Iterative Clustering for Error Correction were polished using Quiver. High-quality
transcripts with post-correction accuracy of >99% were retained for further analysis.
179

The illustration of transcripts obtained from the results of PacBio could provide
reference transcriptional sequences for the assembly of Illumine sequencing data, to
improve sequencing quality. Therefore, Illumina (New England Biolabs) sequencing
was performed on all head kidney samples. The procedure used for the preparation of
the gene library and sequencing of the transcriptome followed previously published
methods (16). Briefly, sequencing libraries were generated using the NEBNext
UltraTM RNA Library Prep Kit for Illumina (New England Biolabs), and the library
quality was assessed on the Agilent Bioanalyzer 2100 system. The library
preparations were sequenced on an Illumina platform, and 150 bp paired-end reads

were generated.

Annotation and Functional Analysis of Transcripts by Public Databases

All reads in the transcriptome data were mapped on the common carp genome

(https://asia.ensembl.org/Cyprinus_carpio_german_mirror/Info/Index) (17) for

annotation. The data from the Illumina platform were also used to check and replace

errors in the data from the PacBio platform. Both the annotation of genes and

lncRNAs were subsequently generated. The revealed common carp DEGs were

annotated by GO terms and KEGG pathways, following a previously published

protocol (16, 18). In brief, functional annotation and the classification of genes were

determined by both employing local genes blasts against GO Consortium

(http://geneontology.org/), Blast2GO (Bioinformatics Department, Prince Felipe

Research Institute, Valencia, Spain), and KEGG

(https://www.kegg.jp/kegg/pathway.html). Enrichment of the KEGG pathways was

carried out for differently expressed immune genes, including both upregulated and

downregulated genes for all comparison groups. For this purpose, the gene list of the

current construct common carp immune gene library was used as follows. For the

comparison groups, DEGs from intra-stain comparison groups A (survivors vs.

controls in the resistant strain) and B (survivors vs. controls in the normal strain), as

well as inter-strain comparison groups C (the comparison between survivors at day 7

for the two strains) and E (the comparison between control fish at day 0 for the two

strains) were used. Vital genes for survival were investigated through group D, in

which the genes without significant differences ($P > 0.05$) between the survivors from

the resistant strain and the normal strain were compared with the transcripts from

dying fish. LncRNAs were also annotated by GO terms.

Construction of the Common Carp Immune Gene Library

The common carp immune library was constructed by following our previously

published method, which was applied to grass carp (16) and tilapia (18), with some

adjustment for viral infection-related immune genes. The modifications were based on

gene information obtained by blasting each sequence to databases, including NCBI

NR database, as well as Swiss-Prot and Pfam databases. The common carp immune

gene library contained information for immune genes at two levels. For the first level,

nine categories of immune processes, namely "acute phase reactions", "pattern

recognition", "antigen processing and regulators", "complement system",

"inflammatory cytokines and receptors", "adapters, effectors and signal transducers",

"innate immune cells related", "T/B cell antigen activation", and "other genes related

to immune response", were proposed. Subsequently, many categories of immune

genes for each immune process (detailed in Table S1) were applied for the second

level. The library was used to filter transcriptome data and obtain details of the

immune processes and particular immune genes for each comparison group, during

the GO term and KEGG pathway enrichment.

Statistics Analysis

The DEG from intra-strain comparison groups A and B, as well as inter-strain

comparison groups C and E were generated by comparing the RPKM (Reads Per

Kilobase of transcript, per Million mapped reads) for each gene during a steady or

inflammatory state, using the DESeq2 R package (1.16.1). To reveal the basic genes

for the fish to survive from CyHV-3 infection, the mRNAs of non-significant

difference ($P > 0.05$) between the survivors of both the resistant and normal strains,

were compared using transcripts from the dying fish, consequently obtaining the DEG

in group D. The resulting P-values were adjusted using the Benjamini and Hochberg

approach for controlling the false discovery rate. Genes with an adjusted $P < 0.05$

found by DESeq2 were assigned as being differentially expressed. The following

bioinformatics analysis was performed to select immune-related transcripts from the

common carp immune gene library, and construct the barplots for the major immune

processes and immune categories. The t-test was used to assess differences, with a

false discovery rate adjusted $P < 0.05$. Qualitative comparisons were performed

between samples by counting the number of DEG. The data were rearranged in

Microsoft EXCEL, and applied to plot charts by ggplot2 (2.2.1) using R language.

Correlation Analysis between LncRNA and Genes Involved in Immune-related

GO

The weighted gene co-expression network analysis (WGCNA) was performed using

the R package "WGCNA" (19). Specifically, all genes with an expression variance

ranked in the top 75 percentile of the data set were retained (20). The R package

WGCNA was used to construct the weighted gene co-expression network (21). A

matrix of signed Pearson correlations between all gene pairs was computed, and the

transformed matrix was used as input for linkage hierarchical clustering. Genes with

similar expression patterns were clustered together as one module. Subsequently,

using R package clusterProfiler (22), enriched GO terms for lncRNA-related protein-

coding genes were generated for the gene list of every module. The immune-related

GO terms were selected, and the immune genes, as well as related lncRNAs, were

classified into different comparison groups, to elucidate the detail lncRNA-mRNA

network for the differential anti-disease mechanism involved in the two analyzed

strains.

Validation through Reverse Transcription-Polymerase Chain Reaction (RT-PCR)

The mRNA samples used for transcriptome sequencing were also subjected to real-

time quantitative RT-PCR validation (n = 3), following a previously published

protocol [14, 16]. After RNA isolation, reverse transcription and quantitative PCR

(qPCR) experiments were carried out for 11 genes. Gene expression levels were

determined using TaqMan-based qPCR (Thermo Fisher Scientific, Waltham, MA,

USA), which was performed on the RotorGene system (Qiagen, Hilden, Germany).

The total RNA was reverse transcribed, and 60 ng were used for the One Taq RT PCR
Kit (New England Biolabs) according to the instructions provided by the
manufacturer for mRNA quantification and reverse transcription (Thermo Fisher
Scientific, Waltham, MA, USA). For mRNA quantification, the resulting cDNA was
pre-amplified using a TaqMan PreAmp Master Mix Kit (Thermo Fisher Scientific,
Waltham, MA, USA). The relative levels of mRNA were determined from diluted
cDNA and pre-amplified cDNA using qPCR and appropriate TaqMan probes. The
expression levels were calculated after an efficiency correction (based on the results
of a serial dilution) relative to those of endogenous controls.

RESULTS

Mortality Rate and Pathological Appearance
The number of dead fish was recorded from day 1 to day 15 after challenge (Table
S1). The mortality rate was calculated for all groups (Table S2). For the challenged
normal strain (Figure 2A), the mortality rate increased daily, except for a decrease
observed on day 5. Mortality peaked at day 7 (69%) and decreased to a markedly low
level thereafter. For the challenged resistant strain (Figure 2A), the mortality rate
remained at a markedly low level, with the highest value (11%) recorded at day 7. The
mortality rates of the two challenged groups were significantly different ($P = 0.000$).
In the two unchallenged controls, there were no dead fish. Additionally, compared
with the organs in survivors of the resistant strain, pathological swelling and
hyperemia of the internal organs (particularly the spleen) was obvious in survivors
from the normal strain on day 7 (Figure 2B).
295

Quality of Transcriptomic Data
Sequencing was performed with both the PacBio RS II and Illumina platforms to
analyze the gene information of the common carp. Through PacBio sequencing, 19.49
G data were obtained. Of the 258,346 ccs, 79.79% were full-length sequences, and
17,769 polished high-quality isoforms were also revealed (Table S3, sheet 1).
Meanwhile, the Illumina data were also characterized by high quality, and the
information for each sample is provided in Table S3 (sheet 2).
303

304 **Validation of Transcripts**
305 A total of 24 reactions were performed to validate the transcripts of 11 genes, most of
306 which were immune genes. The correlation analysis showed that fold changes
307 between transcriptome and qPCR results correlated well (R^2=0.929718564). The gene
308 ID, annotation, primers, and fold-change information are listed in Table S4.
309

GO Analysis of Differentially Expressed Transcripts among Tested Fish
The heatmap of all the tested head kidney samples (Figure S1) revealed significant
differences across samples, including between the normal and resistant strains. Cluster
analysis of the expression levels of immune genes demonstrated that survivors of the

normal strain had significantly differences versus those of the resistant strain.
According to the enriched GO terms (Figures 3 and S2), the immune-related
biological processes were "immune system process" in comparison of A–E and "cell
killing" in comparison between A and C; the immune-related cellular component was
"virion". In comparisons between B and C, the immune-related molecular function
was "antioxidant activity". In comparisons between A, B and C, and "chemoattractant
activity" in comparisons between A and B.
321
KEGG Analysis of Differentially Expressed Transcripts among Tested Fish
After filtering was conducted using the gene list of the common carp immune gene
library (Table S1), the immune-related pathways appeared more clearly (Figure 4).
Compared with group A, there were markedly more genes involved in immune-related
pathways in groups B and C. The immune-related pathways found in different
comparison groups were "phagosome", "regulation of autophagy", "ubiquitin
mediated proteolysis", and "plant-pathogen interaction" in group A (Figure 4A);
"endocytosis", "phagosome", "FoxO signaling pathway", "ubiquitin mediated
proteolysis", "DNA replication", "fructose and mannose metabolism", "oxidative
phosphorylation", and "plant-pathogen interaction" in group B (Figure 4B); and
"endocytosis", "phagosome", "ubiquitin mediated proteolysis", "DNA replication",
"fructose and mannose metabolism", "oxidative phosphorylation", and "plant-
pathogen interaction" in group C (Figure 4C). Since the comparison of C highlighted
the differences between both the survivors involved in comparisons A and B, a bubble
chart was used to further clarify both the enrichment factors and gene numbers for the
related pathways. The results (Figure 4D) showed that the most enriched immune-
related pathways were "p53 signaling pathway", "NOD-like receptor signaling
pathway", "lysosome", "oxidative phosphorylation", and "proteasome". Of note, the
pathways with the greatest number were "phagosome", "proteasome", and "oxidative
phosphorylation". Besides, some immune-related metabolic pathways, such as
"glycolysis/gluconeogenesis" and "fructose and mannose metabolism", were also
revealed through filtering with the gene list of the common carp immune gene library
in the KEGG pathway analysis for the comparison C (Figure 4D). Additionally, the
pathway "dopaminergic synapse" (Figure S3) may indicate alteration of the
dopaminergic signal during viral infection.
347
Classification of Immune-related DEG from Different Comparison Groups into
Immune Process and Immune Gene Category
The current construction of the common carp immune gene library (Table S4) was
used to classify the revealed immune transcripts and refine the involved immune
processes and immune gene categories. To illustrate the immune process, the
information for immune gene categories and the immune-related DEG in all
comparisons were listed in Table 1, with the details for comparison groups A–E
provided in Tables S5–S9, respectively. In group A (Figure 5A and Table S5), most

immune mRNAs were upregulated in the immune processes of "pattern recognition",
"inflammatory cytokines and receptors", and "T/B cell antigen activation". They were
downregulated in the immune process of "inflammatory cytokines and receptors",
"complement system", "adapters, effectors and signal transducers", and "pattern
recognition". Interestingly, there were no genes in "antigen processing and regulators"
for group A. In group A, there were only downregulated genes in "acute phase
reactions", as well as only upregulated genes in "innate immune cells related". In
group B (Figure 5A and Table S6), immune genes were upregulated in the immune
processes, such as "other genes related to immune response", "inflammatory
cytokines and receptors", and "adapters, effectors and signal transducers".
Meanwhile, the downregulated genes were involved in the immune processes, such as
"pattern recognition", "inflammatory cytokines and receptors" and "other genes
related immune response". The largest number of immune mRNAs was observed in
group C (Figure 5A and Table S7). The upregulated genes were mainly involved in
"inflammatory cytokines and receptors", "other genes related to immune response",
"T/B cell antigen activation", "pattern recognition", and "adapters, effectors and
signal transducers", whereas the downregulated genes were mainly involved in "other
genes related to immune response", "adapters, effectors and signal transducers",
"inflammatory cytokines and receptors", "T/B cell antigen activation", "pattern
recognition", and "antigen processing and regulators". In group D (Figure 5B and
Table S8), without "acute phase reaction", "innate immune cells related", and
"complement system", there were only six immune processes. Among those, "T/B cell
antigen activation" and "other genes related to immune response" exhibited the
greatest number of DEG. In group E (Figure 5A and Table S9), there were no genes
involved in "acute phase reaction" and "innate immune cells related". Only
upregulated genes were reported in "antigen processing and regulators" and "T/B cell
antigen activation", while only downregulated genes were noted in "complement
system". The upregulated genes in group E were mostly involved in "adapters,
effectors and signal transducers", and few in "inflammatory cytokines and receptors",
whereas the downregulated genes were mostly involved in "inflammatory cytokines
and receptors".

Venn-regional Analysis of Immune-related DEG from Different Comparison
Groups
A Venn diagram was created to demonstrate the relationship between the immune-
related DEG among different comparison groups (Figure 5C). The region-specific
genes accounted for a large proportion of each comparison group. The region-specific
DEG are listed in Table 2. The results in overlapping regions showed that the greatest
number was observed between the overlap of B and C. There were a few overlapping
genes between groups A and B, and some overlapping genes between A and C. Some
regions had multiple overlaps that contained certain important genes. The overlapping
regional DEGs are listed in Table 3 and details are provided in Table S10.

Immune-related GO Terms and Correlation between Immune-related LncRNAs
and Genes

The results of "WGCNA" were selected to determine the biological processes
involved in CyHV-3-induced modulation of genes by lncRNAs, from an immune
perspective. Among the currently divided 35 gene modules (Figure S4), only 14
modules could be classified according to the immune-related GO terms (Figure S5).
Specifically, "double-stranded DNA binding", "receptor-mediated endocytosis",
"response to bacterium", "lysosome", "defense response to fungus", "killing of cells
of other organism", "oxygen carrier activity", "oxygen binding", "defense response to
bacterium", "killing of cells of other organism", "NuA4 histone acetyltransferase
complex", "defense response to gram-negative bacterium", "defense response to
gram-positive bacterium", "immune response", "regulation of cytokine secretion",
"cytolysis", "execution phase of apoptosis", "cellular response to gamma radiation",
"nucleic acid metabolic process", "TOR signaling", "DNA replication", "ubiquitin-
dependent protein catabolic process", "granulocyte differentiation", and "macrophage
differentiation" were revealed after the "WGCNA" analysis. Furthermore, for the
differentially expressed lncRNAs and their potential modulated genes in comparison
groups (A–C), the immune-related GO terms are also demonstrated in Figure 6.
Notably, "receptor-mediated endocytosis" and "lysosome" were found in both groups
A and B. In addition, "ubiquitin-dependent protein catabolic process", "DNA
replication", and "response to bacterium" were also found in group B. For group C,
besides the GO terms found in groups A and B, "killing of cells of other organism",
"defense response to fungus", and "double-stranded DNA binding" were also
revealed. Specifically, for detailed correlations of lncRNAs and genes in groups A to
C (Table S11), T cell leukemia homeobox 3 (TLX3) and LGALS3 were revealed,
respectively, in the survivors of the resistant strain.

DISCUSSION

This study demonstrated the immune mechanisms of a CyHV-3-resistant strain of
common carp. By comparing the survivors and healthy fish in either resistant or
normal strains, we identified key genes and related lncRNAs involved in different
immune processes and related downstream pathways. The network of these processes
and pathways has been merged to improve clarity. Accordingly, the
immunogenetically insensitive or susceptible factors to CyHV-3 infection were
determined.

The current results indicated the components involved in the entry of the virus.
Firstly, integrin acts as a herpesvirus receptor (23), and the integrin-dependent
signalosome in herpesvirus-infected cells mediated or coactivated numerous
inflammatory responses and signaling transductions (23). Therefore, the current
finding that integrin beta 1 (ITGB1) was significantly downregulated in the resistant

strain compared with the normal strain, even during the steady status, may indicate
that ITGB1 facilitates the binding of herpes virus glycoprotein for entry (24). This
coincides with the finding that integrin signaling promotes the release of intracellular
calcium stores and contributes to viral entry and cell-to-cell spreading via
glycoprotein H, during herpes simplex virus infection in humans (25). Recent reports
have also demonstrated that some integrins on lymphocytes, such as B cells, could
facilitate the mucosa-specific homing (26). Secondly, higher TLR expression in
survivors of the normal strain compared with that of the resistant strain also indicated
its role in virus entry and downstream proinflammatory signaling. Of note, TLR4
signaling leads to the production of proinflammatory cytokines in human lymphatic
endothelial cells (23). In addition, fish-specific and virus-responding TLR18 (27, 28),
(the most overlapping gene among groups) was downregulated in survivors of the
resistant strain, whereas it was upregulated in those of the normal stain. This
suggested its potential ability to facilitate CyHV-3 infection. MAPK signaling, as the
downstream process of pattern recognition receptors, could facilitate the tumor
necrosis factor-alpha (TNFA)-induced suppressor of cytokine signaling 3 (SOCS3)
expression. This can lead to both pro-inflammatory immune response and failure in
growth (29), according to the upregulated genes upon infection observed in the
normal strain in group B.
458
Non-specific binding of the virus by lectins played a protective role in preventing
virus entry. Both upregulated ladderlectins and higher galectin expression were
detected in survivors of resistant strains compared with normal strains. This is
suggestive of their blocking ability for viral proteins (30, 31), such as glycoprotein
(32). Moreover, as the head kidney is one of the major reticuloendothelial systems in
fish (33), mucin was also found to be upregulated as the anti-virus barrier. This was
shown in group B as the gel-forming mucin 5B (MUC5B) (34) was upregulated in the
survivors of the normal strain. However, higher expression levels of membrane-bound
MUC3 (35) were found in survivors of the resistant strain versus the normal strain.
These findings suggested that more mucus was secreted, thereby causing tissue
swelling upon CyHV-3 infection in the normal strain. For the survivors of the resistant
strain, the membrane-bound mucin could effectively bind the virus with no gel.
471
Lectins are involved in the clearance of apoptotic cells via the complement pathway.
Hence, in survivors, the complement components reported in the acute phase in
common carp during CyHV-3 infection (36), could facilitate the phagocytic process
via binding of MRC (Mannose-Receptor C) in fish (37). In the comparison of the
survivors from groups A and B, the observed difference in the phagocytic process
suggested the clearance of apoptosis cells, which was indicated by ubiquitin ligase in
group A. Meanwhile, IL1, IL6, and IL8 were upregulated in group B. With regard to
cytokine signaling, there were less proinflammatory cytokines and immune
regulation-related cytokine receptors upregulated in the survivors of the resistant

strain compared with the normal strain. The interleukin 21 receptor (IL21R), which
was upregulated in group A and downregulated in group B, indicated the control of
inflammation by suppressing STAT3 signaling (38). The upregulated IL23R in group
A can promote cytotoxic ability (39), which may immediately kill infected cells. The
lack of antigen presentation in group A suggested that the infection was overcome
before the amplification of CyHV-3. In contrast, in group B, numerous upregulated
genes were involved in antigen presentation. Therefore, the survivors from breeding
strains overcame the infection mainly through phagocytosis and cytotoxicity at the
cellular level, without activating the typical process of proinflammation as in
survivors from the normal strain.
491
Thus, there are different signatures between the two strains as survival strategies. For
the survivors in the resistant strain, self-repairing related autophagy was detected as
both the KEGG pathways "regulation of autophagy" and autophagy-related fish
antiviral tripartite motif (TRIM) protein (40) were found in group A. These findings
were in line with TRIMs, which were upregulated in group A and downregulated in
group B. In the survivors from the resistant strain, the higher PI3K also suggests
higher autophagy, since the PI3K/AKT/mTOR pathway enhances this process [28].
Fish TRIM may inhibit the activation of IFN and attenuate IFN regulatory factor
(IRF) (41, 42). In addition, in carp, nuclear factor, interleukin 3 regulated (NFIL3)
stimulated both proinflammatory (e.g., NF-kappa B [NF-κB]) and anti-inflammatory
factors (e.g., IL10) (43). This may upregulate NFIL3 in group A, as suggested by the
extensive activation of immune cells, enhancing immune regulation. The directed
lymphocytes response could be present as there was upregulation of IL23R, which
significantly enhances the expression of cytotoxic mediators (39), as well as cathepsin
L (a component of lysosomes) (44). This indicated an enhanced activation of the
cytotoxic ability of T cells in survivors from the resistant strain. For B cells, the
secretion of Ig-related genes (e.g., polymeric immunoglobulin receptor [PIGR], Ig
heavy chain and light chain) was also increased in group A. To fight the virus, the
survivors from the normal strain developed typical inflammatory cascades, including
pro-inflammatory cytokines (e.g., IL1, IL6, and IL8), as well as MAPK signals, which
are profoundly involved in cell survival functions during viral infection (45). For the
control of inflammation, the survivors from the common strain exhibited suppression
of the IFN response, which was also reflected by the upregulated IRF8, the inhibitor
of the MYD88-mediated NF-κB signaling pathway (46). Downstream hypoxia was
also found, as indicated by the "p53 signaling pathway" and "oxidative
phosphorylation". The hypoxic status could also protect the fish from death, since p53
suppresses cell apoptosis (47), and HIF1A regulates virus-induced glycolytic genes
(48).
520
Furthermore, the advantages of disease-resistant genes were reflected by comparing
the survivors from the two strains, which can provide clues for breeding resistance to

CyHV-3. In group C there was more semaphorin, which is related to
immunoregulation (49, 50), in survivors from resistant strains than normal strains.
This indicated a possible tolerance of viral replication, without negative effects on the
proliferation of host cells (51). There was less cyclin-dependent kinase inhibitor 1D
(CDKN1D) in survivors from resistant strains compared with normal strains. This
indicated that there was an interrupted circadian cell-cycle timing in the survivors
from normal strains (52). The higher levels of STAT3 and several integrins in
survivors from the normal strain may facilitate the effect of enhanced IL6 signaling.
This was upregulated in group B through upregulation of integrins by STAT3 (53).
Apart from the typical immune signals in group C, there was relatively higher PI3K
activity, which was involved in the KEGG pathway "dopaminergic synapse". This
indicated that dopamine inhibited inflammation (54) in survivors from the resistant
strain. This is because PI3K is dependent on the accumulation of DOPA
decarboxylase, the enzyme involved in the production of dopamine, which is reduced
by a viral infection [27]. Additionally, genes responsive to the secretion of
immunoglobulin (e.g., PIGR and Ig light chain), which is important for anti-virus
immunity, were common (in group D) between the two strains.
540

The comparison of two strains with healthy status suggested a basal anti-virus
immunity. Among the upregulated genes in group E, the expression of NACHT, LRR,
and PYD domains-containing protein 3 (NALP3) indicated a stronger ability to form
the inflammasome in the resistant strain. The higher levels of NLR family CARD
domain containing 3 (NLRC3), which is a negative regulator of the DNA sensor
STING, suggest less sensitivity to the DNA virus in the resistant strain. The higher
levels of MHC I indicate greater potential to activate response by natural killer cells
upon herpes virus infection (55) in the resistant strain. For the potential T helper cell
differentiation-related JAK/STAT pathway, the upregulation of both STAT3 and
histone deacetylase (HDAC) indicated an easier differentiation of both T helper 17
and regulatory T directions (56), respectively, in the resistant strain even in the steady
state (57, 58). Upregulation of HDAC in group E, that could inhibit the function of
macrophages in fish (59), also suggests a more delicate immune regulation during the
steady state in the resistant strain. However, in group E, among the downregulated
genes, which indicated higher expression in the normal stain at a steady state, TLR18
and ITGB1 suggested susceptibility to virus binding. Moreover, C-C motif chemokine
ligand 4 (CCL4), which can protect infected cells with viral burden (60), may be a
risk factor for the normal strain.
559

The revealed regulation of resistance to CyHV-3 by lncRNA involved numerous
biological processes. Among the revealed GO terms in group B, "DNA replication"
could possibly indicate virus proliferation in fish. In survivors from the normal strain,
the lncRNA-regulated genes were mainly involved in innate immune cell function
(e.g., macrophage protein and cathepsin, as components of lysosomes) and cell

apoptosis. In survivors from the resistant strain, the only lncRNA-modulated gene,
TLX3, has been shown to be strongly methylated. This indicates that TLX3
expression was suppressed during hepatitis B virus-related cancer (61). This suggests
that TLX3 in survivors from a resistant strain may play a protective role, and
participate in lymphocyte proliferation in the head kidney. Meanwhile, echoing recent
findings regarding the participation of zebrafish galectin proteins in immunity against
viral infection (32, 62), the expression of LGALS3 (the only lncRNA regulation-
related immune gene found in group C), was markedly higher than that recorded in
the normal strain upon infection. This suggested an ability of LGALS3 to block viral
attachment.
575
Therefore, based on the present findings, a hypothesis of immune mechanisms
involved in both healthy controls and survivors from infection in both strains, has
been proposed in Figure 7. In conclusion, the resistant strain of common carp showed
a better ability to maintain immune homeostasis in both steady and inflammatory
states, and displayed enhanced blockage of CyHV-3 infection compared with the
normal strain. Since the modulation by lncRNA currently remains a hypothesis,
further molecular evidence is needed to elucidate the mechanism of resistance. In
addition, the finding that both the inhibition of inflammation by dopamine in the
resistant strain and the disrupted bio-clock in the normal strain upon CyHV-3
infection suggested better growth performance for the resistant strain. Thus, the
possible advantage in the resistant strain for growth performance warrants further
study.

AUTHOR CONTRIBUTIONS

Lianyu Shi and Zhiying Jia conceived the project and designed the experiments. Nan
Wu and Zhying Jia wrote the manuscript. Zhying Jia performed the experiments.
Zhying Jia and Jiaxin Sun conducted the CyHV-3 infection experiment and collected
samples. Chitao Li, Xuesong Hu and Yanlong Ge performed fish propagation and
culture. Xiaona Jiang conducted the RT-PCR experiment. Nan Wu and Xiaoqin Xia
coordinated the data analysis tasks. Nan Wu, Heng Li, Mijuan Shi, Xiaona Jiang,
Weidong Ye, Ying Tang, and Yingyin Cheng analyzed the data. The manuscript was
revised and approved by Xiao-Qin Xia and Lianyu Shi.

ACKNOWLEDGMENTS

This work was funded by grants from the National Key R&D Program of China
(2018YFD0900302-6), The Natural Science Foundation of Heilongjiang Province
(TD2019C004), China Aquaculture Research System (CARS-45-06), and Central
Public-interest Scientific Institutional Basal Research Fund (2020TD31).

Conflict of Interest Statement: The authors declare that the research was conducted
in the absence of any commercial or financial relationships that could be construed as
a potential conflict of interest.

CONTRIBUTION TO THE FIELD STATEMENT

Cyprinid herpesvirus-3 (CyHV-3) is the major threat to common carp aquaculture.
Experimental infections of carp from pure lines or crosses have indicated the
existence of genetic background of resistance by divergent survival rates. The
immune response of carp to CyHV-3 involves both innate and adaptive aspects. Even
non-coding RNAs play an important role in anti-CyHV-3 immunity. However, there
is a lack of systemic studies focusing on a detailed network of the immune system for
the anti-CyHV-3 immune mechanism in common carp. Benefiting from a strain of
common carp which has already shown a higher survival rate in CyHV-3 infection,
this study systematically illustrated the immune mechanisms involved in anti-CyHV-3
ability on both mRNA and non-coding RNA levels, demonstrated that the carp may
obtain significant resistance to CyHV-3 through a specific innate immune mechanism,
and revealed the involved biological processes including autophagy, phagocytosis,
cytotoxicity, blocking of the virus by lectins and MUC 3, and the immune suppressive
signal transduction. The immune factors involved in resistant or susceptible to the
disease shed light on prevention of CyHV-3 infection genetically.

DATA AVAILABILITY STATEMENT

The datasets, containing all clean data of this transcriptomic analysis, can be found in
the Genome Sequence Archive (GSA) database (http://gsa.big.ac.cn/index.jsp) with
the BioProject identifier <CRA003105>.

REFERENCES

1.	Neave MJ, Sunarto A, McColl KA. Transcriptomic analysis of common carp
anterior kidney during Cyprinid herpesvirus 3 infection: Immunoglobulin
repertoire and homologue functional divergence. *Sci Rep.* (2017) 7:41531. doi:
10.1038/srep41531.
2.	Adamek M, Steinhagen D, Irnazarow I, Hikima J, Jung TS, Aoki T. Biology
and host response to Cyprinid herpesvirus 3 infection in common carp. *Dev
Comp Immunol.* (2014) 43:151-159. doi: 10.1016/j.dci.2013.08.015.
3.	Sunarto A, McColl KA. Expression of immune-related genes of common carp
during cyprinid herpesvirus 3 infection. *Dis Aquat Organ.* (2015) 113:127-
135. doi: 10.3354/dao02824.
4.	Reichert M, Lukasik A, Zielenkiewicz P, Matras M, Maj-Paluch J, Stachnik M
et al. Host microRNA analysis in cyprinid Herpesvirus-3 (CyHV-3) infected
common carp. *BMC Genomics.* (2019) 20:46. doi: 10.1186/s12864-018-5266-
9.
5.	Rakus KL, Irnazarow I, Adamek M, Palmeira L, Kawana Y, Hirono I *et al*.
Gene expression analysis of common carp (*Cyprinus carpio L.*) lines during
Cyprinid herpesvirus 3 infection yields insights into differential immune
responses. *Dev Comp Immunol.* (2012) 37:65-76. doi:
10.1016/j.dci.2011.12.006.

6. Adamek M, Matras M, Dawson A, Piackova V, Gela D, Kocour M *et al*. Type
I interferon responses of common carp strains with different levels of
resistance to koi herpesvirus disease during infection with CyHV-3 or SVCV.
Fish Shellfish Immunol. (2019) 87:809-819. doi: 10.1016/j.fsi.2019.02.022.

7. Lee X, Yi Y, Weng S, Zeng J, Zhang H, He J *et al*. Transcriptomic analysis of
koi (*Cyprinus carpio*) spleen tissue upon cyprinid herpesvirus 3 (CyHV3)
infection using next generation sequencing. *Fish Shellfish Immunol.* (2016)
49:213-224. doi: 10.1016/j.fsi.2015.12.007.

8. Hwang JA, Kim JE, Kim HS, Lee JH. Immune Response to Koi Herpesvirus
(KHV) of Koi and Koi x Red Common Carp (*Cyprinus carpio*). *Dev Reprod.*
(2017) 21:361-370. doi: 10.12717/DR.2017.21.4.361.

9. Adamek M, Rakus KL, Brogden G, Matras M, Chyb J, Hirono I *et al*.
Interaction between type I interferon and Cyprinid herpesvirus 3 in two
genetic lines of common carp (*Cyprinus carpio*). *Dis Aquat Organ.* (2014)
111:107-118. doi: 10.3354/dao02773.

10. Tadmor-Levi R, Hulata G, David L. Multiple interacting QTLs affect disease
challenge survival in common carp (*Cyprinus carpio*). *Heredity (Edinb).*
(2019) 123:565-578. doi: 10.1038/s41437-019-0224-0.

11. Jia ZY, Chen L, Ge YL, Li SW, Peng WZ, Li CT *et al*. Genetic mapping of
Koi herpesvirus resistance (KHVR) in Mirror carp (*Cyprinus carpio*) revealed
genes and molecular mechanisms of disease resistance. *Aquaculture.* (2020)
519 doi: ARTN 73485010.1016/j.aquaculture.2019.734850.

12. Jia ZY, Wang SH, Bai SS, Ge YL, Li CT, Hu XS *et al*. Survival rate and
immunological responses of mirror carp selective breeding generations to
CyHV-3. *J World Aquacult Soc.* (2018) 49:388-395. doi: 10.1111/jwas.12511.

13. Sun JX, Shi LY, Jiang XN, Li CT, Ge YL, Hu XS *et al*. Research on
expression of virus in cyprinid herpesvirus 3 (CyHV-3) F4 disease resistant
strains. *Journal of Shanghai Ocean University.* (2020) doi: CN 31-2024/S.

14. Conte MA, Gammerdinger WJ, Bartie KL, Penman DJ, Kocher TD. A high
quality assembly of the Nile Tilapia (*Oreochromis niloticus*) genome reveals
the structure of two sex determination regions. *BMC Genomics.* (2017)
18:341. doi: 10.1186/s12864-017-3723-5.

15. Zhang X, Li G, Jiang H, Li L, Ma J, Li H *et al*. Full-length transcriptome
analysis of Litopenaeus vannamei reveals transcript variants involved in the
innate immune system. *Fish Shellfish Immunol.* (2019) 87:346-359. doi:
10.1016/j.fsi.2019.01.023.

16. Wu N, Wang B, Cui ZW, Zhang XY, Cheng YY, Xu X *et al*. Integrative
transcriptomic and microRNAomic profiling reveals immune mechanism for
the resilience to soybean meal stress in fish gut and liver. *Front Physiol.*
(2018) 9:1154. doi: 10.3389/fphys.2018.01154.

17. Xu P, Zhang X, Wang X, Li J, Liu G, Kuang Y *et al*. Genome sequence and
genetic diversity of the common carp, *Cyprinus carpio*. *Nat Genet.* (2014)

46:1212-1219. doi: 10.1038/ng.3098.

18. Wu N, Song YL, Wang B, Zhang XY, Zhang XJ, Wang YL *et al*. Fish gut-liver immunity during homeostasis or inflammation revealed by integrative transcriptome and proteome studies. *Sci Rep.* (2016) 6:36048. doi: 10.1038/srep36048.

19. Wang J, Fu L, Koganti PP, Wang L, Hand JM, Ma H *et al*. Identification and functional prediction of large intergenic noncoding RNAs (lincRNAs) in rainbow trout (*Oncorhynchus mykiss*). *Mar Biotechnol (NY).* (2016) 18:271-282. doi: 10.1007/s10126-016-9689-5.

20. Liao Q, Liu C, Yuan X, Kang S, Miao R, Xiao H *et al*. Large-scale prediction of long non-coding RNA functions in a coding-non-coding gene co-expression network. *Nucleic Acids Res.* (2011) 39:3864-3878. doi: 10.1093/nar/gkq1348.

21. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics.* (2008) 9:559. doi: 10.1186/1471-2105-9-559.

22. Yu GC, Wang LG, Han YY, He QY. clusterProfiler: an R package for comparing biological themes among gene clusters. *Omics.* (2012) 16:284-287. doi: 10.1089/omi.2011.0118.

23. Campadelli-Fiume G, Collins-McMillen D, Gianni T, Yurochko AD. Integrins as herpesvirus receptors and mediators of the host signalosome. *Annu Rev Virol.* (2016) 3:215-236. doi: 10.1146/annurev-virology-110615-035618.

24. Azab W, Gramatica A, Herrmann A, Osterrieder N. Binding of alphaherpesvirus glycoprotein H to surface alpha4beta1-integrins activates calcium-signaling pathways and induces phosphatidylserine exposure on the plasma membrane. *mBio.* (2015) 6:e01552-01515. doi: 10.1128/mBio.01552-15.

25. Cheshenko N, Trepanier JB, Gonzalez PA, Eugenin EA, Jacobs WR, Jr., Herold BC. Herpes simplex virus type 2 glycoprotein H interacts with integrin alphavbeta3 to facilitate viral entry and calcium signaling in human genital tract epithelial cells. *J Virol.* (2014) 88:10026-10038. doi: 10.1128/JVI.00725-14.

26. Delecluse S, Tsai MH, Shumilov A, Bencun M, Arrow S, Beshirova A *et al*. Epstein-Barr virus induces expression of the LPAM-1 integrin in B cells in vitro and in vivo. *J Virol.* (2019) 93 doi: 10.1128/JVI.01618-18.

27. Huang WJ, Shen Y, Xu XY, Hu MY, Li JL. Identification and characterization of the TLR18 gene in grass carp (*Ctenopharyngodon idella*). *Fish Shellfish Immunol.* (2015) 47:681-688. doi: 10.1016/j.fsi.2015.09.052.

28. Lee PT, Zou J, Holland JW, Martin SA, Collet B, Kanellos T *et al*. Identification and characterisation of TLR18-21 genes in Atlantic salmon (*Salmo salar*). *Fish Shellfish Immunol.* (2014) 41:549-559. doi: 10.1016/j.fsi.2014.10.006.

29. Jiang X, He M, Bai J, Chan CB, Wong AOL. Signal transduction for

729   TNFalpha-induced type II SOCS expression and its functional implication in
730   growth hormone resistance in carp hepatocytes. *Front Endocrinol (Lausanne).*
731   (2020) 11:20. doi: 10.3389/fendo.2020.00020.

732 30. Wang W, Cole AM, Hong T, Waring AJ, Lehrer RI. Retrocyclin, an
733   antiretroviral theta-defensin, is a lectin. *J Immunol.* (2003) 170:4708-4716.
734   doi: 10.4049/jimmunol.170.9.4708.

735 31. Zhu D, Fu P, Huang R, Xiong L, Wang Y, He L *et al*. Molecular
736   characterization, tissue distribution and functional analysis of galectin 1-like 2
737   in grass carp (*Ctenopharyngodon idella*). *Fish Shellfish Immunol.* (2019)
738   94:455-463. doi: 10.1016/j.fsi.2019.09.041.

739 32. Ghosh A, Banerjee A, Amzel LM, Vasta GR, Bianchet MA. Structure of the
740   zebrafish galectin-1-L2 and model of its interaction with the infectious
741   hematopoietic necrosis virus (IHNV) envelope glycoprotein. *Glycobiology.*
742   (2019) 29:419-430. doi: 10.1093/glycob/cwz015.

743 33. Wu C, Zhao X, Babu VS, Yuan G, Wang W, Su J *et al*. Distribution of
744   mannose receptor in blunt snout bream (*Megalobrama amblycephala*) during
745   the embryonic development and its immune response to the challenge of
746   *Aeromonas hydrophila. Fish Shellfish Immunol.* (2018) 78:52-59. doi:
747   10.1016/j.fsi.2018.03.049.

748 34. Schneider JR, Shen X, Orlandi C, Nyanhete T, Sawant S, Carias AM *et al*. A
749   MUC16 IgG binding activity selects for a restricted subset of IgG enriched for
750   certain simian immunodeficiency virus epitope specificities. *J Virol.* (2020) 94
751   doi: 10.1128/JVI.01246-19.

752 35. Tang MLK. (2016) "The Physiological Induction of Tolerance to Allergens,
753   Allergy". in Allergy, Immunity and Tolerance in Early Childhood, ed. Wahn U,
754   Sampson HA, Academic Press.

755 36. Pionnier N, Adamek M, Miest JJ, Harris SJ, Matras M, Rakus KL *et al*. C-
756   reactive protein and complement as acute phase reactants in common carp
757   (*Cyprinus carpio*) during CyHV-3 infection. *Dis Aquat Organ.* (2014)
758   109:187-199. doi: 10.3354/dao02727.

759 37. Magnadottir B, Gudmundsdottir S, Lange S. A novel ladder-like lectin relates
760   to sites of mucosal immunity in Atlantic halibut (*Hippoglossus hippoglossus*
761   *L.*). *Fish Shellfish Immunol.* (2019) 87:9-12. doi: 10.1016/j.fsi.2018.12.034.

762 38. Ryu JG, Lee J, Kim EK, Seo HB, Park JS, Lee SY *et al*. Treatment of IL-21R-
763   Fc control autoimmune arthritis via suppression of STAT3 signal pathway
764   mediated regulation of the Th17/Treg balance and plasma B cells. *Immunol*
765   *Lett.* (2015) 163:143-150. doi: 10.1016/j.imlet.2014.09.007.

766 39. Moens E, Brouwer M, Dimova T, Goldman M, Willems F, Vermijlen D. IL-
767   23R and TCR signaling drives the generation of neonatal Vgamma9Vdelta2 T
768   cells expressing high levels of cytotoxic mediators and producing IFN-gamma
769   and IL-17. *J Leukoc Biol.* (2011) 89:743-752. doi: 10.1189/jlb.0910501.

770 40. Langevin C, Levraud JP, Boudinot P. Fish antiviral tripartite motif (TRIM)

proteins. *Fish Shellfish Immunol.* (2019) 86:724-733. doi:
10.1016/j.fsi.2018.12.008.

41.   Wu M, Dan C, Gui JF, Zhang YB. Fish species-specific TRIM gene FTRCA1
negatively regulates interferon response through attenuating IRF7
transcription. *Fish Shellfish Immunol.* (2019) 90:180-187. doi:
10.1016/j.fsi.2019.04.297.

42.   Wu M, Zhao X, Gong XY, Wang Y, Gui JF, Zhang YB. FTRCA1, a species-
specific member of finTRIM family, negatively regulates Fish IFN response
through autophage-lysosomal degradation of TBK1. *J Immunol.* (2019)
202:2407-2420. doi: 10.4049/jimmunol.1801645.

43.   Yu H, Shen Y, Sun J, Xu X, Wang R, Xuan Y *et al*. Molecular cloning and
functional characterization of the NFIL3/E4BP4 transcription factor of grass
carp, *Ctenopharyngodon idella*. *Dev Comp Immunol.* (2014) 47:215-222. doi:
10.1016/j.dci.2014.07.019.

44.   Szymczak M. Distribution of cathepsin D activity between lysosomes and a
soluble fraction of marinating brine. *J Food Sci.* (2016) 81:E1966-1970. doi:
10.1111/1750-3841.13375.

45.   Kumar R, Khandelwal N, Thachamvally R, Tripathi BN, Barua S, Kashyap
SK *et al*. Role of MAPK/MNK1 signaling in virus replication. *Virus Res.*
(2018) 253:48-61. doi: 10.1016/j.virusres.2018.05.028.

46.   Yan X, Zhao X, Huo R, Xu T. IRF3 and IRF8 regulate NF-kappaB signaling
by targeting MyD88 in teleost fish. *Front Immunol.* (2020) 11:606. doi:
10.3389/fimmu.2020.00606.

47.   Feng X, Liu X, Zhang W, Xiao W. p53 directly suppresses BNIP3 expression
to protect against hypoxia-induced cell death. *EMBO J.* (2011) 30:3397-3415.
doi: 10.1038/emboj.2011.248.

48.   Godoy-Lugo JA, Miranda-Cruz MM, Rosas-Rodriguez JA, Adan-Bante NP,
Icedo-Garcia R, Sonanez-Organis JG. Hypoxia inducible factor -1 regulates
WSSV-induced glycolytic genes in the white shrimp Litopenaeus vannamei.
Fish Shellfish Immunol. (2019) 92:165-171. doi: 10.1016/j.fsi.2019.05.040.

49.   Lepelletier Y, Moura IC, Hadj-Slimane R, Renand A, Fiorentino S, Baude C *et*
al. Immunosuppressive role of semaphorin-3A on T cell proliferation is
mediated by inhibition of actin cytoskeleton reorganization. *Eur J Immunol.*
(2006) 36:1782-1793. doi: 10.1002/eji.200535601.

50.   Alamri A, Soussi Gounni A, Kung SKP. View point: semaphorin-3E: an
emerging modulator of natural killer cell functions? *Int J Mol Sci.* (2017) 18
doi: 10.3390/ijms18112337.

51.   Yan Y, Cui H, Guo C, Wei J, Huang Y, Li L *et al*. Singapore grouper
iridovirus-encoded semaphorin homologue (SGIV-sema) contributes to viral
replication, cytoskeleton reorganization and inhibition of cellular immune
responses. *J Gen Virol.* (2014) 95:1144-1155. doi: 10.1099/vir.0.060608-0.

52.   Laranjeiro R, Tamai TK, Peyric E, Krusche P, Ott S, Whitmore D. Cyclin-

dependent kinase inhibitor p20 controls circadian cell-cycle timing. *Proc Natl*
Acad Sci U S A. (2013) 110:6835-6840. doi: 10.1073/pnas.1217912110.

53.   Rivera-Soto R, Dissinger NJ, Damania B. Kaposi's sarcoma-associated
herpesvirus viral interleukin-6 signaling upregulates integrin beta3 levels and
is dependent on STAT3. *J Virol.* (2020) 94 doi: 10.1128/JVI.01384-19.

54.   Kalkman HO, Feuerbach D. Antidepressant therapies inhibit inflammation and
microglial M1-polarization. *Pharmacol Ther.* (2016) 163:82-93. doi:
10.1016/j.pharmthera.2016.04.001.

55.   Finton KA, Strong RK. Structural insights into activation of antiviral NK cell
responses. *Immunol Rev.* (2012) 250:239-257. doi: 10.1111/j.1600-
065X.2012.01168.x.

56.   Zhang H, Xiao Y, Zhu Z, Li B, Greene MI. Immune regulation by histone
deacetylases: a focus on the alteration of FOXP3 activity. *Immunol Cell Biol.*
(2012) 90:95-100. doi: 10.1038/icb.2011.101.

57.   Jin H, Guo X. Valproic acid ameliorates coxsackievirus-B3-induced viral
myocarditis by modulating Th17/Treg imbalance. *Virol J.* (2016) 13:168. doi:
10.1186/s12985-016-0626-z.

58.   Sugimoto K, Itoh T, Takita M, Shimoda M, Chujo D, SoRelle JA *et al.*
Improving allogeneic islet transplantation by suppressing Th17 and enhancing
Treg with histone deacetylase inhibitors. *Transpl Int.* (2014) 27:408-415. doi:
10.1111/tri.12265.

59.   Moreira JD, Koch BEV, van Veen S, Walburg KV, Vrieling F, Guimaraes
TMPD *et al.* Functional Inhibition of Host Histone Deacetylases (HDACs)
Enhances in vitro and in vivo Anti-mycobacterial Activity in Human
Macrophages and in Zebrafish. *Frontiers in Immunology.* (2020) 11: ARTN
3610. doi: 3389/fimmu.2020.00036.

60.   Parekh NJ, Krouse TE, Reider IE, Hobbs RP, Ward BM, Norbury CC. Type I
interferon-dependent CCL4 is induced by a cGAS/STING pathway that
bypasses viral inhibition and protects infected tissue, independent of viral
burden. *Plos Pathog.* (2019) 15: ARTN e1007778. doi:
10.1371/journal.ppat.1007778.

61.   Tao R, Li J, Xin J, Wu J, Guo J, Zhang L *et al.* Methylation profile of single
hepatocytes derived from hepatitis B virus-related hepatocellular carcinoma.
PLoS One. (2011) 6:e19862. doi: 10.1371/journal.pone.0019862.

62.   Nita-Lazar M, Mancini J, Feng C, Gonzalez-Montalban N, Ravindran C,
Jackson S *et al.* The zebrafish galectins Drgal1-L2 and Drgal3-L1 bind in vitro
to the infectious hematopoietic necrosis virus (IHNV) glycoprotein and reduce
viral adhesion to fish epithelial cells. *Dev Comp Immunol.* (2016) 55:241-252.
doi: 10.1016/j.dci.2015.09.007.

FIGURE LEGENDS

Figure 1 Strategy for the identification of regulated immune genes in fish gut and
liver by integrative analysis of both DEG and lncRNA using transcriptomics data. (**A**)
Virus challenge was performed by infecting common carp through addition of sick
fish tissue lysate into the tank. Subsequently, sampling of immune organs was
conducted at day 0 (immediately before challenge) and day 7 post challenge, for
transcriptomics analysis. (**B**) Immune-related annotation was carried out via
bioinformatics analysis of the transcriptomics results using GO annotation and KEGG
pathway enrichment, through filtering using the currently constructed common carp
immune gene library. (**C**) Venn diagrams were generated using immune-related DEG
data from comparisons of groups A–E. Next, Venn-regional analysis was applied to
identify the immune-related anti-CyHV-3 or CyHV-3-resistant protein-coding genes
or lncRNAs.
CyHV-3, cyprinid herpesvirus-3; DEG, differentially expressed gene; GO, Gene
Ontology; KEGG, Kyoto Encyclopedia of Genes and Genomes; lncRNA, long non-
coding RNA
869

Figure 2 Differential appearance of general mortality and pathology between fish
from the resistant and normal strains. (**A**) Comparison of daily mortality between the
resistant and normal strains. The mortality was monitored within 15 days post
challenge. N=300 per group. (**B**) The degree of swelling in the spleen was limited in
the survivors from the resistant strain (a) compared with the markedly enlarged spleen
observed in fish from the normal strain (b). The arrow indicates the spleen region.
876

877   **Figure 3** GO terms comparing groups A, B, and C. The immune-related terms are
878   labeled with stars.
879   GO, Gene Ontology
880

881   **Figure 4** KEGG pathways comparing groups A, B, C, and D. The immune-related
882   terms are labeled with arrows.
883   KEGG, Kyoto Encyclopedia of Genes and Genomes
884

885   **Figure 5** The analysis of immune processes of DEG among different comparison
886   groups.
887   DEG, differentially expressed gene
888

Figure 6 The GO terms involved in immune-related modular, which contain the
matched lncRNA and genes. The arrows indicate the matched lncRNA and genes
containing GO terms in black (**A**), blue (**B**), brown (**C**), grey60 (**D**), and turquoise
(**E**). The star, triangle, and arrow represent the revealed GO term for Group A, Group
B, and Group C, respectively.
GO, Gene Ontology; lncRNA, long non-coding RNA
895

Figure 7 Hypothesis of the carp immune mechanism in both breeding and normal
strains either at the steady state or upon CyHV-3 infection. Group A: healthy control
in the steady state in the breeding strain. Group B: survivors in an inflammatory state
in the breeding strain. Group C: healthy control in the steady state in normal strain.
Group D: survivors in an inflammatory state in the normal strain. The words in italics,
bold, and green represent the biological process, cell type, and lncRNA-modulated
target immune genes, respectively. The arrow denotes facilitation or promotion, and --
-| denotes inhibition. The dashed line indicates a possible correlation.
CyHV-3, lncRNA, long non-coding RNA
905

906   **Table 1** The immune processes and immune gene categories involved in differentially

907   expressed mRNA for each comparison group

Compared group	Immune process	Immune gene category
Upregulated in group A (26)	Pattern recognition (6), complement system, inflammatory cytokines and receptors (5), adapters, effectors and signal transducers (4), innate immune cells related (2), T/B cell antigen activation (5), other genes related to immune response (3).	ladderlectin, LRR-containing proteins, MRC, C3, GVIN, IL21R, IL23R, integrin alpha, NFIL3, CSF, perforin, UBE, MMR, NCAM, Ig heavy chain, Ig light chain, PIGR, SEMA, cathepsin, TRIM, ubiquitin ligase.
Downregulated in group A (23)	Acute phase reactions (2), pattern recognition (4) complement system (4), inflammatory cytokines and receptors (5), adapters, effectors and signal transducers (4), T/B cell antigen activation (2), other genes related to immune response.	Fibrinogen, C-type lectin, galectin, TLR18, C3, CXCL, GVIN, IFI, IL1R, integrin alpha, NALP12, NLRC3, Ig light chain, TIP, myeloid cell related.
Upregulated in group B (155)	Pattern recognition (13), antigen processing and regulators (13), complement system (5), inflammatory cytokines and receptors (34), adapters, effectors and signal transducers (23), innate immune cells related (6), T/B cell antigen activation (15), other genes related to immune response (46).	FBXL, fucolectin, MPR, nattectin, NITR, TLR1, TLR18, TLR22, TLR25, TLR5, CDK, CDK related, CDKIs, HA, MHC I, MHC II, TGFB, TNF, TNFAIP3-interacting protein, C3, C3aR, C7, CCL, CCR, CFLAR, CXCL, CXCR, cytokine receptor, IFI, IFIT, IL1, IL1R, IL1R related, IL6, IL6R, IL8, IRF8, MX, NFIL3, MAPK, MAPKKK, NALP12, NLRC3, Programmed cell death protein, SOCS1, SOCS3, SOCS7, TRAF, TTRAP, UBE, MIF, NCF, NCK and related, NRAMP, BCL, BLNK, CD22, FCR, HDAC, HMG, Ig light chain, PBX, TAGAP, TCIRG1, AP-3, CARD, cathepsin, CEBP, hepcidin, HSP, HSP5, HSP70, lymphocyte antigen, mucin, myeloid cell related, PIN1, platelet related, TMED, ubiquitin ligase, ubiquitin related, UBL.
Downregulated in group B (102)	Acute phase reactions, pattern recognition (21), antigen processing and regulators (6), complement system (2), inflammatory cytokines and receptors (21), adapters, effectors and signal transducers (14), innate immune cells related (3), T/B cell antigen activation (14), other genes related to immune response (20).	Plasminogen, collectin, C-type lectin, galectin, Hepatic lectin, LRR-containing proteins, RBL, SITR, TLR13, CDK, VEGF, VEGFR, C3aR, CD59, ACRs, CCL, CCR, EGFR, FGFR, GVIN, IL17R, IL21R, IL2R, IL6R, IL8, integrin alpha, integrin beta, IRF related, TITIN, granzyme, MAPKKK, NALP12, NALP3, perforin, UBE, TRIM.
Upregulated in group C (357)	Acute phase reactions (3), pattern recognition (57), antigen processing and regulators (11), complement system (4), inflammatory cytokines and receptors (91), adapters, effectors	Fibrinogen, MBL, collectin, C-type lectin, FBXL, fucolectin, galectin, Hepatic lectin, intelectin, ladderlectin, LRR-containing proteins, MRC, NITR, RBL, scavenger receptor, SIGLEC, TLR8, TLR13, TLR3, TLR9, CDK, CDKIs, CIITA, LRMP, MHC I, MHC II, TGBR, VEGF, C3, C3aR, C5, C8, ACRs, CCL, CCR, CXCR, EGF, FGFR, GVIN, IFI, IFIT, IL10R, IL13R, IL17R, IL1R

	and signal transducers (50), innate immune cells related (6), T/B cell antigen activation (64), other genes related to immune response (70).	related, IL21R, IL23R, IL2R, IL6R, IL8, integrin alpha, integrin beta, IRF4, IRF5, IRF7, MX, NFIL3, TITIN, XCR, CBX4, CSF, CSFR, granzyme, MAPK, MAPKKK, NALP12, NALP3, NDRG1, NLRC3, perforin, Programmed cell death protein, TRAF, TRAIL, UBE, MCP, MMR, NCAM, B-cell CLL/lymphoma, CD2, CD22, CD276, CD4, CD6, FCR, GATA and related, HDAC, Ig heavy chain, Ig light chain, immunoglobulin superfamily, JAG, LAG, MAL, NFAT, PIGR, PKC, SEMA, T-bet, VSIG, VTCN, ZAP, ACTININ, cathepsin, CD302, HSPb1, lymphocyte antigen, lymphocyte related, mucin, myeloid cell related, nectin, palladin, PI3K, rootletin, RTK, TIAM, ubiquitin ligase, ubiquitin related, TRIM, UBL.
Downregulated in group C (303)	Acute phase reactions (2), pattern recognition (27), antigen processing and regulators (25), complement system (11), inflammatory cytokines and receptors (55), adapters, effectors and signal transducers (58), innate immune cells related (7), T/B cell antigen activation (37), other genes related to immune response (81).	Macroglobulin, CD209, c-type lectin, FBXL, galectin, intelectin, LPS-anchor protein, LRR-containing proteins, MPR, TLR1, TLR18, TLR22, TLR25, TLR4, TLR5, CDK, CDK related, CDKIs, HA, MHC II, TGFB, TNF, TNFAIP3-interacting protein, C3, C3aR, C4, C7, CD59, ACRs, CCL, CCR, CD11, CFLAR, CXCL, CXCR, cytokine receptor, FAM, FGF, GVIN, HB-EGF, IL1, IL12, IL12R,IL1R, IL1R related, IL2R, IL6, IL6R, IL8, integrin alpha, integrin beta, IRF1, IRF8, NFIL3, CSF, CSFR, JAK1, MAPK, MAPKKK, NALP12, NALP3, NLRC3, Programmed cell death protein, SOCS1, SOCS3, SOCS7, STAT3, UBE, MIF, NCF, NCK and related, NRAMP, BAFF, B-cell CLL/lymphoma, BCL, BLNK, CD22, CD276, FCR, HDAC, HMG, Ig light chain, PBX, PIGR, SEMA, TAGAP, TAL, TCIRG1, TCIRG2, TCIRG3, VSIG, WWP, ACTININ, AP-3, CARD, caspase, cathepsin, CEBP, hepcidin, HSP, HSP4, HSP5, HSP70, HSP90, Kruppel-like factor, lymphocyte antigen, MALT1, mucin, myeloid cell related, PAXILLIN, PIN1, platelet related, TMED, ubiquitin ligase, ubiquitin related, TRIM, UBL.
All in group D (48)	Pattern recognition (4), antigen processing and regulators (7), inflammatory cytokines and receptors (7), adapters, effectors and signal transducers (4) T/B cell antigen activation (14), other genes related to immune response (12).	LRR-containing proteins, VEGFR, VEGF, CDKIs, CDK, NFIL3, LIFR, integrin beta, FAM, CCR, CD166, MAPK, CD97, SEMA, PIGR, NFAT, Ig light chain, HMG, HDAC, BTG, BCL, ubiquitin ligase, platelet related, HIF1a, CEBP.
Upregulated in group E (12)	Pattern recognition, antigen processing and regulators, inflammatory cytokines and receptors (3), adapters, effectors and signal transducers (4), T/B cell antigen activation, other genes related to immune response (2).	MRC, MHC I, GVIN, IL21R, integrin alpha, NALP3, NLRC3, perforin, STAT3, HDAC, platelet related, ubiquitin related.
Downregulated in group E (7)	Pattern recognition, complement system, inflammatory cytokines and receptors (3), adapters, effectors and	TLR18, C3, CCL, IFI, integrin beta, perforin, myeloid cell related.

signal transducers, other genes related

to immune response.

908 Notes: the number (>1) of DEG is shown in the brackets. The full names of all gene

909 abbreviations could be found in the Tables S5–S9.

911 **TABLE 2** Comparison of group-specific DEG in the Venn regional analysis

Comparison group (DEG No.)	Immune process (DEG NO.)	Immune gene category
Up-regulated in group A specific (7)	inflammatory cytokines and receptors, innate immune cells related, pattern recognition, T/B cell antigen activation (3), other genes related to immune response	NFIL3, MMR, cathepsin, MRC, Ig heavy chain, SEMA
Downregulated in group A specific (10)	acute phase reactions (2), inflammatory cytokines and receptors (2), adapters, effectors and signal transducers (3), T/B cell antigen activation (2), other genes related to immune response	Fibrinogen, NLRC3, GVIN, integrin alpha, myeloid cell related, Ig light chain, TIP
Upregulated in group B specific (29)	pattern recognition (5), antigen processing and regulators, complement system, inflammatory cytokines and receptors (5), adapters, effectors and signal transducers (8), T/B cell antigen activation (2), other genes related to immune response (7)	MAPK, NLRC3, programmed cell death protein, SOCS3, TRAF, TTRAP, MHC I, C3, IFI, IFIT, MX, NFIL3, CEBP, mucin, myeloid cell related, ubiquitin ligase, TRIM, ubiquitin related, Fucolectin, nattectin, NITR, TLR1, BCL, HMG
Downregulated in group B specific (36)	acute phase reactions, pattern recognition (9), antigen processing and regulators, complement system, inflammatory cytokines and receptors (7), adapters, effectors and signal transducers (3), innate immune cells related, T/B cell antigen activation (3), other genes related to immune response (10)	Plasminogen, NALP12, perforin, UBE, VEGFR, CD59, CCL, EGFR, integrin alpha, integrin beta, IRF related, NCAM, ACTININ, CD33, HSP90, Kruppel-like factor, MALT1, ubiquitin ligase, UBL, C-type lectin, galectin, Hepatic lectin, LRR-containing proteins, RBL, SITR, B-cell CLL/lymphoma, Ig heavy chain, TCR
Upregulated in group C specific (278)	acute phase reactions (3), pattern recognition (41), antigen processing and regulators (9), complement system (2) inflammatory cytokines and receptors (74), adapters, effectors and signal transducers (35), innate immune cells related (3), T/B cell antigen activation (51), other genes related to immune response (60)	Fibrinogen, MBL, CBX4, CSFR, granzyme, MAPK, MAPKKK, NALP12, NALP3, NDRG1, NLRC3, perforin, Programmed cell death protein, TRAF, TRAIL, UBE, CDK, CDKIs, CIITA, LRMP, MHC I, MHC II, TGBR, C5, C8, CCL, CCR, CXCR, EGF, FGFR, GVIN, IFI, IFIT, IL10R, IL13R, IL17R, IL1R related, IL21R, IL23R, IL2R, IL6R, IL8, integrin alpha, integrin beta, IRF4, IRF5, IRF7, MX, NFIL3, XCR, MCP, MMR, NCAM, cathepsin, CD302, HSPb1, lymphocyte antigen, lymphocyte related, mucin, myeloid cell related, nectin, paladin, PI3K, rootletin, TIAM, TRIM, ubiquitin ligase, ubiquitin related, collectin, C-type lectin, FBXL, Fucolectin, galectin, Hepatic lectin, intelectin, LRR-containing proteins, MRC, NITR, scavenger receptor, SIGLEC, TLR 8, TLR3, TLR9, B-cell CLL/lymphoma, CD2, CD22, CD276, CD4, CD6, FCR, GATA and related, HDAC, Ig heavy chain, immunoglobulin superfamily, JAG, LAG, MAL,

Downregulated in group C specific (172)	acute phase reactions (2), pattern recognition (16), antigen processing and regulators (13), complement system (5), inflammatory cytokines and receptors (26), adapters, effectors and signal transducers (42), innate immune cells related, T/B cell antigen activation (24), other genes related to immune response (43)	NFAT, PIGR, PKC, SEMA, T-bet, VSIG, VTCN, ZAP Macroglobulin, CSF, CSFR, JAK1, MAPK, MAPKKK, NALP3, NLRC3, Programmed cell death protein, SOCS1, SOCS3, STAT3, UBE, CDK, CDK related, CDKIs, MHC II, TNF, TNFAIP3-interacting protein, C3aR, C4, C7, CD59, ACRs, CD11, CXCL, FAM, FGF, GVIN, HB-EGF, IL1, IL12, IL12R, IL1R related, IL2R, IL6, IL6R, integrin alpha, integrin beta, IRF1, NFIL3, NCK and related, ACTININ, AP-3, CARD, caspase, cathepsin, CEBP, HSP4, HSP70, HSP90, Kruppel-like factor, MALT1, mucin, myeloid cell related, PAXILLIN, platelet related, TMED, ubiquitin ligase, ubiquitin related, TRIM, UBL, CD209, C-type lectin, FBXL, intelectin, LPS-anchor protein, LRR-containing proteins, MPR, TLR4, BAFF, B-cell CLL/lymphoma, BCL, BLNK, CD22, CD276, FCR, HDAC, Ig light chain, PBX, PIGR, SEMA, TAL, TCIRG2, TCIRG3, VSIG, WWP
All in group D specific (44)	pattern recognition (4), antigen processing and regulators (4), inflammatory cytokines and receptors (7), adapters, effectors and signal transducers (4), T/B cell antigen activation (14), other genes related to immune response (11)	CD166, CD97, MAPK, CDK, CDKIs, VEGF, VEGFR, CCR, FAM, integrin beta, LIFR, NFIL3, CEBP, HIF1a, platelet related, ubiquitin ligase, LRR-containing proteins, BCL, BTG, HDAC, HMG, Ig light chain, NFAT, PIGR, SEMA
Upregulated in group E specific (9)	pattern recognition, antigen processing and regulators, inflammatory cytokines and receptors, adapters, effectors and signal transducers (3), T/B cell antigen activation, other genes related to immune response (2),	NLRC3, platelet related, GVIN, HDAC, MHC I, MRC, ubiquitin related, NALP3, STAT3
Downregulated in group E specific (2)	inflammatory cytokines and receptors, other genes related to immune response	integrin beta, myeloid cell related

912   Notes: the number (>1) of DEG is shown in the brackets. The full names of all gene
913   abbreviations could be found in the Tables S10.
914

915   **TABLE 3** Overlapping genes in the Venn regional analysis

Overlapping regions (DEG NO.)	Immune process (DEG NO.)	Immune gene category
Group A & C specific (23)	pattern recognition (7), complement system (4), inflammatory cytokines and receptors (3), adapters, effectors and signal transducers (4), innate immune cells related, T/B cell antigen activation (2), other genes related to immune response (2)	CSF, NALP12, UBE, C3, GVIN, IL23R, integrin alpha, NCAM, ubiquitin ligase, C-type lectin, galectin, ladderlectin, LRR-containing proteins, MRC, PIGR
Group A & E specific (1)	inflammatory cytokines and receptors	IFI
Group B & C specific (176)	pattern recognition (18), antigen processing and regulators (14), complement system (4), inflammatory cytokines and receptors (38), adapters, effectors and signal transducers (23), innate immune cells related (8), T/B cell antigen activation (24), other genes related to immune response (47)	Granzyme, MAPK, MAPKKK, NALP12, NALP3, NLRC3, perforin, Programmed cell death protein, SOCS1, SOCS3, SOCS7, UBE, CDK, CDK related, CDKIs, HA, MHC II, TGFB, TNF, TNFAIP3-interacting protein, VEGF, C3aR, C7, ACRs, CCL, CCR, CFLAR, CXCL, CXCR, cytokine receptor, FGFR, GVIN, IL1, IL17R, IL1R related, IL2R, IL6, IL6R, IL8, integrin alpha, IRF8, NFIL3, TITIN, MIF, NCAM, NCF, NCK and related, NRAMP, ACTININ, AP-3, CARD, cathepsin, CEBP, hepcidin, HSP, HSP5, HSP70,lymphocyte antigen, paladin, PIN1, platelet related, RTK, TMED, TRIM, ubiquitin ligase, ubiquitin related, UBL, collectin, C-type lectin, FBXL, galectin, LRR-containing proteins, MPR, RBL, TLR1, TLR13, TLR22, TLR25, TLR5, BCL, BLNK, CD22, FCR, HDAC, HMG, Ig heavy chain, Ig light chain, JAG, NFAT, PBX, SEMA, TAGAP, TCIRG1
Group B & D (4)	antigen processing and regulators (3), other genes related to immune response	CEBP, VEGFR
Group B & E specific (1)	inflammatory cytokines and receptors	integrin alpha
Group C & E specific (1)	inflammatory cytokines and receptors	CCL
Group A & B & C specific (5)	pattern recognition, inflammatory cytokines and receptors (3), adapters, effectors and signal transducers	LRR-containing proteins, CXCL, IL1R, perforin
Group A & B & E (1)	complement system	C3
Group B & C & E (2)	adapters, effectors and signal transducers (2)	perforin
Group A & B & C	pattern recognition, inflammatory cytokines and	TLR18, IL21R

& E (2)	receptors,	

916 Notes: the number (>1) of lncRNAs or mRNAs is shown in the brackets. The full
917 names of all gene abbreviations could be found in the Tables S10.

Stage II : Transcriptomic Analysis

Stage I : Virus Challenge

A

common carp cultivars with vs without breeding (~70g, with nontreated controls)

Day 0 | Day 7

25°C

Immersion with spleen and kidney tissue fluids from diseased fish

Sampling & sequencing

(HK, for DEG analysis) Illumina

Pac-Bio (mixture of LI, SP and HK, as the reference)

B

DEG result: GO annotation & KEGG pathway enrichment

Key regulated or immune related terms or pathways

Constructing a carp immune gene library

Blast all carp genes using NR, Swissprot, Pfam database, and selected immune genes

Flitting

C

Comparation within breed or no-breed population (infected vs Ctrl) , between survivors from different cultivars , and between survivor and near-death fish

Venn-regional analysis immune DEG

Anti-CyHV3 related genes or LncRNAs upon infection or homeostasis

CyHV3-resistance related genes or LncRNAs upon infection

PPI interaction analysis between DEG and lncRNA

Stage III : Hypothesis

Immunologic mechanism involved in common carp anti-CyHV3 breeding

Public database analysis

Bioinformatic analysis

Common carp immune gene library analysis

qPCR validation