SorCS1 binds the insulin receptor to enhance insulin sensitivity.

Mads Kjolby[1,2,3*#], Karen Marie Pedersen[1,3*], Peter Breining[1,3*], Niels Wellner[1,3], Karen Marie Juul Sørensen[1,3,4], Dorthe Maria Vittrup[3], Anette Marianne Prior Gjesing[5], Dovile Januliene[4], Arulmani Manavalan[6], Maarten Rotman[6], Gangadaar Thotakura[6], Sune Skeldal[3], Peder Madsen[3], Søren Thirup[4], Guido Hermey[7], Christian B. Vaegter[3], Arne Möller[4,8], Olof Asplund[9], Rashmi B. Prasad[9], Edwin T. Parlevliet[10], Patrick C.N. Rensen[10], Ernst Martin Füchtbauer[11], Andreas Nygaard Madsen[12], Birgitte Holst[12], Olav M. Andersen[1,3], Torben Hansen[5], and Anders Nykjaer[1,3,6, 13,#].

[1]PROMEMO and DANDRITE, Aarhus University, Denmark; [2]Department of Clinical Pharmacology, Aarhus University Hospital, Denmark and Danish Diabetes Academy; [3]Department of Biomedicine, Aarhus University, Denmark; [4]Department of Molecular Biology, Aarhus University; [5]Novo Nordisk Foundation Center for Basic Metabolic Research, Copenhagen University, Denmark; [6]Department of Neuroscience, Mayo Clinic, Florida, USA; [7]Institute for Molecular and Cellular Cognition, Center for Molecular Neurobiology Hamburg, University Medical Center Hamburg-Eppendorf; [8]Osnabrück University, Frankfurt Am Main Area, Germany; [9]Lund University Diabetes Centre, Malmö, Sweden; [10]Dept. Medicine, Div. Endocrinology, and Einthoven Laboratory for Experimental Vascular Medicine, Leiden University Medical Center (LUMC), Leiden, The Netherlands; [11]Department of Molecular Biology and Genetics, Aarhus University, Denmark; [12]RMPC, Department of Biomedical Sciences, Copenhagen University, Denmark; [13]Department of Neurosurgery, Aarhus University Hospital, Denmark. * Shared first authors. #Corresponding authors

Anders Nykjaer

PROMEMO and DANDRITE

Department of Biomedicine

Hoegh-Guldbergsgade 10,

Aarhus University

DK-8000 Aarhus C

Denmark

Phone: +45 2899 2384

Email:an@biomed.au.dk

Mads Kjolby

DANDRITE

Department of Biomedicine,

Hoegh-Guldbergsgade 10,

Aarhus University

DK-8000 Aarhus C

Denmark

Phone: +45 6086 6653

Kjolby@biomed.au.dk

Word count: 2579 words, excluding title, authors, figure legends and references.

Type II diabetes mellitus (T2DM) has reached endemic levels and comprises a substantial burden

to the patient and the society. Along with lifestyle factors, a number of genetic loci predisposing to

T2DM have been suggested including *SORCS1* that encodes a type-1 transmembrane receptor.

Here we establish SorCS1 as a high-affinity binding partner for the insulin receptor (IR) that

increases insulin affinity, Akt activation, and peripheral glucose uptake. Mice lacking full-length

SorCS1 develop age-dependent insulin resistance characterized by increased plasma glucose and

insulin levels and a blunted response to exogenous insulin. SorCS1 exists in three forms; as a

transmembrane monomer and dimer, and as a truncated soluble form (sol-SorCS1) produced in

adipose tissue and is present in plasma. Whereas dimeric SorCS1 engages the proform of the

insulin receptor (proIR) during biosynthesis and supports its maturation, the monomeric form

stabilizes the mature IR at the plasma membrane. In its soluble form, SorCS1 positively correlates

with body mass index and inversely with plasma glucose in diabetic patients, and in mouse models

of insulin resistance, overexpression or exogenous administration of the monomeric soluble

receptor domain restores insulin sensitivity. We conclude that SorCS1 is a critical regulator of

peripheral insulin sensitivity operating in both a cell autonomous and endocrine manner, and we

propose sol-SorCS1 as a novel adipokine.

The mechanisms controlling insulin sensitivity and glucose metabolism are fundamental to all vertebrates. Skeletal muscle is essential to acutely buffer high plasma glucose and reduced insulin sensitivity is affected very early on in the development of type 2 diabetes[1,2]. The downstream signaling pathways of IR in skeletal muscle (e.g. Akt and mTOR) are well described, but whether insulin sensitivity may be dynamically regulated at the level of the plasma membrane through interaction with co-receptors or other membrane proteins is unknown[3-5]. The existence of such a molecular thermostat to adapt insulin sensitivity to the metabolic state would transform our conceptual understanding on how key metabolic processes can be controlled.

SorCS1 is a member of the Vps10p-domain family of sorting receptors that also comprises sortilin, SorLA, SorCS2, and SorCS3[6,7]. Several studies have identified *SORCS1* as a risk locus for type 2 diabetes mellitus that is conserved in women[8-10], and in female rats[11] and mice[12]. Recently, SorCS1 was shown to facilitate the biogenesis of insulin containing secretory granules, but a function outside the β-cells was not studied[13]. The structurally related receptor sortilin can function in the secretory pathway by assisting hepatic lipoprotein export[14-16], while in neurons it physically interacts with tropomyosin receptor kinases (Trks) to enable their surface exposure and signaling capacity[17]. We hypothesized that SorCS1 likewise may engage a tyrosine kinase receptor to promote its activity. Given the genetic association with T2DM we here asked whether this tyrosine kinase receptor could be IR.

SorCS1 exists in several isoforms comprising an identical extracellular domain encoded by exons 1-24 but with at least four distinct sorting competent cytoplasmic tails (a, b, c, and d)[18,19] resulting from alternative splicing from exons 25 to 28 (exon 25 also encodes the transmembrane domain) (**Extended Data Fig. 1**). Regardless the isoform, the extracellular domain (sol-SorCS1) can be efficiently liberated from the plasma membrane by ADAM17/TACE mediated cleavage[20] but the biological

relevance of this is unknown. To investigate the function specifically of the full-length receptor variants

we generated a mouse model (*Sorcs1^Δ25-28*), in which exons 25-28 had been deleted leaving the first 24

exons intact to allow production of only sol-SorCS1 (**Extended Data Fig. 1**). RT-PCR for *Sorcs1* in

brain, adipose tissue and muscle extracts confirmed that exons 25-28 had been deleted whereas the

extracellular domain encoded by exons 1-24 were transcribed.

Starting from 23 weeks of age, female *Sorcs1^Δ25-28* mice fed a normal chow developed

hyperglycemia. This was paralleled by elevated plasma insulin levels (**Fig. 1a-b**). The hyperinsulinemia

progressively increased as animals aged, reaching 4.6-fold higher insulin levels in 50 weeks old

compared 10 weeks old mice, and twice the concentration in wildtype controls at 50 weeks. Consistent

with genetic studies in mice, rats and humans[8-12], the glycemic phenotype was present only in female

Sorcs1^Δ25-28 mice inasmuch plasma glucose in the male transgenics remained normal up till 60 weeks of

age (**Extended Data Fig. 2**). An oral glucose tolerance test (OGTT) confirmed that glucose handling

was impaired in the females. Hence, in 14 months old *Sorcs1^Δ25-28* mice plasma glucose peaked at 15.7

mM compared to 12. 6 mM for their littermate controls and it returned slower to baseline, corresponding

to a 45% increase in the area under the curve (282 vs. 506 mM*min) (**Fig. 1 c-d**).

The perturbed glucose homeostasis was most pronounced when comparing littermates from

heterozygous breeding on a mixed C57BL/6BomTac x 129sv genetic background (**Fig. 1a-d**). However,

reduced peripheral insulin sensitivity was also evident in 47 weeks old C57BL/6BomTac congenic

female mutant mice when subjected to hyperinsulinemic euglycemic clamp[21,22] (**Fig. 1e-f and Extended**

Figs. 3a-b). At the beginning of the experiment, plasma glucose did not differ between genotypes and

neither did peripheral glucose uptake nor hepatic glucose production. However, when continuously

infusing a similar concentration of insulin (**Extended Data Fig. 3c**) we observed a 37% weaker glucose-

suppressing effect at euglycemic levels in the $Sorcs1^{\Delta25-28}$ mutants compared to wildtype mice (**Fig. 1e**).

This difference was solely accounted for by reduced peripheral glucose uptake since hepatic glucose

production had been fully shutdown (**Fig. 1f and Extended Fig. 3b**).

64   To assess peripheral glucose uptake, we examined the role of SorCS1 on IR signaling in

muscle extracts, in primary differentiated myocytes, and in the myocyte-derived cell line C2C12. In 19

weeks old normoglycemic female mice administration of an insulin bolus potently stimulated IR

phosphorylation (pIRβ) in skeletal muscle of both $Sorcs1^{\Delta25-28}$ mutant animals and their littermate

controls. In marked contrast, at 50 weeks of age, when $Sorcs1^{\Delta25-28}$ mice were diabetic, pIR was now

substantially compromised (**Fig. 2a**). A blunted insulin response was also present in primary $Sorcs1^{\Delta25-28}$

myocyte cultures, as more than 10-fold higher insulin concentrations were required to activate IR in

the mutant cultures compared to those from wildtype mice (**Fig. 2b**). Finally, we studied insulin

stimulated AKT signaling in C2C12 cells after CRISPR/Cas9 knockout of $Sorcs1$. Corroborating the data

in the primary myocytes, C2C12 knockout cells were less insulin responsive as the EC50 curve for pAKT

was right-shifted approximately five-fold (**Fig. 2c**). In the mutant $Sorcs1^{\Delta25-28}$ myocyte cultures we

observed a band with a molecular weight smaller than that of the IRβ-chain (**Fig. 2b**). Surface

biotinylation experiments revealed that this band was restricted to the interior of the cells indicating a

degradation product of proIR or IR. Possibly as a consequence, IRβ expression at the plasma membrane

was reduced by up to 60% in the $Sorcs1^{\Delta25-28}$ cultures (**Fig. 2d**). Impaired N-glycosylation of proIR

prevents two-chain processing and leads to degradation of the pro-receptor[23]. In the mutant myocytes we

observed in addition to mature proIR a species of lower molecular weight (**Fig. 2e**). Treatment with

PNGase F to remove N-linked oligosaccharides identified this protein as immature non-glycosylated

proIR. Taken together the data argue that intracellular SorCS1 facilitates glycosylation and maturation

of proIR and prevents its degradation.

84   We next asked whether surface exposed SorCS1 may also control IR activity. To this end,

IR was overexpressed in HEK 293 cells alone or in combination with SorCS1-b, an isoform that is

enriched at the plasma membrane and incapable of endocytosis and sorting[24]. Co-expression with SorCS1

increased insulin affinity to IR by approximately five-fold (**Fig. 3a**). To assess whether a physical

interaction between SorCS1 and IR could be involved we employed surface plasmon resonance (SPR)

analysis using the extracellular domains of SorCS1 (sol-SorCS1) and IR. The SPR analysis showed that

sol-SorCS1 binds to immobilized IR with an estimated affinity of 5 nM (**Fig. 3b**). Importantly, binding

of insulin to IR did not inhibit its interaction with SorCS1, indicating formation of a trimeric complex

(**Fig. 3c**). No binding was observed for insulin, pro-insulin, glucagon, or glucagon-like peptide 1 (GLP-

1) to SorCS1 (**Extended Data Fig. 4a**). To demonstrate that SorCS1, IR, and insulin can physically

interact at the plasma membrane we performed immunoprecipitation (IP) experiments in HEK293 cells

co-transfected with SorCS1 and IR in the absence or presence of insulin (**Fig. 3d**). Antibodies against IR

co-precipitated SorCS1 and vice versa and this interaction was unaffected by preincubation with 100 nM

insulin. Notably, IP of SorCS1 also co-precipitated proIR, which supports its role in facilitating N-linked

glycosylation of the receptor in the biosynthetic pathway.

99   SorCS1 exists as a non-covalent dimer and in a monomeric form, the conformational state

100   of which is regulated by the level of glycosylation[25]. While unglycosylated receptor is a dimer,

101   maturation with N-linked oligosaccharides converts it into a monomer[25]. We asked whether the

102   stoichiometric state may regulate the preference of SorCS1 for proIR or IR. To elucidate this, HEK293

103   cells expressing endogenous IR were co-transfected with two SorCS1 constructs where the cytoplasmic

104   tail had been substituted by a FLAG- and Myc-tag, respectively. The cells were then subjected to

105 sequential co-immunoprecipitation, first with anti-FLAG and after elution from the beads subsequently

106 with anti-Myc (**Fig. 3e, Extended Data Fig. 4b**). Immunoprecipitation with anti-FLAG-coated beads

107 precipitates receptor populations that comprise a mixture of monomers (SorCS1-FLAG) and dimers

108 (SorCS1-FLAG/FLAG and SorCS1-FLAG/Myc). After elution from the beads followed by the second

109 round of IP using anti-Myc antibodies only dimeric SorCS1 will be left (SorCS1-FLAG/Myc) (**Fig. 3e**).

110 Anti-FLAG immunoprecipitated in addition to the Myc-tagged SorCS1 both proIR and mature IR.

111 However, following the successive co-IP with anti-Myc to select for dimeric SorCS1, only proIR was

112 now co-immunoprecipitated. We conclude that dimeric SorCS1 is specific for proIR whereas monomeric

113 SorCS1 can bind mature IR. To explore the spatial distribution of these interactions, we used a

114 bimolecular fluorescence complementation (BiFC) assay[26]. HEK293-IR cells were transfected with two

115 human SorCS1 constructs each containing one-half of the Venus fluorescent protein. When SorCS1

116 dimerizes the two Venus fragments unite to elicit a green fluorescent signal that echoes the subcellular

117 distribution of dimeric SorCS1. First, we stained the cells with an antibody that recognizes both

118 monomeric and dimeric SorCS1 and found that only monomeric SorCS1 is present at the plasma

119 membrane as the BiFC-positive signal was restricted to intracellular compartments (**Fig. 3f**). We next

120 combined the BiFC signal with proximity ligation assay (PLA) for SorCS1 and IR. In this assay a red

121 fluorescent signal will appear when the receptors are in close proximity (<30 nm). In unpermeabilized

122 cells we observed a strong PLA signal at the plasma membrane that was BiFC-negative, indicating

123 binding of monomeric SorCS1 to mature IR (**Fig. 3g**). In contrast, in permeabilized cells a strong PLA

124 signal was present in BiFC-positive intracellular structures compatible with the Golgi-compartment and

125 the endoplasmatic reticulum that is dispersed throughout the cytosol (**Fig. 3h**). Because these

126 compartments are enriched in proIR, the combined data suggest that monomeric SorCS1 engages IR at

the plasma membrane whereas the dimeric and not fully glycosylated receptor binds proIR in the

secretory pathway[25].

Both humans and mice express SorCS1 in skeletal muscle and adipose tissue (**Figs. 4a and**

Extended Data Fig. 1c). We previously reported that up to 95% of SorCS1 is shed per hour from the

plasma membrane by ADAM17/TACE mediated cleavage[20,24]. Given SorCS1 and IR can bind each other

through an interaction in their extracellular domains (cf. **Fig. 3b and c**), we hypothesized that soluble

SorCS1 could have a glycemic regulatory function. To address this, we studied IR signaling in

differentiated myocyte cultures from *Sorcs1*$^{\Delta 25\text{-}28}$ mice after insulin stimulation in the presence of

increasing concentrations of exogenous sol-SorCS1 (**Fig. 4b**). Sol-SorCS1 augmented insulin induced

phosphorylation of IR and Akt and did so in a concentration dependent manner. No effect was observed

in the absence of insulin demonstrating that sol-SorCS1 is not an insulin mimetic. To establish a function

of sol-SorCS1 *in vivo*, we first compared the plasma concentration in a cohort of T2DM patients with

normoglycemic healthy individuals. In the diabetic group, but not among healthy individuals, plasma

glucose inversely correlated with sol-SorCS1 (**Fig. 4c**). Since differentiated 3T3-L1 adipocytes potently

liberates sol-SorCS1 (**Fig. 4d**) and human and murine adipose tissue express SorCS1 (**Fig. 4a, Extended**

data Fig. 5) alongside ADAM17/TACE[20], we examined, whether body mass index (BMI) associates

with the plasma concentration of sol-SorCS1. We found that BMI positively correlates with measures of

sol-SorCS1 both in diabetic individuals and healthy controls (**Extended Data Fig. 6**). The data suggested

that sol-SorCS1 may act as an adipokine to counteract the increased plasma glucose levels that commonly

follows obesity.

To establish a glucose lowering and insulin sensitizing effect of sol-SorCS1 *in vivo*, we

injected an adenovirus encoding human sol-SorCS1 (AV-sol-SorCS1) or AV-LacZ (encoding β-

galactosidase) as a negative control into the tail vein of obese leptin receptor knockout mice (*db/db*) that

develop severe T2DM. One week later, plasma glucose was increased by 55% and insulin by 65% in the

LacZ group due to the gradual worsening of the diabetic phenotype (**Fig. 4e-f**). Remarkably, in the AV-

sol-SorCS1 treated mice, plasma glucose remained unchanged and this despite a decrease in plasma

insulin. In an intraperitoneal glucose tolerance test (IPGTT), AV-sol-SorCS1 mice responded markedly

stronger with lower plasma glucose levels at all timepoints corresponding to a reduction in the area under

the curve (AUC) by 79.5% (**Fig. 4g-h**). The treated animals also exhibited superior ability in mobilizing

insulin compared to the control group (**Fig. 4i).** A similar glucose and insulin reducing effect was

obtained in adenovirus treated male mice with dietary induced T2DM (**Extended Data Fig. 7**). Next we

performed hyperinsulinemic euglycemic clamp[22]. AV-sol-SorCS1 mice showed lower basal plasma

glucose and increased capacity for glucose uptake compared to AV-LacZ treated animals (**Fig. 4j and**

k). Glucose tracer studies revealed that this effect was accounted for by increased insulin sensitivity in

peripheral tissues (**Fig. 4l, Extended Fig. 8**). Notably, three weeks post injection, glycated hemoglobin

(HbA1c) had increased considerably in the AV-LacZ group but it stayed unchanged in the AV-sol-

SorCS1 group indicating a long-lasting beneficial effect of the soluble receptor in reducing plasma

glucose (**Fig. 4m**).

165   In cells, full-length dimeric SorCS1 binds exclusively to the proIRs whereas monomeric

SorCS1 recognizes mature IR (**Fig 3e-h**). Hence, we speculated that the active form of sol-SorCS1

liberated by ADAM17 mediated cleavage is the monomer. To demonstrate this, we purified monomeric

and dimeric sol-SorCS1 from transfected CHO cells and injected *db/db* mice with each form

intraperitoneally 10 hrs and 1 hr prior to glucose tolerance test. Intriguingly, monomeric sol-SorCS1, but

not the dimeric receptor, profoundly improved glucose tolerance (**Fig. 4n**). Hence, the AUC for plasma

glucose was reduced by approximately 60% (from 1060 to 393 mM*min, P<0.05) in animals treated with

monomeric sol-SorCS1 (**Fig. 4o**).

In conclusion, we have demonstrated that full-length SorCS1, expressed in peripheral

tissues augments insulin sensitivity by engaging proIR and IR; the dimeric SorCS1 binds intracellular

proIR and the monomeric receptor the surface exposed mature IR. As a soluble protein, monomeric

SorCS1 can target tissues remote from where it is being produced (**Extended Data Fig. 9**). Given SorCS1

is expressed in adipose tissue, that it can be shed from adipocytes, and that its plasma concentration

positively correlates with BMI, we propose sol-SorCS1 as a novel adipokine to enhance insulin

sensitivity.

Acknowledgements

The study was funded by the Lundbeck Foundation grant no. R248-2017-431 (AN) , DANDRITE-R248-

2016-2518 (AN), R90-2011-7723 (AN); The Independent Research Fund Denmark grant no. DFF-7016-

00261 (AN); The Novo Nordisk Foundation grant no. NNF16OC0020984 (AN); The Danish National

Research Foundation grant no. DNRF133 (AN); The Danish Diabetes Academy (MK). We thank

Benedicte Vestergaard, Anja Aagaard and Anne Kerstine Thomassen for excellent technical assistance.

Figure 1: Insulin resistance and T2DM in *Sorcs1$^{Δ25-28}$* mice.

a, Plasma glucose and, **b**, insulin concentrations in aging female *Sorcs1$^{Δ25-28}$* mice (wildtype n=4,

Sorcs1$^{Δ25-28}$ n=9). **c**, Impaired glucose tolerance in 50 weeks old female *Sorcs1$^{Δ25-28}$* mice. **d**, AUC of the

data in **c** calculated minus baselines at t=0 (wildtype n=7, *Sorcs1$^{Δ25-28}$* n=10). **e**, In hyperinsulinemic

euglycemic clamp, glucose infusion rate (GIR) is lower in *Sorcs1$^{Δ25-28}$* female mice at steady state (t=60-

90 min). **f**, The insulin resistance is peripheral (wildtype n=7, *Sorcs1$^{Δ25-28}$* n=6). Data are presented as

mean ± SEM, * = P<0.05, ** = P<0.01.

Figure 2: Impaired insulin receptor signaling in *Sorcs1*$^{Δ25-28}$ muscle.

a, IR phosphorylation is reduced in 50 but not in 19 weeks old female mutant mice, 15 min post insulin

injection. **b**, Impaired insulin-induced IR phosphorylation in differentiated primary myocytes from

Sorcs1$^{Δ25-28}$ mice. Expression of desmin confirms the cultures are equally differentiated. **c**, Reduced

phosphorylation of AKT in *Sorcs1*$^{-/-}$ C2C12 cells. **d**, Surface biotinylation of *Sorcs1*$^{Δ25-28}$ primary

myocytes shows reduced IR$_β$ expression at the plasma membrane and accumulation of an intracellular

IR$_β$ immunoreactive fragment. Asterisk indicate the low molecular weight band. **e**, PNGase treatment

reveals reduced N-linked glycosylation of proIR in *Sorcs1*$^{Δ25-28}$ myocyte cultures. DG proIR designates

deglycosylated proIR.

Figure 3: SorCS1 and IR physically interact.

a, SorCS1 overexpression in HEK293 cells increases insulin affinity to IR. **b**, SPR analysis demonstrates dose-dependent binding of sol-SorCS to IR. **c**, Complex formation of SorCS1, IR, and insulin determined by SPR. At 1200 sec, the additive signals for Sol-SorCS1 (*blue*) and insulin (*red*) equals the response units obtained when IR is incubated with both proteins simultaneously (*black*). **d**, Co-immunoprecipitation between SorCS1 and proIR/ IR. The interaction is independent on the presence of saturating concentrations of insulin (100 nM). **e**, Binding of dimeric SorCS1 is restricted to proIR, but monomeric SorCS1 can bind mature IR. **f**, SorCS1 dimers that are BiFC-positive (*green*) are confined to cytosolic compartments, whereas monomeric and BiFC-negative SorCS1 (*red*), is present at the cell surface (**f**, *inset*). **g**, In non-permeabilized HEK293, PLA for SorCS1 and IR (*red*) demonstrates binding

219   between monomeric SorCS1 (BiFC-negative) and IR. **h**, Dimeric SorCS1 (BiFC-positive, *green*)

220   interacts with pro-IR (*red*) in cytosolic compartment.

Figure 4: Sol-SorCS1 is an insulin sensitizer.

a, mRNA expression of SorCS1 in human tissues (n=12/group). **b**, Insulin-induced activation of IR and

AKT in *Sorcs1*$^{Δ25-28}$ primary myocytes correlates with sol-SorCS1 concentration. **c**, In diabetic patients

sol-SorCS1 in plasma negatively correlates with fasting plasma glucose (**β**-value -2.6E-3, P<0.001). **d**,

Expression of full-length SorCS1 and shedding of its soluble form in 3T3-L1 adipocytes. **e**, Treatment

with AV-sol-SorCS1, but not AV-LacZ, reduces plasma glucose levels and, **f**, plasma insulin in 7 weeks

old *db/db* mice. **g**, Enhanced glucose uptake determined by IPGTT in AV-sol-SorCS1 treated *db/db* mice

shown in real-time and, **h**, as AUC (n=4/group). **i**, Plasma insulin in the IPGTT shown in g-h. **j**, Lower

fasting plasma glucose but, **k**, higher infusion rate in euglycemic hyperinsulinemic clamp of AV-sol-

SorCS1 treated *db/db* mice (n=5/group). **l**, The higher infusion rate is accounted for by increased

peripheral glucose uptake. **m**, Reduced HbA1c in *db/db* mice 21 days after infection with AV-sol-SorCS1

234   infection. **n**, Intraperitoneal injection of recombinant monomeric, but not dimeric, sol-SorCS1 increases

235   glucose tolerance (IPGTT) in *db/db* mice. **o**, AUC of the data displayed in panel **n**. Data are presented as

236   means ± SEM, * = *P*<0.05, ** = *P*<0.01, ***=*P*<0.001.

References:

1. Shanik, M.H*., et al.* Insulin resistance and hyperinsulinemia: is hyperinsulinemia the cart or the horse? *Diabetes Care* **31 Suppl 2**, S262-268 (2008).

2. Warram, J.H., Martin, B.C., Krolewski, A.S., Soeldner, J.S. & Kahn, C.R. Slow glucose removal rate and hyperinsulinemia precede the development of type II diabetes in the offspring of diabetic parents. *Ann Intern Med* **113**, 909-915 (1990).

3. Kim, S.J*., et al.* mTOR complex 2 regulates proper turnover of insulin receptor substrate-1 via the ubiquitin ligase subunit Fbw8. *Mol Cell* **48**, 875-887 (2012).

4. Tremblay, F*., et al.* Identification of IRS-1 Ser-1101 as a target of S6K1 in nutrient- and obesity-induced insulin resistance. *Proc Natl Acad Sci U S A* **104**, 14056-14061 (2007).

5. Blagosklonny, M.V. TOR-centric view on insulin resistance and diabetic complications: perspective for endocrinologists and gerontologists. *Cell Death Dis* **4**, e964 (2013).

6. Willnow, T.E., Petersen, C.M. & Nykjaer, A. VPS10P-domain receptors - regulators of neuronal viability and function. *Nat Rev Neurosci* **9**, 899-909 (2008).

7. Willnow, T.E., Kjolby, M. & Nykjaer, A. Sortilins: new players in lipoprotein metabolism. *Curr Opin Lipidol* **22**, 79-85 (2011).

8. He, Y., Fang, Z. & Yu, G. Sortilin-related VPS10 domain containing receptor 1 and Alzheimer's disease-associated allelic variations preferentially exist in female or type 2 diabetes mellitus patients in southern Han Chinese. *Psychogeriatrics* **12**, 215-225 (2012).

9. Paterson, A.D*., et al.* A genome-wide association study identifies a novel major locus for glycemic control in type 1 diabetes, as measured by both A1C and glucose. *Diabetes* **59**, 539-549 (2010).

10.   Goodarzi, M.O.*, et al.* SORCS1: a novel human type 2 diabetes susceptibility gene suggested by

the mouse. *Diabetes* **56**, 1922-1929 (2007).

11.   Granhall, C., Park, H.B., Fakhrai-Rad, H. & Luthman, H. High-resolution quantitative trait locus

analysis reveals multiple diabetes susceptibility loci mapped to intervals<800 kb in the species-

conserved Niddm1i of the GK rat. *Genetics* **174**, 1565-1572 (2006).

12.   Clee, S.M.*, et al.* Positional cloning of Sorcs1, a type 2 diabetes quantitative trait locus. *Nat Genet*

38, 688-693 (2006).

13.   Kebede, M.A.*, et al.* SORCS1 is necessary for normal insulin secretory granule biogenesis in

metabolically stressed beta cells. *J Clin Invest* **124**, 4240-4256 (2014).

14.   Kjolby, M.*, et al.* Sort1, encoded by the cardiovascular risk locus 1p13.3, is a regulator of hepatic

lipoprotein export. *Cell Metab* **12**, 213-223 (2010).

15.   Kjolby, M., Nielsen, M.S. & Petersen, C.M. Sortilin, encoded by the cardiovascular risk gene

SORT1, and its suggested functions in cardiovascular disease. *Curr Atheroscler Rep* **17**, 496

(2015).

16.   Mortensen, M.B.*, et al.* Targeting sortilin in immune cells reduces proinflammatory cytokines

and atherosclerosis. *J Clin Invest* **124**, 5317-5322 (2014).

17.   Vaegter, C.B.*, et al.* Sortilin associates with Trk receptors to enhance anterograde transport and

neurotrophin signaling. *Nat Neurosci* **14**, 54-61 (2011).

18.   Hermey, G.*, et al.* SorCS1 variants and amyloid precursor protein (APP) are co-transported in

neurons but only SorCS1c modulates anterograde APP transport. *J Neurochem* **135**, 60-75 (2015).

19.   Savas, J.N.*, et al.* The Sorting Receptor SorCS1 Regulates Trafficking of Neurexin and AMPA

Receptors. *Neuron* **87**, 764-780 (2015).

20.   Hermey, G., Sjogaard, S.S., Petersen, C.M., Nykjaer, A. & Gliemann, J. Tumour necrosis factor

282   alpha-converting enzyme mediates ectodomain shedding of Vps10p-domain receptor family

283   members. *Biochem J* **395**, 285-293 (2006).

284   21.   Ayala, J.E.*, et al.* Standard operating procedures for describing and performing metabolic tests

285   of glucose homeostasis in mice. *Dis Model Mech* **3**, 525-534 (2010).

286   22.   Graae, A.S.*, et al.* ADAMTS9 Regulates Skeletal Muscle Insulin Sensitivity Through

287   Extracellular Matrix Alterations. *Diabetes* **68**, 502-514 (2019).

288   23.   Hwang, J.B. & Frost, S.C. Effect of alternative glycosylation on insulin receptor processing. *J

289   Biol Chem* **274**, 22813-22820 (1999).

290   24.   Hermey, G.*, et al.* Characterization of sorCS1, an alternatively spliced receptor with completely

291   different cytoplasmic domains that mediate different trafficking in cells. *J Biol Chem* **278**, 7390-

292   7396 (2003).

293   25.   Januliene, D.*, et al.* Hidden Twins: SorCS Neuroreceptors Form Stable Dimers. *J Mol Biol* **429**,

294   2907-2917 (2017).

295   26.   Kerppola, T.K. Design and implementation of bimolecular fluorescence complementation (BiFC)

296   assays for the visualization of protein interactions in living cells. *Nat Protoc* **1**, 1278-1286 (2006).

Figure 1

Figure 2

a Murine Muscle

Sorcs1$^{+/+}$ Sorcs1$^{Δ25-28}$

IR$_β$
pIR$_β$
19 Weeks

Sorcs1$^{+/+}$ Sorcs1$^{Δ25-28}$

IR$_β$
pIR$_β$
50 Weeks

b Primary Myocytes

Sorcs1$^{+/+}$ Sorcs1$^{Δ25-28}$

Insulin (nM)   0   0.1   1.0   10   0   0.1   1.0   10

pIR$_β$
IR$_β$
Desmin

c C2C12 cells

Insulin (nM)   0 1.0 2.5 5.0 7.5 10 25 50 100 200 500 1000

pAKT   Sorcs1$^{+/+}$
pAKT   Sorcs1$^{-/-}$

EC50
Sorcs1$^{+/+}$ = 14.19
Sorcs1$^{-/-}$ = 3.05

d IR degradation myocytes

Sorcs1$^{+/+}$ Sorcs1$^{Δ25-28}$

Total   Intra   Surface   Total   Intra   Surface

IR$_β$
*

e proIR glycosylation myocytes

Sorcs1$^{+/+}$ Sorcs1$^{Δ25-28}$ Sorcs1$^{Δ25-28}$ Sorcs1$^{Δ25-28}$ PNGase

proIR
DG proIR

Figure 3

Figure 4

a *SORCS1* tissue expression

b *Sorcs1*$^{\Delta 25-28}$ myocytes

c Plasma SorCS1

d 3T3-L1 SorCS1 protein

e Plasma glucose

f Plasma insulin

g IPGTT glucose

h IPGTT AUC

i IPGTT insulin

j Clamp plasma glucose

k Glucose infusion rate

l Peripheral tissue

m HbA1c

n IPGTT glucose

o IPGTT AUC